Genome-wide association study of water use patterns of common bean (*Phaseolus vulgaris*
L.) genotypes in response to drying soil

Henry Cordoba-Novoa[1], James D. Kelly[2], Valerio Hoyos-Villegas[1]*

[1]McGill University, Montreal, Canada

[2]Michigan State University, East Lansing, MI, USA

Corresponding author: valerio.hoyos-villegas@mcgill.ca

Abstract

Common bean (*Phaseolus vulgaris* L.) is the main legume crop for direct human consumption
worldwide, especially in developing countries. Among the abiotic factors affecting common
bean production, drought is the most limiting condition. Capacity for the accurate identification
and selection of drought-tolerant genotypes is a paramount step in the improvement of new
cultivars. This study is aimed at characterizing the genetic variability and genetic architecture of
transpiration, stomatal regulation and whole plant water use within the Mesoamerican common
bean germplasm. Three greenhouse experiments were conducted, and water use was recorded
gravimetrically. Transpiration of water-stressed plants was normalized against well-watered
plants to obtain a normalized transpiration rate (NTR). A critical fraction of transpirable soil
water (FTSWc) was estimated as the inflection point at which NTR starts decreasing linearly.
Genome-wide association (GWA) analyses for average NTR and FTSWc were performed. High
variation in average NTR and FTSWc was found among genotypes. Genomic signals (QTL)
controlling the variation of these traits were identified on *Pv*01 and *Pv*07 some located in
intergenic, intronic and exonic regions. A set of novel candidate genes and putative regulatory
elements located in these QTL were identified. Most of the genes have been previously reported
to be involved in abiotic tolerance, including some of the five transcription factors identified.
Candidate genes can be used for marker-assisted selection and further study of drought tolerance
mechanisms in common bean. Our results provide tools to accelerate the drought tolerance
improvement in the crop.

Keywords

Breeding, abiotic stress, drought, GWAS, transcription factor

Word count: 4880

Introduction

Pulses are a rich source of dietary fiber, micronutrients, vitamins, carbohydrates, and mono- and poly-unsaturated fatty acids (McCrory et al., 2010). Among pulses, common bean (*Phaseolus vulgaris* L.) is the most cultivated pulse crop for direct human consumption (FAOSTAT, 2019). This crop represents a valuable commodity to ensure food security worldwide, mainly in developing countries. In climate change scenarios, increasing temperatures and the reduction of water availability pose serious threats to food production (Fereres and Soriano, 2007; OECD, 2014). Common bean yield is affected by drought (Beebe et al., 2014), therefore the development of drought-tolerant cultivars is an urgent need in breeding programs (Miklas et al., 2006).

One of the main plant responses to water limitations is reducing transpiration to avoid excessive water loss (Osakabe et al., 2014). Transpiration is limited by partial stomatal closure in response to increased atmospheric vapor pressure deficit (VPD) and reduction of the fraction of transpirable soil water (FTSW) (Sanchez et al., 2021). An early plant response to drying soil will conserve water content for late in the season, especially during grain filling, preventing the wilting genotype and improving plant drought-resistance (Jafarikouhini et al., 2020; Chiango et al., 2022). The selection of genotypes with improved capacity to maintain water status is crucial to developing new drought-tolerant cultivars (Ray and Sinclair, 1997; Egan et al., 2021). The limited-transpiration trait is now included in commercial soybean (*Glycine max* Merr. L.) (Carter et al., 2016)and maize (*Zea mays* L.) (Gaffney et al., 2015)cultivars for increased yield under water limiting conditions.

The identification of differences in traits such as transpiration among genotypes can be challenging as they show high plasticity across environments in multiple plant species (Ritchie, 1980). A methodology to identify patterns in water use is to analyze the normalized transpiration rate (NTR) based on soil water content, and the critical fraction of transpirable soil water (FTSWc). Comparisons based on soil water content are more reliable to detect differences between genotypes attributable to actual variation in plant physiology (Ray and Sinclair, 1997). The FTSWc indicates the plant-transpirable water content in soil at which the plant starts closing stomata and reducing transpiration rates. Therefore, FTSWc allows the characterization of genotypes based on their response. FTSW is centered between 0 and 1, where relatively low values of FTSWc indicate that the plant maintains transpiration rates longer and may continue growing under long periods of stress. In contrast, germplasm with relatively high FTSWc thresholds are more sensitive to water deficit and close stomata earlier, making them suitable for intermittent drought stress periods (Egan et al., 2021). High FTSWc may also be evidence for germplasm with faster recovery rates between intermittent drought periods.

The analysis of NTR and FTSWc has been used to characterize stomatal regulation dynamics in maize (Ray and Sinclair, 1997; Devi and Reddy, 2020b); genotypic variation in cotton (Devi and Reddy, 2020a), sorghum (Choudhary and Sinclair, 2014), white clover (Egan et al., 2021); and pearl miller cultivars (*Pennisetum glaucum* (L.) R. Br.) (Kholová et al., 2010a). To date, no GWA have been performed to attempt to identify candidate genes involved in the regulation of these constitutive water-conserving mechanisms.

75　The identification of candidate genes is a valuable resource to design and implement marker-
76　assisted selection (MAS) strategies that allow a rapid screening and selection of genotypes. The
77　overall objective of this study was to characterize the response of transpiration in Mesoamerican
78　bean accessions to drying soil and identify candidate genes using GWA.

Materials and methods

Plant material and experiment design

A subset of 82 Mesoamerican bean genotypes (hereafter drought diversity panel, DDP) from the
larger Mesoamerican diversity panel (MDP, 498 individuals) was used in the present study. The
DDP is a selected group of varieties, lines, and landraces that differ in their response to drought
(Mukeshimana et al., 2014). Three independent greenhouse experiments were conducted at
Michigan State University in 2013 (repetitions one and two) and 2014 (repetition three), with a
duration 26, 37, and 32 days after treatment (DAT), respectively.

For each repetition trial, treatments consisted of irrigation (control), and drought with three
replicates in a randomized complete design. Prior to the beginning of the treatments, pots were
saturated and allowed to drain overnight. The next morning, pots were sealed at the bottom and
top to avoid any water loss from runoff and evaporation. At the beginning of the experiment the
saturated pot weights were determined, and daily pot weights were recorded. Plants under well-
watered conditions were replenished with the exact amount of water that had been transpired in a
24-hr period. Drought treatments underwent a progressive drydown with no water replenishment.
The drought treatment finished once all the cultivars had died. Further details on this
methodology can be found in Ray and Sinclair (1997), Egan et al. (2021), and Chiango et al.
(2022).

Transpiration rate and FTSW

Transpiration was analyzed using the combined data from the three experiments. The
transpiration was analyzed following the methodology proposed by Ray and Sinclair (1997) and
measured as the difference in pot weights every 24 hours. In days when there was no deficit (e.g.
due to cloudy conditions between days), values were eliminated for that replicate. The
transpiration rate (TR) was calculated as the ratio between the weight of droughted pots and
well-watered pots on each day. A second normalization was applied to TR using the average TR
between days two and five (well-watered period) according to the formula:

$$NTR = \frac{TR}{Day\ 2 - 5\ average\ TR}$$

105　During days 2 – 5, all the plants were still well-watered, as sufficient soil water was available.
106　With this normalization, plants at full transpiration would have a normalized TR (NTR) near to
107　1.

108　To determine the extent of water extracted by plants, the fraction of transpirable soil water
109　(FTSW) was calculated as:

$$FTSW = \frac{(Daily\ pot\ weigth - Final\ pot\ weight)}{(Pot\ weigth\ on\ 1\ DAT - Final\ pot\ weight)}$$

110 The relationship between NTR and FTSW was obtained by fitting the equation previously
111 described by Muchow and Sinclair, (1991):

$$NTR = \frac{1}{[1 + A \times \exp(B \times FTSW)]}$$

112 Where A and B are empirical coefficients generated for each genotype through curve fitting. A
113 linear-plateau regression was used to determine the critical FTSW (FTSWc) where NTR starts
114 decreasing linearly and indicates the point at which stomata start to close in stressed plants.
115 FTSWc was determined for each genotype in each replicate.

Statistical analysis

117 The data from the three experiments were combined and analyzed together in a randomized
118 complete block design (RCBD) with average values across replications within repetitions as
119 replicates. One-way analysis of variance (ANOVA) was conducted using the following model:

$$y_{ijk} = \mu + \alpha_i + \beta_j + e_{ij}$$

120 Where μ is the grand mean, α_i is the effect of the *ith* genotype, β_j is the effect of the *jth*
121 repetition experiment, and e_{ij} is the random error. Best linear unbiased predictors (BLUPs) were
122 estimated with the restricted maximum likelihood (REML) method using the *lme4* package in R
123 v4.1 (Bates et al., 2015) and BLUP-adjusted means were calculated. BLUP-adjusted means were
124 compared by the least-squares method (α=0.05). Broad-sense heritability (H^2) was estimated by:

$$H^2 = \frac{\hat{\sigma}_g^2}{\hat{\sigma}_g^2 + \frac{\hat{\sigma}_e^2}{n}}$$

125 Where $\hat{\sigma}_g^2$ is the genotypic variance, $\hat{\sigma}_e^2$ is the error variance, and n is the number of replicates
126 (experiments).

Genome-Wide Association Analyses

Genotypic data

The DDP was previously genotyped (Moghaddam et al., 2016). The genotypic data contains
around 200K single nucleotide polymorphisms (SNPs) and can be found on the BeanCAP
website (http://arsftfbean.uprm.edu/beancap/research/, Accessed Jun. 2022). SNPs with a minor
allele frequency (MAF) < 0.02 were filtered. This threshold was chosen to facilitate the

identification of rare genotypes. Linkage disequilibrium (LD) between markers was calculated
using all the SNPs within a sliding window of 50 SNPs in TASSEL 5 software (Bradbury et al.,
2007).

The LD decay pattern for each chromosome and whole genome was analyzed by nonlinear
regression fitting the expectation of the parameter r^2 and the distance between each pair of
markers (Hill and Weir, 1988; Remington et al., 2001). Population structure was evaluated using
the variational Bayesian inference in fastSTRUCTURE (Raj et al., 2014) with two populations
(K = 2) based on previous knowledge of the subpanel (Hoyos-Villegas et al., 2017).

GWAS

A GWAS analysis was performed using 182,243 SNPs with a MAF > 0.02, and the BLUP-
adjusted means of the FTSWc and average NTR under water deficit for each genotype. Analyses
were carried out using multiple models, including Mixed Linear Model (MLM), compressed
MLM (CMLM), Settlement of MLM Under Progressively Exclusive Relationship (SUPER),
Fixed and random model Circulating Probability Unification (FarmCPU), and Bayesian-
information and LD Iteratively Nested Keyway (BLINK) implemented in GAPIT v.2 (Tang et
al., 2016). Individual relatedness was also considered with the kinship matrix that is
automatically calculated within GAPIT using the VanRaden method (VanRaden, 2008). Peak
SNPs with *p*-values above a Benjamini-Hochberg false discovery rate (FDR) threshold of 0.05
were considered significant and were further examined. The variance explained by significant
SNPs was estimated by the coefficient of determination (R^2) of a linear regression model. The
model included all the five PCs from PC analysis (PCA), the SNP marker parameterized in
accordance with the gene action determined using the general, additive, and simplex dominance
models in the R package GWASpoly (Rosyara et al., 2016), and a vector of random residuals
(Wallace et al., 2016).

Haplotype analysis and candidate genes

Haplotype blocks were defined using the solid spine of LD method implemented in Haploview
v.4.2 software with default parameters (Barrett et al., 2005). The proportion of the phenotypic
variance explained by significant SNPs from GWAS (R^2) was recalculated after including the
SNPs present in the same haplotype block.

To identify candidate genes, JBrowse on Phytozome v13 (Goodstein et al., 2012) was used to
browse the common bean reference genome v.2.1 (Schmutz et al., 2014) within a ± 100-Kb
window centered on each associated SNP. Candidate genes were selected based on their
annotation and literature review related to drought and other abiotic stresses.

Results

Transpiration rate response to drying soil and heritability

The average NTR per genotype during the evaluation period ranged from 0.33 in Yolano to 0.96
in ABCP-8, with a global mean of 0.52 (Table 1) and an H^2 of 0.53.

There was a consistent relationship between NTR and FTSW. The NTR value decreased with a linear tendency below the FTSWc for each genotype. The FTSW threshold varied from 0.55 in the pink bean genotype Harold to 0.97 in the black bean T-39 with an overall mean of 0.76 and significant differences between the maximum and minimum ($p<0.05$; Table 1). Figure 1 depicts the average response of NTR to a decreasing FTSW for the bottom (average FTSWc = 0.58; Figure 1A) and top five (average FTSWc = 0.94; Figure 1B) genotypes. Using the genotypic and error variances, the estimated H^2 for FTSWc was 0.38.

Population structure and LD decay

Two subpopulations were identified in the DDP (K=2) with varying probabilities of ancestry among genotypes. Admixture between the two subpopulations was also observed. The chromosome-wise analysis of the LD decay pattern showed that in the DDP, LD decayed at a range from 135.5 Kbp in *Pv04* to 848.7 Kbp in *Pv09* with a genome-wise decay distance of 363.3 Kbp.

Genome-Wide Association and haplotype analysis

Based on the adjustment of the models in the QQ plots, all the tested models performed similarly with a slight improvement in Blink for FTSWc (Figure S1) and FarmCPU for average NTR (Figure S2). Blink and FarmCPU were used for the analysis of marker-trait associations for FTSWc and average NTR, respectively. FarmCPU and Blink have also shown improved statistical power with both simulated and real datasets (Huang et al., 2019; Kaler et al., 2020).

In FTSWc, one SNP was identified on *Pv01* at 33,516,386 bp (C/T; *Pv01*FTSWc) and another on *Pv07* at 9,224,979 bp (C/T; *Pv07*FTSWc) (Figure 2A). Each SNP for FTSWc had an effect of +0.26 units and explained the 31% of the phenotypic variation. For average NTR, two significant SNPs were identified in *Pv01* at 28,451,384 bp (C/A; *Pv01*NTR_1) and 35,489,136 bp (A/G; *Pv01*NTR_2) with an effect of -0.22 and +0.19 units, respectively (Figure 2B). *Pv01*NTR_1 explained the 23% of the phenotypic variation (R^2), and *Pv01*NTR_2 the 24%. The four SNPs were detected using the multiple models and had a MAF of 0.02.

The genotype T-39 that had the highest FTSWc was homozygous for the alternative allele (TT) of *Pv01*FTSWc and *Pv07*FTSWc. NW-63 was heterozygous for the significant SNPs of FTSWc and was among the top five values. Harold, lowest FTSWc (0.55) was homozygous for the reference alleles (CC). For *Pv01*NTR_1, ABCP-8 is homozygous (AA) for the alternative allele and had the highest average NTR (0.96), followed by A-55 which is heterozygous (A/G). ABCP-8 and A-55 are homozygous (GG) for the alternative allele of *Pv01*NTR_2. Yolano is homozygous for the reference allele (GG) and had the lowest average NTR.

In the haplotype analysis for Pv01, the SNP *Pv*01FTSWc was grouped with 29 additional SNPs in a haplotype block of 308 Kbp. Out of the 30 SNPs in the haplotype, 15 were in genic regions and the others in intergenic regions. The haplotype containing *Pv*01FTSWc explained a higher variation (42%) compared to the single SNP (31%). In aberage NTR, *Pv01*NTR_1 was grouped with 48 additional SNPs where the marker ss01_28185985 is in a coding region. The block had a size of 462 Kbp and explained a higher total phenotypic variation (36%) compared to the single

SNP (23%). *Pv01*NTR_2 was in a block of 454 Kbp with four SNPs, but no increases in the R^2
were observed for the haplotype compared to the individual significant SNP.

For haplotypes on Pv07, *Pv07*FTSWc was grouped with 6 additional markers in a haplotype of
464 Kbp with the SNP ss07_9279035 located on a genic region. For this haplotype, no increases
in the variation explained were detected.

Candidate genes

Using a 100 Kbp window, four genes were associated with the haplotype of *Pv01*FTSWc, 12
with the haplotype for *Pv07*FTSWc, three genes with the haplotype containing *Pv01*NTR_1, and
four with the haplotype containing *Pv01*NTR_2. Considering their annotations and potential role
in abiotic stress responses, genes were filtered, and seven candidate genes were identified on
Pv01, and seven on *Pv07* for limited transpiration in common bean (Table 2). Interestingly, four
of the genes are annotated as transcription factors (TFs). A TF on *Pv01* was bHLH106-related
(*Phvul.001G121200*); and on *Pv07*, TFs were UPBEAT1 (*Phvul.007G089100*), No apical
meristem (NAM) protein (*Phvul.007G089600*) on *Pv07*, and C2H2-type Zn-finger protein
(*Phvul.007G090500*).

Discussion

The present study investigated the transpiration rate response of common bean and genotypic
differences among Mesoamerican accessions. Under drought-like conditions, stomata closure is
among the first physiological responses to avoid water loss through transpiration (Osakabe et al.,
2014). Stomatal closure under drought is mainly triggered by an increase in the abscisic acid
(ABA) concentration in leaves (Bharath et al., 2021). However, it can also be induced by
hormones like salicylic acid (SA), methyl jasmonate (MJ) and ethylene; metabolites such as
proline and polyamines; and gaseous signals such as hydrogen sulfide (H_2S) (Agurla et al.,
2018). Stomatal closure triggered by these components is mediated by hydrogen peroxide
($H2O_2$), calcium (Ca^{2+}), and nitric oxide (NO) (Agurla et al., 2018). Likewise, drought causes the
ABA-dependent and independent activation of key transcription factors (TFs) (Nakashima et al.,
2014).

Stomatal limitations restrict gas exchange rate, photosynthetic activity, and plant growth and
development. Drought tolerant varieties are expected to minimize physiological and yield
penalties despite water shortage and can be identified based on a slow leaf wilting phenotype.
The soybean cultivar PI 471938 is an example of a slow wilting genotype that exhibits water-
conservative traits by maintaining TR (Fletcher et al., 2007). However, a reduced TR is usually
observed in drought tolerant genotypes compared to drought sensitive materials (Kholová et al.
2010) for short drought periods. Reductions in TR are associated with early stomatal closure and
high FTSW thresholds (Ray and Sinclair, 1997; Egan et al., 2021). For PI 471938 has been ruled
out that its resistance is due to differences in FTSWc values compared to non-resistant genotypes
(Sadok et al., 2012). Implying additional mechanisms involved in water conservation and may
not be fully detected by differences in the critical FTSW.

Among the genotypes with the lowest average NTR (Table 1), Othello has been reported as a
drought resistant cultivar, but US-1140 and UI-59 as low yielding cultivars (Muñoz-Perea et al.,
2006), which suggests that the stomatal responses of common bean to water deficit have varying
impacts on plant performance and is influenced by other traits depending on the cultivar.

A lower TR indicates that the plant favors a conservative water use mechanism and survival for
longer periods (Kholová et al. 2010). Devi and Reddy (2020) observed a positive correlation
between TR and FTSWc when studying different cotton accessions. In contrast, we observed no
relationship between average NTR and FTSWc since the genotypes with the lower average NTR
did not have a higher FTSWc (Table 1). This can be explained by genotypic differences in
stomatal and hydraulic conductance among the evaluated genotypes. Hydraulic conductance
affects stomatal conductivity, transpiration rate, and water potential, and its variation depends on
anatomic changes in membrane thickness, structure, and permeability (Hacke and Way, 2014;
Simonin et al., 2015). Adjustments in hydraulic conductance may also depend on changes in
water transport in roots and shoot via aquaporins (Choudhary and Sinclair, 2014). In French bean
leaves, Sôber, (1998) observed that high stomatal conductance rates were accompanied by a high
hydraulic conductivity. Additionally, Choudhary and Sinclair (2014) found that differences in
TR among sorghum genotypes result from a varying hydraulic conductance where some
genotypes with low TR can maintain high rates of both stomatal and hydraulic conductance. The
slow wilting soybean PI 471938 has been found to have a limited hydraulic conductance in
leaves while maintaining constant TR(Sinclair et al., 2008). It is possible then, that the genetic
variation in the evaluated panel allows some cultivars to have an early stomatal closure (high
FTSWc) with an improved transpiration and hydraulic conductance. In contrast, low FTSWc
may provide a plant with the ability to conserve water under intermittent drought conditions, but
this has to be coupled with an equivalent rate of stomatal reopening upon rehydration. In
addition, growth and development (as they pertain to seed yield) must not be severely
compromised, placing further emphasis on efficiencies between photosynthetic rate and CO_2
exchange under intermittent and terminal drought conditions.

Plants with high TRs may keep stomata open longer in drying soil, which results in lower
FTSWc values (Ray and Sinclair, 1997). Genotypes with low FTSWc maximize water use and
are desirable for short-term water deficit periods (Egan et al., 2021). Harold had the lowest
threshold (0.55) and has been reported to have a moderate to high level of resistance to drought
(Singh et al., 2001). Common Red Mexican (FTSWc=0.58) is a landrace that exhibits high level
of water stress tolerance (Muñoz-Perea et al., 2007), despite having a moderate average NTR in
the present study (0.54). As explained above, a varying dynamic between TR and FTSWc may
be explained by genetic differences in stomatal and hydraulic conductance. We did observe that
ABCP-8 had the highest average NTR (0.96) and one of the lowest FTSWc (0.58; Table 1).
ABCP-8 is a pinto bean line bred for common bacterial blight resistance by crossing the donor
XAN 159 and the recurrent parent 'Chase' pinto (Mutlu et al., 2005). It is worth noting that the
line XAN 159 was developed using interspecific crosses with tepary bean (*Phaseolus acutifolius*
A. Gray) (Thomas and Giles Waines, 1984), which has been referred as a source of resistance for
heat and drought stress (Souter et al., 2017; Moghaddam et al., 2021). It is possible that genetic
elements involved in the response of the cultivar to drying soil and response to drought were
introgressed into ABCP-8 from tepary bean accessions.

Drought tolerance is a complex trait with high variation in multiple structural, physiological, and metabolic components (Bowles et al., 2021). Similar to our results, previous studies on drought responses in common bean have identified QTL on Chr01 and Chr07 (Mukeshimana et al., 2014; Briñez et al., 2017; Dramadri et al., 2019). In a biparental population of Andean genotypes, Sedlar et al. (2020) identified QTL for water potential mainly in linkage groups one and seven. Leitão et al. (2021) found significant associations for stomatal conductance under different water regimes in Portuguese accessions. However, they found no associations on Chr07, and one SNP was located on Chr01 in a different locus compared to our results.

Depending on the breeding purpose, once validated, the identified SNPs can be used for marker-assisted selection (MAS). For short drought periods, selections favoring the alternative alleles (T) for *Pv01*FTSW and *Pv07*FTSW would increase the FTSWc and allow an early stomatal closure. In contrast, if breeding for long drought periods, selection against the alternative allele would be desirable since the plant will keep stomata open, reducing the impact on photosynthesis activity and growth. Nevertheless, this selection must be done considering that a limited NTR would be necessary to avoid excessive loss of water. In this case, a positive selection for the alternative allele (A) of *Pv01*NTR_1 and a selection against the alternative allele (G) in *Pv01*NTR_2 should favor a reduced NRT.

The candidate gene *Phvul.001G108800* encodes for an aquaporin SIP2-1-Related protein (Table 2). In Arabidopsis, the aquaporin SIP2-1 protein is localized in the endoplasmic reticulum (ER) and has been associated with the alleviation of ER stress (Sato and Maeshima, 2019). ER stress results from the accumulation of misfolded or unfolded proteins in the ER during environmental stresses, including drought (Deng et al., 2013).

Thioredoxins (Trx) are thiol:disulfide oxidoreductase proteins involved in the formation or reduction of disulfide bonds (Eklund et al., 1991). This class of proteins can be induced by water deficit in *Solanum tuberosum* (Rey et al., 1998) and have been proposed as candidate genes for salt tolerance in bread wheat (Chaurasia et al., 2020). The candidate gene encoding for a Thioredoxin-Like 1-1 protein (*Phvul.001G109200*) may provide drought tolerance in common bean by regulating the redox state of chloroplast proteins. Recently, Yokochi et al. (2021) demonstrated in Arabidopsis that Trx and Trx-like proteins are involved in the regulation of chloroplast functions, including photosynthesis-related proteins.

*Pv01*NTR_2 had three candidate genes related to abiotic stress (Table 2). Extensins (EXTs) are proline-rich proteins that play a paramount role in cell wall synthesis and modification during root and pollen development (Kishor et al., 2015). Proline is an amino acid that accumulates under stress conditions as a beneficial solute to reduce the osmotic potential and help cope with the effects of drought (Chul et al., 2018). The candidate Phosphatidylinositol N-acetylglucosaminyltransferase subunit Q protein (PIG-Q) (*Phvul.001G127800*) is part of the GPI-GlcNAc-transferase (GPI-GnT) multiprotein complex involved in the glycosylphosphatidylinositol (GPI)-anchor biosynthesis in plants (Beihammer et al., 2020). The GPI-anchor is a post-transcriptional modification of diverse protein families implicated in cell wall composition, cell expansion, biotic and abiotic stress response, stomata development, pollen tube elongation, and hormone signaling (Beihammer et al., 2020). Variations in PIG-Q may

change the modification of key proteins in the water deficit response. However, further
investigation would be necessary.

Water stress increases reactive oxygen species (ROS) leading to oxidative stress. Nitric oxide
(NO) limits the accumulation of ROS by inhibiting the Nicotinamide adenine dinucleotide
phosphate (NADPH) oxidase enzyme complex, reacting with ROS, and inducing the expression
of antioxidant enzymes genes (del Río, 2015). Furthermore, NO plays a key role in the abscisic
acid (ABA)-induced stomatal closure during stress (Perlikowski et al., 2022). The nitric-oxide
synthase (NOS) identified (candidate gene *Phvul.001G128000*) is one of the main enzymes
involved in NO production (Neill et al., 2003) that may alleviate the oxidative stress and regulate
an early stomatal closure. Cao et al. (2019) showed that an early NOS-mediated nitric oxidative
increase mitigates oxidative damage in rice roots under drought.

Alkaline ceramidases have been poorly studied in plants. However, the overexpression of an
alkaline ceramidase in Arabidopsis promotes autophagy under abiotic stresses (Zheng et al.,
2018). Autophagy is a conserved pathway that plays a critical role in the adaptation of plants to
drastically changing environmental stresses (Han et al., 2011). Particularly, the alkaline
ceramidase AtACER is involved in disease resistance and salt tolerance (Wu et al., 2015). The
Alkaline ceramidase-related candidate gene (*Phvul.007G090000*) is a novel target in common
bean to study the role of alkaline ceramidases in crop plants, and also for drought tolerance
improvement.

The transport of auxins plays an important role in plant growth and development, and is
regulated by influx (AUXIN1/LIKE-AUX1 family) and efflux (mainly PIN-Formed (PIN))
carries (Swarup and Bhosale, 2019). The candidate Auxin efflux carrier component 1
(*Phvul.001G121100,* Table 2) is encoded by the PIN1 gene belonging to the type-I PIN proteins
(Armengot et al., 2016). The downexpression of PIN1 and other members of the type-I PIN
subfamily has shown to affect root meristem size during salt stress by increasing NO levels
(Korver et al., 2018). The potential role of PIN1 in common bean is in accordance with the
candidate NOS gene detected also on Chr01. Furthermore, Zhang et al., (2012) showed that in
rice PIN3t, another type-I PIN, is involved in drought stress response and tolerance.

On Pv07, the candidate gene *Phvul.007G088800* encodes the ATP-binding Cassette Transporter
G family member 25 (ABCG25). ABCG25 is an ABA exporter expressed in vascular tissue that
controls cellular ABA levels (Park et al., 2016) and influences stomatal regulation (Kuromori et
al., 2010). ABA is the main plant hormone involved in abiotic stress responses, seed processes,
and senescence. Particularly, long-distance ABA signaling and its interplay with other processes
are involved in the detection of drying soil and plant responses (Davies et al., 2005), which
agrees with our experiments. ABCG25 and other transporters have been shown to respond to
abiotic stresses including temperature (Baron et al., 2012) and drought (Chen et al., 2011).

Phosphorylation is one of the main signaling mechanisms involved in a myriad of plant
responses. The candidate genes *Phvul.007G089200* and *Phvul.007G089301* encoding for a
Protein Tyrosine kinase (PTK) and a Calcium-dependent protein kinase (CPK) 11-related
protein, respectively, may be involved in the response to drought stress. The role of PTKs has
been mostly studied in animals and humans and remains poorly understood in plants. Recently,

(Allimuthu et al., 2020) characterized a set of PTKs in rice, identifying key genes implicated in
the osmotic and drought stress response. CPKs also phosphorylate Tyr residues of beta-tubulin
and several transcription factors (TF) in Arabidopsis (Nemoto et al., 2015). Interestingly, many
of these TFs are related to ethylene and ABA, and have shown to be relevant for plant response
to abiotic stresses (reviewed in Miyamoto et al., 2019).

Five candidate genes involved in gene expression regulation were identified. Mitochondrial
transcription termination factor (mTERF) proteins (*Phvul.007G089900*) are key components of
organellar gene expression (OGE) machinery in chloroplasts and mitochondria (Barkan, 2011).
Despite plants having more mTERF genes compared with animals, their role is still poorly
understood. Studies in maize and Arabidopsis have shown that mTERFs regulate different
developmental processes, including the response to abiotic stresses. For instance, in Arabidopsis,
mTERF mutants display varying sensitivity to heat and saline stress, as well as response to ABA
(Robles and Quesada, 2021). This poses an interesting role for this candidate gene in the drought
tolerance of common bean.

The TFs bHLH106-related (*Phvul.001G121200*), UPBEAT1 (*Phvul.007G089100*), C2H2-type
Zn-finger protein (*Phvul.007G090500*), and No apical meristem (NAM) protein
(*Phvul.007G089600*) are promising candidate genes for drought tolerance improvement in
common bean. bHLH TFs are one of the main TF families in angiosperms with paramount roles
in abiotic stresses (Khan et al., 2018). In Arabidopsis, Ahmad et al. (2015) showed that
bHLH106 is a major regulator that integrates the function of multiple salt tolerance-related
genes. The homolog gene in common bean could be a main target for crop improvement.
UPBEAT1 is a TF also belonging to the bHLH family. In Arabidopsis, UPBEAT1 regulates
peroxidases and is involved in the accumulation of ROS and primary root elongation
(Tsukagoshi et al., 2010). C2H2 zinc finger proteins have key regulatory roles in abiotic stress
resistance in plants. The CH2H-type proteins have a significant binding diversity characterized
by their ability to interact with DNA, RNA, and other proteins (reviewed in (Han et al., 2020).
Finally, the NAM proteins are part of the NAC TF family, which has been described in multiple
crops (Shao et al., 2015), and associated with stress response in solanaceous (Tweneboah and
Oh, 2017) and cereals (Dudhate et al., 2021).

Conclusions

The present study represents the largest FTSW study reported. The high variation in average
NTR and FTSWc among Mesoamerican beans shows that variation for the improvement of these
traits exists and may enable breeding cultivars with an improved capacity to preserve water
status during short or long water deficit periods. The identified genotypes with the extreme
values for average NTR and FTSWc such as Othello, Harold, and ABCP-8 can be used as
parents for future breeding strategies in either short or long drought periods. The four candidate
QTL identified on Pv01 and Pv07 allowed the identification of 15 new candidate genes that can
be targets for marker-assisted selection and for further research aimed at identifying causal
polymorphisms in gene sequences. Interestingly, five candidate genes corresponded to
transcription factors (e.g., *Phvul.001G121200*) which constitute major target genes due to their

role in controlling multiple genes. Some of the candidate genes have been poorly studied in crop plans, which opens new opportunities for further validation in beans and other crop species.

References

Agurla, S., Gahir, S., Munemasa, S., Murata, Y., and Raghavendra, A. S. (2018). Mechanism of stomatal closure in plants exposed to drought and cold stress. *Adv Exp Med Biol* 1081, 215–232. doi: 10.1007/978-981-13-1244-1_12/TABLES/3.

Ahmad, A., Niwa, Y., Goto, S., Ogawa, T., Shimizu, M., Suzuki, A., et al. (2015). BHLH106 integrates functions of multiple genes through their g-box to confer salt tolerance on Arabidopsis. *PLoS One* 10. doi: 10.1371/journal.pone.0126872.

Allimuthu, E., Dalal, M., Kumar, K. G., Sellathdurai, D., Kumar, R. R., Sathee, L., et al. (2020). Characterization of atypical protein tyrosine kinase (PTK) genes and their role in abiotic stress response in rice. *Plants* 9. doi: 10.3390/plants9050664.

Armengot, L., Marquès-Bueno, M. M., and Jaillais, Y. (2016). Regulation of polar auxin transport by protein and lipid kinases. *J Exp Bot* 67, 4015–4037. doi: 10.1093/jxb/erw216.

Barkan, A. (2011). Studying the structure and processing of chloroplast transcripts. *Methods in Molecular Biology* 774, 183–197. doi: 10.1007/978-1-61779-234-2_12/FIGURES/1_12.

Baron, K. N., Schroeder, D. F., and Stasolla, C. (2012). Transcriptional response of abscisic acid (ABA) metabolism and transport to cold and heat stress applied at the reproductive stage of development in Arabidopsis thaliana. *Plant Science* 188–189, 48–59. doi: 10.1016/j.plantsci.2012.03.001.

Barrett, J. C., Fry, B., Maller, J., and Daly, M. J. (2005). Haploview: Analysis and visualization of LD and haplotype maps. *Bioinformatics* 21, 263–265. doi: 10.1093/bioinformatics/bth457.

Bates, D., Mächler, M., Bolker, B. M., and Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *J Stat Softw* 67. doi: 10.18637/jss.v067.i01.

Beebe, S. E., Rao, I. M., Devi, M. J., Polania, J., Beebe, S. E., Rao, I. M., et al. (2014). Common beans, biodiversity, and multiple stresses: challenges of drought resistance in tropical soils. *Crop Pasture Sci* 65, 667–675. doi: 10.1071/CP13303.

Beihammer, G., Maresch, D., Altmann, F., and Strasser, R. (2020). Glycosylphosphatidylinositol-Anchor Synthesis in Plants: A Glycobiology Perspective. *Front Plant Sci* 11. doi: 10.3389/fpls.2020.611188.

Bharath, P., Gahir, S., and Raghavendra, A. S. (2021). Abscisic Acid-Induced Stomatal Closure: An Important Component of Plant Defense Against Abiotic and Biotic Stress. *Front Plant Sci* 12, 324. doi: 10.3389/FPLS.2021.615114/XML/NLM.

Bowles, A. M. C., Paps, J., and Bechtold, U. (2021). Evolutionary Origins of Drought Tolerance in Spermatophytes. *Front Plant Sci* 0, 1068. doi: 10.3389/FPLS.2021.655924.

Bradbury, P. J., Zhang, Z., Kroon, D. E., Casstevens, T. M., Ramdoss, Y., and Buckler, E. S. (2007). TASSEL: Software for association mapping of complex traits in diverse samples. *Bioinformatics* 23, 2633–2635. doi: 10.1093/bioinformatics/btm308.

Briñez, B., Perseguini, J. M. K. C., Rosa, J. S., Bassi, D., Gonçalves, J. G. R., Almeida, C., et al. (2017). Mapping QTLs for drought tolerance in a SEA 5 x AND 277 common bean cross with SSRs and SNP markers. *Genet Mol Biol* 40, 813–823. doi: 10.1590/1678-4685-gmb-2016-0222.

Cao, X., Zhu, C., Zhong, C., Zhang, J., Wu, L., Jin, Q., et al. (2019). Nitric oxide synthase-mediated early nitric oxide burst alleviates water stress-induced oxidative damage in ammonium-supplied rice roots. *BMC Plant Biol* 19. doi: 10.1186/s12870-019-1721-2.

Carter, T. E., Todd, S. M., and Gillen, A. M. (2016). Registration of 'USDA-N8002' Soybean Cultivar with High Yield and Abiotic Stress Resistance Traits. *J Plant Regist* 10, 238–245. doi: 10.3198/JPR2015.09.0057CRC.

Chaurasia, S., Singh, A. K., Songachan, L. S., Sharma, A. D., Bhardwaj, R., and Singh, K. (2020). Multi-locus genome-wide association studies reveal novel genomic regions associated with vegetative stage salt tolerance in bread wheat (Triticum aestivum L.). *Genomics* 112, 4608–4621. doi: 10.1016/j.ygeno.2020.08.006.

Chen, G., Komatsuda, T., Ma, J. F., Nawrath, C., Pourkheirandish, M., Tagiri, A., et al. (2011). An ATP-binding cassette subfamily G full transporter is essential for the retention of leaf water in both wild barley and rice. *Proc Natl Acad Sci U S A* 108, 12354–12359. doi: 10.1073/pnas.1108444108.

Chiango, H., Jafarikouhini, N., Pradhan, D., Figueiredo, A., Silva, J., Sinclair, T. R., et al. (2022). Drought resilience in CIMMYT maize lines adapted to Africa resulting from transpiration sensitivity to vapor pressure deficit and soil drying. *J Crop Improv* 36, 301–315. doi: 10.1080/15427528.2021.1961334/FORMAT/EPUB.

Choudhary, S., and Sinclair, T. R. (2014). Hydraulic conductance differences among sorghum genotypes to explain variation in restricted transpiration rates. *Functional Plant Biology* 41, 270–275. doi: 10.1071/FP13246.

Chul, S. C., Paramasivan, M., and Chandrasekaran, M. (2018). Proline Accumulation Influenced by Osmotic Stress in Arbuscular Mycorrhizal Symbiotic Plants. doi: 10.3389/fmicb.2018.02525.

Davies, W. J., Kudoyarova, G., and Hartung, W. (2005). Long-distance ABA signaling and its relation to other signaling pathways in the detection of soil drying and the mediation of the plant's response to drought. *J Plant Growth Regul* 24, 285–295. doi: 10.1007/s00344-005-0103-1.

del Río, L. A. (2015). ROS and RNS in plant physiology: an overview. doi: 10.1093/jxb/erv099.

Deng, Y., Srivastava, R., and Howell, S. H. (2013). Endoplasmic Reticulum (ER) Stress Response and Its Physiological Roles in Plants. *International Journal of Molecular Sciences 2013, Vol. 14, Pages 8188-8212* 14, 8188–8212. doi: 10.3390/IJMS14048188.

Devi, M. J., and Reddy, V. (2020a). Cotton genotypic variability for transpiration decrease with progressive soil drying. *Agronomy* 10. doi: 10.3390/agronomy10091290.

Devi, M. J., and Reddy, V. R. (2020b). Stomatal closure response to soil drying at different vapor pressure deficit conditions in maize. *Plant Physiology and Biochemistry* 154, 714–722. doi: 10.1016/j.plaphy.2020.07.023.

Dramadri, I. O., Nkalubo, S. T., and Kelly, J. D. (2019). Identification of QTL Associated with Drought Tolerance in Andean Common Bean. *Crop Sci* 59, 1007–1020. doi: 10.2135/CROPSCI2018.10.0604.

Dudhate, A., Shinde, H., Yu, P., Tsugama, D., Gupta, S. K., Liu, S., et al. (2021). Comprehensive analysis of NAC transcription factor family uncovers drought and salinity stress response in pearl millet (Pennisetum glaucum). *BMC Genomics* 22. doi: 10.1186/s12864-021-07382-y.

Egan, L., Hofmann, R., Nichols, S., Hadipurnomo, J., and Hoyos-Villegas, V. (2021). Transpiration Rate of White Clover (Trifolium repens L.) Cultivars in Drying Soil. *Front Plant Sci* 12. doi: 10.3389/fpls.2021.595030.

Eklund, H., Gleason, F. K., and Holmgren, A. (1991). Structural and functional relations among thioredoxins of different species. *Proteins: Structure, Function, and Bioinformatics* 11, 13–28. doi: 10.1002/PROT.340110103.

FAOSTAT (2019). Dry bean production. http://www.fao.org/faostat/en/#data/QC. *Rome: Food and Agriculture Organization of the United Nations.*

Fereres, E., and Soriano, M. A. (2007). Deficit irrigation for reducing agricultural water use. *J Exp Bot* 58, 147–159. doi: 10.1093/JXB/ERL165.

Fletcher, A. L., Sinclair, T. R., and Allen, L. H. (2007). Transpiration responses to vapor pressure deficit
in well watered 'slow-wilting' and commercial soybean. *Environ Exp Bot* 61, 145–151. doi:
10.1016/J.ENVEXPBOT.2007.05.004.

Gaffney, J., Schussler, J., Löffler, C., Cai, W., Paszkiewicz, S., Messina, C., et al. (2015). Industry-Scale
Evaluation of Maize Hybrids Selected for Increased Yield in Drought-Stress Conditions of the US
Corn Belt. *Crop Sci* 55, 1608–1618. doi: 10.2135/CROPSCI2014.09.0654.

Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., et al. (2012). Phytozome: A
comparative platform for green plant genomics. *Nucleic Acids Res* 40, 1178–1186. doi:
10.1093/nar/gkr944.

Hacke, U. G., and Way, D. (2014). Variable plant hydraulic conductance. *Tree Physiol* 34, 105–108. doi:
10.1093/TREEPHYS/TPU007.

Han, G., Lu, C., Guo, J., Qiao, Z., Sui, N., Qiu, N., et al. (2020). C2H2 Zinc Finger Proteins: Master
Regulators of Abiotic Stress Responses in Plants. *Front Plant Sci* 11. doi: 10.3389/fpls.2020.00115.

Han, S., Yu, B., Wang, Y., and Liu, Y. (2011). Role of plant autophagy in stress response. *Protein Cell* 2,
784–791. doi: 10.1007/s13238-011-1104-4.

Hill, W. G., and Weir, B. S. (1988). Variances and covariances of squared linkage disequilibria in finite
populations. *Theor Popul Biol* 33, 54–78. doi: 10.1016/0040-5809(88)90004-4.

Hoyos-Villegas, V., Song, Q., and Kelly, J. D. (2017). Genome-wide Association Analysis for Drought
Tolerance and Associated Traits in Common Bean. *Plant Genome* 10, plantgenome2015.12.0122.
doi: 10.3835/plantgenome2015.12.0122.

Huang, M., Liu, X., Zhou, Y., Summers, R. M., and Zhang, Z. (2019). BLINK: a package for the next
level of genome-wide association studies with both individuals and markers in the millions.
Gigascience 8, 1–12. doi: 10.1093/GIGASCIENCE/GIY154.

Jafarikouhini, N., Pradhan, D., and Sinclair, T. R. (2020). Basis of limited-transpiration rate under
elevated vapor pressure deficit and high temperature among sweet corn cultivars. *Environ Exp Bot*
179. doi: 10.1016/J.ENVEXPBOT.2020.104205.

Kaler, A. S., Gillman, J. D., Beissinger, T., and Purcell, L. C. (2020). Comparing Different Statistical
Models and Multiple Testing Corrections for Association Mapping in Soybean and Maize. *Front*
Plant Sci 10, 1794. doi: 10.3389/FPLS.2019.01794/BIBTEX.

Khan, S. A., Li, M. Z., Wang, S. M., and Yin, H. J. (2018). Revisiting the role of plant transcription
factors in the battle against abiotic stress. *Int J Mol Sci* 19. doi: 10.3390/ijms19061634.

Kholová, J., Hash, C. T., Kakkera, A., Koová, M., and Vadez, V. (2010a). Constitutive water-conserving
mechanisms are correlated with the terminal drought tolerance of pearl millet [Pennisetum glaucum
(L.) R. Br.]. *J Exp Bot* 61, 369–377. doi: 10.1093/jxb/erp314.

Kholová, J., Hash, C. T., Kakkera, A., Koová, M., and Vadez, V. (2010b). Constitutive water-conserving
mechanisms are correlated with the terminal drought tolerance of pearl millet [Pennisetum glaucum
(L.) R. Br.]. *J Exp Bot* 61, 369–377. doi: 10.1093/jxb/erp314.

Kishor, P. B. K., Hima Kumari, P., Sunita, M. S. L., and Sreenivasulu, N. (2015). Role of proline in cell
wall synthesis and plant development and its implications in plant ontogeny. *Front Plant Sci* 6, 1–
17. doi: 10.3389/fpls.2015.00544.

Korver, R. A., Koevoets, I. T., and Testerink, C. (2018). Out of Shape During Stress: A Key Role for
Auxin. *Trends Plant Sci* 23, 783–793. doi: 10.1016/j.tplants.2018.05.011.

Kuromori, T., Miyaji, T., Yabuuchi, H., Shimizu, H., Sugimoto, E., Kamiya, A., et al. (2010). ABC
transporter AtABCG25 is involved in abscisic acid transport and responses. *Proc Natl Acad Sci U S*
A 107, 2361–2366. doi: 10.1073/pnas.0912516107.

Leitão, S. T., Bicho, M. C., Pereira, P., Paulo, M. J., Malosetti, M., Araújo, S. de S., et al. (2021).
Common bean SNP alleles and candidate genes affecting photosynthesis under contrasting water
regimes. *Horticulture Research 2020 8:1* 8, 1–14. doi: 10.1038/s41438-020-00434-6.

McCrory, M. A., Hamaker, B. R., Lovejoy, J. C., and Eichelsdoerfer, P. E. (2010). Pulse Consumption,
Satiety, and Weight Management. *Advances in Nutrition* 1, 17–30. doi: 10.3945/AN.110.1006.

Miklas, P. N., Kelly, J. D., Beebe, S. E., and Blair, M. W. (2006). Common bean breeding for resistance
against biotic and abiotic stresses: From classical to MAS breeding. *Euphytica 2006 147:1* 147,
105–131. doi: 10.1007/S10681-006-4600-5.

Miyamoto, T., Uemura, T., Nemoto, K., Daito, M., Nozawa, A., Sawasaki, T., et al. (2019). Tyrosine
kinase-dependent defense responses against herbivory in arabidopsis. *Front Plant Sci* 10. doi:
10.3389/fpls.2019.00776.

Moghaddam, S. M., Mamidi, S., Osorno, J. M., Lee, R., Brick, M., Kelly, J., et al. (2016). Genome-Wide
Association Study Identifies Candidate Loci Underlying Agronomic Traits in a Middle American
Diversity Panel of Common Bean. *Plant Genome* 9, 1–21. doi: 10.3835/plantgenome2016.02.0012.

Moghaddam, S. M., Oladzad, A., Koh, C., Ramsay, L., Hart, J. P., Mamidi, S., et al. (2021). The tepary
bean genome provides insight into evolution and domestication under heat stress. *Nature
Communications 2021 12:1* 12, 1–14. doi: 10.1038/s41467-021-22858-x.

Muchow, R. C., and Sinclair, T. R. (1991). Water Deficit Effects on Maize Yields Modeled under Current
and "Greenhouse" Climates. *Agron J* 83, 1052–1059. doi:
10.2134/agronj1991.00021962008300060023x.

Mukeshimana, G., Lasley, A. L., Loescher, W. H., and Kelly, J. D. (2014). Identification of shoot traits
related to drought tolerance in common bean seedlings. *Journal of the American Society for
Horticultural Science* 139, 299–309. doi: 10.21273/jashs.139.3.299.

Muñoz-Perea, C. G., Allen, R. G., Westermann, D. T., Wright, J. L., and Singh, S. P. (2007). Water use
efficiency among dry bean landraces and cultivars in drought-stressed and non-stressed
environments. *Euphytica* 155, 393–402. doi: 10.1007/S10681-006-9340-Z/TABLES/2.

Muñoz-Perea, C. G., Terán, H., Allen, R. G., Wright, J. L., Westermann, D. T., and Singh, S. P. (2006).
Selection for drought resistance in dry bean landraces and cultivars. *Crop Sci* 46, 2111–2120. doi:
10.2135/cropsci2006.01.0029.

Mutlu, N., Miklas, P. N., Steadman, J. R., Vidaver, A. K., Lindgren, D. T., Reiser, J., et al. (2005).
Registration of Common Bacterial Blight Resistant Pinto Bean Germplasm Line ABCP-8. *Crop Sci*
45, 806–807. doi: 10.2135/CROPSCI2005.0806.

Nakashima, K., Yamaguchi-Shinozaki, K., Shinozaki, K., Jain, M., and Discovery, G. (2014). The
transcriptional regulatory network in the drought response and its crosstalk in abiotic stress
responses including drought, cold, and heat. doi: 10.3389/fpls.2014.00170.

Neill, S. J., Desikan, R., and Hancock, J. T. (2003). Nitric oxide signalling in plants. *New Phytologist*
159, 11–35. doi: 10.1046/j.1469-8137.2003.00804.x.

Nemoto, K., Takemori, N., Seki, M., Shinozaki, K., and Sawasaki, T. (2015). Members of the plant CRK
superfamily are capable of trans- and autophosphorylation of tyrosine residues. *Journal of
Biological Chemistry* 290, 16665–16677. doi:
10.1074/JBC.M114.617274/ATTACHMENT/46236EFA-031C-407A-87BE-
61BE30CA3CDB/MMC1.DOCX.

OECD (2014). *Climate Change, Water and Agriculture: Towards Resilient Systems*. doi:
10.2166/9781780406619.

Osakabe, Y., Osakabe, K., Shinozaki, K., and Tran, L.-S. P. (2014). Response of plants to water stress.
Front Plant Sci 5. doi: 10.3389/fpls.2014.00086.

Park, Y., Xu, Z. Y., Kim, S. Y., Lee, J., Choi, B., Lee, J., et al. (2016). Spatial regulation of ABCG25, an ABA exporter, is an important component of the mechanism controlling cellular aba levels. *Plant Cell* 28, 2528–2544. doi: 10.1105/tpc.16.00359.

Perlikowski, D., Lechowicz, K., Pawłowicz, I., Arasimowicz-Jelonek, M., and Kosmala, A. (2022). Scavenging of nitric oxide up-regulates photosynthesis under drought in Festuca arundinacea and F. glaucescens but reduces their drought tolerance. *Sci Rep* 12. doi: 10.1038/s41598-022-10299-5.

Raj, A., Stephens, M., and Pritchard, J. K. (2014). FastSTRUCTURE: Variational inference of population structure in large SNP data sets. *Genetics* 197, 573–589. doi: 10.1534/genetics.114.164350.

Ray, J. D., and Sinclair, T. R. (1997). Stomatal closure of maize hybrids in response to drying soil. *Crop Sci* 37, 803–807. doi: 10.2135/cropsci1997.0011183X003700030018x.

Remington, D. L., Thornsberry, J. M., Matsuoka, Y., Wilson, L. M., Whitt, S. R., Doebley, J., et al. (2001). Structure of linkage disequilibrium and phenotypic associations in the maize genome. *Proc Natl Acad Sci U S A* 98, 11479–11484. doi: 10.1073/pnas.201394398.

Rey, P., Pruvot, G., Lle Becuwe, N., Oise Eymery, F., Rumeau, D., and Peltier, G. (1998). A novel thioredoxin-like protein located in the chloroplast is induced by water deficit in Solanum tuberosum L. plants.

Ritchie, J. T. (1980). "Climate and Soil Water," in *Moving Up the Yield Curve: Advances and Obstacles*, eds. L. S. Murphy and et al. (ASA Spec. Publ.), 1–24. doi: 10.2134/ASASPECPUB39.C1.

Robles, P., and Quesada, V. (2021). Research progress in the molecular functions of plant mTERF proteins. *Cells* 10, 1–16. doi: 10.3390/cells10020205.

Rosyara, U. R., De Jong, W. S., Douches, D. S., and Endelman, J. B. (2016). Software for Genome-Wide Association Studies in Autopolyploids and Its Application to Potato. *Plant Genome* 9, 1–10. doi: 10.3835/plantgenome2015.08.0073.

Sadok, W., Gilbert, M. E., Raza, M. A. S., and Sinclair, T. R. (2012). Basis of Slow-Wilting Phenotype in Soybean PI 471938. *Crop Sci* 52, 1261–1269. doi: 10.2135/CROPSCI2011.11.0622.

Sanchez, M., Sinclair, T. R., and Pradhan, D. (2021). Transpiration response to vapor pressure deficit and soil drying among quinoa genotypes (Chenopodium quinoa Willd.). *J Crop Improv* 35, 291–302. doi: 10.1080/15427528.2020.1817221/FORMAT/EPUB.

Sato, R., and Maeshima, M. (2019). The ER-localized aquaporin SIP2;1 is involved in pollen germination and pollen tube elongation in Arabidopsis thaliana. *Plant Mol Biol* 100, 335–349. doi: 10.1007/s11103-019-00865-3.

Schmutz, J., McClean, P. E., Mamidi, S., Wu, G. A., Cannon, S. B., Grimwood, J., et al. (2014). A reference genome for common bean and genome-wide analysis of dual domestications. *Nat Genet* 46, 707–713. doi: 10.1038/ng.3008.

Sedlar, A., Zupin, M., Maras, M., Razinger, J., Šuštar-Vozlič, J., Pipan, B., et al. (2020). QTL Mapping for Drought-Responsive Agronomic Traits Associated with Physiology, Phenology, and Yield in an Andean Intra-Gene Pool Common Bean Population. *Agronomy 2020, Vol. 10, Page 225* 10, 225. doi: 10.3390/AGRONOMY10020225.

Shao, H., Wang, H., and Tang, X. (2015). NAC transcription factors in plant multiple abiotic stress responses: Progress and prospects. *Front Plant Sci* 6. doi: 10.3389/fpls.2015.00902.

Simonin, K. A., Burns, E., Choat, B., Barbour, M. M., Dawson, T. E., and Franks, P. J. (2015). Increasing leaf hydraulic conductance with transpiration rate minimizes the water potential drawdown from stem to leaf. *J Exp Bot* 66, 1303–1315. doi: 10.1093/JXB/ERU481.

Sinclair, T. R., Zwieniecki, M. A., and Holbrook, N. M. (2008). Low leaf hydraulic conductance associated with drought tolerance in soybean. *Physiol Plant* 132, 446–451. doi: 10.1111/J.1399-3054.2007.01028.X.

Singh, S., Hayes, R., Robinson, C., Dennis, M., and Powers, E. (2001). Response of Bean (Phaseolus
vulgaris L.) Cultivars to Drought Stress. *Annu. Rep. Bean Improv. Coop.* 44, 45–46.

Sôber, A. (1998). Hydraulic conductance, stomatal conductance, and maximal photosynthetic rate in bean
leaves. *Photosynthetica 1997 34:4* 34, 599–603. doi: 10.1023/A:1006834119588.

Souter, J. R., Gurusamy, V., Porch, T. G., and Bett, K. E. (2017). Successful introgression of abiotic
stress tolerance from wild tepary bean to common bean. *Crop Sci* 57, 1160–1171. doi:
10.2135/cropsci2016.10.0851.

Swarup, R., and Bhosale, R. (2019). Developmental Roles of AUX1/LAX Auxin Influx Carriers in
Plants. *Front Plant Sci* 10. doi: 10.3389/fpls.2019.01306.

Tang, Y., Liu, X., Wang, J., Li, M., Wang, Q., Tian, F., et al. (2016). GAPIT Version 2: An Enhanced
Integrated Tool for Genomic Association and Prediction. *Plant Genome* 9. doi:
10.3835/plantgenome2015.11.0120.

Thomas, C. v, and Giles Waines, J. (1984). Fertile backcross and allotetraploid plants from crosses
between tepary beans and common beans. Available at: https://academic.oup.com/jhered/article-
abstract/75/2/93/765293.

Tsukagoshi, H., Busch, W., and Benfey, P. N. (2010). Transcriptional regulation of ROS controls
transition from proliferation to differentiation in the root. *Cell* 143, 606–616. doi:
10.1016/J.CELL.2010.10.020/ATTACHMENT/F4808132-6242-40E4-A4F7-
A40762FFE4AE/MMC3.XLS.

Tweneboah, S., and Oh, S. K. (2017). Biological roles of NAC transcription factors in the regulation of
biotic and abiotic stress responses in solanaceous crops. *J Plant Biotechnol* 44, 1–11. doi:
10.5010/JPB.2017.44.1.001.

VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. *J Dairy Sci* 91, 4414–4423.
doi: 10.3168/jds.2007-0980.

Wallace, J. G., Zhang, X., Beyene, Y., Semagn, K., Olsen, M., Prasanna, B. M., et al. (2016). Genome-
wide association for plant height and flowering time across 15 tropical maize populations under
managed drought stress and well-watered conditions in Sub-Saharan Africa. *Crop Sci* 56, 2365–
2378. doi: 10.2135/cropsci2015.10.0632.

Wu, J. X., Li, J., Liu, Z., Yin, J., Chang, Z. Y., Rong, C., et al. (2015). The Arabidopsis ceramidase
AtACER functions in disease resistance and salt tolerance. *Plant Journal* 81, 767–780. doi:
10.1111/tpj.12769.

Yokochi, Y., Fukushi, Y., Wakabayashi, K.-I., Yoshida, K., and Hisabori, T. (2021). Oxidative regulation
of chloroplast enzymes by thioredoxin and thioredoxin-like proteins in Arabidopsis thaliana. *Proc
Natl Acad Sci U S A* 118. doi: 10.1073/pnas.2114952118/-/DCSupplemental.

Zhang, Q., Li, J., Zhang, W., Yan, S., Wang, R., Zhao, J., et al. (2012). The putative auxin efflux carrier
OsPIN3t is involved in the drought stress response and drought tolerance. *Plant Journal* 72, 805–
816. doi: 10.1111/j.1365-313X.2012.05121.x.

Zheng, P., Wu, J. X., Sahu, S. K., Zeng, H. Y., Huang, L. Q., Liu, Z., et al. (2018). Loss of alkaline
ceramidase inhibits autophagy in Arabidopsis and plays an important role during environmental
stress response. *Plant Cell Environ* 41, 837–849. doi: 10.1111/pce.13148.

683
684 **Figure 1.** Average normalized transpiration rate (NTR) in response to the fraction of transpirable
685 soil water (FTSW) in the five genotypes with the lowest FTSWc (A) and the five genotypes with
686 the highest FTSWc (B).
687

688
689 **Figure 2.** Manhattan plots of QTL identified for NTR (A) using FarmCPU model, and for
690 FTSWc (B) using Blink model in the subpanel of 82 Mesoamerican beans. SNPs above the FDR-
691 corrected threshold were considered as significant. The haplotype block for each significant SNP
692 is depicted in the bottom of each Manhattan plot. The Q-Q plot on the right depicts the
693 adjustment of the model for each trait.

Parsed Citations

Agurla, S., Gahir, S., Munemasa, S., Murata, Y., and Raghavendra, A. S. (2018). Mechanism of stomatal closure in plants exposed to drought and cold stress. Adv Exp Med Biol 1081, 215–232. doi: 10.1007/978-981-13-1244-1_12/TABLES/3.
  Google Scholar: Author Only Title Only Author and Title

Ahmad, A, Niwa, Y., Goto, S., Ogawa, T., Shimizu, M., Suzuki, A, et al. (2015). BHLH106 integrates functions of multiple genes through their g-box to confer salt tolerance on Arabidopsis. PLoS One 10. doi: 10.1371/journal.pone.0126872.
  Google Scholar: Author Only Title Only Author and Title

Allimuthu, E., Dalal, M., Kumar, K. G., Sellathdurai, D., Kumar, R. R., Sathee, L., et al. (2020). Characterization of atypical protein tyrosine kinase (PTK) genes and their role in abiotic stress response in rice. Plants 9. doi: 10.3390/plants9050664.
  Google Scholar: Author Only Title Only Author and Title

Armengot, L., Marquès-Bueno, M. M., and Jaillais, Y. (2016). Regulation of polar auxin transport by protein and lipid kinases. J Exp Bot 67, 4015–4037. doi: 10.1093/jxb/erw216.
  Google Scholar: Author Only Title Only Author and Title

Barkan, A. (2011). Studying the structure and processing of chloroplast transcripts. Methods in Molecular Biology 774, 183–197. doi: 10.1007/978-1-61779-234-2_12/FIGURES/1_12.
  Google Scholar: Author Only Title Only Author and Title

Baron, K. N., Schroeder, D. F., and Stasolla, C. (2012). Transcriptional response of abscisic acid (ABA) metabolism and transport to cold and heat stress applied at the reproductive stage of development in Arabidopsis thaliana. Plant Science 188–189, 48–59. doi: 10.1016/j.plantsci.2012.03.001.
  Google Scholar: Author Only Title Only Author and Title

Barrett, J. C., Fry, B., Maller, J., and Daly, M. J. (2005). Haploview: Analysis and visualization of LD and haplotype maps. Bioinformatics 21, 263–265. doi: 10.1093/bioinformatics/bth457.
  Google Scholar: Author Only Title Only Author and Title

Bates, D., Mächler, M., Bolker, B. M., and Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. J Stat Softw 67. doi: 10.18637/jss.v067.i01.
  Google Scholar: Author Only Title Only Author and Title

Beebe, S. E., Rao, I. M., Devi, M. J., Polania, J., Beebe, S. E., Rao, I. M., et al. (2014). Common beans, biodiversity, and multiple stresses: challenges of drought resistance in tropical soils. Crop Pasture Sci 65, 667–675. doi: 10.1071/CP13303.
  Google Scholar: Author Only Title Only Author and Title

Beihammer, G., Maresch, D., Altmann, F., and Strasser, R. (2020). Glycosylphosphatidylinositol-Anchor Synthesis in Plants: A Glycobiology Perspective. Front Plant Sci 11. doi: 10.3389/fpls.2020.611188.
  Google Scholar: Author Only Title Only Author and Title

Bharath, P., Gahir, S., and Raghavendra, A. S. (2021). Abscisic Acid-Induced Stomatal Closure: An Important Component of Plant Defense Against Abiotic and Biotic Stress. Front Plant Sci 12, 324. doi: 10.3389/FPLS.2021.615114/XML/NLM.
  Google Scholar: Author Only Title Only Author and Title

Bowles, A. M. C., Paps, J., and Bechtold, U. (2021). Evolutionary Origins of Drought Tolerance in Spermatophytes. Front Plant Sci 0, 1068. doi: 10.3389/FPLS.2021.655924.
  Google Scholar: Author Only Title Only Author and Title

Bradbury, P. J., Zhang, Z, Kroon, D. E., Casstevens, T. M., Ramdoss, Y., and Buckler, E. S. (2007). TASSEL: Software for association mapping of complex traits in diverse samples. Bioinformatics 23, 2633–2635. doi: 10.1093/bioinformatics/btm308.
  Google Scholar: Author Only Title Only Author and Title

Briñez, B., Perseguini, J. M. K. C., Rosa, J. S., Bassi, D., Gonçalves, J. G. R., Almeida, C., et al. (2017). Mapping QTLs for drought tolerance in a SEA 5 x AND 277 common bean cross with SSRs and SNP markers. Genet Mol Biol 40, 813–823. doi: 10.1590/1678-4685-gmb-2016-0222.
  Google Scholar: Author Only Title Only Author and Title

Cao, X., Zhu, C., Zhong, C., Zhang, J., Wu, L., Jin, Q., et al. (2019). Nitric oxide synthase-mediated early nitric oxide burst alleviates water stress-induced oxidative damage in ammonium-supplied rice roots. BMC Plant Biol 19. doi: 10.1186/s12870-019-1721-2.
  Google Scholar: Author Only Title Only Author and Title

Carter, T. E., Todd, S. M., and Gillen, A. M. (2016). Registration of 'USDA-N8002' Soybean Cultivar with High Yield and Abiotic Stress Resistance Traits. J Plant Regist 10, 238–245. doi: 10.3198/JPR2015.09.0057CRC.
  Google Scholar: Author Only Title Only Author and Title

Chaurasia, S., Singh, A. K., Songachan, L. S., Sharma, A. D., Bhardwaj, R., and Singh, K. (2020). Multi-locus genome-wide association studies reveal novel genomic regions associated with vegetative stage salt tolerance in bread wheat (Triticum

aestivum L.). Genomics 112, 4608–4621. doi: 10.1016/j.ygeno.2020.08.006.

Google Scholar: Author Only Title Only Author and Title

Chen, G., Komatsuda, T., Ma, J. F., Nawrath, C., Pourkheirandish, M., Tagiri, A, et al. (2011). An ATP-binding cassette subfamily G full transporter is essential for the retention of leaf water in both wild barley and rice. Proc Natl Acad Sci U S A 108, 12354–12359. doi: 10.1073/pnas.1108444108.

Google Scholar: Author Only Title Only Author and Title

Chiango, H., Jafarikouhini, N., Pradhan, D., Figueiredo, A, Silva, J., Sinclair, T. R., et al. (2022). Drought resilience in CIMMYT maize lines adapted to Africa resulting from transpiration sensitivity to vapor pressure deficit and soil drying. J Crop Improv 36, 301–315. doi: 10.1080/15427528.2021.1961334/FORMAT/EPUB.

Google Scholar: Author Only Title Only Author and Title

Choudhary, S., and Sinclair, T. R. (2014). Hydraulic conductance differences among sorghum genotypes to explain variation in restricted transpiration rates. Functional Plant Biology 41, 270–275. doi: 10.1071/FP13246.

Google Scholar: Author Only Title Only Author and Title

Chul, S. C., Paramasivan, M., and Chandrasekaran, M. (2018). Proline Accumulation Influenced by Osmotic Stress in Arbuscular Mycorrhizal Symbiotic Plants. doi: 10.3389/fmicb.2018.02525.

Google Scholar: Author Only Title Only Author and Title

Davies, W. J., Kudoyarova, G., and Hartung, W. (2005). Long-distance ABA signaling and its relation to other signaling pathways in the detection of soil drying and the mediation of the plant's response to drought. J Plant Growth Regul 24, 285–295. doi: 10.1007/s00344-005-0103-1.

Google Scholar: Author Only Title Only Author and Title

del Río, L. A. (2015). ROS and RNS in plant physiology: an overview. doi: 10.1093/jxb/erv099.

Google Scholar: Author Only Title Only Author and Title

Deng, Y., Srivastava, R., and Howell, S. H. (2013). Endoplasmic Reticulum (ER) Stress Response and Its Physiological Roles in Plants. International Journal of Molecular Sciences 2013, Vol. 14, Pages 8188-8212 14, 8188–8212. doi: 10.3390/IJMS14048188.

Google Scholar: Author Only Title Only Author and Title

Devi, M. J., and Reddy, V. (2020a). Cotton genotypic variability for transpiration decrease with progressive soil drying. Agronomy 10. doi: 10.3390/agronomy10091290.

Google Scholar: Author Only Title Only Author and Title

Devi, M. J., and Reddy, V. R. (2020b). Stomatal closure response to soil drying at different vapor pressure deficit conditions in maize. Plant Physiology and Biochemistry 154, 714–722. doi: 10.1016/j.plaphy.2020.07.023.

Google Scholar: Author Only Title Only Author and Title

Dramadri, I. O., Nkalubo, S. T., and Kelly, J. D. (2019). Identification of QTL Associated with Drought Tolerance in Andean Common Bean. Crop Sci 59, 1007–1020. doi: 10.2135/CROPSCI2018.10.0604.

Google Scholar: Author Only Title Only Author and Title

Dudhate, A, Shinde, H., Yu, P., Tsugama, D., Gupta, S. K., Liu, S., et al. (2021). Comprehensive analysis of NAC transcription factor family uncovers drought and salinity stress response in pearl millet (Pennisetum glaucum). BMC Genomics 22. doi: 10.1186/s12864-021-07382-y.

Google Scholar: Author Only Title Only Author and Title

Egan, L., Hofmann, R., Nichols, S., Hadipurnomo, J., and Hoyos-Villegas, V. (2021). Transpiration Rate of White Clover (Trifolium repens L.) Cultivars in Drying Soil. Front Plant Sci 12. doi: 10.3389/fpls.2021.595030.

Google Scholar: Author Only Title Only Author and Title

Eklund, H., Gleason, F. K., and Holmgren, A (1991). Structural and functional relations among thioredoxins of different species. Proteins: Structure, Function, and Bioinformatics 11, 13–28. doi: 10.1002/PROT.340110103.

Google Scholar: Author Only Title Only Author and Title

FAOSTAT (2019). Dry bean production. http://www.fao.org/faostat/en/#data/QC. Rome: Food and Agriculture Organization of the United Nations.

Google Scholar: Author Only Title Only Author and Title

Fereres, E., and Soriano, M. A (2007). Deficit irrigation for reducing agricultural water use. J Exp Bot 58, 147–159. doi: 10.1093/JXB/ERL165.

Google Scholar: Author Only Title Only Author and Title

Fletcher, A L., Sinclair, T. R., and Allen, L. H. (2007). Transpiration responses to vapor pressure deficit in well watered 'slow-wilting' and commercial soybean. Environ Exp Bot 61, 145–151. doi: 10.1016/J.ENVEXPBOT.2007.05.004.

Google Scholar: Author Only Title Only Author and Title

Gaffney, J., Schussler, J., Löffler, C., Cai, W., Paszkiewicz, S., Messina, C., et al. (2015). Industry-Scale Evaluation of Maize Hybrids Selected for Increased Yield in Drought-Stress Conditions of the US Corn Belt. Crop Sci 55, 1608–1618. doi: 10.2135/CROPSCI2014.09.0654.
Google Scholar: Author Only Title Only Author and Title

Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., et al. (2012). Phytozome: A comparative platform for green plant genomics. Nucleic Acids Res 40, 1178–1186. doi: 10.1093/nar/gkr944.
Google Scholar: Author Only Title Only Author and Title

Hacke, U. G., and Way, D. (2014). Variable plant hydraulic conductance. Tree Physiol 34, 105–108. doi: 10.1093/TREEPHYS/TPU007.
Google Scholar: Author Only Title Only Author and Title

Han, G., Lu, C., Guo, J., Qiao, Z., Sui, N., Qiu, N., et al. (2020). C2H2 Zinc Finger Proteins: Master Regulators of Abiotic Stress Responses in Plants. Front Plant Sci 11. doi: 10.3389/fpls.2020.00115.
Google Scholar: Author Only Title Only Author and Title

Han, S., Yu, B., Wang, Y., and Liu, Y. (2011). Role of plant autophagy in stress response. Protein Cell 2, 784–791. doi: 10.1007/s13238-011-1104-4.
Google Scholar: Author Only Title Only Author and Title

Hill, W. G., and Weir, B. S. (1988). Variances and covariances of squared linkage disequilibria in finite populations. Theor Popul Biol 33, 54–78. doi: 10.1016/0040-5809(88)90004-4.
Google Scholar: Author Only Title Only Author and Title

Hoyos-Villegas, V., Song, Q., and Kelly, J. D. (2017). Genome-wide Association Analysis for Drought Tolerance and Associated Traits in Common Bean. Plant Genome 10, plantgenome2015.12.0122. doi: 10.3835/plantgenome2015.12.0122.
Google Scholar: Author Only Title Only Author and Title

Huang, M., Liu, X., Zhou, Y., Summers, R. M., and Zhang, Z. (2019). BLINK: a package for the next level of genome-wide association studies with both individuals and markers in the millions. Gigascience 8, 1–12. doi: 10.1093/GIGASCIENCE/GIY154.
Google Scholar: Author Only Title Only Author and Title

Jafarikouhini, N., Pradhan, D., and Sinclair, T. R. (2020). Basis of limited-transpiration rate under elevated vapor pressure deficit and high temperature among sweet corn cultivars. Environ Exp Bot 179. doi: 10.1016/J.ENVEXPBOT.2020.104205.
Google Scholar: Author Only Title Only Author and Title

Kaler, A. S., Gillman, J. D., Beissinger, T., and Purcell, L. C. (2020). Comparing Different Statistical Models and Multiple Testing Corrections for Association Mapping in Soybean and Maize. Front Plant Sci 10, 1794. doi: 10.3389/FPLS.2019.01794/BIBTEX.
Google Scholar: Author Only Title Only Author and Title

Khan, S. A., Li, M. Z., Wang, S. M., and Yin, H. J. (2018). Revisiting the role of plant transcription factors in the battle against abiotic stress. Int J Mol Sci 19. doi: 10.3390/ijms19061634.
Google Scholar: Author Only Title Only Author and Title

Kholová, J., Hash, C. T., Kakkera, A., Koová, M., and Vadez, V. (2010a). Constitutive water-conserving mechanisms are correlated with the terminal drought tolerance of pearl millet [Pennisetum glaucum (L.) R. Br.]. J Exp Bot 61, 369–377. doi: 10.1093/jxb/erp314.
Google Scholar: Author Only Title Only Author and Title

Kholová, J., Hash, C. T., Kakkera, A., Koová, M., and Vadez, V. (2010b). Constitutive water-conserving mechanisms are correlated with the terminal drought tolerance of pearl millet [Pennisetum glaucum (L.) R. Br.]. J Exp Bot 61, 369–377. doi: 10.1093/jxb/erp314.
Google Scholar: Author Only Title Only Author and Title

Kishor, P. B. K., Hima Kumari, P., Sunita, M. S. L., and Sreenivasulu, N. (2015). Role of proline in cell wall synthesis and plant development and its implications in plant ontogeny. Front Plant Sci 6, 1–17. doi: 10.3389/fpls.2015.00544.
Google Scholar: Author Only Title Only Author and Title

Korver, R. A., Koevoets, I. T., and Testerink, C. (2018). Out of Shape During Stress: A Key Role for Auxin. Trends Plant Sci 23, 783–793. doi: 10.1016/j.tplants.2018.05.011.
Google Scholar: Author Only Title Only Author and Title

Kuromori, T., Miyaji, T., Yabuuchi, H., Shimizu, H., Sugimoto, E., Kamiya, A., et al. (2010). ABC transporter AtABCG25 is involved in abscisic acid transport and responses. Proc Natl Acad Sci U S A 107, 2361–2366. doi: 10.1073/pnas.0912516107.
Google Scholar: Author Only Title Only Author and Title

Leitão, S. T., Bicho, M. C., Pereira, P., Paulo, M. J., Malosetti, M., Araújo, S. de S., et al. (2021). Common bean SNP alleles and candidate genes affecting photosynthesis under contrasting water regimes. Horticulture Research 2020 8:1 8, 1–14. doi: 10.1038/s41438-020-00434-6.
Google Scholar: Author Only Title Only Author and Title

McCrory, M. A, Hamaker, B. R., Lovejoy, J. C., and Eichelsdoerfer, P. E. (2010). Pulse Consumption, Satiety, and Weight Management. Advances in Nutrition 1, 17–30. doi: 10.3945/AN.110.1006.
Google Scholar: Author Only Title Only Author and Title

Miklas, P. N., Kelly, J. D., Beebe, S. E., and Blair, M. W. (2006). Common bean breeding for resistance against biotic and abiotic stresses: From classical to MAS breeding. Euphytica 2006 147:1 147, 105–131. doi: 10.1007/S10681-006-4600-5.
Google Scholar: Author Only Title Only Author and Title

Miyamoto, T., Uemura, T., Nemoto, K., Daito, M., Nozawa, A, Sawasaki, T., et al. (2019). Tyrosine kinase-dependent defense responses against herbivory in arabidopsis. Front Plant Sci 10. doi: 10.3389/fpls.2019.00776.
Google Scholar: Author Only Title Only Author and Title

Moghaddam, S. M., Mamidi, S., Osorno, J. M., Lee, R., Brick, M., Kelly, J., et al. (2016). Genome-Wide Association Study Identifies Candidate Loci Underlying Agronomic Traits in a Middle American Diversity Panel of Common Bean. Plant Genome 9, 1–21. doi: 10.3835/plantgenome2016.02.0012.
Google Scholar: Author Only Title Only Author and Title

Moghaddam, S. M., Oladzad, A, Koh, C., Ramsay, L., Hart, J. P., Mamidi, S., et al. (2021). The tepary bean genome provides insight into evolution and domestication under heat stress. Nature Communications 2021 12:1 12, 1–14. doi: 10.1038/s41467-021-22858-x.
Google Scholar: Author Only Title Only Author and Title

Muchow, R. C., and Sinclair, T. R. (1991). Water Deficit Effects on Maize Yields Modeled under Current and "Greenhouse" Climates. Agron J 83, 1052–1059. doi: 10.2134/agronj1991.00021962008300060023x.
Google Scholar: Author Only Title Only Author and Title

Mukeshimana, G., Lasley, A. L., Loescher, W. H., and Kelly, J. D. (2014). Identification of shoot traits related to drought tolerance in common bean seedlings. Journal of the American Society for Horticultural Science 139, 299–309. doi: 10.21273/jashs.139.3.299.
Google Scholar: Author Only Title Only Author and Title

Muñoz-Perea, C. G., Allen, R. G., Westermann, D. T., Wright, J. L., and Singh, S. P. (2007). Water use efficiency among dry bean landraces and cultivars in drought-stressed and non-stressed environments. Euphytica 155, 393–402. doi: 10.1007/S10681-006-9340-Z/TABLES/2.
Google Scholar: Author Only Title Only Author and Title

Muñoz-Perea, C. G., Terán, H., Allen, R. G., Wright, J. L., Westermann, D. T., and Singh, S. P. (2006). Selection for drought resistance in dry bean landraces and cultivars. Crop Sci 46, 2111–2120. doi: 10.2135/cropsci2006.01.0029.
Google Scholar: Author Only Title Only Author and Title

Mutlu, N., Miklas, P. N., Steadman, J. R., Vidaver, A. K., Lindgren, D. T., Reiser, J., et al. (2005). Registration of Common Bacterial Blight Resistant Pinto Bean Germplasm Line ABCP-8. Crop Sci 45, 806–807. doi: 10.2135/CROPSCI2005.0806.
Google Scholar: Author Only Title Only Author and Title

Nakashima, K., Yamaguchi-Shinozaki, K., Shinozaki, K., Jain, M., and Discovery, G. (2014). The transcriptional regulatory network in the drought response and its crosstalk in abiotic stress responses including drought, cold, and heat. doi: 10.3389/fpls.2014.00170.
Google Scholar: Author Only Title Only Author and Title

Neill, S. J., Desikan, R., and Hancock, J. T. (2003). Nitric oxide signalling in plants. New Phytologist 159, 11–35. doi: 10.1046/j.1469-8137.2003.00804.x.
Google Scholar: Author Only Title Only Author and Title

Nemoto, K., Takemori, N., Seki, M., Shinozaki, K., and Sawasaki, T. (2015). Members of the plant CRK superfamily are capable of trans- and autophosphorylation of tyrosine residues. Journal of Biological Chemistry 290, 16665–16677. doi: 10.1074/JBC.M114.617274/ATTACHMENT/46236EFA-031C-407A-87BE-61BE30CA3CDB/MMC1.DOCX.
Google Scholar: Author Only Title Only Author and Title

OECD (2014). Climate Change, Water and Agriculture: Towards Resilient Systems. doi: 10.2166/9781780406619.
Google Scholar: Author Only Title Only Author and Title

Osakabe, Y., Osakabe, K., Shinozaki, K., and Tran, L.-S. P. (2014). Response of plants to water stress. Front Plant Sci 5. doi: 10.3389/fpls.2014.00086.
Google Scholar: Author Only Title Only Author and Title

Park, Y., Xu, Z. Y., Kim, S. Y., Lee, J., Choi, B., Lee, J., et al. (2016). Spatial regulation of ABCG25, an ABA exporter, is an important component of the mechanism controlling cellular aba levels. Plant Cell 28, 2528–2544. doi: 10.1105/tpc.16.00359.
Google Scholar: Author Only Title Only Author and Title

Perlikowski, D., Lechowicz, K., Pawłowicz, I., Arasimowicz-Jelonek, M., and Kosmala, A. (2022). Scavenging of nitric oxide up-regulates photosynthesis under drought in Festuca arundinacea and F. glaucescens but reduces their drought tolerance. Sci Rep 12. doi: 10.1038/s41598-022-10299-5.

Google Scholar: Author Only Title Only Author and Title

Raj, A, Stephens, M., and Pritchard, J. K. (2014). FastSTRUCTURE: Variational inference of population structure in large SNP data sets. Genetics 197, 573–589. doi: 10.1534/genetics.114.164350.
  Google Scholar: Author Only Title Only Author and Title

Ray, J. D., and Sinclair, T. R. (1997). Stomatal closure of maize hybrids in response to drying soil. Crop Sci 37, 803–807. doi: 10.2135/cropsci1997.0011183X003700030018x.
  Google Scholar: Author Only Title Only Author and Title

Remington, D. L., Thornsberry, J. M., Matsuoka, Y., Wilson, L. M., Whitt, S. R., Doebley, J., et al. (2001). Structure of linkage disequilibrium and phenotypic associations in the maize genome. Proc Natl Acad Sci U S A 98, 11479–11484. doi: 10.1073/pnas.201394398.
  Google Scholar: Author Only Title Only Author and Title

Rey, P., Pruvot, G., Lle Becuwe, N., Oise Eymery, F., Rumeau, D., and Peltier, G. (1998). A novel thioredoxin-like protein located in the chloroplast is induced by water deficit in Solanum tuberosum L. plants.
  Google Scholar: Author Only Title Only Author and Title

Ritchie, J. T. (1980). "Climate and Soil Water," in Moving Up the Yield Curve: Advances and Obstacles, eds. L. S. Murphy and et al. (ASA Spec. Publ.), 1–24. doi: 10.2134/ASASPECPUB39.C1.
  Google Scholar: Author Only Title Only Author and Title

Robles, P., and Quesada, V. (2021). Research progress in the molecular functions of plant mTERF proteins. Cells 10, 1–16. doi: 10.3390/cells10020205.
  Google Scholar: Author Only Title Only Author and Title

Rosyara, U. R., De Jong, W. S., Douches, D. S., and Endelman, J. B. (2016). Software for Genome-Wide Association Studies in Autopolyploids and Its Application to Potato. Plant Genome 9, 1–10. doi: 10.3835/plantgenome2015.08.0073.
  Google Scholar: Author Only Title Only Author and Title

Sadok, W., Gilbert, M. E., Raza, M. A S., and Sinclair, T. R. (2012). Basis of Slow-Wilting Phenotype in Soybean PI 471938. Crop Sci 52, 1261–1269. doi: 10.2135/CROPSCI2011.11.0622.
  Google Scholar: Author Only Title Only Author and Title

Sanchez, M., Sinclair, T. R., and Pradhan, D. (2021). Transpiration response to vapor pressure deficit and soil drying among quinoa genotypes (Chenopodium quinoa Willd.). J Crop Improv 35, 291–302. doi: 10.1080/15427528.2020.1817221/FORMAT/EPUB.
  Google Scholar: Author Only Title Only Author and Title

Sato, R., and Maeshima, M. (2019). The ER-localized aquaporin SIP2;1 is involved in pollen germination and pollen tube elongation in Arabidopsis thaliana. Plant Mol Biol 100, 335–349. doi: 10.1007/s11103-019-00865-3.
  Google Scholar: Author Only Title Only Author and Title

Schmutz, J., McClean, P. E., Mamidi, S., Wu, G. A, Cannon, S. B., Grimwood, J., et al. (2014). A reference genome for common bean and genome-wide analysis of dual domestications. Nat Genet 46, 707–713. doi: 10.1038/ng.3008.
  Google Scholar: Author Only Title Only Author and Title

Sedlar, A, Zupin, M., Maras, M., Razinger, J., Šuštar-Vozlič, J., Pipan, B., et al. (2020). QTL Mapping for Drought-Responsive Agronomic Traits Associated with Physiology, Phenology, and Yield in an Andean Intra-Gene Pool Common Bean Population. Agronomy 2020, Vol. 10, Page 225 10, 225. doi: 10.3390/AGRONOMY10020225.
  Google Scholar: Author Only Title Only Author and Title

Shao, H., Wang, H., and Tang, X. (2015). NAC transcription factors in plant multiple abiotic stress responses: Progress and prospects. Front Plant Sci 6. doi: 10.3389/fpls.2015.00902.
  Google Scholar: Author Only Title Only Author and Title

Simonin, K. A, Burns, E., Choat, B., Barbour, M. M., Dawson, T. E., and Franks, P. J. (2015). Increasing leaf hydraulic conductance with transpiration rate minimizes the water potential drawdown from stem to leaf. J Exp Bot 66, 1303–1315. doi: 10.1093/JXB/ERU481.
  Google Scholar: Author Only Title Only Author and Title

Sinclair, T. R., Zwieniecki, M. A, and Holbrook, N. M. (2008). Low leaf hydraulic conductance associated with drought tolerance in soybean. Physiol Plant 132, 446–451. doi: 10.1111/J.1399-3054.2007.01028.X.
  Google Scholar: Author Only Title Only Author and Title

Singh, S., Hayes, R., Robinson, C., Dennis, M., and Powers, E. (2001). Response of Bean (Phaseolus vulgaris L.) Cultivars to Drought Stress. Annu. Rep. Bean Improv. Coop. 44, 45–46.
  Google Scholar: Author Only Title Only Author and Title

Sôber, A. (1998). Hydraulic conductance, stomatal conductance, and maximal photosynthetic rate in bean leaves. Photosynthetica 1997 34:4 34, 599–603. doi: 10.1023/A:1006834119588.

Google Scholar: Author Only Title Only Author and Title

Souter, J. R., Gurusamy, V., Porch, T. G., and Bett, K. E. (2017). Successful introgression of abiotic stress tolerance from wild tepary bean to common bean. Crop Sci 57, 1160–1171. doi: 10.2135/cropsci2016.10.0851.
Google Scholar: Author Only Title Only Author and Title

Swarup, R., and Bhosale, R. (2019). Developmental Roles of AUX1/LAX Auxin Influx Carriers in Plants. Front Plant Sci 10. doi: 10.3389/fpls.2019.01306.
Google Scholar: Author Only Title Only Author and Title

Tang, Y., Liu, X., Wang, J., Li, M., Wang, Q., Tian, F., et al. (2016). GAPIT Version 2: An Enhanced Integrated Tool for Genomic Association and Prediction. Plant Genome 9. doi: 10.3835/plantgenome2015.11.0120.
Google Scholar: Author Only Title Only Author and Title

Thomas, C. v, and Giles Waines, J. (1984). Fertile backcross and allotetraploid plants from crosses between tepary beans and common beans. Available at: https://academic.oup.com/jhered/article-abstract/75/2/93/765293.
Google Scholar: Author Only Title Only Author and Title

Tsukagoshi, H., Busch, W., and Benfey, P. N. (2010). Transcriptional regulation of ROS controls transition from proliferation to differentiation in the root. Cell 143, 606–616. doi: 10.1016/J.CELL.2010.10.020/ATTACHMENT/F4808132-6242-40E4-A4F7-A40762FFE4AE/MMC3.XLS.
Google Scholar: Author Only Title Only Author and Title

Tweneboah, S., and Oh, S. K. (2017). Biological roles of NAC transcription factors in the regulation of biotic and abiotic stress responses in solanaceous crops. J Plant Biotechnol 44, 1–11. doi: 10.5010/JPB.2017.44.1.001.
Google Scholar: Author Only Title Only Author and Title

VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. J Dairy Sci 91, 4414–4423. doi: 10.3168/jds.2007-0980.
Google Scholar: Author Only Title Only Author and Title

Wallace, J. G., Zhang, X., Beyene, Y., Semagn, K., Olsen, M., Prasanna, B. M., et al. (2016). Genome-wide association for plant height and flowering time across 15 tropical maize populations under managed drought stress and well-watered conditions in Sub-Saharan Africa. Crop Sci 56, 2365–2378. doi: 10.2135/cropsci2015.10.0632.
Google Scholar: Author Only Title Only Author and Title

Wu, J. X., Li, J., Liu, Z, Yin, J., Chang, Z. Y., Rong, C., et al. (2015). The Arabidopsis ceramidase AtACER functions in disease resistance and salt tolerance. Plant Journal 81, 767–780. doi: 10.1111/tpj.12769.
Google Scholar: Author Only Title Only Author and Title

Yokochi, Y., Fukushi, Y., Wakabayashi, K.-I., Yoshida, K., and Hisabori, T. (2021). Oxidative regulation of chloroplast enzymes by thioredoxin and thioredoxin-like proteins in Arabidopsis thaliana. Proc Natl Acad Sci U S A 118. doi: 10.1073/pnas.2114952118/-/DCSupplemental.
Google Scholar: Author Only Title Only Author and Title

Zhang, Q., Li, J, Zhang, W., Yan, S., Wang, R., Zhao, J., et al. (2012). The putative auxin efflux carrier OsPIN3t is involved in the drought stress response and drought tolerance. Plant Journal 72, 805–816. doi: 10.1111/j.1365-313X.2012.05121.x.
Google Scholar: Author Only Title Only Author and Title

Zheng, P., Wu, J. X., Sahu, S. K., Zeng, H. Y., Huang, L. Q., Liu, Z, et al. (2018). Loss of alkaline ceramidase inhibits autophagy in Arabidopsis and plays an important role during environmental stress response. Plant Cell Environ 41, 837–849. doi: 10.1111/pce.13148.
Google Scholar: Author Only Title Only Author and Title