Diel fluctuations in *in-vivo* SnRK1 activity in Arabidopsis rosettes during light-dark cycles

Omri Avidan[1], Thiago A. Moraes[1, a], Virginie Mengin[1, b], Regina Feil[1], Filip Rolland[2,3], Mark Stitt[1,*] and

John E. Lunn[1,*]

[1]Max Planck Institute of Molecular Plant Physiology, Am Muehlenberg 1, 14476 Potsdam-Golm,

Germany

[2]Laboratory of Molecular Plant Biology, KU Leuven, Leuven, Belgium

[3] KU Leuven Plant Institute (LPI), Leuven, Belgium

*Authors for correspondence: mstitt@mpimp-golm.mpg.de, lunn@mpimp-golm.mpg.de

[a]Current address: Crop Science Centre, Lawrence Weaver Road, Cambridge CB3 0LE, UK

[b]Current address: Department of Biological Sciences, University of Essex, Colchester, CO4 3SQ, UK.

ORCID

OA: 0000-0002-1844-1280 (avidan@mpimp-golm.mpg.de)

TAM: 0000-0002-7156-6947 (ta507@cam.ac.uk)

VM: 0000-0002-4958-4209 (v.mengin@essex.ac.uk)

RF: 0000-0002-9936-1337 (feil@mpimp-golm.mpg.de)

FR: 0000-0003-2601-0752 (filip.rolland@kuleuven.be)

MS: 0000-0002-4900-1763 (mstitt@mpimp-golm.mpg.de)

JEL: 0000-0001-8533-3004 (lunn@mpimp-golm.mpg.de)

Running Title: Diel fluctuations in SnRK1 activity in leaves

One sentence summary: *In vivo* SnRK1 activity shows an unexpected diel response and a complex

relationship with trehalose 6-phosphate and other possible metabolic regulators.

Abstract

SUCROSE-NON-FERMENTING1 (SNF1)-RELATED KINASE1 (SnRK1) is a central hub in carbon and energy signalling in plants, and is orthologous with SNF1 in yeast and the AMP-ACTIVATED PROTEIN KINASE (AMPK) in animals. Previous studies of SnRK1 relied on *in-vitro* activity assays or on monitoring the expression of putative marker genes. Neither approach gives unambiguous information about *in-vivo* SnRK1 activity. We have monitored *in-vivo* SnRK1 activity using Arabidopsis (*Arabidopsis thaliana*) reporter lines that express a chimeric polypeptide with a SNF1/SnRK1/AMPK-specific phosphorylation site. We investigated responses during an equinoctial diel cycle, and after perturbing this cycle. As expected, *in vivo* SnRK1 activity rose towards the end of the night and rose even further when the night was extended. Unexpectedly, although sugars rose after dawn, SnRK1 activity did not decline until about 12 hours into the light period. The sucrose signal trehalose 6-phosphate (Tre6P) has been shown to inhibit SnRK1 *in vitro*. We introduced the SnRK1 reporter into lines that harboured an inducible *TREHALOSE-6-PHOSPHATE SYNTHASE* construct. Elevated Tre6P decreased *in-vivo* SnRK1 activity in the light period, but not at the end of the night. Reporter polypeptide phosphorylation was sometimes negatively correlated with Tre6P, but a stronger and more widespread negative correlation was observed with glucose 6-phosphate. We propose that SnRK1 operates within a network that controls carbon utilization and maintains diel sugar homeostasis, and that Tre6P, hexose phosphates and the circadian clock contribute to regulation of SnRK1 activity in a context-dependent manner, and SnRK1-signalling is further modulated by factors that act downstream of SnRK1.

Keywords: *Arabidopsis thaliana*, SnRK1, trehalose 6-phosphate, glucose 6-phosphate, diel, circadian clock, starch

Abbreviations: GEN, general SnRK1 reporter; Glc1P, glucose 1-phosphate; Glc6P, glucose 6-phosphate; NUC, nuclear SnRK1 reporter; SnRK1, SUCROSE-NON-FERMENTING1-RELATED KINASE1; TPP, trehalose-6-phosphate phosphatase; TPS, trehalose-6-phosphate synthase; Tre6P, trehalose 6-phosphate; UDP-Glc, UDP-glucose.

Introduction

Plant SUCROSE-NON-FERMENTING1-RELATED PROTEIN KINASES 1 (SnRK1) belongs to a eukaryotic family of Ca^{+2}-independent Ser/Thr protein kinases. SnRK1 is implicated in maintaining cellular homeostasis and energy balance alongside linking carbon (C) availability and growth, and has been extensively studied in recent years (Polge and Thomas, 2006; Jossier *et al.*, 2009; Hulsmans *et al.*, 2016; Baena-González *et al.*, 2007; Paul *et al.*, 2018; 2020; Crepin and Rolland, 2019; Baena-González and Lunn, 2020). These functions are broadly similar to those observed for its yeast and animal orthologues, SUCROSE-NON-FERMENTING1 (SNF1) and AMP-ACTIVATED PROTEIN KINASE (AMPK) but our understanding of the functions and regulation of SnRK1 in plants is much less complete.

SnRK1 is a heterotrimer consisting of a catalytic (α) subunit and two regulatory (β and βγ) subunits that modify complex stability, activity, interactions and cellular localization (Polge *et al.*, 2008; Nietzsche *et al.*, 2016; Wang *et al.*, 2020). It is located in the nucleus and cytosol (Nietzsche *et al.*, 2014; Blanco *et al.*, 2019; Ramon *et al.*, 2019) and has also been reported to be present in plastids (Ávila-Castañeda *et al.*, 2014; Ruiz-Gayosso *et al.*, 2018). According to current knowledge, the mode of regulation differs from that of SNF1 and AMPK. The regulatory properties of these kinases have been studied mainly by *in-vitro* assays, which monitor phosphorylation of peptides containing an evolutionarily conserved SNF1/SnRK1/AMPK-specific recognition sequence: [MLVFI].X.[RKH].X.X.**S**.X.X.X.[LFIMV] or the less favoured [MLVFI].[RKH].X.X.X.**S**.X.X.X.[LFIMV] (Dale *et al.*, 1995). Such studies have demonstrated that animal AMPK, and to a certain degree also yeast SNF1, are directly regulated by AMP and/or ADP (Hardie and Carling, 1997; Hardie *et al.*, 1998; Mayer *et al.*, 2011). In mammals, these adenylates inhibit T-loop dephosphorylation, thereby activating the catalytic (α) subunit in low energy conditions and/or in response to low C availability (García-Salcedo *et al.*, 2014; Crozet *et al.*, 2014). In contrast, plant SnRK1 is not activated by adenylates in *in-vitro* assays. Instead, it is inhibited *in vitro* by sugar phosphates, such as trehalose 6-P (Tre6P), glucose 6-phosphate (Glc6P) and glucose 1-phosphate (Glc1P) (Toroser *et al.*, 2000; Lu *et al.*, 2007; Zhang *et al.*, 2009; Emanuelle *et al.*, 2015; Broeckx *et al.*, 2016; Zhai *et al.*, 2017; 2018; Ruiz-Gayosso *et al.*, 2018). Inhibition by sugar phosphates could potentially link SnRK1 activity to C availability rather than energy status (Hara *et al.*, 2013; Nunes *et al.*, 2013; Herzig and Shaw, 2017). These *in-vitro* studies also indicated that inhibition by Tre6P requires an unknown proteinaceous factor that is present in growing tissues but not in mature source leaves (Toroser *et al.*, 2000; Zhang *et al.*, 2009; Emanuelle *et al.*, 2015). Subsequent studies have shown that Tre6P can bind directly to the SnRK1 catalytic α subunit, thereby interfering with binding of the SnRK1-activating kinases GRIK1/2, leading to inhibition of SnRK1 activity (Glab *et al.*, 2017; Zhai *et al.*, 2018; Hwang et al., 2019). However, GRIK1/2 might not be the only mechanism by which Tre6P inhibits SnRK1 activity. As GRIK1/2 are widely expressed in

source leaves, Zhai et al. (2018) concluded that they were unlikely to be the unknown proteinaceous

factor from young tissues that was referred to by Zhang et al. (2009).

Glc6P and Glc1P are intermediates in glycolysis, gluconeogenesis and in the synthesis and

degradation of major carbohydrate reserves like starch and sucrose. Tre6P is an intermediate in the

pathway of trehalose biosynthesis. It is synthesized from UDP-Glc and Glc6P by TREHALOSE-6-

PHOSPHATE SYNTHASE (TPS) and is dephosphorylated to trehalose by TREHALOSE-6-PHOSPHATE

PHOSPHATASE (TPP) (Cabib and Leloir, 1958). Trehalose is a major C storage metabolite, osmolyte and

stress protectant in fungi, but has been largely replaced by sucrose in vascular plants, allowing neo-

functionalization of the trehalose biosynthesis pathway (Goddijn and van Dun, 1999; Lunn *et al.*, 2014;

Lunn, 2016). In addition to catalytically active class I TPS proteins), plants possess a large family of

catalytically inactive class II TPS proteins (Leyman *et al*., 2001; Harthill *et al*., 2006; Lunn, 2007;

Vandesteene *et al.*, 2010; Lunn et al., 2014; Delorge *et al.*, 2015) as well as a large family of TPP

proteins (Leyman *et al*., 2001; Vandesteene *et al.*, 2012; Kretzschmar *et al.*, 2015). Tre6P is considered

to serve as a signal of sucrose availability (Lunn *et al.*, 2006), acting to regulate C allocation between

sucrose and other photosynthesis products in source leaves, and to link sucrose supply and growth in

sink organs (Schluepmann *et al.*, 2003; Martins *et al.*, 2013; Wahl *et al.*, 2013; Figueroa *et al.*, 2016;

Paul *et al.*, 2018; 2020; Fichtner and Lunn, 2021).

The possible contribution of Glc6P and Glc1P to regulation of SnRK1 activity *in planta* has not

been addressed in detail. Interactions between Tre6P and SnRK1 have been more extensively studied,

mainly by investigating changes in SnRK1 marker transcript abundance (Zhang *et al.*, 2009; 2015;

Henry *et al.*, 2015; Bledsoe *et al.*, 2017) and post-translational regulation of target enzymes Baena-

González and Sheen, 2008; Figueroa *et al.*, 2016). The former builds on the work of Baena-González

et al. (2007) who, using a protoplast overexpression system, identified over 1000 genes as

downstream targets of SnRK1, such as *DIN6*, *DIN1*, *BCAT2* and *EXP10*, whose expression is regulated

by SnRK1-mediated phosphorylation of upstream transcription factors, such as bZIP11 (Ma et al.,

2011) and bZIP63 (Mair et al., 2016). Their expression has been widely used as a sensitive readout of

SnRK1 activity, either via promoter activation assays (Rodrigues *et al.*, 2013; Ramon *et al.*, 2019) or by

analyses of transcript abundance. Abundance of SnRK1-induced transcripts is typically low in a diel

cycle but rises strongly when plants are starved by transfer to continuous darkness (Baena-González

et al., 2007; Baena-González and Sheen, 2008; Usadel *et al.*, 2008; Flis *et al.*, 2016) as expected if

SnRK1 is activated in C starvation conditions (Mair *et al.*, 2015; Pedrotti *et al.*, 2018; Ramon *et al.*,

2019). Their abundance often changes reciprocally to Tre6P levels in starvation treatments and other

perturbations in several species and tissues (Martínez-Barajas *et al.*, 2011; Wingler *et al.*, 2012; Nunes

et al., 2013; Henry *et al.*, 2015; Nuccio *et al.*, 2015; Baena-González and Lunn, 2020; Peixoto *et al.*,

2021) providing correlative evidence for the idea that Tre6P inhibits SnRK1 *in vivo*. Application of

permeable Tre6P analogues to Arabidopsis led to changes in transcript abundance for a subset of

these SnRK1 downstream target genes that were consistent with Tre6P acting to inhibit SnRK1

(Griffiths *et al*., 2016). Further support for partial convergence of the SnRK1 and Tre6P signalling

pathways was provided by the identification of several class II TPSs as transcriptional and post-

translational SnRK1 targets (Glinski and Weckwerth, 2005; Harthill *et al*., 2006; Cho *et al.*, 2016;

Nukarinen *et al.*, 2016). However, the functional significance of this observation remains unclear

because the precise role of the TPS class II proteins has not yet been defined (Schluepmann and Paul,

2009; Ponnu *et al.*, 2011; Yang *et al*., 2012; Lunn et al. 2014; Delorge *et al.*, 2015; Figueroa and Lunn,

2016). More recently, it was demonstrated that mutant lines with increased or reduced expression of

SnRK1 exhibit an altered relationship between sucrose and Tre6P in diel cycles, as well as modified

anaplerotic flux of carbon to the tricarboxylic acid cycle that is reminiscent of responses to Tre6P

(Peixoto *et al.*, 2021). Furthermore, mutations in the *SnRK1α1* and *SnRK1βγ* genes were shown to

suppress developmental defects in the Arabidopsis *tps1* mutant (Zacharaki *et al*., 2022).

Taken together, these results point to SnRK1 and Tre6P signalling being closely intertwined.

They also indicate that SnRK1 signalling is involved not only in starvation responses, but also in the

coordination of metabolism in benign conditions where plants are not C-starved or facing a large C

excess (see also Baena-González and Lunn, 2020). However, many open questions remain; for

example, negative correlations between Tre6P levels and the transcript abundance of SnRK1-induced

genes are seen in mature tissues even though Tre6P does not inhibit SnRK1 activity in *in-vitro* assays

with material from mature tissues (see above and Baena-González and Lunn, 2020). Moreover, there

is an inconsistent correlation between Tre6P levels and transcript abundance of SnRK1 target genes

in *in-vitro* cultured excised maize kernels (Bledsoe *et al.*, 2017, see also Baena-González and Lunn,

2020). It remains possible that Tre6P also acts on signalling downstream of SnRK1, as is the case in

yeast (Deroover *et al*., 2016), resulting in changes in downstream transcript abundance that are not

directly due to inhibition of SnRK1 itself. Interpretation is further complicated by potential interactions

between SnRK1 and TARGET OF RAPAMYCIN (TOR) signalling (Lastdrager *et al.*, 2014; Margalha and

Confraria, 2019; Ryabova *et al*., 2019).

The two main approaches employed in plants to study SnRK1 regulation have shortcomings

which have not been addressed to date. The *in-vitro* assay is direct and highly specific, but the use of

crude extracts or purified proteins may bring about other complications due to loss of the *in-vivo*

context, such as artificial interactions or potential loss of regulatory elements that otherwise would

have an effect on activity (Baena-González and Lunn, 2020). Monitoring changes in expression of

SnRK1 marker genes is sensitive, but gene expression is an indirect readout that may be prone to

secondary effects that are unrelated to SnRK1 activity. Indeed, there is often poor agreement between
SnRK1 activity in the *in-vitro* phosphorylation assay and changes in SnRK1 target gene expression
(Debast *et al.*, 2011; Martínez-Barajas *et al.*, 2011; Baena-González and Lunn, 2020).

Working in yeast, Deroover *et al.* (2016) developed a specific *in-vivo* SNF1 reporter that
harbours an SNF1/SnRK1/AMPK-specific recognition sequence-containing peptide, based on rat
ACETYL COA CARBOXYLASE 1 (ACC1), an established AMPK target (Dale *et al.*, 1995). The
phosphorylation status of this peptide can be monitored by extraction and immunoblot analysis with
a validated commercial phospho-specific antibody, providing a direct measure of *in-vivo* SNF1 activity.
Subsequently, modified ACC1 reporters were used in Arabidopsis to study the regulation of SnRK1
activity by nitrogen status (Sanagi *et al.*, 2021) and in other responses (Muralidhara *et al*.,
2021;cHenninger *et al*., 2022; Belda-Palazon *et al*., 2022).

We have used Arabidopsis ACC1 reporter lines to investigate *in-vivo* SnRK1 activity during a
diel cycle in benign conditions, when the C supply is only slightly restricting growth, and after changing
the light regime to modify the C supply. We have also introduced the reporter construct into
Arabidopsis lines in which Tre6P levels can be increased in an inducible manner (Martins *et al*., 2013;
Figueroa *et al*., 2016). We show that SnRK1 activity does not change reciprocally to overall C
availability, with a peak at dawn and a minimum at dusk. Instead, SnRK1 activity appears to be
maintained in the light period, declines after darkening and rises towards the end of the night.
Furthermore, SnRK1 activity *in vivo* is not strongly or consistently inhibited by an induced increase in
Tre6P. Multiple regression analyses of metabolite levels indicate that the link between *in-vivo* SnRK1
activity and the level of Tre6P is context-dependent and point to SnRK1 activity being regulated *in vivo*
by additional mechanisms, including inhibition by Glc6P.

Results

Reporter constructs to quantify in vivo *SnRK1 activity*

To obtain a direct *in-vivo* readout of SnRK1 activity in Arabidopsis we employed transgenic SnRK1-
reporter lines expressing a peptide derived from the Ser79 phosphorylation site of ACC1 fused to
GREEN FLUORESCENT PROTEIN (GFP), which was used to normalize for the level of expression of the
reporter polypeptide (see Supplementary Figure S1 for full description) and to monitor its localization
(see below). The level of phosphorylation of the ACC1 peptide (pACC) was assessed using a specific
anti-phospho-site antibody, and an anti-GFP antibody was used to normalize for expression level and
gel loading. To increase statistical and biological robustness, we used three-four biological replicates
per time point. An additional control sample (extended night) was loaded in all gels, and used to

correct for between-gel variation. We took the normalized pACC/GFP signal ratio as a readout of *in-*
vivo SnRK1 activity at a given time point (for details see Methods and Supplementary Figure S1).

Two transgenic lines were used: (i) expressing a nuclear reporter (NUC), that contains the SV40
nuclear location signal (NLS) and (ii) expressing a general reporter (GEN), in which GFP is observed in
both the nucleus and the cytoplasm (Supplementary Figure S2).

Validation of the reporter construct in Arabidopsis rosettes
The specificity of the ACC1 reporter peptide was previously tested in yeast by incubating WT and *snf1*
mutant cells with fermentable glycerol and ethanol to induce SNF1 or with glucose to inhibit SNF1
activity (Deroover *et al.*, 2016). The peptide was phosphorylated under fermenting conditions but not
following glucose addition in the wild type background, while no phosphorylation was observed in the
snf1 background, suggesting that the reporter is indeed SNF1-specific in yeast. Sanagi *et al.* (2021)
validated an ACC1-based SnRK1 reporter in Arabidopsis mesophyll protoplasts, by showing that
phosphorylation is increased by transient expression of wild-type SnRK1α1 but not by expression of a
mutated form, SnRK1α1^{K48A}, which is defective in the SnRK1 ATP-binding pocket (K48M). They also
showed that there is less phosphorylation in protoplasts prepared from inducible *snrk1α* knockdown
plants than in protoplasts prepared from wild-type plants.

We took two approaches to further validate the reporters for use in whole Arabidopsis
rosettes. In the first approach, we subjected GEN and NUC lines to 3-h of extended darkness, starting
at the end of the night (Supplementary Figure S3A; Supplementary Dataset S1 – Exp.1). Extended
darkness is known to activate SnRK1 (see Introduction). As expected, phosphorylation of both
reporters increased in extended darkness, with phosphorylation being significantly higher at ZT25
(zeitgeber time, i.e. 25 h after the previous dawn and into the extended night) and rising further at
ZT27 in NUC, and being significantly higher by ZT27 in GEN. In the second approach, the NUC reporter
was crossed to a 35:SnRK1α1 line that has elevated SnRK1 activity and to the 35S:SnRK1α1^{K48M} line
expressing a catalytically inactive mutant protein (Jossier *et al.*, 2009; Crozet *et al*., 2014; Cho *et al.*,
2016) (Supplementary Figure S3B; Supplementary Dataset S1 – Exp.2). The crossed lines and a NUC
control were grown in equinoctial growth conditions for 22 days after sowing (DAS), and then
harvested at ZT0, 8, 16, 24 and 28 (i.e. 4-h of extended darkness). We confirmed expression of SnRK1
proteins in these lines by immunoblotting with anti-SnRK1α1 antibody (Supplementary Figure S3C).
Phosphorylation of the NUC polypeptide was consistently higher in NUC x 35S:SnRK1α1 than in NUC x
35S:SnRK1^{K48M} (significant at ZT0, 8, 16, 24 and 28). Compared to the NUC control, reporter
phosphorylation in NUC x 35S:SnRK1^{K48M} was significantly lower at ZT0, 8, 24 and 28 (with SnRK1^{K48M}
apparently functioning as a dominant negative mutant protein; Cho et al., 2016), and reporter

phosphorylation in NUC x 35S:SnRK1α1 was essentially control-like at ZT0, 8, 24 and 28 and

significantly higher at ZT16. This experiment also confirmed the rise in reporter phosphorylation in the

extended night, further validating the reporter.

Diel changes in SnRK1 activity in equinoctial growth conditions

The diel response of the NUC control in Supplementary Figure S3B revealed a somewhat unexpected

pattern, with the signal being quite high throughout the light period, decreasing rapidly at the

beginning of night, and then rising in the last part of the night. This was unexpected as photosynthesis

provides a higher C supply in the light, which is expected to inhibit SnRK1. We investigated if this

unexpected diel response was robust and reproducible. Independent experiment with equinoctial

light-dark regimes (three with NUC, two with GEN) yielded a robust and reproducible diel response

with SnRK1 activity at dawn being maintained during the light period, followed by a decline after

darkening to a minimum at about ZT16 and an increase in the last part of the night (Figure 1A). The

NUC and GEN lines exhibited similar responses.

We explored diel changes in metabolic status and in potential metabolite effectors of SnRK1

(Figure 1B) in the same plant material. Obvious candidates include sugars (glucose, sucrose) and starch

as indicators of C status, and sugar phosphates, such as Tre6P, Glc6P and Glc1P, all of which were

suggested previously to be directly or indirectly involved in the regulation of SnRK1 (Toroser *et al.*,

2000; Lu *et al.*, 2007; Zhang *et al.*, 2009; Emanuelle *et al.*, 2015; Broeckx *et al.*, 2016; Hulsmans *et al.*,

2016; Zhai *et al.*, 2017; 2018; Ruiz-Gayosso *et al.*, 2018). The diel responses of metabolites were

reproducible and resembled previous reports (Lunn *et al.*, 2006; Sulpice *et al.*, 2014; Mengin *et al.*,

2017; Flis *et al.*, 2019; Moraes *et al.*, 2019). In brief, starch accumulated in a near-linear manner during

the light period and was mobilized in a near-linear manner at night, sucrose gradually increased in the

light to a peak at ZT12 (ED) and gradually decreased during the night, and glucose levels showed a 1.5-

to 3-fold transient increase at ZT4 but otherwise remained relatively low throughout the diel cycle

(see also Mengin *et al.*, 2017). Tre6P changed in parallel with sucrose, as expected from previous

studies (Martins *et al.*, 2013; Figueroa *et al.*, 2016; Peixoto *et al.*, 2021). Glc6P and Glc1P levels

remained fairly stable throughout the day, rose at the beginning of the night until ZT16 and then

decreased.

These experiments did not reveal pronounced reciprocal changes in Tre6P levels and SnRK1

activity during the diel cycle, except late in the night (between ZT16-24) when Tre6P levels decreased

and phosphorylation of both NUC and GEN polypeptides increased. In addition, no reciprocal changes

were observed between NUC or GEN phosphorylation and sugar levels, except for sucrose in the last

part of the night. However, Glc6P and Glc1P did show a reciprocal relationship with SnRK1 activity

through the night; Glc6P and Glc1P initially increased after dusk (from ZT12-16), and then decreased

(from ZT16-24), while phosphorylation of NUC and GEN decreased during the first part of the night

and then increased (see later for more analysis).

Perturbations of the light regime

To further explore how SnRK1 is regulated by changes in metabolites, by light and/or by endogenous

(e.g. circadian clock) rhythms, we next investigated NUC and GEN phosphorylation during

perturbations of the light regime, including standard equinoctial light treatments as a control.

265   The first treatment was based on a perturbation described in Moraes *et al.* (2019). Plants were

grown in equinoctial conditions under 160 µmol m^{-2} s^{-1} irradiance for 19 days, and then transferred to

60 µmol m^{-2} s^{-1} for one light period (treatment light period), darkened at ZT12 at the same time as in

the initial growth conditions (treatment night) and then returned to 160 µmol m^{-2} s^{-1} on the next day

(recovery light period). Control plants were left in the original growth conditions (Figure 2). The

response of starch and sugars closely resembled that in Moraes *et al.* (2019). Compared to control

plants, a single low-irradiance day led to slower starch accumulation and a decrease in the levels of

sucrose and glucose during the treatment light period, and slower starch mobilization and lower sugar

levels in the following night (Figure 2A; Supplementary Dataset S1 – Exp.3). During the recovery light

period, glucose was higher and starch accumulation was faster than in the light period in control

plants, indicating that growth was restricted (see also Gibon *et al.*, 2004; Sulpice *et al.*, 2014b; Mengin

et al., 2017).

277   Phosphorylation of NUC and GEN polypeptides (Figure 2B) showed a combination of expected

and unexpected features, which were partially shared between the NUC and GEN lines but also

showed some differences. Rather surprisingly, a sudden decrease in irradiance had no immediate

impact on NUC or GEN phosphorylation in the treatment light period. However, it did have significant

effects in the following night and during the recovery day. For NUC, there was no initial drop in

phosphorylation after darkening, with the result that phosphorylation was significantly higher than

the control (where it decreased in the first part of the night). However, after ZT20, there was a sharp

decrease in NUC phosphorylation and this persisted during the recovery light period. For GEN,

phosphorylation increased significantly compared to the control by ZT16, and increased even further

as the night progressed, followed by a decrease to control levels in the recovery light period.

287   It is rather surprising that, after a shift to low light, the decrease of sugar levels was not

accompanied by an increase in NUC or GEN phosphorylation in the light period. This indicates that at

this time in the diel cycle, nuclear and extra-nuclear SnRK1 activity are not responding to current C

availability as reflected by overall sugar levels. At night, NUC and GEN phosphorylation were higher in

plants exposed to low irradiance during the preceding light period than in the control, reflecting the
lower C availability, but this response was much more marked for GEN than NUC. Indeed, NUC signal
fell at the end of the night, even though sugars remained low. In the recovery day, when sugars were
higher than in the control, NUC phosphorylation was lower than the control as would be expected if
it were responding to current C availability. However, GEN phosphorylation was high, indicating that
extra-nuclear SnRK1 activity was not responding to current C availability. Indeed, considering that GEN
is a composite of nuclear and of cytoplasmic activity, it is likely that cytoplasmic SnRK1 activity was
actually increased since NUC signal activity was decreased at this time (see above).

The second treatment was based on an experiment reported by Fernandez *et al.* (2017), in
which plants were grown in equinoctial conditions with 160 µmol m^{-2} s^{-1} irradiance for 19 days and
then transferred to continuous lower irradiance. The original study of Fernandez *et al*. (2017)
investigated starch degradation in the light, and the low irradiance was used to avoid over-
accumulation of starch in continuous light. The rationale for the current experiment was to test
whether the diel changes in phosphorylation activity of SnRK1 are driven by the external light regime
or by endogenous cues.

We grew plants in standard equinoctial conditions at 160 µmol m^{-2} s^{-1}, then transferred them,
at dawn on day 20 after sowing, to continuous light with an irradiance of 90 µmol m^{-2} s^{-1}, and sampled
them at 4-h intervals for 32 h (Figure 3). Control plants were left in the original growth conditions.
After transfer to continuous low light, starch accumulated for about 12 h, although more slowly than
in control plants, and starch levels then plateaued or even declined slightly in the subjective night from
ZT12 onwards (Figure 3A; Supplementary Dataset S1 – Exp.4). Sugar levels were lower than in the
control in the subjective day period (ZT0-12) but rose to reach higher levels from ZT16 onwards. These
changes of starch and sucrose reflect a gradual but progressive increase in the rate of starch
mobilization from about ZT10 onwards (see Fernandez *et al.*, 2017 for details). Compared to the
control, Glc6P was only marginally and non-significantly lower in the subjective day, and was
consistently lower in the subjective night (Supplementary Figure S4). Once again, we observed that
lower sugar levels during the first 12 h of the 24-h cycle were not accompanied by a significant increase
in phosphorylation of NUC or GEN, which essentially exhibited a control-like response until ZT12
(Figure 3; Supplementary Dataset S1 – Exp.4). In both lines, SnRK1 activity in continuous low light was
higher in the subjective night than in the control during the night. Nevertheless, the decline in SnRK1
activity that is observed after darkening control plants at ZT12 was also observed under low
continuous light, although in a weakened form. NUC phosphorylation showed a gradual significant
decrease (relative to ZT12) from ZT14 to ZT24, while GEN phosphorylation started to decrease by ZT12

and declined until ZT16 (Figure 3B). In the next subjective day, phosphorylation levels of both NUC

and GEN resembled those in the control.

These observations confirmed that a fall in C availability due to decreased irradiance does not

lead to increased SnRK1 activity in the first part of the 24-h cycle. They also point to an endogenous

component contributing to the decline in SnRK1 activity from about ZT12 onwards. Furthermore, we

noted a reciprocal relationship between diel changes in Glc6P and SnRK1 activity, particularly with

NUC phosphorylation.

Response in a T28 cycle

The Arabidopsis circadian clock is mainly entrained by light at dawn and imposes an endogenous 24-

h rhythmicity even when the external cycle deviates from 24 h (Millar *et al.*, 1995; Locke *et al.*, 2005;

Graf *et al.*, 2010). Growing plants in a light-dark cycle that deviates from 24 h is therefore a good way

to search for oscillatory responses that are driven directly or indirectly by the clock.

We grew plants in a 14-h light / 14-h dark (T28) or a standard 12-h light / 12-h dark (T24) cycle

for 19 days with 160 μmol m^{-2} s^{-1} irradiance and harvested them at 4-h intervals on day 20 after sowing,

including a 4-h or 8-h period of extended darkness for the T28 and T24 control, respectively (Figure

4). As previously seen (Graf *et al.*, 2010), in a T28 cycle starch is exhausted at about ZT24, resulting in

transient C starvation and low sucrose levels from ZT24 to ZT28 (i.e., the four hours preceding the next

dawn) (Figure 4A; Supplementary Dataset S1 – Exp.5) as well as lower Glc6P levels through the night

(Supplementary Figure S5A). Correspondingly, phosphorylation of NUC and GEN at ZT0 (dawn) was

almost 6- and 3-fold higher in the T28 cycle than the T24 cycle, comparable to the increase observed

after extending the night in a T24 cycle (Figure 4B; Supplementary Dataset S1 – Exp.5, see also

Supplementary Fig. S3A). After illumination, NUC and GEN phosphorylation decreased strongly in a

T28 cycle, falling to levels that were lower than in a T24 cycle for most of the remaining light-dark

cycle (significant for both NUC and GEN at ZT8, ZT14 and ZT24). Importantly, NUC and GEN

phosphorylation decreased between ZT12 and ZT14 in a similar manner in both the T28 and T24 cycle

plants, even though the former were still in the light while the latter were already in the dark. Together

with the response in continuous light (see above), these observations point to a component with 24-

h rhythmicity contributing to the regulation of SnRK1 activity.

Manipulation of Tre6P levels

Tre6P is a signal of sucrose levels (Lunn *et al.*, 2006; 2014; Figueroa and Lunn, 2016; Fichtner and Lunn,

2021) and is widely accepted as an inhibitor of SnRK1, at least *in vitro* (Zhang *et al.*, 2009; Zhai *et al.*,

2018). However, the evidence that these interactions occur *in vivo* is largely correlative (Martínez-

Barajas *et al.*, 2011; Wingler *et al.*, 2012; Nunes *et al.*, 2013; Henry *et al.*, 2015; Nuccio *et al.*, 2015;

Griffiths *et al*., 2016; Peixoto *et al.*, 2021; Zacharaki *et al*., 2022). We exploited our SnRK1 reporter

lines to develop a genetic approach to test whether Tre6P inhibits SnRK1 activity *in vivo*. We have

previously used inducible-TPS (iTPS) lines, expressing the *Escherichia coli* TPS under the control of the

Aspergillus nidulans ethanol-inducible ALCOHOL DEHYDROGENASE REGULATOR (alcR) and alcA

promoter to investigate the impact of a short-term induced increase in Tre6P on starch, organic acid

and nitrogen metabolism (Martins et al.,2013; Figueroa et al., 2016). We now introgressed the NUC

and GEN reporters into a well characterized iTPS line (TPS29.2; Martins et al., 2013) and into its

corresponding empty-vector control, alcR. Four homozygous lines were generated: two lines

containing both the alcR gene and the inducible *TPS* (NUC x iTPS #6.3 and GEN x iTPS #2.4), and two

control lines with only the alcR gene (NUC x alcR #1.2 and GEN x alcR #1.1).

These lines were grown in long-day conditions (16-h photoperiod, 160 µmol m^{-2} s^{-1} irradiance).

On day 21 after sowing, plants were sprayed at ZT1 with either water (control) or 2% (v/v) ethanol to

induce the expression of the bacterial TPS (Figure S6). Two whole rosettes were harvested and pooled

at 4, 6 and 8h after spraying. Transient expression of the bacterial TPS protein was detectable within

4 h of induction (Supplementary Figure S6B), consistent with previous experiments with the parental

iTPS line (Martins *et al.*, 2013; Figueroa *et al.*, 2016). After ethanol induction, there was a strong and

consistent decrease in SnRK1 nuclear activity (in NUC x iTPS) and a small decrease in overall SnRK1

activity after 4 and 6 but not after 10-h in GEN x iTPS (Supplementary Figure S6A; Supplementary

Dataset S1 – Exp.6). This led us to focus on the NUC x iTPS line (#6.3) in subsequent experiments. There

was no consistent effect of ethanol on NUC or GEN phosphorylation in the control alcR lines, in which

alcR is expressed but there is no bacterial *TPS* gene to be expressed under the control of the alcA

promoter. This control excludes any risk of off-target effects in the presence of ethanol or due to

expression of alcR (Randall, 2021).

To confirm the inhibitory effect of Tre6P induction in the light period, we conducted a more

extensive experiment. Line #6.3 was grown in equinoctial conditions under 160 µmol m^{-2} s^{-1} irradiance

for 20 days (the standard conditions of previous experiments), sprayed on 21 DAS with either water

or 2% (v/v) ethanol at ZT4, and triplicate samples were harvested at ZT7, 9 and 11 and separately

analysed (Figure 5; Supplementary Dataset S1 – Exp.7). Ethanol spraying led to a 41, 169 and 130%

increase in Tre6P levels at ZT7, ZT9 and ZT11, compared to control plants (Figure 5). Starch

accumulation was marginally enhanced, sucrose levels decreased at ZT7 and ZT11 (Figure 5), reducing

sugars were not consistently or significantly affected, and the levels of many organic acids increased

(Supplementary Figure S7; Supplementary Dataset S1 – Exp.7), as previously reported by Figueroa *et*

al. (2016). NUC phosphorylation showed a significant but small decrease (12-23%) following ethanol

spraying, compared to the water sprayed control (Figure 5). This provides evidence that Tre6P indeed

inhibits nuclear SnRK1 activity *in vivo*, but the impact seems to be rather small at the whole-rosette

level.

We hypothesized that if Tre6P were not the only regulator of SnRK1, other factors might

counteract, and thereby mask, the effect of Tre6P. Therefore, we carried out a further experiment in

which TPS was induced in conditions when sugars and other metabolites were low, and might not

mask the expected inhibition by Tre6P. Plants were grown in short-day conditions (6-h light / 18-h

dark) for 21 DAS with 160 μmol m^{-2} s^{-1} irradiance, sprayed with either water or 2% (v/v) ethanol

towards the end of the night (ZT20) and harvested at ZT20, 22, 24, 25 and 27 (Figure 6). We included

an alcR control line, and all comparisons were made between ethanol-induced NUC x iTPS (#6.3) and

ethanol-induced NUC x alcR #1.2 as the control. Under these conditions, starch was almost completely

exhausted and soluble sugars were at low levels (Supplementary Figure S8A). Tre6P levels declined

strongly between ZT20 and ZT27 in the control. Induction of TPS slowed down this decline (Figure 6),

with Tre6P levels at ZT24, 25 and 27 being significantly higher in ethanol-sprayed NUC x iTPS #6.3 than

in the NUC x alcR (#1.2) control at the same time (by 59, 112 and 33% , respectively). Phosphorylation

of NUC in the control rose between ZT20 and ZT24, and rose further at ZT25 and ZT27 (Figure 6). This

resembles the response of wild-type plants at the end of the night and in an extended night (see

above). Unexpectedly, induction of TPS did not attenuate but instead promoted this increase in

nuclear SnRK1 activity.

The increase in Tre6P levels was accompanied by a slight slowing of starch breakdown and

lower glucose levels (Supplementary Figure S8A). This confirms the response previously reported in

Martins *et al.* (2013), but in the present case for induction at a later time in the night. The increase in

Tre6P levels was also accompanied by increases in malate, pyruvate and glycerol 3-phosphate (Gly3P)

and a decrease in phospho*enol*pyruvate (PEP) levels (Supplementary Figure S8B; Supplementary

Dataset S1 – Exp.8), indicating that the stimulation by Tre6P of anaplerotic flux that was previously

reported in the light (Figueroa *et al.*, 2016) may also occur at the end of the night. Some of these

downstream responses to elevated Tre6P may lead to the unexpected increase in nuclear SnRK1

activity after iTPS induction.

Overall, these results show that Tre6P is not the sole regulator of SnRK1, and its inhibitory

effect appears to be context dependent. Indeed, in some conditions, secondary changes caused by

Tre6P induction may lead, indirectly, to a stronger positive effect on SnRK1 activity than the direct

inhibitory effect of elevated Tre6P itself.

Correlation analysis of relationships between SnRK1 activity and metabolite levels

Taken together, our experiments point to complex and somewhat flexible regulation of SnRK1 activity

in conditions where C status is changing in a range above that prevailing during starvation. We

therefore investigated more systematically the relationship between diel changes of metabolite levels

and NUC or GEN phosphorylation.

We did this first in the five control experiments in equinoctial conditions of Figure 1. We

performed regression analysis separately for each experiment on all biological replicates and time

points across the diel cycle (Table I, for plots see Supplementary Dataset S2). Tre6P levels were not

correlated with either NUC (R^2 = 0.007, 0.004 and 0.004) or GEN phosphorylation (R^2 = 0.001 and

0.046). Glc6P levels were not consistently related to NUC phosphorylation with R^2 values of 0.048, 0.21

(negative and significant, P = 0.022) and 0.02 but were negatively correlated with GEN

phosphorylation with R^2 values of 0.22 (P = 0.03) and 0.58 (P = 2.24x10^{-5}). Similar relationships were

observed for Glc1P which, due to the near-equilibrium reaction catalysed by phosphoglucomutase,

can be expected to change in parallel with Glc6P. Starch was not correlated with either NUC or GEN

phosphorylation. Sucrose was not correlated with NUC and was negatively correlated with GEN

phosphorylation in one but not the other experiment. Glucose was not correlated with NUC, and

positively correlated with GEN phosphorylation in one but not the other experiment.

We also performed regression analysis separately on the light period (ZT4-12) and night (ZT16-

24) datasets (Supplementary Table S1). There were almost no significant relationships in the light

period. This can be expected as even though the levels of most sugars, sugar-phosphates and many

other metabolites change, SnRK1 activity is rather constant during the light period. During the night

there were many negative relationships between NUC/GEN phosphorylation and many metabolites,

including Tre6P, Glc6P, Glc1P, starch and sucrose. These negative relationships often reflect falling

levels of many metabolites towards the end of the night, as starch is gradually exhausted. They may

individually or collectively contribute to the rise in NUC and GEN activity at this time.

Pairwise correlation analyses between metabolites (Supplementary Table S2) revealed that

most of the measured metabolites are positively correlated with each other. This reflects the highly

coordinated nature of the changes in metabolism in diel cycles, and underlines that some of the

observed correlations between metabolites and NUC or GEN phosphorylation may be secondary.

The diel changes in SnRK1 activity and metabolite levels include abrupt changes after

illumination or darkening. These may not be well represented in correlations based on changes across

the entire cycle. To place more weight on such transient changes, we calculated the difference

between the average absolute levels at consecutive time points and divided this by the time difference

to obtain the direction and rate of change of a given trait in a given time interval. Compared to

absolute values, derivative values provide more information about changes in the balance in

metabolism. These derivative values were used for linear regression analyses as described above,
except that analyses were only performed for the entire diel cycle (Supplementary Table S3;
Supplementary Dataset S2). Changes in both NUC and GEN phosphorylation are significantly and
negatively correlated to changes in Glc6P and Glc1P, with changes in Glc6P showing the strongest
reciprocal agreement with changes in GEN (highest R^2 and lowest p-values).

We also performed correlation analyses in the experiments where plants were transferred to
continuous light, or grown in T28 cycles, or in which iTPS lines were used to elevate Tre6P levels
(Supplementary Dataset S2; Supplementary Tables S4-5, Supplementary Text S1). These analyses
again detected correlations between NUC or GEN phosphorylation and Glc6P and Glc1P, and also with
Tre6P. Further, in the experiments in which Tre6P was induced in the light period, (Fig 5), NUC
phosphorylation correlated negatively with Tre6P in the control plants ($R^2 = 0.67$, p = 0.76) but not
when ethanol was sprayed to increase Tre6P ($R^2 = 0.12$, p = 0.004) (Supplementary Table S6). This is
consistent with the idea that SnRK1 activity is regulated by a network including Tre6P and that this
network adjusts to decrease the sensitivity to Tre6P when Tre6P levels are artificially elevated.

Discussion

Monitoring in vivo *SnRK1 activity in plants by expression of a chimeric protein reporter*

Two approaches have been widely utilized to study the regulation of SnRK1 activity in plants. One is
in-vitro phosphorylation assays using a peptide with a conserved SNF1/SnRK1/AMPK recognition
sequence (Dale *et al.*, 1995). *In-vitro* studies are useful for identifying potential ligands and regulators,
including interacting proteins, and can reveal changes in maximum catalytic activity of SnRK1.
However, removal of the enzyme from its *in-vivo* context means that activity measurements on *in-*
vitro extracts provide only a limited information about activities *in vivo* The other approach is to
monitor *in vivo* readouts of SnRK1 activity, such as changes in the abundance of marker gene
transcripts (Baena-González *et al.*, 2007; Zhang *et al.*, 2009; Ramon *et al.*, 2019; Crepin and Rolland,
2019). However, while relatively easy to quantify, transcripts are only indirect readouts of SnRK1
activity. Their abundance reflects changes in the activity of transcriptional regulators that are the
direct targets for phosphorylation by SnRK1, but can also be influenced by SnRK1-independent
changes in transcriptional or post-transcriptional regulation. The ACC1-based polypeptide reporters
developed by Deroover *et al*. (2016) have been used to monitor *in-vivo* SnRK1 activities in Arabidopsis
(Sanagi et al., 2021; Muralidhara et al., 2021; Henninger et al., 2021; Belda-Palazon et al., 2022), and
we applied this approach to study *in-vivo* SnRK1 activity during the diel cycle in Arabidopsis rosettes.
This approach does have some potential limitations. The ACC1-based polypeptide might be
phosphorylated by other protein kinases in addition to SnRK1, or the extent of phosphorylation may

be affected by changes in the activity of endogenous protein phosphatases. Further, the approach

detects phosphorylation of a heterologous polypeptide, and so might not capture modifications of

SnRK1 function that alter its ability to recognize different endogenous protein targets. In addition to

the tests performed in leaf mesophyll protoplasts by Sanagi et al. (2021), we validated our approach

by confirming that phosphorylation of the reporter increased when SnRK1 activity *in planta* was

enhanced by extended darkness (Supplementary Figure S3A) or overexpression of SnRK1α1, and that

phosphorylation of the reporter decreased when SnRK1 activity was suppressed by expression of a

dominant negative form (35S:SnRK1α1[K48M]; Supplementary Figure S3B). Nevertheless, the potential

complications noted above should be borne in mind when interpreting the results.

SnRK1 protein is located in the nucleus and in various extra-nuclear locations, and probably

has different targets in each of these cellular locations (see Introduction and Crozet *et al.*, 2014; Ruiz-

Gayosso *et al.*, 2018; Ramon *et al.*, 2019; Baena-González and Lunn, 2020). We introduced GFP-tagged

reporter constructs with a nuclear localization sequence (NUC) and without a nuclear localization

sequence (GEN), and confirmed that the former is localized to the nucleus, whereas the latter has a

more general location inside and outside the nucleus (Supplementary Figure S2). In many treatments,

including unperturbed diel cycles (Figure 1) and after transfer to extended darkness (Supplementary

Figure S3A), the phosphorylation state of the NUC and GEN peptides changed in a qualitatively similar

manner. However, in some perturbations they changed in a dissimilar or even opposite manner (see

Figures 2-3), presumably reflecting differing or even opposing changes in SnRK1 activity inside and

outside the nucleus, which might be due to differential regulation or to altered subcellular distribution

of SnRK1.

Diel changes in SnRK1 activity point to it operating in a network that buffers C status

It is well known that SnRK1 activity increases in stress and starvation conditions (Baena-Gonzalez *et*

al., 2007; Baena-Gonzalez and Sheen, 2008; Mair *et al.*, 2015; Cho et al., 2016; Pedrotti *et al.*, 2018;

Ramon *et al.*, 2019). However, it has emerged that SnRK1 also plays a role in benign conditions (Ramon

et al., 2019; Baena-González and Lunn, 2020; Peixoto *et al.*, 2021). After establishing that our *in vivo*

assay captures the expected increase in SnRK1 activity in C-depleted conditions, we focused on

monitoring SnRK1 activity in conditions where C availability is changing but without stressful extremes

of C starvation or surplus. To do this, we grew Arabidopsis in a 12-h photoperiod with 160 μmol m^{-2} s^{-1}

^{1} irradiance as a standard condition in which growth is restricted by the C supply, but not strongly

(Sulpice *et al.*, 2014).

One major finding was that *in-vivo* SnRK1 activity was relatively high in the light period, fell

after ZT12 (coinciding with darkening in our standard growth conditions) to a minimum at ZT16 and

then rose until the end of the night (Figure 1). Activity during the unperturbed diel cycle was several

times lower than in extended darkness or other treatments that led to C depletion (see Figure 4, Figure

6, Supplementary Fig 3A). Nevertheless, it was unexpected to observe higher SnRK1 activity in the light

period than during the first part of the night, because photosynthesis provides far more C in the light

than can be provided by mobilizing C reserves like starch in the night. Plants face a daily challenge to

balance the utilization of newly fixed C for immediate growth with the need to lay down C reserves

(e.g. starch) and to use these prudently to survive through the night. Therefore, plants must carefully

regulate the rate of starch accumulation in the light period, the rate of starch mobilization at night,

and the rate at which C is used for growth at different times in the diel cycle (Ishihara *et al.*, 2015;

Ivakov *et al.*, 2017; Mengin *et al.*, 2017). The diel pattern of SnRK1 activity that we observed

presumably reflects coordinated regulation of C utilization for reserve formation and growth

throughout the diel cycle. Incidentally, the relatively low SnRK1 activity during the night adds to the

evidence that C homeostasis is maintained across the diel light-dark cycle. In agreement, sucrose levels

remained relatively high for much of the night.

A second unexpected finding was that SnRK1 activity did not increase in response to a

decrease in overall C availability in the light. Illumination from dawn onwards at a lower irradiance

than growth irradiance did not lead to an increase in the phosphorylation status of either the NUC or

GEN reporter polypeptides, compared to that at growth irradiance (Figures 2 and 3). Although

photosynthesis was not measured in these experiments, the low irradiance treatments did lead to a

consistent decrease in the levels of sucrose and reducing sugars (Figures 2B and 3B; Supplementary

Dataset S1 – Exp.3-4; see also Moraes e*t al*., 2019). Further, these changes in irradiance occurred in

the range where Arabidopsis photosynthesis is light-limited (see Borghi *et al.*, 2019), and previous

studies have shown that similar drops in irradiance lead to a large decrease in photosynthetic rate

(see Fernandez *et al*., 2017; Moraes *et al*., 2019). On the other hand, treatments that led to a

decreased dusk starch content and slower rate of starch mobilization during the night did lead to a

consistent increase *in-vivo* SnRK1 activity in the first part of the night.

These observations point to SnRK1 being regulated in a more flexible manner in the light than

in the dark. This flexibility is also revealed by the response when plants are transferred to low

irradiance for one light period and then returned to growth irradiance. This perturbation led to higher

levels of sugars and faster accumulation of starch when plants were returned to growth irradiance,

compared to control plants that had been left at growth irradiance (Figure 2B). These observations

indicate that low irradiance results in a restriction of growth on the following day after returning plants

to previous growth irradiance (see also Moraes et al., 2019). SnRK1 activity showed a complex pattern

after returning to the previous growth irradiance (Figure 2); phosphorylation of NUC was lower than

in control plants, reflecting the higher levels of sugars, whereas phosphorylation of GEN resembled

that in control plants implying that extra-nuclear SnRK1 activity was higher than in control plants

despite sugar levels also being higher. It is known that growth in the light period is restricted when

Arabidopsis experiences a shortfall in C in the preceding 24-h cycle (Gibon et al., 2004; Sulpice *et al*.,

2014; Flis et al., 2016; Mengin *et al*., 2017; Moraes *et al*., 2019). Our observations indicate that

increased extranuclear SnRK1 activity might contribute to this restriction of growth and the resulting

increase in starch accumulation.

There may also be more ways for the plant to cope with a lower rate of photosynthesis in the

light than there are to adjust to a shortfall of starch at night. For example, the impact of decreased

irradiance on Glc6P levels in the light period was less marked than the impact on sugar levels

(Supplementary Figure S4, see also Moraes et al., 2019). This may reflect compensatory responses

that buffer the level of this key metabolic intermediate in the light, for example, by decreasing starch

synthesis, and by decreasing consumption of hexose phosphates for sucrose synthesis in the cytosol

by allosteric and post-transitional inhibition of sucrose-phosphate synthase (Huber and Huber, 1996;

Smith and Stitt, 2007; Lunn, 2016).

A third finding was that an endogenous circadian rhythm may contribute to the diel changes

in SnRK1 activity, at least in the middle of the 24-h cycle, when SnRK1 activity decreased between

about ZT12 and TT16. The decrease after ZT12 was partly retained in low continuous light (Figure 3)

and was also observed in plants growing in a T28 cycle, even though they were in the light between

ZT12 and ZT14 (Figure 4). The parallel changes in sugars and predicted extra-nuclear SnRK1 activity

after returning plants from low to growth irradiance may also point to an endogenous signal that

relays information about C status in the previous 24-h cycle.

SnRK1 activity may be regulated by multiple metabolic inputs including Tre6P and hexose phosphates

The unexpected diel response pointed to SnRK1 operating within a network that controls C utilization

in the daytime and at night to maintain C homeostasis. SnRK1 activity integrates various inputs

including multiple signals from metabolism (Polge and Thomas, 2007; Baena- González and Sheen,

2008; Wurzinger *et al.*, 2018). We asked to what extent diel changes in SnRK1 activity are accompanied

by reciprocal changes in the levels of sugars that might be viewed as indicators of C availability, or

reciprocal changes in the levels of metabolites, like Tre6P, Glc6P and Glc1P, that are known to inhibit

SnRK1 *in vitro* (Toroser *et al.*, 2000; Zhang *et al.*, 2009; Debast *et al.*, 2011; Delatte *et al.*, 2011; Piattoni

et al., 2011; Nunes *et al.*, 2013a; Coello and Martínez-Barajas, 2014; Emanuelle *et al.*, 2015; Griffiths

et al., 2016; Bledsoe *et al.*, 2017; Henry *et al.*, 2020; Baena-González and Lunn, 2020).

The role of Tre6P in regulation of SnRK1 has attracted wide attention (Zhang *et al.*, 2009;
Nunes *et al.*, 2013; Henry *et al.*, 2015; Zhang *et al.*, 2015; Bledsoe *et al.*, 2017; Baena-González and
Lunn, 2020). To directly test whether an induced increase in Tre6P leads to inhibition of *in-vivo* SnRK1
activity, we crossed our reporter constructs into lines carrying an inducible bacterial TPS construct. An
induced 1.4- to 2-fold increase in Tre6P in the light period indeed led to 12-23% decrease in
phosphorylation of the NUC reporter polypeptide (Figure 5), and a weak decrease for the GEN reporter
polypeptide (Supplementary Figure S6A). On the other hand, during diel cycles in wild-type plants
Tre6P levels did not show a consistent negative relationship to NUC or GEN phosphorylation (Figure
1; Table 1). Whereas Tre6P levels rose 3- to 5-fold during the light period, peaked at ZT12 (ED) and
then fell, (Figure 1B), SnRK1 activity did not change during the light period and, after darkening, fell
by 2- to 3-fold to a minimum at around ZT16 (Figure 1A). Also, comparison of the magnitude of the
changes in Figure 5 and Figure 1 indicates that the diel changes of Tre6P are too small to explain the
diel changes in NUC phosphorylation. In the last part of the night, where there was a negative
correlation between Tre6P levels and SnRK1 activity (Supplementary Table S1), an induced increase in
Tre6P resulted in an increase rather than a decrease in SnRK1 activity (Figure 6). Whilst these findings
provide evidence that Tre6P can contribute to the regulation of SnRK1 *in vivo*, they also show that the
contribution of Tre6P is context-dependent and that Tre6P is not the only or even the major factor
that regulates SnRK1 activity.

The possible role of hexose phosphates like Glc6P and Glc1P in the *in-vivo* regulation of SnRK1
activity has attracted less attention, despite evidence from *in-vitro* assays that they can inhibit SnRK1
(Toroser et al., 2000; Zhang *et al.*, 2009; Piattoni *et al.*, 2011; Nunes *et al.*, 2013). Several features of
our study point to them playing a role *in vivo*. First, visual inspection of diel changes under control
conditions reveal that Glc6P and Glc1P often change in a reciprocal manner to SnRK1 activity (Figure
1), peaking in the first part of the night at ZT16 at the same time as the minimum in SnRK1
phosphorylation activity. A negative relationship between SnRK1 and Glc6P was also observed when
plants were exposed to a sudden low-light day (Figure 2; see also Moraes *et al.*, 2019) or transferred
to continuous low light (Figure 3; Supplementary Figure S4). Second, there was a significant negative
correlation between Glc6P or Glc1P levels and NUC or GEN phosphorylation in some, though not all,
full diel cycles (Table 1) and highly consistent negative correlations in the night (Supplementary Table
S1). This conclusion was supported by comparison of the rates of change of Glc6P or Glc1P levels and
SnRK1 activity at different times in the diel cycle (Supplementary Table S3). Glc6P and Glc1P are linked
by the near-equilibrium reaction catalysed by phosphoglucomutase, with both changing in parallel
and Glc6P being about 10-fold higher than Glc1P, therefore, the observed negative relationship
between SnRK1 and Glc1P might be secondary to the relationship with Glc6P.

The question also arises what endogenous factor might be responsible for the decline in SnRK1

activity between ZT12 and ZT16, which occurred even when plants were not darkened at ZT12 (see

above). One possible explanation is that this decline is driven by diel changes in starch turnover. As

shown in Fernandez *et al.* (2017) and Ishihara *et al.* (2022), starch mobilization occurs simultaneously

with starch synthesis in the light and speeds up with time in the light, leading from about ZT10

onwards to a marked rise in sucrose and related metabolites, including Tre6P. This might contribute

to the decline in SnRK1 activity after ZT12 in continuous light. As suggested in Fernandez *et al*. (2017)

and Ishihara *et al*. (2022), this gradual rise in the rate of starch mobilization is itself under circadian

control. However, it would be premature to exclude the possibility that the circadian clock exerts a

more direct influence on SnRK1 activity, mirroring the recent discovery that SnRK1 is involved in

transcriptional regulation of the circadian clock (Jeong *et al.*, 2015; Sánchez-villarreal *et al.*, 2018; Shin

et al., 2017; Frank *et al.*, 2018). A possible link between starch mobilization and SnRK1 activity is also

of interest in view of the report that the dark-expressed SnRK1β3 subunit binds maltose, pointing to

a possible role in regulating starch degradation or of starch degradation, via maltose, regulating SnRK1

activity (Ruiz-Gayosso *et al.*, 2018).

Taken together, our results point to both Tre6P and hexose phosphates contributing to the

regulation of SnRK1 *in vivo*. These metabolites provide complementary information about C status.

Tre6P provides information about the amount of sucrose (Lunn *et al.*, 2006; Figueroa *et al.*, 2016;

Figueroa and Lunn, 2016; Fichtner *et al.*, 2017; Fichtner and Lunn, 2021), which is a key long-distance

transport metabolite and often a C-storage molecule (Lunn, 2016). Hexose phosphate levels reflect

the balance between several central metabolic pathways, for example, between photosynthetic C

fixation and sucrose and starch synthesis in the light, and between starch mobilization, respiration

and biosynthesis in the dark. Further, Tre6P and hexose phosphates may have differing subcellular

and intercellular compartmentation. TPS1 protein, and by implication Tre6P, is partly localized to the

nucleus (Fichtner *et al.*, 2020) whereas hexose phosphates are likely to have a more general

subcellular distribution with high levels in the cytosol (Stitt *et al.*, 1980; Gerhardt *et al.*, 1987). Such

differences in metabolite compartmentation might underlie some of the differing responses of the

NUC and GEN reporters. Further, TPS1, and by implication Tre6P, is very abundant in companion cells

and surrounding phloem parenchyma (Fichtner *et al.*, 2020) whereas hexose phosphates are probably

distributed more generally across cell types, including high levels in photosynthetic mesophyll cells.

This may introduce important nuances into regulation of SnRK1 activity, whose detection would

require cell-specific expression of the reporter constructs.

Changes in complex composition might impact on diel SnRK1 activity

663 An additional regulatory mechanism that might contribute to observed diel changes in SnRK1 activity

664 would be changes in subunit composition, including expression or stability of its subunits, such as by

665 SUMOylation and ubiquitination of SnRK1α1, SnRK1β1, SnRK1β2 SnRK1β3, and by myristoylation of

666 SnRK1β1 and SnRK1β2 (Ruiz-Gayosso *et al.*, 2018; Ramon *et al.*, 2019; Blanco *et al.*, 2019; Wang et al.,

667 2020). These were shown to affect cellular localization, primarily by translocating SnRK1 in and out of

668 the nucleus. We cannot yet be certain whether changes in SnRK1 composition contribute to the diel

669 changes in SnRK1 activity detected using the ACC reporter constructs. In the future it would be of

670 interest to compare changes in SnRK1 composition and SnRK1 location with the relative rates of NUC

671 and GEN phosphorylation.

673 *Comparison of* in-vivo *SnRK1 activity with responses of SnRK1 downstream targets highlights potential*

674 *flexibility in downstream signalling pathways*

675 As already mentioned, downstream target genes identified in protoplast overexpression studies have

676 been widely used to make inferences about SnRK1 activity *in vivo*. We inspected published data to

677 learn if the diel changes in transcript abundance of these genes is consistent with the diel changes of

678 *in-vivo* SnRK1 activity, as monitored by reporter peptide phosphorylation. To do this, we focused on

679 studies in which Arabidopsis was grown in the same conditions as those used in the current study. As

680 shown in Supplementary Figure S9A, which replots data from Usadel *et al.* (2008), there was a fairly

681 consistent pattern with the abundance of SnRK1-induced transcripts falling in the light period and

682 rising during the night, and the abundance of SnRK1-repressed transcripts rising in the light and falling

683 during the night (two exceptions were *AXP* and *EXP10*). This pattern contrasts with the response of

684 the NUC/GEN reporter polypeptide phosphorylation, which was relatively high in the light period and

685 fell in the first part of the night (Figure 1A). In a second comparison, we inspected the response of a

686 set of SnRK1 target transcripts after a sudden decrease in irradiance (Moraes *et al.*, 2019), similar to

687 the treatment in Figure 2 and the first 12 h after transfer to continuous low light in Figure 3

688 (Supplementary Figure S9B). Transfer from 160 to 60 µmol m^{-2} s^{-1} led, for SnRK1-induced genes, to a

689 small but consistent increase in *DIN6* and *BCAT2* and a large increase in *DIN1* transcript abundance

690 and, for SnRK1-repressed genes, to a small but consistent decrease in *CPN60α* and *Asp.P* transcript

691 abundance. This contrasts with NUC/GEN reporter polypeptide phosphorylation, which did not change

692 after transfer to low irradiance (Figures 2-3). In a recent study, Peixoto *et al*. (2021) reported that

693 abundance of SnRK1-induced transcripts was maximal at the end of the night when Tre6P levels were

694 lowest, and minimal at the end of the day when Tre6P levels peaked. However, even though the

695 changes of Tre6P levels in Peixoto *et al*. (2021) resembled those in the current study, the responses of

696 transcript abundance did not match the changes in NUC/GEN reporter polypeptide phosphorylation

reported in the current study. Furthermore, whilst transient elevation of Tre6P in an inducible

bacterial TPS line led to decreased abundance of transcripts for three SnRK1-induced genes (*BGAL,*

DIN10, PYL5) (Peixoto *et al*., 2021), consistent with involvement of Tre6P in diel regulation of SnRK1

signalling, it did not decrease transcript abundance for many other SnRK1-induced genes, such as

DIN1, *DIN6*, *AXP* and *EXP10*.

There are various possible explanations for discrepancies between SnRK1 activity as

monitored by phosphorylation of the NUC/GEN reporter polypeptides and SnRK1 activity as inferred

from the abundance of downstream transcripts. It is possible that in some cases gene expression is

driven by a fraction of the total SnRK1 pool, and that this fraction may remain undetectable when

assaying total SnRK1 activities (Baena-González and Lunn, 2020). The discrepancy may also reflect

multiple levels of regulation, for example, not only via changes in SnRK1 activity itself but also via

inputs that modify the affinity of SnRK1 for different immediate downstream targets or via inputs that

potentiate or attenuate different downstream signalling pathways. Such modifying factors might also

explain the varying responses of different SnRK1 downstream transcripts within a given experiment

(see above, also Griffiths *et al*., 2016), as well as between different studies. Indeed, of the top SnRK1

target genes, only a few changed in a reproducible manner following SnRK1 induction in protoplasts

versus whole plants (Baena-González *et al.*, 2007; Ramon *et al.*, 2019; Peixoto *et al.*, 2021). For

example, *DIN1*, *DIN6* and *CAT1*, whose expression is commonly used as a readout of SnRK1 activity,

are strongly upregulated in response to transient expression of SnRK1α1 in protoplasts (Baena-

González *et al.*, 2007), but remain unaffected in response to constitutive overexpression of SnRK1α1

in whole plants (Peixoto *et al.*, 2021).

Functionally, operation of SnRK1 in a benign diel cycle as well as during starvation mirrors the

differing responses of two sub-sets of C-regulated genes. Many of these show large changes in

expression during a regular equinoctial diel cycle (Smith and Stitt, 2007; Usadel *et al*., 2008; Cookson

et al., 2016, Flis et al., 2016). These genes may contribute to modifications of metabolism and slowing

down of growth, and serve to maintain C homeostasis and avoid transient periods of C starvation and

the resulting wasteful alternation between growth and catabolism (Ishihara et al., 2015, 2017). Other

C-regulated genes do not show changes in expression until starch is exhausted, and may serve to

orchestrate catabolic responses in conditions when the plant can no longer avoid C starvation.

In conclusion, by employing an *in-vivo* reporter polypeptide assay we have uncovered

unexpected changes in SnRK1 activity under benign light-dark cycles. Continued SnRK1 activity in the

light period may contribute to maintenance of diel sugar homeostasis, by restricting C utilization for

growth and supporting build-up of C reserves. Further, whilst demonstrating that Tre6P can inhibit

SnRK1 activity *in vivo*, we also show that its contribution is context dependent and that hexose

phosphates, and probably also other factors, interact to regulate SnRK1 activity and signalling.

Materials and Methods

Plant material

All Arabidopsis (*Arabidopsis thaliana* [L.] Heynh.) genotypes were in the Columbia-0 (Col-0)

background. Lines expressing the NUC or GEN SnRK1 reporters were generated as described in Sanagi

et al. (2021). The reporters comprise a 57-amino acid peptide derived from rat (*Rattus norvegicus*)

acetyl CoA carboxylase (rACC1), containing the AMPK recognition motif and Ser79 phosphorylation

site, fused at the C-terminus to eGFP and a double haemagglutinin tag (Supplementary Figure S1;

Deroover *et al.*, 2016). The NUC reporter includes an N-terminal SV40 nuclear localization signal that

targets the reporter specifically to the nucleus, while the GEN reporter has no added nuclear

localization signal and is localized in both the nucleus and cytoplasm (Supplementary Figure S2). The

Arabidopsis (Col-0) ethanol-inducible TPS-overexpressor (iTPS 29.2) and empty-vector control AlcR

lines were the same as those described in Martins *et al.*, 2013. The Arabidopsis (Col-0) 35S:SnRK1α1

over-expression line (Jossier *et al.*, 2009) and the 35S:SnRK1α1^{K48M} line, expressing the SnRK1α1

subunit with a mutation that disrupts ATP binding (Cho *et al.*, 2016), were kindly provided by Elena

Baena-González (Instituto Gulbenkian de Ciência, Oeiras, Portugal) and Hsing-Yi Cho (National Chung-

Hsing University, Academia Sinica, Taiwan), respectively.

Introgression of NUC and GEN SnRK1 reporters

The NUC and GEN SnRK1 reporters were introgressed into established lines with altered SnRK1 activity

or inducible expression of TPS by crossing. The respective homozygous parental lines were grown in

soil for 4-5 weeks (see below). Pollen from the parental parent was transferred by rubbing anthers

from mature flowers against exposed stigmas of emasculated maternal plants under a binocular

microscope, and the inflorescence meristems and unopened flower buds were removed. After

crossing, the maternal plants were grown on, with regular watering, for 2-3 weeks until siliques had

fully developed and started to dry. Siliques were collected, left to dry for a further 3-4 days at room

temperature and then kept at 10 °C until seeds were germinated on selective media. After surface

sterilization by treatment with 2% (w/v) sodium hypochlorite for 5 min at room temperature and

washing in five changes of water, seeds were placed on 0.5x Murashige and Skoog medium (Murashige

and Skoog, 1962) containing 1% (w/v) sucrose and appropriate antibiotics for selection of the

respective transgenes from each parent: NUC/GEN phosphinothricin (Basta); 35S:SnRK1α1,

kanamycin; 35S: SnRK1α1^{K48M}, hygromycin; iTPS/alcR; kanamycin. The seeds were stratified for 2 days

at 4°C in the dark and then transferred to 22°C in continuous light at 100 µmol m^{-2} s^{-1}. After 7 days,

surviving seedlings were transferred to soil. The F1 plants were selfed and F2 progeny were screened

on selective media, as above, to identify lines that exhibit the expected susceptible: resistant ratio of

7:9. Individual F2 plants were selfed and the F3 progeny were screened on selective media, as above,

to identify lines that were homozygous for both transgenic loci.

Plant growth and harvest

Seeds   were sown on a 1:1 mixture of compost (Stender AG, Schermbeck, Germany;

https://www.stender.de) and vermiculite in 6-cm diameter pots, covered and kept at 4°C in the dark

for 48 h then transferred to a controlled environment chamber (Percival E-36 L chamber, CLF Plant

Climatics GmbH, Weringen, Germany; https://www.plantclimatics.de/) with a 16-h photoperiod (160

µmol m^{-2} s^{-1} irradiance provided by white LEDs) and day/night temperatures of 21°C/19°C. After

germination, seedlings were transferred to 10-cm diameter pots (four or five seedlings per pot), and

grown as above or as described for individual experiments. Whole rosettes were harvested *in situ* and

immediately quenched in liquid nitrogen, with four or five plants from the same pot being pooled for

each biological replicate. Frozen plant material was ground to a fine powder and stored at -80°C until

use.

Metabolite extraction and analysis

Soluble sugars and starch were extracted and assayed enzymatically as described in Gibon *et al.* (2002)

with three-four biological replicates per sample type. For each biological replicate, values were

calculated as the average of two measurements. Tre6P, other phosphorylated intermediates and

organic acids were extracted with chloroform-methanol and quantified by anion-exchange high-

performance liquid chromatography coupled to tandem mass spectrometry as described in Lunn *et al.*

(2006), with modifications as described in Figueroa *et al*. (2016).

Ethanol-inducible iTPS experiments

After introgression of the NUC/GEN SnRK1 reporters, ethanol-inducible TPS (iTPS) and control alcR

lines (Martins *et al.*, 2013) were grown as described above. On the day of harvest, plants were sprayed

with either water (mock induction control) or with 2% (v/v) ethanol to induce expression of the

bacterial TPS and thereby increase Tre6P levels. Samples were collected and harvested as described

above.

SDS-PAGE and immunoblot assay of SnRK1 reporter phosphorylation

Proteins were extracted from aliquots (10 mg FW) of frozen, finely powdered plant material by

suspension in 100 µl of pre-warmed (90°C) sample buffer (66 mM Tris-HCl, pH 6.8, 10% (v/v) glycerol,

2% (w/v) sodium dodecyl sulphate, 0.13% (w/v) bromophenol blue) quickly vortex mixed, incubated

at 90°C for 5 min, then chilled on ice and centrifuged at 13,500 x *g* (4°C) for 3 min. Aliquots of the

supernatant containing approx. 12 µg of soluble protein were electrophoresed on a 12% (w/v)

polyacrylamide-SDS gel. Proteins were electro-blotted onto nitrocellulose membrane, visualized by

staining with 0.2% (w/v) Ponceau Red in 1% (v/v) acetic acid and imaged for use as a protein-loading

reference. After blocking in Tris-buffered saline containing 5% (w/v) milk powder, the membrane was

incubated overnight at 4°C with rabbit anti-phospho-acetyl CoA carboxylase (Ser79) antibody (#3661;

Cell Signaling Technology Europe BV, Leiden, The Netherlands; https://www.cellsignal.de) at a dilution

of 1:1500. After washing, NUC/GEN phosphorylation was detected by incubation with IRDye 800CW

goat anti-rabbit IgG (H+L) secondary antibody (Li-COR Biosciences GmbH, Bad Homburg, Germany;

www.licor.com) at a dilution of 1:9000 and scanning with an Odyssey Imaging Scanner (Li-COR), and

quantified using Image Studio Live 5.2 software (Li-COR). Total NUC/GEN polypeptide abundance was

detected by probing duplicate blots with a mouse anti-GFP antibody (#11814460001; Sigma Aldrich

(Roche), Darmstadt, Germany; www.sigmaaldrich.com) at a dilution of 1:1500, followed by incubation

with IRDye 680RD goat anti-mouse IgG (H+L) secondary antibody (Li-COR) at a dilution of 1:9000, and

quantified as above.

To minimize the impact of gel-to-gel variation, aliquots from an extended night sample were

loaded onto each gel and the signal from this sample was used to normalize the signals from the other

samples on the corresponding immunoblot. Two aliquots of the protein extract from each biological

replicate were analyzed on separate gels and, after normalization as described above, the average of

the two technical replicates was calculated as the reported value for that biological replicate. The

signal ratio pACC/GFP was used to normalize for any differences in reporter protein abundance in the

tissue and differences in gel loading. See Supplementary Figure S1 for further details.

Confocal microscopy

To determine the subcellular localization of the NUC/GEN SnRK1 reporter proteins, GFP was visualized

by laser confocal microscopy of 6-day-old seedlings. Imaging was performed on a Leica TCS SP5

microscope (Leica Microsystems, Wetzlar, Germany; https://www.leica-microsystems.com) using a

20x objective. To visualize nuclei, NUC samples were stained with 10 µg/ml propidium iodide (PI)

solution for 5 to 10 min. A 488 nm laser was used to excite GFP and 561 nm laser to excite PI and their

fluorescence emission was detected in the 524-555 nm and 571-640 nm ranges, respectively. Confocal

Z sections were acquired every 0.5 μm.

Statistical analysis

Technical replicates were always averaged to generate a single value for each biological replicate,

statistical analysis was restricted to biological replicates and performed using Sigma-Plot 14.5

software (Systat Software GmbH, Düsseldorf, Germany; http://www.systat.de). Significance of

changes in metabolite and SnRK1 activity was tested by one-way ANOVA using a pairwise multiple

comparison procedure, with post-hoc testing by the Holm-Sidak method ($p<0.05$), as described in each

figure legend.

Acknowledgements

We thank Sofie Deroover and Tom Broeckx for help with generation and selection of the NUC and GEN

SnRK1 reporter lines and Arun Sampathkumar for help with laser confocal microscopy.

Funding

This work was supported by the Max Planck Society (O.A., T.A.M., V.M., R.F., M.S., and J.E.L.) and by

KU Leuven (F.R.).

Author contributions: O.A. performed all experiments, statistically analysed data, and contributed to

paper writing; T.A.M contributed to experimental design, harvesting, measurement of metabolites

and data analysis. V.M. contributed to harvesting and measurement of metabolites, R.F. performed

LC-MS/MS metabolite analysis; F.R. conceived and generated the SnRK1 reporter constructs, M.S.

assisted data analysis and contributed to writing the paper. J.E.L. conceived the project, assisted data

analysis, and contributed to writing the paper.

Supplementary Data

Supplementary Figure S1: *In vivo* SnRK1 activity assay

Supplementary Figure S2: Localization of reporter peptides by confocal microscopy

Supplementary Figure S3: Validation of the NUC and GEN constructs in whole Arabidopsis rosettes

Supplementary Figure S4: Comparison of different low light treatments (supplementary to Figure 2 and 3)

Supplementary Figure S5: T28 and T24 cycles; diel Glc6P levels and comparison of SnRK1 activity excluding time points affected by C starvation (supplementary to Figure 4).

Supplementary Figure S6: Response of NUC and GEN phosphorylation in crosses with a line containing inducible bacterial TPS.

Supplementary Figure S7: Impact of transiently elevated Tre6P in the light period on the levels of selected metabolites (supplementary to Figure 5).

Supplementary Figure S8: Transient elevation of Tre6P at the end of the night slows down starch mobilization and leads to low glucose and high organic acid levels (supplementary to Figure 6)

Supplementary Figure S9: Transcript abundance for SnRK1 target genes during a diel cycle, after an extension of the night and after a sudden decrease in light intensity for one day.

Supplementary Table S1: Correlations between metabolite levels and NUC or GEN phosphorylation, performed separately for the light period and the night (supplementary to Table 1 and Figure 1).

Supplementary Table S2: Correlations between metabolite levels (supplementary to Table 1 and Figure 1)

Supplementary Table S3: Correlations between changes in metabolite levels and changes in NUC or GEN phosphorylation (supplementary to Table 1 and Figure 1).

Supplementary Table S4: Correlations between metabolite levels and NUC or GEN phosphorylation in growth regimes that differed from the equinoctial T24 cycle (supplementary to Figures 3 and 4).

Supplementary Table S5: Summary of correlation analysis between changes in SnRK1 NUC and GEN activity to changes in metabolite levels (supplementary to Figures 3 and 4).

Supplementary Table S6: Correlation analysis between metabolite levels and NUC phosphorylation following transient elevation of Tre6P in the light period (supplementary to Figure 5, Supplementary Dataset S2 Exp.7).

Supplementary Text S1: Correlations between in vivo SnRK1 activity and metabolites in continuous light, in T28 cycles and in experiments where TPS is induced (supplementary to Supplementary Tables S4-S6)

Supplementary Dataset S1: *In-vivo* SnRK1 activity and metabolite levels.

Supplementary Dataset S2: Regressions between *in-vivo* SnRK1 activity and metabolite levels.

References

Ávila-Castañeda A, Gutiérrez-Granados N, Ruiz-Gayosso A, Sosa-Peinado A, Martínez-Barajas E,

Coello P (2014) Structural and functional basis for starch binding in the SnRK1 subunits AKINβ2

and AKINβγ. Front Plant Sci **5:** 199

Baena-González E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription

networks in plant stress and energy signalling. Nature **448**: 938–942

Baena-González E, Lunn JE (2020) SnRK1 and trehalose 6-phosphate – two ancient pathways converge

to regulate plant metabolism and growth. Curr Opin Plant Biol **55**: 52–59

Belda-Palazón, B., Costa, M., Beeckman, T., Rolland, F., Baena-González, E. (2022) ABA represses TOR

and root meristem activity through nuclear exit of the SnRK1 kinase. Proc Natl Acad Sci USA **119:**

2204862119.

Blanco NE, Liebsch D, Díaz MG, Strand Å, Whelan J (2019) Dual and dynamic intracellular localization

of Arabidopsis thaliana SnRK1.1. J Exp Bot **70**: 2325–2338

Bledsoe SW, Henry C, Griffiths CA, Paul MJ, Feil R, Lunn JE, Stitt M, Lagrimini LM (2017) The role of

Tre6P and SnRK1 in maize early kernel development and events leading to stress-induced kernel

abortion. BMC Plant Biol **17**: 1–17

Borghi GL, Moraes TA, Günther M, Feil R, Mengin V, Lunn JE, Stitt M, Arrivault S (2019) Relationship

between irradiance and levels of Calvin–Benson cycle and other intermediates in the model

eudicot Arabidopsis and the model monocot rice. J Exp Bot **70:** 5809–5825

Broeckx T, Hulsmans S, Rolland F (2016) The plant energy sensor: evolutionary conservation and

divergence of SnRK1 structure, regulation, and function. J Exp Bot **67**, 6215–6252

Cabib E, Leloir LF (1958) The biosynthesis of trehalose phosphate. J Biol Chem **231:** 259-75

Cho H, Wen T, Wang Y, Shih M (2016) Quantitative phosphoproteomics of protein kinase SnRK1

regulated protein phosphorylation in Arabidopsis under submergence. J Exp Bot **67**: 2745–2760

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation

of *Arabidopsis thaliana*. Plant J **16**: 735–743

Coello P, Martínez-Barajas E (2014) The activity of SnRK1 is increased in *Phaseolus vulgaris* seeds in

response to a reduced nutrient supply. Front Plant Sci **5**: 1–7

Cookson S, Yadav UP, Klie S, Morcuende RM, Usadel B, Lunn JE, Stitt M (2016) Temporal kinetics of

the transcriptional response to carbon depletion and sucrose readdition in Arabidopsis seedlings.

Plant Cell Environ **39:** 768-786

Crepin N, Rolland F (2019a) SnRK1 activation , signaling , and networking for energy homeostasis.

Curr Opin Plant Biol **51:** 29–36

Crozet P, Margalha L, Confraria A, Rodrigues A, Martinho C, Adamo M, Elias CA, Baena-González E
(2014) Mechanisms of regulation of SNF1/AMPK/SnRK1 protein kinases. Front Plant Sci **5:** 190

Dale S, Wilson WA, Edelman AM, Hardie DG (1995) Similar substrate recognition motifs for
mammalian AMP-activated protein kinase, higher plant HMG-CoA reductase kinase-A, yeast
SNF1, and mammalian calmodulin-dependent protein kinase I. FEBS Lett **361**: 191–195

Debast S, Nunes-Nesi A, Hajirezaei MR, Hofmann J, Sonnewald U, Fernie AR, Börnke F (2011) Altering
trehalose-6-phosphate content in transgenic potato tubers affects tuber growth and alters
responsiveness to hormones during sprouting. Plant Physiol **156:** 1754-1771

Delatte TL, Sedijani P, Kondou Y, et al. (2011) Growth arrest by trehalose-6-phosphate : an
astonishing case of primary metabolite control over growth by way of the SnRK1 signaling
pathway. Plant Physiol **157**: 160–174

Delorge I, Figueroa CM, Feil R, Lunn JE, van Dijck P (2015) Trehalose-6-phosphate synthase 1 is not
the only active TPS in *Arabidopsis thaliana*. Biochem J **290:** 283–290

Deroover S, Ghillebert R, Broeckx T, Winderickx J, Rolland F (2016) Trehalose-6-phosphate synthesis
controls yeast gluconeogenesis downstream and independent of SNF1. FEMS Yeast Res **16:**36

Emanuelle S, Hossain MI, Moller IE, et al. (2015) SnRK1 from *Arabidopsis thaliana* is an atypical AMPK.
Plant J **82**: 183–192

Fernandez O, Ishihara H, George GM, et al. (2017) Leaf starch turnover occurs in long days and in
falling light at the end of the day. Plant Physiol **174**: 2199–2212

Fichtner F, Barbier FF, Feil R, Watanabe M, Annunziata MG, Stitt M, Beveridge CA, Lunn JE (2017)
Trehalose 6-phosphate is involved in triggering axillary bud outgrowth in garden pea (*Pisum*
sativum L .). Plant J **92:** 611–623

Fichtner F, Olas JJ, Feil R, Watanabe M, Krause U, Hoefgen R, Stitt M, Lunn JE (2020) Functional
features of TREHALOSE-6-PHOSPHATE SYNTHASE1, an essential enzyme in Arabidopsis. Plant Cell
32: 1949–1972

Fichtner F, Lunn JE (2021) The Role of trehalose 6-phosphate (Tre6P) in plant metabolism and
development. Annu Rev Plant Biol **72:** 737-760

Figueroa CM, Feil R, Ishihara H, et al. (2016) Trehalose 6-phosphate coordinates organic and amino
acid metabolism with carbon availability. Plant J **85**: 410–423

Figueroa CM, Lunn JE (2016) A tale of two sugars: trehalose 6-phosphate and sucrose. Plant Physiol
172: 7–27

Flis A, Sulpice R, Abel C, Ivakov A, Liput M, Millar AJ, Stitt M (2016) Photoperiod-dependent changes
in the phase of core clock transcripts, and in global transcriptional outputs at dusk and dawn in
Arabidopsis. Plant Cell Environ **39**: 1955-1981

Flis A, Mengin V, Ivakov AA, et al. (2019) Multiple circadian clock outputs regulate diel turnover of
carbon and nitrogen reserves. Plant Cell Environ **42**: 549–573

Frank A, Matiolli CC, Viana AJC, et al. (2018) Circadian entrainment in arabidopsis by the sugar-
responsive transcription factor bZIP63. Curr Biol **28:** 2597-2606

García-Salcedo R, Lubitz T, Beltran G, et al. (2014) Glucose de-repression by yeast AMP-activated
protein kinase SNF1 is controlled via at least two independent steps. FEBS J **281**: 1901–1917

Gerhardt R, Stitt M, Heldt HW (1987) Subcellular metabolite levels in spinach leaves : regulation of
sucrose synthesis during diurnal alterations in photosynthetic partitioning. Plant Physiol **83**: 399-
407

Gibon Y, Vigeolas H, Tiessen A, Geigenberger P, Stitt M (2002) Sensitive and high throughput
metabolite assays for inorganic pyrophosphate, ADPGlc, nucleotide phosphates, and glycolytic
intermediates based on a novel enzymic cycling system. Plant J **30**: 221–235

Gibon Y, Bläsing OE, Palacios-Rojas N, Pankovic D, Hendriks JHM, Fisahn J, Höhne M, Günther M,
Stitt M (2004) Adjustment of diurnal starch turnover to short days: depletion of sugar during the
night leads to a temporary inhibition of carbohydrate utilization, accumulation of sugars and
post-translational activation of ADP-glucose pyrophosphorylase in the following light period.
Plant J **39**: 847–862

Glab N, Oury C, Guérinier T, Domenichini S, Crozet P, Thomas M, Vidal J, Hodges M (2017) The impact
of *Arabidopsis thaliana* SNF1-related-kinase 1 (SnRK1)-activating kinase 1 (SnAK1) and SnAK2 on
SnRK1 phosphorylation status: characterization of a SnAK double mutant. Plant J **89:** 1031–1041

Glinski M, Weckwerth W (2005) Differential multisite phosphorylation of the trehalose-6-phosphate
synthase gene family in *Arabidopsis thaliana*: a mass spectrometry-based process for
multiparallel peptide library phosphorylation analysis. Mol Cell Proteomics **4**: 1614-1625

Goddijn OJM, van Dun K (1999) Trehalose metabolism in plants. Trends Plant Sci **4:** 315–319

Graf A, Schlereth A, Stitt M, Smith AM (2010) Circadian control of carbohydrate availability for growth
in Arabidopsis plants at night. Proc Natl Acad Sci USA **107**: 9458–9463

Griffiths CA, Sagar R, Geng Y, et al. (2016) Chemical intervention in plant sugar signalling increases
yield and resilience. Nature **540**: 574–578

Hara LEO, Paul MJm Wingler A (2013) How do sugars regulate plant growth and development? New
insight into the role of trehalose-6-phosphate. Mol Plant **6**, 261–274

Hardie DG, Carling D (1997) The AMP-activated protein kinase – fuel gauge of the mammalian cell?
Eur J Biochem **246**: 259–273

Hardie DG, Carling D, Carlson M (1998) The AMP-activated/SNF1 protein kinase subfamily: metabolic
sensors of the eukaryotic cell? Annu Rev Biochem **67**: 821–855

Harthill JE, Meek SEM, Morrice N, Peggie MW, Borch J, Wong BHC, Mackintosh C (2006)
Phosphorylation and 14-3-3 binding of Arabidopsis trehalose-phosphate synthase 5 in response
to 2-deoxyglucose. Plant J **47:** 211–223

Henninger M, Pedrotti L, Krischke M, Draken J, Wildenhain T, Fekete A, Rolland F, Müller MJ,
Fröschel C, Weiste C, Dröge-Laser W (2022) The evolutionarily conserved kinase SnRK1
orchestrates resource mobilization during Arabidopsis seedling establishment. Plant Cell **34**: 616-
632

Henry C, Bledsoe SW, Griffiths CA, Kollman A, Paul MJ, Sakr S, Lagrimini LM (2015) Differential role
for trehalose metabolism in salt-stressed maize. Plant Physiol **169**: 1072–1089

Henry C, Watson-Lazowski A, Oszvald M, Griffiths C, Paul MJ, Furbank RT, Ghannoum O (2020) Sugar
sensing responses to low and high light in leaves of the C_4 model grass *Setaria viridis*. J Exp Bot
71: 1039–1052

Herzig S, Shaw RJ (2017) AMPK: guardian of metabolism and mitochondrial homeostasis. Nat Rev Mol
Cell Biol **19:** 121-135

Huber SC, Huber JL (1996) Role and regulation of sucrose-phosphate synthase in higher plants. Annu
Rev Plant Physiol Plant Mol Biol **47**: 431–444

Hulsmans S, Rodriguez M, de Coninck B, Rolland F (2016) The SnRK1 energy sensor in plant biotic
interactions. Trends Plant Sci **21**: 648–661

Hwang G, Kim S, Cho J-Y, Paik I, Kim J-I, Oh E (2019) Trehalose-6-phosphate signaling regulates
thermoresponsive hypocotyl growth in *Arabidopsis thaliana.* EMBO Rep **20:** 47828

Ishihara H, Obata T, Sulpice R, Fernie AR, Stitt M (2015) Quantitative determination of the rates of
protein synthesis and turnover in Arabidopsis rosettes using dynamic $^{13}CO_2$ labelling and analysis
of isotope enrichment of individual amino acids in free pools and in protein. Plant Physiol **168:**
74-93

Ishihara H, Obata T, Sulpice R, Fernie AR, Stitt M (2015) Quantifying protein synthesis and
degradation in arabidopsis by dynamic $^{13}CO_2$ labeling and analysis of enrichment in individual
amino acids in their free pools and in protein. Plant Physiol **168:** 74–93

Ishihara H, Moraes T, Pyl E-T, Schulze WX, Obata T, Scheffel A, Fernie AR, Sulpice R, Stitt M (2017)
Growth rate correlates negatively with protein turnover in Arabidopsis accessions. Plant J **91**:
416-429

Ishihara H, Alseekh S, Feil R, et al. (2022) Rising rates of starch degradation with time in the light and
trehalose 6-phosphate levels optimize carbon availability and thereby buffer growth. Plant
Physiol **189**: 1976-2000

Ivakov A, Flis A, Apelt F, et al. (2017) Cellulose synthesis and cell expansion are regulated by different

mechanisms in growing arabidopsis hypocotyls. Plant Cell **29:** 1305-1315

Jeong EY, Seo PJ, Woo JC, Park CM (2015) AKIN10 delays flowering by inactivating IDD8 transcription

factor through protein phosphorylation in Arabidopsis. BMC Plant Biol **15**: 1–13

Jossier M, Bouly JP, Meimoun P, Arjmand A, Lessard P, Hawley S, Hardie GD, Thomas M (2009) SnRK1

(SNF1-related kinase 1) has a central role in sugar and ABA signalling in *Arabidopsis thaliana*.

Plant J **59**: 316–328

Kretzschmar T, Pelayo MAF, Trijatmiko KR, et al. (2015) A trehalose-6-phosphate phosphatase

enhances anaerobic germination tolerance in rice. Nat Plants **1:** 15124

Lastdrager J, Hanson J, Smeekens S (2014) Sugar signals and the control of plant growth and

development. J Exp Bot **65:** 799–807

Leyman B, van Dijck P, Thevelein JM (2001) An unexpected plethora of trehalose biosynthesis genes

in Arabidopsis thaliana. Trends Plant Sci **6:** 510–513

Locke JCW, Millar AJ, Turner MS (2005) Modelling genetic networks with noisy and varied

experimental data: the circadian clock in *Arabidopsis thaliana*. J Theor Biol **234**: 383–393

Lu C, Lin C, Lee K, Chen J, Huang L, Ho S, Liu H, Hsing Y, Yu S (2007) The SnRK1A protein kinase plays

a key role in sugar signaling during germination and seedling growth of rice. Plant Cell **19**: 2484–

2499

Lunn JE, Feil R, Hendriks JHMM, et al. (2006) Sugar-induced increases in trehalose 6-phosphate are

correlated with redox activation of ADPglucose pyrophosphorylase and higher rates of starch

synthesis in *Arabidopsis thaliana*. Biochem J **148**: 139–148

Lunn JE (2007) Gene families and evolution of trehalose metabolism in plants. Funct Plant Biol **34**:

550–563

Lunn JE, Delorge I, Mar C, van Dijck P, Stitt M (2014) Trehalose metabolism in plants. Plant J **4**: 544–

567

Lunn JE (2016) Sucrose Metabolism. Encyclopedia of Life Sciences (ELS). John Wiley Sons, 1–9.

Ma J, Hanssen M, Lundgren K, et al. (2011) The sucrose-regulated Arabidopsis transcription factor

bZIP11 reprograms metabolism and regulates trehalose metabolism. New Phytol **191**: 733-745

Mair A, Pedrotti L, Wurzinger B, et al. (2015) SnRK1-triggered switch of bZIP63 dimerization mediates

the low-energy response in plants. Elife **4**: 05828

Margalha L, Confraria A (2019) SnRK1 and TOR : modulating growth – defense trade-offs in plant

stress responses. J Exp Bot **70:** 2261–2274

Martínez-Barajas E, Delatte T, Schluepmann H, et al. (2011) Wheat grain development is

characterized by remarkable trehalose 6-phosphate accumulation pregrain filling: tissue

distribution and relationship to SNF1-related protein kinase1 activity. Plant Physiol **156**: 373–381

Martins MCM, Hejazi M, Fettke J, et al. (2013) Feedback inhibition of starch degradation in arabidopsis leaves mediated by trehalose 6-phosphate. Plant Physiol **163**: 1142–1163

Mayer F V, Heath R, Underwood E, et al. (2011) ADP regulates SNF1, the *Saccharomyces cerevisiae* homolog of AMP-activated protein kinase. Cell Metab **14**: 707–714

Mengin V, Pyl ET, Moraes TA, Sulpice R, Krohn N, Encke B, Stitt M (2017) Photosynthate partitioning to starch in *Arabidopsis thaliana* is insensitive to light intensity but sensitive to photoperiod due to a restriction on growth in the light in short photoperiods. Plant Cell Environ **40**: 2608–2627

Millar AJ, Carré IA, Strayer CA, Chua NH, Kay SA (1995) Circadian clock mutants in arabidopsis identified by luciferase imaging. Science **267**: 1161–1163

Moraes TA, Mengin V, Annunziata MG, Encke B, Krohn N, Höhne M, Stitt M (2019) Response of the circadian clock and diel starch turnover to one day of low light or low CO_2. Plant Physiol **179**: 1457–1478

Muralidhara P, Weiste C, Collani S, et al. (2021) Perturbations in plant energy homeostasis prime lateral root initiation via SnRK1-bZIP63-ARF19 signalling. Proc Natl Acad Sci USA **118**: 210696111.

Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco tissue cultures. Physiol. Plant **15**: 473–497

Nietzsche M, Landgraf R, Tohge T, Börnke F (2016) A protein-protein interaction network linking the energy-sensor kinase SnRK1 to multiple signaling pathways in *Arabidopsis thaliana*. Curr Opin Plant Biol **5**: 36–44

Nietzsche M, Schießl I, Börnke F, Griffiths CA (2014) The complex becomes more complex : protein-protein interactions of SnRK1 with DUF581 family proteins provide a framework for cell- and stimulus type-specific SnRK1 signaling in plants. Front Plant Sci **5**: 1–13

Nuccio ML, Wu J, Mowers R, et al. (2015) Expression of trehalose-6-phosphate phosphatase in maize ears improves yield in well-watered and drought conditions. Nat Biotech **33**: 862–869

Nukarinen E, Nägele T, Pedrotti L, Wurzinger B, Baena-González E, Dröge-Laser W, Weckwerth W (2016) Quantitative phosphoproteomics reveals the role of the AMPK plant ortholog SnRK1 as a metabolic master regulator under energy deprivation. Sci Rep **6:** 31697

Nunes C, Hara LEO, Primavesi LF, et al. (2013) The trehalose 6-phosphate/SnRK1 signaling pathway primes growth recovery following relief of sink limitation. Plant Physiol **162:** 1720–1732

Nunes C, Primavesi LF, Patel MK, Martinez-Barajas E, Powers SJ, Sagar R, Fevereiro PS, Davis BG, Paul MJ (2013) Inhibition of SnRK1 by metabolites: Tissue-dependent effects and cooperative inhibition by glucose 1-phosphate in combination with trehalose 6-phosphate. Plant Physiol Biochem **63**: 89–98

Oszvald M, Primavesi LF, Grif A, Cohn J, Basu S, Nuccio ML, Paul MJ (2018) Trehalose 6- phosphate

regulates photosynthesis and assimilate partitioning in reproductive tissue. Plant Physiol **176:**

2623–2638

Paul MJ, Gonzalez-Uriarte A, Grif CA, Hassani-Pak K (2018) The role of trehalose 6-phosphate in crop

yield and resilience. Plant Physiol **177:** 12–23

Paul MJ, Watson A, Griffiths CA (2020) Trehalose 6-phosphate signalling and impact on crop yield.

Biochem Soc Trans **48**: 2127-2137

Pedrotti L, Weiste C, Nägele T, et al. (2018) Snf1-RELATED KINASE1-controlled C / S 1 -bZIP signaling

activates alternative mitochondrial metabolic pathways to ensure plant survival in extended

darkness. Plant Cell **30**: 495–509

Peixoto B, Moraes TA, Mengin V, et al. (2021) Impact of the SnRK1 protein kinase on sucrose

homeostasis and the transcriptome during the diel cycle. Plant Physiol **187**: 1357–1373

Piattoni CV, Bustos DM, Guerrero SA, Iglesias AÁ (2011) Nonphosphorylating glyceraldehyde-3-

phosphate dehydrogenase is phosphorylated in wheat endosperm at serine-404 by an SNF1-

related protein kinase. Plant Physiol **156**: 1337–1350

Polge C, Jossier M, Crozet P, Gissot L, Thomas M (2008) β-Subunits of the SnRK1 complexes share a

common ancestral function together with expression and function specificities; physical

interaction with nitrate reductase specifically occurs via AKINβ1-subunit. Plant Physiol **148**: 1570-

1582

Polge C, Thomas M (2006) SNF1 / AMPK / SnRK1 kinases, global regulators at the heart of energy

control? Trends Plant Sci **12**: 20–28

Ponnu J, Wahl V, Schmid M (2011) Trehalose-6-phosphate: Connecting plant metabolism and

development. Front Plant Sci **2:** 70

Ramon M, Dang TVT, Broeckx T, Hulsmans S, Crepin N, Sheen J, Rolland F (2019) Default activation

and nuclear translocation of the plant cellular energy sensor SnRK1 regulate metabolic stress

responses and development. Plant Cell **31:** 1614–1632

Randall RS (2021) The plant AlcR-pAlcA ethanol-inducible system displays gross growth artefacts

independently of downstream pAlcA-regulated inducible constructs. Sci Rep **11:** 1–5

Ryabova LA, Robaglia C, Meyer C (2019) Target of Rapamycin kinase: central regulatory hub for plant

growth and metabolism. J Exp Bot **70:** 2211-2216

Rodrigues A, Adamo M, Crozet P, et al. (2013) ABI1 and PP2CA phosphatases are negative regulators

of Snf1-related protein kinase1 signaling in Arabidopsis. Plant Cell **25:** 3871–3884

Ruiz-Gayosso A, Rodríguez-Sotres R, Martínez-Barajas E, Coello P (2018) A role for the carbohydrate-

binding module (CBM) in regulatory SnRK1 subunits: the effect of maltose on SnRK1 activity.

Plant J **96:** 163–175

Sanagi M, Aoyama S, Kubo A, et al. (2021) Low nitrogen conditions accelerate flowering by
modulating the phosphorylation state of FLOWERING BHLH 4 in Arabidopsis. Proc Natl Acad Sci
USA **118**: 2022942118

Sánchez-Villarreal A, Davis AM, Davis SJ (2018) AKIN10 activity as a cellular link between metabolism
and circadian-clock entrainment in *Arabidopsis thaliana*. Plant Signal Behav **13**: 1411448

Schluepmann H, Pellny T, van Dijken A, Smeekens S, Paul M (2003) Trehalose 6-phosphate is
indispensable for carbohydrate utilization and growth in *Arabidopsis thaliana*. Proc Natl Acad Sci.
USA **100**: 6849–6854

Schluepmann H, Paul M (2009) Trehalose metabolites in Arabidopsis—elusive, active and central.
Arabidopsis Book **7:** 0122

Seki M, Ohara T, Hearn TJ, Frank A, da Silva VCH, Caldana C, Webb AAR, Satake A (2017) Adjustment
of the Arabidopsis circadian oscillator by sugar signalling dictates the regulation of starch
metabolism. Sci Rep **7**: 8305

Sheen J, Baena-González E (2008) Convergent energy and stress signaling. Trends Plant Sci **13**: 474-
482

Shin J, Sánchez-Villarreal A, Davis AM, Du SX, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ (2017)
The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent
manner. Plant Cell Environ **40:** 997–1008

Smith AM, Stitt M (2007) Coordination of carbon supply and plant growth. Plant Cell Environ **30**:
1126–1149

Stitt M, Wirtz W, Heldt HW (1980) Metabolite levels during induction in the chloroplast and
extrachloroplast compartments of spinach protoplasts. Biochim Biophys Acta Bioenerg **593**: 85–
102

Sulpice R, Flis A, Ivakov AA, et al. (2014) Arabidopsis coordinates the diurnal regulation of carbon
allocation and growth across a wide range of photoperiods. Mol Plant **7**: 137–155

Toroser D, Plaut Z, Huber SC (2000) Regulation of a plant SNF1-related protein kinase by glucose-6-
phosphate. Plant Physiol **123:** 403–411

Usadel B, Bläsing OE, Gibon Y, Retzlaff K, Höhne M, Günther M, Stitt M (2008) Global transcript levels
respond to small changes of the carbon status during progressive exhaustion of carbohydrates
in arabidopsis rosettes. Plant Physiol **146**: 1834–1861

Vandesteene L, Ramon M, Le K, van Dijck P, Rolland F (2010) A single active trehalose-6-P synthase
(TPS) and a family of putative regulatory TPS-like proteins in Arabidopsis. Mol Plant **3**: 406–419

Vandesteene L, López-Galvis L, Vanneste K, et al. (2012) Expansive evolution of the *TREHALOSE-6-*
PHOSPHATE PHOSPHATASE gene family in Arabidopsis. Plant Physiol **160**: 884–896

Wahl V, Ponnu J, Schlereth A, et al. (2013) Regulation of flowering by trehalose-6-phosphate signaling

in *Arabidopsis thaliana*. Science **339**: 704–707

Wang Y, Wang L, Micallef BJ, Tetlow IJ, Mullen RT, Feil R, Lunn JE, Emes MJ (2020) AKINβ1, a subunit

of SnRK1, regulates organic acid metabolism and acts as a global modulator of genes involved in

carbon, lipid, and nitrogen metabolism. J Exp Bot **71**: 1010–1028

Wingler A, Delatte TL, Hara LEO, Primavesi LF, Jhurreea D, Paul MJ, Schluepmann H (2012) Trehalose

6-phosphate is required for the onset of leaf senescence associated with high carbon availability.

Plant Physiol **158:** 1241–1251

Wurzinger B, Nukarinen E, Nägele T, Weckwerth W, Teige M (2018) The SnRK1 kinase as central

mediator of energy signaling between different organelles. Plant Physiol **176:** 1085–1094

Yang HL, Liu YJ, Wang CL, Zeng QY (2012) Molecular evolution of trehalose-6-phosphate synthase

(TPS) gene family in Populus, Arabidopsis and rice. PLoS One **7**: 42438

Zacharaki V, Ponnu J, Crepin N, Langenecker T, Hagmann J, Skorzinski N, Musialak-Lange M, Wahl

V, Rolland F, Schmid M (2022) Impaired KIN10 function restores developmental defects in the

Arabidopsis *trehalose 6-phosphate synthase1* (*tps1*) mutant. New Phytol **235**: 220-233

Zhai Z, Liu H, Shanklin J (2017) Phosphorylation of WRINKLED1 by KIN10 results in its proteasomal

degradation, providing a link between energy homeostasis and lipid biosynthesis. Plant Cell **29**,

871–889.

Zhai Z, Keereetaweep J, Liu H, Feil R, Lunn JE, Shanklin J (2018) Trehalose 6- phosphate positively

regulates fatty acid synthesis by stabilizing WRINKLED1. Plant Cell **30:** 2616–2627

Zhang Y, Primavesi LF, Jhurreea D, et al. (2009) Inhibition of SNF1- related protein kinase1 activity

and regulation of metabolic pathways by trehalose-6-phosphate. Plant Physiol **149**: 1860–1871

Zhang ZP, Deng Y, Song X, Miao M (2015) Trehalose-6-phosphate and SNF1-related protein kinase 1

are involved in the first-fruit inhibition of cucumber. J Plant Physiol **177**: 110–120

1187

1188   **Figure legends**

1189

Figure 1: Reproducible diel response in phosphorylation of SnRK1 reporters under equinoctial

growth conditions. SnRK1 reporter lines were grown in a 12-h photoperiod (irradiance 160 µmol m^{-2}

s^{-1}) for 22 days as controls in three independent experiments, in which additional plants grown in

parallel were subjected to different treatments (see Figs 2-4). Whole rosettes were harvested at 4-h

intervals through the diel cycle. (A) Phosphorylation of the NUC/GEN polypeptide was quantified by

immunoblotting as described in Methods and in Supplementary Figure S1. (B) Selected metabolites.

Data are from the control samples from the following experiments: (i) Control 1: Supplementary

Dataset S1 – Exp.3 (blue symbols; NUC and GEN lines); (ii) Control 2: Supplementary Dataset S1 – Exp.4

and Supplementary Figure S4 (red symbols; NUC and GEN lines); (iii) Control 3: Supplementary Dataset

S1 – Exp.2 and Supplementary Figure S3b (grey symbols; NUC line only). Data are shown as mean ±

S.D. (n = 3-4 biological replicates, each containing 3-5 pooled rosettes). Statistical analyses are

provided for each time series in Figs 2, 3 and 4, respectively.

1202

Figure 2: SnRK1 activity in Arabidopsis plants transferred to low irradiance for one light period. NUC

and GEN lines were grown in a 12-h photoperiod at 160 µmol m^{-2} s^{-1} irradiance for 19 days. At 20 DAS,

starting at dawn (ZT0), irradiance was reduced to 60 µmol m^{-2} s^{-1} for the following 12 h (treatment

light period), plants were darkened at ZT12 (treatment night), and at ZT24 were re-illuminated at an

irradiance of 160 µmol m^{-2} s^{-1} (recovery day), as described in Moraes *et al*. (2019). Samples from both

genotypes and from the corresponding control treatment (plants left in the original growth light

regime) were harvested at 4-h intervals from ZT0-ZT36 (with respect to dawn on the treatment day).

(A) Soluble sugars and starch. (B) Phosphorylation of NUC (top) and GEN (bottom) SnRK1 reporter

polypeptides. The dashed grey background denotes the low-light treatment day. Results are shown as

mean ± S.D. (n=3-4 biological replicates). Statistical analysis: letters indicate significant (*P*<0.05)

changes between different times for a given genotype and asterisks indicate significant (*P*<0.05)

differences between lines at a given time point, according to one-way ANOVA and pairwise multiple

comparison post-testing using the Holm-Sidak method. The original and additional data are available

in Supplementary Dataset S1 – Exp.3. The control time-series is also shown in Figure 1 (blue symbols).

1217

Figure 3: SnRK1 activity in Arabidopsis plants transferred to continuous low light. NUC and GEN lines

were grown in a 12-h photoperiod at 160 µmol m^{-2} s^{-1} irradiance for 19 days. At 20 DAS, plants were

shifted at dawn (ZT0) to continuous low light (90 µmol m^{-2} s^{-1}) for 32 h, while corresponding controls

were kept in the original growth conditions (12-h photoperiod, 160 µmol m^{-2} s^{-1}). (A) Diel changes of

sugars and starch. (B) Phosphorylation of NUC (top) and GEN (bottom) SnRK1 reporter polypeptides.

The dashed grey background represents the subjective night. Results are shown as mean ± S.D. (n = 3

biological replicates). Statistical analysis: letters indicate significant (P<0.05) changes between

different times for a given treatment and asterisks represent significant (P<0.05) differences between

the low light treatment and the control at a given time point according to one-way ANOVA with

pairwise multiple comparison post-testing using the Holm-Sidak method. The original and additional

data are available in Supplementary Dataset S1 – Exp.4. The control time-series is also shown in Figure

1 (red symbols).

1230

Figure 4: SnRK1 activity in Arabidopsis plants grown in T24 and T28 light-dark cycles. NUC and GEN

lines were grown from sowing in a 14-h light / 14 h dark (T28) cycle under 160 µmol m^{-2} s^{-1} irradiance

for 19 days, in parallel with control plants that were grown in a 12-h light / 12-h dark (T24) cycle. At

20 DAS (as defined by the T24 cycle), starting at dawn, plants were harvested at 4-h intervals

throughout the light-dark cycle (open symbols). At the end of the light-dark cycle, additional plants

were left in darkness and harvested for two further time points for the plants grown in a T24 cycle (i.e.

at ZT28 and ZT32), and for one further time point for plants grown in a T28 cycle (i.e. at ZT32). (A)

Soluble sugars and starch. (B) Phosphorylation of NUC (top) and GEN (bottom) SnRK1 reporter

polypeptides. The white background represents light in both T cycles and the grey shaded areas

represent darkness (ZT12-ZT24 for the T24 cycle and ZT14-ZT28 for the T28 cycle) in one or both T

cycles: pale grey denotes darkness in the T24 and light in the T28 cycle, mid-grey denotes darkness in

both T-cycles, and dark grey represents extended night in both T cycles. Black closed symbols

represent time points in extended darkness. Results are shown as mean ± S.D. (n = 3 biological

replicates). Statistical analysis: letters indicate significant (P<0.05) changes between different times

for a given genotype and asterisks represent significant (P<0.05) differences between T28 vs T24 at a

given time point according to a one-way ANOVA with pairwise multiple comparison post-testing using

the Holm-Sidak method. The original and additional data are available in Supplementary Dataset S1-

Exp.5. The control time-series is also shown in Figure 1 (grey symbols).

1249

Figure 5: Elevated Tre6P in the light period inhibits nuclear SnRK1 activity. Line #6.3 (NUC x iTPS) was

grown in a 12-h photoperiod at 160 µmol m^{-2} s^{-1} irradiance for 20 days. At 21 DAS, plants were sprayed

at ZT4 (red arrow) with 2% (v/v) ethanol to induce the expression of bacterial TPS (otsA) or with water

as a mock-induction control, and then harvested at ZT7, ZT9 and ZT11 for measurement of Tre6P (top

panel), sucrose (middle) and the phosphorylation status of the NUC polypeptide (bottom) as a readout

of nuclear SnRK1 activity *in vivo*. Results are shown as mean ± S.D. (n = 3-4 biological replicates).

Statistical analysis: asterisks indicate significant ($P<0.05$) differences between the ethanol-induced

plants and water controls according to a one-way ANOVA with pairwise multiple comparison post-

testing using the Holm-Sidak method. Additional metabolites are shown in Supplementary Figure S7.

The original data are available in Supplementary Dataset S1 – Exp.7.

1260

Figure 6: Transiently high Tre6P at the end of the night leads to an increase in nuclear SnRK1 activity.

Line #6.3 (NUC x iTPS) and #1.2 control (NUC x alcR) were grown in short-day conditions (6 h light / 18

h dark) at 160 µmol m^{-2} s^{-1} irradiance for 25 days. At 26 DAS, plants were sprayed with 2% (v/v) ethanol

at ZT20 (red arrow) to induce the expression of bacterial TPS (otsA). Rosettes were harvested at ZT20,

ZT22 and ZT24 (end of normal night), and at ZT25 and ZT27 (extended darkness)for measurement of:

Tre6P (top panel) and the phosphorylation of the NUC polypeptide (bottom) as a readout of nuclear

SnRK1 activity *in vivo*. Results are shown as mean ± S.D. (n=3-4 biological replicates). Statistical

analysis: letters indicate significant ($P<0.05$) changes between different times for a given genotype

and asterisks indicate significant ($P<0.05$) differences between the #6.3 (NUC x iTPS) and #1.2 control

(NUC x alcR) lines according to a one-way ANOVA with pairwise multiple comparison post-testing using

the Holm-Sidak method. Additional metabolites are shown in Supplementary Figure S8. The original

data are available in Supplementary Dataset S1 – Exp.7.

1273

Table 1: Correlation between metabolite levels and NUC or GEN phosphorylation. Presented are R^2
and p-values of linear regression, performed separately for the three independent control
experiments compiled in Figure 1. Linear regression analysis was performed with the "least squares"
method, using individual replicates for the complete diel cycle (ZT0 to ZT24). The column +/- provides
information about the slope of the regression: positive (+), negative (-). Significant correlations ($p <$
0.05) are highlighted in blue (positive) or red (negative). n.d. not determined. Individual plots are
provided in Supplementary Dataset S2.

Experiment	Metabolite	NUC			GEN		
		R^2	p-value	+/-	R^2	p-value	+/-
1	Tre6P	0.007	0.673	+	0.001	0.965	+
	Glc6P	0.048	0.270	-	**0.225**	**0.030**	-
	Glc1P	0.056	0.236	-	0.036	0.412	-
	Starch	0.006	0.706	+	0.051	0.324	-
	Sucrose	0.085	0.139	-	0.058	0.295	-
	Glucose	0.121	0.076	+	**0.272**	**0.015**	+
2	Tre6P	0.004	0.778	+	0.046	0.325	-
	Glc6P	**0.215**	**0.022**	-	**0.583**	**2.24×10^{-5}**	-
	Glc1P	**0.418**	**0.001**	-	**0.683**	**1.17×10^{-6}**	-
	Starch	0.081	0.190	-	0.081	0.189	-
	Sucrose	1×10^{-8}	0.685	-	**0.324**	**0.005**	-
	Glucose	0.054	0.687	-	2×10^{-5}	0.985	-
3	Tre6P	0.004	0.233	-	n.d.	n.d.	
	Glc6P	0.002	0.766	-	n.d.	n.d.	
	Glc1P	0.02	0.373	+	n.d.	n.d.	
	Starch	0.032	0.364	-	n.d.	n.d.	
	Sucrose	0.007	0.678	+	n.d.	n.d.	
	Glucose	0.013	0.561	-	n.d.	n.d.	

1281

A.

B.

Figure 1: Reproducible diel response in phosphorylation of SnRK1 reporters under equinoctial growth conditions. SnRK1 reporter lines were grown in a 12-h photoperiod (irradiance 160 µmol m^{-2} s^{-1}) for 22 days as controls in three independent experiments, in which additional plants grown in parallel were subjected to different treatments (see Figs 2-4). Whole rosettes were harvested at 4-h intervals through the diel cycle. (A) Phosphorylation of the NUC/GEN polypeptide was quantified by immunoblotting as described in Methods and in Supplementary Figure S1. (B) Selected metabolites. Data are from the control samples from the following experiments: (i) Control 1: Supplementary Dataset S1 – Exp.3 (blue symbols; NUC and GEN lines); (ii) Control 2: Supplementary Dataset S1 – Exp.4 and Supplementary Figure S4 (red symbols; NUC and GEN lines); (iii) Control 3: Supplementary Dataset S1 – Exp.2 and Supplementary Figure S3b (grey symbols; NUC line only). Data are shown as mean ± S.D. (n = 3-4 biological replicates, each containing 3-5 pooled rosettes). Statistical analyses are provided for each time series in Figs 2, 3 and 4, respectively.

Figure 2: SnRK1 activity in Arabidopsis plants transferred to low irradiance for one light period. NUC and GEN lines were grown in a 12-h photoperiod at 160 µmol m^{-2} s^{-1} irradiance for 19 days. At 20 DAS, starting at dawn (ZT0), irradiance was reduced to 60 µmol m^{-2} s^{-1} for the following 12 h (treatment light period), plants were darkened at ZT12 (treatment night), and at ZT24 were re-illuminated at an irradiance of 160 µmol m^{-2} s^{-1} (recovery day), as described in Moraes *et al.* (2019). Samples from both genotypes and from the corresponding control treatment (plants left in the original growth light regime) were harvested at 4-h intervals from ZT0-ZT36 (with respect to dawn on the treatment day). (A) Soluble sugars and starch. (B) Phosphorylation of NUC (top) and GEN (bottom) SnRK1 reporter polypeptides. The dashed grey background denotes the low-light treatment day. Results are shown as mean ± S.D. (n=3-4 biological replicates). Statistical analysis: letters indicate significant (*P*<0.05) changes between different times for a given genotype and asterisks indicate significant (*P*<0.05) differences between lines at a given time point, according to one-way ANOVA and pairwise multiple comparison post-testing using the Holm-Sidak method. The original and additional data are available in Supplementary Dataset S1 – Exp.3. The control time-series is also shown in Figure 1 (blue symbols).

Figure 3: SnRK1 activity in Arabidopsis plants transferred to continuous low light. NUC and GEN lines were grown in a 12-h photoperiod at 160 µmol m^{-2} s^{-1} irradiance for 19 days. At 20 DAS, plants were shifted at dawn (ZT0) to continuous low light (90 µmol m^{-2} s^{-1}) for 32 h, while corresponding controls were kept in the original growth conditions (12-h photoperiod, 160 µmol m^{-2} s^{-1}). (A) Diel changes of sugars and starch. (B) Phosphorylation of NUC (top) and GEN (bottom) SnRK1 reporter polypeptides. The dashed grey background represents the subjective night. Results are shown as mean ± S.D. (n = 3 biological replicates). Statistical analysis: letters indicate significant (*P*<0.05) changes between different times for a given treatment and asterisks represent significant (*P*<0.05) differences between the low light treatment and the control at a given time point according to one-way ANOVA with pairwise multiple comparison post-testing using the Holm-Sidak method. The original and additional data are available in Supplementary Dataset S1 – Exp.4. The control time-series is also shown in Figure 1 (red symbols).

Figure 4: SnRK1 activity in Arabidopsis plants grown in T24 and T28 light-dark cycles. NUC and GEN lines were grown from sowing in a 14-h light / 14 h dark (T28) cycle under 160 µmol m⁻² s⁻¹ irradiance for 19 days, in parallel with control plants that were grown in a 12-h light / 12-h dark (T24) cycle. At 20 DAS (as defined by the T24 cycle), starting at dawn, plants were harvested at 4-h intervals throughout the light-dark cycle (open symbols). At the end of the light-dark cycle, additional plants were left in darkness and harvested for two further time points for the plants grown in a T24 cycle (i.e. at ZT28 and ZT32), and for one further time point for plants grown in a T28 cycle (i.e. at ZT32). (A) Soluble sugars and starch. (B) Phosphorylation of NUC (top) and GEN (bottom) SnRK1 reporter polypeptides. The white background represents light in both T cycles and the grey shaded areas represent darkness (ZT12-ZT24 for the T24 cycle and ZT14-ZT28 for the T28 cycle) in one or both T cycles: pale grey denotes darkness in the T24 and light in the T28 cycle, mid-grey denotes darkness in both T-cycles, and dark grey represents extended night in both T cycles. Black closed symbols represent time points in extended darkness. Results are shown as mean ± S.D. (n = 3 biological replicates). Statistical analysis: letters indicate significant (*P*<0.05) changes between different times for a given genotype and asterisks represent significant (*P*<0.05) differences between T28 vs T24 at a given time point according to a one-way ANOVA with pairwise multiple comparison post-testing using the Holm-Sidak method. The original and additional data are available in Supplementary Dataset S1- Exp.5. The control time-series is also shown in Figure 1 (grey symbols).

Figure 5: Elevated Tre6P in the light period inhibits nuclear SnRK1 activity. Line #6.3 (NUC x iTPS) was grown in a 12-h photoperiod at 160 μmol m^{-2} s^{-1} irradiance for 20 days. At 21 DAS, plants were sprayed at ZT4 (red arrow) with 2% (v/v) ethanol to induce the expression of bacterial TPS (otsA) or with water as a mock-induction control, and then harvested at ZT7, ZT9 and ZT11 for measurement of Tre6P (top panel), sucrose (middle) and the phosphorylation status of the NUC polypeptide (bottom) as a readout of nuclear SnRK1 activity *in vivo*. Results are shown as mean ± S.D. (n = 3-4 biological replicates). Statistical analysis: asterisks indicate significant (*P*<0.05) differences between the ethanol-induced plants and water controls according to a one-way ANOVA with pairwise multiple comparison post-testing using the Holm-Sidak method. Additional metabolites are shown in Supplementary Figure S7. The original data are available in Supplementary Dataset S1 – Exp.7.

Figure 6: Transiently high Tre6P at the end of the night leads to an increase in nuclear SnRK1 activity. Line #6.3 (NUC x iTPS) and #1.2 control (NUC x alcR) were grown in short-day conditions (6 h light / 18 h dark) at 160 µmol m^{-2} s^{-1} irradiance for 25 days. At 26 DAS, plants were sprayed with 2% (v/v) ethanol at ZT20 (red arrow) to induce the expression of bacterial TPS (otsA). Rosettes were harvested at ZT20, ZT22 and ZT24 (end of normal night), and at ZT25 and ZT27 (extended darkness)for measurement of: Tre6P (top panel) and the phosphorylation of the NUC polypeptide (bottom) as a readout of nuclear SnRK1 activity *in vivo*. Results are shown as mean ± S.D. (n=3-4 biological replicates). Statistical analysis: letters indicate significant (*P*<0.05) changes between different times for a given genotype and asterisks indicate significant (*P*<0.05) differences between the #6.3 (NUC x iTPS) and #1.2 control (NUC x alcR) lines according to a one-way ANOVA with pairwise multiple comparison post-testing using the Holm-Sidak method. Additional metabolites are shown in Supplementary Figure S8. The original data are available in Supplementary Dataset S1 – Exp.7.