SAGA histone acetyltransferase module facilitates chromatin accessibility to SMC5/6

Mahrik L. [1,2], Stefanovie B. [1,2], Maresova A. [3], Princova J. [3], Kolesar P. [1], Lelkes E. [1],

Helmlinger D. [4], Prevorovsky M. [3*], Palecek J.J. [1,2*]

[1] National Centre for Biomolecular Research, Faculty of Science, Masaryk University,

Kotlarska 2, 61137 Brno, Czech Republic

[2] Mendel Centre for Plant Genomics and Proteomics, Central European Institute of

Technology, Masaryk University, Kamenice 5, 62500 Brno, Czech Republic

[3] Department of Cell Biology, Faculty of Science, Charles University, Vinicna 7, 12800

Prague, Czech Republic

[4] Centre de Recherche en Biologie cellulaire de Montpellier, University of Montpellier,

CNRS, 1919 route de Mende, 34293 Montpellier Cedex 05, France

*Corresponding addresses: jpalecek@sci.muni.cz; National Centre for Biomolecular

Research, Faculty of Science, Masaryk University, Kamenice 5, 62500 Brno, Czech Republic,

Tel.: +420-54949-6128, FAX: +420-54949-6351;

prevorov@natur.cuni.cz, Department of Cell Biology, Faculty of Science, Charles University,

Vinicna 7, 12800 Prague, Czech Republic, Tel.: +420-22195-1769, FAX: +420-22195-1758

Abstract

Structural Maintenance of Chromosomes (SMC) complexes are molecular machines driving chromatin organization at higher levels. In eukaryotes, three SMC complexes (cohesin, condensin, and SMC5/6) play key roles in cohesion, condensation, replication, transcription and DNA repair. Here, we performed a genetic screen in fission yeast to identify novel factors required for SMC5/6 binding to DNA. We identified 79 genes of which histone acetyltransferases (HATs) were the most represented. Genetic and phenotypic analyses suggested a particularly strong functional relationship between the SMC5/6 and SAGA complexes. Furthermore, several SMC5/6 subunits physically interacted with SAGA HAT module components Gcn5 and Ada2. As Gcn5-dependent acetylation facilitates the accessibility of chromatin to DNA repair proteins, we first analysed the formation of DNA damage-induced SMC5/6 foci in the Δ*gcn5* mutant. The SMC5/6 foci formed normally in Δ*gcn5,* suggesting SAGA-independent SMC5/6 localization to DNA-damaged sites. Next, we used Nse4-FLAG chromatin-immunoprecipitation (ChIP-seq) analysis in unchallenged cells to assess SMC5/6 distribution. A significant portion of SMC5/6 accumulated within gene regions in wild-type cells, which was reduced in Δ*gcn5* and Δ*ada2* mutants. The drop in SMC5/6 levels was also observed in *gcn5*-E191Q acetyltransferase-dead mutant. Altogether, our data suggest that the SAGA HAT module may facilitate chromatin accessibility to SMC5/6 at gene regions.

Author Summary

Genomes of all eukaryotes must be folded and packed into their relatively small nuclear spaces. Histones first pack free genomic DNA into nucleosomes and their arrays. Other complexes like histone modifiers and remodelers can regulate nucleosome positions and their packing within chromatin fibres. They assist in the relative opening or condensation of

chromatin fibres and facilitate their accessibility to DNA-binding proteins. The highly

conserved Structural Maintenance of Chromosomes (SMC) complexes (cohesin, condensin,

and SMC5/6) compact further chromatin fibres at higher levels. These molecular machines

can loop chromatin fibres, which need access to segments of free DNA for their physical

binding to DNA. Here, we studied genetic and physical interactions between histone-

modifying SAGA complex and SMC5/6. We show that the SAGA histone acetyltransferase

module may facilitate chromatin access to SMC5/6.

Introduction

Chromatin is composed of DNA and protein complexes structured at multiple levels to ensure its spatial and functional organization [1]. Histone proteins pack DNA into nucleosomes and their arrays at the basic level. At the higher levels, Structural Maintenance of Chromosomes (SMC) complexes (cohesin, condensin and SMC5/6) assist in the formation of higher-order structures like topologically associated domains or condensed mitotic chromosomes [2]. Chromatin compaction affects DNA accessibility at each level. Histone chaperones, modifiers, and remodelers can loosen, move or remodel nucleosomes to modulate essential processes like transcription or DNA repair [3, 4]. For example, the histone-modifying SAGA complex acetylates H3 histones at promoter regions, contributing to chromatin opening and facilitating the assembly of transcription initiation complexes onto core promoters and the recruitment of factors that directly interact with DNA [5].

The SMC complexes play roles in all key chromatin processes, including cohesion, condensation, replication, transcription, and DNA repair. Their cores comprise the long-armed SMC, kleisin, and kleisin-associated (KITE or HAWK) subunits [6, 7]. Uniquely, SMC5/6 complexes contain the highly conserved Nse1 and Nse2 ubiquitin- and SUMO-ligases, respectively [8, 9]. The Smc subunits are primarily built of head ATPase domains, long anti-parallel coiled-coil arms, and hinges [10-12]. Two Smc molecules form stable dimers via their hinge domains, and without ATP, their arms align into rod-like structures [13, 14]. The binding of ATP molecules to the ATPase head domains promotes the formation of large annular structures [15, 16]. The ATP binding-hydrolysis cycle drives ring-to-rod dynamic changes and promotes DNA translocation or loop extrusion [17-20].

SMC complexes were believed to interact only topologically with chromatin fibres via their large annular structures, which can embrace and traverse large chromatin complexes,

including nucleosomes [12]. However, growing evidence suggests their requirement for open

chromatin and direct physical binding to DNA [21-23]. Moreover, Piazza et al. [22] described

the preferential binding of condensin kleisin-associated HAWK subunits to free DNA over

nucleosomal DNA. Recent cryoEM analyses showed the formation of K-compartments within

all SMC complexes, consisting of ATP-bound Smc heads, kleisin, and kleisin-associated

subunits, which can accommodate only free DNA [16, 24-26]. In line with these findings, the

SAGA complex assists in loading condensin at open chromatin regions of highly transcribed

genes in fission yeast [23]. Similarly, the RSC chromatin remodelling complex recruits the

Scc2-Scc4 factor to nucleosome-free regions, assisting in cohesin loading at these sites [27-

29].

Recently, it was shown that the SMC5/6 K-compartment, composed of Smc5-Smc6

heads, Nse4 kleisin, and Nse1-Nse3 kleisin-associated KITE subunits, binds free DNA [16,

21]. In our previous study, we described the essential role of Nse3-DNA interaction for

SMC5/6 loading or accumulation [21]. Here, we performed a genetic screen with a *nse3-*

R254E fission yeast mutant that exhibits reduced DNA-binding affinity to identify new

factors required for its viability. We found strong genetic interactions with the SAGA and

NuA4 histone acetyltransferase (HAT) complexes. Using chromatin-immunoprecipitation

(ChIP-seq) analysis, we observed a significant portion of SMC5/6 accumulated within gene

regions, which was reduced in SAGA HAT deletion (Δ*ada2* and Δ*gcn5*) mutants. The

magnitude of the decrease in SMC5/6 occupancy correlated with the SAGA-modified

H3K9ac levels around the transcription start sites. The SMC5/6 reduced levels were also

observed in *gcn5-E191Q* acetyltransferase-dead mutant, suggesting that the SAGA HAT

module may facilitate the accessibility of chromatin for SMC5/6 loading.

Results

Genetic screen with a DNA-binding defective allele of the SMC5/6 complex

To identify factors that facilitate the loading or accumulation of the SMC5/6 complex

on chromatin, we performed a genetic search for genes affecting the survival of cells with

compromised DNA-binding ability of the SMC5/6 complex [21]. First, we created a query

fission yeast strain with the DNA-binding defective *nse3-R254E* mutation in the PEM2

background (Suppl. Fig. S1A; [30]) and crossed it against the whole gene deletion yeast

collection from BIONEER [31]. Using yeast colony size phenotypic readout [32], we

identified 79 deletion strains that exhibited a negative genetic interaction with *nse3-R254E*

(Suppl. Table ST1).

To validate our results, we randomly selected 19 of the 79 strains and crossed them

with the original *nse3-R254E* strain [21]. Tetrad analysis confirmed that these mutations are

synthetically sick or lethal with the *nse3-R254E* mutation (Suppl Fig. S1B). Analysis of all 79

interacting genes for the Biological process category using Gene ontologies (GO; [33])

showed the highest scores for DNA repair, chromatin organization, meiosis, and replication

processes (Fig. 1A and Suppl Table ST2), in line with previous studies (reviewed in [34, 35]).

Reassuringly, several *nse3-R254E* genetic interactions overlapped with the genetic

interactions of other *smc5/6* mutants [21, 36-41], supporting the validity of our screen results.

Genetic interactions between SMC5/6 and histone acetyltransferase complexes

The analysis of the Cellular components GO of the 79 interacting genes showed the

highest scores for histone acetyltransferase (HAT) complexes (SAGA and NuA4; Fig. 1B and

Suppl Table ST3). Three (Ada2, Ada3/Ngg1, Gcn5) out of four HAT module subunits and

one (Ubp8) out of four DUB module subunits of the SAGA complex were identified as hits in

our screen [5]. We verified their genetic interactions using tetrad analysis (Figs. 1C and S1B)

and also tested the other non-essential SAGA subunits not detected in our screen. The mating

defects of the Δ*ada1,* Δ*spt7,* Δ*spt8,* and Δ*spt20* deletion mutants hampered the double mutant

preparation [42]. However, the tetrad analysis of the other non-essential SAGA subunits

(Suppl. Fig. S1C) showed negative genetic interactions with the *nse3-R254E* mutation (Suppl.

Fig. S1D), suggesting a strong functional relationship between the SMC5/6 and SAGA

complexes.

Interestingly, the temperature-sensitive (*ts*) phenotypes of the SAGA HAT module

Δ*ada2*, Δ*ada3*, and Δ*gcn5* mutants were enhanced by *nse3-R254E* (Fig. 1D; [42]). The *ts*

phenotypes were also enhanced by the *smc6-74* and *smc6-X* hypomorphic mutations (Suppl.

Fig. S2A; [37, 43]). We also used *nse1-R188E* and *nse2-SA* mutations, which specifically

abrogate DNA-repair function (ubiquitin- and SUMO- ligase activity, respectively), but not

the SMC5/6 essential function [8, 44]. However, these mutations did not affect the *ts*

phenotype of Δ*gcn5* (Suppl. Fig. S2A; not shown). These data suggest that the SMC5/6

essential function supports cell survival at higher temperatures in the absence of the HAT

module and, conversely, that the viability of the *nse3-R254E* mutant is compromised in the

absence of a functional SAGA HAT module.

SMC5/6 and SAGA physically interact

The strong genetic relationship between the SAGA HAT module and the SMC5/6

complex prompted us to test their mutual physical interactions. First, we performed co-

immunoprecipitation of Gcn5-myc and Nse4-FLAG kleisin subunit to show the association

between the SAGA and SMC5/6 complexes *in vivo* (Fig. 2A). The fission yeast cells carrying

Nse4-FLAG (with or without Gcn5-myc) were lysed, and proteins were precipitated with an

anti-myc antibody. A small amount of Nse4-FLAG was specifically recovered in the Gcn5-

myc precipitates but not in the control experiment without myc-tagged Gcn5, suggesting a

weak or transient association between SAGA and SMC5/6 in the yeast cells.

Next, we used our panel of yeast two-hybrid (Y2H) SMC5/6 subunits to test their

interactions with the HAT module (Ada2, Ada3, Gcn5, and Sgf29) subunits. Two of the HAT

module subunits, Ada2 and Gcn5, bound the Nse3 subunit (Figs. 2B and S2B). The fragment

analysis showed that Gcn5 bound the N-terminal part of Nse3(aa1-190), while Ada2

interacted with its C-terminal WHB domain (Nse3(aa200-307); Figs. 2B and S2C; [45]). In

addition, Ada2 bound the N-terminal region of Nse2 and the coiled-coil arm of SMC5 next to

the SMC5-Nse2 binding interface [46]. Altogether, our data suggest several physical

interactions underlying a functional relationship between SAGA and SMC5/6.

SMC5/6 and SAGA are required for efficient DNA repair

The SAGA complex plays an important role in transcription regulation and DNA

repair by facilitating the accessibility of chromatin to transcription factors and repair proteins

[42, 47-50]. Deletion mutants of most SAGA genes were sensitive to hydroxyurea (HU; [42])

and SAGA mutations increased the sensitivity of the *smc5/6* hypomorphic mutants to HU and

other DNA-damaging agents (Figs. 3A, S2A, and S3A). These data suggest either a direct role

of the SAGA complex in facilitating SMC5/6 access to chromatin at sites of DNA damage or

its indirect, independent role.

Upon DNA damage, SMC5/6 accumulates in foci in a Brc1-dependent way [51, 52].

To assess whether SMC5/6 chromatin accessibility at these DNA damage sites is directly

facilitated by SAGA, we analysed the Nse4-GFP foci in the Δ*gcn5* and Δ*ada2* mutants (Fig.

3B). We observed no difference between the frequency of cells with foci in the wild-type

(WT), Δ*gcn5,* and Δ*ada2* after treatment with either MMS (methyl methane sulfonate) or HU.

In contrast, the number of foci in the Δ*brc1* mutant was strongly reduced (Fig. 3C). These

results indicate no direct involvement of SAGA in SMC5/6 localization to sites of DNA

damage. Instead, they suggest that the observed HU phenotype results from additive effects of

the SAGA and SMC5/6 complexes during DNA damage repair.

SAGA facilitates chromatin accessibility to SMC5/6 in gene regions

In addition to the accessibility of chromatin to DNA repair and transcription factors, SAGA facilitates chromatin accessibility to condensin complexes [23]. Therefore, we determined the SMC5/6 localization in the unchallenged WT, Δ*gcn5*, Δ*ada2*, and Δ*ubp8* cells using chromatin immunoprecipitation of Nse4-FLAG followed by deep sequencing (ChIP-seq). In the WT cells, most SMC5/6 localized to the repetitive regions (like rDNA, centromeres, or tDNA copies), with the highest occupancy at the rDNA repeats (Fig. 4A) consistent with previous reports [53-58]. Interestingly, we identified 331 Nse4-FLAG peaks (representing ¼ of the total Nse4 occupancy; Fig. 4A) that localized to gene regions.

In Δ*gcn5* and Δ*ada2* mutants, SMC5/6 distribution was altered, while SMC5/6 occupancy in the Δ*ubp8* mutant was similar to that in WT cells. A heatmap clustering analysis showed that the Nse4-FLAG peaks were either enhanced, unchanged, or reduced in the Δ*gcn5* and Δ*ada2* mutants (Fig. 4B). The peaks were mainly enhanced at the repetitive sequences (Fig. 4C), while most peaks in the intergenic regions were not changed (Fig. 4D). Strikingly, most peaks within the gene regions showed reduced Nse4 occupancy (252 out of 331; Figs. 4C and D), suggesting that the SAGA HAT module may facilitate chromatin accessibility to SMC5/6 at gene loci.

The SAGA HAT module acetylates histone H3 at its lysine K9 and K14 residues [59, 60]. To determine the H3K9ac distribution in our fission yeast cells, we performed ChIP-seq using an anti-H3K9ac antibody. A heatmap clustering analysis showed that the magnitude of decrease in SMC5/6 occupancy in gene bodies in the Δ*gcn5* mutant (Δ*gcn5*-WT plot; Figs. 5A and B) correlated with H3K9ac levels around the transcription start site and with transcript levels [61]. It suggests that the SAGA-dependent H3K9 acetylation may facilitate the accessibility of chromatin to SMC5/6 at gene regions.

To assess the role of the SAGA-dependent acetylation further, we used the *gcn5-*

E191Q acetyltransferase-dead mutant [50]. First, we found that the *gcn5-E191Q ts* phenotype

was exacerbated by *nse3-R254E,* similar to Δ*gcn5* (Suppl. Fig. S3D). Second, the *gcn5-*

E191Q nse3-R254E double mutant was as sensitive to DNA-damaging agents as the Δ*gcn5*

nse3-R254E double mutant. It suggests that the Gcn5 acetyltransferase activity plays a crucial

role in the SAGA-SMC5/6 genetic interactions. Finally, using ChIP-qPCR analysis, we

observed reduced Nse4-FLAG occupancy in the *gcn5-E191Q* mutant at selected gene regions

similar to Δ*gcn5* (Fig. 5C), although some loci exhibited only a modest drop in SMC5/6

levels. Altogether, our results suggest an important role for the SAGA HAT module in

facilitating the accessibility of chromatin to SMC5/6 at gene loci.

Discussion

Here, we discovered the SAGA HAT module as a new chromatin factor assisting in the

localization of SMC5/6 to specific genomic loci. In line with SAGA's best-known function in

transcription [5], Δ*gcn5* deletion reduced SMC5/6 occupancy specifically at the gene loci

(Fig. 4). In addition, the magnitude of decrease in SMC5/6 occupancy in gene bodies

correlated with the H3K9 acetylation and transcript levels (Fig. 5). These data suggest that the

SAGA-dependent H3K9 acetylation may facilitate the accessibility of chromatin to SMC5/6.

Consistent with this notion, SMC5/6 occupancy was also reduced in the *gcn5-E191Q*

acetyltransferase-dead mutant. The fact that occupancy reduction was less pronounced in the

gcn5-E191Q mutant could be explained by the physical interactions between SAGA and

SMC5/6. Such interactions may target SMC5/6 to SAGA-rich regions, while H3K9

acetylation opening chromatin would enable DNA entrapment. In the *gcn5-E191Q* mutant,

SMC5/6 could still be targeted to gene regions, although (unmodified) inaccessible chromatin

would hamper DNA entrapment. As the Nse3-mediated binding to free DNA is essential for

the SMC5/6 function, the DNA-unbound SMC5/6 in the *gcn5-E191Q* mutant would likely

cause a similar defect as in the Δ*gcn5* mutant and result in the same phenotypes of *gcn5-*

E191Q nse3-R254E and Δ*gcn5 nse3-R254E* double mutants (Suppl. Fig. S3D and E; [16,

21]).

The additive phenotypes of the double mutants (Fig. 3) probably stem from the non-

redundant functions of SMC5/6 and SAGA complexes. For example, SMC5/6 is essential for

replication processes (like replication restart; [34]), while SAGA is necessary for the timely

transcription of S-phase specific factors [59]. Defects in these functions probably combine in

double mutants to cause high HU sensitivity (Figs. 3, S3A, and D). Similarly, the reduced

DNA-binding affinity of the *nse3-R254E* mutant synergizes with the reduced binding to gene

regions upon *gcn5* deletion, resulting in the slow growth of double mutants. Interestingly,

enhanced SMC5/6 accumulation at repetitive regions (Fig. 4) could not rescue these growth

defects suggesting an important role of SMC5/6 binding to the gene regions.

It was recently shown that plant ADA2b binds SMC5 and assists in the localization of

SMC5/6 to DNA-damage sites [62]. Later studies uncovered an ADA2b-diRNA specific

pathway targeting SMC5/6 to DNA-damage foci in plants [63]. However, whether this

pathway involves other SAGA components and acetyltransferase activity is unclear. In fission

yeast, we and others showed that the SMC5/6 targeting to DNA-damage sites depends on the

BRCT domain-containing Brc1 protein but not on Ada2 or Gcn5 (Fig. 3; [64]), suggesting a

plant-specific function of the ADA2b-SMC5 interaction in DNA-damage response. Future

studies may show if SAGA likewise facilitates the accessibility of chromatin to SMC5/6 at

gene loci in plants. Interestingly, we found an interaction between human hADA2b and

hNSE4a/b subunits (Suppl. Fig. S2D), suggesting physical interaction between the SAGA and

SMC5/6 complexes is conserved in humans. It will be interesting to examine the functional

relationships between human SAGA and SMC5/6 complexes, particularly in facilitating

SMC5/6 targeting to gene regions.

Consistent with the lack of effect of SAGA on SMC5/6 localization to sites of DNA

damage, the SMC5/6 levels at repetitive regions, which are prone to DNA damage even

without any genotoxic treatment, were not reduced in Δ*gcn5* and Δ*ada2* deletion mutants

(Fig. 4). Furthermore, the SMC5/6 accumulation at repetitive regions depends on H3K9

methyltransferase Clr4 in fission yeast [53]. In our ChIP-seq experiments, the repetitive loci

were enriched for SMC5/6 upon SAGA HAT deletion (Fig. 4). We speculate that reduced

acetylation levels promoted an increase in methylation levels, indirectly stimulating the

SMC5/6 localization to repetitive heterochromatic regions [53, 65]. Altogether, our findings

support previous views that different factors target SMC5/6 to different genomic regions [34,

52, 53] and show a new role of SAGA in SMC5/6 targeting to gene regions.

Materials and methods

Yeast techniques

Standard fission yeast genetic techniques were used [66]. Yeast strains were crossed and

sporulated either at 25˚C (*ts* mutants) or 28˚C (non-*ts* mutants). Tetrad analysis was carried

out on Singer MSM300 (Singer, UK). The deletion integrations were verified on both ends by

PCR with specific primers to the G418 cassette and genomic sequence of a deleted gene

(approximately 600-800bp from start or end). The PCR products were sequenced.

S. pombe cultures were grown to the mid-log phase, and serial 10-fold dilutions were

spotted onto rich media with the indicated dose of DNA-damaging agent (hydroxyurea or

methyl methane sulfonate). Subsequently, plates were incubated at the indicated temperatures

(25, 28, or 37°C) for 3–4 days. Selective media were supplemented with Nourseothricin

(cloNAT, 100 μg/ml, Jena Bioscience), G418 (100 μg/ml; Applichem), and/or cycloheximide

(100 μg/ml; Sigma).

Yeast genetic screens

The pAW8-Nse3 integration construct [21] was modified for use in the PEM2 strain as

follows. The *Sph*I site was mutated to *Xho*I using a site-directed mutagenesis kit (Agilent

Technologies; primers: LJ48 and LJ49; Suppl. Table ST4). The cloNAT cassette was

amplified (LJ42 and LJ43) and inserted into the *Xho*I site in front of the Nse3 gene using the

In-fusion cloning kit (Takara). A 650bp-long genomic sequence (upstream of Nse3; LJ44 and

LJ45) was inserted in front of the cloNAT cassette (using *Xho*I) to ensure its proper

integration into the *S. pombe* genome. The mutant cloNAT-*nse3-R254E* construct was created

using site-directed mutagenesis (R254E_F and R254E_R primers; [21]). For the yeast

transformation, the WT and mutant cloNAT constructs were cleaved by *Spe*I, and the 3246bp

long fragment was purified from agarose gel by Gel Extraction Kit (Qiagen). Approximately

1 µg of purified DNA was transformed into the PEM2 strain [30] by standard LiAc protocol.

The proper integration of the cloNAT-Nse3 cassette and *rlp42* mutation was checked by PCR

and sequencing. The *nse3-R254E* PEM2 strain phenotypes were compared with the original

nse3-R254E strain (Suppl. Fig. S1A; [21])

The WT and *nse3-R254E* mutant PEM2 strains (YLJ222 and YLJ228; Suppl. Table

ST5) were crossed with the *S. pombe* haploid deletion library (BIONEER, version 5,

us.bioneer.com) according to the published protocol [30]. The screen was repeated twice

using the Rotor HDA robot (Singer, UK). The plate images were taken by a Canon EOS

Rebel T3i camera, and the individual colony size was measured. The viability of single

deletion mutants (control WT plates) against double mutants (test *nse3-R254E* plates) was

compared using SGAtools online platform [32]. Genes with a score less than -0.25 were

chosen as potential negative interactors (Suppl. Tab. ST1).

The resulting group of 79 genes was analysed by the Gene ontology tool BiNGo [67],

which is a plugin of the Cytoscape online platform [33]. The genes were classified according

to the Pombase GO database into Biological processes and Cellular component categories,

respectively [68]. The default parameters with a 0.05 significance level were applied for both categories.

Yeast two-hybrid analysis

The Gal4-based Y2H system was used to analyse SMC5/6-SAGA interactions [69]. *S. pombe* *ada2*, *ada3*, *gcn5*, and *sgf29* genes were PCR amplified from genomic DNA (primers used for *ada2* and *gcn5* cloning are listed in Suppl. Table ST4). All inserts were cloned into respective sites of the pGBKT7 or pGADT7 vectors using the In-Fusion cloning system. pGBKT7-Nse2(aa2-178) was described in [8]. pGADT7-Nse3(aa1-190) was prepared by mutagenesis of 191[st] aa to STOP codon in pGADT7-Nse3(aa1-328) [70]. The Nse3(aa200-307) fragment was cut out from pTriEx4-Nse3(aa200-307) [45] by *Nco*I-*Xho*I enzymes and cloned into pGADT7. The Smc5(aa170-225+837-910) fragment was amplified from the Smc5(aa2-225+837-1065) construct [71] and inserted into the *Nco*I-*Not*I sites of pGBKT7.

The pairs of pGBKT7 and pGADT7 constructs were co-transformed into the *Saccharomyces cerevisiae* PJ69–4a strain by standard LiAc transformation protocol and selected on SD -Leu, -Trp plates. Drop tests were carried out on SD -Leu, -Trp, -His (with 0.3, 0.5, 1, 3, 5, or 10 mM 3-aminotriazole) plates at 28°C. Each combination of partners was co-transformed and tested at least twice.

Co-immunoprecipitation of *S. pombe* proteins

Logarithmically growing YLJ507 and MMP21 cells (Suppl. Table ST5; Suppl. Fig. S3B) were cultivated in a rich medium at 28°C (OD$_{595}$ = 0.4 – 0.7). 5×10^8 cells were harvested by centrifugation (3 min, 4°C, 5000 rpm) and washed with 10 ml of ice-cold PBS. Pellets were stored in the 2 ml screw cup tubes at -80°C. The crude yeast extracts were prepared in 400 μl CHIP lysis buffer (50 mM HEPES, pH 7.5, 140 mM NaCl, 1 mM EDTA, 1% Triton X-100, Complete EDTA-free protease inhibitor cocktail tablets, Roche) with half volume of glass

beads (Sigma) in 2 ml low binding tubes using FASTprep-24 (MP Biomedicals; 5 times, 30

sec, 6.5 speed). The suspension was recovered by piercing the bottom of the tube with a

needle, placing it into a new 2 ml tube, and centrifugation (3 min, 4°C, 5000 rpm). The beads

were washed with 200 µl of CHIP lysis buffer (3 min, 4°C, 5000 rpm). The collected

suspensions were clarified by centrifugation (15 min, 4˚C, 15000 rpm), and the supernatant

was transferred to new low-binding tubes. 40 µl of cell extract was taken for input control.

Immunoprecipitation was carried out by adding 2 µl mouse anti-myc antibody (2276S, Cell

Signalling) to the cell extracts and incubation for 2 hours at 4°C. 20 µl of protein G coated

Dynabeads (Invitrogen) were washed twice with 1 ml CHIP lysis buffer and resuspended in

60 µl of CHIP lysis buffer, then added to the extract with anti-myc antibody and incubated

overnight at 4°C. The beads were pelleted using a magnetic rack, and the unbound fraction

(40 µl) was taken. Beads were washed four times with 1 ml CHIP lysis buffer, and proteins

were eluted by 40 µl of 1xSDS loading buffer. After 15 min incubation at room temperature,

the supernatant (bound fraction) was recovered. All fractions were analysed by western

blotting using mouse anti-myc-HRP (R951-25, Thermo Fisher) and mouse anti-FLAG-HRP

(F1804-1MG, Sigma) antibodies, respectively.

Chromatin immunoprecipitation analysis (ChIP)

Nse4-FLAG ChIP-seq

All strains (Suppl. Table ST5) were cultivated into the mid-log phase (OD = 0.4 – 0.6) and

incubated with 1 % formaldehyde for 15 min at room temperature to cross-link DNA–protein

complexes. Glycine was added to a final concentration of 125 mM, and the incubation

continued for 5 min. $5x10^8$ cells were harvested and washed with 10 ml of ice-cold PBS. The

yeast cell wall breakage was performed in 400 µl CHIP lysis buffer with half the volume of

glass beads in 2 ml low binding tubes using FASTprep-24. The suspension was washed two

times with CHIP lysis buffer (15 min, 4°C, 15000 rpm), and 300 μl of the extract was

sonicated with Bioruptor (Diagenode, 30 sec ON/30 sec OFF, High Power, 25 times) and

clarified by centrifugation (15 min, 4°C, 15000 rpm), resulting in an average DNA fragment

size of 300-500 bp. 5 μl of the sonicated precleared extract was taken as an input control

sample.

Monoclonal anti-FLAG M2 antibody (F1804, Sigma) was diluted 1:150, incubated

with precleared cell extract in 1.5 ml low binding tube for 2 h on ice, and precipitated over-

night with Dynabeads protein G (Invitrogen). Precipitates were washed with 1 ml of CHIP

lysis buffer, 1 ml of High Salt buffer (CHIP lysis buffer with 500 mM NaCl), 1 ml of Wash

buffer (10 mM Tris-HCl at pH 8.0, 0.25 M LiCl, 0.5% NP-40, 1 mM EDTA), and 1 ml of TE

buffer (20 mM Tris-HCl at pH 8.0, 1 mM EDTA). After elution (50 mM Tris at pH 8, 0.1%

SDS, 10 mM EDTA) and de-crosslinking overnight at 65°C, the DNA was purified using

QIAquick PCR Purification Kit (Qiagen).

For ChIP-seq analysis, the input DNA samples were tested for DNA fragmentation

and determination of DNA concentration by the Fragment analyser (Agilent). Input and

immunoprecipitation (IP) samples with the best fragmentation and high concentration were

used for the creation of NGS libraries (NEBNEXT ULTRA II DNA Library Prep kit, NEB)

and sequencing (Illumina Next seq 500, Illumina).

H3K9ac/H3 ChIP-seq

Two independent replicates were performed. Cells were grown to the exponential phase

(OD_{600} = 0.5) in the complex YES medium and fixed by adding formaldehyde to the final

concentration of 1 %. After 30 min incubation, the remaining formaldehyde was quenched by

125 mM glycine. Cells were washed with PBS and broken with glass beads. Extracted

chromatin was sheared with the Bioruptor sonicator (Diagenode) using 15 or 30 cycles (for

biological replicate 1 and 2, respectively) of 30 sec ON/30 sec OFF at high power settings.

For all immunoprecipitations (IP) within a biological replicate, the same amount of chromatin

extract was used (2.5 or 3.7 mg of total protein); 1/10 of the total chromatin extract amount

was kept for input DNA control. For each IP, 5 μg of antibody (H3: Ab1791, H3K9ac:

Ab4441, all Abcam) were incubated with the chromatin extract for 1 hour at 4°C with

rotation. Then, 50 μl of BSA-blocked Protein A-coated magnetic beads (10002D,

ThermoFisherScientific) were added to the chromatin extract-antibody suspension and

incubated for additional 4 hours at 4°C with rotation. The precipitated material and input

chromatin extract were de-crosslinked and treated with RNase A and proteinase K. DNA was

purified using phenol-chloroform extraction and sodium acetate/ethanol precipitation. In

biological replicate 2, DNA purification on AMPure XP beads (AC63880, Beckman Coulter)

was performed after the phenol-chloroform extraction to remove low-molecular fragments

and RNA. DNA concentration was measured using the Quantus fluorometer (Promega), and

fragment size distribution was checked on Agilent Bioanalyser using the High Sensitivity

DNA Assay. Library construction and sequencing (50 nt SE) were performed by BGI Tech

Solutions (Hong Kong) using the BGISEQ-500 sequencing system.

NGS data analysis

The reference fission yeast *S. pombe* genome (2018-09-04) and annotation (2019-11-15) were

downloaded from PomBase (https://www.pombase.org/; [68, 72]). Read quality was checked

using FastQC version 0.11.8 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/),

and reads were aligned to the *S. pombe* genome using HISAT2 2.1.0 [73] and SAMtools 1.9

[74, 75]. Read coverage tracks (i.e., target protein occupancy) were then computed and

normalized to the respective mapped library sizes using deepTools 3.5.1 [76]. The raw ChIP-

seq data are available from the ArrayExpress (https://www.ebi.ac.uk/) database under the

accession number E-MTAB-11081.

WT fission yeast RNA-seq data were obtained from the NCBI Sequence Read Archive

(https://www.ncbi.nlm.nih.gov/sra; datasets SRR8742773-SRR8742775; [61]). Reads were

processed and analysed with the same tools as above. All relevant scripts for (ChIP-seq and

RNA-seq) data processing and analysis are available from

https://github.com/mprevorovsky/Palecek-Nse-SAGA.

ChIP-qPCR

The Nse4-FLAG strains (crossed with the Δ*gcn5* or *gcn5-E191Q* strain; Suppl. Table ST5)

were used. The untagged wild-type strain was used as a negative control. All cells were

cultivated into the mid-log phase. Cells were then incubated with 1 % formaldehyde for 10

min at room temperature to cross-link DNA–protein complexes. Chromatin

immunoprecipitation was performed using a protocol described above for H3K9ac ChIP with

the following modifications. Monoclonal anti-FLAG M2 antibody (F1804; Sigma) was

diluted at 1:350 (5 µg /sample), incubated with 2 mg of total cell extract for 2 h at 4°C with

rotation, and precipitated with Dynabeads protein G (Invitrogen). After overnight incubation,

several washes, elution, and de-crosslinking, the DNA was purified using Phenol/Chloroform

method.

The relative amount of PCR product was quantified by qPCR using SensiFASTTM

SYBR® Hi -ROX Kit (Bioline). The sequences of primers used for the quantitative detection

of the chromosomal loci are listed in Suppl Table ST6. Input DNA recovery was calculated as

2 squared [CT(input)– CT(immunoprecipitate)] and normalized to a negative locus *slx9*. Melt

curve analysis was performed for each sample after PCR amplification to ensure that a single

product was obtained.

Microscopy

For the Nse4-GFP foci number determination, cells were grown in YES medium overnight,

diluted to OD = 0.4 in the morning, and treated with 0.03 % MMS or 20 mM HU for 5 h at 30

°C. 2.5 µl of cell culture was mounted on the slides, and GFP fluorescence was observed.

Pictures were taken on the Axio Imager Z1 microscope, using a Plan-Apochromat 63x oil

objective, the Axiocam CCD camera, and processed with the AxioVision software (all by

Zeiss). A minimum of 500 cells were counted in three independent experiments. For statistical

evaluation, p-values were calculated using the $\chi 2$ test.

Acknowledgements

We thank N. J. Krogan and M. N. Boddy for providing the yeast strains. We are grateful to C.

Haering for his help with the Rotor HDA robot and M. Adamus for the critical reading of our

manuscript.

Funding

Funding from the Ministry of Education, Youth, and Sports of the Czech Republic (project

INTER-COST LTC20033 to JJP) is gratefully acknowledged. BŠ was supported by the

Masaryk University grant MUNI/R/1142/2021. MP, AM, and JP were supported by the

Charles University grant PRIMUS/MED/26. We duly acknowledge the assistance of the

GeneCore facility (EMBL Heidelberg) and Genomics CF (CEITEC; project LM2018132

funded by MEYS CR) with the NGS part of our project. The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

Conceptualization: J.J.P., M.P., and D.H**.**; Data Curation, Formal Analysis, and Funding

Acquisition, Supervision: J.J.P. and MP**;** Investigation: L.M., BŠ., A.M., J.P., P.K., and E.L.;

Visualization: J.J.P., M.P. B.Š., and L.M.; Writing – Original Draft Preparation: J.J.P., M.P.,

P.K., and L.M.

References

1.   Uhlmann F. SMC complexes: from DNA to chromosomes. Nat Rev Mol Cell Biol.

2016;17(7):399-412.   Epub   2016/04/14.   doi: 10.1038/nrm.2016.30.   PubMed   PMID:

27075410.

2.   Davidson IF, Peters JM. Genome folding through loop extrusion by SMC complexes.

Nat Rev Mol Cell Biol. 2021;22(7):445-64. Epub 20210325. doi: 10.1038/s41580-021-00349-

7. PubMed PMID: 33767413.

3.   Ransom M, Dennehey BK, Tyler JK. Chaperoning histones during DNA replication

and   repair.   Cell.   2010;140(2):183-95.   doi: 10.1016/j.cell.2010.01.004.   PubMed   PMID:

20141833; PubMed Central PMCID: PMCPMC3433953.

4.   Morrison AJ, Shen XT. Chromatin remodelling beyond transcription: the INO80 and

SWR1   complexes.   Nature   Reviews   Molecular   Cell   Biology.   2009;10(6):373-84.   doi:

10.1038/nrm2693. PubMed PMID: WOS:000266270900012.

5.   Helmlinger D, Tora L. Sharing the SAGA. Trends Biochem Sci. 2017;42(11):850-61.

Epub   2017/09/27.   doi: 10.1016/j.tibs.2017.09.001.   PubMed   PMID: 28964624;   PubMed

Central PMCID: PMCPMC5660625.

6.   Palecek JJ, Gruber S. Kite Proteins: a Superfamily of SMC/Kleisin Partners

Conserved Across Bacteria, Archaea, and Eukaryotes. Structure. 2015;23(12):2183-90. doi:

10.1016/j.str.2015.10.004. PubMed PMID: WOS:000366171500001.

7.   Wells JN, Gligoris TG, Nasmyth KA, Marsh JA. Evolution of condensin and cohesin

complexes driven by replacement of Kite by Hawk proteins. Curr Biol. 2017;27(1):R17-R8.

doi: 10.1016/j.cub.2016.11.050.   PubMed   PMID: 28073014;   PubMed   Central   PMCID:

PMCPMC5228436.

8.   Andrews E, Palecek J, Sergeant J, Taylor E, Lehmann A, Watts F. Nse2, a component

of the Smc5-6 complex, is a SUMO ligase required for the response to DNA damage. Mol

Cell Biol. 2005;25(1):185-96. doi: 10.1128/MCB.25.1.185-196.2005. PubMed PMID:

WOS:000226236900016.

9.   Zhao X, Blobel G. A SUMO ligase is part of a nuclear multiprotein complex that

affects DNA repair and chromosomal organization. Proc Natl Acad Sci U S A.

2005;102(13):4777-47782. PubMed PMID: 15738391.

10.   Gligoris T, Löwe J. Structural Insights into Ring Formation of Cohesin and Related

Smc Complexes. Trends Cell Biol. 2016;26(9):680-93. Epub 2016/04/28. doi:

10.1016/j.tcb.2016.04.002. PubMed PMID: 27134029; PubMed Central PMCID:

PMCPMC4989898.

11.   Hassler M, Shaltiel IA, Haering CH. Towards a Unified Model of SMC Complex

Function. Curr Biol. 2018;28(21):R1266-R81. doi: 10.1016/j.cub.2018.08.034. PubMed

PMID: 30399354.

12.   Nasmyth K, Haering CH. The structure and function of SMC and kleisin complexes.

Annu Rev Biochem. 2005;74:595-648.

13.   Alt A, Dang HQ, Wells OS, Polo LM, Smith MA, McGregor GA, et al. Specialized

interfaces of Smc5/6 control hinge stability and DNA association. Nat Commun.

2017;8:14011. Epub 2017/01/30. doi: 10.1038/ncomms14011. PubMed PMID: 28134253;

PubMed Central PMCID: PMCPMC5290277.

14.   Hallett ST, Campbell Harry I, Schellenberger P, Zhou L, Cronin NB, Baxter J, et al.

Cryo-EM structure of the Smc5/6 holo-complex. Nucleic Acids Res. 2022. Epub 20220822.

doi: 10.1093/nar/gkac692. PubMed PMID: 35993814; PubMed Central PMCID:

PMCPMC9458440.

15.   Adamus M, Lelkes E, Potesil D, Ganji SR, Kolesar P, Zabrady K, et al. Molecular

Insights into the Architecture of the Human SMC5/6 Complex. Journal of Molecular Biology.

2020;432(13):3820-37.   doi: 10.1016/j.jmb.2020.04.024.   PubMed   PMID:

WOS:000541931500007.

16.   Yu Y, Li S, Ser Z, Kuang H, Than T, Guan D, et al. Cryo-EM structure of DNA-

bound Smc5/6 reveals DNA clamping enabled by multi-subunit conformational changes. Proc

Natl   Acad   Sci   U   S   A.   2022;119(23):e2202799119.   Epub   20220601.   doi:

10.1073/pnas.2202799119. PubMed PMID: 35648833.

17.   Ganji M, Shaltiel IA, Bisht S, Kim E, Kalichava A, Haering CH, et al. Real-time

imaging of DNA loop extrusion by condensin. Science. 2018;360(6384):102-5. Epub

2018/02/22. doi: 10.1126/science.aar7831. PubMed PMID: 29472443.

18.   Davidson IF, Bauer B, Goetz D, Tang W, Wutz G, Peters JM. DNA loop extrusion by

human   cohesin.   Science.   2019;366(6471):1338-45.   Epub   2019/11/21.   doi:

10.1126/science.aaz3418. PubMed PMID: 31753851.

19.   Wang XD, Hughes AC, Brandao HB, Walker B, Lierz C, Cochran JC, et al. In Vivo

Evidence for ATPase-Dependent DNA Translocation by the Bacillus subtilis SMC Condensin

Complex. Molecular Cell. 2018;71(5):841-7. doi: 10.1016/j.molcel.2018.07.006. PubMed

PMID: WOS:000443829100016.

20.   Pradhan B, Kanno K, Igarashi MU, Baaske MD, Wong JSK, Jeppsson K, et al. The

Smc5/6   complex   is   a   DNA   loop   extruding   motor.   bioRxiv.   2022.   doi:

10.1101/2022.05.13.491800.

21.   Zabrady K, Adamus M, Vondrova L, Liao C, Skoupilova H, Novakova M, et al.

Chromatin association of the SMC5/6 complex is dependent on binding of its NSE3 subunit

to DNA. Nucleic Acids Research. 2016;44(3):1064-79. doi: 10.1093/nar/gkv1021. PubMed

PMID: WOS:000371268700017.

22.   Piazza I, Rutkowska A, Ori A, Walczak M, Metz J, Pelechano V, et al. Association of

condensin with chromosomes depends on DNA binding by its HEAT-repeat subunits. Nat

Struct Mol Biol. 2014;21(6):560-8. Epub 20140518. doi: 10.1038/nsmb.2831. PubMed PMID: 24837193.

23.   Toselli-Mollereau E, Robellet X, Fauque L, Lemaire S, Schiklenk C, Klein C, et al. Nucleosome eviction in mitosis assists condensin loading and chromosome condensation. EMBO J. 2016;35(14):1565-81. Epub 2016/06/06. doi: 10.15252/embj.201592849. PubMed PMID: 27266525; PubMed Central PMCID: PMCPMC4946138.

24.   Shaltiel IA, Datta S, Lecomte L, Hassler M, Kschonsak M, Bravo S, et al. A hold-and-feed mechanism drives directional DNA loop extrusion by condensin. Science. 2022;376(6597):1087-94. Epub 20220602. doi: 10.1126/science.abm4012. PubMed PMID: 35653469.

25.   Shi ZB, Gao HS, Bai XC, Yu HT. Cryo-EM structure of the human cohesin-NIPBL-DNA complex. Science. 2020;368(6498):1454-+. doi: 10.1126/science.abb0981. PubMed PMID: WOS:000545264600036.

26.   Bürmann F, Funke LFH, Chin JW, Löwe J. Cryo-EM structure of MukBEF reveals DNA loop entrapment at chromosomal unloading sites. Mol Cell. 2021;81(23):4891-906.e8. Epub 20211104. doi: 10.1016/j.molcel.2021.10.011. PubMed PMID: 34739874; PubMed Central PMCID: PMCPMC8669397.

27.   Lopez-Serra L, Kelly G, Patel H, Stewart A, Uhlmann F. The Scc2-Scc4 complex acts in sister chromatid cohesion and transcriptional regulation by maintaining nucleosome-free regions. Nature Genetics. 2014;46(10):1147-51. doi: 10.1038/ng.3080. PubMed PMID: WOS:000342554100021.

28.   Munoz S, Minamino M, Casas-Delucchi CS, Patel H, Uhlmann F. A Role for Chromatin Remodeling in Cohesin Loading onto Chromosomes. Molecular Cell. 2019;74(4):664-+.   doi: 10.1016/j.molcel.2019.02.027.   PubMed   PMID: WOS:000468094300005.

29. Munoz S, Passarelli F, Uhlmann F. Conserved roles of chromatin remodellers in

cohesin loading onto chromatin. Current Genetics. 2020;66(5):951-6. doi: 10.1007/s00294-

020-01075-x. PubMed PMID: WOS:000558454500001.

30. Roguev A, Wiren M, Weissman JS, Krogan NJ. High-throughput genetic interaction

mapping in the fission yeast Schizosaccharomyces pombe. Nat Methods. 2007;4(10):861-6.

Epub 2007/09/23. doi: 10.1038/nmeth1098. PubMed PMID: 17893680.

31. Kim DU, Hayles J, Kim D, Wood V, Park HO, Won M, et al. Analysis of a genome-

wide set of gene deletions in the fission yeast Schizosaccharomyces pombe. Nat Biotechnol.

2010;28(6):617-23. Epub 2010/05/16. doi: 10.1038/nbt.1628. PubMed PMID: 20473289;

PubMed Central PMCID: PMCPMC3962850.

32. Wagih O, Usaj M, Baryshnikova A, VanderSluis B, Kuzmin E, Costanzo M, et al.

SGAtools: one-stop analysis and visualization of array-based genetic interaction screens.

Nucleic Acids Res. 2013;41(Web Server issue):W591-6. Epub 2013/05/15. doi:

10.1093/nar/gkt400. PubMed PMID: 23677617; PubMed Central PMCID:

PMCPMC3692131.

33. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a

software environment for integrated models of biomolecular interaction networks. Genome

Res. 2003;13(11):2498-504. doi: 10.1101/gr.1239303. PubMed PMID: 14597658; PubMed

Central PMCID: PMCPMC403769.

34. Palecek JJ. SMC5/6: Multifunctional Player in Replication. Genes. 2019;10(1):E7.

doi: 10.3390/genes10010007. PubMed PMID: WOS:000459743800007.

35. Aragon L. The Smc5/6 Complex: New and Old Functions of the Enigmatic Long-

Distance Relative. In: Bonini NM, editor. Annual Review of Genetics, Vol 52. Annual

Review of Genetics. 52. Palo Alto: Annual Reviews; 2018. p. 89-107.

36.   Lee KM, Nizza S, Hayes T, Bass KL, Irmisch A, Murray JM, et al. Brc1-mediated

rescue of Smc5/6 deficiency: requirement for multiple nucleases and a novel Rad18 function.

Genetics. 2007;175(4):1585-95.

37.   Lehmann AR, Walicka M, Griffiths DJF, Murray JM, Watts FZ, McCready S, et al.

The rad18 gene of Schizosaccharomyces pombe defines a new subgroup of the SMC

superfamily involved in DNA repair. Mol Cell Biol. 1995;15:7067-80.

38.   Morikawa H, Morishita T, Kawane S, Iwasaki H, Carr AM, Shinagawa H. Rad62

protein functionally and physically associates with the smc5/smc6 protein complex and is

required for chromosome integrity and recombination repair in fission yeast. Mol Cell Biol.

2004;24(21):9401-13.

39.   Dixon SJ, Fedyshyn Y, Koh JLY, Prasad TSK, Chahwan C, Chua G, et al. Significant

conservation of synthetic lethal genetic interaction networks between distantly related

eukaryotes. Proceedings of the National Academy of Sciences of the United States of

America. 2008;105(43):16653-8. doi: 10.1073/pnas.0806261105. PubMed PMID:

WOS:000260913500046.

40.   Ryan CJ, Roguev A, Patrick K, Xu J, Jahari H, Tong Z, et al. Hierarchical modularity

and the evolution of genetic interactomes across species. Mol Cell. 2012;46(5):691-704. doi:

10.1016/j.molcel.2012.05.028. PubMed PMID: 22681890; PubMed Central PMCID:

PMCPMC3380636.

41.   Pebernard S, McDonald WH, Pavlova Y, Yates JR, Boddy MN. Nse1, Nse2, and a

novel subunit of the Smc5-Smc6 complex, Nse3, play a crucial role in meiosis. Mol Biol Cell.

2004;15(11):4866-76. doi: 10.1091/mbc.E04-05-0436. PubMed PMID: 15331764; PubMed

Central PMCID: PMCPMC524734.

42.   Helmlinger D, Marguerat S, Villén J, Swaney DL, Gygi SP, Bähler J, et al. Tra1 has

specific regulatory roles, rather than global functions, within the SAGA co-activator complex.

EMBO J. 2011;30(14):2843-52. Epub 2011/06/03. doi: 10.1038/emboj.2011.181. PubMed

PMID: 21642955; PubMed Central PMCID: PMCPMC3160243.

43.   Verkade HM, Bugg SJ, Lindsay HD, Carr AM, O'Connell MJ. Rad18 is required for

DNA repair and checkpoint responses in fission yeast. Mol Biol Cell. 1999;10(9):2905-18.

44.   Kolesar P, Stejskal K, Potesil D, Murray JM, Palecek JJ. Role of Nse1 Subunit of

SMC5/6 Complex as a Ubiquitin Ligase. Cells. 2022;11(1):13. doi: 10.3390/cells11010165.

PubMed PMID: WOS:000758383300001.

45.   Hudson JJR, Bednarova K, Kozakova L, Liao CY, Guerineau M, Colnaghi R, et al.

Interactions between the Nse3 and Nse4 Components of the SMC5-6 Complex Identify

Evolutionarily Conserved Interactions between MAGE and EID Families. Plos One.

2011;6(2):14. doi: 10.1371/journal.pone.0017270. PubMed PMID: WOS:000287764100039.

46.   Duan X, Sarangi P, Liu X, Rangi GK, Zhao X, Ye H. Structural and functional

insights into the roles of the Mms21 subunit of the Smc5/6 complex. Mol Cell.

2009;35(5):657-68. PubMed PMID: 19748359.

47.   Clouaire T, Rocher V, Lashgari A, Arnould C, Aguirrebengoa M, Biernacka A, et al.

Comprehensive Mapping of Histone Modifications at DNA Double-Strand Breaks Deciphers

Repair Pathway Chromatin Signatures. Mol Cell. 2018;72(2):250-62.e6. Epub 2018/09/27.

doi: 10.1016/j.molcel.2018.08.020. PubMed PMID: 30270107; PubMed Central PMCID:

PMCPMC6202423.

48.   Deshpande GP, Hayles J, Hoe KL, Kim DU, Park HO, Hartsuiker E. Screening a

genome-wide S. pombe deletion library identifies novel genes and pathways involved in

genome stability maintenance. DNA Repair (Amst). 2009;8(5):672-9. Epub 2009/03/04. doi:

10.1016/j.dnarep.2009.01.016. PubMed PMID: 19264558; PubMed Central PMCID:

PMCPMC2675035.

49.   Pan X, Lei B, Zhou N, Feng B, Yao W, Zhao X, et al. Identification of novel genes

involved in DNA damage response by screening a genome-wide Schizosaccharomyces pombe

deletion library. BMC Genomics. 2012;13:662. Epub 2012/11/23. doi: 10.1186/1471-2164-

13-662. PubMed PMID: 23173672; PubMed Central PMCID: PMCPMC3536581.

50.   Helmlinger D, Marguerat S, Villén J, Gygi SP, Bähler J, Winston F. The S. pombe

SAGA complex controls the switch from proliferation to sexual differentiation through the

opposing roles of its subunits Gcn5 and Spt8. Genes Dev. 2008;22(22):3184-95. doi:

10.1101/gad.1719908.   PubMed   PMID: 19056896;   PubMed   Central   PMCID:

PMCPMC2593614.

51.   Oravcová M, Gadaleta MC, Nie M, Reubens MC, Limbo O, Russell P, et al. Brc1

Promotes the Focal Accumulation and SUMO Ligase Activity of Smc5-Smc6 During

Replication Stress. Mol Cell Biol. 2018. Epub 2018/10/22. doi: 10.1128/MCB.00271-18.

PubMed PMID: 30348841.

52.   Oravcová M, Boddy MN. Recruitment, loading, and activation of the Smc5-Smc6

SUMO ligase. Curr Genet. 2019;65(3):669-76. Epub 2019/01/02. doi: 10.1007/s00294-018-

0922-9. PubMed PMID: 30600397; PubMed Central PMCID: PMCPMC6511331.

53.   Pebernard S, Schaffer L, Campbell D, Head SR, Boddy MN. Localization of Smc5/6

to centromeres and telomeres requires heterochromatin and SUMO, respectively. EMBO J.

2008;27(22):3011-23.

54.   Lindroos HB, Strom L, Itoh T, Katou Y, Shirahige K, Sjogren C. Chromosomal

association of the Smc5/6 complex reveals that it functions in differently regulated pathways.

Mol Cell. 2006;22(6):755-67.

55.   Irmisch A, Ampatzidou E, Mizuno K, O'Connell MJ, Murray JM. Smc5/6 maintains

stalled   replication   forks   in   a   recombination-competent   conformation.   EMBO   J.

2009;28(2):144-55. Epub 2009/01/23. doi: emboj2008273 [pii] 10.1038/emboj.2008.273.

PubMed PMID: 19158664.

56.   Peng XP, Lim S, Li SB, Marjavaara L, Chabes A, Zhao XL. Acute Smc5/6 depletion

reveals its primary role in rDNA replication by restraining recombination at fork pausing

sites. Plos Genetics. 2018;14(1):20. doi: 10.1371/journal.pgen.1007129. PubMed PMID:

WOS:000423718600006.

57.   Torres-Rosell J, Machin F, Farmer S, Jarmuz A, Eydmann T, Dalgaard JZ, et al.

SMC5 and SMC6 genes are required for the segregation of repetitive chromosome regions.

Nat Cell Biol. 2005;7(4):412-9.

58.   Torres-Rosell J, Sunjevaric I, De Piccoli G, Sacher M, Eckert-Boulet N, Reid R, et al.

The Smc5-Smc6 complex and SUMO modification of Rad52 regulates recombinational repair

at the ribosomal gene locus. Nat Cell Biol. 2007;9(8):923-31. Epub 2007/07/22. doi:

10.1038/ncb1619. PubMed PMID: 17643116.

59.   González-Medina A, Hidalgo E, Ayté J. Gcn5-mediated acetylation at MBF-regulated

promoters induces the G1/S transcriptional wave. Nucleic Acids Res. 2019;47(16):8439-51.

doi: 10.1093/nar/gkz561. PubMed PMID: 31260531; PubMed Central PMCID:

PMCPMC6895280.

60.   Nugent RL, Johnsson A, Fleharty B, Gogol M, Xue-Franzén Y, Seidel C, et al.

Expression profiling of S. pombe acetyltransferase mutants identifies redundant pathways of

gene regulation. BMC Genomics. 2010;11:59. Epub 20100122. doi: 10.1186/1471-2164-11-

59. PubMed PMID: 20096118; PubMed Central PMCID: PMCPMC2823694.

61.   Elías-Villalobos A, Toullec D, Faux C, Séveno M, Helmlinger D. Chaperone-

mediated ordered assembly of the SAGA and NuA4 transcription co-activator complexes in

yeast. Nat Commun. 2019;10(1):5237. Epub 2019/11/20. doi: 10.1038/s41467-019-13243-w.

PubMed PMID: 31748520; PubMed Central PMCID: PMCPMC6868236.

62.   Lai J, Jiang J, Wu Q, Mao N, Han D, Hu H, et al. The transcriptional coactivator

ADA2b recruits a structural maintenance protein to double-strand breaks during DNA repair

in plants. Plant Physiol. 2018. Epub 2018/02/20. doi: 10.1104/pp.18.00123. PubMed PMID:

29463775.

63.   Jiang J, Ou X, Han D, He Z, Liu S, Mao N, et al. A diRNA-protein scaffold module

mediates SMC5/6 recruitment in plant DNA repair. Plant Cell. 2022. Epub 20220701. doi:

10.1093/plcell/koac191. PubMed PMID: 35775944.

64.   Oravcova M, Gadaleta MC, Nie MH, Reubens MC, Limbo O, Russell P, et al. Brc1

Promotes the Focal Accumulation and SUMO Ligase Activity of Smc5-Smc6 during

Replication Stress. Molecular and Cellular Biology. 2019;39(2):15. doi: 10.1128/mcb.00271-

18. PubMed PMID: WOS:000454772000001.

65.   Chiolo I, Minoda A, Colmenares SU, Polyzos A, Costes SV, Karpen GH. Double-

Strand Breaks in Heterochromatin Move Outside of a Dynamic HP1a Domain to Complete

Recombinational Repair. Cell. 2011;144(5):732-44. Epub 2011 Feb 25.

66.   Moreno S, Klar A, Nurse P. Molecular genetic analysis of fission yeast

Schizosaccharomyces pombe. Methods Enzymol. 1991;194:795-823. doi: 10.1016/0076-

6879(91)94059-l. PubMed PMID: 2005825.

67.   Maere S, Heymans K, Kuiper M. BiNGO: a Cytoscape plugin to assess

overrepresentation of gene ontology categories in biological networks. Bioinformatics.

2005;21(16):3448-9. Epub 2005/06/24. doi: 10.1093/bioinformatics/bti551. PubMed PMID:

15972284.

68.   Lock A, Rutherford K, Harris MA, Hayles J, Oliver SG, Bähler J, et al. PomBase

2018: user-driven reimplementation of the fission yeast database provides rapid and intuitive

access to diverse, interconnected information. Nucleic Acids Res. 2019;47(D1):D821-D7. doi:

10.1093/nar/gky961. PubMed PMID: 30321395; PubMed Central PMCID:

PMCPMC6324063.

69. Paleček JJ, Vondrová L, Zábrady K, Otočka J. Multicomponent Yeast Two-Hybrid

System: Applications to Study Protein-Protein Interactions in SMC Complexes. Methods Mol

Biol. 2019;2004:79-90. doi: 10.1007/978-1-4939-9520-2_7. PubMed PMID: 31147911.

70. Sergeant J, Taylor E, Palecek J, Fousteri M, Andrews E, Sweeney S, et al.

Composition and architecture of the Schizosaccharomyces pombe Rad18 (Smc5-6) complex.

Mol Cell Biol. 2005;25(1):172-84. doi: 10.1128/MCB.25.1.172-184.2005. PubMed PMID:

WOS:000226236900015.

71. Palecek J, Vidot S, Feng M, Doherty AJ, Lehmann AR. The SMC5-6 DNA repair

complex: Bridging of the SMC5-6 heads by the Kleisin, NSE4, and non-Kleisin subunits. J

Biol Chem. 2006;281:36952-9. PubMed PMID: 17005570.

72. Wood V, Gwilliam R, Rajandream MA, Lyne M, Lyne R, Stewart A, et al. The

genome sequence of Schizosaccharomyces pombe. Nature. 2002;415(6874):871-80.

73. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory

requirements. Nat Methods. 2015;12(4):357-60. Epub 20150309. doi: 10.1038/nmeth.3317.

PubMed PMID: 25751142; PubMed Central PMCID: PMCPMC4655817.

74. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. Epub 20090608.

doi: 10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed Central PMCID:

PMCPMC2723002.

75. Bonfield JK, Marshall J, Danecek P, Li H, Ohan V, Whitwham A, et al. HTSlib: C

library for reading/writing high-throughput sequencing data. Gigascience. 2021;10(2). doi:

10.1093/gigascience/giab007. PubMed PMID: 33594436; PubMed Central PMCID:

PMCPMC7931820.

76.   Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al.

deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids

Res. 2016;44(W1):W160-5. Epub 20160413. doi: 10.1093/nar/gkw257. PubMed PMID:

27079975; PubMed Central PMCID: PMCPMC4987876.

77.   Tapia-Alveal C, Lin SJ, Yeoh A, Jabado OJ, O'Connell MJ. H2A.Z-Dependent

Regulation of Cohesin Dynamics on Chromosome Arms. Mol Cell Biol. 2014;34(11):2092-

104.

78.   Bonnet J, Wang CY, Baptista T, Vincent SD, Hsiao WC, Stierle M, et al. The SAGA

coactivator complex acts on the whole transcribed genome and is required for RNA

polymerase   II   transcription.   Genes   Dev.   2014;28(18):1999-2012.   doi:

10.1101/gad.250225.114.   PubMed   PMID: 25228644;   PubMed   Central   PMCID:

PMCPMC4173158.

79.   Bavner A, Matthews J, Sanyal S, Gustafsson JA, Treuter E. EID3 is a novel EID

family member and an inhibitor of CBP-dependent co-activation. Nucleic Acids Res.

2005;33(11):3561-9. PubMed PMID: 15987788.

80.   Schneider M, Hellerschmied D, Schubert T, Amlacher S, Vinayachandran V, Reja R,

et al. The Nuclear Pore-Associated TREX-2 Complex Employs Mediator to Regulate Gene

Expression. Cell. 2015;162(5):1016-28. doi: 10.1016/j.cell.2015.07.059. PubMed PMID:

26317468; PubMed Central PMCID: PMCPMC4644235.

81.   Luthra R, Kerr SC, Harreman MT, Apponi LH, Fasken MB, Ramineni S, et al.

Actively transcribed GAL genes can be physically linked to the nuclear pore by the SAGA

chromatin modifying complex. J Biol Chem. 2007;282(5):3042-9. Epub 2006/12/06. doi:

10.1074/jbc.M608741200. PubMed PMID: 17158105.

82. Brown CE, Howe L, Sousa K, Alley SC, Carrozza MJ, Tan S, et al. Recruitment of

HAT complexes by direct activator interactions with the ATM-related Tra1 subunit. Science.

2001;292(5525):2333-7. doi: 10.1126/science.1060214. PubMed PMID: 11423663.

83. Rivosecchi J, Jost D, Vachez L, Gautier FDR, Bernard P, Vanoosthuyse V. RNA

polymerase backtracking results in the accumulation of fission yeast condensin at active

genes. Life Science Alliance. 2021;4(6):12. doi: 10.26508/lsa.202101046. PubMed PMID:

WOS:000654748200015.

84. Nakazawa N, Arakawa O, Yanagida M. Condensin locates at transcriptional

termination sites in mitosis, possibly releasing mitotic transcripts. Open Biology.

2019;9(10):9. doi: 10.1098/rsob.190125. PubMed PMID: WOS:000521587300003.

85. Hocquet C, Robellet X, Modolo L, Sun XM, Burny C, Cuylen-Haering S, et al.

Condensin controls cellular RNA levels through the accurate segregation of chromosomes

instead of directly regulating transcription. Elife. 2018;7:31. doi: 10.7554/eLife.38517.

PubMed PMID: WOS:000446417500001.

86. Kim KD, Tanizawa H, Iwasaki O, Noma K. Transcription factors mediate condensin

recruitment and global chromosomal organization in fission yeast. Nat Genet.

2016;48(10):1242-52. Epub 2016/08/22. doi: 10.1038/ng.3647. PubMed PMID: 27548313;

PubMed Central PMCID: PMCPMC5042855.

87. Lengronne A, Katou Y, Mori S, Yokobayashi S, Kelly GP, Itoh T, et al. Cohesin

relocation from sites of chromosomal loading to places of convergent transcription. Nature.

2004;430(6999):573-8. Epub 2004 Jun 30.

88. D'Ambrosio C, Schmidt CK, Katou Y, Kelly G, Itoh T, Shirahige K, et al.

Identification of cis-acting sites for condensin loading onto budding yeast chromosomes.

Genes Dev. 2008;22(16):2215-27. doi: 10.1101/gad.1675708. PubMed PMID: 18708580;

PubMed Central PMCID: PMCPMC2518811.

89.   Kakui Y, Barrington C, Barry DJ, Gerguri T, Fu X, Bates PA, et al. Fission yeast

condensin contributes to interphase chromatin organization and prevents transcription-

coupled DNA damage. Genome Biol. 2020;21(1):272. Epub 2020/11/05. doi:

10.1186/s13059-020-02183-0. PubMed PMID: 33153481; PubMed Central PMCID:

PMCPMC7643427.

90.   Borrie MS, Campor JS, Joshi H, Gartenberg MR. Binding, sliding, and function of

cohesin during transcriptional activation. Proc Natl Acad Sci U S A. 2017;114(7):E1062-e71.

Epub 20170130. doi: 10.1073/pnas.1617309114. PubMed PMID: 28137853; PubMed Central

PMCID: PMCPMC5320966.

91.   Kong M, Cutts EE, Pan D, Beuron F, Kaliyappan T, Xue C, et al. Human Condensin I

and II Drive Extensive ATP-Dependent Compaction of Nucleosome-Bound DNA. Mol Cell.

2020;79(1):99-114.e9. Epub 20200522. doi: 10.1016/j.molcel.2020.04.026. PubMed PMID:

32445620; PubMed Central PMCID: PMCPMC7335352.

92.   Johnsson A, Durand-Dubief M, Xue-Franzén Y, Rönnerblad M, Ekwall K, Wright A.

HAT-HDAC interplay modulates global histone H3K14 acetylation in gene-coding regions

during stress. EMBO Rep. 2009;10(9):1009-14. Epub 2009/07/24. doi:

10.1038/embor.2009.127. PubMed PMID: 19633696; PubMed Central PMCID:

PMCPMC2750064.

93.   Garcia-Luis J, Lazar-Stefanita L, Gutierrez-Escribano P, Thierry A, Cournac A, Garcia

A, et al. FACT mediates cohesin function on chromatin. Nature Structural & Molecular

Biology. 2019;26(10):970-+. doi: 10.1038/s41594-019-0307-x. PubMed PMID:

WOS:000488970400021.

94.   Shintomi K, Takahashi TS, Hirano T. Reconstitution of mitotic chromatids with a

minimum set of purified factors. Nat Cell Biol. 2015;17(8):1014-23. Epub 20150615. doi:

10.1038/ncb3187. PubMed PMID: 26075356.

95.   Nonaka N, Kitajima T, Yokobayashi S, Xiao G, Yamamoto M, Grewal SI, et al.

Recruitment of cohesin to heterochromatic regions by Swi6/HP1 in fission yeast. Nat Cell

Biol. 2002;4(1):89-93. doi: 10.1038/ncb739. PubMed PMID: 11780129.

96.   Choppakatla P, Dekker B, Cutts EE, Vannini A, Dekker J, Funabiki H. Linker histone

H1.8 inhibits chromatin binding of condensins and DNA topoisomerase II to tune

chromosome length and individualization. Elife. 2021;10:37. doi: 10.7554/eLife.68918.

PubMed PMID: WOS:000693094000001.

97.   Sánchez A, Roguev A, Krogan NJ, Russell P. Genetic Interaction Landscape Reveals

Critical Requirements for Schizosaccharomyces pombe Brc1 in DNA Damage Response

Mutants. G3 (Bethesda). 2015;5(5):953-62. Epub 2015/03/19. doi: 10.1534/g3.115.017251.

PubMed PMID: 25795664; PubMed Central PMCID: PMCPMC4426379.

Figure legends

Figure 1. Strong genetic relation between the SMC5/6 and histone acetyltransferases

(**A**) Summary of significantly enriched GO categories for Biological processes of genes with

genetic interactions with *nse3-R254E* (P < 0.005). (**B**) Negative log10 (P-values) evaluating

the significance of the main GO Cellular component terms identified in the set of the 79

genes. Only the molecular complexes are shown. (**C**) Tetrad analysis of the heterozygous

diploid fission yeast strains. The colony size of the *nse3-R254E*, Δ*gcn5* and *nse3-R254E*,

Δ*ada2* double mutant is significantly reduced (triangle). Single and double mutant alleles are

indicated. (**D**) Ten-fold serial dilutions of the indicated strains were plated onto YES media

and grown at 25°C (control) or 37°C. The *nse3-R254E* (*nse3/RE*) mutation enhanced the

sensitivity of the Δ*ada2*, Δ*ada3*, and Δ*gcn5* mutants to the higher temperature.

Figure 2. Interactions between the SMC5/6 and SAGA complexes

(**A**) Extracts from fission yeast strains MMP21 (Nse4-FLAG) and YLJ507 (Nse4-FLAG and

Gcn5-myc) were immunoprecipitated using the anti-myc antibody. The input (I), unbound (U)

and bound (B) fractions were separated by 12% SDS-PAGE. The Nse4-FLAG and Gcn5-myc

proteins were analysed on a western blot using anti-FLAG-HRP and anti-myc-HRP,

respectively. (**B**) The yeast two-hybrid (Y2H) system was used to determine individual

protein-protein interactions between SMC5/6 and SAGA HAT module subunits. The Gal4AD

or Gal4BD domains fused to the full-length Ada2 or Gcn5 subunits were co-transformed

together with the fragments of SMC5/6 subunits into the PJ69 cells and grown on the plates

without Leu, Trp (-L, W; control plates). The protein-protein interactions between SMC5/6

and SAGA were scored by the growth of the yeast PJ69 transformants on the plates without

Leu, Trp, and His, containing 3-Amino-1,2,4-triazole (0.5 mM AT or 10 mM AT plates). The

fragments were as follows: Gal4BD-Nse2 (aa2-178), Gal4AD-Nse3 (aa1-190), Gal4AD-Nse3

(aa200-307), and Gal4BD-SMC5 CC arm (aa170-225 + 837-910). In control experiments,

respective empty pGADT7 (AD) or pGBKT7 (BD) vector was co-transformed with either

SAGA or SMC5/6 construct. Note that the Gal4BD-SMC5 CC arm construct self-activated

(SMC5-vector combination) and was therefore grown on 10 mM AT plates to assess its

binding to Ada2 (SMC5-Ada2 combination).

Figure 3. SMC5/6 and SAGA are required for efficient DNA repair

(**A**) Sensitivity of the SMC5/6 and SAGA mutants to genotoxins. Ten-fold serial dilutions of

the yeast strains were plated onto YES media containing indicated concentrations of the

hydroxyurea (HU) or methyl methane sulfonate (MMS). The double mutants were more

sensitive than their respective single mutant counterparts, suggesting the non-redundant

functions of SMC5/6 and SAGA in DNA repair. (**B**) Live-cell microscopy of endogenous

Nse4-GFP upon HU and MMS treatment, respectively. The Nse4-GFP foci were present in

the WT and Δ*gcn5* cells but largely absent in Δ*brc1* cells. (**C**) Quantification of the data in

panel B suggests that the localization of SMC5/6 to the DNA-damage foci is independent of

SAGA.

Figure 4. SMC5/6 distribution is dependent on the SAGA HAT module

(**A**) The pie chart shows the distribution of the Nse4-FLAG peak areas in the different

genome regions in the WT fission yeast cells. Most SMC5/6 is localized to the repetitive

regions (like rDNA and centromeres), with the highest occupancy of the rDNA repeats. A

significant portion of the Nse4-FLAG is localized within the intergenic regions or genes. (**B**)

The heatmap diagrams compare the occupancy of Nse4-FLAG peaks in the WT, Δ*gcn5,*

Δ*ada2,* and Δ*ubp8* mutant cells (as identified in the WT). The top part shows peaks enhanced

in Δ*gcn5* and Δ*ada2* (enhanced)*,* while the bottom part clusters peaks reduced in the SAGA

HAT module deficient cells (reduced). Peaks in the rDNA repeats are shown separately as

these chromosome regions are not fully assembled and annotated in the *S. pombe* reference

genome and exert a different range of coverage values (rDNA). The Nse4-FLAG peaks

(normalized to median 760 bp width) and their surrounding regions (200 bp upstream and 200

bp downstream) are shown. (**C**) The pie charts show the distribution of the enhanced (top) and

reduced (bottom) Nse4-FLAG peak areas in Δ*gcn5*. The SMC5/6 accumulation is mainly

enhanced at the repetitive loci. The SMC5/6 localization is primarily reduced in gene regions.

(**D**) The box plot graph compares the Nse4-FLAG occupancy in WT and Δ*gcn5* cells. The

paired two-sided Wilcoxon statistical test was used: ns, non-significant; ***, $p < 0.001$.

Figure 5. The SMC5/6 accumulation correlates with the H3K9 acetylation status

(**A**) Heatmap statistical analysis of the loci with reduced SMC5/6 occupancy upon the *gcn5*

deletion. The Nse4-FLAG signals from WT and Δ*gcn5* are compared, and their differential

plot is shown in the middle panel (Δ*gcn5*-WT). Results from ChIP-seq analysis of H3 and

H3K9ac are shown (H3K9ac/H3). In addition, transcriptomic (RNA-seq) data are included

(mRNA). The genes (normalized to 1 kb width) and their surrounding regions (500bp

upstream of TSS and 1 kb downstream of TTS) are shown. (**B**) Scatter plot analysis shows a

strong correlation between the drop of SMC5/6 accumulation upon *gcn5* deletion (X-axis) and

H3K9-acetylation status (Y-axis) of gene regions (and their transcription levels; colour scale).

(**C**) Results of chromatin immunoprecipitation followed by quantitative PCR (ChIP–qPCR) at

selected gene loci are shown. Strains 503 (neg. control), MMP21 (WT, containing Nse4-

FLAG), Nse4-FLAG Δ*gcn5* (Δ*gcn5*), and Nse4-FLAG *gcn5-E191Q (gcn5-E191Q)* were

analysed. The fold enrichment was calculated against the negative *slx9* locus (mean ±

standard deviation, n ≥3 biological replicates). The unpaired Wilcoxon statistical test was

used: ns, non-significant; *, $p < 0.05$.

A.

BIOLOGICAL PROCESSES

p-value	Description
3.27E-13	DNA metabolic process
1.69E-12	response to DNA damage stimulus
4.47E-08	DNA repair
1.79E-07	cell cycle checkpoint
1.66E-06	post-translational protein modification
7.32E-06	chromosome organization
2.50E-05	histone acetylation
6.27E-05	meiosis
1.15E-04	DNA replication
1.20E-04	chromatin remodeling
2.61E-04	base-excision repair
7.50E-04	nucleotide-excision repair
1.33E-03	telomere organization
4.50E-03	positive regulation of ubiquitin-protein ligase activity

B.

MOLECULAR COMPLEXES

C.

Δgcn5 x nse3/RE

Δada2 x nse3/RE

○ WT

□ *nse3/R254E* mutant

⬡ single deletion mutant

△ double mutant

D.

Ctrl (25°C)   37°C

WT
Δgcn5
Δada2
Δada3
Δsgf29
nse3/RE
Δgcn5, nse3/RE
Δada2, nse3/RE
Δada3, nse3/RE
Δsgf29, nse3/RE

A.

control			Gcn5-myc			
I	U	B	I	U	B	

Gcn5-myc

Nse4-FLAG

B.

A.

	Ctrl	1mM HU	2mM HU	3mM HU	0.004% MMS

WT
Δgcn5
Δada2
Δada3
Δsgf29
nse3/RE
Δgcn5, nse3/RE
Δada2, nse3/RE
Δada3, nse3/RE
Δsgf29, nse3/RE

B.

Nse4-GFP

	WT	Δgcn5	Δada2	Δbrc1

MMS

HU

C.

MMS

HU

A. Nse4 distribution

- rDNA repeats
- centromeric regions
- intergenic regions
- gene regions

B.

WT | Δgcn5 | Δada2 | Δubp8

rDNA

enhanced

unchanged

reduced

0.2 peak 0.2 | 0.2 peak 0.2 | 0.2 peak 0.2 | 0.2 peak 0.2

C.

enhanced

reduced

- gene regions
- repeats
- intergenic reg.

D. Nse4 occupancy

normalized peak coverage

10^7
10^6
10^5
10^4

□ WT
■ Δgcn5

*** | *** | ns | ***

rDNA repeats | centromeric repeats | intergenic regions | gene regions

A.

Nse4-FLAG

| WT | Δ*gcn5* | Δ*gcn5*-WT | H3K9ac/H3 | mRNA |

gene regions with reduced Nse4 occupancy in Δ*gcn5*

0.5 gene 1.0Kb   0.5 gene 1.0Kb   0.5 gene 1.0Kb   0.5 gene 1.0Kb   0.5 gene 1.0Kb

0 15 30 45   0 15 30 45   -10 -5 0   0.0 1.5   0 15 30 45

B.

n=234, Pearson's r=-0.296, p=3.97e^{-6}

H3K9ac/H3

Nse4-FLAG (Δ*gcn5*-WT)

mRNA

C.

ChIP-qPCR

Fold enrichment

ns

vht1   *rps502*   *fip1*   *ser3*   *tef3*

neg. control
WT
Δ*gcn5*
gcn5-E191Q