FLAIRR-seq: A novel method for single molecule resolution of near full-length

immunoglobulin heavy chain repertoires

Authors:

Easton E. Ford[1,2*], David Tieri[2], Oscar Rodriguez[2], Nancy Francoeur[3], Juan Soto[3], Justin Kos[2],

William Gibson[2], Catherine A. Silver[2], Gintaras Deikus[3], Elizabeth Hudson[2], Cassandra R.

Woolley[1], Noam Beckmann[3], Alexander Charney[3], Thomas C. Mitchell[1], Robert P. Sebra[3], Corey

T. Watson[2*], Melissa L. Smith[2*]

Affiliations:

* denotes equal author contribution to the manuscript

[1]Department of Microbiology and Immunology University of Louisville School of Medicine,

Louisville, Kentucky, USA

[2]Department of Biochemistry and Molecular Genetics University of Louisville School of

Medicine, Louisville, Kentucky, USA

[3]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New

York City, New York, USA

Corresponding Author:

Melissa L. Smith, PhD

ml.smith@louisville.edu

24
25

Abstract:

Current Adaptive Immune Receptor Repertoire Sequencing (AIRR-seq) strategies resolve immunoglobulin (IG) variable region transcripts with limited resolution of the constant region. Here we present a novel near full-length AIRR-seq (FLAIRR-Seq) method that utilizes targeted heavy chain amplification by 5' rapid amplification of cDNA ends (RACE), combined with single molecule, real-time (SMRT) sequencing to generate highly accurate (>Q40, 99.99%), immunoglobulin heavy chain transcripts. FLAIRR-seq was benchmarked by comparing V, D, and J gene usage, complementarity-determining region 3 (CDR3) length, and the extent of total variable gene somatic hypermutation to matched datasets generated with standard 5' RACE AIRR-seq and full-length isoform sequencing methods. Together these data demonstrate robust, unbiased FLAIRR-seq performance using RNA samples derived from peripheral blood mononuclear cells, purified B cells, and whole blood, which recapitulated results generated by commonly used methods, while resolving novel constant region features. FLAIRR-seq data provides, for the first time, simultaneous, single-molecule characterization of variable and constant region genes and alleles, allele-resolved subisotype definition, and identification of class-switch recombination within a clonal lineage. FLAIRR-seq, in conjunction with IGHC genotyping, of the IgM- and IgG repertoires from 10 individuals resulted in the identification of 32 unique IGHC alleles; 28/32 (87%) that were not represented by sequences curated in the IMmunoGeneTics Information System (IMGT) database. Together, these data demonstrate the capabilities of FLAIRR-seq to characterize IGHV, D, J and C gene diversity for the most comprehensive view of B cell repertoires to date.

Introduction

Antibodies or immunoglobulins (IG) are the primary effectors of humoral immunity and are found as both membrane-bound receptors on B cells and circulating, secreted proteins (1). Membrane-bound IG (also known as the B cell receptor (BCR)) and secreted IG act to recognize and bind specific antigen. All IG are composed of a heavy chain and a light chain that are post-translationally associated to complete the correctly folded IG. The heavy chain is comprised of two distinct domains: (i) the variable domain (Fab), which allows for antigen binding, and (ii) the constant domain (Fc), which modulates downstream effector functions (1, 2). The IG light chain also includes a variable domain that, once post-translationally associated with the heavy chain variable domain in order to interact with cognate antigen; (3). Heavy chain constant region genes separate IG into discrete isotypes (i.e., IgM, IgD, IgG, IgA, IgE). Each isotype demonstrates unique effector properties that together represent the wide diversity of IG-mediated functions, including binding of Fc receptors (FCR), activation of complement, opsonization, antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP) (4, 5).

To recognize the wide range of pathogens humans encounter, the IG loci are highly polymorphic and harbor diverse and complex sets of genes that recombine in each B cell to encode up to 10^{13} unique IG specificities (6). B cells create this expansive catalog of specificities through somatic recombination of the variable (V), diversity (D), and joining (J) genes in the IG heavy chain locus (IGH), and V and J genes from the corresponding light chain loci, IG lambda (IGL) and kappa (IGK) (7). First, a single D and J gene are recombined, while the intervening and unused D and J gene sequences are removed by RAG recombinase (8). After D and J genes are recombined, further recombination of a specific V gene to the DJ gene cassette completes the formation of the full VDJ exon. After transcription of the recombined variable region, unique constant region genes are spliced with the VDJ cassette to generate the heavy chain transcript (7). Recombination of the light chain occurs similarly, recombining V and J genes only. Heavy

and light chain transcripts are independently translated and linked via covalent cysteine bonds

resulting in a fully functional IG protein prior to B cell cell-surface expression or secretion (**Figure**

1A) (9). Naïve B cells, which develop in the bone marrow from hematopoietic stem cell

progenitors, have undergone VDJ recombination but have not yet encountered antigen, and solely

express IgM and IgD (10). These naïve B cells then migrate to B cell zones in secondary lymphoid

tissues where they encounter antigen, driving further maturation and class switch recombination

(CSR) to enable the most effective humoral response (11). CSR allows for further excision of

constant genes at the DNA level, shifting the utilization of constant genes in the heavy chain locus

and resulting in different IG isotypes and subisotypes linked to the same VDJ (11).

The IgG isotype family is a composite of four subisotypes: IgG1, IgG2, IgG3, and IgG4.

Each IgG subisotype circulates at varied frequencies and facilitates unique immune functions. For

example, IgG1 is typically the most abundant circulating IgG and mediates proinflammatory

responses; IgG2 targets bacterial polysaccharides, conferring general protection from bacterial

pathogens; IgG3 confers protection against intracellular bacterial infections and enables clearing

of parasites; and IgG4 contains exclusive structural and functional characteristics often resulting

in anti-inflammatory and tolerance-inducing effects (5). Multiple studies have identified antibody-

mediated subisotype-specific pathogenicity in the context of autoimmune diseases and cancer

highlighting the need for further investigation of subisotype-specific IG repertoires (12-15).

Current AIRR-seq methods aim to resolve IG variable region transcripts to differing

extents. Profiling of this region, even in part, defines V, D and J gene usage while also providing

characterization of complementarity determining regions (CDR) 1, 2, and 3, which are

hypervariable and directly interact with target antigen (**Figure 1B**) (16). CDR3-targeted profiling

approaches, such as those used by Adaptive Biotechnologies (17), allow for V, D, and J gene

assignments but do not provide complete resolution of the entire variable region (17). Multiplexed

primer-based AIRR-seq strategies generate full variable region content but can be restrictive due

to the use of complex primer pools that can lead to inconsistent sequence coverage and will not

capture unknown genes and alleles for which there are no primers included. 5' RACE AIRR-seq

methods capture the full-length VDJ exon without multiplexed primer pools therefore limiting the

impact of primer bias and enabling discovery of novel genes and alleles (18). Crucially, 5' RACE

methods can also prime from the first constant region exon (CH1), allowing for determination of

isotype with the variable region. Additional methods have been developed that shift amplification

strategies by capturing more of the CH1 exon to enable subisotype resolution; however, these

methods sacrifice full and contiguous characterization of the V gene segment (19). All commonly

used AIRR-seq methods are further technically limited by the length restrictions (≤ 600nt) of short-

read next generation sequencing. As a result, no current AIRR-seq strategy resolves the complete

IG heavy chain transcript from the start of the constant exons to the end of the VDJ exon. Our

team has recently shown that population-based polymorphisms within the IGHV, IGHD, IGHJ and

IGL loci are far more extensive than previously known; the constant region has also been shown

to contain genomic diversity (20-22). Although it is understood that the Fc domain mediates IG

effector function, there is limited knowledge as to how genetic variation in this region may impact

functional capabilities (5, 23, 24). As such, there is a growing need to understand genomic

variation across the complete IG molecule. To address these limitations, we have developed

novel, end-to-end pipeline to target, profile, and characterize the IG heavy chain repertoire in the

context of isotype (IgG, IgM) and subisotype (IgG1, IgG2, IgG3, IgG4) groups.

Here, we present FLAIRR-seq, a novel targeted 5' RACE-based amplification of near full-

length IgG and IgM IGH transcripts, paired with single molecule real time (SMRT) sequencing,

resulting in highly accurate (mean read accuracy as high as Q60, 99.9999%), near full-length IG

sequences from RNA derived from whole blood, isolated PBMC, and purified B cells. When

analyzed with the Immcantation AIRR-seq tool suite, we demonstrate that FLAIRR-seq performs

comparably to standard 5'RACE AIRR-seq methods and single-molecule isoform sequencing

(IsoSeq) strategies for characterizing adaptive immune receptor repertoires. We further highlight

the novel features of FLAIRR-seq data, including phased identification of V, D, J, and constant

genes, facilitating the profiling of subisotype- and IGHC allele-specific repertoires and the

characterization of class switch recombination.

Materials and Methods

Sample collections

Experiments were conducted using healthy donor peripheral blood mononuclear cells (PBMC), purified B cells from healthy donors, or whole blood collected from hospitalized COVID-19 patients (Supplementary Table 1). Commercially available healthy donor PBMC (STEMCELL Technologies) and a subset of matched purified B cells were utilized to generate AIRR-seq and FLAIRR-seq validation datasets. Full-length isoform sequencing (IsoSeq) was performed using B cells isolated from the PBMC of a healthy, consented 57-year-old male donor at the University of Louisville (UofL) School of Medicine. The UofL Institutional Review Board approved sample collection (IRB 14.0661). For COVID-19 affected patient samples (n=5), whole blood was collected from the Mount Sinai COVID-19 biobank cohort of hospitalized COVID-19 patients, approved by the Institutional Review Board at the Icahn School of Medicine at Mount Sinai as previously described (25).

PBMC isolation and B cell purification

Frozen healthy donor PBMCs were purchased, thawed, and aliquoted for use in downstream experiments. For IsoSeq analyses, 175mL of venous blood was collected in a final concentration of 6mM K_3EDTA using standard phlebotomy. PBMCs were isolated using Sepmate PBMC Isolation Tubes (STEMCELL Technologies) as previously described (26), with an additional granulocyte depletion step using the RosetteSep Human Granulocyte Depletion Cocktail (STEMCELL Technologies) as directed by the manufacturer. B cells from the freshly collected and frozen healthy donor PBMC were isolated using the EasySep Human Pan-B Cell Enrichment Kit, as described by the manufacturer (STEMCELL Technologies). Briefly, B cells, including plasma cells, were isolated by negative selection using coated magnetic particles. First, the B cell enrichment cocktail was added to the sample and mixed for a 5-minute incubation at room temperature, followed by addition of magnetic particles and further incubation for 5 minutes

182   on the benchtop. The sample tube was then placed on an EasySep magnet (STEMCELL

183   Technologies), and purified B cells were carefully eluted from the magnetic particles and

184   immediately used for RNA extraction.

Genomic DNA and RNA extraction

For the healthy frozen PBMC and matched purified B cells, genomic DNA (gDNA) and

RNA were co-extracted using the AllPrep DNA/RNA Mini Kit (Qiagen) according to the

manufacturer's instructions. For the freshly processed UofL healthy donor PBMC, purified Pan-B

cells were lysed in Buffer RLT Plus, and RNA was extracted with the RNeasy Plus Mini Kit

(Qiagen) per the manufacturer's protocol; no gDNA was collected from this sample. COVID-19

whole blood-derived RNA was extracted from samples collected in Tempus Blood RNA tubes

using the Tempus Spin RNA Isolation Kit (ThermoFisher) as described by the manufacturer. For

all samples, concentrations of RNA and gDNA (when appropriate) were assessed using the Qubit

4.0 fluorometer, with the RNA HS Assay Kit and Qubit DNA HS Assay Kit, respectively

(ThermoFisher Scientific). RNA and gDNA integrity were evaluated using the Bioanalyzer RNA

Nano Kit and DNA 1200 Kit, respectively (Agilent Technologies). Extracted RNA and gDNA were

stored at -80℃ and -20℃, respectively, until used.

FLAIRR-seq targeted amplification of heavy chain transcripts

Extracted RNA was thawed on ice and converted to first strand complementary DNA

(cDNA) using the SMARTer RACE 5'/3' Kit (Takara Bio USA), as described by the manufacturer

and a custom oligonucleotide that contained the template switch oligo and a unique molecular

identifier (5' TSO-UMI) for template switch during first strand cDNA synthesis. The following

reaction conditions were used: (i) a primary master mix was prepared with 4.0 µL 5X First-Strand

Buffer, 0.5 µL DTT (100 mM), and 1.0 µL dNTPs (20 mM) per reaction and set aside until needed;

(ii) in a separate 0.2-mL PCR tube, 10 µL of sample RNA and 1 µL 5'-CDS Primer A were

combined and incubated in a thermal cycler at 72°C (lid temperature: 105°C) for 3 minutes,

followed by cooling to 42°C for 2 minutes; (iii) after cooling, tubes were spun briefly to collect

contents and 1μL (12μM) of the 5' TSO-UMI was added to the RNA; (iv) 0.5 μL of RNase inhibitor

and 2.0 μL of SMARTScribe Reverse Transcriptase were added to the primary master mix tube

per sample and 8 μL of the combined master mix was then added to each RNA-containing sample

tube. First-strand cDNA synthesis reactions were incubated in a thermal cycler at 42°C (lid

temperature: 105°C) for 90 mins, followed by heat inactivation at 70°C for 10 minutes. Total first

strand cDNA generated in this reaction was diluted 1:2 with Tricine-EDTA Buffer before moving

onto targeted heavy chain transcript amplification.

To specifically amplify heavy chain transcripts from total first-strand cDNA, targeted IgG

and IgM transcript amplification reactions were performed using barcoded IgG (3' primer binding

in the constant region exon 3, CH3) or IgM (3' primer binding in the constant region exon 4, CH4)-

specific primers (Supplementary Table 2) and the following conditions: (i) 5 μL of diluted first-

strand cDNA was added to 0.2-mL PCR tubes; (ii) a master mix was generated using 10 μL 5X

PrimeSTAR GXL Buffer, 4 μL GXL dNTP mixture, 28 μL PCR-grade water, 1 μL PrimeSTAR GXL

Polymerase and 1 μL 10x UPM form the SMARTer RACE 5'/3' Kit per reaction; (iii) 44 μL master

mix was added to each reaction tube followed by 1 μL of the appropriate barcoded IgG (CH3) or

IgM (CH4) primer. Different temperatures were used for annealing of IgG- (63.5°C) and IgM-

specific primers (60°C) to account for primer specific melting temperatures and to enhance

targeted amplification specificity. Amplification conditions for full-length IgG were: 1 minute at

95°C, followed by 35 amplification cycles of 95°C for 30 sec, 63.5°C for 20 sec, and 2 minutes at

68°C, followed by a final extension for 3 minutes at 68°C and hold at 4°C. Amplification conditions

for full-length IgM were: 1 minute at 95°C, followed by 35 amplification cycles of 95°C for 30 sec,

60°C for 20 sec., and 2 minutes at 68°C, followed by a final extension for 3 minutes at 68°C and

hold at 4°C. Final amplification reactions were purified using a 1.1x (vol:vol) cleanup with ProNex

magnetic beads (Promega) . Successfully amplified products were quantified with Qubit dsDNA

HS assay (ThermoFisher Scientific) and length was evaluated with the Fragment Analyzer

Genomic DNA HS assay (Agilent). Samples were equimolar pooled in 8-plexes for SMRTbell

library preparation and sequencing.

FLAIRR-seq SMRTbell library preparation and sequencing

Eight-plex pools of targeted IgG or IgM amplicons were prepared into SMRTbell

sequencing templates according to the "Procedure and Checklist for Iso-Seq Express Template

for Sequel and Sequel II systems" protocol starting at the "DNA Damage Repair" step and using

the SMRTbell Express Template Prep Kit 2.0, with some modifications (Pacific Biosciences).

Briefly, targeted IgG and IgM amplicons underwent enzymatic DNA damage and end repair,

followed by ligation with overhang SMRTbell adapters as specified in the protocol. To increase

consistency in SMRTbell loading on the Sequel IIe system, we further treated the SMRTbell

libraries with a nuclease cocktail to remove unligated amplified products, using the SMRTbell

Enzyme Cleanup Kit, as recommended by the manufacturer (Pacific Biosciences). Briefly, after

heat-killing the ligase with an incubation at 65°C, samples were treated with a nuclease cocktail

at 37°C for 1 hour, and then purified with a 1.1X Pronex cleanup. Final SMRTbell libraries were

evaluated for quantity and quality using the Qubit dsDNA HS assay and Fragment Analyzer

Genomic DNA assay, respectively. Sequencing of each 8-plex, barcoded sample pool was

performed on one SMRTcell 8M using primer v4 and polymerase v2.1 on the Sequel IIe system

with 30 hr movies. Demultiplexed, high-fidelity circular consensus sequence reads ("HiFi reads")

were generated on the instrument for downstream analyses.

AIRR-seq SMARTer Human BCR IgG/IgM IGH sequencing

Matched healthy donor RNA was used to generate targeted IgG and IgM AIRR-seq

libraries using the SMARTer Human BCR IgG IgM H/K/L Profiling Kit (Takara Bio USA) according

to the manufacturer's instructions with no modifications. Briefly, for each sample, proprietary IgG

and IgM primers were used to amplify heavy chain transcripts following a 5'RACE reaction. AIRR-

seq libraries were then quality controlled using the 2100 Bioanalyzer High Sensitivity DNA Assay

Kit (Agilent) and the Qubit 3.0 Fluorometer dsDNA High Sensitivity Assay Kit. Sequencing on the

MiSeq platform using 300bp paired-end reads was performed using the 600-cycle MiSeq Reagent

Kit v3 (Illumina) according to the manufacturer's instructions, and FASTQ reads were generated

using the associated DRAGEN software package (Illumina).

B cell IsoSeq

RNA extracted from healthy sorted B cells was used to generate IsoSeq SMRTbell

libraries following the "Procedure & Checklist Iso-Seq Express Template Preparation for the

Sequel II System" with minor adaptations compared to the manufacturer's instructions. Briefly,

Iso-Seq libraries were generated using 500 ng high-quality (RIN > 8) RNA as input into oligo-dT

primed cDNA synthesis (NEB). Barcoded primers were incorporated into the cDNA during second

strand synthesis. Following double-stranded cDNA amplification, transcripts from two samples

sourced from purified B cells and NK cells were equimolar pooled as previously described (26).

SMRTbells were generated from the pooled cDNA as described above for the FLAIRR-seq

amplification products, including the addition of a nuclease digestion step. Quantity and quality of

the final IsoSeq libraries were performed with the Qubit dsDNA High Sensitivity Assay Kit and

Agilent Fragment Analyzer Genomic DNA assay, respectively. This 2-plex IsoSeq pool was

sequenced using primer v4 and polymerase v2.1 on the Sequel IIe system with a 30-hour movie.

HiFi reads were generated on instrument before analyses. Demultiplexing of barcoded samples

and generation of full-length non-concatemer (FLNC) predicted transcripts were performed using

the IsoSeq v3 pipeline available through SMRTLink (v.10.2). B-cell-derived FLNC reads were

mapped to the human genome using the GMAP reference database, and reads derived from

chromosome 14 were extracted for downstream IGH transcript characterization via Immcantation,

as described below.

Immcantation analyses of IgG and/or IgM IGH variable region repertoires

Analyses of FLAIRR-seq, AIRR-seq, and Iso-seq datasets were performed using

Immcantation tools to characterize IG variable regions (27, 28). Demultiplexed barcoded HiFi (for

SMRT sequencing data) or FASTQ (for AIRR-seq) reads were first processed using the pRESTO

tool for quality control, UMI processing, and error profiling (28). For AIRR-seq, pRESTO analysis

data from paired-end reads ("R1" and "R2") were trimmed to remove bases with < Q20 read

quality and/or <125 bp length using the "FilterSeq trimqual" and "FilterSeq length", respectively.

IgG and IgM CH3 or CH4 primer sequences were identified with an error rate of 0.2, and primers

identified were then noted in FASTQ headers using "MaskPrimers align". Next, 12 basepair (bp)

UMIs were located and extracted using "Maskprimers extract". Sequences found to have the

same UMIs were grouped and aligned using "AlignSets muscle," with a consensus sequence

generated for each UMI using "BuildConsensus". Mate pairing of AIRR-seq reads was conducted

using a reference-guided alignment requiring a minimum of a 5 bp overlap via "AssemblePairs

sequential". After collapsing consensus reads with the same UMI ("conscount") using

"CollapseSeq"," reads with < 2 supporting sequences were removed from downstream analysis.

For pRESTO processing of FLAIRR-seq, single HiFi reads ("R1") reads did not require trimming

due to > Q20 sequence quality across all bases. 5'TSO-UMI and CH3 or CH4 region primers were

identified along with a 22bp UMI with an error rate of 0.3 using "MaskPrimers align". Reads were

then grouped and aligned using "AlignSets muscle". Due to the single molecule nature of FLAIRR-

seq reads, no mate pairing was required. Consensus reads were then generated as described

above, including removal of sequences with < 2 supporting reads.

pRESTO-filtered reads for both AIRR-seq and FLAIRR-seq data were then input into the

Change-O tool (Table 1). Iso-seq required no initial processing from pRESTO and was input into

Change-O for IG gene reference alignment along with AIRR-seq and FLAIRR-seq data using

"igblastn", clonal clustering using "DefineClones", and germline reconstruction and conversion

using "CreateGermlines.py" and the GRCh38 chromosome 14 germline reference (27). Fully

processed and annotated data was then converted into a TSV format for use in downstream

analyses. The Alakazam Immcantation tool suite was then used to quantify gene usage analysis,

calculate CDR3 length, assess somatic hypermutation frequencies, and analyze clonal diversity

(27). Scoper clonal assignment by spectral clustering was conducted for COVID-19 patient time

course samples (29, 30). For clonal lineage tree analysis, the Dowser tool was used to examine

clonal diversity and CSR over time (31).

Targeted IG gDNA capture, long-read sequencing, and IGenotyper analyses.

FLAIRR-seq validation samples also underwent IGHC targeted enrichment and long-read

sequencing as previously described (32). Briefly, gDNA was mechanically sheared and size

selected to include 5-9kb fragments using the BluePippin (Sage Science). Samples were then

end repaired and A-tailed using the standard KAPA library preparation protocol (Roche).

Universal priming sequences and barcodes were then ligated onto samples for multiplexing

(Pacific Biosciences). Barcoded gDNA libraries were captured using IGH-specific probes

following a SeqCap protocol (Roche). 26 IGH-enriched samples were purified and pooled together

for SMRTbell library prep as described above, including the final nuclease digestion step. Pooled

SMRTbells were annealed to primer v4, bound to polymerase v2.0, and sequenced on the Sequel

IIe system with 30h movies. After sequencing, HiFi reads were generated and analyzed by the

IGenotyper pipeline (32). In brief, IGenotyper was used to detect single nucleotide variants and

assemble sequences into haplotype assemblies for downstream constant region genotyping.

Constant region genotyping with FLAIRR-seq

To genotype constant region genes and alleles from FLAIRR-seq data, productive reads

were filtered by constant region length (900bp–1100bp) and aligned to the chromosome 14 hg38

reference using minimap2 along with SAMtools to generate sorted and indexed bam files (33,

34). WhatsHap was used to identify, genotype, and phase single nucleotide variants (SNV) (35).

Phased SNVs were used to assign each read to a haplotype using MsPAC. Reads from each

haplotype and gene were clustered using CD-HIT using a 100% identity clustering threshold

parameter, and a single representative read from the largest cluster was BLASTed to the

ImMunoGeneTics (IMGT) reference database (downloaded on 2/21/22) to determine the constant

region allele usage (36, 37). The representative read was selected based on 100% identity to all

other sequences in that cluster. Constant region alleles observed by FLAIRR-seq were

designated as "novel" if there was no match to the IMGT database or "novel, extended" if a match

was detected to a partial allele found in IMGT, but the IMGT allele was a substring of the FLAIRR-

seq identified allele. In cases for which FLAIRR-seq resulted in ambiguous allele calls,

visualization of IGenotyper DNA haplotype assemblies and FLAIRR-seq data were performed

using the Integrated Genomics Viewer (IGV v2.12.3) software to manually confirm allele

assignment based on IGenotyper ground truth.

Results

Validation and benchmarking of the FLAIRR-seq method

Current methods for AIRR-Seq utilize targeted amplification of the variable region and, in

some cases, a small portion of the first constant region exon (CH1), in conjunction with short-read

sequencing to characterize IG repertoires. However, this minimal examination of CH1 is primarily

used to define IG isotype groups. No current method defines both the heavy chain variable and

constant regions with allele-level resolution allowing for subisotype classification. To address

these technical limitations, we developed the FLAIRR-Seq method (Figure 1C), a targeted 5'

RACE approach combined with SMRT sequencing to generate highly accurate, near full-length

IgG (~1500 bp) and/or IgM (2000 bp) sequences, allowing for direct, simultaneous analysis of the

heavy chain variable and constant regions, including gene/allele identification for V, D, J and C

segments, and isotype- and subisotype-specific repertoire profiling.

To evaluate and validate the capabilities of FLAIRR-seq, matched FLAIRR-seq and AIRR-

seq analyses were performed on ten healthy donor PBMC samples. FLAIRR-seq data were

filtered from the initial HiFi reads (>Q20) to include only >Q40 reads. The average read quality of

these filtered reads was >Q50 (99.999%), with a pass filter rate ranging from 88%-93% of total

reads. AIRR-seq FASTQ reads were filtered for those with an average quality of >Q30 (99.9%).

These filtered reads were used as input into the Immcantation suite, specifically the pREST-O

and Change-O tools, for V, D, and J gene assignment, and different repertoire feature analyses,

such as identification of clones, extent of somatic hypermutation, and evaluation of CDR3 lengths.

As shown in **Table 1,** fewer overall FLAIRR-seq reads were used as input into the Immcantation

analyses, after required filtration and read assembly steps (which were not needed for the high-

quality single-molecule FLAIRR-seq reads). FLAIRR-seq resulted in comparable or, in many

cases, increased number of unique V, D, and J gene allele pairs (VDJ), clones, and CDR3

sequences identified compared to the matched AIRR-seq-derived samples in both the IgG and

IgM repertoires (schematized in Figure 2A). These basic sequencing and initial analysis metrics

demonstrated that FLAIRR-seq could produce high-quality variable region data for detailed IG

repertoire analyses. Furthermore, the ability to utilize Immcantation shows that current AIRR-seq

tools are capable of analyzing FLAIRR-seq data. While input RNA mass used for FLAIRR-seq

was often more than for AIRR-seq, ongoing and future optimization of the method is aimed at

reducing RNA input. Furthermore, a comparative cost analysis was performed. To obtain the

closest metric to a direct comparison, we calculated the cost per "actionable read", defined as the

read number per sample and per method after read and length quality filtration and assembly, but

prior to cluster consensus, performed in the pRESTO pipeline. This method was used to represent

the total unique single molecule or assembled variable region templates captured by either the

FLAIRR-seq or AIRR-seq methods that passed all necessary quality control criteria for

downstream annotation and analyses irrespective of biologic repertoire diversity or clonality,

removing technology/platform biases. Lastly, to remove the impact of pooling differences, we

used this "per actionable read" cost to calculate the cost for the generation of 15,000 "actionable

reads" as our standard price. For AIRR-seq this cost was $25.50 per sample, whereas for

FLAIRR-seq, this cost was $33.57 per sample. These costs reflect needed reagents and

consumables only, assume instrumentation access and do not include labor. Future optimization

for FLAIRR-seq will include integrating a multiplexed array sequencing (MAS) step to concatenate

reads and enhance overall depth of sequencing and multiplexing capacity per pool, resulting in

decreasing costs per sample (38).

While optimizing FLAIRR-seq sample preparation, we examined whether upstream

isolation of B cells before FLAIRR-seq molecular preparation would enhance the ability to detect

unique variable region gene usage. To do this, aliquots of PBMC (n=4) were split into two groups

for RNA extraction: (i) RNA derived from bulk PBMC, and (ii) RNA isolated from purified pan B-

cells, followed by FLAIRR-seq preparation, SMRT sequencing, and Immcantation analysis of both

groups. V gene usage correlations between groups are shown in Figure 2B and demonstrate a

significant association (p-values ranging from 0.033 to $4.1e^{-16}$) strongly supporting the conclusion

that B cell isolation before RNA extraction was not necessary to achieve comparable gene usage

metrics. The limited differences that were observed could be explained by template sampling

differences between the two experiments. Due to the strong associations observed and the ease

of processing PBMC in bulk, we moved forward with bulk-derived PBMC material for the

remainder of our analyses.

We established FLAIRR-seq performance by comparing its output to the commonly used

5' RACE AIRR-seq method. 5' RACE AIRR-seq was chosen as it provides the most extensive

AIRR resolution of standard methods in the field, capturing both the variable region and a small

portion of the constant region CH1 exon, allowing for isotype differentiation. Following matched

preparation of both FLAIRR-seq and AIRR-seq libraries from healthy donor PBMC samples

(n=10), we compared many variable region features to benchmark FLAIRR-seq performance.

First, we evaluated variable region gene usage frequencies. We observed strongly significant

associations (p-values ranging from 0.033 to $<2.2e^{-16}$) between V, D and J gene usage for both

IgG and IgM isotypes (Figure 2C), indicating that FLAIRR-seq accurately and comparably

resolves variable region genes. To note, J genes showed less significance (larger p-value) across

all comparisons due to the relatively few genes that make up the J gene family compared to the

more diverse V and D families. We next investigated the performance of both methods in terms

of resolving CDR3 lengths (Figure 2C) and somatic hypermutation frequencies contained in

productive sequences (Figure 2E), which are often used as measures of evaluating B cell affinity

maturation. FLAIRR-seq and AIRR-seq performed robustly, with significantly longer CDR3

regions detected with FLAIRR-seq across all samples in both the IgM and IgG groups. The

characterization of unusually long CDR3 regions (> 40 nt) in the IgG sequences with FLAIRR-seq

that were not observed in the matched AIRR-seq data, could be due the higher contiguity and

quality afforded from longer read length when using FLAIRR-seq. Single-molecule resolution of

near full-length heavy chain transcripts would allow for the inclusion of unusually long variable

regions that may have been missed by short-read 2x 300 bp paired-end sequencing strategies if

reads 1 and 2 did not maintain enough overlap in this region to assemble. Overall, these data

demonstrate that FLAIRR-seq achieves unambiguous and comparable, if not improved,

resolution of the variable region genes and features, and improved in terms of CDR3 length

resolution, and can be used with publicly available AIRR-seq analyses tools to provide commonly

used metrics for repertoire analysis.

Others have recognized the power of long-read sequencing to resolve B cell repertoires

but have used bulk Iso-Seq methods to examine full-length transcripts from isolated B cells (39).

The IsoSeq method captures full-length transcripts expressing a poly(A) tail without bias through

oligo dT-based priming. The trade-offs of this approach are throughput, depth, and cost, as

IsoSeq processing generates a complete transcriptome per sample without enrichment of heavy

chain sequences, which then need to be filtered out and analyzed, resulting in a considerable

amount of non-repertoire data to be discarded. To investigate whether the untargeted

transcriptome-wide IsoSeq method would resolve a qualitatively different repertoire than FLAIRR-

seq, which would have indicated FLAIRR-seq-driven primer bias, we performed matched Iso-Seq

and FLAIRR-seq on purified B-cell derived RNA. Variable gene usage frequencies between

IsoSeq- and FLAIRR-seq-derived V, D, and J calls were examined by Spearman ranked

correlation and showed highly significant associations between methods (V genes: r^2 = 0.94,

p=2.2e^{-16}; D genes: r^2 = 0.92, p =1.4e^{-13}; J genes: r^2 = 1.0, p = 0.0028). These data strongly

suggest that FLAIRR-seq has very limited to no primer-driven bias compared to whole

transcriptome data. Collectively, this benchmarking dataset confirmed that FLAIRR-seq is robust

and accurate, providing complete characterization of IG variable genes and associated commonly

used IG repertoire metrics, with limited increases in per sample cost.

IGenotyper and FLAIRR-seq provide constant region gene and allele identification

Although confirmed that FLAIRR-seq successfully characterized IG variable region

transcripts, the true value added by FLAIRR-seq is the novel capabilities it provides in resolving

the constant region, including gene and allele usage, subisotype identification, and phasing of

variable and constant regions for comprehensive repertoire analysis. To evaluate the capabilities

and accuracy of constant region gene and allele identification with FLAIRR-seq, we first

established a ground truth dataset of IGHC alleles for all 10 samples by targeted sequencing of

the germline IGH locus (Figure 3A) using IGenotyper, as previously described (32). Constant

region alleles called by IGenotyper were assigned to one of three categories, schematized in

Figure 3B: (i) "exact match" - known constant region alleles found in the IMGT database; (ii)

"novel not in IMGT" – a constant region allele not found in the IMGT database; or (iii) "extended"

– constant region alleles that appeared to match partial alleles in the IMGT database, but

IGenotyper data extended the sequence beyond what has been previously characterized.

IGenotyper identified a total of 32 alleles across all individuals and classifications, as schematized

in Figure 3C. Furthermore, we discovered IGHG4 gene duplications in in 4/10 samples.

Identification of these constant region gene duplications further highlights the structural

complexity and diversity of IGH.

Given the extensive additional constant region gene content resolved by IGenotyper

compared to IMGT, we used the IGenotyper-derived IGHC gene database as the ground-truth for

evaluation of the capabilities of FLAIRR-seq data to identify and resolve constant region alleles.

FLAIRR-seq data alone identified a total of 23/32 (72%) alleles at 100% identity with no additional

alleles (false-positives) detected as compared to the IGenotyper ground-truth database (Figures

3D and E). Due to the internal placement of the CH3- and CH4-specific primers used to target IG

transcripts, the entirety of these exons were not resolved and, in some cases, resulted in slightly

truncated transcripts that resulted in ambiguous allele calls when aligned to our ground-truth

IGenotyper constant region database (Figure 3C). For example, in IGHG1 alleles the complete

sequence of the CH3 exon was often necessary to distinguish between two closely related alleles,

as well as determine if an individual was homozygous or heterozygous at these positions. In these

cases, FLAIRR-seq data were able to substantially narrow the field of possible alleles at that

position in an individual, but the IGenotyper data was used to identify the exact allele. Using

FLAIRR-seq data alone also limited phasing capabilities for some constant region genes due to

missing allele-defining SNVs in the truncated transcripts resulting in false homozygous calls,

primarily in IGHG1 genes (30%). Further improvements to FLAIRR-seq will aim to iterate in primer

design to capture the remaining 3' region of CH3 to increase genotype capabilities of FLAIRR-

seq as a stand-alone method.

In total, IGenotyper data identified 32 constant region alleles across the 10 healthy donors

examined, including 11 alleles (34%) that were completely novel and 17 (53%) alleles that were

extended compared to those that were curated in IMGT at the time of publication. FLAIRR-seq

was also able to resolve 23/32 (72%) these alleles when assessed as a stand-alone genotyping

method. Given the relatively small size of this proof-of-concept healthy donor cohort, the

identification of 28 (87%) novel or novel-extended alleles here underscores the extensive

polymorphism in this region, despite a historical assumption that these genes are "constant" and

reflects the paucity of information regarding this locus in existing immunogenomics databases.

FLAIRR-seq enables subisotype-specific repertoire analyses

Once subisotypes were identified, we performed subisotype-specific repertoire analyses

(Figure 4). We initially examined V gene family usage as a function of IgG subisotype across all

10 healthy donors. From this analysis we observed V gene family usage diversity within each

subisotype across all seven V gene families (Figure 4A). For example, significantly disparate V

family gene usage patterns are seen across subisotypes for the IGHV1 genes where IGHV1

genes are most highly expressed in the IgG4-specific repertoire and are seen the least within the

505 IgG2-specific repertoire (ANOVA p=1.3e^{-07}). We chose to investigate subisotype-specific V, D,

506 and J gene usage for samples 1008 and 602c in detail as a demonstration of this capability.

507 Characterization of constant region genes and alleles within FLAIRR-seq data allowed for the

508 relative frequency of heavy chain transcripts per isotype class, subisotype, and subisotype-

509 specific allele to be assigned. Once classified by subisotype, detailed analysis of variable region

510 repertoire per subisotype-specific allele was possible (Figure 4B). For samples 1008 and 602c,

511 we observed that IGHV3 genes were highly expressed across all subisotypes, whereas IGHV1

512 genes are relatively elevated in the IgG4 population compared to the IgG1, IgG2, and IgG3

513 compartments (Figure 4C and D). However, samples 1008 and 602c differ in their expression of

514 IGHV1 family genes, which are expressed at relatively higher levels in sample 602c compared to

515 sample 1008 (Figure 4C and D). In the context of standard AIRR-seq analyses, we would not be

516 able to identify allele-resolved V gene usage or enrichment within subisotype populations, which

517 could be essential for linking subisotype functionality with VDJ clonotypes.

518 **FLAIRR-seq identifies subisotype-specific clonal expansion and CSR in longitudinal**

519 **samples.**

520  It was critical to demonstrate the feasibility of performing FLAIRR-seq from clinically

521 relevant sample types, such as whole blood. To do this, we evaluated FLAIRR-seq performance

522 in a small subset of samples collected in April 2020 from hospitalized COVID-19 patients (n=5)

523 from the Mount Sinai COVID-19 biobank cohort at the Icahn School of Medicine at Mount Sinai

524 (25). RNA was directly extracted from whole blood, as outlined in the Materials and Methods, and

525 robust FLAIRR-seq amplification of targeted IgG and IgM repertoires was observed from all

526 samples (data not shown).

527  IG responses are highly dynamic, with specific IG clones expanding upon activation by

528 cognate antigen. We were interested to know if these dynamics could be captured by FLAIRR-

529 seq, given the capability to identify clonotypes with the subisotype resolved. We had the

opportunity to evaluate FLAIRR-seq resolved repertoires over time in four samples collected from

one individual over their >13-day hospitalization. Patient PICR7356 was hospitalized with severe

COVID-19 disease, and blood draws were taken on days 1, 4, 8, and 13 post-hospitalization

(Figure 5A) and analyzed with FLAIRR-seq across all time points. After initial FLAIRR-seq

processing and analysis, we defined unique clones using SCOPer, which clusters groups based

on CDR3 similarity and mutations in V and J genes and provided clone counts across the four

timepoints examined. Overall, we observed that most clones were expressed within the IgG1

compartment, but longitudinal clone distribution across subisotypes fluctuated over time.

Specifically, the IgG3-specific repertoire expanded from day 1 to day 4, but then contracted in

overall frequency from day 8 to day 13 (Figure 5B). To assess clonal diversity within each

subisotype repertoire across time, we calculated the Simpson's diversity index (q=2) using

Alakazam (27). All subisotype-specific repertoires became less diverse from day 1 to day 4,

suggesting clonal expansion across the IgG repertoire (Figure 5C). To note, IgG4 was not

included in diversity calculations because IgG4 would have required higher sequencing depth to

ascertain diversity in this individual, given the overall lower expression of IgG4 transcripts in this

individual. Subisotype-specific IG repertoire polarity was also calculated by counting the number

of clones needed to represent 80% of the total repertoire and then expressing this number as a

percentage of the total (Figure 5D). In this metric, the lower the clone number that is needed to

represent 80% of the repertoire, the higher the extent of clonal expansion observed. Polarity

values confirmed those seen with the diversity index, demonstrating an increase of clonal

expansion (i.e., decreased polarity) at day 4 across IgG1, IgG2, and IgG3 compartments, which

returned to baseline at later timepoints. Notably, although expressed at low levels, polarity of the

IgG4-specific repertoire was examined and was an exception to this trend, demonstrating

indications of clonal expansion at day 8 and day 13.

CSR is a critical immunological phenomenon allowing for the replacement of constant

region genes with others and, therefore, altering associated effector functions while maintaining

antigen-specific variable regions. Given the ability of FLAIRR-seq to resolve variable clonotypes

and constant region gene usage, we evaluated whether this method could identify signatures of

CSR in a longitudinal repertoire. To do this, we examined clones shared across timepoints and

asked how these clones were distributed within each subisotype-specific repertoire over time. In

total, 34 unique clonal lineages were shared across all timepoints and could be examined for

signatures of CSR. As a proof-of-concept, we examined the relative frequency of the dominant

clone 13611 within each subisotype repertoire across all time points (Figure 5E). To establish that

we were examining possible CSR, we confirmed that all 13611 clonal lineage sequences shared

the same variable region alleles. On day 1 post-hospitalization, clone 13611 sequences were

found equally distributed within the IgM and IgG2 repertoire, indicating that these B cells had not

undergone complete maturation. By day 4, we see clear evidence of B cell maturation and CSR

by the dominance of IgG2 usage by clone 13611 sequences. Residual low-level IgM sequences

rebounded on day 8, while also showing low-level CSR from IgM to IgG1 at that same time point.

To visualize these dynamics, we built a phylogeny using the Dowser pipeline via Immcantation.

Highlighted in the red box on the phylogenetic tree shown in Figure 5F, we observe a clustered

IgG1 transcript emerging from IgM-specific clade, indicating CSR from IgM to IgG1. Together,

these data provide demonstrative proof-of-concept evidence that FLAIRR-seq profiling performs

robustly on clinical samples, including RNA directly extracted from whole blood. In addition, these

data provide novel repertoire resolution than what would have been provided with standard AIRR-

seq methods, including analysis of subisotype-specific repertories, evaluation of clonal

expansion, and characterization of CSR in the IgG and IgM repertoires.

Discussion

Here we present the development, validation, and application of FLAIRR-seq, a novel

method to resolve near full Ig IGH transcripts from bulk PBMC-, isolated B cell- and whole blood-

derived total RNA. FLAIRR-seq enables highly accurate, simultaneous resolution of Ig variable

and constant regions and suggests that constant region polymorphism is far more extensive than

previously assumed. FLAIRR-seq performed equivalent to or with increased resolution compared

to existing standard 5'RACE AIRR-seq methods when resolving variable gene calls, CDR3

lengths, and somatic hypermutation signatures, suggesting that our CH3/CH4 targeting strategies

did not compromise variable region characterization while simultaneously adding the capability to

resolve extensive constant region genomic variation. Little to no primer bias was observed when

compared to IG repertoire profiling from total mRNA Iso-seq methods. FLAIRR-seq provides the

novel ability to use constant region gene usage to identify IG subisotypes and genotype heavy

chain transcripts, linking these data back to evaluate subisotype-specific repertoires, clonal

expansion, and CSR. Underscoring the underappreciated extent of constant region genomic

variation, our FLAIRR-seq profiling of a restricted cohort of only 10 individuals from relatively

homogenous backgrounds still identified 4 and 6 completely novel constant region alleles in IgM

and IgG, respectively, and extended an additional 16 alleles beyond which had been available in

the IMGT database.

The unique capabilities of FLAIRR-seq will allow for novel examination of IG repertoires,

including the characterization of variable gene usage and clonotype distribution within unique

subisotype subsets. This perspective has the potential to provide key insights into dynamic IG

responses in diseases known to be mediated by subisotype-specific IG or have skewed

subisotype distribution as predictive markers of disease, including Myasthenia gravis (mediated

by pathogenic autoantibodies across subisotypes that give rise to varied disease pathologies),

Acute Rheumatic Fever (associated with elevated IgG3), and melanoma (skewing towards IgG4

in late-stage disease thought to be indicative of tolerogenic responses and poor prognosis)(12-

14). These subisotype-specific repertoire profiling approaches may be the first step toward

identification of unique IG clones that mediate disease pathogenicity or serve as high resolution

biomarkers to disease progression, as well as open the door for potential functional experiments

on subisotype clones of interest, including examining the functional impact of the novel constant

region alleles identified here. Expanded population-based FLAIRR-seq profiling and curation of

novel constant region alleles, particularly in conjunction with IGenotyper targeted genomic

assembly efforts in IGH, will be a significant first step in defining the full extent of variation in a

region too long assumed to be relatively invariant.

The Fc region is known to be critical for modulating differential IG effector functions. These

differential functionalities are currently understood to be regulated by differential posttranslational

modifications, such as variable glycosylation (40-43). Future FLAIRR-seq profiling will be a

valuable tool to investigate how genomic variation across constant region genes impacts residue

usage and resultant Fc receptor binding, signaling potential, crosslinking, and potential for

posttranslational modification, all of which would be expected to alter downstream effector

function potential, such as ADCC, ADCP, and complement fixation.

We further demonstrate that FLAIRR-seq can effectively examine clonal expansion and

CSR in longitudinal samples, demonstrating the feasibility of using FLAIRR-seq to resolve IG

repertoire dynamics. This increased resolution will further our understanding of IG repertoire

evolution in the transition of acute to chronic disease states, many of which are associated with

overall IgG subisotype distribution changes that are thought to reflect the inflammatory milieu (14,

44). One example is advanced melanoma, where late-stage disease is characterized by elevated

IgG4 compared to IgG1, which is believed to reflect a more tolerizing, pro-tumor environment (12).

FLAIRR-seq examination of these samples may identify specific IG distribution patterns that could

act as biomarkers of disease progression. Moving forward it is critical to account for all variability

within the IG repertoire for the most comprehensive understanding of repertoire dynamics and

the myriad factors impacting antibody effector function. Together, the data presented here

demonstrate that the FLAIRR-seq method provides a comprehensive characterization of allele-

resolved IgG and IgM repertoires, detailing variable region gene usage and measurements of

maturation, isotype and subisotype identification, and the unappreciated extent of constant region

variation, which will be necessary to fully appreciate the impact of IG genomic variation in health

and disease.

Figure Legends:

Figure 1. Overview of IG structure and FLAIRR-seq molecular method. (A) Schematic

representation of the IGH locus, heavy chain transcript structure, and functional IG protein. (B)

Comparative coverage across the heavy chain transcript of commonly used AIRR-seq methods

compared with FLAIRR-seq. (C) FLAIRR-seq molecular pipeline: RNA (brown) was converted to

first-strand cDNA (red) using the 5' RACE method, incorporating a 5' TSO-UMI (pink) via template

switch. Second strand amplification specifically targeted IgG and IgM molecules through priming

of the 5' TSO-UMI and the 3' constant region IGH exon 3 (CH3) for IgG, or CH4 for IgM. A 16bp

barcode was incorporated into the 3' CH3/CH4 primers to enable sample multiplexing post-

amplification. (D) Fragment Analyzer traces of final FLAIRR-seq products at ~1500bp (IgG) and

~2000bp (IgM).

Figure 2. FLAIRR-seq shows robust characterization of IG variable regions. (A) Schematic

representation analyses metrics evaluated including clones, unique VDJ, CDR3 length, and

somatic hypermutation. (B) Spearman ranked correlations and p-values of V, D, and J gene usage

frequencies identified by FLAIRR-seq performed on matched total PBMC and purified B cells. (C)

Heatmap of Spearman ranked correlations and p-values of V, D and J gene usage frequencies

between FLAIRR-seq and AIRR-seq processed samples. (D) Boxplots of CDR3 length defined

by FLAIRR-seq or AIRR-seq analysis in both IgM- and IgG-specific repertoires. (E) Boxplots of

somatic hypermutation frequencies defined by FLAIRR-seq or AIRR-seq in IgM- and IgG-specific

repertoires. (F) Spearman ranked correlation of V, D, and J gene usage frequencies between

FLAIRR-seq-based and Iso-Seq-based repertoire profiling. Significant differences between

FLAIRR-seq and AIRR-seq data indicated by * ($p < 0.05$) or ** ($p < 0.01$)

Figure 3. FLAIRR-seq provides novel constant region resolution for allelic discovery. (A)

Overview of experimental design and pipeline overviews of genotyping by IGenotyper (gDNA)

and FLAIRR-seq (RNA). (B) Schematic depicting constant region allele types identified by

IGenotyper, including identification of (i) exact match to IMGT, (ii) novel, not in IMGT and (iii)

extended. (C) Pie chart and stacked bar graph representing total alleles and their types (exact

match, novel, not in IMGT, or extended) per IGHC gene as identified by IGenotyper. * indicates

additional allele found due to IGHG4 duplicaton (D) Table summarizing of novel (pink) and

extended (diamond) alleles resolved in FLAIRR-seq data alone. E) Pie chart and stacked bar

graph of total alleles and their types (exact match, novel, not in IMGT, extended, or ambiguous)

resolved by FLAIRR-seq data alone.

Figure 4. FLAIRR-seq resolves subisotype specific repertoire diversity. (A) Boxplots of V

gene family usage frequencies in expressed IgG subisotypes across all ten individuals. (B) Bar

plots showing distribution of unique VDJ sequences across isotypes, subisotypes, and subisotype

alleles in two representative samples, 1008 and 602c, characterized by FLAIRR-seq. (C) Circos

plots showing V family gene usage frequency within each subisotype allele fraction for samples

1008 and 602c. (D) Table showing percentage of V gene family usage within each constant region

allele as represented in the circus plots in (C).

Figure 5. FLAIRR-seq resolves subisotype-specific clonal expansion and evidence of CSR

in a patient hospitalized for COVID-19. (A) Overview of experimental design: whole blood-

derived RNA was collected on days 1, 4, 8 and 13 post-hospitalization and used for FLAIRR-seq

profiling. (B) Bar plot showing percentage of unique clones per subisotype across all timepoints.

(C) Simpson's diversity index (q=2) for all clones in each subisotype across four timepoints; IgG4

not included due to low sequence number that did not pass filtering criteria. Decreased diversity

indicates increased clonal expansion. (D) Polarization, or number clones needed to comprise

80% of the repertoire, reported as percentage of total subisotype-specific repertoires across time.

Decreased polarity indicates increased clonal expansion. (E) Distribution of clone 13611 across

isotypes and subisotypes over time, suggesting CSR of this clone. (F) Phylogenetic tree of clone

13611 showing emergence of CSR transcript over time (red box).

References:

1. Schroeder, H. W., and L. Cavacini. 2010. Structure and function of immunoglobulins. *Journal of Allergy and Clinical Immunology* 125: S41-S52.

2. Janda, A., A. Bowen, N. S. Greenspan, and A. Casadevall. 2016. Ig Constant Region Effects on Variable Region Structure and Function. *Front Microbiol* 7: 22.

3. Nakano, T., M. Matsui, I. Inoue, T. Awata, S. Katayama, and T. Murakoshi. 2011. Free immunoglobulin light chain: Its biology and implications in diseases. *Clinica Chimica Acta* 412: 843-849.

4. Lu, L. L., T. J. Suscovich, S. M. Fortune, and G. Alter. 2018. Beyond binding: antibody effector functions in infectious diseases. *Nature Reviews Immunology* 18: 46-61.

5. Vidarsson, G., G. Dekkers, and T. Rispens. 2014. IgG subclasses and allotypes: from structure to effector functions. *Front Immunol* 5: 520-520.

6. Greiff, V., C. R. Weber, J. Palme, U. Bodenhofer, E. Miho, U. Menzel, and S. T. Reddy. 2017. Learning the High-Dimensional Immunogenomic Features That Predict Public and Private Antibody Repertoires. *The Journal of Immunology* 199: 2985.

7. Tonegawa, S. 1983. Somatic generation of antibody diversity. *Nature* 302: 575-581.

8. Nishana, M., and S. C. Raghavan. 2012. Role of recombination activating genes in the generation of antigen receptor diversity and beyond. *Immunology* 137: 271-281.

9. Tong, P., A. Granato, T. Zuo, N. Chaudhary, A. Zuiani, S. S. Han, R. Donthula, A. Shrestha, D. Sen, J. M. Magee, M. P. Gallagher, C. E. van der Poel, M. C. Carroll, and D. R. Wesemann. 2017. IgH isotype-specific B cell receptor expression influences B cell fate. *Proc Natl Acad Sci U S A* 114: E8411-e8420.

10. Noviski, M., J. L. Mueller, A. Satterthwaite, L. A. Garrett-Sinha, F. Brombacher, and J. Zikherman. 2018. IgM and IgD B cell receptors differentially respond to endogenous antigens and control B cell fate. *eLife* 7: e35074.

11.   Stavnezer, J., J. E. Guikema, and C. E. Schrader. 2008. Mechanism and regulation of

class switch recombination. *Annu Rev Immunol* 26: 261-292.

12.   Karagiannis, P., A. E. Gilbert, D. H. Josephs, N. Ali, T. Dodev, L. Saul, I. Correa, L.

Roberts, E. Beddowes, A. Koers, C. Hobbs, S. Ferreira, J. L. Geh, C. Healy, M. Harries,

K. M. Acland, P. J. Blower, T. Mitchell, D. J. Fear, J. F. Spicer, K. E. Lacy, F. O. Nestle,

and S. N. Karagiannis. 2013. IgG4 subclass antibodies impair antitumor immunity in

melanoma. *J Clin Invest* 123: 1457-1474.

13.   Chung, A. W., T. K. Ho, P. Hanson-Manful, S. Tritscheller, J. M. Raynes, A. L. Whitcombe,

M. L. Tay, R. McGregor, N. Lorenz, J. R. Oliver, J. K. Gurney, C. G. Print, N. J. Wilson,

W. J. Martin, D. A. Williamson, M. G. Baker, and N. J. Moreland. 2020. Systems

immunology reveals a linked IgG3-C4 response in patients with acute rheumatic fever.

Immunol Cell Biol 98: 12-21.

14.   Vander Heiden, J. A., P. Stathopoulos, J. Q. Zhou, L. Chen, T. J. Gilbert, C. R. Bolen, R.

J. Barohn, M. M. Dimachkie, E. Ciafaloni, T. J. Broering, F. Vigneault, R. J. Nowak, S. H.

Kleinstein, and K. C. O'Connor. 2017. Dysregulation of B Cell Repertoire Formation in

Myasthenia Gravis Patients Revealed through Deep Sequencing. *J Immunol* 198: 1460-

1473.

15.   Huijbers, M. G., L. A. Querol, E. H. Niks, J. J. Plomp, S. M. van der Maarel, F. Graus, J.

Dalmau, I. Illa, and J. J. Verschuuren. 2015. The expanding field of IgG4-mediated

neurological autoimmune disorders. *European Journal of Neurology* 22: 1151-1161.

16.   Polonelli, L., J. Pontón, N. Elguezabal, M. D. Moragues, C. Casoli, E. Pilotti, P. Ronzi, A.

S. Dobroff, E. G. Rodrigues, M. A. Juliano, D. L. Maffei, W. Magliani, S. Conti, and L. R.

Travassos. 2008. Antibody complementarity-determining regions (CDRs) can display

differential antimicrobial, antiviral and antitumor activities. *PLoS One* 3: e2371.

17.   Liu, H., W. Pan, C. Tang, Y. Tang, H. Wu, A. Yoshimura, Y. Deng, N. He, and S. Li. 2021.

752   The methods and advances of adaptive immune receptors repertoire sequencing.

753   *Theranostics* 11: 8945-8963.

754   18.   Trück, J., A. Eugster, P. Barennes, C. M. Tipton, E. T. Luning Prak, D. Bagnara, C. Soto,

755   J. S. Sherkow, A. S. Payne, M.-P. Lefranc, A. Farmer, A. C. The, M. Bostick, and E.

756   Mariotti-Ferrandiz. 2021. Biological controls for standardization and interpretation of

757   adaptive immune receptor repertoire profiling. *eLife* 10: e66274.

758   19.   Horns, F., C. Vollmers, D. Croote, S. F. Mackey, G. E. Swan, C. L. Dekker, M. M. Davis,

759   and S. R. Quake. 2016. Lineage tracing of human B cells reveals the in vivo landscape of

760   human antibody class switching. *eLife* 5: e16578.

761   20.   Calonga-Solís, V., D. Malheiros, M. H. Beltrame, L. d. B. Vargas, R. M. Dourado, H. C.

762   Issler, R. Wassem, M. L. Petzl-Erler, and D. G. Augusto. 2019. Unveiling the Diversity of

763   Immunoglobulin Heavy Constant Gamma (IGHG) Gene Segments in Brazilian

764   Populations Reveals 28 Novel Alleles and Evidence of Gene Conversion and Natural

765   Selection. *Front Immunol* 10.

766   21.   Jonsson, S., G. Sveinbjornsson, A. L. de Lapuente Portilla, B. Swaminathan, R. Plomp,

767   G. Dekkers, R. Ajore, M. Ali, A. E. H. Bentlage, E. Elmér, G. I. Eyjolfsson, S. A.

768   Gudjonsson, U. Gullberg, A. Gylfason, B. V. Halldorsson, M. Hansson, H. Holm, Å.

769   Johansson, E. Johnsson, A. Jonasdottir, B. R. Ludviksson, A. Oddsson, I. Olafsson, S.

770   Olafsson, O. Sigurdardottir, A. Sigurdsson, L. Stefansdottir, G. Masson, P. Sulem, M.

771   Wuhrer, A.-K. Wihlborg, G. Thorleifsson, D. F. Gudbjartsson, U. Thorsteinsdottir, G.

772   Vidarsson, I. Jonsdottir, B. Nilsson, and K. Stefansson. 2017. Identification of sequence

773   variants influencing immunoglobulin levels. *Nature Genetics* 49: 1182-1191.

774   22.   Buck, D., E. Albrecht, M. Aslam, A. Goris, N. Hauenstein, A. Jochim, S. Cepok, V.

775   Grummel, B. Dubois, A. Berthele, P. Lichtner, C. Gieger, J. Winkelmann, and B. Hemmer.

2013. Genetic variants in the immunoglobulin heavy chain locus are associated with the

IgG index in multiple sclerosis. *Ann Neurol* 73: 86-94.

23.   van Erp, E. A., W. Luytjes, G. Ferwerda, and P. B. van Kasteren. 2019. Fc-Mediated

Antibody Effector Functions During Respiratory Syncytial Virus Infection and Disease.

Front Immunol 10.

24.   Jefferis, R., J. Lund, and J. D. Pound. 1998. IgG-Fc-mediated effector functions: molecular

definition of interaction sites for effector ligands and the role of glycosylation.

Immunological Reviews 163: 59-76.

25.   Charney, A. W., N. W. Simons, K. Mouskas, L. Lepow, E. Cheng, J. Le Berichel, C. Chang,

R. Marvin, D. M. Del Valle, S. Calorossi, A. Lansky, L. Walker, M. Patel, H. Xie, N. Yi, A.

Yu, G. Kang, A. Mendoza, L. E. Liharska, E. Moya, M. Hartnett, S. Hatem, L. Wilkins, M.

Eaton, H. Jamal, K. Tuballes, S. T. Chen, A. Tabachnikova, J. Chung, J. Harris, C.

Batchelor, J. Lacunza, M. Yishak, K. Argueta, N. Karekar, B. Lee, G. Kelly, D. Geanon, D.

Handler, J. Leech, H. Stefanos, T. Dawson, I. Scott, N. Francoeur, J. S. Johnson, A. Vaid,

B. S. Glicksberg, G. N. Nadkarni, E. E. Schadt, B. D. Gelb, A. Rahman, R. Sebra, G.

Martin, C. Agashe, P. Agrawal, A. Akyatan, K. Alesso-Carra, E. Alibo, K. Alvarez, A.

Amabile, S. Ascolillo, R. Bailey, P. Begani, P. B. Correra, S.-A. Brown, M. Buckup, L.

Burka, L. Cambron, G. Carrara, S. Chang, J. Chien, M. Chowdhury, C. C. Bozkus, P.

Comella, D. Cosgrove, F. Cossarini, L. Cotter, A. Dave, B. Dayal, M. Dhainaut, R.

Dornfeld, K. Dul, N. Eber, C. Elaiho, F. Fabris, J. Faith, D. Falci, S. Feng, B. Fennessy, M.

Fernandes, S. Gangadharan, J. Grabowska, G. Gyimesi, M. Hamdani, M. Herbinet, E.

Herrera, A. Hochman, G. E. Hoffman, J. Hook, L. Horta, E. Humblin, S. Karim, J. Kim, D.

Lebovitch, G. Lee, G. H. Lee, J. Lee, M. Leventhal, K. Lindblad, A. Livanos, R. Machado,

Z. Mahmood, K. Mar, S. Maskey, P. Matthews, K. Meckel, S. Mehandru, C. Mercedes, D.

Meyer, G. Mollaoglu, S. Morris, K. Nie, M. Nisenholtz, G. Ofori-Amanfo, K. Onel, M.

Ounadjela, V. Patel, C. Pruitt, S. Rathi, J. Redes, I. Reyes-Torres, A. Rodrigues, A.

Rodriguez, V. Roudko, E. Ruiz, P. Scalzo, P. Silva, A. S. Schanoski, M. Straw, S.

Tabachnikova, C. Teague, B. Upadhyaya, V. Van Der Heide, N. Vaninov, D. Wacker, H.

Walsh, C. M. Wilk, J. Wilson, K. M. Wilson, L. Xue, N.-a. Yeboah, S. Young, N. Zaks, R.

Zha, T. Marron, N. Beckmann, S. Kim-Schulze, S. Gnjatic, M. Merad, and C.-B. T. The

Mount Sinai. 2020. Sampling the host response to SARS-CoV-2 in hospitals under siege.

Nature Medicine 26: 1157-1158.

26.   Woolley, C., J. Chariker, E. Rouchka, E. Ford, E. Hudson, S. Waigel, M. Smith, and T.

Mitchell. 2022. Reference long-read isoform-aware transcriptomes of four human

peripheral blood lymphocyte subsets

. G3: Genes, Genomes, Genetics

.

27.   Gupta, N. T., J. A. Vander Heiden, M. Uduman, D. Gadala-Maria, G. Yaari, and S. H.

Kleinstein. 2015. Change-O: a toolkit for analyzing large-scale B cell immunoglobulin

repertoire sequencing data. *Bioinformatics* 31: 3356-3358.

28.   Vander Heiden, J. A., G. Yaari, M. Uduman, J. N. H. Stern, K. C. O'Connor, D. A. Hafler,

F. Vigneault, and S. H. Kleinstein. 2014. pRESTO: a toolkit for processing high-throughput

sequencing raw reads of lymphocyte receptor repertoires. *Bioinformatics* 30: 1930-1932.

29.   Nouri, N., and S. H. Kleinstein. 2018. A spectral clustering-based method for identifying

clones from high-throughput B cell repertoire sequencing data. *Bioinformatics* 34: i341-

i349.

30.   Nouri, N., and S. H. Kleinstein. 2020. Somatic hypermutation analysis for improved

identification of B cell clonal families from next-generation sequencing data. *PLOS*

Computational Biology 16: e1007977.

31.   Hoehn, K. B., O. G. Pybus, and S. H. Kleinstein. 2022. Phylogenetic analysis of migration,

differentiation, and class switching in B cells. *PLOS Computational Biology* 18: e1009885.

32. Rodriguez, O. L., W. S. Gibson, T. Parks, M. Emery, J. Powell, M. Strahl, G. Deikus, K.

Auckland, E. E. Eichler, W. A. Marasco, R. Sebra, A. J. Sharp, M. L. Smith, A. Bashir, and

C. T. Watson. 2020. A Novel Framework for Characterizing Genomic Haplotype Diversity

in the Human Immunoglobulin Heavy Chain Locus. *Front Immunol* 11.

33. Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:

3094-3100.

34. Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis,

R. Durbin, and G. P. D. P. Subgroup. 2009. The Sequence Alignment/Map format and

SAMtools. *Bioinformatics* 25: 2078-2079.

35. Martin, M., M. Patterson, S. Garg, S. O Fischer, N. Pisanti, G. W. Klau, A. Schöenhuth,

and T. Marschall. 2016. WhatsHap: fast and accurate read-based phasing. *bioRxiv*:

085050.

36. Fu, L., B. Niu, Z. Zhu, S. Wu, and W. Li. 2012. CD-HIT: accelerated for clustering the next-

generation sequencing data. *Bioinformatics (Oxford, England)* 28: 3150-3152.

37. Lefranc, M. P. 2001. IMGT, the international ImMunoGeneTics database. *Nucleic Acids*

Res 29: 207-209.

38. Al'Khafaji, A. M., J. T. Smith, K. V. Garimella, M. Babadi, M. Sade-Feldman, M. Gatzen,

S. Sarkizova, M. A. Schwartz, V. Popic, E. M. Blaum, A. Day, M. Costello, T. Bowers, S.

Gabriel, E. Banks, A. A. Philippakis, G. M. Boland, P. C. Blainey, and N. Hacohen. 2021.

High-throughput RNA isoform sequencing using programmable cDNA concatenation.

bioRxiv: 2021.2010.2001.462818.

39. Brochu, H. N., E. Tseng, E. Smith, M. J. Thomas, A. M. Jones, K. R. Diveley, L. Law, S.

G. Hansen, L. J. Picker, M. Gale, and X. Peng. 2020. Systematic Profiling of Full-Length

Ig and TCR Repertoire Diversity in Rhesus Macaque through Long Read Transcriptome

Sequencing. *The Journal of Immunology* 204: 3434.

40. Irvine, E. B., and G. Alter. 2020. Understanding the role of antibody glycosylation through

the lens of severe viral and bacterial diseases. *Glycobiology* 30: 241-253.

41. Alter, G., T. H. M. Ottenhoff, and S. A. Joosten. 2018. Antibody glycosylation in

inflammation, disease and vaccination. *Seminars in Immunology* 39: 102-110.

42. Collin, M. 2020. Antibody glycosylation as an immunological key in health and disease.

Glycobiology 30: 200-201.

43. Plomp, R., L. R. Ruhaak, H.-W. Uh, K. R. Reiding, M. Selman, J. J. Houwing-Duistermaat,

P. E. Slagboom, M. Beekman, and M. Wuhrer. 2017. Subclass-specific IgG glycosylation

is associated with markers of inflammation and metabolic health. *Scientific Reports* 7:

12325.

44. Trampert, D. C., L. M. Hubers, S. F. J. van de Graaf, and U. Beuers. 2018. On the role of

IgG4 in inflammatory conditions: lessons for IgG4-related disease. *Biochimica et*

Biophysica Acta (BBA) - Molecular Basis of Disease 1864: 1401-1409.

FIGURE 1.

FIGURE 2.

FIGURE 3.

A.

B.

D.

C.

E.

FIGURE 4.

FIGURE 5.

A. Severe COVID-19 hospitalization

BD BD BD BD

Day1 Day4 Day8 Day13 (Post hospitalization)

Whole blood RNA isolation over four timepoints

IgG and IgM FLAIRR-seq

B. Clones per IgG sub-isotype

C. IgG Sub-Isotype diversity over time

D. IgG sub-Isotype polarity over time

E. Clone 13611

Sequence frequency per isotype/sub-isotype

	IgM	IgG1	IgG2	IgG3	IgG4
Day 1	46%	0%	54%	0%	0%
Day 4	0%	0%	100%	0%	0%
Day 8	66%	2%	32%	0%	0%
Day13	50%	0%	50%	0%	0%

F. Clone 13611

State
- IGHM
- IGHG2
- IGHG1
- Germline

0.01

Table 1: Comparative analysis metrics between AIRR-seq and FLAIRR-seq on matched samples

Sample ID	Isotype	Method	RNA input (ng)	Input reads into Immcantation[a]	Unique VDJ[b]	Unique Clones[c]	Unique CDR3[d]
0007	IgG	AIRR	100	**841627**	3164	1008	1187
		FLAIRR	335	181109	4880	1358	1520
201c	IgG	AIRR	100	**1509063**	4568	1221	1371
		FLAIRR	296	221301	10819	2145	2388
203c	IgG	AIRR	100	**1006965**	5793	1436	1705
		FLAIRR	106	475703	3778	993	1138
602c	IgG	AIRR	100	**1275902**	7736	1542	2217
		FLAIRR	149	344906	9225	1828	2357
705c	IgG	AIRR	100	**1113660**	1362	557	576
		FLAIRR	268	279601	2328	1063	1081
1008	IgG	AIRR	100	**1310350**	12217	3005	3307
		FLAIRR	335	238820	18235	3505	3813
1013	IgG	AIRR	100	**1158222**	5027	1795	1964
		FLAIRR	222	396322	7487	2565	2744
2008	IgG	AIRR	100	**835928**	3900	1235	1368
		FLAIRR	240	418906	4258	1308	1380
4002	IgG	AIRR	100	**632272**	1112	727	772
		FLAIRR	335	398838	5377	1719	1887
5001	IgG	AIRR	100	**695884**	1885	1134	1244
		FLAIRR	335	313843	12146	2226	2381
0007	IgM	AIRR	100	**841627**	9154	7924	8163
		FLAIRR	335	331380	14926	12367	12851
201c	IgM	AIRR	100	**1509063**	17888	14281	14622
		FLAIRR	296	148355	12803	9353	9573
203c	IgM	AIRR	100	**1006965**	18189	16295	16629
		FLAIRR	106	509536	10636	9315	9419
602c	IgM	AIRR	100	**1275902**	8378	7704	7833
		FLAIRR	149	273103	16406	14495	14745
705c	IgM	AIRR	100	**1113660**	14273	12936	13413
		FLAIRR	268	435391	17417	15706	16249
1008	IgM	AIRR	100	**1310350**	12710	7878	8368
		FLAIRR	335	181063	15284	8745	9276
1013	IgM	AIRR	100	**1158222**	9012	8057	8241
		FLAIRR	222	262617	17812	15291	15759
2008	IgM	AIRR	100	**835928**	8551	5838	6206
		FLAIRR	240	502354	14132	10225	10674
4002	IgM	AIRR	100	**632272**	11607	11184	11367
		FLAIRR	335	489287	22379	19000	21532
5001	IgM	FLAIRR	335	**313843**	12146	2226	2381
		FLAIRR	355	171688	6164	4866	5011

([a]) AIRR-seq data reflects total number of unassembled paired end reads (R1 and R2 bold) used as input into analyses.
([b]) Counts of unique sequences after Immcantation annotation
([c]) Counts of unique clones after clonal clustering
([d]) Counts of sequences containing unique CDR3 sequences