FLAIRR-seq: A novel method for single molecule resolution of near full-length

immunoglobulin heavy chain repertoires

Authors:

Easton E. Ford[1,2], David Tieri[2], Oscar Rodriguez[2], Nancy Francoeur[3], Juan Soto[3], Justin Kos[2],

Ayelet Peres[4,5], William Gibson[2], Catherine A. Silver[2], Gintaras Deikus[3], Elizabeth Hudson[2],

Cassandra R. Woolley[1], Noam Beckmann[3], Alexander Charney[3], Thomas C. Mitchell[1], Gur

Yaari[4,5], Robert P. Sebra[3], Corey T. Watson[2*], Melissa L. Smith[2*]

Affiliations:

* denotes equal author contribution to the manuscript

[1]Department of Microbiology and Immunology University of Louisville School of Medicine,

Louisville, Kentucky, USA

[2]Department of Biochemistry and Molecular Genetics University of Louisville School of

Medicine, Louisville, Kentucky, USA

[3]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New

York City, New York, USA

[4] Faculty of Engineering, Bar Ilan University, 5290002 Ramat Gan, Israel

[5] Bar Ilan Institute of Nanotechnology and Advanced Materials, Bar Ilan University, 5290002,

Ramat Gan, Israel

Corresponding Author:

Melissa L. Smith, PhD

ml.smith@louisville.edu

Abstract:

Current Adaptive Immune Receptor Repertoire Sequencing (AIRR-seq) strategies resolve

expressed antibody (Ab) transcripts with limited resolution of the constant region. Here we present

a novel near full-length AIRR-seq (FLAIRR-Seq) method that utilizes targeted amplification by 5'

rapid amplification of cDNA ends (RACE), combined with single molecule, real-time sequencing

to generate highly accurate (>Q40, 99.99%) IG heavy chain transcripts. FLAIRR-seq was

benchmarked by comparing IG heavy chain variable (IGHV), diversity (IGHD), and joining (IGHJ)

gene usage, complementarity-determining region 3 (CDR3) length, and somatic hypermutation to

matched datasets generated with standard 5' RACE AIRR-seq and full-length isoform

sequencing. Together these data demonstrate robust, unbiased FLAIRR-seq performance using

RNA samples derived from peripheral blood mononuclear cells, purified B cells, and whole blood,

which recapitulated results generated by commonly used methods, while additionally resolving

novel IG heavy chain constant (IGHC) gene features. FLAIRR-seq data provides, for the first time,

simultaneous, single-molecule characterization of IGHV, IGHD, IGHJ, and IGHC region genes

and alleles, allele-resolved subisotype definition, and high-resolution identification of class-switch

recombination within a clonal lineage. In conjunction with genomic sequencing and genotyping of

IGHC genes, FLAIRR-seq of the IgM and IgG repertoires from 10 individuals resulted in the

identification of 32 unique IGHC alleles, 28 (87%) of which were previously uncharacterized.

Together, these data demonstrate the capabilities of FLAIRR-seq to characterize IGHV, IGHD,

IGHJ, and IGHC gene diversity for the most comprehensive view of bulk expressed Ab repertoires

to date.

Introduction

Antibodies (Abs) or immunoglobulins (IGs) are the primary effectors of humoral immunity

and are found as both membrane-bound receptors on B cells and circulating, secreted proteins

(1). Both membrane-bound B cell receptors (BCRs) and secreted Abs act to recognize and bind

antigen. All Abs and BCRs are composed of two identical heavy and light chains that are post-

translationally associated. The heavy chain is comprised of two distinct domains: (i) the variable

domain (Fab), which allows for antigen binding, and (ii) the constant domain (Fc), which

modulates downstream effector functions (1, 2). The light chain also includes a variable domain

that, once post-translationally associated with the heavy chain variable domain, interacts with

cognate antigen (3). In humans, Abs are grouped into discrete isotypes and subisotypes (i.e.,

IgM, IgD, IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, and IgE), based on the expression of specific

constant (C) genes within the IG heavy chain locus (IGH). Each isotype and subisotype has

unique effector properties that together represent the wide diversity of Ab-mediated functions,

including binding of Fc receptors (FCR), activation of complement, opsonization, antibody-

dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP) (4,

5).

To facilitate the development of diverse Ab repertoires capable of recognizing the wide

range of pathogens humans encounter, the IG genomic loci are highly polymorphic and harbor

diverse and complex sets of genes that recombine in each B cell to encode up to 10^{13} unique

specificities (6). B cells create this expansive catalog of specificities through somatic

recombination of the variable (V), diversity (D), and joining (J) genes in IGH, and V and J genes

from the corresponding light chain loci, lambda (IGL) and kappa (IGK) (7). During VDJ

recombination in IGH, a single D and J gene are first recombined, while the intervening and

unselected D and J gene sequences are removed by RAG recombinase (8). After D and J genes

are joined, further recombination of a specific V gene to the DJ gene cassette completes the

formation of the full VDJ rearrangement. Following transcription of the recombined VDJ, a single

constant region gene is spliced together with the VDJ cassette to generate the completed heavy

chain transcript (7). Recombination at IGL and IGK occurs similarly, recombining V and J genes

only. Heavy and light chain transcripts are independently translated and linked via covalent

cysteine bonds resulting in a fully functional protein prior to B cell cell-surface expression or

secretion (Figure 1A) (9). Naïve B cells, which develop in the bone marrow from hematopoietic

stem cell progenitors, have undergone VDJ recombination but have not yet encountered antigen,

and solely express IgM and IgD (10). These naïve B cells then migrate to B cell zones in

secondary lymphoid tissues where they encounter antigen, driving further maturation and class

switch recombination (CSR) to enable the most effective humoral responses (11). CSR mediates

the excision of IGHC genes at the DNA level, which leads to the utilization and linkage of different

IGHC genes to the same VDJ, ultimately resulting in class switching to alternate isotypes and

subisotypes (11).

The IgG isotype class is represented by four subisotypes: IgG1, IgG2, IgG3, and IgG4.

Each IgG subisotype circulates at varied frequencies and facilitates unique immune functions. For

example, IgG1 is typically the most abundant circulating IgG and mediates proinflammatory

responses; IgG2 targets bacterial polysaccharides, providing protection from bacterial pathogens;

IgG3 confers protection against intracellular bacterial infections and enables clearing of parasites;

and IgG4 contains exclusive structural and functional characteristics often resulting in anti-

inflammatory and tolerance-inducing effects (5). Multiple studies have identified Ab-mediated

subisotype-specific pathogenicity in the context of autoimmune diseases and cancer highlighting

the need for further investigation of subisotype-specific repertoires (12-15).

Current Adaptive Immune Receptor Repertoire sequencing (AIRR-seq) methods aim to

resolve variable and constant region transcripts to differing extents. Profiling of the variable

region, even in part, defines V, D and J gene usage while also providing characterization of

complementarity determining regions (CDR) 1, 2, and 3, which are hypervariable and directly

interact with target antigen (Figure 1B) (16). CDR3-targeted profiling approaches, such as those

used by Adaptive Biotechnologies (17), allow for V, D, and J gene assignments but do not provide

complete resolution of the entire variable region (17). Multiplexed primer-based AIRR-seq

strategies generate full variable region content but require specific primers to known targets and

therefore may miss novel genes and alleles. 5' RACE AIRR-seq methods capture the full-length

VDJ exon without variable region-targeted multiplexed primer pools therefore limiting the impact

of primer bias and enabling discovery of novel IGHV, IGHD, and IGHJ genes and alleles (18). 5'

RACE methods also often prime from the first IGHC exon (CH1), allowing for determination of

isotype. Additional methods have been developed that shift amplification strategies by capturing

additional IGHC region sequence to enable subisotype resolution; however, these methods

sacrifice full and contiguous characterization of the IGHV gene (19). All commonly used AIRR-

seq methods are further technically limited by the length restrictions (≤ 600nt) of short-read next

generation sequencing. As a result, no current AIRR-seq strategy resolves the complete heavy

chain transcript, including all IGHC exons alongside the recombined IGHV, IGHD, and IGHJ

genes. Our team has recently shown that population-based polymorphisms within the IGHV,

IGHD, IGHJ and IGL loci are far more extensive than previously known; the IGHC region has also

been shown to contain genomic diversity, although the extent of this diversity has likely not been

fully explored (20-27). Although it is understood that the Fc domain mediates Ab effector

functions, there is limited knowledge as to how genetic variation in this region may impact

functional capabilities or posttranslational modification (5, 28, 29). As such, there is a growing

need to understand genomic variation across the complete Ab molecule. To address these

limitations, we have developed novel, end-to-end pipeline to target, profile, and characterize the

Ab heavy chain repertoire in the context of isotype (IgG, IgM) and subisotype (IgG1, IgG2, IgG3,

IgG4).

Here, we present FLAIRR-seq, a novel targeted 5' RACE-based amplification of near full-

length IgG and IgM heavy chain transcripts, paired with single molecule real time (SMRT)

sequencing, resulting in highly accurate (mean read accuracy ~Q60, 99.9999%), near full-length

Ab sequences from RNA derived from whole blood, isolated PBMC, and purified B cells. When

analyzed with the Immcantation AIRR-seq tool suite (30, 31), we demonstrate that FLAIRR-seq

performs comparably to standard 5'RACE AIRR-seq methods and single-molecule isoform

sequencing (Iso-Seq) strategies for characterizing the expressed Ab repertoire. We further

highlight the novel features of FLAIRR-seq data, including phased identification of IGHV, IGHD,

IGHJ, and IGHC genes, facilitating the profiling of subisotype- and IGHC allele-specific repertoires

and CSR characterization.

Materials and Methods

Sample collections

Experiments were conducted using healthy donor peripheral blood mononuclear cells

(PBMC), purified B cells from healthy donors, or whole blood collected from hospitalized COVID-

19 patients (Supplementary Table 1). Commercially available healthy donor PBMC (STEMCELL

Technologies) and a subset of matched purified B cells were utilized to generate AIRR-seq and

FLAIRR-seq validation datasets. Full-length isoform sequencing (Iso-Seq) was performed using

B cells isolated from the PBMC of a healthy, consented 57-year-old male donor at the University

of Louisville (UofL) School of Medicine. The UofL Institutional Review Board approved sample

collection (IRB 14.0661). For COVID-19 affected patient samples (n=5), whole blood was

collected from the Mount Sinai COVID-19 biobank cohort of hospitalized COVID-19 patients,

approved by the Institutional Review Board at the Icahn School of Medicine at Mount Sinai as

previously described (32).

PBMC isolation and B cell purification

Frozen healthy donor PBMCs were purchased, thawed, and aliquoted for use in

downstream experiments (STEMCELL Technologies). For Iso-Seq analyses, 175mL of venous

blood was collected in a final concentration of 6mM K_3EDTA using standard phlebotomy. PBMCs

were isolated using Sepmate PBMC Isolation Tubes (STEMCELL Technologies) as previously

described (33), with an additional granulocyte depletion step using the RosetteSep Human

Granulocyte Depletion Cocktail (STEMCELL Technologies) as directed by the manufacturer. B

cells from the freshly collected and frozen healthy donor PBMC were isolated using the EasySep

Human Pan-B Cell Enrichment Kit, as described by the manufacturer (STEMCELL Technologies).

Briefly, B cells, including plasma cells, were isolated by negative selection using coated magnetic

particles. First, the B cell enrichment cocktail was added to the sample and mixed for a 5-minute

incubation at room temperature, followed by addition of magnetic particles and further incubation

for 5 minutes on the benchtop. The sample tube was then placed on an EasySep magnet

(STEMCELL Technologies), and purified B cells were carefully eluted from the magnetic particles

and immediately used for RNA extraction.

Genomic DNA and RNA extraction

For the healthy frozen PBMC and matched purified B cells, genomic DNA (gDNA) and

RNA were co-extracted using the AllPrep DNA/RNA Mini Kit (Qiagen) according to the

manufacturer's instructions. For the freshly processed UofL healthy donor PBMC, purified Pan-B

cells were lysed in Buffer RLT Plus, and RNA was extracted with the RNeasy Plus Mini Kit

(Qiagen) per the manufacturer's protocol; no gDNA was collected from this sample. COVID-19

whole blood-derived RNA was extracted from samples collected in Tempus Blood RNA tubes

using the Tempus Spin RNA Isolation Kit (ThermoFisher) as described by the manufacturer. For

all samples, concentrations of RNA and gDNA (when appropriate) were assessed using the Qubit

4.0 fluorometer, with the RNA HS Assay Kit and Qubit DNA HS Assay Kit, respectively

(ThermoFisher Scientific). RNA and gDNA integrity were evaluated using the Bioanalyzer RNA

Nano Kit and DNA 1200 Kit, respectively (Agilent Technologies). Extracted RNA and gDNA were

stored at -80℃ and -20℃, respectively, until used.

FLAIRR-seq targeted amplification of heavy chain transcripts

Extracted RNA was thawed on ice and converted to first strand complementary DNA

(cDNA) using the SMARTer RACE 5'/3' Kit (Takara Bio USA), as described by the manufacturer

and a custom oligonucleotide that contained the template switch oligo and a unique molecular

identifier (5' TSO-UMI) for template switch during first strand cDNA synthesis. The following

reaction conditions were used: (i) a primary master mix was prepared with 4.0 µL 5X First-Strand

Buffer, 0.5 µL DTT (100 mM), and 1.0 µL dNTPs (20 mM) per reaction and set aside until needed;

(ii) in a separate 0.2-mL PCR tube, 10 µL of sample RNA and 1 µL 5'-CDS Primer A were

combined and incubated in a thermal cycler at 72°C (lid temperature: 105°C) for 3 minutes,

followed by cooling to 42°C for 2 minutes; (iii) after cooling, tubes were spun briefly to collect

contents and 1µL (12µM) of the 5' TSO-UMI was added to the RNA; (iv) 0.5 µL of RNase inhibitor

and 2.0 µL of SMARTScribe Reverse Transcriptase were added to the primary master mix tube

per sample and 8 µL of the combined master mix was then added to each RNA-containing sample

tube. First-strand cDNA synthesis reactions were incubated in a thermal cycler at 42°C (lid

temperature: 105°C) for 90 mins, followed by heat inactivation at 70°C for 10 minutes. Total first

strand cDNA generated in this reaction was diluted 1:2 with Tricine-EDTA Buffer before moving

onto targeted heavy chain transcript amplification.

To specifically amplify heavy chain transcripts from total first-strand cDNA, targeted IgG

and IgM transcript amplification reactions were performed using barcoded IgG (3' primer binding

in the constant region exon 3, CH3) or IgM (3' primer binding in the constant region exon 4, CH4)-

specific primers (Supplementary Table 2) and the following conditions: (i) 5 µL of diluted first-

strand cDNA was added to 0.2-mL PCR tubes; (ii) a master mix was generated using 10 µL 5X

PrimeSTAR GXL Buffer, 4 µL GXL dNTP mixture, 28 µL PCR-grade water, 1 µL PrimeSTAR GXL

Polymerase and 1 µL 10x UPM form the SMARTer RACE 5'/3' Kit per reaction; (iii) 44 µL master

mix was added to each reaction tube followed by 1 µL of the appropriate barcoded IgG (CH3) or

IgM (CH4) primer. Different temperatures were used for annealing of IgG- (63.5°C) and IgM-

specific primers (60°C) to account for primer specific melting temperatures and to enhance

targeted amplification specificity. Amplification conditions for full-length IgG were: 1 minute at

95°C, followed by 35 amplification cycles of 95°C for 30 sec, 63.5°C for 20 sec, and 2 minutes at

68°C, followed by a final extension for 3 minutes at 68°C and hold at 4°C. Amplification conditions

for full-length IgM were: 1 minute at 95°C, followed by 35 amplification cycles of 95°C for 30 sec,

60°C for 20 sec., and 2 minutes at 68°C, followed by a final extension for 3 minutes at 68°C and

hold at 4°C. Final amplification reactions were purified using a 1.1x (vol:vol) cleanup with ProNex

magnetic beads (Promega) . Successfully amplified products were quantified with Qubit dsDNA

HS assay (ThermoFisher Scientific) and length was evaluated with the Fragment Analyzer

Genomic DNA HS assay (Agilent). Samples were equimolar pooled in 8-plexes for SMRTbell

library preparation and sequencing.

FLAIRR-seq SMRTbell library preparation and sequencing

Eight-plex pools of targeted IgG or IgM amplicons were prepared into SMRTbell

sequencing templates according to the "Procedure and Checklist for Iso-Seq Express Template

for Sequel and Sequel II systems" protocol starting at the "DNA Damage Repair" step and using

the SMRTbell Express Template Prep Kit 2.0, with some modifications (Pacific Biosciences).

Briefly, targeted IgG and IgM amplicons underwent enzymatic DNA damage and end repair,

followed by ligation with overhang SMRTbell adapters as specified in the protocol. To increase

consistency in SMRTbell loading on the Sequel IIe system, we further treated the SMRTbell

libraries with a nuclease cocktail to remove unligated amplified products, using the SMRTbell

Enzyme Cleanup Kit, as recommended by the manufacturer (Pacific Biosciences). Briefly, after

heat-killing the ligase with an incubation at 65°C, samples were treated with a nuclease cocktail

at 37°C for 1 hour, and then purified with a 1.1X Pronex cleanup. Final SMRTbell libraries were

evaluated for quantity and quality using the Qubit dsDNA HS assay and Fragment Analyzer

Genomic DNA assay, respectively. Sequencing of each 8-plex, barcoded sample pool was

performed on one SMRTcell 8M using primer v4 and polymerase v2.1 on the Sequel IIe system

with 30 hr movies. Demultiplexed, high-fidelity circular consensus sequence reads ("HiFi reads")

were generated on the instrument for downstream analyses.

AIRR-seq SMARTer Human BCR IgG/IgM sequencing

Matched healthy donor RNA was used to generate targeted IgG and IgM AIRR-seq

libraries using the SMARTer Human BCR IgG IgM H/K/L Profiling Kit (Takara Bio USA) according

to the manufacturer's instructions with no modifications. Briefly, for each sample, proprietary IgG

and IgM primers were used to amplify heavy chain transcripts following a 5'RACE reaction. AIRR-

seq libraries were then quality controlled using the 2100 Bioanalyzer High Sensitivity DNA Assay

Kit (Agilent) and the Qubit 3.0 Fluorometer dsDNA High Sensitivity Assay Kit. Sequencing on the

MiSeq platform using 300 bp paired-end reads was performed using the 600-cycle MiSeq

Reagent Kit v3 (Illumina) according to the manufacturer's instructions, and FASTQ reads were

generated using the associated DRAGEN software package (Illumina).

B cell Iso-Seq

RNA extracted from healthy sorted B cells was used to generate Iso-Seq SMRTbell

libraries following the "Procedure & Checklist Iso-Seq Express Template Preparation for the

Sequel II System" with minor adaptations compared to the manufacturer's instructions. Briefly,

Iso-Seq libraries were generated using 500 ng high-quality (RIN > 8) RNA as input into oligo-dT

primed cDNA synthesis (NEB). Barcoded primers were incorporated into the cDNA during second

strand synthesis. Following double-stranded cDNA amplification, transcripts from two samples

sourced from purified B cells and NK cells were equimolar pooled as previously described (33).

SMRTbells were generated from the pooled cDNA as described above for the FLAIRR-seq

amplification products, including the addition of a nuclease digestion step. Quantity and quality of

the final Iso-Seq libraries were performed with the Qubit dsDNA High Sensitivity Assay Kit and

Agilent Fragment Analyzer Genomic DNA assay, respectively. This 2-plex Iso-Seq pool was

sequenced using primer v4 and polymerase v2.1 on the Sequel IIe system with a 30-hour movie.

HiFi reads were generated on instrument before analyses. Demultiplexing of barcoded samples

and generation of full-length non-concatemer (FLNC) predicted transcripts were performed using

the Iso-Seq v3 pipeline available through SMRTLink (v.10.2). B-cell-derived FLNC reads were

mapped to the human genome using the GMAP reference database and reads derived from

chromosome 14 were extracted for downstream IGH transcript characterization via Immcantation,

as described below.

Immcantation analyses of IgG and IgM repertoires

Analyses of FLAIRR-seq, AIRR-seq, and Iso-seq datasets were performed using

Immcantation tools (30, 31). Demultiplexed barcoded HiFi (for SMRT sequencing data) or FASTQ

(for AIRR-seq) reads were first processed using the pRESTO tool for quality control, UMI

processing, and error profiling (30). For AIRR-seq, pRESTO analysis data from paired-end reads

("R1" and "R2") were trimmed to remove bases with < Q20 read quality and/or <125 bp length

using the "FilterSeq trimqual" and "FilterSeq length", respectively. IgG and IgM CH3 or CH4 primer

sequences were identified with an error rate of 0.2, and primers identified were then noted in

FASTQ headers using "MaskPrimers align". Next, 12 basepair (bp) UMIs were located and

extracted using "Maskprimers extract". Sequences found to have the same UMIs were grouped

and aligned using "AlignSets muscle," with a consensus sequence generated for each UMI using

"BuildConsensus". Mate pairing of AIRR-seq reads was conducted using a reference-guided

alignment requiring a minimum of a 5 bp overlap via "AssemblePairs sequential". After collapsing

consensus reads with the same UMI ("conscount") using "CollapseSeq"," reads with < 2

supporting sequences were removed from downstream analysis. For pRESTO processing of

FLAIRR-seq, single HiFi reads ("R1") reads did not require trimming due to > Q20 sequence

quality across all bases. 5'TSO-UMI and CH3 or CH4 region primers were identified along with a

22 bp UMI with an error rate of 0.3 using "MaskPrimers align". Reads were then grouped and

aligned using "AlignSets muscle". Due to the single molecule nature of FLAIRR-seq reads, no

mate pairing was required. Consensus reads were then generated as described above, including

removal of sequences with < 2 supporting reads. Read counts following each step of data filtration

for AIRR-seq and FLAIRR-seq are represented in Supplementary Tables 3 and 4, respectively.

pRESTO-filtered reads for both AIRR-seq and FLAIRR-seq data were then input into the

Change-O tool (Table 1). Iso-seq required no initial processing from pRESTO and was input into

Change-O for IG gene reference alignment along with AIRR-seq and FLAIRR-seq data using

"igblastn", clonal clustering using "DefineClones", and germline reconstruction and conversion

using "CreateGermlines.py" and the GRCh38 chromosome 14 germline reference (31). Fully

processed and annotated data was then converted into a TSV format for use in downstream

analyses. The Alakazam Immcantation tool suite was then used to quantify gene usage analysis,

calculate CDR3 length, assess somatic hypermutation frequencies, and analyze clonal diversity

(31). SCOPer clonal assignment by spectral clustering was conducted for COVID-19 patient time

course samples (34, 35). For clonal lineage tree analysis, the Dowser tool was used to examine

clonal diversity and CSR over time (36).

Targeted IG gDNA capture, long-read sequencing, and IGenotyper analyses.

FLAIRR-seq validation samples also underwent IGHC targeted enrichment and long-read

sequencing as previously described (37). Briefly, gDNA was mechanically sheared and size

selected to include 5-9kb fragments using the BluePippin (Sage Science). Samples were then

end repaired and A-tailed using the standard KAPA library preparation protocol (Roche).

Universal priming sequences and barcodes were then ligated onto samples for multiplexing

(Pacific Biosciences). Barcoded gDNA libraries were captured using IGH-specific probes

following a SeqCap protocol (Roche). 26 IGH-enriched samples were purified and pooled together

for SMRTbell library prep as described above, including the final nuclease digestion step. Pooled

SMRTbells were annealed to primer v4, bound to polymerase v2.0, and sequenced on the Sequel

IIe system with 30h movies. After sequencing, HiFi reads were generated and analyzed by the

IGenotyper pipeline (37). In brief, IGenotyper was used to detect single nucleotide variants and

assemble sequences into haplotype-specific assemblies for downstream IGHC gene genotyping.

Alleles were then extracted from assemblies using a bed file containing coordinates for each

IGHC gene exon. After sequences were extracted, reads were then aligned to the IMGT database

(downloaded on 2/21/22) and assigned as an exact match to IMGT, "novel" if there was no match

to the IMGT database or "novel, extended" if a match was detected to a partial allele found in

IMGT, but the IMGT allele was a substring of the IGenotyper identified allele (38). This set of

alleles was then used as a ground truth dataset.

IGHC gene genotyping with FLAIRR-seq

To genotype IGHC genes and alleles from FLAIRR-seq data, productive reads were

filtered by IGHC length (900bp–1100bp) and aligned to the chromosome 14 hg38 reference using

minimap2 along with SAMtools to generate sorted and indexed bam files (39, 40). WhatsHap was

used to identify, genotype, and phase single nucleotide variants (SNV) (41). Phased SNVs were

used to assign each read to a haplotype using MsPAC. Reads from each haplotype and gene

were clustered using CD-HIT using a 100% identity clustering threshold parameter, and a single

representative read from the largest cluster was aligned to the IGenotyper curated alleles using

BLAST (42) to determine the closest matching IGHC gene and allele (38, 43). The representative

read was selected based on 100% identity to all other sequences in that cluster.

Inference of IGHV, IGHD, and IGHJ gene haplotypes from FLAIRR-seq data using IGHC

gene anchors

To test the ability of IGHC genes to be used for the inference of IGHV, IGHD, and IGHJ haplotypes

from FLAIRR-seq data, we chose one sample that was heterozygous for both IGHM and IGHJ6

(IGHJ6 is standardly used for AIRR-seq haplotype inference). For this sample, we employed

TigGER (31, 44-46) to infer novel IGHV alleles, and generate sample-level IGHV genotypes using

a Bayesian approach. Rearranged sequences within the Change-O table were then reannotated

taking into account sample genotype and detected novel alleles. Updated annotations were then

used to infer haplotypes using RAbHIT version 0.2.4 (47). Both IGHJ6 and IGHM were used as

anchor points for haplotyping, and the resulting haplotypes were compared.

Results

Gene usage, CDR3, and SHM profiles characterized from FLAIRR-seq data are comparable

to AIRR-seq and Iso-Seq

Current methods for commercially available 5' RACE AIRR-Seq utilize targeted

amplification of the variable region and, in some cases, a small portion of the first constant region

exon (CH1), in conjunction with short-read sequencing to characterize IG repertoires. However,

this minimal examination of the IGHC gene sequence is primarily used to define isotypes. No

current method defines both the heavy chain variable and constant regions allowing for both

subisotype classification and IGHC allele-level resolution. To address these technical limitations,

we developed the FLAIRR-Seq method (Figure 1C), a targeted 5' RACE approach combined with

SMRT sequencing to generate highly accurate, near full-length IgG (~1500 bp) and/or IgM (2000

bp) sequences, allowing for direct, simultaneous analysis of the heavy chain variable and constant

regions (Figure 1D), including gene/allele identification for IGHV, IGHD, IGHJ and IGHC

segments, and isotype- and subisotype-specific repertoire profiling.

To evaluate and validate the capabilities of FLAIRR-seq, matched FLAIRR-seq and AIRR-

seq analyses were performed on ten healthy donor PBMC samples. FLAIRR-seq data were

filtered from the initial HiFi reads (>Q20) to include only >Q40 reads. The average read quality of

these filtered reads was >Q60 (99.9999%), with a pass filter rate ranging from 88%-93% of total

reads. AIRR-seq FASTQ bases were trimmed to retain sequences with an average quality of Q20

(99%). These filtered reads were used as input into the Immcantation suite, specifically the

pREST-O and Change-O tools, for IGHV, IGHD, and IGHJ gene assignment, and repertoire

feature analyses, including identification of clones, extent of somatic hypermutation, and

evaluation of CDR3 lengths. As shown in **Table 1,** fewer overall FLAIRR-seq reads were used as

input into the Immcantation analyses, after required filtration and read assembly steps (which

were not needed for the high-quality single-molecule FLAIRR-seq reads). FLAIRR-seq resulted

in comparable or, in many cases, increased number of unique VDJ sequences, clones, and CDR3

sequences identified compared to the matched AIRR-seq-derived samples in both the IgG and

IgM repertoires. These basic sequencing and initial analysis metrics demonstrated that FLAIRR-

seq produced high-quality variable region data for detailed Ab repertoire analyses and is

amenable to analysis using existing AIRR-seq analysis tools. While input RNA mass used for

FLAIRR-seq was often more than for AIRR-seq, ongoing and future optimization of the method is

aimed at reducing RNA input. Furthermore, a comparative cost analysis was performed. To obtain

the closest metric to a direct comparison, we calculated the cost per "actionable read", defined as

the read number per sample and per method after read and length quality filtration and assembly,

but prior to cluster consensus, performed in the pRESTO pipeline. This method was used to

represent the total unique single molecule or assembled templates captured by either the

FLAIRR-seq or AIRR-seq methods, respectively, that passed all necessary quality control criteria

for downstream annotation and analyses irrespective of biologic repertoire diversity or clonality.

Lastly, to remove the impact of pooling differences, we used this "per actionable read" cost to

calculate the cost for the generation of 15,000 "actionable reads" as our standard price. For AIRR-

seq this cost was $25.50 per sample, whereas for FLAIRR-seq, this cost was $33.57 per sample.

These costs reflect needed reagents and consumables only, assume instrumentation access and

do not include labor. Future optimization for FLAIRR-seq will include integrating a multiplexed

array sequencing (MAS) step to concatenate reads and enhance overall depth of sequencing and

multiplexing capacity per pool, resulting in decreasing costs per sample (48).

While optimizing FLAIRR-seq sample preparation, we examined whether upstream

isolation of B cells before FLAIRR-seq molecular preparation would enhance the ability to detect

IGHV, IGHD, and IGHJ gene usage. To do this, aliquots of PBMC (n=4) were split into two groups

for RNA extraction: (i) RNA derived from bulk PBMC, and (ii) RNA isolated from purified pan B-

cells, followed by FLAIRR-seq preparation, SMRT sequencing, and Immcantation analysis of both

groups. IGHV, IGHD, and IGHJ gene usage correlations between groups are shown in Figure 2A

and demonstrate a significant association (p-values ranging from 0.033 to $4.1e^{-16}$) strongly

supporting the conclusion that B cell isolation before RNA extraction was not necessary to achieve

comparable gene usage metrics. The limited differences that were observed could be explained

by template sampling differences between the two experiments. Due to the strong associations

observed and the ease of processing PBMC in bulk, we moved forward with RNA derived directly

from PBMC aliquots for the remainder of our analyses.

We established FLAIRR-seq performance by comparing its output to the commonly used

5' RACE AIRR-seq method. 5' RACE AIRR-seq was chosen as it provides resolution of the

complete variable region and a small portion of IGHC, allowing for isotype differentiation.

Following matched preparation of both FLAIRR-seq and AIRR-seq libraries from healthy donor

PBMC samples (n=10), we compared multiple repertoire features to benchmark FLAIRR-seq

performance. First, we evaluated IGHV, IGHD, and IGHJ gene usage frequencies. We observed

significant correlations between FLAIRR-seq and AIRR-seq datasets in IGHV, IGHD and IGHJ

gene usage for both IgM (V genes: r = 0.93-0.97, p=<$2.2e^{-16}$; D genes: r = 0.98-0.99, p=<$2.2e^{-16}$;

J genes: r = 0.94-1.0, p = 0.0028-0.017) and IgG isotypes (V genes: r = 0.90-0.96, p=<$2.2e^{-16}$; D

genes: r = 0.87-0.99, p=<$2.2e^{-16}$-$6.1e^{-14}$;J genes: r = 0.89-1.0, p = 0.0028-0.033) (Figure 2B),

indicating that FLAIRR-seq comparably resolves IGHV, IGHD, and IGHJ gene usage profiles. To

note, IGHJ genes showed lower levels of significance (larger p-values) across all comparisons

due to the relatively few genes that make up the IGHJ gene family compared to the more diverse

IGHV and IGHD families. We next investigated the performance of both methods in terms of

resolving somatic hypermutation (SHM) frequencies (Figure 2C), and complementarity

determining region 3 (CDR3) lengths (Figure 2D), which are often used as measures of evaluating

B cell affinity maturation. Although we did observe occasional statistically significant differences

in the SHM frequency between AIRR-seq and FLAIRR-seq data using the same samples, these

differences were not seen across all samples, suggesting that sample-to-sample variation may

drive this observation rather than technology-based discrepancies. We found that CDR3 lengths

were consistently longer in the FLAIRR-seq datasets for both the IgM and IgG isotypes in most

donors. The characterization of unusually long CDR3 regions (> 40 nt) in the IgG sequences with

FLAIRR-seq is likely due to the higher contiguity and quality afforded by the longer read lengths,

which are less likely to be spanned by short-read 2x300 bp paired-end sequencing strategies.

Together, these data demonstrate that FLAIRR-seq achieves comparable gene usage profiles,

and improved resolution of long CDR3 sequences.

Others have recognized the power of long-read sequencing to resolve B cell repertoires

using bulk Iso-Seq methods, allowing for the examination of full-length transcripts from isolated B

cells (49). The Iso-Seq method captures full-length transcripts expressing a poly(A) tail without

bias through oligo dT-based priming. The trade-offs of this approach are throughput, depth, and

cost, as Iso-Seq processing generates a complete transcriptome per sample without enrichment

of heavy chain sequences, which then need to be filtered out and analyzed, resulting in a

considerable amount of non-repertoire data that is discarded. To investigate whether the

untargeted transcriptome-wide Iso-Seq method would resolve a qualitatively different repertoire

than FLAIRR-seq, which would have indicated FLAIRR-seq-driven primer bias, we performed

matched Iso-Seq and FLAIRR-seq on purified B-cell derived RNA. IGHV, IGHD, and IGHJ gene

usage frequencies were compared between Iso-Seq and FLAIRR-seq datasets (Spearman's rank

correlation), revealing significant correlations between usage profiles (V genes: r = 0.94, p=2.2e⁻

[16]; D genes: r = 0.92, p =1.4e^{-13}; J genes: r = 1.0, p = 0.0028; Figure 2E). These data strongly

suggest that FLAIRR-seq has very limited to no primer-driven bias compared to whole

transcriptome data. Collectively, this benchmarking dataset confirmed that FLAIRR-seq is

comparable other state-of-the-art methods, providing robust characterization of commonly used

repertoire metrics, with limited increases in per sample cost.

IGenotyper and FLAIRR-seq provide constant region gene allele identification and allow

for haplotyping of variable genes

The novel value added by FLAIRR-seq is improved resolution of IGHC, including

estimation of IGHC gene and allele usage, subisotype identification, and phasing of variable and

constant regions for comprehensive repertoire analysis**.** To evaluate the capabilities and accuracy

of IGHC gene and allele identification with FLAIRR-seq, we first established a ground truth dataset

of IGHC alleles for all 10 samples by targeted sequencing of the germline IGH locus (Figure 3A)

using IGenotyper, as previously described (37). IGHG1, IGHG2, IGHG3, IGHG4 and IGHM alleles

called by IGenotyper (see Methods) were assigned to one of three categories, schematized in

Figure 3B: (i) "exact match" - alleles documented in the IMGT database; (ii) "novel not in IMGT"

– alleles not documented in the IMGT database; or (iii) "extended" – alleles that matched partial

alleles in the IMGT database (i.e., those only spanning a subset of exons), but were extended by

sequences in our dataset. IGenotyper identified a total of 32 unique IGHG1, IGHG2, IGHG3,

IGHG4 and IGHM alleles across all individuals, as schematized in Figure 3C. Among these 32

alleles, only 4 were documented in IMGT, the remaining represented novel alleles (n=11) or

extensions (n=17) of known alleles. In aggregate, we observed a greater number of IGHG4 alleles

than for any of the IGHG genes. Among these alleles were 4 sequences represented by

suspected duplications of IGHG4. In fact, we observed 3 IGHG4 gene alleles in 4/10 samples,

indicating the presence of gene duplications in these donors; in all cases, these alleles were also

identified in the FLAIRR-seq data (see below). Given the relatively small size of this proof-of-

concept healthy donor cohort, the identification of 28 (87%) novel or extended alleles underscores

the extensive polymorphism In this region and reflects the paucity of information regarding this

locus in existing immunogenomics databases.

We next used the iGenotyper-derived IGHC gene database as the ground-truth for

evaluating the capability of FLAIRR-seq to identify and resolve IGHC gene alleles. Based on our

analysis workflow for identifying IGHC alleles from FLAIRR-seq data, we resolved 19/32 (59%)

iGenotyper alleles at 100% identity; no additional false-positive alleles were identified. Of the

alleles that were not unambiguously resolved by our FLAIRR-seq pipeline, 8 had allele defining

single nucleotide variants (SNVs) 3' of the FLAIRR-seq primers. The rate of true-positive allele

calls using FLAIRR-seq across all 10 samples ranged from 5% for IGHG1 to 90% for IGHM

(Figures 3C and 3D). As a result, the IGHC genotypes inferred by FLAIRR-seq have some

limitations, but on the whole allow for much greater resolution of IGHC variation in the expressed

repertoire than currently used methods. Future iterations of FLAIRR-seq will include primer

optimization to facilitate better sequence coverage in the 3' regions of the IGHC alleles and

improve the direct genotyping capabilities of the FLAIRR-seq method.

Previous studies have demonstrated the use of IGHJ6 heterozygosity to infer haplotypes

of V and D genes from AIRR-seq data (46, 50). However, the frequency of IGHJ6 heterozygotes

in the population can vary. Therefore, we wanted to assess the utility of leveraging IGHC

polymorphism resolved by FLAIRR-seq for haplotyping IGHV alleles with the publicly available

tool RAbHIT(46, 50, 51). We selected a single donor ("1013") from our cohort that was

heterozygous for both IGHJ6 (*02 and *03) and IGHM (FL_2 and FL_4). Importantly, we were

able to associate each IGHJ6 allele to the respective IGHM allele from the corresponding

haplotype (Figure 3E). After defining germline IGHV alleles using TIgGER (31), we generated and

compared IGHV haplotype inferences using either IGHJ6 or IGHM alleles as anchor genes using

RAbHIT (47)(Figure 3E). Although some allele assignments were ambiguous ("unknown") using

both methods, we observed a strong consensus between haplotype inferences using the two

anchor genes. For haplotype 1, represented by IGHJ6*03 and IGHM_FL_4, the IGHJ6*03-derived

haplotype had 35 IGHV genes for which either an allele or deletion call was made. When using

IGHM_FL_4, the same allele/deletion calls were made for 33 of these genes; in addition, using

IGHM as the anchor gene, assignments were made for an additional 5 IGHV genes that had

"unknown" designations using IGHJ6. Similarly, of the allele/deletion calls made for 36 IGHV

genes on haplotype 2 assigned to IGHJ6*02, 33 of these gene had identical assignments to

IGHM_FL_2. Together these results indicate that IGHC variants can be utilized for haplotype

inference from repertoire data when commonly used IGHJ or IGHD genes are homozygous in

individuals of interest.

FLAIRR-seq enables isotype-, subisotype-, and allele-specific repertoire analyses

IGHG and IGHM alleles identified in each sample were used to annotate reads in each

respective repertoire. These assignments allowed for partitioning of the repertoire by isotype,

subisotype and IGHC allele (Figure 4). To demonstrate this, we utilized the same representative

sample ("1013") that was heterozygous for all IGHC genes. As shown in Figure 4A, IGHC gene

assignments allow for subisotype and allele level frequencies to be estimated as a proportion of

the overall IgG and IgM repertoires. In addition, detailed analyses of the repertoire can be

conducted within each of these compartments. For example, Figure 4B shows the frequencies of

IGHV gene subfamilies for each IGHG and IGHM allele identified in this sample. Using standard

AIRR-seq analyses, we would not be able to identify allele-resolved V gene usage or enrichment

within subisotype populations, which is important for linking subisotype functionality to particular

antigen-specific VDJ clones.

Through the partitioning of repertoire sequences by subisotype and IGHC allele, we found

that FLAIRR-seq also allowed for trends to be assessed in aggregate across donors. To

demonstrate this, we further examined V gene family usage partitioned by IgG subisotype across

all 10 healthy donors. This analysis revealed expected patterns, in that IGHV1, IGHV3 and IGHV4

subfamily genes were dominant across the 4 subisotypes (Figure 4C). However, we did observe

significant variation in subfamily proportions between subisotypes, associated with distinct

profiles in specific subisotypes (Figure 4C). Specifically, the estimated frequencies of IGHV1 and

IGHV3 were statistically different between subisotypes (P<0.01, ANOVA); IGHV1 usage was

elevated in IGHG1 and IGHG4, whereas IGHV3 was elevated in IGHG2 and IGHG3. These

analyses demonstrate the unique capability of FLAIRR-seq to examine variation in the expressed

repertoire at the level of isotype, subisotype, and IGHC allele. As samples sizes increase, we

expect that a multitude of additional repertoire features will become accessible to this kind of

analysis leading to novel discoveries linking VDJ and IGHC genetic signatures.

FLAIRR-seq identifies subisotype-specific clonal expansion and CSR in longitudinal

samples.

We wanted to investigate the utility of FLAIRR-seq in clinical samples, particularly to

observe changes in immune repertoires over time. Ab responses are highly dynamic, with specific

Ab clones expanding upon activation by antigen. We were interested to know if class switch

recombination could be captured by FLAIRR-seq, given the capability to identify clones with the

subisotype resolved. We had the opportunity to evaluate FLAIRR-seq resolved repertoires over

time in four samples collected from one individual over their >13-day hospitalization for severe

COVID-19 disease. Blood draws were taken on days 1, 4, 8, and 13 post-hospitalization (Figure

5A) and analyzed with FLAIRR-seq across all time points. After initial FLAIRR-seq processing

and analysis, we defined unique clones using SCOPer, which clusters sequences based on CDR3

similarity and mutations in IGHV and IGHJ genes (34). This analysis allowed for the estimation of

subisotype-specific clone counts across the four timepoints examined. Overall, we observed IgG1

dominated the repertoire at all four time points, but the proportion of subisotypes fluctuated over

time (Figure 5B). Specifically, the IGHG2 and IGHG3-specific repertoires expanded from day 1 to

day 4, but then contracted in overall frequency from day 8 to day 13 (Figure 5B). To assess clonal

diversity within each subisotype repertoire across time, we calculated the Simpson's diversity

index (q=2) using Alakazam (31). All subisotype-specific repertoires became less diverse from

day 1 to day 4, suggesting clonal expansion across the IGHG repertoire (Figure 5C). To note,

IGHG4 was not included in diversity calculations because IGHG4 would have required higher

sequencing depth to ascertain diversity, given the overall lower expression of IgG4 transcripts in

this individual. Subisotype-specific repertoire polarity was also assessed by calculating the

fraction of clones needed to represent 80% of the total repertoire (Figure 5D), with lower fractions

representing more polarized and clonally expended repertoires. Results of this analysis were

consistent with the diversity index, demonstrating an increase in clonal expansion (i.e., decreased

polarity) at day 4 across IGHG1, IGHG2, and IGHG3 compartments, which returned to baseline

at later timepoints.

CSR mediates the switching of Abs from one sub/isotype to another. This occurs through

the somatic recombination of IGHC genes, which brings the switched/selected IGHC genes

adjacent to the recombined IGHV, IGHD, and IGHJ segments, facilitating transcription. The

switching of isotypes and subisotypes can result in changes to associated effector functions of

the Ab while maintaining antigen-specific variable regions. Given the ability of FLAIRR-seq to

resolve clones with subisotype and IGHC allele resolution, as proof-of-concept, we sought to

assess whether FLAIRR-seq could allow for more detailed haplotype-level analysis of CSR

through the course of infection. To do this, we identified the largest clones in our dataset that were

both represented by multiple isotypes and found across timepoints. In total, using SCOper (34),

we identified 19 unique clonal lineages that met this criteria. We focused our detailed analysis on

one of the largest clones, "9900" (Figure 5E), comprising IGHM, IGHG1, and IGHG2 sequences.

On day 1 post-hospitalization, clone 9900 sequences were represented by both IGHM and

IGHG2. At day 4, IGHG2 was the only isotype observed, whereas again at day 8, both IGHM and

IGHG2 were observed, as well as IGHG1. To visualize CSR, we built a phylogeny using Dowser

(36). Highlighted in the red box on the phylogenetic tree shown in Figure 5F, we observe a single

subclade that is represented by IGHM, IGHG1, and IGHG2.

We were also able to resolve IGHC alleles from this individual, with the exception of

IGHG1 alleles which were ambiguous. Critically, both IGHG1 and IGHG2 were heterozygous

(Figure 5G). Through the assignment IGHG alleles to haplotypes within this individual using

heterozygous V genes (IGHV3-7 and IGHV3-48), we were able to determine that sequences from

clone 9900 (Figure 5F) utilized IGHG1 and IGHG2 alleles from the same haplotype, associated

with IGHV3-7*07 and IGHV3-48*03 (31). This observation offers direct characterization of CSR

events occurring on the same chromosome. When we looked across the remaining clones (n=8)

in this dataset that spanned time points and were represented by IGHG1 and IGHG2 subisotypes,

we were able to confirm that all of these clones used IGHG alleles from the same haplotype.

Together, these data provide demonstrative proof-of-concept evidence that FLAIRR-seq

profiling performs robustly on clinical samples, including RNA directly extracted from whole blood.

In addition, these data provide novel repertoire resolution extending what would have been

possible with standard AIRR-seq methods, including analysis of subisotype-specific repertoires,

evaluation of clonal expansion, and characterization of CSR in the IgG and IgM repertoires.

Discussion

Here we present the development, validation, and application of FLAIRR-seq, a novel

method to resolve near full-length Ab transcripts from bulk PBMC-, isolated B cell- and whole

blood-derived total RNA. FLAIRR-seq enables highly accurate, simultaneous resolution of

variable and constant regions and suggests that IGHC polymorphism is far more extensive than

previously assumed. FLAIRR-seq performed equivalent to or with increased resolution compared

to existing standard 5'RACE AIRR-seq methods when resolving V, D, and J gene calls, CDR3

lengths, and SHM signatures, suggesting that our CH3/CH4 targeting strategies did not

compromise variable region characterization while simultaneously adding the capability to resolve

IGHC variation. Little to no primer bias was observed when compared to Ab repertoire profiling

from total mRNA Iso-seq methods. FLAIRR-seq provides the novel ability to use IGHC gene

usage to identify subisotypes and genotype heavy chain transcripts, linking these data back to

evaluate subisotype-specific repertoires, clonal expansion, and CSR. Underscoring the

underappreciated extent of IGHC variation, our profiling of a restricted cohort of only 10 individuals

from relatively homogenous backgrounds still identified 4 and 7 completely novel IGHC alleles in

IgM and IgG, respectively, and extended an additional 17 alleles beyond which had been available

in the IMGT database.

The unique capabilities of FLAIRR-seq will allow for novel examination of Ab repertoires,

including the characterization of variable gene usage and clonotype distribution within unique

subisotype subsets. This perspective has the potential to provide key insights into dynamic Ab

responses in diseases known to be mediated by subisotype-specific processes or have skewed

subisotype distribution as predictive markers of disease, including Myasthenia gravis (mediated

by pathogenic autoantibodies across subisotypes that give rise to varied disease pathologies),

Acute Rheumatic Fever (associated with elevated IgG3), and melanoma (skewing towards IgG4

in late-stage disease thought to be indicative of tolerogenic responses and poor prognosis)(12-

14). These subisotype-specific repertoire profiling approaches may be the first step toward

identification of unique clones that mediate disease pathogenicity or serve as high-resolution

biomarkers to disease progression, as well as open the door for potential functional experiments

on subisotype clones of interest, including examining the functional impact of the novel IGHC

alleles identified here. Expanded population-based FLAIRR-seq profiling and curation of novel

IGHC alleles, particularly in conjunction with IGenotyper targeted genomic assembly efforts in

IGH, will be a significant first step in defining the full extent of variation in a region too long

assumed to be relatively invariant.

The Fc region is known to be critical for modulating differential Ab effector functions. These

differential functionalities are currently understood to be regulated by differential posttranslational

modifications, such as variable glycosylation (52-55). Future FLAIRR-seq profiling will be a

valuable tool to investigate how genomic variation across IGHC genes impacts residue usage

and resultant Fc receptor binding, signaling potential, crosslinking, and potential for

posttranslational modification, all of which would be expected to alter downstream effector

functions, such as ADCC, ADCP, and complement fixation.

We further demonstrate that FLAIRR-seq can effectively examine clonal expansion and

CSR in longitudinal samples, demonstrating the feasibility of using FLAIRR-seq to resolve Ab

repertoire dynamics. This increased resolution will further our understanding of Ab repertoire

evolution in the transition of acute to chronic disease states, many of which are associated with

overall IgG subisotype distribution changes that are thought to reflect the inflammatory milieu (14,

56). One example is advanced melanoma, where late-stage disease is characterized by elevated

IgG4 compared to IgG1, which is believed to reflect a more tolerizing, pro-tumor environment (12).

FLAIRR-seq examination of these samples may identify specific repertoire distribution patterns

that could act as biomarkers of disease progression. Moving forward it is critical to account for all

variability within the Ab repertoire for the most comprehensive understanding of repertoire

dynamics and the myriad factors impacting Ab effector function. Future efforts will expand

FLAIRR-seq methods to target IGHA and IGHE repertoires, as well as implement multiplexed

arrays (MAS) sequencing to considerably increase throughput and lower cost (48). Together, the

data presented here demonstrate that the FLAIRR-seq method provides a comprehensive

characterization of allele-resolved IgG and IgM repertoires, detailing variable region gene usage

and measurements of maturation, isotype and subisotype identification, and the unappreciated

extent of constant region variation, which will be necessary to fully appreciate the impact of IG

genomic variation in health and disease.

Acknowledgements:

The authors would like to thank Kamille Rasche, Kaitlyin Shields, Uddalok Jana, and the staff of

the University of Louisville Sequencing Technology Center for assistance in the sequencing and

analysis of FLAIRR-seq samples. They would also like to acknowledge the support and

assistance from Gur Yaari and Ayelet Peres for their assistance with the RabHIT workflow. This

work was funded, in part, by P20 GM135004-02 (EEF, DT, CTW, and MLS).

References:

1. Schroeder, H. W., and L. Cavacini. 2010. Structure and function of immunoglobulins. *Journal of Allergy and Clinical Immunology* 125: S41-S52.

2. Janda, A., A. Bowen, N. S. Greenspan, and A. Casadevall. 2016. Ig Constant Region Effects on Variable Region Structure and Function. *Front Microbiol* 7: 22.

3. Nakano, T., M. Matsui, I. Inoue, T. Awata, S. Katayama, and T. Murakoshi. 2011. Free immunoglobulin light chain: Its biology and implications in diseases. *Clinica Chimica Acta* 412: 843-849.

4. Lu, L. L., T. J. Suscovich, S. M. Fortune, and G. Alter. 2018. Beyond binding: antibody effector functions in infectious diseases. *Nature Reviews Immunology* 18: 46-61.

5. Vidarsson, G., G. Dekkers, and T. Rispens. 2014. IgG subclasses and allotypes: from structure to effector functions. *Front Immunol* 5: 520-520.

6. Greiff, V., C. R. Weber, J. Palme, U. Bodenhofer, E. Miho, U. Menzel, and S. T. Reddy. 2017. Learning the High-Dimensional Immunogenomic Features That Predict Public and Private Antibody Repertoires. *The Journal of Immunology* 199: 2985.

7. Tonegawa, S. 1983. Somatic generation of antibody diversity. *Nature* 302: 575-581.

8. Nishana, M., and S. C. Raghavan. 2012. Role of recombination activating genes in the generation of antigen receptor diversity and beyond. *Immunology* 137: 271-281.

9. Tong, P., A. Granato, T. Zuo, N. Chaudhary, A. Zuiani, S. S. Han, R. Donthula, A. Shrestha, D. Sen, J. M. Magee, M. P. Gallagher, C. E. van der Poel, M. C. Carroll, and D. R. Wesemann. 2017. IgH isotype-specific B cell receptor expression influences B cell fate. *Proc Natl Acad Sci U S A* 114: E8411-e8420.

10. Noviski, M., J. L. Mueller, A. Satterthwaite, L. A. Garrett-Sinha, F. Brombacher, and J. Zikherman. 2018. IgM and IgD B cell receptors differentially respond to endogenous antigens and control B cell fate. *eLife* 7: e35074.

11.   Stavnezer, J., J. E. Guikema, and C. E. Schrader. 2008. Mechanism and regulation of

class switch recombination. *Annu Rev Immunol* 26: 261-292.

12.   Karagiannis, P., A. E. Gilbert, D. H. Josephs, N. Ali, T. Dodev, L. Saul, I. Correa, L.

Roberts, E. Beddowes, A. Koers, C. Hobbs, S. Ferreira, J. L. Geh, C. Healy, M. Harries,

K. M. Acland, P. J. Blower, T. Mitchell, D. J. Fear, J. F. Spicer, K. E. Lacy, F. O. Nestle,

and S. N. Karagiannis. 2013. IgG4 subclass antibodies impair antitumor immunity in

melanoma. *J Clin Invest* 123: 1457-1474.

13.   Chung, A. W., T. K. Ho, P. Hanson-Manful, S. Tritscheller, J. M. Raynes, A. L. Whitcombe,

M. L. Tay, R. McGregor, N. Lorenz, J. R. Oliver, J. K. Gurney, C. G. Print, N. J. Wilson,

W. J. Martin, D. A. Williamson, M. G. Baker, and N. J. Moreland. 2020. Systems

immunology reveals a linked IgG3-C4 response in patients with acute rheumatic fever.

Immunol Cell Biol 98: 12-21.

14.   Vander Heiden, J. A., P. Stathopoulos, J. Q. Zhou, L. Chen, T. J. Gilbert, C. R. Bolen, R.

J. Barohn, M. M. Dimachkie, E. Ciafaloni, T. J. Broering, F. Vigneault, R. J. Nowak, S. H.

Kleinstein, and K. C. O'Connor. 2017. Dysregulation of B Cell Repertoire Formation in

Myasthenia Gravis Patients Revealed through Deep Sequencing. *J Immunol* 198: 1460-

1473.

15.   Huijbers, M. G., L. A. Querol, E. H. Niks, J. J. Plomp, S. M. van der Maarel, F. Graus, J.

Dalmau, I. Illa, and J. J. Verschuuren. 2015. The expanding field of IgG4-mediated

neurological autoimmune disorders. *European Journal of Neurology* 22: 1151-1161.

16.   Polonelli, L., J. Pontón, N. Elguezabal, M. D. Moragues, C. Casoli, E. Pilotti, P. Ronzi, A.

S. Dobroff, E. G. Rodrigues, M. A. Juliano, D. L. Maffei, W. Magliani, S. Conti, and L. R.

Travassos. 2008. Antibody complementarity-determining regions (CDRs) can display

differential antimicrobial, antiviral and antitumor activities. *PLoS One* 3: e2371.

17.   Liu, H., W. Pan, C. Tang, Y. Tang, H. Wu, A. Yoshimura, Y. Deng, N. He, and S. Li. 2021.

686   The methods and advances of adaptive immune receptors repertoire sequencing.

687   *Theranostics* 11: 8945-8963.

688   18.   Trück, J., A. Eugster, P. Barennes, C. M. Tipton, E. T. Luning Prak, D. Bagnara, C. Soto,

689   J. S. Sherkow, A. S. Payne, M.-P. Lefranc, A. Farmer, A. C. The, M. Bostick, and E.

690   Mariotti-Ferrandiz. 2021. Biological controls for standardization and interpretation of

691   adaptive immune receptor repertoire profiling. *eLife* 10: e66274.

692   19.   Horns, F., C. Vollmers, D. Croote, S. F. Mackey, G. E. Swan, C. L. Dekker, M. M. Davis,

693   and S. R. Quake. 2016. Lineage tracing of human B cells reveals the in vivo landscape of

694   human antibody class switching. *eLife* 5: e16578.

695   20.   Calonga-Solís, V., D. Malheiros, M. H. Beltrame, L. d. B. Vargas, R. M. Dourado, H. C.

696   Issler, R. Wassem, M. L. Petzl-Erler, and D. G. Augusto. 2019. Unveiling the Diversity of

697   Immunoglobulin Heavy Constant Gamma (IGHG) Gene Segments in Brazilian

698   Populations Reveals 28 Novel Alleles and Evidence of Gene Conversion and Natural

699   Selection. *Front Immunol* 10.

700   21.   Jonsson, S., G. Sveinbjornsson, A. L. de Lapuente Portilla, B. Swaminathan, R. Plomp,

701   G. Dekkers, R. Ajore, M. Ali, A. E. H. Bentlage, E. Elmér, G. I. Eyjolfsson, S. A.

702   Gudjonsson, U. Gullberg, A. Gylfason, B. V. Halldorsson, M. Hansson, H. Holm, Å.

703   Johansson, E. Johnsson, A. Jonasdottir, B. R. Ludviksson, A. Oddsson, I. Olafsson, S.

704   Olafsson, O. Sigurdardottir, A. Sigurdsson, L. Stefansdottir, G. Masson, P. Sulem, M.

705   Wuhrer, A.-K. Wihlborg, G. Thorleifsson, D. F. Gudbjartsson, U. Thorsteinsdottir, G.

706   Vidarsson, I. Jonsdottir, B. Nilsson, and K. Stefansson. 2017. Identification of sequence

707   variants influencing immunoglobulin levels. *Nature Genetics* 49: 1182-1191.

708   22.   Buck, D., E. Albrecht, M. Aslam, A. Goris, N. Hauenstein, A. Jochim, S. Cepok, V.

709   Grummel, B. Dubois, A. Berthele, P. Lichtner, C. Gieger, J. Winkelmann, and B. Hemmer.

2013. Genetic variants in the immunoglobulin heavy chain locus are associated with the

IgG index in multiple sclerosis. *Ann Neurol* 73: 86-94.

23.   Keyeux, G., G. Lefranc, and M.-P. Lefranc. 1989. A multigene deletion in the human IGH

constant region locus involves highly homologous hot spots of recombination. *Genomics*

5: 431-441.

24.   Bashirova, A. A., W. Zheng, M. Akdag, D. G. Augusto, N. Vince, K. L. Dong, C. O'HUigin,

and M. Carrington. 2021. Population-specific diversity of the immunoglobulin constant

heavy G chain (IGHG) genes. *Genes Immun* 22: 327-334.

25.   Lefranc, M. P., G. Lefranc, G. de Lange, T. A. Out, P. J. van den Broek, J. van Nieuwkoop,

J. Radl, A. N. Helal, H. Chaabani, E. van Loghem, and et al. 1983. Instability of the human

immunoglobulin heavy chain constant region locus indicated by different inherited

chromosomal deletions. *Mol Biol Med* 1: 207-217.

26.   Lefranc, M.-P., and G. Lefranc. 2012. Human Gm, Km, and Am Allotypes and Their

Molecular Characterization: A Remarkable Demonstration of Polymorphism. In

Immunogenetics: Methods and Applications in Clinical Practice. F. T. Christiansen, and B.

D. Tait, eds. Humana Press, Totowa, NJ. 635-680.

27.   Lefranc, M.-P., G. Lefranc, and T. H. Rabbitts. 1982. Inherited deletion of immunoglobulin

heavy chain constant region genes in normal human individuals. *Nature* 300: 760-762.

28.   van Erp, E. A., W. Luytjes, G. Ferwerda, and P. B. van Kasteren. 2019. Fc-Mediated

Antibody Effector Functions During Respiratory Syncytial Virus Infection and Disease.

Front Immunol 10.

29.   Jefferis, R., J. Lund, and J. D. Pound. 1998. IgG-Fc-mediated effector functions: molecular

definition of interaction sites for effector ligands and the role of glycosylation.

Immunological Reviews 163: 59-76.

30. Vander Heiden, J. A., G. Yaari, M. Uduman, J. N. H. Stern, K. C. O'Connor, D. A. Hafler,

F. Vigneault, and S. H. Kleinstein. 2014. pRESTO: a toolkit for processing high-throughput

sequencing raw reads of lymphocyte receptor repertoires. *Bioinformatics* 30: 1930-1932.

31. Gupta, N. T., J. A. Vander Heiden, M. Uduman, D. Gadala-Maria, G. Yaari, and S. H.

Kleinstein. 2015. Change-O: a toolkit for analyzing large-scale B cell immunoglobulin

repertoire sequencing data. *Bioinformatics* 31: 3356-3358.

32. Charney, A. W., N. W. Simons, K. Mouskas, L. Lepow, E. Cheng, J. Le Berichel, C. Chang,

R. Marvin, D. M. Del Valle, S. Calorossi, A. Lansky, L. Walker, M. Patel, H. Xie, N. Yi, A.

Yu, G. Kang, A. Mendoza, L. E. Liharska, E. Moya, M. Hartnett, S. Hatem, L. Wilkins, M.

Eaton, H. Jamal, K. Tuballes, S. T. Chen, A. Tabachnikova, J. Chung, J. Harris, C.

Batchelor, J. Lacunza, M. Yishak, K. Argueta, N. Karekar, B. Lee, G. Kelly, D. Geanon, D.

Handler, J. Leech, H. Stefanos, T. Dawson, I. Scott, N. Francoeur, J. S. Johnson, A. Vaid,

B. S. Glicksberg, G. N. Nadkarni, E. E. Schadt, B. D. Gelb, A. Rahman, R. Sebra, G.

Martin, C. Agashe, P. Agrawal, A. Akyatan, K. Alesso-Carra, E. Alibo, K. Alvarez, A.

Amabile, S. Ascolillo, R. Bailey, P. Begani, P. B. Correra, S.-A. Brown, M. Buckup, L.

Burka, L. Cambron, G. Carrara, S. Chang, J. Chien, M. Chowdhury, C. C. Bozkus, P.

Comella, D. Cosgrove, F. Cossarini, L. Cotter, A. Dave, B. Dayal, M. Dhainaut, R.

Dornfeld, K. Dul, N. Eber, C. Elaiho, F. Fabris, J. Faith, D. Falci, S. Feng, B. Fennessy, M.

Fernandes, S. Gangadharan, J. Grabowska, G. Gyimesi, M. Hamdani, M. Herbinet, E.

Herrera, A. Hochman, G. E. Hoffman, J. Hook, L. Horta, E. Humblin, S. Karim, J. Kim, D.

Lebovitch, G. Lee, G. H. Lee, J. Lee, M. Leventhal, K. Lindblad, A. Livanos, R. Machado,

Z. Mahmood, K. Mar, S. Maskey, P. Matthews, K. Meckel, S. Mehandru, C. Mercedes, D.

Meyer, G. Mollaoglu, S. Morris, K. Nie, M. Nisenholtz, G. Ofori-Amanfo, K. Onel, M.

Ounadjela, V. Patel, C. Pruitt, S. Rathi, J. Redes, I. Reyes-Torres, A. Rodrigues, A.

Rodriguez, V. Roudko, E. Ruiz, P. Scalzo, P. Silva, A. S. Schanoski, M. Straw, S.

Tabachnikova, C. Teague, B. Upadhyaya, V. Van Der Heide, N. Vaninov, D. Wacker, H.

Walsh, C. M. Wilk, J. Wilson, K. M. Wilson, L. Xue, N.-a. Yeboah, S. Young, N. Zaks, R. Zha, T. Marron, N. Beckmann, S. Kim-Schulze, S. Gnjatic, M. Merad, and C.-B. T. The Mount Sinai. 2020. Sampling the host response to SARS-CoV-2 in hospitals under siege. *Nature Medicine* 26: 1157-1158.

33. Woolley, C., J. Chariker, E. Rouchka, E. Ford, E. Hudson, S. Waigel, M. Smith, and T. Mitchell. 2022. Reference long-read isoform-aware transcriptomes of four human peripheral blood lymphocyte subsets *G3: Genes, Genomes, Genetics*

34. Nouri, N., and S. H. Kleinstein. 2018. A spectral clustering-based method for identifying clones from high-throughput B cell repertoire sequencing data. *Bioinformatics* 34: i341-i349.

35. Nouri, N., and S. H. Kleinstein. 2020. Somatic hypermutation analysis for improved identification of B cell clonal families from next-generation sequencing data. *PLOS Computational Biology* 16: e1007977.

36. Hoehn, K. B., O. G. Pybus, and S. H. Kleinstein. 2022. Phylogenetic analysis of migration, differentiation, and class switching in B cells. *PLOS Computational Biology* 18: e1009885.

37. Rodriguez, O. L., W. S. Gibson, T. Parks, M. Emery, J. Powell, M. Strahl, G. Deikus, K. Auckland, E. E. Eichler, W. A. Marasco, R. Sebra, A. J. Sharp, M. L. Smith, A. Bashir, and C. T. Watson. 2020. A Novel Framework for Characterizing Genomic Haplotype Diversity in the Human Immunoglobulin Heavy Chain Locus. *Front Immunol* 11.

38. Lefranc, M. P. 2001. IMGT, the international ImMunoGeneTics database. *Nucleic Acids Res* 29: 207-209.

39. Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34: 3094-3100.

40. Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, and G. P. D. P. Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078-2079.

41.   Martin, M., M. Patterson, S. Garg, S. O Fischer, N. Pisanti, G. W. Klau, A. Schöenhuth,
and T. Marschall. 2016. WhatsHap: fast and accurate read-based phasing. *bioRxiv*:
085050.
42.   Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local
alignment search tool. *J Mol Biol* 215: 403-410.
43.   Fu, L., B. Niu, Z. Zhu, S. Wu, and W. Li. 2012. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics (Oxford, England)* 28: 3150-3152.
44.   Gadala-Maria, D., G. Yaari, M. Uduman, and S. H. Kleinstein. 2015. Automated analysis
of high-throughput B-cell sequencing data reveals a high frequency of novel
immunoglobulin V gene segment alleles. *Proceedings of the National Academy of*
Sciences 112: E862-E870.
45.   Gadala-Maria, D., M. Gidoni, S. Marquez, J. A. Vander Heiden, J. T. Kos, C. T. Watson,
K. C. O'Connor, G. Yaari, and S. H. Kleinstein. 2019. Identification of Subject-Specific
Immunoglobulin Alleles From Expressed Repertoire Sequencing Data. *Front Immunol* 10.
46.   Gidoni, M., O. Snir, A. Peres, P. Polak, I. Lindeman, I. Mikocziova, V. K. Sarna, K. E. A.
Lundin, C. Clouser, F. Vigneault, A. M. Collins, L. M. Sollid, and G. Yaari. 2019. Mosaic
deletion patterns of the human antibody heavy chain gene locus shown by Bayesian
haplotyping. *Nature Communications* 10: 628.
47.   Peres, A., M. Gidoni, P. Polak, and G. Yaari. 2019. RAbHIT: R Antibody Haplotype
Inference Tool. *Bioinformatics* 35: 4840-4842.
48.   Al'Khafaji, A. M., J. T. Smith, K. V. Garimella, M. Babadi, M. Sade-Feldman, M. Gatzen,
S. Sarkizova, M. A. Schwartz, V. Popic, E. M. Blaum, A. Day, M. Costello, T. Bowers, S.
Gabriel, E. Banks, A. A. Philippakis, G. M. Boland, P. C. Blainey, and N. Hacohen. 2021.
High-throughput RNA isoform sequencing using programmable cDNA concatenation.
bioRxiv: 2021.2010.2001.462818.

49.   Brochu, H. N., E. Tseng, E. Smith, M. J. Thomas, A. M. Jones, K. R. Diveley, L. Law, S.

G. Hansen, L. J. Picker, M. Gale, and X. Peng. 2020. Systematic Profiling of Full-Length

Ig and TCR Repertoire Diversity in Rhesus Macaque through Long Read Transcriptome

Sequencing. *The Journal of Immunology* 204: 3434.

50.   Kidd, M. J., Z. Chen, Y. Wang, K. J. Jackson, L. Zhang, S. D. Boyd, A. Z. Fire, M. M.

Tanaka, B. A. Gaëta, and A. M. Collins. 2012. The Inference of Phased Haplotypes for

the Immunoglobulin H Chain V Region Gene Loci by Analysis of VDJ Gene

Rearrangements. *The Journal of Immunology* 188: 1333.

51.   Kirik, U., L. Greiff, F. Levander, and M. Ohlin. 2017. Parallel antibody germline gene and

haplotype analyses support the validity of immunoglobulin germline gene inference and

discovery. *Mol Immunol* 87: 12-22.

52.   Irvine, E. B., and G. Alter. 2020. Understanding the role of antibody glycosylation through

the lens of severe viral and bacterial diseases. *Glycobiology* 30: 241-253.

53.   Alter, G., T. H. M. Ottenhoff, and S. A. Joosten. 2018. Antibody glycosylation in

inflammation, disease and vaccination. *Seminars in Immunology* 39: 102-110.

54.   Collin, M. 2020. Antibody glycosylation as an immunological key in health and disease.

Glycobiology 30: 200-201.

55.   Plomp, R., L. R. Ruhaak, H.-W. Uh, K. R. Reiding, M. Selman, J. J. Houwing-Duistermaat,

P. E. Slagboom, M. Beekman, and M. Wuhrer. 2017. Subclass-specific IgG glycosylation

is associated with markers of inflammation and metabolic health. *Scientific Reports* 7:

12325.

56.   Trampert, D. C., L. M. Hubers, S. F. J. van de Graaf, and U. Beuers. 2018. On the role of

IgG4 in inflammatory conditions: lessons for IgG4-related disease. *Biochimica et*

Biophysica Acta (BBA) - Molecular Basis of Disease 1864: 1401-1409.

 Figure Legends:

Figure 1. Overview of Ab structure and FLAIRR-seq molecular method. (A) Schematic

representation of the IGH locus, heavy chain transcript structure, and functional IG protein. (B)

Comparative coverage across the heavy chain transcript of commonly used AIRR-seq methods

compared with FLAIRR-seq. (C) FLAIRR-seq molecular pipeline: RNA (brown) was converted to

first-strand cDNA (red) using the 5' RACE method, incorporating a 5' TSO-UMI (pink) via template

switch. Second strand amplification specifically targeted IgG and IgM molecules through priming

of the 5' TSO-UMI and the 3' constant region IGH exon 3 (CH3) for IgG, or CH4 for IgM. A 16bp

barcode was incorporated into the 3' CH3/CH4 primers to enable sample multiplexing post-

amplification. (D) IGV screenshot showing near full-length single molecule structure of IGHG4

FLAIRR-seq transcripts.

Figure 2. FLAIRR-seq shows robust characterization of V, D, and J genes. (A) Spearman

ranked correlations and p-values of V, D, and J gene usage frequencies identified by FLAIRR-

seq performed on matched total PBMC and purified B cells. (B) Heatmap of Spearman ranked

correlations and p-values of V, D and J gene usage frequencies between FLAIRR-seq and AIRR-

seq processed samples. (C) Boxplots of somatic hypermutation frequencies defined by FLAIRR-

seq or AIRR-seq in IgM- and IgG-specific repertoires. (D) Boxplots of CDR3 length defined by

FLAIRR-seq or AIRR-seq analysis in both IgM- and IgG-specific repertoires. (E) Spearman

ranked correlation of V, D, and J gene usage frequencies between FLAIRR-seq-based and Iso-

Seq-based repertoire profiling. Significant differences between FLAIRR-seq and AIRR-seq data

indicated by * ($p < 0.05$) or ** ($p < 0.01$).

Figure 3. FLAIRR-seq provides novel IGHC resolution for allelic discovery and allows

variable gene haplotyping. (A) Overview of experimental design and pipeline overviews of

genotyping by IGenotyper (gDNA) and FLAIRR-seq (RNA). (B) Schematic depicting IGHC alleles

identified by IGenotyper, partitioned by identification as (i) exact matches to documented IMGT

alleles, (ii) novel alleles that are not in IMGT, or (iii) extended alleles. (C) Pie chart and stacked

bar graph representing the total number of alleles and fraction of each category identified per

IGHC gene as identified by either IGenotyper or FLAIRR-seq. Bar charts showing number of

IGHC alleles from FLAIRR-seq that were resolved, ambiguous or unresolved when compared to

IGenotyper alleles. * Indicates additional allele found due to IGHG4 duplication. (D) Table

summarizing novel and extended alleles resolved by IGenotyper data. Extended alleles are

denoted by *(allele number)-FL and novel alleles are denoted by FL_(number) alleles resolved

by FLAIRR-seq are marked with a dot (•). (E) Venn diagrams showing number of IGHV haplotype

allele/deletion calls when using IGHJ6 or IGHM anchors for each haplotype. Tile plots showing

IGHV gene haplotypes inferred using either IGHJ6 anchors or IGHM anchors for one sample.

Dark gray represents a deletion (DEL), off-white a non-reliable allele annotation (NRA), and light

gray represents an unknown allele (Unk). Non-reliable alleles are annotated with an asterisk (*).

Figure 4. FLAIRR-seq resolves subisotype specific repertoire diversity. (A) Bar plots

showing distribution of unique VDJ sequences across isotypes, subisotypes, and subisotype

alleles in one representative sample, 1013, characterized by FLAIRR-seq. (B) Circos plots

showing V family gene usage frequency within each subisotype allele for sample 1013. (C)

Boxplots of V gene family usage frequencies within IGHG1, IGHG2, IGHG3, and IGHG4

repertoires across all ten individuals. (D) Principal component analysis of V gene family usage by

subisotype; plot includes the first two principal components, and individual repertoires are colored

by IGHG subisotype. (E) Boxplots showing sequence frequency of IGHV1 and IGHV3 family

genes by subisotype across all 10 samples.

Figure 5. FLAIRR-seq resolves subisotype-specific clonal expansion and facilitates

haplotype analysis of CSR in a patient hospitalized for COVID-19. (A) Overview of

experimental design: whole blood-derived RNA was collected on days 1, 4, 8 and 13 post-

hospitalization and used for FLAIRR-seq profiling. (B) Bar plot showing the percentage of clones

represented by each subisotype across timepoints. (C) Simpson's diversity index (q=2) for all

clones in each subisotype across four timepoints; IgG4 not included due to low sequence counts.

(D) Polarity, or the fraction of clones needed to comprise 80% of the repertoire, reported as

fraction of total subisotype-specific repertoires across time. (E) Distribution of a single clone

"9900" across isotypes and subisotypes over time, suggesting CSR of this clone. (F) Phylogenetic

tree constructed from sequences/members of clone 9900, with the inferred germline sequence

as the outgroup (star). Shapes and colors of tips (sequences) indicate time point and

isotype/subisotype. The scale bar represents the number of mutations between each node in the

tree. The subclade within the red box is represented by multiple time points and subisotypes,

providing evidence of CSR. (G) Tile plot showing the assignment of IGHC alleles to their

respective haplotypes, based on the frequency of observations in which each IGHC allele was

linked to each respective allele of heterozygous V genes, IGHV3-7 and IGHV3-48; light gray

denotes IGHC alleles for which haplotype assignment was not possible. Analysis of sequences

in (F) revealed that the IGHG1 and IGHG2 alleles represented in the phylogeny came from the

same haplotype (IGHG1*02/*07, IGHG2*08, IGHM_FL_2).

Table 1: Comparative analysis metrics between AIRR-seq and FLAIRR-seq on matched samples

Sample ID	Isotype	Method	RNA input (ng)	Input reads into Immcantation[a]	Post assembly /Single reads[b]	Unique VDJ[c]	Unique Clones[d]	Unique CDR3[e]
0007	IgG	AIRR	100	841627	15217	3164	1008	1187
		FLAIRR	335	181109	17881	4880	1358	1520
201c	IgG	AIRR	100	1509063	25015	4568	1221	1371
		FLAIRR	296	221301	36053	10819	2145	2388
203c	IgG	AIRR	100	1006965	23085	5793	1436	1705
		FLAIRR	106	475703	14923	3778	993	1138
602c	IgG	AIRR	100	1275902	38078	7736	1542	2217
		FLAIRR	149	344906	33903	9225	1828	2357
705c	IgG	AIRR	100	1113660	15358	1362	557	576
		FLAIRR	268	279601	18627	2328	1063	1081
1008	IgG	AIRR	100	1310350	56848	12217	3005	3307
		FLAIRR	335	238820	57020	18235	3505	3813
1013	IgG	AIRR	100	1158222	25250	5027	1795	1964
		FLAIRR	222	396322	27871	7487	2565	2744
2008	IgG	AIRR	100	835928	21681	3900	1235	1368
		FLAIRR	240	418906	16441	4258	1308	1380
4002	IgG	AIRR	100	632272	8949	1112	727	772
		FLAIRR	335	398838	19692	5377	1719	1887
5001	IgG	AIRR	100	695884	13930	1885	1134	1244
		FLAIRR	335	313843	42845	12146	2226	2381
0007	IgM	AIRR	100	841627	12777	9154	7924	8163
		FLAIRR	335	331380	46384	14926	12367	12851
201c	IgM	AIRR	100	1509063	26850	17888	14281	14622
		FLAIRR	296	148355	98069	12803	9353	9573
203c	IgM	AIRR	100	1006965	24623	18189	16295	16629
		FLAIRR	106	509536	37527	10636	9315	9419
602c	IgM	AIRR	100	1275902	12306	8378	7704	7833
		FLAIRR	149	273103	48897	16406	14495	14745
705c	IgM	AIRR	100	1113660	19484	14273	12936	13413
		FLAIRR	268	435391	79251	17417	15706	16249
1008	IgM	AIRR	100	1310350	20264	12710	7878	8368
		FLAIRR	335	181063	47531	15284	8745	9276
1013	IgM	AIRR	100	1158222	12769	9012	8057	8241
		FLAIRR	222	262617	52508	17812	15291	15759
2008	IgM	AIRR	100	835928	11608	8551	5838	6206
		FLAIRR	240	502354	47596	14132	10225	10674
4002	IgM	AIRR	100	632272	18411	11607	11184	11367
		FLAIRR	335	489287	91732	22379	19000	21532
5001	IgM	FLAIRR	335	313843	10018	12146	2226	2381
		FLAIRR	355	171688	18944	6164	4866	5011

FIGURE 3.

A.

Severe COVID-19 hospitalization

BD   BD   BD   BD

Day1   Day4   Day8   Day13   (Post hospital admission)

Whole blood RNA isolation over four timepoints

IGHG and IGHM FLAIRR-seq

B.

Percentage of clones represented in each subisotype

C.

IGHG subisotype diversity over time

D.

IGHG subisotype polarity over time

E.

Clone 9900

F.

Clone 9900

Day1 ▲
Day4 ■
Day8 ●
Day13 ◆
Germline ★
IGHM ●(orange)
IGHG1 ●(blue)
IGHG2 ●(red)

0.01

G.

Alleles

*02/*07
*03/*06
*02
*08
*11
FL_2
Unk

IGHV3-7*01
IGHV3-7*03
IGHV3-48*01
IGHV3-48*03

IGHG1 IGHG2 IGHG3 IGHG4 IGHM