VOCAL SIMILARITY AND DIFFERENCES IN FOUR SPECIES OF THE GENUS *ANTHUS*: ACOUSTIC FEATURE ANALYSIS OF SOME COMMON CALLS

Marco Dragonetti

Gruppo Ornitologico Maremmano, c/o Museo di Storia Naturale di Grosseto, Strada Corsini 5, 58100, Grosseto, Italy. Email: dragom@quipo.it ORCiD: 0000-0001-8368-4150

ABSTRACT

Birds of the same genus often share similar call repertoires, the aim of this paper is to find species-specific and common acoustic features across species. This can be a useful tool for identification purposes and for studying intra-interspecific communication. Similar flight call (*tsip*) in two closely related species (*Anthus pratensis*, *Anthus spinoletta*) were studied to find characteristics, that allows to discriminate the two species with acoustic means. Three different call types (*tsip*, *soft* and *alarm*) of four species of the genus Anthus (*A. pratensis*, *A. spinoletta*, *A. petrosus*, *A. cervinus*) were also studied to find whether these common call types show different degrees of similarity. Discriminant function analysis correctly classified 98.4% of *A. pratensis* and *A. spinoletta* flight calls. Three acoustic parameters showed the highest discrimination power: the frequency modulations, the maximum frequency value and the minimum frequency value of the peak frequency contour. Using these three values I proposed a simpler procedure for recognizing these two species, that allowed a correct classification of 96% of calls.The three call types of the four *Anthus* species were studied using cross correlation among spectrogram contours. *Alarm* calls of the four species showed stronger similarity, while the other call types were more distinctive, with *soft* call seeming to have a lower similarity between species and hence a higher distinctive power. These results suggest the hypothesis that *alarm* call is similar, because it retains features of a common ancestor easing heterospecific communication, while the other calls showed decreasing similarity and more species-specific features.

KEYWORDS: call similarity - acoustic recognition - genus *Anthus* - vocal behaviour - taxonomy

INTRODUCTION

Investigating the acoustic features of the vocalisations among species of the same genus may help to better understand their communication strategies and to shed light about the variable acoustic structures across related taxa. Moreover, acoustic analysis can be an useful tool for identification of morphologically similar species. Common call types used all year round offer a fruitful field of investigation for these purposes.

Meadow pipits (*Anthus pratensis*) and Water pipits (*Anthus spinoletta*) are morphologically similar species, they have partly overlapping wintering range and often share feeding areas (Cramp 1983; Alström and Mild 2003; Brichetti and Fracasso 2007). These congeneric similar species cope with problems of intraspecific and interspecific communication, therefore they should have finely tuned capability of call discrimination between species as found by Elfström (1992) for *A. pratensis* and *A. petrosus*, but in some cases they might

also benefit from interspecific communication, e.g. in mixed flock with contact calls and with common

danger signals. Outside the breeding season the vocal repertoire of these two species rely mainly on two call

types, a *tsip* call used more often in flight and a *soft* call typically produced between members of a flock as a

contact signal (Elfström 1992; Alström and Mild 2003). The former shows a very similar spectrographic

structure for both species (Alström and Mild 2003; Bergmann et al. 2008), although it can sound slightly

different for a trained ear (Brichetti and Fracasso 2007).

Elfström (1992) showed that calls of *A. pratensis* and *A. petrosus* contain species-specific characteristics and

that these species are able to discriminate between conspecific and heterospecific calls of similar

configuration at a distance. Finding comparable species-specific differences also for *tsip* call of *A. pratensis*

and *A. spinoletta* would support the hypothesis that this call type is mainly used for intraspecific

communication.

In addition, finding species-specific features of a common flight call can enable vocal identification of

species that have similar winter plumage and behaviour. This can be a useful tool for field work when the

two sister species share the same habitat and in the case of passive recording of migratory birds. Indeed,

flight call recording and identification of nocturnal migrant birds is a method increasingly used (Hamilton

1962; Gal 2003; Farnsworth 2005). Some evidence suggest that birds of the genus *Anthus*, although mainly

diurnal migrants (Elkins 2005), can migrate also by night (Ebenhöh 1998; Gal 2003; Wang 2019; Briedis et

al. 2020), therefore acoustic identification turns to be interesting for these species too.

For years bird calls were considered innate (see for example Hamilton 1962), because subjects raised away

from their own species developed abnormal songs but normal calls. After the discovery of developmental

plasticity in the calls of several bird species (Mudinger 1970; Mudinger 1979; Marler and Slabbekoorn

2004), learning appeared to play a role in the call repertoire of some passerines. Particularly flight calls of

cardueline finches showed remarkable plasticity and dialects were discovered for other call types (Baker et

al. 2000; Marler and Slabbekoorn 2004). Although bird calls have not been studied as extensively as bird

song, today it is commonly accepted that some calls are learned while others appear to be innate (Zann 1985;

Williams 2008). Some calls can share very similar characteristics in several species of related taxa (Stefanski

and Falls 1972; Fallow et al. 2011; Gayk et. al 2021), while others can be more or less distinctive for

intraspecific recognition and/or communication (Elfström 1992; Kroodsma 2000; Fijen 2014). Genetic

relatedness should favour a stronger similarity mainly for those innate calls that convey crucial information

involved with immediate issues of life and death. On the other hand there are also calls, either genetically

determined or developed by learning processes, which contains distinctive features and are important for

species-specific recognition (Nowicki 1989, Marler and Slabbekoorn 2004).

In this paper I have investigated the acoustic features of a common flight call (*tsip*) of two sister species (*A.*

pratensis, A. spinoletta). The aim is to find whether there are objective measurable differences among vocal

parameters of *tsip* call, which is fairly similar for these two species. If consistent quantitative differences are

found, it is interesting to find a measurable vocal threshold that allows a safe and relatively easy

identification of the species. Finding quantitative differences for one call type poses the question whether

this pattern is also found for other common call types in closely related species of the genus *Anthus*.

Interestingly, the acoustic similarity of some call types in different species can be the result of retaining

features from a common ancestor (De Kort and Ten Cate 2001) or convergence in acoustic structure on the

most effective design for a particular environment (Marler 1955; Marler and Slabbekoorn 2004). Broadening

the view on other species of the genus *Anthus* and investigating the acoustic similarity and differences of

different call types seemed me an useful approach to shed some light on these topics. Therefore three

different call types (*tsip*, *soft* and *alarm*) of four species (*A. pratensis, A. spinoletta, A. petrosus, A. cervinus*)

were studied. The results of this analysis, although have to be considered preliminar given the limitations of

the dataset, can be useful for the study of intra/interspecific vocal communication and genetic relatedness

among sister species of the genus *Anthus*.

METHODS

I recorded calls of *Anthus pratensis* and *Anthus spinoletta* in central Italy with a Fostex FR2 digital recorder

in wav uncompressed format (48 kHz sampling rate, 16 bit resolution). The microphone was a Sennheiser

MKH20 in the focus of a 22" Telinga parabolic dish. The species identification was made visually (with

binoculars, cameras and spotting scopes) before starting the recording session. Moreover almost all

recordings of *A. spinoletta* were made in nesting areas of this species, where *A. pratensis* is absent.

I made 34 good quality recordings of *A. pratensis* and 31 of *A. spinoletta* from 2004 to 2021 in different

places of Central Italy. The minimum distance between sampling places for each species, when recordings

were made during the same season, was 25 km; the minimum time gap, when recordings were made in the

same zone, was 26 months, therefore it is very unlikely that the same birds were recorded more than once,

given that both species are not resident in Central Italy. The average of the acoustic parameters of all good

quality calls was calculated from each recording, yelding a dataset of 65 independent measures. The *tsip* call

type (in flight) was selected from these recordings following the classification and spectrographic

representation available in the literature (Cramp 1983; Elfström 1992; Alström and Mild 2003; Bergmann

2008; Fijen 2014; Garner et al. 2015). All spectrograms were made with Raven Pro 1.6.1 (K. Lisa Yang

Center for Conservation Bioacoustics 2019): window and DFT size 1024 samples, overlap 75%, window

type Hann (3dB filter bandwidth 67.5 Hz).

One *tsip* call template of the two species (Fig. 1) was chosen among good quality recordings, with a signal

to noise ratio above 10 calculated on the amplitude values expressed by Raven in arbitrary units. This call

type has at least three visible harmonics and the second is always the dominant one, therefore a bandpass

filter was used for limiting the analysis to this harmonic. As shown in Figure 1 most sound energy of the

template *tsip* calls is concentrated between 4 and 8.5 kHz, for this reason I used the band limited energy

detector of Raven Pro to select all the *tsip* calls to be analysed. The detector parameters were: minimum

frequency 4000 Hz, maximum frequency 8500 Hz, minimum duration 0.03 s, maximum duration 0.2 s,

minimum separation 0.02 s, minimum occupancy 50%, SNR (signal to noise ratio) threshold 5 dB. The

detections showing unwanted signals (e.g. overlapped calls of the target species or of other species, various

noises) were discarded. A detailed acoustic analysis was performed on 212 *A. pratensis* and 165 *A.*

spinoletta tsip calls and data were averaged for each recording, yielding 34 and 31 independent data for the

two species; the following parameters were calculated by means of the appropriate Raven Pro functions:

Center Frequency (CF), Frequency Contour Percentile 75 (FC75), Peak Frequency Contour Max Frequency

(PFCMa), Peak Frequency Contour Min Frequency (PFCMi), Duration 90% (DUR90). The difference

between PFCMa and PFCMi was calculated as a measure of the frequency bandwidth (BW). I counted for

each call the number of relative maximum plus minimum values (RMM) found by the FC75 function. The

time interval between the beginnings of consecutive call selections (TI) was measured. In 2020/2021 further

5 recordings of *A. pratensis tsip* calls were made in Northern and Central Italy and other 5 recordings of *A.*

spinoletta tsip calls from Central Europe and Northern Italy were provided by Xeno-Canto recordists in wav

uncompressed original format. These 10 recordings (for a total of 45 calls) were used to test the ability of

DFA to discriminate the two species.

The three different call types (*tsip*, *soft* and *alarm*) of the four *Anthus* species (*A. pratensis, A. spinoletta, A.*

petrosus, A. cervinus) were compared following the classification and spectrographic representation of

Elfström (1992) and Alström and Mild (2003). Figure 1 shows the template calls chosen for each call type

of the four species. 13 recordings of *tsip* call and 12 recordings of *soft* calls for *A. pratensis* and *A. spinoletta*

were randomly selected from the dataset above described, while 13 recordings of *tsip* call for *A. cervinus* and

A. petrosus, 12 recordings of *soft* call for *A. cervinus* and *A. petrosus* and 10 recordings of *alarm* call for *A.*

cervinus, *A. petrosus*, *A. pratensis* were audio files published on Xeno-Canto (randomly selected among all

those available), 10 recordings of *alarm* call for *A. spinoletta* were made in Central Italy by the author as

specified above. Unfortunately most of Xeno-Canto recordings were not available in an original non-

compressed (wav) format, but only as mp3 files. Since all the recordings were not homogeneous as file

format, bitrate, recording tools and editing processes, a detailed analysis of all the acoustic parameters, as

made for *A. pratensis* and *A. spinoletta tsip* call, could have yielded unreliable results. To avoid this

drawback I preliminarily tested the capability of Raven Pro FC75 function to assess in a sufficiently precise

manner general spectral and temporal characteristics of the same calls recorded in three different ways: wav

original uncompressed format, mp3 compressed file of medium quality (170/210 kbps), mp3 compressed

format of low quality (145/185 kbps). The results of this test (shown in the supplementary material)

suggested that FC75 function was able to assess correctly general bioacoustic characteristics for all the

different quality levels. Similarly Araya-Salas et al.(2019) found that compression did not significantly bias

the acoustic measurements of similarity. Therefore the comparison among the three call types in the four

Anthus species was made using only the FC75 parameter. From each recording of *alarm* call 7 to 9 call

contours were averaged yelding 10 independent data for each *Anthus* species, from each recording of *tsip* call

4 to 8 call contours were averaged yelding 13 data, from each recording of *soft* call 3 to 9 call contours were

averaged yelding 12 data. The total number of contours averaged for each call type and for each species was

always 84. The call selection was made with the same method described above, using the band limited

energy detector of Raven Pro (band limits: 4000 - 8500 Hz, 3000 - 7000 Hz, 3500 - 7000 Hz for *tsip*, *soft* and

alarm call respectively). All recordings were filtered with a bandpass filter for limiting the analysis to the

dominant harmonic where most of sound energy was found.

Statistical analysis.

Where not specified otherwise all the statistical tests were performed using PAST v. 4.09 package (Hammer

et al. 2001). Multivariate normality of data was checked with omnibus test by Doornik and Hansen (2008).

The comparison of *A. pratensis* versus *A. spinoletta tsip* call was performed with PERMANOVA based on

Euclidean distance measure (Anderson 2001). Principal components analysis (PCA) was used to find

hypothetical variables (components) accounting for as much as possible of the variance in these multivariate

data (Legendre and Legendre 1998). The PCA and PERMANOVA were performed including all the seven

varaiables shown in Table 1. Coefficient of variation (CV) was calculated with the following formula: CV

$=100*(1+1/4n)*sd/m$ (Scherrer 1984; Aubin et al. 2004), where n = number of observations, SD = standard

deviation, m = mean. I have also calculated a CVr as the ratio of interspecific (CVis) versus average

intraspecific coefficient of variation (CVin). CVr is a measure of the identification power of each vocal

parameter, if it is higher than 1 the interspecific differences are heavier than the intraspecific ones and of

course the higher is CVr value the better is the discrimination power of the parameter. The equivalence of the

variance/covariance on multivariate samples was checked with Box's M test (Rencher 2002); this is a

prerequisite to perform the discriminant function analysis (DFA), used to find the differences among

acoustic parameters of *A. pratensis* versus *A. spinoletta tsip* calls.

The comparison among four *Anthus* species (*A. pratensis, A. cervinus, A. spinoletta, A. petrosus*) was

performed using FC75 to find similarity and differences in the three different call types (*alarm*, *tsip*, *soft*).

The randomly selected call contours were averaged for each recording and an averaged contour was then

calculated for the four species in the three different call types. Each averaged contour represented a series of

frequency data evenly spaced in time (5 ms apart). Mantel correlogram (Legendre and Legendre 1998), that

is a multivariate extension to autocorrelation based on any similarity or distance measure, made a first

comparison among all four species contours of each call type. The correlation index of Mantel test showed

the similarity between the averaged call contours shown in Figure 3. A second approach was made with a

more classical cross-correlation comparing the time series obtained with the FC75 function. All paired

comparisons were performed between average call contours from every recording of the four species for

each different call type with the following constraints: each recording must be used only once, all recordings

must be used, each pair of recording comparison was randomly chosen. This scheme yielded for example 10

comparisons for *alarm* call of *A. pratensis* versus *A. spinoletta*, 10 comparisons of *A. pratensis* versus *A.*

cervinus ...and so on. These values of cross correlation were averaged for each pair of comparisons among

the four species and for each call type. This way it was obtained a similarity value for each possible

comparison among the four species and a mean similarity value for all the comparisons among the four

species for each call type (Table 1 row 8). The average correlation index was calculated at two different lags

(lag 0 and lag m of maximum correlation index); the average value of the lag accounting the maximum

correlation index was also calculated. Of course a higher correlation index at lag 0 or near 0 expresses a

higher similarity among different species for that particular call type. The cross-correlation was calculated

using the R function ccf (R core Team 2020).

RESULTS

Comparison between *tsip* call of *A. pratensis* versus *A. spinoletta*. Table 1 shows *tsip* call parameters of

A. pratensis (n= 34; each data is average of 3 to 10 calls, mean 6.2) and *A. spinoletta* (n= 31; each data is

average of 2 to 9 calls, mean 5.3). Doornik and Hansen (2008) omnibus test indicates a highly significant

(p<0.001) departure from normality of these *tsip* call data. Therefore the two species were compared by

means of non parametric PERMANOVA test, which showed a highly significant difference (F=45; P<0.001)

of acoustic features.

Principal component analysis (PCA) of all multivariate data (Figure 2) was performed. The three first

components accounted for 89.4% of the variance and PCA loadings of the variables showed values ranging

between 0.33 (DUR90 and RMM), 0.34 (BW) and 0.38 (CF, PCFMa, PCFMi), but TI showed a lower value

(0.28). Since TI has several missing data and also a very high CVin (= 78.5) with a quite low CVr (Table 1),

it was excluded from further analyses because of its unreliability.

CVr was definitely higher than 1 for all acoustic parameters (except BW that had the lower value of 1.03,

Table 1) showing a good potential predictor power of these acoustic features for the recognition of the two

sister species.

The Box's M test (Rencher 2002) for equality of variance-covariance of multivariate data (excluding TI

parameter) yielded a not significant result (F=1,38). Moreover the normality test (Doornik and Hansen 2008)

showed a slightly significant value (p=0.045), accounting for a slight departure from normality of the data.

Hence the prerequisites to perform DFA were satisfied, since Sever et al. (2005) showed the DFA to be

highly robust to non-normal data.

DFA correctly classified 98.4% of calls with jack-knifed validation. Only one *A. pratensis* recording was

misclassified as *A. spinoletta*. Further 5 recordings of *A. spinoletta* and 5 of *A. pratensis*, included in the

same discriminant function as unknown specimens, were all correctly classified. The acoustic parameters

with higher DFA loadings, accounting for a better discriminant power, were RMM, PCFMa, PCFMi and BW

(Table 1). DUR90 has a very low loading and CF a near zero value.

I have calculated threshold values that can discriminate the *tsip* call of the two species (Table 1), as median

of range between mean + SD of lower value parameter and mean - SD of higher value parameter. I tested

the acoustic data using threshold values of three parameters (RMM, PCFMa and PCFMi) with the following

rule: i) at least two parameters must be in agreement; ii) one of the two must be the RMM. This procedure

was able to classify correctly 96% of all available recordings (n=75).

Similarity and differences among four *Anthus* species calls. Figure 3 shows the average contours of

alarm, *tsip* and *soft* calls for *A. spinoletta*, *A. pratensis*, *A. cervinus* and *A. petrosus*. An objective measure of

similarities among species for each different call type was calculated with two approaches. A multivariate

Mantel correlogram, computed with correlation index, was applied on the averaged time series of Figure 3.

This test showed a high similarity of *alarm* call of the four species (0.981) and lower values for *tsip* (0.660)

and *soft* calls (0.421).

Table 2 shows the results of a classical cross-correlation applied to the six possible comparisons between the

four species for each of the three different call type. Average value of *alarm* call is again by far higher

(0.943) than *tsip* and *soft* calls (0.441 and 0.329). Cross-correlation indexes, either at lag(0) or lag(m), were

not normally distributed and variance-covariance was significantly unequal. Therefore I performed the

comparisons among the three call types with Kruskall-Wallis test. Cross-correlation indexes, both at lag (0)

and at lag (m) of maximum correlation (Table 2), showed a significant difference among the three call types

(Hc=75.9 p<0.001 at lag 0 and Hc=78.3 p<0.001 at lag m). Mann Whitney post hoc test (with continuity

correction and a correction for ties) was significant for the comparisons *alarm* vs *tsip* call (p<0.001) and

alarm vs *soft* call (p<0.001), while the comparison *tsip* vs *soft* call was not significant. The average lag of

maximum correlation was 0.05 for *alarm* call, but shifted to 1.71 for *tsip* and to 3.31 for *soft* call. Again

Kruskall-Wallis test showed a significant difference among these lag data (Hc=54.5 p<0.001). Mann

Whitney post hoc test showed a significant shift of lag for *tsip* and *soft* versus *alarm* call (p<0.001), but also

a significantly higher shift for *soft* versus *tsip* call (p<0.01). An outlier data was produced by the

comparisons between *A. spinoletta* versus *A. petrosus* both for *tsip* and *soft* calls (Table 2), which yielded a

maximum correlation coefficient of 0.889 and 0.860 at lag 0. These two data, although significantly lower

than *alarm* call coefficient (p<0.05), were clearly by far higher than the other coefficients calculated for

these two call types. Moreover, for *A. spinoletta* versus *A. petrosus* comparison, the average lag of maximum

correlation (avg. lag m) did not shift significantly from zero also for *tsip* and *soft* calls.

DISCUSSION

This paper showed that a similar flight call (*tsip*) of two sister species, *A. pratensis* and *A. spinoletta*,

contains species-specific characteristics, that allows to discriminate the two species only with acoustic

means. Some results of my research are similar to those reported by Elfström (1992) for *A.pratensis* and *A.*

petrosus: for both studies, although the parameters are measured in different ways, *A. pratensis* has a lower

call duration, a reduced frequency range (BW of Table1) and a less frequency modulated call, represented by

RMM parameter in this paper and by pulse number in Elfström's work (1992). The top frequency figure of

A.pratensis (PFCMa Table 1) is quite consistent with that reported by Elfström (1992). Higher and lower

frequencies of *A. spinoletta* are close to those reported by Fijen (2014). Multivariate analysis of all acoustic

features showed a highly significant difference (Table 1) between the two species. The first question is: is it

possible to reduce the parameters that are effective and reliable to discriminate the two species? The low

loading value in the PCA analysis, the relatively low CVr (Table 1), the very high intraspecific variability

(CVin = 78.5, that means a fairly unreliable parameter) and several missing data, lead to discard time

interval between calls (TI). Frequency bandwidth (BW) proved to be less reliable because of its very low

CVr (1.05) and high variability (CVin = 29.7). Therefore the acoustic parameters were now reduced to five

(RMM, CF, PCFMa, PCFMi, DUR90).

The second question is: what is a suitable method to recognize acoustically the species and to classify new

unknown data? DFA is a commonly used method in bioacoustics to classify multivariate parameters

extracted from bird vocalizations (Galeotti and Sacchi 2001; Lengagne 2001; Christie et al. 2004; Terry *et al.*

2005) and it allows classifying new unknown specimens that are not part of the dataset used to create the

discriminant function, so that future observations can be correctly grouped. Moreover DFA helps to

understand what are the most important acoustic features to distinguish the two sister species. In this paper I

found that the discriminant function is able to correctly classify 98.4% of the recordings and group

assignment is validated by a leave-one-out cross-validation (jack-knifing) procedure. Only one bird out of 65

was misclassified. The DFA correctly classified all further 10 recordings of the two *Anthus* species that were

added to the discriminant function as unknown specimens. An analysis of the DFA loadings (Table 1) shows

that two parameters (central frequency CF and call duration DUR90) have very low values compared with

the other ones, therefore their discrimination power is relatively low. Since BW has a good DFA loading,

but, as stated above, it is less reliable than the other parameters, the most important acoustic features in the

discrimination process are the remaining three parameters: RMM representing the frequency modulations of

the tsip call, PFCMa that is the maximum frequency value of the peak frequency contour and PFCMi that is

the minimum frequency value of the peak frequency contour. Among these last three parameters RMM

shows the highest loading value for the DFA and can thus be considered the most critical acoustic feature for

the recognition of these two bird species. I also tested a simpler and lighter procedure to discriminate the *tsip*

calls of *A. pratensis* and *A. spinoletta*, which needs only the three above mentioned parameters. Table 1 at

the sixth row shows for each acoustic parameter the lower and higher values yielded by the mean minus the

SD and the mean plus SD: these ranges can reasonably used to find a threshold that separates the two

species. The threshold for the three most important parameters (RMM, PFCMa and PFCMi) was calculated

as the median of these ranges (Table 1). As shown above in the Results, these threshold values allowed a

correct classification of 96% of recordings, where only 2 *A. pratensis* and 1 *A. spinoletta* were misclassified

over 75 recordings. Of course either this procedure - that is relatively straigthtfoward and simpler than DFA -

or the DFA itself may be successfully used only if the methods of acoustic analysis exposed in this paper are

carefully followed.

This paper showed that two closely related species, with an overall similar vocalization, nevertheless

possess fine, but significant differences of acoustic features that may be important to transmit species-

specific information and to discriminate between conspecific and heterospecific vocalizations, like found by Elfström (1992) for *A. petrosus* and *A. pratensis*. This finding is confirmed by other authors for some species of the genus *Anthus* (Fijen 2014; Garner et al. 2015) and for bird species belonging to other taxa (Stiffler et al. 2018; Tietze et al. 2008). On the other hand, many bioacoustics researches found interspecific communication mediated by various bird call types (Johnson et al. 2003; Goodale and Kotagama 2005; Magrath et al. 2007; Templeton and Greene 2007; Magrath et al. 2009; Fallow et al. 2011; Fallow et al. 2013). The advantages of interspecific eavesdropping and communication by means of calls are documented in the literature (Nuechterlein 1981; Burger 1984; Forsmann and Mönkkönen 2001; Goodale and Kotagama 2005; Goodale et al. 2010) and this raises the question of how birds recognize heterospecific calls and their meaning. Learning has been suggested to explain avian recognition and response to heterospecific calls (Curio 1971; Hurd 1996; Magrath 2009), a second suggested hypothesis is that similar acoustic features facilitate heterospecific recognition (Marler 1957; Stefanski and Falls 1972; Johnson *et al.* 2003; Fallow *et al.* 2011) and a third hypothesis combines both previous ones: Hurd (1996) suggested that interspecific recognition results from both associative learning and common acoustic properties. Therefore it was useful to broaden my acoustic study to three different call types of four *Anthus* species, to study the heterospecific communication by means of some common call types in this genus.

The choice to analyse only the call contour (see Methods) was made for the following reasons: my aim was to represent objectively the overall acoustic features of the calls, underlining the trend of dominant frequency, which seems particularly involved in heterospecific recognition (Johnson et al. 2003); since most recordings were from Xeno-Canto mp3 files, with unknown quality level, I previously tested the reliability of contour representation (see Methods), that proved to be enough for the aims of the present work; finally, call representation by mean of average spectrogram contour is particularly suitable to be combined with cross-correlation as an objective method of evaluating vocal similarity. Combining cross-correlation with frequency contour is considered a method that may capture more precisely how signals vary over the duration of an acoustic unit (Odom *et al.* 2021).

The results of the comparison among *A. spinoletta*, *A. pratensis*, *A. cervinus* and *A. petrosus* on three different call types (*tsip*, *soft* and *alarm*) show that similarity of *alarm* calls in the four species is significantly and by far higher than the other two call types, with a tendency of a decrease in similarity from *tsip* to *soft* call. Indeed average correlation coefficient of the four species is significantly higher for *alarm* call either at lag 0 or at lag of the maximum correlation (Table 2). Since a shift from lag 0 proves a decreased similarity on the overall time shape of the calls, it is worth to analyse the average lag values where the correlation coefficient reach its maximum. For *alarm* it is 0.05, for *tsip* is 1.71 and for *soft* call is 3.31, where the differences of these values are highly significant, showing a decrease of similarity for the three call types among the four species. While the similarity among *alarm* calls of the four species is very strong and by far higher than the other two call types, the mean values of the correlation coefficients of *tsip* and *soft* calls don't differ significantly, although the lag 0 values show a higher similarity for *tsip* than for *soft* call. This slight difference is confirmed by Mantel correlogram index: 0.660 for *tsip* and 0.421 for *soft* call. I can conclude

that *alarm* calls of the four species show a very strong similarity, while the other two call types are more distinctive showing a much lower similarity, with *soft* call seeming to have a slightly lower similarity between species and hence a higher distinctive power. Since all four species are closely related from a taxonomic point of view, the results of my investigation support the hypothesis that *alarm* calls retain features of a common ancestor, while the other two call types (*tsip* and *soft*) show a decreasing similarity between the four *Anthus* species, resulting in a higher content of species-specific characteristics. The *alarm* call type studied in the present paper is only given in the vicinity of nest or in the breeding territory (Alström and Mild 2003; Bergmann et al. 2021; Dragonetti pers. obs.) therefore it is a very strong signal of danger, that the evolutive process may have maintained almost unaltered for these *Anthus* species, because it eases heterospecific communication. On the other hand *tsip* call is the most common call uttered in many different contexts mainly in flight and *soft* call is typically uttered between members of a flock as a contact signal (Elfström 1992; Alström and Mild 2003), therefore for these two calls the needs of intraspecific communication increase and evolution may have increased their acoustic divergence, as shown in the present paper.

Interestingly the comparison *A. petrosus* versus *A. spinoletta* yields a somewhat outlier data (see Table 2 row 5): indeed both *tsip* and *soft* calls show values (0.889 and 0.860 respectively) very close to that of *alarm* call (0.960) and a nearly null shift from lag 0, accounting for a greater similarity of this couple of species than the other ones. The taxonomy of "Water Pipit complex" has been much debated and until recently *A. petrosu*s and *A. spinoletta* were treated as a single species (Ali and Ripley 1998). Arctander et al. (1996) found only 1.2 % difference in mitochondrial cytochrome b sequences between *A. spinoletta* and *A. petrosus*, for this reason they expressed their reluctance to treat them as separate species. Today they are commonly treated as separate species, but there is no doubt that they are more closely related from a genetic point of view than the other species studied in this paper (see Voelker 1999) and this may explain the higher similarity indexes reported in Table 2.

A further consideration can be made about *alarm* call similarity in the genus *Anthus*, checking the eighteen species living in Europe, Asia and North America and visually comparing specimen spectrograms of *alarm* call available in the literature (Cramp and Simmons 1983; Alström and Mild 2003; Pieplow 2019; Bergmann et al. 2021). The *alarm* calls of six species were unknown (*A. nilghiriensis*, *A. silvanus*, *A. spragueii*, *A. rufulus*, *A. similis*, *A. campestris*) the remaining twelve might be classified as follows: three (*A. berthelotii*, *A. godlewskii*, *A. richardi*) show a completely different *alarm* call compared to those studied in this paper, seven (*A. trivialis*, *A. gustavi*, *A. roseatus*, *A. cervinus*, *A. petrosus*, *A. spinoletta*, *A. pratensis*) show a highly similar call and two (*A. hodgsoni*, *A. rubescens*) show a lower degree of similarity but are very reminiscent of the *alarm* described in this paper. If we look at the maximum likelihood tree of the genus *Anthus* based on the mitochondrial cytochrome b gene (from Voelker 1999), we can find all the former three species in the 4th group (the "large" pipits), while the remaining nine are all in the 3rd group (the "small" pipits).

The results of the present paper support the hypothesis that *alarm* call of four *Anthus* species is highly

similar, because it retains features of a common ancestor and may ease heterospecific communication. The

call similarities of closely related species can be important for phylogenetic comparative studies, as

suggested by the comparisons made in the *Anthus* genus. The method commonly used in phylogenetic

studies is integrative taxonomy (combination and integration of multiple types of evidence, see Sangster

2018): together with genetic and morphological information, acoustic data play a major role in taxonomic

designation of birds. The song of passerines, considered to be one of the most important traits promoting

differentiation among species (Price 2008), was extensively investigated in phylogenetic comparative

studies, but the data I reported here suggest that also the comparisons of call repertoires may be a useful tool

for taxonomic studies.

Conflict of Interest: the author declare that he has no conflict of interest.

ACKNOWLEDGEMENTS

I am extremely grateful to Martin Boesman, Fernand Deroussen, Ivan Farronato, Niels Krabbe, Lars

Lachmann, Dimitri Marrone, Nicolas Martinez, Jarek Matusiak, Marco Pesente, Maurizio Sighele, Bodo

Sonnenburg, Stanislav Wroza who provided original recordings (either in uncompressed or compressed

format) of *Anthus* species.Without their help this work could not be done. The Xeno-Canto collection was an

invaluable resource for my work. The credits for all recordings I have used from Xeno-Canto platform can

be accessed in the supplementary material. I am indebted to Gianni Pavan and Orlando Tommasini for their

comments and suggestions on a first draft of the manuscript. I also thank two anonymous reviewers for their

helpful comments on the manuscript.

REFERENCES

Ali S, Ripley S D (1998) Handbook of the birds of India and Pakistan. Volume 9 second edition Oxford

University Press, Delhi

Alström P, Mild K (2003) Pipits and Wagtails of Europe, Asia and North America. Christopher Helm,

London

Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance. Australian Ecology

26:32-46

Araya-Salas M, Smith-Vidaurre G, Webster M (2017) Assessing the effect of sound file compression and

background noise on measures of acoustic signal structure. Bioacoustics.

https://doi.org/10.1080/09524622.2017.1396498

Arctander P, Folmer O, Fjeldså J (1996) The phylogenetic relationships of Berthelot's Pipit *Anthus*

Berthelotii illustrated by DNA sequence data, with remarks on the genetic distance between Rock and Water

Pipits *A. spinoletta*. Ibis 138:263-272

Aubin T, Mathevon N, Da Silva ML, Vielliard JME, Sebe F (2004) How a simple and stereotyped acoustic

signal transmits individual information: the song of White-browed warbler Basileuterus leucoblepharus. An.

Acad Bras de Ciências 76:335-344

Baker MC, Howard TM, Sweet PW (2000) Microgeographic variation and sharing of the gargle vocalization

and its component syllable in black-capped chickadee (Aves, Paridae*, Poecile atricapillus*) populations.

Ethology 106:819-838

Bergmann H H, Helb HW, Baumann S (2008) Die stimmen der Vögel Europas. Aula Verlag GmbH,

Wiebelsheim

Bergmann H H, Engländer W, Baumann S (2021) Die stimmen der Vögel Europas. Aula Verlag GmbH,

Wiebelsheim

Brichetti P, Fracasso G (2007) Ornitologia Italiana. Volume 4 Oasi Alberto Perdisa Editore, Bologna

Briedis M, Beran V, Adamic P, Hanh S (2020) Integrating light□level geolocation with activity tracking

reveals unexpected nocturnal migration patterns of the tawny pipit. Journal of Avian Biology.

https://doi.org/10.1111/jav.02537

Burger J (1984) Grebes nesting in gull colonies: protective association and early warning. The American

Naturalist 123:327-337

Center for Conservation Bioacoustics (2019) Raven Pro: interactive sound analysis software. Version 1.6.1.

The Cornell Lab of Ornithology, Ithaca, NY

Christie PJ, Mennill DJ, Ratcliffe LM (2004) Chickadee song structure is individually distinctive over long

broadcasting distances. Behaviour 141:101-124

Cramp S, Simmons KEL (1983) The Birds of the Western Palearctic. Volume 3 Oxford University Press,

Oxford.

Curio E (1971) Die akustische Wirkung von Feindalarmen auf einige Singvogel. Journal für Ornithologie

112:365-372

De Kort SR, Ten Cate (2001) Response to interspecific vocalizations is affected by degree of phylogenetic

relatedness in Streptopelia doves. Animal Behaviour 61:239-247

Doornik J, Hansen H (2008) An Omnibus Test for Univariate and Multivariate Normality. Oxford Bulletin of

Economics and Statistics 70(s1):927-939

Ebenhöh H, Hoffrichter O (1998) Beobachtungen an einem winterlichen Schlafplatz des Bergpiepers1

(*Anthus spinoletta*) bei Köndringen, Landkreis Emmendingen. Naturschutz südl. Oberrhein 2:181-194.

Elfström ST (1992) Tsip calls of meadow and rock pipits: interspecific comparisons and playback

experiments. Animal Behaviour 43:795-802

Elkins N (2005) Weather and bird migration. British Birds 98:238-256

Farnsworth A (2005) Flight calls and their value for future ornithological studies and conservation research.

The Auk 122(3):733-746

Fallow P, Gardner JL, Magrath RD (2011) Sound familiar? Acoustic similarity provokes responses to

unfamiliar heterospecific alarm calls. Behavioral Ecology 22:401-410

Fallow P, Pitcher BJ, Magrath RD (2013) Alarming features: birds use specific acoustic properties to identify

heterospecific alarm calls. Proceedings of The Royal Society B. https://doi.org/10.1098/rspb.2012.2539

Fijen TPM (2014) Flight call identification of Rock Pipit and Water Pipit. Dutch Birding 36:87-95

Forsman JT, Mönkkönen M (2001) Responses by breeding birds to heterospecific song and mobbing call

playbacks under varying predation risk. Animal Behaviour 62:1067-1073

Gal A, (2003) A New Technique for Studying Nocturnal Bird Migration. Semantic Scholar.

https://www.semanticscholar.org/paper/A-New-Technique-for-Studying-Nocturnal-Bird-

Gal/eb621c8c3cbba823ca9b9133cb4f1e75d3bd621c

Galeotti P, Sacchi R (2001) Turnover of territorial Scops Owls Otus scops as estimated by spectrographic

analyses of male hoots. Journal of Avian Biology 32:256-262

Garner M, Perlman Y, Kiat Y, Collison MJ (2015.) Water Pipits: three species rather than one? British Birds

108:42-48

Gayk ZG, Simpson RK, Mennill DJ (2021) The evolution of wood warbler flight calls: Species with similar

migrations produce acoustically similar calls. Evolution 75(3):719-730

Goodale E, Kotagama S (2005) Alarm calling in Sri Lankan mixed-species bird flocks. The Auk 122(1):108-

120

Goodale E, Beauchamp G, Magrath RD, Nieh JC, Ruxton GD (2010) Interspecific information transfer

influences animal community structure. Trends in Ecology and Evolution 25:354-361

Hamilton WJ III (1962) Evidence concerning the function of nocturnal call notes of migratory birds. Condor

64:390-401

Hammer Ø, Harper DAT, Ryan PD (2001) PAST: Paleontological Statistics Software Package for Education
and Data Analysis. Palaeontologia Electronica 4(1):1-9

Hurd CR (1996) Interspecific attraction to the mobbing calls of black capped chickadees (*Parus*
atricapillus). Behavioural Ecology and Sociobiology 38:287-292

Johnson FR, McNaughton EJ, Shelley CD, Blumstein DT (2003) Mechanism of heterospecific recognition
in avian mobbing calls. Australian Journal of Zoology 51:577-585

Kroodsma DE (2000) Vocal behavior. In Home Study Course in Bird Biology. Cornell Laboratory of
Ornithology (ed), Ithaca, New York, Chapter 7 pp 1-18

Legendre P, Legendre L (1998) Numerical Ecology. 2nd English ed. Elsevier, Amsterdam

Lengagne T (2001) Temporal stability in the individual features in the calls of Eagle Owl (*Bubo bubo*).
Behaviour 138:1407-1419

Magrath RD, Pitcher BJ, Gardner JL (2007) A mutual understanding? Interspecific

responses by birds to each other's aerial alarm calls. Behavioral Ecology 18(5):944-951

Magrath RD, Pitcher BJ, Gardner JL (2009) Recognition of other species' aerial alarm calls: speaking the
same language or learning another? Proceedings of The Royal Society B 276:769-774

Marler P (1955) Characteristics of some animal calls. Nature 176:6-8

Marler P (1957) Specific distinctiveness in the communication signals of birds. Behaviour 11:13-38

Marler P, Slabbekoorn H (2004) Nature's music. The science of birdsong. Elsevier Academic Press, San
Diego

Mudinger PC (1970) Vocal imitation and individual recognition of finch calls. Science 168:480-482

Mudinger PC (1979) Call learning in the Carduelinae: Ethological and systematic considerations. Systematic
Zoology 28:270-283

Nowicki S (1989) Vocal plasticity in captive black-capped chickadees: the acoustic basis and rate of call
convergence. Animal Behaviour 37:64-73.

Nuechterlein GL (1981) 'Information parasitism' in mixed colonies of western grebes and Forster's terns.
Animal Behaviour 29:985-989

Odom KJ, Araya-Salas M, Morano JL et al (2021) Comparative bioacoustics: a roadmap for quantifying and
comparing animal sounds across diverse taxa. Biological Reviews 96(4):1135-1159

Pieplow N (2019) Peterson Field Guide to Bird Sound of Western North America. Houghton Mifflin

Harcourt, New York

Price T (2008) Speciation in Birds. Roberts and Company, Greenwood Village, CO

R Core Team (2020) R: A language and environment for statistical computing. R Foundation for Statistical

Computing

Rencher AC (2002) Methods of multivariate analysis. 2nd ed. Wiley, New York

Sangster G (2018) Integrative taxonomy of birds: The nature and delimitation of species. In: Tietze DT (ed)

Bird Species, Springer International Publishing, Cham, pp 9-37

Scherrer B (1984) Biostatistique. Gaetan Morin Press, Quebec

Sever M, Lajovic J, Rajer B (2005) Robustness of the Fisher's Discriminant Function to Skew-Curved

Normal Distribution. Metodoloski Zvezki 2(2):231-242

Stefanski RA, Falls JB (1972) A study of distress calls of song, swamp, and white-throated sparrows (Aves:

Fringillidae). II. Interspecific responses and properties used in recognition. Canadian Journal of Zoology

50:1513-1525.

Stiffler LL, Schroeder KM, Anderson JT, McRae SB, Katzner TE (2018) Quantitative acoustic

differentiation of cryptic species illustrated with King and Clapper rails. Ecology and Evolution 8:12821–

12831.

Templeton CN, Greene E (2007) Nuthatches eavesdrop on variations in heterospecific chickadee mobbing

calls. Proceedings of the National Academy of Sciences USA 104:5479-5482

Terry AM, Peake TM, McGregor PK (2005) The role of vocal individuality in conservation. Frontiers in

Zoology https://doi.org/10.1186/1742-9994-2-10

Tietze DT, Martensa J, Sun Y, Packert M (2008) Evolutionary history of treecreeper vocalisations (Aves:

Certhia). Organism, Diversity and Evolution 8:305-324

Voelker G (1999) Molecular evolutionary relationships in the Avian genus *Anthus* (pipits: Motacillidae).

Molecular Phylogenetics and Evolution 11:84-94

Wang Y, Shi Y et al (2019) Patterns of bird nocturnal migration at Shenyang Taoxian International Airport,

Northeast China. The Journal of applied ecology 30(1):292-300

Williams H (2008) Birdsongs and singing behavior. In: Neuroscience of Birdsongs, Zeigler HP, Marler P

(eds), Cambridge University Press, Cambridge, pp 5-32

Zann R (1985) Ontogeny of the zebra finch distance call. I. Effects of cross fostering to bengalese finches. Z.

Tierpsychol 68:1-23

512   **TABLES AND FIGURES**

	RMM (n)	**CF (Hz)**	**PFCMa (Hz)**	**PFCMi (Hz)**	**BW (Hz)**	**DUR90 (ms)**	**TI (ms)**
A. pratensis	5.01 +/- 0.62	6643 +/- 280	7271 +/- 273	5928 +/- 448	1343 +/- 461	56 +/- 6	293 +/- 305
A. spinoletta	7.24 +/- 0.91	6078 +/- 370	6655 +/- 408	5037 +/- 386	1618 +/- 396	72 +/- 8	1255 +/- 638
CVis	22.08	6.82	6.23	11.14	30.62	16.36	92.58
CVr	1.75	1.31	1.25	1.45	1.03	1.53	1.18
DFA loadings	76.07	0.20	47.91	47.55	47.68	7.88	n.c.
T/range	5.63 – 6.33	n.c.	6998 – 7063	5423 – 5480	1222 – 1804	62 - 64	n.c.
Threshold	5.98	n.c.	7030	5451	1513	63	n.c.

Table 1. Comparison of *tsip* call vocal parameters between *A. pratensis* and *A. spinoletta*. **CV** = coefficient of variation (see Methods); **CVis** = interspecific coefficient of variation; **CVr** = ratio between CVis and CV intraspecific of the two species. **DFA** = discriminant function analysis. **T/range** = range between mean + SD of lower value parameter and mean - SD of higher value parameter. **Threshold** = median of T/range values taken as separation threshold between the two species. **RMM** = number of relative minimum and maximum values of call frequency contour; **CF** = center frequency; **PFCMa** = maximum of the peak frequency contour measured from spectrogram slices; **PFCMi** = minimum of the peak frequency contour; **BW** = call frequency bandwidth calculated from (PFCMa - PFCMi); **DUR90** = call duration containing 90% of the call energy; **TI** = time interval between start of subsequent calls; n.c. = not calculated. *A. pratensis* n=34, *A. spinoletta* n=31

	n (a)	alarm(0)	alarm(m)	avg. lag (m)	n (t)	tsip(0)	tsip(m)	avg. lag (m)	n (s)	soft(0)	soft(m)	avg. lag (m)
pr vs sp	10	0.869+/-0.151	0.905+/-0.085	0.29+/-0.49	13	0.433+/-0.173	0.502+/-0.128	1.14+/-0.90	12	0.463+/-0.132	0.650+/-0.089	1.86+/-0.38
pr vs ce	10	0.962+/-0.021	0.962+/-0.021	0.00	13	0.313+/-0.196	0.447+/-0.070	2.71+/-3.15	12	-0.06+/-0.394	0.711+/-0.140	3.71+/-1.38
pr vs pe	10	0.935+/-0.041	0.935+/-0.041	0.00	13	0.367+/-0.234	0.455+/-0.145	1.29+/-1.25	12	0.388+/-0.118	0.582+/-0.106	2.00+/-0.82
sp vs ce	10	0.963+/-0.045	0.963+/-0.045	0.00	13	0.318+/-0.194	0.462+/-0.085	2.43+/-1.40	12	0.175+/-0.450	0.481+/-0.182	4.86+/-3.76
sp vs pe	10	0.960+/-0.034	0.960+/-0.034	0.00	13	0.889+/-0.034	0.889+/-0.034	0.00	12	0.860+/-0.077	0.871+/-0.068	0.43+/-0.053
ce vs pe	10	0.968+/-0.027	0.968+/-0.027	0.00	13	0.325+/-0.131	0.383+/-0.061	2.71+/-4.03	12	0.146+/-0.371	0.437+/-0.212	7.00+/-3.37
Mean	60	0.943+/-0.074	0.949+/-0.049	0.05+/-0.22	78	0.441+/-0.263	0.523+/-0.192	1.71+/-2.34	72	0.329+/-0.407	0.622+/-0.198	3.31+/-3.00

Table 2. Cross correlation results (average +/- SD) among contours of three different call type (*alarm* call, *tsip* call, *soft* call) in four *Anthus* species; **pr** = *A.*
pratensis; **sp** = *A. spinoletta*; **ce** = *A. cervinus*; **pe** = *A. petrosus*. **Mean** = average of all the cross-correlations of all comparisons. **(0)** = cross-correlation results at lag
0; **(m)** = cross-correlation results at lag of maximum correlation coefficient; **avg. lag(m)** = average lag value corresponding to the maximum correlation coefficient;
n(a) = number of cross-correlations performed for *alarm* calls; **n(t)** = number of cross-correlations for *tsip* calls; **n(s)** = number of cross-correlations for *soft* calls.
Each contour submitted to cross-correlation is an average of minimum 3 and maximum 9 randomly selected calls from each recording of a given species. The total
number of randomly selected calls is always 84 for each recording set and for every call type of the four species.
520

Figure 1. Spectrograms of three different call types of *A. spinoletta* (Asp), *A. pratensis* (Apr), *A. cervinus* (Ace), *A. petrosus* (Ape). See Methods for call type classification

527　**Figure 2 .** Comparison of *tsip* call vocal parameters between *A. pratensis* (cubes) and *A. spinoletta* (crosses).
528　Scatter plot in the coordinate system given by the first three PCA components that account for the 89.4 % of
529　the variance.
530

Figure 3 . Average frequency contours (FCP75, see Methods) of three different calls in four *Anthus* species.
Alarm call: average of 10 recordings for each species. **Tsip call**: average of 13 recordings for each species.
Soft call : average of 12 recordings for each species. The total number of randomly selected calls for every
call type of each species is always 84. Time series (x axis) are made by points evenly spaced 5 ms apart