Title: The *trans*-zeatin-type side-chain modification of cytokinins controls rice growth

Takatoshi Kiba[1,2], Kahori Mizutani[1], Aimi Nakahara[1], Yumiko Takebayashi[2], Mikiko Kojima[2],

Tokunori Hobo[3], Yuriko Osakabe[4,5], Keishi Osakabe[5], and Hitoshi Sakakibara[1,2]

1. Graduate School of Bioagricultural Sciences, Nagoya University, Furocho, Chikusa-ku,

Nagoya, 464-8601, Japan

2. RIKEN Center for Sustainable Resource Science, 1-7-22, Suehiro, Tsurumi, Yokohama

230-0045, Japan

3. Biosci. Biotech Center, Nagoya University., Furocho, Chikusa-ku, Nagoya, 464-8601,

Japan

4. Department of Life Science and Technology, Tokyo Institute of Technology, Yokohama,

226-8501, Japan

5. Graduate School of Technology, Industrial and Social Sciences, Tokushima University,

Tokushima, 770-8503, Japan,

Corresponding Author: Takatoshi Kiba: e-mail: kiba@agr.nagoya-u.ac.jp, TEL:

81-52-789-5511; FAX: 81-52-789-4095

Running head: Roles of *trans*-zeatin-type cytokinins in rice

Abstract

Cytokinins (CKs), a class of phytohormones with vital roles in growth and

development, occur naturally with various side-chain structures, including

N^6-(Δ^2-isopentenyl)adenine-, *cis*-zeatin- and *trans*-zeatin (tZ)-types. Recent studies in a model

dicot plant *Arabidopsis* demonstrated that tZ-type CKs are biosynthesized via cytochrome

P450 monooxygenase (P450) CYP735A, and have a specific function in shoot growth

promotion. Although the function of some of these CKs has been demonstrated in a few

dicotyledonous plant species, the significance of these variations and their biosynthetic

mechanism and function in monocots and in plants with distinctive side-chain profiles than

Arabidopsis, such as *Oryza sativa* (rice), remain elusive. In this study, we characterized

CYP735A3 and *CYP735A4* to investigate the role of tZ-type CKs in rice. Complementation

test of the *Arabidopsis* CYP735A-deficient mutant and CK profiling of loss-of-function rice

mutant, *cyp735a3 cyp735a4*, demonstrated that *CYP735A3* and *CYP735A4* encode P450s

required for tZ-type side-chain modification in rice. *CYP735As* are expressed in both roots

and shoots. The *cyp735a3 cyp735a4* mutants exhibited growth retardation concomitant with

reduction in CK activity in both roots and shoots, indicating that tZ-type CKs function in

growth promotion of both organs. Expression analysis revealed that tZ-type CK biosynthesis

is negatively regulated by auxin, abscisic acid, and cytokinin and positively by dual nitrogen

nutrient signals, namely glutamine-related and nitrate-specific signals. These results suggest

that the physiological role of tZ-type CKs in rice is different from that in *Arabidopsis* and

they control growth of both roots and shoots in response to internal and environmental cues in

rice.

Introduction

Cytokinins (CKs), a group of plant hormones, are involved in the regulation of various aspects of plant growth and development, including cell division and differentiation, organogenesis, root and shoot growth, and environmental responses (Mok and Mok, 2001; Sakakibara, 2006; Werner and Schmülling, 2009; Kiba et al., 2011; Kieber and Schaller, 2018; Kiba et al., 2019; Sakakibara, 2021). Natural CKs are mostly derivatives of adenine with a prenyl side chain at the N^6 position. There are variations in the side-chain structure of CKs and the most common structures in plants are N^6-(Δ^2-isopentenyl)adenine (iP)-, *trans*-zeatin (tZ)-, and *cis*-zeatin (cZ)-types, which differ in the presence and stereoisomeric position of the terminal hydroxyl group on the prenyl side chain (Shaw, 1994; Mok and Mok, 2001; Gajdošová et al., 2011).

Our understanding of biosynthetic pathways of the side-chain variants has greatly progressed, thanks mostly to studies in *Arabidopsis*. The initial step of *de novo* iP- and tZ-type CK biosynthesis is catalyzed by adenosine phosphate-isopentenyltransferase (IPT) to produce iP-type CK nucleotide precursors, iP ribotides (iPRPs) (Kakimoto, 2001; Takei et al., 2001a). tZ-type side chain is formed by the enzyme cytochrome P450 monooxygenase CYP735A, which is encoded by *CYP735A1* and *CYP735A2* in *Arabidopsis* (Takei et al., 2004; Kiba et al., 2013). CYP735A *trans*-hydroxylates the prenyl side chain of iPRPs to produce tZ ribotides (tZRPs). Finally, the cytokinin-activating enzyme LONELY GUY (LOG) converts the nucleotide precursors to their active nucleobase forms, iP and tZ (Kurakawa et al., 2007; Kuroha et al., 2009; Tokunaga et al., 2012). The cZ-type CKs are biosynthesized via the prenylation of tRNA by tRNA-isopentenyltransferase (tRNA-IPT) and the degradation of the prenylated tRNA (Miyawaki et al., 2006; Gajdošová et al., 2011).

Genes responsible for prenylated tRNA degradation for cZ-type CK production have not been

identified yet.

CK biosynthesis is regulated by various internal and environmental factors (Hirose

et al., 2008; Zwack and Rashotte, 2015). Nitrogen nutrition, which is available mostly as

nitrate and ammonium in soil (Miller and Cramer, 2004; Kirk and Kronzucker, 2005; Xu et al.,

2012; Kiba and Krapp, 2016; Coskun et al., 2017), is one of the major factors. Studies in

Arabidopsis showed that *IPT3* and *CYP735As* are induced by a nitrate-specific signal to boost

iP- and tZ-type CK accumulation in *Arabidopsis* (Miyawaki et al., 2004; Takei et al., 2004;

Engelsberger and Schulze, 2012; Maeda et al., 2018). *OsIPT4* and *OsIPT5* are not induced by

nitrate itself but by ammonium and glutamine with accompanying accumulation of iP-type

and tZ-type CKs, indicating that *de novo* iP- and tZ-type CK biosynthesis is regulated by

glutamine-related signals in rice (Kamada-Nobusada et al., 2013). A *CYP735A* gene in *Oryza*

longistaminata, a wild rice, was shown to be upregulated by ammonium nitrate but the signal

responsible for the upregulation is not known (Shibasaki et al., 2021). Thus, plants have dual

systems, nitrate-specific and glutamine-related systems, to modulate the iP-type and tZ-type

CK biosynthesis in response to nitrogen availability. Although the underlying mechanism is

not understood, cZ-type CK levels are generally increased in response to biotic and abiotic

stresses, such as phosphate starvation, low temperature, drought, and salt treatment

(Vyroubalová et al., 2009; Macková et al., 2013; Vanková et al., 2014; Silva-Navas et al.,

2019).

Previous studies have suggested that there is biological relevance to side-chain

variations of CKs (Schmitz and Skoog, 1972; Mok et al., 1978; Werner et al., 2003;

Spíchal et al., 2004; Gajdošová et al., 2011; Stolz et al., 2011; Choi et al., 2012;

Osugi and Sakakibara, 2015; Schäfer et al., 2015). However, it is only recently that

studies in *Arabidopsis* demonstrated it. The abundance of iP- and tZ-type CKs is higher than

that of cZ-type (Osugi and Sakakibara, 2015; Osugi et al., 2017; Kiba et al., 2019)

and tZ-type CKs are the most active, while cZ-type CKs are the least active in bioassays in

Arabidopsis (Gajdošová et al., 2011; Kudo et al., 2012). *Arabidopsis* CK receptors have

different affinities toward side-chain variants (Spíchal et al., 2004; Romanov et al., 2006;

Stolz et al., 2011). For example, AHK2 and AHK4/WOL/CRE1 have high affinity to both

iP and tZ, while AHK3 binds to iP with a ten times lower affinity than tZ in *in vitro* binding

assays. Consistent with the bioassay, the affinity of AHKs to cZ is significantly lower

compared with that of iP and tZ (Spíchal et al., 2004; Romanov et al., 2006; Stolz et al.,

2011). Furthermore, characterization of the *Arabidopsis cyp735a1 cyp735a2* mutant that is

impaired in tZ-type side-chain modification has unequivocally shown that the function of

tZ-type CKs is shoot growth promotion, and it is different from iP-type and cZ-type CKs

(Kiba et al., 2013). Thus, tZ-type CKs are the major active variants possessing a specific

function in shoot growth, while cZ-type CKs are the weakly active or inactive forms with

minor roles in *Arabidopsis*.

Although the occurrence of the three side-chain variants is ubiquitous among plants

(Gajdošová et al., 2011), several lines of evidence suggest that their functional

differentiation may not be as universal. For instance, the tZ-type CKs is not consistently more

abundant than the cZ-type CKs among all plants. Plants that contain higher levels of cZ-type

CKs than tZ-type can be found across the whole evolutionary tree of land plants (Gajdošová

et al., 2011). Interestingly, many crops, including pea, potato, oats, maize and rice, are

cZ-type-abundant plants (Gajdošová et al., 2011; Schäfer et al., 2015). Furthermore,

cZ-type CKs are shown to be as active as tZ-type in some bioassays, including

cytokinin-responsive gene expression assay in maize cultured cell (Yonekura-Sakakibara

et al., 2004) and seminal root growth inhibition assay in rice (Kudo et al., 2012).

Consistently, cytokinin receptors of maize (ZmHK1) and rice (OsHK3 and OsHK4) were

shown to have similar affinity toward cZ and tZ (Lomin et al., 2011; Choi et al., 2012).

However, the exact function of each of these variants in plants other than *Arabidopsis*,

including monocot and cZ-type-abundant plants, is still unknown and remains to be

determined.

Rice is a monocot plant possessing a significantly distinctive CK side-chain profile

from *Arabidopsis*; cZ-type CKs comprise more than 80% of total CK in various tissues

(Kojima et al., 2009; Kudo et al., 2012; Osugi and Sakakibara, 2015). In this study,

we characterized the consequence of disruption of rice *CYP735A*s (*CYP735A3* and

CYP735A4) and demonstrated that these genes are vital for tZ-type CK biosynthesis. Our

results also revealed that tZ-type CK biosynthesis is regulated by phytohormones and dual

nitrogen signals and that tZ-type CKs act to promote root and shoot growth in rice. These

results suggest that the physiological role of tZ-type CKs in rice is different from that in

Arabidopsis despite the fact that the biosynthetic mechanism is conserved. The function of

tZ-type CKs might be optimized for each plant species depending on the body plan and/or

growth environments.

Results

***CYP735A3* and *CYP735A4* complement the Arabidopsis *cyp735a1 cyp735a2* mutant**

In *Oryza sativa* genome, *CYP735A3* and *CYP735A4* encode proteins with high

amino acid sequence homology to CYP735A1 and CYP735A2, which are cytochrome P450

monooxygenases for tZ-type CK biosynthesis in *Arabidopsis* (Fig. S1) (Hansen et al.,

2021). The *cyp735a1 cyp735a2* mutant (*atcypDM)* is deficient in tZ-type CKs resulting in

shoot growth retardation (Kiba et al. 2013). To assess whether these genes encode enzymes

that catalyze the same reaction as CYP735A1 and CYP735A2, we expressed them in

atcypDM under the control of cauliflower mosaic virus (CaMV) 35S promoter. Accumulation

of *CYP735A3* or *CYP735A4* transcripts was confirmed by quantitative RT-PCR analysis

(qRT-PCR) in two independent lines of transgenic plants CYP735A3-ox or CYP735A4-ox,

respectively (Fig. 1A). When grown on soil, all the transgenic lines developed a rosette with

significantly greater diameter than *atcypDM* (Fig. 1B). Both *CYP735A3* and *CYP735A4*

expression resulted in an increase of tZ-type CK (tZ and its conjugates) concentration with a

concomitant decrease of iP-type CK (iP and its conjugates) concentration compared with

atcypDM, but no significant differences in cZ-type CK (cZ and its conjugates) concentration

were observed (Fig. 1C, Table S1). These results demonstrate that CYP735A3 and

CYP735A4 catalyze the same reaction as CYP735A1 and CYP735A2, which is the

*tran*s-hydroxylation of CK side chain to synthesize tZ-type CKs from iP-type CKs.

The spatial expression pattern of *CYP735A3* and *CYP735A4*

Expression levels of *CYP735A3* and *CYP735A4* were analyzed by qRT-PCR in

various organs of rice in seedling to early reproductive phases (Fig. 2). At seedling phase,

transcripts of *CYP735A*s were detected both in roots and shoots but they were more abundant

in roots as reported previously (Tsai et al., 2012) (Fig. 2A). In later phases tested, the

accumulation of the *CYP735A3* transcript was the highest around the vegetative shoot apex,

while that of *CYP735A4* transcript was prominent in the leaf blade of plants at vegetative and

early reproductive phases (Fig. 2B).

In order to localize *CYP735A*s expression at the tissue level, we used GUS staining

technique. For this purpose, we generated transgenic plants harboring a fusion of the 3.6 kb

CYP735A upstream sequence and the *GUS* reporter gene (proCYP735A3:GUS and

proCYP735A4:GUS). Similar staining patterns were observed in multiple independent T1

lines of proCYP735A3:GUS (Figs. 2C-E). However, no staining was detected in any of the

proCYP735A4:GUS lines (data not shown), suggesting that the 3.6 kb upstream sequence of

CYP735A4 was not sufficient to drive expression. Consistent with the qRT-PCR results (Fig.

2A), staining was detected both in roots and shoots of proCYP735A3:GUS seedlings (Fig. 2C).

The GUS activity was restricted to the vascular bundle (Fig. 2D) and observation of

cross-sections revealed that the staining was pronounced in vascular parenchyma cells in the

shoot, and vascular parenchyma and pericycle cells in the root (Fig. 2E).

***CYP735A3* and *CYP735A4* play a central role in tZ-type cytokinin biosynthesis in rice**

To examine the role of *CYP735A*s in CK side-chain modification in rice, we

disrupted *CYP735A3* and *CYP735A4* by CRISPR-Cas9 system. Two transfer

RNA-based-multiplex CRISPR-Cas9 vectors, each harboring three different guides targeting

unique sequences of *CYP735A3* and *CYP735A4,* were used to edit *CYP735A3* and *CYP735A4*

(Figs. S2 and S3). Two independent lines with mutations in both *CYP735A3* and *CYP735A4*

were identified. In a T1 line, the *cyp735a3-1* and *cyp735a4-1* alleles were found as

homozygote and heterozygote, respectively (Fig. S2). In another T1 line, the *cyp735a3-2* and

cyp735a4-2 alleles were found as heterozygote and homozygote, respectively (Fig. S3). The

cyp735a3-1 single, *cyp735a4-2* single, *cyp735a3-1 cyp735a4-1* (*a3a4-1*) double and

cyp735a3-2 cyp735a4-2 (*a3a4-2*) double mutants were isolated from progenies of these lines.

The *cyp735a3-1* and *cyp735a3-2* are one-base insertions that shift the reading frame and yield

predicted proteins with no similarity to any Non-redundant UniProtKB/SwissProt sequences

and without the cytochrome P450 domain (Fig. S4). The *cyp735a4-1* and *cyp735a4-2* mutants

are 263- and 298-base deletions, respectively. The 263- and 298-base deletions resulted in a

predicted protein lacking "A helix", a secondary structure conserved among cytochrome P450s

(Midlik et al., 2021), and an in-frame stop codon at 4 codons after the mutation, respectively

(Fig. S5). Since the proteins predicted to be produced in all the mutants (except for

cyp735a4-1) are apparently inactive as cytochrome P450, we concluded that *cyp735a3-1*,

cyp735a3-2, and *cyp735a4-2* are null alleles.

CK concentrations in roots and shoots were measured in wild type (WT) strain

Nipponbare (NB), vector control (VC), and mutant strains *cyp735a3-1*, *cyp735a4-2*, *a3a4-1*

and *a3a4-2* seedlings grown in hydroponic culture (Fig. 3, Tables S2 and S3). The

concentrations of tZ-type CK and iP-type CK were comparable to WT in *cyp735a3-1* (Fig. 3A).

In contrast, *cyp735a4-2* had slightly decreased tZ-type CK and slightly increased iP-type CK

concentrations in roots and shoots as compared to the WT (Fig. 3B). In *a3a4-1* and *a3a4-2,* the

concentration of tZ-type CKs reduced dramatically to 3.5 to 8.8% of the wild-type levels in

roots and shoots, while that of iP-type CK increased more than 2 -fold. In the root of double

mutants, iP-type CK levels were increased so dramatically that the loss of tZ-type CK was

compensated in terms of the sum of iP- and tZ-type CKs (iP-/tZ-type CK quantity). On the

other hand, in the shoot, iP-type CK levels were increased but the iP-/tZ-type CK quantity

was significantly lower compared with WT (Fig. 3; Tables S2 and S3). The cZ-type CK levels

were not altered in any of the mutants (Fig. 3, Tables S2 and S3). Since the CK profiles of

a3a4-1 and *a3a4-2* are distorted in a similar manner, *cyp735a4-1* seems to be a null allele.

These results show that tZ-type CKs are mostly synthesized by CYP735A3 and CYP735A4 in

rice.

Disruption of *CYP735A3* and *CYP735A4* retards growth of rice

To explore the physiological role of tZ-type CKs in rice, we examined the growth

and development of *cyp735a* single and double mutants from seedling phase to maturation

phases. Although *cyp735a3-1* and *cyp735a4-2* single mutants were indistinguishable from

control plants (WT and VC) on our growth conditions, similar significant alterations in growth

and development were observed in *a3a4-1* and *a3a4-2* double mutants (Figs. 4, 5 and S6).

The double mutant seedlings showed a decrease in the length and fresh weight of

the shoot and root compared with control (Figs. 4A, 4B, S6A, S6B). To determine the cause

of the decrease in shoot length and fresh weight, leaf number and leaf size were measured

(Figs. 4C-E). Leaf number of double mutant seedlings at 15 days after germination was

reduced compared to control (Fig. 4C), indicating that the rate of leaf formation is slower in

the double mutant. Leaf blade and sheath length of fully expanded third leaf was similar for

the double mutant and control (Fig. 4D), while leaf blade width was decreased in the double

mutant (Fig. 4E). These results indicated that a decrease in leaf number accounts for the

reduction in shoot length and a decrease in leaf number and leaf width contributes to the

reduction in shoot fresh weight. Reduced shoot length phenotype of the double mutant

continued into later vegetative growth stage (Figs. S6C and S6D), but it cannot be attributed

only to a decrease in leaf number because leaf size, namely blade length, blade width and

sheath length, was significantly reduced at this stage as observed in fully expanded ninth leaf

(Fig. 4F). In addition, the double mutant developed significantly fewer tillers compared with

control (Figs. 4G and 5C), suggesting that the activities of shoot apical and axillary meristems

are impaired.

235   All control plants headed about 83 days after germination (n=19) but double mutant

plants either died without heading from the main stem (16 out of 24 plants) or showed

delayed heading (8 out of 24 plants) (Figs. 5A and 5B). Panicle length, panicle weight and

number of primary panicle branch were all significantly reduced in the double mutant

compared to the control (Figs. 5D-G), suggesting that inflorescence meristem activity is

reduced. The double mutant set few but viable seeds, indicating that reproductive organ

development is not severely affected. Together, these results suggest that the tZ-type CKs

regulate both root and shoot growth, most possibly through controlling meristem activity.

The cytokinin activity is reduced both in roots and shoots of the *cyp735a3 cyp735a4*

mutant

246   The growth defects observed in *a3a4* double mutant are reminiscent of those

observed in rice mutants with diminished CK activities such as type-A *RESPONSE*

REGULATOR (*RR*) *OsRR6* overexpressing lines (Hirose et al., 2007), type-B *RR* triple

mutant *rr21/22/23* (Worthen et al., 2019) and cytokinin receptor double mutant *hk5 hk6*

(Burr et al., 2021). To examine whether the CK activity is reduced in the double mutant, we

analyzed the expression levels of immediate-early CK-inducible type-A *RR*s by qRT-PCR

(Fig. 6). The expression level of all tested type-A *RR*s, except for that of *OsRR1* in roots, was

significantly lower in the mutants compared with WT in both roots and shoots (Figs. 6A and

6B). On the other hand, the expression of non-CK-inducible type-B *RR*s, *OsRR21* and

OsRR23, was not reduced (Figs. 6C and 6D). Thus, cytokinin activity in roots and shoots of

a3a4 is diminished, suggesting that the *a3a4* growth phenotype is caused by decreased CK

activity.

Regulation of *CYP735A3* and *CYP735A4* expression by phytohormones

To obtain insights into the physiological relevance of tZ-type CK biosynthesis in rice,

we analyzed the responses of *CYP735A*s to phytohormones. In *Arabidopsis*, the expression of

CYP735A1 and *CYP735A2* was shown to be upregulated by CK and downregulated by auxin

and abscisic acid (Takei et al., 2004). Tsai et al (2012) analyzed the effect of these

phytohormones on *CYP735A3* and *CYP735A4* expression by the NanoString nCounter system

but could not detect *CYP735A3* under some conditions using this system. To fully characterize

the response of *CYP735A*s to these phytohormones in rice, we analyzed their expression in

roots by qRT-PCR. Nipponbare seedings treated with phytohormones were used for the

analysis. *OsRR4*/*OsRR6* (Kudo et al., 2012; Tsai et al., 2012), *OsSIPP2C1* (Li et al.,

2013) and *OsGH3.2* (Du et al., 2012) were used as positive controls for CK, IAA and ABA

responses, respectively. Consistent with the NanoString analysis (Tsai et al., 2012), we found

that the *CYP735A4* transcript was decreased by CK (iP and tZ), IAA and ABA (Fig. 7). Our

qRT-PCR analysis showed that *CYP735A3* is also downregulated by CK, IAA and ABA (Fig.

7). Thus, rice *CYP735A*s and *Arabidopsis CYP735A*s are regulated similarly by IAA and ABA,

but differently by CK.

Regulation of *CYP735A3* and *CYP735A4* expression by nitrogen sources

Nitrogen nutrition is one of the major environmental factors regulating CK

biosynthesis, including tZ-type CK biosynthesis, for plant growth optimization (Takei et al.,

2001b; Kamada-Nobusada et al., 2013; Ohashi et al., 2017; Osugi et al., 2017;

Maeda et al., 2018; Sakakibara, 2021; Shibasaki et al., 2021; Kawai et al., 2022).

In *Arabidopsis*, *CYP735A*s are upregulated by the nitrate-specific signal and play a crucial role

in nitrate-induced accumulation of tZ-type CKs (Engelsberger and Schulze, 2012; Maeda

et al., 2018). To test whether *CYP735A*s of rice are nitrogen-inducible, we analyzed the

expression levels of *CYP735A*s in roots supplemented with ammonium or nitrate (Fig. 8).

OsIPT4 and *OsNIA1* were used as indicators for glutamine-related and nitrate-specific

responses, respectively (Kamada-Nobusada et al., 2013). *CYP735A3* was induced by both

ammonium and nitrate in a manner similar to *OsIPT4*, while the expression of *CYP735A4* was

upregulated only by nitrate like *OsNIA1* (Fig. 8A). To clarify whether *CYP735A*s are regulated

by glutamine-related and/or nitrate-specific signal, we examined the effect of methionine

sulfoximine (MSX), an inhibitor of glutamine synthetase (Fig. 8B) (Lam et al., 1996). MSX

pre-treatment inhibited the nitrate-triggered induction of *CYP735A3* but did not affect

CYP735A4. On the other hand, *CYP735A3* was upregulated by glutamine supplement in the

presence of MSX, while *CYP735A4* was not. These responses of *CYP735A3* and *CYP735A4*

to nitrogen sources and MSX are similar to those of *OsIPT4* and *OsNIA1*, respectively,

indicating that *CYP735A3* and *CYP735A4* are regulated by glutamine-related and

nitrate-specific signals, respectively. Together, these results suggested that tZ-type CK

biosynthesis in rice is controlled by dual nitrogen signals, which is different from that in

Arabidopsis.

Discussion

Angiosperms can be divided into four main categories in terms of their CK side-chain profile and number of cotyledons; iP/tZ-type-abundant dicot, cZ-type-abundant dicot, iP/tZ-type-abundant monocot and cZ-type-abundant monocot (Gajdošová et al., 2011). The function of some of these CKs has been demonstrated in a few dicotyledonous plant species. For example, in *Arabidopsis*, an iP/tZ-type-abundant dicot plant, CYP735As catalyze the biosynthesis of tZ-type CKs, which plays a specific role in shoot growth promotion (Kiba et al., 2013). Recently, it was suggested that CYP735A plays a similar role in *Jatropha curcas,* which is also an iP/tZ-type-abundant dicot plant (Cai et al., 2018). However, the function of CYP735A and tZ-type CKs in plants in other categories remains unknown. Here, we investigated the biosynthetic mechanism and function of tZ-type CKs in cZ-type-abundant monocot plant rice. Characterization of the *a3a4* double mutant demonstrated that *CYP735A*s play a central role in tZ-type CK biosynthesis in rice as in the iP/tZ-type-abundant dicot plants but that the physiological function of tZ-type CKs in rice is not the same as that in *Arabidopsis*, providing new insights into the biological significance of side-chain variations in plants.

Arabidopsis CYP735As catalyze the *trans*-hydroxylation of iP-type CKs to form tZ-type CKs (Takei et al., 2004; Kiba et al., 2013). Complementation test of *atcypDM* and CK quantification of *a3a4* demonstrated that CYP735A3 and CYP735A4 are the functional orthologs of *Arabidopsis* CYP735As (Figs. 1 and 3; Tables S1 to S3). The tZ-type CKs in *a3a4* was scarce but it did not completely disappear (Fig. 3; Tables S2 and S3). This is similar to what has been observed in the CYP735A-null mutants (*atcypDM*) of *Arabidopsis* and *J. curcas* (Kiba et al., 2013; Cai et al., 2018), suggesting the existence of minor

alternative pathway(s) in both iP/tZ-type-abundant and cZ-type-abundant plants. The

isomerization of cZ-type CKs is a possible pathway because *cis*-to-*trans* isomerization

activity has been detected in various plants, including rice (Bassil et al., 1993; Gajdošová

et al., 2011; Kudo et al., 2012). Although the isomerization pathway might be important

under certain biological contexts, given that the pathway is minor even in the

cZ-type-abundant rice, and that *CYP735A* homologs are found in all angiosperms whose

genome is completely sequenced (Hansen et al., 2021), the CYP735A pathway seems to be

the predominant pathway for tZ-type CK biosynthesis in angiosperms.

The *a3a4* mutants displayed severe growth retardation in vegetative and maturation

phases (Figs 4, 5, and S5). We concluded that this phenotype is the result of diminished CK

activity caused by a reduction in tZ-type CKs based on the following reasons: (i) the double

mutant had severely reduced level of tZ-type CKs (Fig. 3; Tables S2 and S3); (ii) the cytokinin

activity, measured by expression levels of immediate-early CK-inducible *RR* genes, was

reduced in the double mutant (Fig. 6); (iii) The phenotypes of the double mutant is partly

similar to those observed in rice mutants with severely reduced CK activity, such as *OsRR6*

overexpressors (Hirose et al., 2007), *rr21/22/23* (Worthen et al., 2019) and *hk5 hk6*

(Burr et al., 2021); (iv) our results were similar to the positive role of tZ-type CKs in rice

growth suggested through the analysis of OsABCG18, a transporter involved in root-to-shoot

CK translocation (Zhao et al., 2019); (v) similar conclusion was drawn from the data

obtained from equivalent experiments on *CYP735A1* and *CYP735A2* in *Arabidopsis* (Kiba et

al., 2013).

The cZ-type CKs are the dominant variants in terms of quantity comprising more

than 80% of total CK in various rice tissues and are suggested to be an active CK that plays a

role in normal growth and development in rice (Kojima et al., 2009; Choi et al., 2012;

Kudo et al., 2012; Osugi and Sakakibara, 2015). Our CK quantification results showed

that the levels of any of the cZ-type CK derivatives were not consistently altered in *a3a4* (Fig

3; Tables S2 and S3), indicating that cZ-type CKs are not directly relevant to the *a3a4*

phenotype. The iP-type CKs are also considered to be an important variant in rice because of

the identification of a receptor (OsHK6) with preferential affinity for iP (Choi et al., 2012).

In *a3a4*, the iP-type CK levels were increased to compensate for the loss of tZ-type CK in the

root but not sufficiently increased in the shoot (Fig. 3; Tables S2 and S3). From these results it

could be interpreted that the cause of the *a3a4* phenotype is an alteration in CK side-chain

profile (CK quality) but not CK quantity in the root and alteration in both CK quality and/or

quantity in the shoot. Together, our results suggest that the function of tZ-type CKs is

different from that of iP-type and cZ-type CKs at least in rice roots.

The phenotypes of *a3a4* were observed both in roots and shoots, including reduced

root proliferation, delayed plastochron and decreased leaf size, panicle size and tiller number

(Figs. 4, 5 and S5), indicating the extensive role of tZ-type CKs as growth enhancers in

various organs in rice. Although it is difficult to simply compare the phenotypes of *a3a4* with

those of *atcypDM* because monocot and dicot have remarkably different body plan, the major

difference lies in the existence of phenotype in the root. The *a3a4* displayed reduced

cytokinin activity and proliferation in the root (Figs. 4, 6 and S5), while no apparent

difference was detected in *atcypDM* root (Kiba et al., 2013), suggesting that the role of

tZ-type CKs in rice varies from that in *Arabidopsis*. The shoot specific-function of tZ-type

CKs in *Arabidopsis* is partly explained by the ligand preference of cytokinin receptors

(Spíchal et al., 2004; Romanov et al., 2006; Stolz et al., 2011; Kiba et al., 2013).

Among the five genes encoding cytokinin receptors (OsHK3-6, CHARK) in rice, so far, the

ligand preference of OsHK3, OsHK4 and OsHK6 has been investigated (Choi et al., 2012).

However, no receptor with preferential affinity for tZ-type CKs has been identified and the

mechanism underlying the functional specificity of tZ-type CKs in rice remains to be

elucidated.

Consistent with the functional distinction of tZ-type CKs between rice and

Arabidopsis, we also showed differences in responses to some internal and external cues of

*CYP735A*s (Figs. 2, 7 and 8). Rice *CYP735A*s are repressed by CKs (Tsai et al. 2012), while

*Arabidopsis CYP735A*s are induced (Takei et al., 2004), suggesting that rice and

Arabidopsis have negative feedback and positive feedback regulation, respectively, on tZ-type

CK function. The expression level of *CYP735A4* is high in leaves (Fig. 2) but that of

*Arabidopsis CYP735A*s is not (Takei et al., 2004; Kiba et al., 2013). Responses of these

genes to nitrogen supplement were also found to be different. *CYP735A3* and *CYP735A4* are

upregulated by glutamine-related and nitrate-specific signals, respectively, in rice (Fig. 8)

while *CYP735A*s of *Arabidopsis* are induced only by nitrate-specific signal (Engelsberger

and Schulze, 2012; Maeda et al., 2018). Although ammonium is known to be the major

form of inorganic nitrogen source in paddy field, rice utilizes not only ammonium but also

nitrate generated by nitrification on the root surface as inorganic nitrogen sources (Kirk and

Kronzucker, 2005; Yan et al., 2011). For *Arabidopsis*, which grows on aerobic soil, nitrate

is the major inorganic nitrogen source (Nacry et al., 2013), indicating that *CYP735As* of

rice and *Arabidopsis* respond to nitrogen in a way relevant to the nitrogen source availability

of the plants. Nitrogen-dependent tZ-type CK accumulation has been proposed to be involved

in growth enhancement in response to nitrogen supply in various plants (Takei et al.,

2001b; Kamada-Nobusada et al., 2013; Ohashi et al., 2017; Maeda et al., 2018;

Sakakibara, 2021; Shibasaki et al., 2021). Thus, the dual nitrogen-signal-dependent

upregulation of tZ-type CK biosynthesis might play a similar role in rice. Consistent to that,

there are reports suggesting that rice grows better with simultaneous supply of nitrate and

ammonium than exclusive supply of nitrate or ammonium (Chanch and Ohira, 1982;

Ying-Hua et al., 2006). Together, these differences between rice and *Arabidopsis* might be the

result of evolution and/or domestication to optimize growth depending on their body plan and

growth environments, though further investigation into the physiological function of tZ-type

CKs in other plant species is necessary to clarify this issue.

Materials and Methods

Plant material and growth conditions

Arabidopsis thaliana ecotype Columbia (Col-0) was used as the wild type. The

cyp735a1-2 cyp735a2-2 double mutant was characterized previously (Kiba et al., 2013).

Arabidopsis plants were grown on half-strength Murashige-Skoog (1/2x MS) agar plates (1%

agar and 1% sucrose) placed vertically or on soil (Supermix A, Sakata, Japan) at 22°C under

fluorescent light (100 µmol m^{-2} s^{-1}, 16 h light/8 h dark).

Oryza sativa cultivar Nipponbare (NB) was used as the wild type. Rice seeds were

germinated on moist filter paper at 30°C for 2-3 days in the dark and then grown under

hydroponic culture or soil culture conditions. For hydroponic culture, germinated seeds were

sown on mesh trays floating on the nutrient solution (Makino et al., 1985) with or without

nitrogen source (NH$_4$NO$_3$ was replaced with KCl) and grown in an environment-controlled

greenhouse at 28°C with a 15-h light (500 µmol m^{-2} s^{-1})/9-h dark cycle. The nutrient solution

was renewed every 3-4 days. To analyze *CYP735A3* and *CYP735A4* expression in response to

nitrogen, 12-day-old seedlings grown on the nutrient solution without nitrogen source were

incubated with the nutrient solution without nitrogen source supplemented with KCl, NH_4Cl,

KNO_3, or methionine sulfoximine (MSX). For soil culture, germinated seeds were sown on a

synthetic soil (SunAgro, No. 3) and grown in an environment-controlled greenhouse at 28°C

with a 15-h light (500 µmol m^{-2} s^{-1})/9- h dark cycle. Fifteen-day-old seedlings were

transplanted in perforated plastic pots ($10 \times 12 \times 15$ cm) filled with nutrient-free soil (Aichi

Medel) supplemented with slow-release fertilizer (N, P, and K at 0.3, 0.3, and 0.3 g/kg soil,

respectively) and grown in a greenhouse at Nagoya University with supplemental lighting

(13-h light/11-h dark cycle).

Generation of transgenic lines overexpressing *CYP735A3* or *CYP735A4* in *cyp735a1*

cyp735a2

The *CYP735A3* and *CYP735A4* cDNA were amplified with the specific primer sets

oxCYP735A3-F/R and oxCYP735A4-F/R, respectively (Table S4). The amplified fragments

were cloned into the *Xba* I/*Sac* I site of pBI121 (Clontech) located downstream of the

cauliflower mosaic virus (CaMV) 35S promoter and introduced into *cyp735a1-2 cyp735a2-2*

double mutant by floral-dipping (Clough and Bent, 1998).

CRISPR-Cas9 mutagenesis of *CYP735A3* and *CYP735A4*

The mutants of *CYP735A3* and *CYP735A4* were generated using the transfer

RNA-based-multiplex CRISPR-Cas9 vector, pMgPoef4_129-2A-GFP (Toda et al., 2019).

Guide sequences for *CYP735A3* and *CYP735A4* were designed by CHOPCHOP (Labun et al.,

2019) and two multiplex CRISPR-Cas9 vectors pMg129_A3A4-1 and pMg129_A3A4-2,

each harboring three guide sequences for *CYP735A3* and *CYP735A4* sequences, were

constructed as described (Toda et al., 2019). pMg129_A3A4-1 and pMg129_A3A4-2

contained guide sequences g3-1, g4-1-1, and g4-1-2 (Fig. S1), and g3-2, g4-1-2, and g4-1-2

(Fig. S2), respectively. Transgenic rice plants were generated by the *Agrobacterium*

tumefaciens-mediated method (Hirose et al., 2005) using EHA105 strain harboring the

vectors. Mutations were identified by DNA sequencing of PCR products amplified with

specific primer sets, gCYP735A3-F2/R2 and gCYP735A4-F3R3 (Table S4), and genomic

DNA prepared from T1 plants. A T1 line transformed using pMg129_A3A4-2 but without any

mutations in *CYP735A3* and *CYP735A* was used as the vector control line. For assays,

cyp735a3-1, *cyp735a4-2*, *cyp735a3-1 cyp735a4-1*, *cyp735a3-2 cyp735a4-2* and vector control

identified by genotyping progenies of these lines were used.

Quantification of plant hormones

Cytokinin levels were determined as described previously (Kojima et al., 2009;

Kojima and Sakakibara, 2012) using an ultra-performance liquid chromatograph coupled

with a tandem quadrupole mass spectrometer (ACQUITY UPLC™ System/XEVO-TQS;

Waters, Milford, MA, USA) with an octadecylsilyl (ODS) column (ACQUITY UPLC HSS

T3, 1.8 μm, 2.1 mm × 100 mm, Waters). In the results, iP, iP-riboside, iP-riboside

5'-phosphates, iP-7-*N*-glucoside, and iP-9-*N*-glucoside are collectively referred to as "iP-type

cytokinin"; tZ, tZ-riboside, tZ-riboside 5'-phosphates, tZ-7-*N*-glucoside, tZ-9-*N*-glucoside,

tZ-*O*-glucoside, and tZR-*O*-glucoside are collectively referred to as "tZ-type cytokinin"; cZ,

cZ-riboside, cZ-riboside 5'-phosphates, cZ-*O*-glucoside, and cZR-*O*-glucoside are

collectively referred to as "cZ-type cytokinin".

Gene expression analysis

Total RNA was extracted from root and shoot samples using the RNeasy Plant Mini

kit (QIAGEN) in combination with the RNase-Free DNase set (QIAGEN). Total RNA was

used for first strand cDNA synthesis by the ReverTra Ace qPCR RT Master Mix (Toyobo).

Quantitative reverse transcription-PCR (qRT-PCR) was performed on a Quant Studio 3

Real-Time PCR system (Thermo Fisher) with the KAPA SYBR Fast qPCR kit (KAPA

Biosystems). *ZIURP1* (LOC_Os03g08010/Os03g0234200) was used as an internal control

because this gene has been shown to be one of the most stably expressed genes in rice

(Pabuayon et al., 2016). Primer sets are listed in Table S4.

Promoter-GUS analysis

The *CYP735A3* promoter (proCYP735A3; -3636 to +21 bp relative to the inferred initiation

codon) and *CYP735A4* promoter (proCYP735A4; -3623 to +21 bp) were amplified with

PrimeSTAR GXL DNA polymerase (Takara) and the specific primer sets proCYP735A3-F/R

and proCYP735A4-F/R, respectively (Table S4). The fragments were cloned into

pENTR/D-TOPO vector (Thermo Fisher), sequenced, and then integrated into the GATEWAY

binary vector pCAMBIA-GW-GUS (Kudo et al., 2012) to generate

pCAMBIA1390-proCYP735A3:GUS and pCAMBIA1390-proCYP735A4:GUS vectors.

Transgenic rice plants were generated by the *Agrobacterium tumefaciens*-mediated method

(Hirose et al., 2005) using EHA105 strain harbouring the vectors. Histochemical analysis of

GUS activity was conducted as described previously (Hirose et al., 2005; Kiba et al.,

2018). Fresh and GUS-stained samples were fixed and then embedded in Technovit 7100 resin

(Heraeus Kulzer) as instructed by the manufacturer. Sections were produced using a Leica

RM2165 microtome (Leica) and observed under an Olympus BX51 microscope (Olympus).

Morphological analyses

The rosette diameter, shoot length, root length, leaf blade length, leaf sheath length,

leaf width, panicle length, and culm length were measured manually or determined from

pictures using ImageJ (https://imagej.nih.gov/ij/). The days leading to the appearance of a

panicle in main stem is defined as "days to heading".

Treatment with phytohormones and nitrogen sources

For phytohormone treatments, 12-day-old Nipponbare seedlings grown in hydroponic

culture with the nutrient solution (Makino et al., 1985) containing 1 mM NH_4NO_3 were

used. Phytohormone stock solutions were prepared by dissolving N^6-(Δ^2-isopentenyl)adenine

(iP), *trans*-zeatin (tZ), indole-3-acetic acid (IAA), and abscisic acid (ABA) in dimethyl

sulfoxide (DMSO) at 100 mM and added to the nutrient solution. Roots of the seedlings were

submerged in the nutrient solution containing 1 µM iP, 1 µM tZ, 10 µM IAA, 50 µM ABA,

0.001% DMSO (mock for iP and tZ treatment), or 0.05% DMSO (mock for IAA and ABA

treatment) and harvested after 2 h.

For nitrogen treatments, Nipponbare seedlings were grown in hydroponic culture

with the nitrogen-free nutrient solution (1 mM NH_4NO_3 was replaced with 1 mM KCl) for 12

days. Roots of the seedlings were incubated with the nitrogen-free nutrient solution

supplemented with 5 mM KCl, 5 mM NH$_4$Cl, or 5 mM KNO$_3$ for 2 h or 4 h. In case of MSX

and nitrogen co-treatment, roots were pre-treated with the nitrogen-free nutrient solution

containing 2 mM MSX for 2 h and then incubated with the MSX containing nitrogen-free

nutrient solution supplemented with 5 mM KCl, 5 mM KNO$_3$ or 50 mM glutamine for 2 h.

Statistical analysis

Data are shown as means ± standard deviation (SD) of one representative

experiment. In order to examine whether hormone concentration, gene expression, or

morphological data were significantly different between treatments, Student's *t*-test, and

Tukey's honestly significant difference (HSD) test were performed using KaleidaGraph ver.

4.1 software (Synergy Software).

Accession numbers

Sequence data for the genes described in this article can be found in The

Arabidopsis Information Resource database (http://www.arabidopsis.org) and Oryzabase

(https://shigen.nig.ac.jp/rice/oryzabase/) under the following accession numbers: *CYP735A1*

(At5g38450),   *CYP735A2*   (At1g67110),   *ACT8*   (AT1G49240),   *CYP735A3*

(LOC_Os08g33300/Os08g0429800),   *CYP735A4*   (LOC_Os09g23820/Os09g0403300),

ZIURP1 (LOC_Os03g08010/Os03g0234200), *OsRR1* (LOC_Os04g36070/Os04g0442300),

OsRR2 (LOC_Os02g35180/Os02g0557800), *OsRR4* (LOC_Os01g72330/Os01g0952500),

OsRR6 (LOC_Os04g57720/Os04g0673300), *OsRR9* (LOC_Os11g04720/Os11g0143300),

OsRR10 (LOC_Os12g04500/Os12g0139400), *OsRR21* (LOC_Os03g12350/Os03g0224200),

OsRR23   (LOC_Os02g55320/Os02g0796500),   *OsSIPP2C1*

(LOC_Os09g15670/Os09g0325700), *OsGH3.2* (LOC_Os01g55940/Os01g0764800), *OsIPT4*

(LOC_Os03g59570/Os03g0810100), *OsNIA1* (LOC_Os08g36480/Os08g0468100).

Acknowledgments

This work was supported by JSPS KAKENHI (No. JP18H04793 and JP20H02888

for T.K., JP17H06473 for H.S.).

Author contributions

T.K. and H.S. designed the experiments; T.K., K.M., A.M., Y.T., M.K., T.H., Y.O.,

K.O., H.S. performed the experiments, T.K., K.M., A.M. analyzed the data, and T.K. and H.S.

wrote the paper.

References

Bassil N V, Mok D, Mok MC (1993) Partial purification of a cis-trans-isomerase of zeatin
from immature seed of *Phaseolus vulgaris* L. Plant Physiol **102**: 867–872

Burr CA, Sun J, Yamburenko M V., Willoughby A, Hodgens C, Boeshore SL, Elmore A,
Atkinson J, Nimchuk ZL, Bishopp A, et al (2021) The HK5 and HK6 cytokinin
receptors mediate diverse developmental pathways in rice. Development **147**: dev191734

Cai L, Zhang L, Fu Q, Xu ZF (2018) Identification and expression analysis of cytokinin
metabolic genes IPTs, CYP735A and CKXs in the biofuel plant *Jatropha curca*s. PeerJ
2018: e4812

Chanch T, Ohira K (1982) Comparison of the uptake and assimilation of ammonium and
nitrate in indica and japonica rice plants using the tracer 15N method. Soil Sci Plant Nutr
28: 79–90

Choi J, Lee J, Kim K, Cho M, Ryu H, An G, Hwang I (2012) Functional identification of
OsHk6 as a homotypic cytokinin receptor in rice with preferential affinity for iP. Plant
Cell Physiol **53**: 1334–1343

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated

transformation of Arabidopsis thaliana. Plant J **16**: 735–743

Coskun D, Britto DT, Shi W, Kronzucker HJ (2017) Nitrogen transformations in modern

agriculture and the role of biological nitrification inhibition. Nat Plants **3**: 17074

Du H, Wu N, Fu J, Wang S, Li X, Xiao J, Xiong L (2012) A GH3 family member,

OsGH3-2, modulates auxin and abscisic acid levels and differentially affects drought and

cold tolerance in rice. J Exp Bot **63**: 6467–6480

Engelsberger WR, Schulze WX (2012) Nitrate and ammonium lead to distinct global

dynamic phosphorylation patterns when resupplied to nitrogen-starved Arabidopsis

seedlings. Plant J **69**: 978–995

Gajdošová S, Spíchal L, Kamínek M, Hoyerová K, Novák O, Dobrev PI, Galuszka P,

Klíma P, Gaudinová A, Žižková E, et al (2011) Distribution, biological activities,

metabolism, and the conceivable function of *cis*-zeatin-type cytokinins in plants. J Exp

Bot **62**: 2827–2840

Hansen CC, Nelson DR, Møller BL, Werck-Reichhart D (2021) Plant cytochrome P450

plasticity and evolution. Mol Plant **14**: 1244–1265

Hirose N, Makita N, Kojima M, Kamada-Nobusada T, Sakakibara H (2007)

Overexpression of a type-A response regulator alters rice morphology and cytokinin

metabolism. Plant Cell Physiol **48**: 523–539

Hirose N, Makita N, Yamaya T, Sakakibara H (2005) Functional characterization and

expression analysis of a gene, OsENT2, encoding an equilibrative nucleoside transporter

in rice suggest a function in cytokinin transport. Plant Physiol **138**: 196–206

Hirose N, Takei K, Kuroha T, Kamada-Nobusada T, Hayashi H, Sakakibara H (2008)

Regulation of cytokinin biosynthesis, compartmentalization and translocation. J Exp Bot

59: 75–83

Kakimoto T (2001) Identification of plant cytokinin biosynthetic enzymes as dimethylallyl

diphosphate : ATP/ADP isopentenyltransferases. Plant Cell Physiol **42**: 677–685

Kamada-Nobusada T, Makita N, Kojima M, Sakakibara H (2013) Nitrogen-dependent

regulation of *de novo* cytokinin biosynthesis in rice: the role of glutamine metabolism as

an additional signal. Plant Cell Physiol **54**: 1881–1893

Kawai M, Tabata R, Ohashi M, Honda H, Kamiya T, Kojima M, Takebayashi Y, Oishi S,

Okamoto S, Hachiya T, et al (2022) Regulation of ammonium acquisition and use in

Oryza longistaminata ramets under nitrogen source heterogeneity. Plant Physiol **188**:

2364–2376

Kiba T, Inaba J, Kudo T, Ueda N, Konishi M, Mitsuda N, Takiguchi Y, Kondou Y,

Yoshizumi T, Ohme-Takagi M, et al (2018) Repression of nitrogen-starvation

responses by members of the Arabidopsis GARP-type transcription factor NIGT1/HRS1 subfamily. Plant Cell **30**: 925–945

Kiba T, Kudo T, Kojima M, Sakakibara H (2011) Hormonal control of nitrogen acquisition: roles of auxin, abscisic acid, and cytokinin. J Exp Bot **62**: 1399–1409

Kiba T, Takebayashi Y, Kojima M, Sakakibara H (2019) Sugar-induced *de novo* cytokinin biosynthesis contributes to Arabidopsis growth under elevated CO_2. Sci Rep **9**: 1–15

Kiba T, Takei K, Kojima M, Sakakibara H (2013) Side-chain modification of cytokinins controls shoot growth in *Arabidopsis*. Dev Cell **27**: 452–461

Kieber JJ, Schaller GE (2018) Cytokinin signaling in plant development. Development **145**: dev149344

Kirk GJ, Kronzucker HJ (2005) The potential for nitrification and nitrate uptake in the rhizosphere of wetland plants: a modelling study. Ann Bot **96**: 639–646

Kojima M, Kamada-Nobusada T, Komatsu H, Takei K, Kuroha T, Mizutani M, Ashikari M, Ueguchi-Tanaka M, Matsuoka M, Suzuki K, et al (2009) Highly sensitive and high-throughput analysis of plant hormones using MS-probe modification and liquid chromatography tandem mass spectrometry: an application for hormone profiling in *Oryza sativa*. Plant Cell Physiol **50**: 1201–1214

Kojima M, Sakakibara H (2012) Highly sensitive high-throughput profiling of six phytohormones using MS-probe modification and liquid chromatography-tandem mass spectrometry. Methods Mol Biol **918**: 151–164

Kudo T, Makita N, Kojima M, Tokunaga H, Sakakibara H (2012) Cytokinin activity of *cis*-zeatin and phenotypic alterations induced by overexpression of putative *cis*-Zeatin-*O*-glucosyltransferase in rice. Plant Physiol **160**: 319–331

Kurakawa T, Ueda N, Maekawa M, Kobayashi K, Kojima M, Nagato Y, Sakakibara H, Kyozuka J (2007) Direct control of shoot meristem activity by a cytokinin-activating enzyme. Nature **445**: 652–655

Kuroha T, Tokunaga H, Kojima M, Ueda N, Ishida T, Nagawa S, Fukuda H, Sugimoto K, Sakakibara H (2009) Functional analyses of LONELY GUY cytokinin-activating enzymes reveal the importance of the direct activation pathway in *Arabidopsis*. Plant Cell **21**: 3152–3169

Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E (2019) CHOPCHOP v3: Expanding the CRISPR web toolbox beyond genome editing. Nucleic Acids Res **47**: W171–W174

Lam HM, Coschigano KT, Oliveira IC, Melo-Oliveira R, Coruzzi GM (1996) The molecular-genetics of nitrogen assimilation into amino acids in higher plants. Annu Rev

627 Plant Physiol Plant Mol Biol **47**: 569–593

628 **Li YS, Sun H, Wang ZF, Duan M, Huang SD, Yang J, Huang J, Zhang HS** (2013) A novel

629  nuclear protein phosphatase 2C negatively regulated by ABL1 is involved in abiotic

630  stress and panicle development in rice. Mol Biotechnol **54**: 703–710

631 **Lomin SN, Yonekura-Sakakibara K, Romanov GA, Sakakibara H** (2011) Ligand-binding

632  properties and subcellular localization of maize cytokinin receptors. J Exp Bot **62**: 5149–

633  5159

634 **Macková H, Hronková M, Dobrá J, Turečková V, Novák O, Lubovská Z, Motyka V,**

635  **Haisel D, Hájek T, Prášil IT, et al** (2013) Enhanced drought and heat stress tolerance of

636  tobacco plants with ectopically enhanced cytokinin oxidase/dehydrogenase gene

637  expression. J Exp Bot **64**: 2805–2815

638 **Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y, Sakakibara H,**

639  **Yanagisawa S** (2018) A NIGT1-centred transcriptional cascade regulates nitrate

640  signalling and incorporates phosphorus starvation signals in *Arabidopsis*. Nat Commun

641  **9**: 1–14

642 **Makino A, Mae T, Ohira K** (1985) Photosynthesis and ribulose-1,5-bisphosphate

643  carboxylase oxygenase in rice leaves from emergence through senescence -

644  quantitative-analysis by carboxylation oxygenation and regeneration of ribulose

645  1,5-bisphosphate. Planta **166**: 414–420

646 **Midlik A, Navrátilová V, Moturu TR, Koča J, Svobodová R, Berka K** (2021) Uncovering

647  of cytochrome P450 anatomy by SecStrAnnotator. Sci Rep **11**: 12345

648 **Miyawaki K, Matsumoto-Kitano M, Kakimoto T** (2004) Expression of cytokinin

649  biosynthetic isopentenyltransferase genes in *Arabidopsis*: tissue specificity and

650  regulation by auxin, cytokinin, and nitrate. Plant J **37**: 128–138

651 **Miyawaki K, Tarkowski P, Matsumoto-Kitano M, Kato T, Sato S, Tarkowska D, Tabata**

652  **S, Sandberg G, Kakimoto T** (2006) Roles of Arabidopsis ATP/ADP

653  isopentenyltransferases and tRNA isopentenyltransferases in cytokinin biosynthesis.

654  Proc Natl Acad Sci USA **103**: 16598–16603

655 **Mok DW, Mok MC** (2001) Cytokinin metabolism and action. Annu Rev Plant Physiol Plant

656  Mol Biol **52**: 89–118

657 **Mok MC, Mok DW, Armstrong DJ** (1978) Differential cytokinin structure-activity

658  relationships in Phaseolus. Plant Physiol **61**: 72–75

659 **Nacry P, Bouguyon E, Gojon A** (2013) Nitrogen acquisition by roots: physiological and

660  developmental mechanisms ensuring plant adaptation to a fluctuating resource. Plant

661  Soil **370**: 1–29

Ohashi M, Ishiyama K, Kojima S, Kojima M, Sakakibara H, Yamaya T, Hayakawa T
(2017) Lack of cytosolic glutamine synthetase1;2 activity reduces nitrogen-dependent
biosynthesis of cytokinin required for axillary bud outgrowth in rice seedlings. Plant Cell
Physiol **58**: 679–690

Osugi A, Kojima M, Takebayashi Y, Ueda N, Kiba T, Sakakibara H (2017) Systemic
transport of *trans*-zeatin and its precursor have differing roles in Arabidopsis shoots. Nat
Plants **3**: 1–6

Osugi A, Sakakibara H (2015) Q&A: How do plants respond to cytokinins and what is their
importance? BMC Biol **13**: 102

Pabuayon IM, Yamamoto N, Trinidad JL, Longkumer T, Raorane ML, Kohli A (2016)
Reference genes for accurate gene expression analyses across different tissues,
developmental stages and genotypes in rice for drought tolerance. Rice **9**: 32

Romanov GA, Lomin SN, Schmülling T (2006) Biochemical characteristics and
ligand-binding properties of Arabidopsis cytokinin receptor AHK3 compared to
CRE1/AHK4 as revealed by a direct binding assay. J Exp Bot **57**: 4051–4058

Sakakibara H (2006) Cytokinins: activity, biosynthesis, and translocation. Annu Rev Plant
Biol **57**: 431–449

Sakakibara H (2021) Cytokinin biosynthesis and transport for systemic nitrogen signaling.
Plant J **105**: 421–430

Schäfer M, Brütting C, Meza-Canales ID, Großkinsky DK, Vankova R, Baldwin IT,
Meldau S (2015) The role of cis-zeatin-type cytokinins in plant growth regulation and
mediating responses to environmental interactions. J Exp Bot. Oxford University Press,
pp 4873–4884

Schmitz RY, Skoog F (1972) Cytokinins: Synthesis and biological activity of geometric and
position isomers of zeatin. Plant Physiol **50**: 702–705

Shaw G (1994) Chemistry of adenine cytokinins. *In* DWS Mok, MC Mok, eds, Cytokinins:
Chemistry, Activity, and Function. CRC Press, Boca Rato, FL, pp 15–34

Shibasaki K, Takebayashi A, Makita N, Kojima M, Takebayashi Y, Kawai M, Hachiya T,
Sakakibara H (2021) Nitrogen nutrition promotes rhizome bud outgrowth via
regulation of cytokinin biosynthesis genes and an *Oryza longistaminata* ortholog of
FINE CULM 1. Front Plant Sci **12**: 788

Spíchal L, Rakova NY, Riefler M, Mizuno T, Romanov GA, Strnad M, Schmülling T
(2004) Two cytokinin receptors of *Arabidopsis thaliana*, CRE1/AHK4 and AHK3, differ
in their ligand specificity in a bacterial assay. Plant Cell Physiol **45**: 1299–1305

Stolz A, Riefler M, Lomin SN, Achazi K, Romanov GA, Schmülling T (2011) The

specificity of cytokinin signalling in *Arabidopsis thaliana* is mediated by differing ligand
affinities and expression profiles of the receptors. Plant J **67**: 157–168

Takei K, Sakakibara H, Sugiyama T (2001a) Identification of genes encoding adenylate
isopentenyltransferase, a cytokinin biosynthesis enzyme, in *Arabidopsis thaliana*. J Biol
Chem **276**: 26405–26410

Takei K, Sakakibara H, Taniguchi M, Sugiyama T (2001b) Nitrogen-dependent
accumulation of cytokinins in root and the translocation to leaf: implication of cytokinin
species that induces gene expression of maize response regulator. Plant Cell Physiol **42**:
85–93

Takei K, Yamaya T, Sakakibara H (2004) Arabidopsis CYP735A1 and CYP735A2 encode
cytokinin hydroxylases that catalyze the biosynthesis of *trans*-zeatin. J Biol Chem **279**:
41866–41872

Toda E, Koiso N, Takebayashi A, Ichikawa M, Kiba T, Osakabe K, Osakabe Y,
Sakakibara H, Kato N, Okamoto T (2019) An efficient DNA- and
selectable-marker-free genome-editing system using zygotes in rice. Nat Plants **5**: 363–
368

Tokunaga H, Kojima M, Kuroha T, Ishida T, Sugimoto K, Kiba T, Sakakibara H (2012)
Arabidopsis lonely guy (LOG) multiple mutants reveal a central role of the
LOG-dependent pathway in cytokinin activation. Plant J **69**: 355–365

Tsai Y-C, Weir NR, Hill K, Zhang W, Kim HJ, Shiu S-H, Schaller GE, Kieber JJ (2012)
Characterization of Genes Involved in Cytokinin Signaling and Metabolism from Rice.
Plant Physiol **158**: 1666–1684

Vanková R, Kosová K, Dobrev P, Vítámvás P, Trávníčková A, Cvikrová M, Pešek B,
Gaudinová A, Prerostová S, Musilová J, et al (2014) Dynamics of cold acclimation
and complex phytohormone responses in *Triticum monococcum* lines G3116 and DV92
differing in vernalization and frost tolerance level. Environ Exp Bot **101**: 12–25

Vyroubalová S, Václavíková K, Turečková V, Novák O, Šmehilová M, Hluska T,
Ohnoutková L, Frébort I, Galuszka P (2009) Characterization of new maize genes
putatively involved in cytokinin metabolism and their expression during osmotic stress
in relation to cytokinin levels. Plant Physiol **151**: 433–447

Werner T, Motyka V, Laucou V, Smets R, Van Onckelen H, Schmülling T (2003)
Cytokinin-deficient transgenic Arabidopsis plants show multiple developmental
alterations indicating opposite functions of cytokinins in the regulation of shoot and root
meristem activity. Plant Cell **15**: 2532–2550

Werner T, Schmülling T (2009) Cytokinin action in plant development. Curr Opin Plant Biol

12: 527–538

Worthen JM, Yamburenko M V, Lim J, Nimchuk ZL, Kieber JJ, Schaller GE (2019)

Type-B response regulators of rice play key roles in growth, development and cytokinin

signaling. Development **146**: dev174870

Xu GH, Fan XR, Miller AJ (2012) Plant nitrogen assimilation and use efficiency. Annu Rev

Plant Biol **63**: 153–182

Yan M, Fan XR, Feng HM, Miller AJ, Shen QR, Xu GH (2011) Rice OsNAR2.1 interacts

with OsNRT2.1, OsNRT2.2 and OsNRT2.3a nitrate transporters to provide uptake over

high and low concentration ranges. Plant Cell Environ **34**: 1360–1372

Yonekura-Sakakibara K, Kojima M, Yamaya T, Sakakibara H (2004) Molecular

characterization of cytokinin-responsive histidine kinases in maize. differential ligand

preferences and ¥response to cis-zeatin. Plant Physiol **134**: 1654–1661

Zhao J, Yu N, Ju M, Fan B, Zhang Y, Zhu E, Zhang M, Zhang K (2019) ABC transporter

OsABCG18 controls the shootward transport of cytokinins and grain yield in rice. J Exp

Bot **70**: 6277–6291

Zwack PJ, Rashotte AM (2015) Interactions between cytokinin signalling and abiotic stress

responses. J Exp Bot **66**: 4863–4871

Legends to Figures

Figure 1. Recovery of *trans*-zeatin-type cytokinin deficiency in the *Arabidopsis* mutant

cyp735a1 cyp735a2* by *CYP735A3* or *CYP735A4

(A) Quantitative RT-PCR analysis of *CYP735A3* (left) or *CYP735A4* (right) expression in

whole seedlings of Col-0, *cyp735a1 cyp735a2* mutant (*atcypDM*), and transgenic plants

constitutively expressing *CYP735A3* (*A3*-ox) or *CYP735A4* (*A4*-ox) in *atcypDM* background.

Seedlings were grown on 1/2x MS agar plates for 12 days before harvest. Expression levels

were normalized using *ACT8* as an internal control. Error bars represent standard deviation

(n=4 independent pools of more than ten plants). N.D., under the quantification detection

limit. (B) Rosette diameter of 28-day-old Col-0, *atcypDM,* and transgenic plants

constitutively expressing *CYP735A3* (*A3*-ox) or *CYP735A4* (*A4*-ox) in *atcypDM* background

grown on soil. A photograph of representative plants is shown. Error bars represent standard

deviation (n=7). Asterisks indicate statistically significant differences compared with

atcypDM ($p<0.01$, Student's *t*-test). Scale bar, 3 cm. (C) Cytokinin concentrations in whole

seedlings of Col-0, *atcypDM* and transgenic plants constitutively expressing *CYP735A3*

(*A3*-ox) or *CYP735A4* (*A4*-ox) in *atcypDM* background. Seedlings were grown on 1/2x MS

agar plates for 12 days before harvest. Error bars represent standard deviation (n=4

independent pools of more than ten plants). Asterisks indicate statistically significant

differences in tZ-type cytokinin concentration compared with *atcypDM* ($p<0.01$, Student's

t-test). The concentration of each cytokinin molecular species in roots and shoots is shown in

Table S1. gFW, gram fresh weight; tZ-type, *trans*-zeatin and its conjugates; cZ-type,

cis-zeatin and its conjugates; iP-type, N^6-(Δ^2-isopentenyl)adenine and its conjugates.

Figure 2. Expression pattern of *CYP735A3* and *CYP735A4* in various tissues

(A) Expression of *CYP735A3* and *CYP735A4* in Nipponbare seedlings. Coleoptiles and

seminal roots, and roots and shoots were harvested 4 days and 14 days, respectively, after

imbibition. Error bars represent standard deviation of biological replicates (n=4-5 independent

pools of more than 2 plants). (B) Expression of *CYP735A3* and *CYP735A4* in Nipponbare

plants in vegetative or early reproductive stages. 42-day-old (vegetative stage) and older

plants (reproductive stage) were used. The shoot apex parts of 42-day-old plants were

collected as shoot apical meristem (SAM). Leaf blade, leaf sheath and lamina joint were

harvested from the youngest fully expanded leaf. The young panicle is a mixture of 2 mm to

21 mm long young panicles collected from plants at the reproductive stage. Error bars

represent standard deviation of biological replicates (n=4 except for SAM and young panicle,

n=4 and n=7 independent pools of more than 2 plants for SAM and young panicle,

respectively). N.D., under the quantification detection limit. (C-E) Representative images of

GUS staining obtained from *proCYP735A3:GUS* transgenic seedlings grown for 8 days after

imbibition. Pictures of a whole seedling (C), first leaf (D, upper), crown root (D, lower), cross

section of the first leaf (E, upper), and cross section of seminal root (E, lower) are presented.

VBS, vascular bundle sheath; XY, xylem; PH, phloem; EN, endodermis; PE, pericycle. Scale

bars, (C) 1 cm; (D) 200 μm; (E) 50 μm.

Figure 3. The *cyp735a3 cyp735a4* double mutants show deficiency in *trans*-zeatin-type

cytokinins

Cytokinin concentrations in roots and shoots of WT strain Nipponbare (NB), vector control

(VC), and mutant strains *cyp735a3-1* (*a3-1*), *cyp735a4-2* (*a4-2*), *cyp735a3-1 cyp735a4-1*

(*a3a4-1*), and *cyp735a3-2 cyp735a4-2* (*a3a4-2*). Seedlings were grown hydroponically for 14

days (A) or 12 days (B) before shoots and roots were harvested. Error bars represent standard

deviation of biological replicates (n=3-6 independent pools of more than ten plants for A,

n=6-8 independent pools of more than ten plants for B). Asterisks indicate statistically

significant differences in tZ-type cytokinin concentration compared with NB ($p<0.05$,

Student's *t*-test). The concentration of each cytokinin molecular species in roots and shoots is

shown in Table S2 and Table S3. gFW, gram fresh weight; tZ-type, *trans*-zeatin and its

conjugates; cZ-type, *cis*-zeatin and its conjugates; iP-type, N^6-(Δ^2-isopentenyl)adenine and its

conjugates.

Figure 4. Disruption of *CYP735As* alters rice growth in seedling to vegetative phases

(A) A representative image of Nipponbare (NB), vector control (VC), and *cyp735a3-2*

cyp735a4-2 (*a3a4-2*) seedlings grown 12 days in hydroponic culture. Scale bar, 5 cm. (B)

Quantification of shoot length, shoot fresh weight, root length, and root fresh weight in

12-day-old seedlings. NB, VC, *cyp735a4-2* (*a4-2*), and *a3a4-2* seedlings were grown in

hydroponic culture. (C) Leaf number at 15 days after germination (D, E) Leaf sheath and leaf

blade length (D) and leaf width (E) of the fully expanded third leaf of NB, VC, *cyp735a3-1*

(*a3-1*), *cyp735a4-2* (*a4-2*), *cyp735a3-1 cyp735a4-1* (*a3a4-1*), and *a3a4-2* seedlings grown on

soil. (F) Quantification of leaf blade length, leaf sheath length, and leaf width of the fully

expanded ninth leaf in NB, VC, *a3a4-1*, and *a3a4-2* grown on soil. (G) Tiller number of NB,

VC, *a3a4-1*, and *a3a4-2* grown on soil for 35 days. Error bars represent standard deviation of

biological replicates (B, n=6-8; C, n=8-12; D, n=8-10; E, n=8-10; F, n=8-12; G, n=4-8).

Different lowercase letters indicate statistically significant differences as indicated by Tukey's

HSD test ($p<0.05$).

Figure 5. Disruption of *CYP735As* alters rice growth in reproductive to maturation

phases

(A) A representative image of Nipponbare (NB), vector control (VC), and *cyp735a3-1* (*a3-1*),

cyp735a4-1 (*a4-2*), and *cyp735a3-1 cyp735a4-1* (*a3a4-2*) plants grown 15 weeks on soil.

Scale bar, 20 cm. (B-D) Heading date (B), panicle and culm length (C), and tiller number (D)

of NB, VC, *a3-1*, *a4-2*, *cyp735a3-1 cyp735a4-1* (*a3a4-1*), and *a3a4-2*. (E) A representative

picture of mature panicles from NB and *a3a4-2*. Scale bar, 5 cm. (F, G) Panicle weight on the

main stem (F) and number of primary branches per panicle (G). Error bars represent standard

deviation of biological replicates (B, n=3-12; C, n=5-12; D, n=9-12; F, n=5-10; G, n=5-10).

Different lowercase letters indicate statistically significant differences as indicated by Tukey's

HSD test ($p<0.05$).

Figure 6. Expression of cytokinin inducible type-A response regulator genes is altered in

the *cyp735a3 cyp735a4* mutant

(A, B) Transcript levels of cytokinin inducible type-A response regulator genes, *OsRR1*,

OsRR2, *OsRR4*, *OsRR6*, and *OsRR9/10*, in shoots (A) and roots (B) of Nipponbare (NB),

vector control (VC), *cyp735a3-1 cyp735a4-1* (*a3a4-1*) and *cyp735a3-2 cyp735a4-2* (*a3a4-2*)

seedlings. *OsRR9/10* indicates that the primer set used could not distinguish *OsRR9* and

OsRR10. (C, D) Transcript levels of type-B response regulator genes, *OsRR21* and *OsRR23*,

in the shoot. (C) and in the root (D) of NB, VC, *a3a4-1* and *a3a4-2* seedlings. Seedlings were

grown hydroponically for 10 days before roots and shoots were harvested. Expression levels

were quantified by qRT-PCR analysis, normalized to *ZIURP1* as an internal control, and are

shown relative to the value of NB. Error bars represent standard deviation of biological

replicates (n=4 independent pools of 2 plants). Asterisks indicate statistically significant

differences compared to NB ($p<0.05$, Student's *t*-test).

Figure 7. Expression of *CYP735A3* and *CYP735A4* in response to phytohormones

(A) Expression of *CYP735A*s in response to cytokinin. Nipponbare (NB) seedlings grown for

12 days were treated with 0.001% DMSO (Mock), 1 µM iP (iP), or 1 µM tZ (tZ) and roots

were harvested after 2 h. (B) Expression of *CYP735A*s in response to abscisic acid (ABA) and

auxin. NB seedlings grown for 12 days were treated with 0.05% DMSO (Mock), 10 µM

indole-3-acetic acid (IAA) or 50 µM ABA and roots were harvested after 2 h. Expression

levels were quantified by qRT-PCR analysis, normalized to *ZIURP1* as an internal control,

and are shown relative to the value in mock treatment. Asterisks indicate statistically

significant differences compared with Mock ($p<0.05$, Student's *t*-test). Error bars represent

standard deviation (n=4 independent pools of more than 2 plants).

Figure 8. Expression of *CYP735A3* and *CYP735A4* in response to nitrogen sources

(A) Expression of *CYP735A*s in response to ammonium or nitrate supplement.

Twelve-day-old NB seedlings hydroponically grown in nitrogen-free nutrient solution were

incubated with 5 mM KCl (-N), 5 mM NH_4Cl (NH_4^+), or 5 mM KNO_3 (NO_3^-) for 2 h or 4 h.

Expression levels were quantified by qRT-PCR analysis, normalized to *ZIURP1* as an internal

control, and are shown relative to the value in 2 h of -N treatment. Error bars represent

standard deviation (n=4 independent pools of 3 plants). (B) Effect of methionine sulfoximine

on the induction of *CYP735A*s by nitrogen. Twelve-day-old NB seedlings hydroponically

grown in nitrogen-free nutrient solution were pre-treated with 2 mM MSX for 2 h and then

incubated with 5 mM KCl, 5 mM KNO_3 or 50 mM glutamine (GLN) for 2 h. Asterisks

indicate statistically significant differences compared with KCl or KCl+MSX ($p<0.05$,

Student's *t*-test). Expression levels were quantified by qRT-PCR analysis, normalized to

ZIURP1 as an internal control, and are shown relative to the value in KCl treatment. Error

bars represent standard deviation (n=4 independent pools of 3 plants).

Supplemental Figure 1. Phylogenetic tree of *Arabidopsis* cytochrome P450s, CYP735A3

and CYP735A4

(A) A phylogenetic tree of 293 *Arabidopsis* P450s, CYP735A3

(LOC_Os08g33300/Os08g0429800) and CYP735A4 (LOC_Os09g23820/Os09g0403300).

(B) Enlargement of a subclade with CYP735A3 and CYP735A4. Full-length amino acid

sequences were obtained from Plant P450 database

(https://erda.dk/public/vgrid/PlantP450/index.html). A phylogenetic tree was inferred by the

Neighbour-Joining method using MEGAX (https://www.megasoftware.net/). The tree is

drawn to scale, with branch lengths in the same units as those of the evolutionary distances

used to infer the phylogenetic tree. The evolutionary distances were computed using the JTT

matrix-based method and are in the units of the number of amino acid substitutions per site.

All positions containing gaps and missing data were eliminated. The values at the nodes

indicate the bootstrap values (using 500 replications).

Supplemental Figure 2. The *cyp735a3-1 cyp735a4-1* mutant generated by the

CRISPR/Cas9 system

(A) Schematic representation of CRISPR target sites to generate *cyp735a3-1* (*a3-1*) and

cyp735a4-1 (*a4-1*) alleles. Boxes represent exons; horizontal bars, introns; triangles, CRISPR

target sites. The blue box represents the "A helix". The bar indicates a 200 bp scale. (B)

Partial sequences of wild-type *CYP735A3* and *cyp735a3-1* (*a3-1*) mutant. The sequence

corresponding to a guide RNA used to generate the *a3-1* mutation (g3-1) is highlighted in

yellow. (C) Partial sequences of wild-type *CYP735A4* and *cyp735a4-1* (*a4-1*) mutant. The

sequence corresponding to guide RNAs used to generate the *a4-1* mutation (g4-1-1 and

g4-1-2) is highlighted in yellow. Numbers in (B) and (C) represent positions in a genome

sequence when the first nucleotide of the putative start codon is counted as 1. The red letter

and red dash indicate an inserted and deleted sequence, respectively.

Supplemental Figure 3. The *cyp735a3-2 cyp735a4-2* mutant generated by CRISPR/Cas9

system

(A) Schematic representation of CRISPR target sites to generate *cyp735a3-2* (*a3-2*) and *cyp735a4-2* (*a4-2*) alleles. Boxes represent exons; horizontal bars, introns; triangles, CRISPR target sites. The blue box represents the "A helix". The bar indicates a 200 bp scale. (B) Partial sequences of wild-type *CYP735A3* and *cyp735a3-2* (*a3-2*) mutant. The sequence corresponding to a guide RNA used to generate the *a3-2* mutation (g3-2) is highlighted in yellow. (C) Partial sequences of wild-type *CYP735A4* and *cyp735a4-2* (*a4-2*) mutant. The sequence corresponding to guide RNAs used to generate the *a4-2* mutation (g4-2-1 and g4-2-2) is highlighted in green. Numbers in (B) and (C) represent positions in a genome sequence when the first nucleotide of the putative start codon is counted as 1. The red letter and red dash indicate an inserted and deleted sequence, respectively.

Supplemental Figure 4. Deduced amino acid sequences of *CYP735A3*, *cyp735a3-1* and

cyp735a3-2

Deduced amino acid sequences of *CYP735A3*, *cyp735a3-1* (*a3-1*), and *cyp735a3-2* (*a3-2*) were aligned by Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/). Triangles indicate CRISPR target sites.

Supplemental Figure 5. Deduced amino acid sequences of *CYP735A4*, *cyp735a4-1* and

cyp735a4-2

Deduced amino acid sequences of *CYP735A4*, *cyp735a4-1* (*a4-1*), and *cyp735a4-2* (*a4-2*)

were aligned by Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/). Triangles

indicate CRISPR target sites. The "A helix" is highlighted in blue.

Supplemental Figure 6. Effect of disruption of *CYP735A*s on vegetative growth

(A) A representative image of Nipponbare (NB), vector control (VC), *cyp735a3-1 cyp735a4-1*

(*a3a4-1*), and *cyp735a3-1* (*a3-1*) seedlings grown 14 days in hydroponic culture. Scale bar, 5

cm. (B) Quantification of shoot length, shoot fresh weight, root length, and root fresh weight

of 14-day-old seedlings. NB, VC, *cyp735a3-1* (*a3-1*), and *a3a4-1* seedlings were grown in

hydroponic culture. (C) A representative image of NB, VC, *a3-1*, *cyp735a4-1* (*a4-2*),

cyp735a3-2 cyp735a4-2 (*a3a4-2*) seedlings grown for 43 days on soil. Scale bar, 20 cm. (D)

Plant height and leaf number of NB, VC, *a3-1*, *a4-2*, *a3a4-1*, and *cyp735a3-2 cyp735a4-2*

(*a3a4-2*) grown for 30 days on soil. Error bars represent standard deviation of biological

replicates (B, n=6-13; D, n=9-12). Different lowercase letters indicate statistically significant

differences as indicated by Tukey's HSD test ($p<0.05$).

Figure 1. Recovery of *trans*-zeatin-type cytokinin deficiency in the *Arabidopsis* mutant *cyp735a1 cyp735a2* by *CYP735A3* or *CYP735A4*

(A) Quantitative RT-PCR analysis of *CYP735A3* (left) or *CYP735A4* (right) expression in whole seedlings of Col-0, *cyp735a1 cyp735a2* mutant (*atcypDM*), and transgenic plants constitutively expressing *CYP735A3* (*A3-*ox) or *CYP735A4* (*A4-*ox) in *atcypDM* background. Seedlings were grown on 1/2x MS agar plates for 12 days before harvest. Expression levels were normalized using *ACT8* as an internal control. Error bars represent standard deviation (n=4 independent pools of more than ten plants). N.D., under the quantification detection limit. (B) Rosette diameter of 28-day-old Col-0, *atcypDM,* and transgenic plants constitutively expressing *CYP735A3* (*A3-*ox) or *CYP735A4* (*A4-*ox) in *atcypDM* background grown on soil. A photograph of representative plants is shown. Error bars represent standard deviation (n=7). Asterisks indicate statistically significant differences compared with *atcypDM* (*p*<0.01, Student's *t*-test). Scale bar, 3 cm. (C) Cytokinin concentrations in whole seedlings of Col-0, *atcypDM* and transgenic plants constitutively expressing *CYP735A3* (*A3-*ox) or *CYP735A4* (*A4-*ox) in *atcypDM* background. Seedlings were grown on 1/2x MS agar plates for 12 days before harvest. Error bars represent standard deviation (n=4 independent pools of more than ten plants). Asterisks indicate statistically significant differences in tZ-type cytokinin concentration compared with *atcypDM* (*p*<0.01, Student's *t*-test). The concentration of each cytokinin molecular species in roots and shoots is shown in Table S1. gFW, gram fresh weight; tZ-type, *trans*-zeatin and its conjugates; cZ-type, *cis*-zeatin and its conjugates; iP-type, N^6-(Δ^2-isopentenyl)adenine and its conjugates.

Figure 2. Expression pattern of *CYP735A3* and *CYP735A4* in various tissues
(A) Expression of *CYP735A3* and *CYP735A4* in Nipponbare seedlings. Coleoptiles and seminal roots, and roots and shoots were harvested 4 days and 14 days, respectively, after imbibition. Error bars represent standard deviation of biological replicates (n=4-5 independent pools of more than 2 plants). (B) Expression of *CYP735A3* and *CYP735A4* in Nipponbare plants in vegetative or early reproductive stages. 42-day-old (vegetative stage) and older plants (reproductive stage) were used. The shoot apex parts of 42-day-old plants were collected as shoot apical meristem (SAM). Leaf blade, leaf sheath and lamina joint were harvested from the youngest fully expanded leaf. The young panicle is a mixture of 2 mm to 21 mm long young panicles collected from plants at the reproductive stage. Error bars represent standard deviation of biological replicates (n=4 except for SAM and young panicle, n=4 and n=7 independent pools of more than 2 plants for SAM and young panicle, respectively). N.D., under the quantification detection limit. (C-E) Representative images of GUS staining obtained from *proCYP735A3:GUS* transgenic seedlings grown for 8 days after imbibition. Pictures of a whole seedling (C), first leaf (D, upper), crown root (D, lower), cross section of the first leaf (E, upper), and cross section of seminal root (E, lower) are presented. VBS, vascular bundle sheath; XY, xylem; PH, phloem; EN, endodermis; PE, pericycle. Scale bars, (C) 1 cm; (D) 200 μm; (E) 50 μm.

Figure 3. The *cyp735a3 cyp735a4* double mutants show deficiency in *trans*-zeatin-type cytokinins
Cytokinin concentrations in roots and shoots of WT strain Nipponbare (NB), vector control (VC), and mutant strains *cyp735a3-1* (*a3-1*), *cyp735a4-2* (*a4-2*), *cyp735a3-1 cyp735a4-1* (*a3a4-1*), and *cyp735a3-2 cyp735a4-2* (*a3a4-2*). Seedlings were grown hydroponically for 14 days (A) or 12 days (B) before shoots and roots were harvested. Error bars represent standard deviation of biological replicates (n=3-6 independent pools of more than ten plants for A, n=6-8 independent pools of more than ten plants for B). Asterisks indicate statistically significant differences in tZ-type cytokinin concentration compared with NB ($p < 0.05$, Student's *t*-test). The concentration of each cytokinin molecular species in roots and shoots is shown in Table S2 and Table S3. gFW, gram fresh weight; tZ-type, *trans*-zeatin and its conjugates; cZ-type, *cis*-zeatin and its conjugates; iP-type, N^6-(Δ^2-isopentenyl)adenine and its conjugates.

Figure 4. Disruption of *CYP735A*s alters rice growth in seedling to vegetative phases

(A) A representative image of Nipponbare (NB), vector control (VC), and *cyp735a3-2 cyp735a4-2* (*a3a4-2*) seedlings grown 12 days in hydroponic culture. Scale bar, 5 cm. (B) Quantification of shoot length, shoot fresh weight, root length, and root fresh weight in 12-day-old seedlings. NB, VC, *cyp735a4-2* (*a4-2*), and *a3a4-2* seedlings were grown in hydroponic culture. (C) Leaf number at 15 days after germination (D, E) Leaf sheath and leaf blade length (D) and leaf width (E) of the fully expanded third leaf of NB, VC, *cyp735a3-1* (*a3-1*), *cyp735a4-2* (*a4-2*), *cyp735a3-1 cyp735a4-1* (*a3a4-1*), and *a3a4-2* seedlings grown on soil. (F) Quantification of leaf blade length, leaf sheath length, and leaf width of the fully expanded ninth leaf in NB, VC, *a3a4-1*, and *a3a4-2* grown on soil. (G) Tiller number of NB, VC, *a3a4-1*, and *a3a4-2* grown on soil for 35 days. Error bars represent standard deviation of biological replicates (B, n=6-8; C, n=8-12; D, n=8-10; E, n=8-10; F, n=8-12; G, n=4-8). Different lowercase letters indicate statistically significant differences as indicated by Tukey's HSD test (*p*<0.05).

Figure 5. Disruption of *CYP735A*s alters rice growth in reproductive to maturatation phases
(A) A representative image of Nipponbare (NB), vector control (VC), and *cyp735a3-1* (*a3-1*), *cyp735a4-1* (*a4-2*), and *cyp735a3-1 cyp735a4-1* (*a3a4-2*) plants grown 15 weeks on soil. Scale bar, 20 cm. (B-D) Heading date (B), panicle and culm length (C), and tiller number (D) of NB, VC, *a3-1*, *a4-2*, *cyp735a3-1 cyp735a4-1* (*a3a4-1*), and *a3a4-2*. (E) A representative picture of mature panicles from NB and *a3a4-2*. Scale bar, 5 cm. (F, G) Panicle weight on the main stem (F) and number of primary branches per panicle (G). Error bars represent standard deviation of biological replicates (B, n=3-12; C, n=5-12; D, n=9-12; F, n=5-10; G, n=5-10). Different lowercase letters indicate statistically significant differences as indicated by Tukey's HSD test (*p*<0.05).

Figure 6. Expression of cytokinin inducible type-A response regulator genes is altered in the _cyp735a3 cyp735a4_ mutant

(A, B) Transcript levels of cytokinin inducible type-A response regulator genes, _OsRR1_, _OsRR2_, _OsRR4_, _OsRR6_, and _OsRR9/10_, in shoots (A) and roots (B) of Nipponbare (NB), vector control (VC), _cyp735a3-1 cyp735a4-1_ (_a3a4-1_) and _cyp735a3-2 cyp735a4-2_ (_a3a4-2_) seedlings. _OsRR9/10_ indicates that the primer set used could not distinguish _OsRR9_ and _OsRR10_. (C, D) Transcript levels of type-B response regulator genes, _OsRR21_ and _OsRR23_, in the shoot. (C) and in the root (D) of NB, VC, _a3a4-1_ and _a3a4-2_ seedlings. Seedlings were grown hydroponically for 10 days before roots and shoots were harvested. Expression levels were quantified by qRT-PCR analysis, normalized to _ZIURP1_ as an internal control, and are shown relative to the value of NB. Error bars represent standard deviation of biological replicates (n=4 independent pools of 2 plants). Asterisks indicate statistically significant differences compared to NB ($p<0.05$, Student's t-test).

Figure 7. Expression of *CYP735A3* and *CYP735A4* in response to phytohormones
(A) Expression of *CYP735A*s in response to cytokinin. Nipponbare (NB) seedlings grown for 12 days were treated with 0.001% DMSO (Mock), 1 μM iP (iP), or 1 μM tZ (tZ) and roots were harvested after 2 h. (B) Expression of *CYP735A*s in response to abscisic acid (ABA) and auxin. NB seedlings grown for 12 days were treated with 0.05% DMSO (Mock), 10 μM indole-3-acetic acid (IAA) or 50 μM ABA and roots were harvested after 2 h. Expression levels were quantified by qRT-PCR analysis, normalized to *ZIURP1* as an internal control, and are shown relative to the value in mock treatment. Asterisks indicate statistically significant differences compared with Mock (p<0.05, Student's t-test). Error bars represent standard deviation (n=4 independent pools of more than 2 plants).

Figure 8. Expression of *CYP735A3* and *CYP735A4* in response to nitrogen sources in roots
(A) Expression of *CYP735A*s in response to ammonium or nitrate supplement. Twelve day-old NB seedlings hydroponically grown in nitrogen-free nutrient solution were incubated with 5 mM KCl (-N), 5 mM NH₄Cl (NH₄⁺), or 5 mM KNO₃ (NO₃⁻) for 2 h or 4 h. Expression levels were quantified by qRT-PCR analysis, normalized to *ZIURP1* as an internal control, and are shown relative to the value in 2 h of -N treatment. Error bars represent standard deviation (n=4 independent pools of 3 plants). (B) Effect of methionine sulfoximine on the induction of *CYP735A*s by nitrogen. Twelve day-old NB seedlings hydroponically grown in nitrogen-free nutrient solution were pre-treated with 2 mM MSX for 2 h and then incubated with 5 mM KCl, 5 mM KNO₃ or 50 mM glutamine (GLN) for 2 h. Asterisks indicate statistically significant differences compared with KCl or KCl+MSX ((p<0.05, Student's t-test)). Expression levels were quantified by qRT-PCR analysis, normalized to *ZIURP1* as an internal control, and are shown relative to the value in KCl treatment. Error bars represent standard deviation (n=4 independent pools of 3 plants).

Parsed Citations

Bassil N V, Mok D, Mok MC (1993) Partial purification of a cis-trans-isomerase of zeatin from immature seed of Phaseolus vulgaris L. Plant Physiol 102: 867–872
  Google Scholar: Author Only Title Only Author and Title

Burr CA, Sun J, Yamburenko M V., Willoughby A, Hodgens C, Boeshore SL, Elmore A, Atkinson J, Nimchuk ZL, Bishopp A, et al (2021) The HK5 and HK6 cytokinin receptors mediate diverse developmental pathways in rice. Development 147: dev191734
  Google Scholar: Author Only Title Only Author and Title

Cai L, Zhang L, Fu Q, Xu ZF (2018) Identification and expression analysis of cytokinin metabolic genes IPTs, CYP735A and CKXs in the biofuel plant Jatropha curcas. PeerJ 2018: e4812
  Google Scholar: Author Only Title Only Author and Title

Chanch T, Ohira K (1982) Comparison of the uptake and assimilation of ammonium and nitrate in indica and japonica rice plants using the tracer 15N method. Soil Sci Plant Nutr 28: 79–90
  Google Scholar: Author Only Title Only Author and Title

Choi J, Lee J, Kim K, Cho M, Ryu H, An G, Hwang I (2012) Functional identification of OsHk6 as a homotypic cytokinin receptor in rice with preferential affinity for iP. Plant Cell Physiol 53: 1334–1343
  Google Scholar: Author Only Title Only Author and Title

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J 16: 735–743
  Google Scholar: Author Only Title Only Author and Title

Coskun D, Britto DT, Shi W, Kronzucker HJ (2017) Nitrogen transformations in modern agriculture and the role of biological nitrification inhibition. Nat Plants 3: 17074
  Google Scholar: Author Only Title Only Author and Title

Du H, Wu N, Fu J, Wang S, Li X, Xiao J, Xiong L (2012) A GH3 family member, OsGH3-2, modulates auxin and abscisic acid levels and differentially affects drought and cold tolerance in rice. J Exp Bot 63: 6467–6480
  Google Scholar: Author Only Title Only Author and Title

Engelsberger WR, Schulze WX (2012) Nitrate and ammonium lead to distinct global dynamic phosphorylation patterns when resupplied to nitrogen-starved Arabidopsis seedlings. Plant J 69: 978–995
  Google Scholar: Author Only Title Only Author and Title

Gajdošová S, Spíchal L, Kamínek M, Hoyerová K, Novák O, Dobrev PI, Galuszka P, Klíma P, Gaudinová A, Žižková E, et al (2011) Distribution, biological activities, metabolism, and the conceivable function of cis-zeatin-type cytokinins in plants. J Exp Bot 62: 2827–2840
  Google Scholar: Author Only Title Only Author and Title

Hansen CC, Nelson DR, Møller BL, Werck-Reichhart D (2021) Plant cytochrome P450 plasticity and evolution. Mol Plant 14: 1244–1265
  Google Scholar: Author Only Title Only Author and Title

Hirose N, Makita N, Kojima M, Kamada-Nobusada T, Sakakibara H (2007) Overexpression of a type-A response regulator alters rice morphology and cytokinin metabolism. Plant Cell Physiol 48: 523–539
  Google Scholar: Author Only Title Only Author and Title

Hirose N, Makita N, Yamaya T, Sakakibara H (2005) Functional characterization and expression analysis of a gene, OsENT2, encoding an equilibrative nucleoside transporter in rice suggest a function in cytokinin transport. Plant Physiol 138: 196–206
  Google Scholar: Author Only Title Only Author and Title

Hirose N, Takei K, Kuroha T, Kamada-Nobusada T, Hayashi H, Sakakibara H (2008) Regulation of cytokinin biosynthesis, compartmentalization and translocation. J Exp Bot 59: 75–83
  Google Scholar: Author Only Title Only Author and Title

Kakimoto T (2001) Identification of plant cytokinin biosynthetic enzymes as dimethylallyl diphosphate : ATP/ADP isopentenyltransferases. Plant Cell Physiol 42: 677–685
  Google Scholar: Author Only Title Only Author and Title

Kamada-Nobusada T, Makita N, Kojima M, Sakakibara H (2013) Nitrogen-dependent regulation of de novo cytokinin biosynthesis in rice: the role of glutamine metabolism as an additional signal. Plant Cell Physiol 54: 1881–1893
  Google Scholar: Author Only Title Only Author and Title

Kawai M, Tabata R, Ohashi M, Honda H, Kamiya T, Kojima M, Takebayashi Y, Oishi S, Okamoto S, Hachiya T, et al (2022) Regulation of ammonium acquisition and use in Oryza longistaminata ramets under nitrogen source heterogeneity. Plant Physiol 188: 2364–2376

Google Scholar: Author Only Title Only Author and Title

Kiba T, Inaba J, Kudo T, Ueda N, Konishi M, Mitsuda N, Takiguchi Y, Kondou Y, Yoshizumi T, Ohme-Takagi M, et al (2018) Repression of nitrogen-starvation responses by members of the Arabidopsis GARP-type transcription factor NIGT1/HRS1 subfamily. Plant Cell 30: 925–945
Google Scholar: Author Only Title Only Author and Title

Kiba T, Kudo T, Kojima M, Sakakibara H (2011) Hormonal control of nitrogen acquisition: roles of auxin, abscisic acid, and cytokinin. J Exp Bot 62: 1399–1409
Google Scholar: Author Only Title Only Author and Title

Kiba T, Takebayashi Y, Kojima M, Sakakibara H (2019) Sugar-induced de novo cytokinin biosynthesis contributes to Arabidopsis growth under elevated CO2. Sci Rep 9: 1–15
Google Scholar: Author Only Title Only Author and Title

Kiba T, Takei K, Kojima M, Sakakibara H (2013) Side-chain modification of cytokinins controls shoot growth in Arabidopsis. Dev Cell 27: 452–461
Google Scholar: Author Only Title Only Author and Title

Kieber JJ, Schaller GE (2018) Cytokinin signaling in plant development. Development 145: dev149344
Google Scholar: Author Only Title Only Author and Title

Kirk GJ, Kronzucker HJ (2005) The potential for nitrification and nitrate uptake in the rhizosphere of wetland plants: a modelling study. Ann Bot 96: 639–646
Google Scholar: Author Only Title Only Author and Title

Kojima M, Kamada-Nobusada T, Komatsu H, Takei K, Kuroha T, Mizutani M, Ashikari M, Ueguchi-Tanaka M, Matsuoka M, Suzuki K, et al (2009) Highly sensitive and high-throughput analysis of plant hormones using MS-probe modification and liquid chromatography tandem mass spectrometry: an application for hormone profiling in Oryza sativa. Plant Cell Physiol 50: 1201–1214
Google Scholar: Author Only Title Only Author and Title

Kojima M, Sakakibara H (2012) Highly sensitive high-throughput profiling of six phytohormones using MS-probe modification and liquid chromatography-tandem mass spectrometry. Methods Mol Biol 918: 151–164
Google Scholar: Author Only Title Only Author and Title

Kudo T, Makita N, Kojima M, Tokunaga H, Sakakibara H (2012) Cytokinin activity of cis-zeatin and phenotypic alterations induced by overexpression of putative cis-Zeatin-O-glucosyltransferase in rice. Plant Physiol 160: 319–331
Google Scholar: Author Only Title Only Author and Title

Kurakawa T, Ueda N, Maekawa M, Kobayashi K, Kojima M, Nagato Y, Sakakibara H, Kyozuka J (2007) Direct control of shoot meristem activity by a cytokinin-activating enzyme. Nature 445: 652–655
Google Scholar: Author Only Title Only Author and Title

Kuroha T, Tokunaga H, Kojima M, Ueda N, Ishida T, Nagawa S, Fukuda H, Sugimoto K, Sakakibara H (2009) Functional analyses of LONELY GUY cytokinin-activating enzymes reveal the importance of the direct activation pathway in Arabidopsis. Plant Cell 21: 3152–3169
Google Scholar: Author Only Title Only Author and Title

Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E (2019) CHOPCHOP v3: Expanding the CRISPR web toolbox beyond genome editing. Nucleic Acids Res 47: W171–W174
Google Scholar: Author Only Title Only Author and Title

Lam HM, Coschigano KT, Oliveira IC, Melo-Oliveira R, Coruzzi GM (1996) The molecular-genetics of nitrogen assimilation into amino acids in higher plants. Annu Rev Plant Physiol Plant Mol Biol 47: 569–593
Google Scholar: Author Only Title Only Author and Title

Li YS, Sun H, Wang ZF, Duan M, Huang SD, Yang J, Huang J, Zhang HS (2013) A novel nuclear protein phosphatase 2C negatively regulated by ABL1 is involved in abiotic stress and panicle development in rice. Mol Biotechnol 54: 703–710
Google Scholar: Author Only Title Only Author and Title

Lomin SN, Yonekura-Sakakibara K, Romanov GA, Sakakibara H (2011) Ligand-binding properties and subcellular localization of maize cytokinin receptors. J Exp Bot 62: 5149–5159
Google Scholar: Author Only Title Only Author and Title

Macková H, Hronková M, Dobrá J, Turečková V, Novák O, Lubovská Z, Motyka V, Haisel D, Hájek T, Prášil IT, et al (2013) Enhanced drought and heat stress tolerance of tobacco plants with ectopically enhanced cytokinin oxidase/dehydrogenase gene expression. J Exp Bot 64: 2805–2815
Google Scholar: Author Only Title Only Author and Title

Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y, Sakakibara H, Yanagisawa S (2018) A NIGT1-centred transcriptional cascade regulates nitrate signalling and incorporates phosphorus starvation signals in Arabidopsis. Nat Commun

9: 1–14

Google Scholar: Author Only Title Only Author and Title

Makino A, Mae T, Ohira K (1985) Photosynthesis and ribulose-1,5-bisphosphate carboxylase oxygenase in rice leaves from emergence through senescence - quantitative-analysis by carboxylation oxygenation and regeneration of ribulose 1,5-bisphosphate. Planta 166: 414–420

Google Scholar: Author Only Title Only Author and Title

Midlik A, Navrátilová V, Moturu TR, Koča J, Svobodová R, Berka K (2021) Uncovering of cytochrome P450 anatomy by SecStrAnnotator. Sci Rep 11: 12345

Google Scholar: Author Only Title Only Author and Title

Miyawaki K, Matsumoto-Kitano M, Kakimoto T (2004) Expression of cytokinin biosynthetic isopentenyltransferase genes in Arabidopsis: tissue specificity and regulation by auxin, cytokinin, and nitrate. Plant J 37: 128–138

Google Scholar: Author Only Title Only Author and Title

Miyawaki K, Tarkowski P, Matsumoto-Kitano M, Kato T, Sato S, Tarkowska D, Tabata S, Sandberg G, Kakimoto T (2006) Roles of Arabidopsis ATP/ADP isopentenyltransferases and tRNA isopentenyltransferases in cytokinin biosynthesis. Proc Natl Acad Sci USA 103: 16598–16603

Google Scholar: Author Only Title Only Author and Title

Mok DW, Mok MC (2001) Cytokinin metabolism and action. Annu Rev Plant Physiol Plant Mol Biol 52: 89–118

Google Scholar: Author Only Title Only Author and Title

Mok MC, Mok DW, Armstrong DJ (1978) Differential cytokinin structure-activity relationships in Phaseolus. Plant Physiol 61: 72–75

Google Scholar: Author Only Title Only Author and Title

Nacry P, Bouguyon E, Gojon A (2013) Nitrogen acquisition by roots: physiological and developmental mechanisms ensuring plant adaptation to a fluctuating resource. Plant Soil 370: 1–29

Google Scholar: Author Only Title Only Author and Title

Ohashi M, Ishiyama K, Kojima S, Kojima M, Sakakibara H, Yamaya T, Hayakawa T (2017) Lack of cytosolic glutamine synthetase1;2 activity reduces nitrogen-dependent biosynthesis of cytokinin required for axillary bud outgrowth in rice seedlings. Plant Cell Physiol 58: 679–690

Google Scholar: Author Only Title Only Author and Title

Osugi A, Kojima M, Takebayashi Y, Ueda N, Kiba T, Sakakibara H (2017) Systemic transport of trans-zeatin and its precursor have differing roles in Arabidopsis shoots. Nat Plants 3: 1–6

Google Scholar: Author Only Title Only Author and Title

Osugi A, Sakakibara H (2015) Q&A: How do plants respond to cytokinins and what is their importance? BMC Biol 13: 102

Pabuayon IM, Yamamoto N, Trinidad JL, Longkumer T, Raorane ML, Kohli A (2016) Reference genes for accurate gene expression analyses across different tissues, developmental stages and genotypes in rice for drought tolerance. Rice 9: 32

Google Scholar: Author Only Title Only Author and Title

Romanov GA, Lomin SN, Schmülling T (2006) Biochemical characteristics and ligand-binding properties of Arabidopsis cytokinin receptor AHK3 compared to CRE1/AHK4 as revealed by a direct binding assay. J Exp Bot 57: 4051–4058

Google Scholar: Author Only Title Only Author and Title

Sakakibara H (2006) Cytokinins: activity, biosynthesis, and translocation. Annu Rev Plant Biol 57: 431–449

Google Scholar: Author Only Title Only Author and Title

Sakakibara H (2021) Cytokinin biosynthesis and transport for systemic nitrogen signaling. Plant J 105: 421–430

Google Scholar: Author Only Title Only Author and Title

Schäfer M, Brütting C, Meza-Canales ID, Großkinsky DK, Vankova R, Baldwin IT, Meldau S (2015) The role of cis-zeatin-type cytokinins in plant growth regulation and mediating responses to environmental interactions. J Exp Bot. Oxford University Press, pp 4873–4884

Google Scholar: Author Only Title Only Author and Title

Schmitz RY, Skoog F (1972) Cytokinins: Synthesis and biological activity of geometric and position isomers of zeatin. Plant Physiol 50: 702–705

Google Scholar: Author Only Title Only Author and Title

Shaw G (1994) Chemistry of adenine cytokinins. In DWS Mok, MC Mok, eds, Cytokinins: Chemistry, Activity, and Function. CRC Press, Boca Rato, FL, pp 15–34

Google Scholar: Author Only Title Only Author and Title

Shibasaki K, Takebayashi A, Makita N, Kojima M, Takebayashi Y, Kawai M, Hachiya T, Sakakibara H (2021) Nitrogen nutrition promotes rhizome bud outgrowth via regulation of cytokinin biosynthesis genes and an Oryza longistaminata ortholog of FINE

CULM 1. Front Plant Sci 12: 788

Google Scholar: Author Only Title Only Author and Title

Spíchal L, Rakova NY, Riefler M, Mizuno T, Romanov GA, Strnad M, Schmülling T (2004) Two cytokinin receptors of Arabidopsis thaliana, CRE1/AHK4 and AHK3, differ in their ligand specificity in a bacterial assay. Plant Cell Physiol 45: 1299–1305

Google Scholar: Author Only Title Only Author and Title

Stolz A, Riefler M, Lomin SN, Achazi K, Romanov GA, Schmülling T (2011) The specificity of cytokinin signalling in Arabidopsis thaliana is mediated by differing ligand affinities and expression profiles of the receptors. Plant J 67: 157–168

Google Scholar: Author Only Title Only Author and Title

Takei K, Sakakibara H, Sugiyama T (2001a) Identification of genes encoding adenylate isopentenyltransferase, a cytokinin biosynthesis enzyme, in Arabidopsis thaliana. J Biol Chem 276: 26405–26410

Google Scholar: Author Only Title Only Author and Title

Takei K, Sakakibara H, Taniguchi M, Sugiyama T (2001b) Nitrogen-dependent accumulation of cytokinins in root and the translocation to leaf: implication of cytokinin species that induces gene expression of maize response regulator. Plant Cell Physiol 42: 85–93

Google Scholar: Author Only Title Only Author and Title

Takei K, Yamaya T, Sakakibara H (2004) Arabidopsis CYP735A1 and CYP735A2 encode cytokinin hydroxylases that catalyze the biosynthesis of trans-zeatin. J Biol Chem 279: 41866–41872

Google Scholar: Author Only Title Only Author and Title

Toda E, Koiso N, Takebayashi A, Ichikawa M, Kiba T, Osakabe K, Osakabe Y, Sakakibara H, Kato N, Okamoto T (2019) An efficient DNA- and selectable-marker-free genome-editing system using zygotes in rice. Nat Plants 5: 363–368

Google Scholar: Author Only Title Only Author and Title

Tokunaga H, Kojima M, Kuroha T, Ishida T, Sugimoto K, Kiba T, Sakakibara H (2012) Arabidopsis lonely guy (LOG) multiple mutants reveal a central role of the LOG-dependent pathway in cytokinin activation. Plant J 69: 355–365

Google Scholar: Author Only Title Only Author and Title

Tsai Y-C, Weir NR, Hill K, Zhang W, Kim HJ, Shiu S-H, Schaller GE, Kieber JJ (2012) Characterization of Genes Involved in Cytokinin Signaling and Metabolism from Rice. Plant Physiol 158: 1666–1684

Google Scholar: Author Only Title Only Author and Title

Vanková R, Kosová K, Dobrev P, Vítámvás P, Trávníčková A, Cvikrová M, Pešek B, Gaudinová A, Prerostová S, Musilová J, et al (2014) Dynamics of cold acclimation and complex phytohormone responses in Triticum monococcum lines G3116 and DV92 differing in vernalization and frost tolerance level. Environ Exp Bot 101: 12–25

Google Scholar: Author Only Title Only Author and Title

Vyroubalová S, Václavíková K, Turečková V, Novák O, Šmehilová M, Hluska T, Ohnoutková L, Frébort I, Galuszka P (2009) Characterization of new maize genes putatively involved in cytokinin metabolism and their expression during osmotic stress in relation to cytokinin levels. Plant Physiol 151: 433–447

Google Scholar: Author Only Title Only Author and Title

Werner T, Motyka V, Laucou V, Smets R, Van Onckelen H, Schmülling T (2003) Cytokinin-deficient transgenic Arabidopsis plants show multiple developmental alterations indicating opposite functions of cytokinins in the regulation of shoot and root meristem activity. Plant Cell 15: 2532–2550

Google Scholar: Author Only Title Only Author and Title

Werner T, Schmülling T (2009) Cytokinin action in plant development. Curr Opin Plant Biol 12: 527–538

Google Scholar: Author Only Title Only Author and Title

Worthen JM, Yamburenko MV, Lim J, Nimchuk ZL, Kieber JJ, Schaller GE (2019) Type-B response regulators of rice play key roles in growth, development and cytokinin signaling. Development 146: dev174870

Google Scholar: Author Only Title Only Author and Title

Xu GH, Fan XR, Miller AJ (2012) Plant nitrogen assimilation and use efficiency. Annu Rev Plant Biol 63: 153–182

Google Scholar: Author Only Title Only Author and Title

Yan M, Fan XR, Feng HM, Miller AJ, Shen QR, Xu GH (2011) Rice OsNAR2.1 interacts with OsNRT2.1, OsNRT2.2 and OsNRT2.3a nitrate transporters to provide uptake over high and low concentration ranges. Plant Cell Environ 34: 1360–1372

Google Scholar: Author Only Title Only Author and Title

Yonekura-Sakakibara K, Kojima M, Yamaya T, Sakakibara H (2004) Molecular characterization of cytokinin-responsive histidine kinases in maize. differential ligand preferences and ¥response to cis-zeatin. Plant Physiol 134: 1654–1661

Google Scholar: Author Only Title Only Author and Title

Zhao J, Yu N, Ju M, Fan B, Zhang Y, Zhu E, Zhang M, Zhang K (2019) ABC transporter OsABCG18 controls the shootward

transport of cytokinins and grain yield in rice. J Exp Bot 70: 6277–6291
  Google Scholar: Author Only Title Only Author and Title

Zwack PJ, Rashotte AM (2015) Interactions between cytokinin signalling and abiotic stress responses. J Exp Bot 66: 4863–4871
  Google Scholar: Author Only Title Only Author and Title