1 Recombination rate variation shapes genomic variability of phylogeographic structure in a
2 widespread North American songbird (Aves: *Certhia americana*)
3

4 Joseph D. Manthey[1]*, Garth M. Spellman[2]
5
6 [1]Department of Biological Sciences, Texas Tech University. Lubbock, Texas, USA.
7 [2]Department of Zoology, Denver Museum of Nature & Science, Denver, Colorado, USA.
8 *Corresponding author email: jdmanthey@gmail.com

9
10 **ABSTRACT**
11

The nonrandom distribution of chromosomal characteristics and functional elements—genomic
architecture—impacts the relative strengths and impacts of population genetic processes across
the genome. Due to this relationship, genomic architecture has the potential to shape variation
in population genetic structure across the genome. Population genetic structure has been
shown to vary across the genome in a variety of taxa, but this body of work has largely focused
on pairwise population genomic comparisons between closely related taxa. Here, we used
whole genome sequencing of seven populations of a North American songbird, the Brown
Creeper (*Certhia americana*), to determine the impacts of genomic architecture on regional
genomic variability in phylogeographic structure. Using multiple methods to infer
phylogeographic structure—summary statistics, clustering methods, and phylogenomics—we
found that recombination rate variation explained a large proportion of phylogeographic
structure variation. Genomic regions with low recombination showed phylogeographic structure
consistent with the genome-wide pattern. In regions with high recombination, we found strong
phylogeographic structure, but with discordant patterns relative to the genome-wide pattern. In
regions with high recombination rate, we found that populations with small effective population
sizes evolve more rapidly than larger populations, leading to discordant signatures of
phylogeographic structure. These results suggest that the interplay between recombination rate
variation and effective population sizes shape the relative impacts of linked selection and
genetic drift in different parts of the genome. Overall, the combined interactions of population
genetic processes, genomic architecture, and effective population sizes shapes patterns of
variability in phylogeographic structure across the genome.

40 **Manuscript Type:** Research Article
41
42 **Running head:** Recombination rate variation shapes local genomic phylogeographic structure
43
44

INTRODUCTION

The formation of population structure is implicit in the speciation process (Allmon 1992; Mayr, et al. 1963). Speciation is a continuum; early differentiating populations may have few genomic regions with evidence for genetic differentiation or barriers to gene flow (e.g., Toews, et al. 2016) while later in the speciation process two diverging populations may exhibit strong genetic differentiation across their entire genomes (e.g., Ravinet, et al. 2018). Across the genome, variation in local evolutionary history (i.e., population genetic structure) forms a complex relationship with the relative strengths of the population genomic processes of natural selection, gene flow, and genetic drift. For example, population genetic structure may differ in genomic regions with strong natural selection or differential patterns of gene flow relative to the rest of the genome (Louis, et al. 2020; Whiting, et al. 2021).

Variable impacts of selection, drift, and gene flow across the genome may be partially attributed to genomic architecture—the nonrandom distribution of chromosomal characteristics and functional elements (Koonin 2009)—whereby variation in gene content, recombination rates, or other genomic characteristics impact the relative effects of different population genomic processes. For example, we expect that genetic differentiation will accumulate faster in genomic regions with relatively lower recombination rates due to increased effects of linked selection; the effects of selection may also be positively correlated with gene density (Cruickshank and Hahn 2014; Haenel, et al. 2018; Hey and Kliman 2002; Nachman and Payseur 2012). Indeed, speciation and population genomic studies have often found a negative relationship between recombination rates and genetic differentiation across the genome (Kulathinal, et al. 2008; Roesti, et al. 2013; Stankowski, et al. 2019; Vijay, et al. 2016).

Genetic clustering and phylogenetic methods allow the investigation of population genetic structure across many sampled populations (i.e., not just population pairwise comparisons). Variation in phylogenetic signal across the genome may provide evidence for differential lineage sorting or gene flow among lineages (e.g., Hobolth, et al. 2011; White, et al. 2009). Similarly, genetic principal components analyses (PCA) across the genome may exhibit shifts in local evolutionary history due to the effects of recombination rate variation or gene density on population genomic processes leading to genetic structure (Li and Ralph 2019). As such, we may expect that variation in genomic architecture impacts regional genomic patterns of genetic clustering and phylogenetic signal in taxonomic groups with lineages at different stages of the speciation continuum. Exemplar taxa to better understand the impacts of genomic architecture on population genetic structure are species (or species complexes) with (1) multiple

lineages on distinct evolutionary trajectories and (2) genomes exhibiting heterogeneous

genomic architecture.

The Brown Creeper (*Certhia americana*) is an excellent study organism to understand

the impacts of genomic architecture on phylogeographic structure; its genome is highly

heterogeneous in both structure and content, and it has seven distinct phylogeographic clades.

The genome of the Brown Creeper, as like other birds, has chromosomes that (1) span two

orders of magnitude in size, (2) vary in their gene and repetitive element content, and (3) exhibit

local recombination rates that can vary by an order of magnitude or more (Dutoit, et al. 2017a;

Ellegren 2010; Kapusta and Suh 2017; Kawakami, et al. 2014; Manthey, et al. 2021).

Additionally, the Brown Creeper has strong phylogeographic structure, with two main lineages

that have been diverging around one million years (Fig. 1A), and multiple phylogeographically

structured clades within each of these two main lineages (Manthey, et al. 2015, 2011a, 2021,

2011b).

Using the Brown Creeper as our study taxon, we used phylogenetic methods, multiple

population genetic clustering methods, and population genetic summary statistics to

characterize phylogeographic structure across the genome. Our goals were threefold. First, we

sought to identify how the nonrandom distributions of coding content, repetitive content, and

variation in recombination rate shape the regional genomic signal of population genetic

structure. Second, we characterized whether effective population size interacts with genomic

architecture in shaping genomic patterns of population differentiation. Lastly, when we observed

population genetic structure that varies across the genome due to genomic architecture and

interactions with effective population size, we discuss the implications for phylogeographic

studies more broadly, such as those that use subgenomic datasets.

METHODS

Study organism, sampling, and sequencing. The Brown Creeper is a songbird found in

forested regions of the Americas; it has a widespread distribution ranging from Honduras in the

south to Alaska in the north (Fig. 1A) (Poulin, et al. 2020). The Brown Creeper is aptly named; it

creeps up trees in search of its largely invertebrate diet and has dark coloration on its back and

sides (Fig. 1A inset) that makes for excellent camouflage on trees with dark bark. In previous

work using subgenomic datasets, we have identified phylogeographic structure in the Brown

Creeper, with up to seven distinct lineages (Manthey, et al. 2015, 2011a, b). Here, we aimed to

obtain population genomic sequencing data for seven populations with three individuals

sampled per population (Fig 1, Table S1). We downloaded previously published sequencing

data from two of these populations and one outgroup sample of *C. familiaris* (Table S1)

(Manthey, et al. 2021). For the other five populations, we obtained representative tissue

samples from natural history museums and extracted genomic DNA using QIAGEN DNeasy

blood and tissue kits following manufacturer guidelines. DNA from extractions was used to

create standard Illumina sequencing libraries, followed by sequencing on an Illumina

NovaSeq6000 at the Texas Tech University Center for Biotechnology and Genomics with the

goal of obtaining ~15-30× coverage per individual.

Reference genome and genomic architecture. For the reference genome, we used the

chromosome-scale *Certhia americana* genome we assembled for a prior study (NCBI assembly

ASM1869719v1; Manthey et al. 2021). From data reported in this previous study (Manthey et al.

2021), we summarized mean recombination rates for each of the two main Brown Creeper

lineages, transposable element (TE) and gene density, and GC content across the genome in

50kbp and 100 kbp non-overlapping sliding windows (Fig. 2).

Sequencing data filtering and genotyping. We used the program bbduk (Bushnell 2014) to

quality filter raw sequencing data. We then use BWA v0.7.17 (Li and Durbin 2009) to align

filtered reads to the *Certhia americana* reference genome. Next, we used samtools v1.4.1 (Li, et

al. 2009) to convert the BWA output SAM file to BAM format, followed by using the Genome

Analysis Toolkit (GATK) v4.1.0.0 (McKenna, et al. 2010) to clean, sort, add read groups to, and

remove duplicates from BAM files. We used the samtools 'depth' command to measure

sequencing coverage across the reference genome for each individual (Fig. S1, Table S1).

We then used GATK in three steps to genotype all individuals: we used the

'HaplotypeCaller' function to call preliminary genotypes for each individual, built a database for

all samples using the 'GenomicsDBImport' function, and lastly used the 'GenotypeGVCFs'

function to group genotype all individuals together for both variant and invariant sites. We

filtered the genotype data using VCFtools v0.1.14 (Danecek, et al. 2011) with the following

restrictions: (1) minimum site quality of 20, (2) minimum genotype quality of 20, (3) minimum

depth of coverage of 6, (4) maximum mean depth of coverage of 50, and (5) removal of indels.

For downstream analyses, we used various settings for (1) amount of missing data allowed, (2)

whether the outgroup was included, (3) inclusion or exclusion of invariant sites, and (4) thinning

between sites (see Table 1 for exact dataset characteristics used for all seven datasets used for

analyses). Although it has been suggested that variation in minor allele frequency (MAF) filtering

may impact estimates of genetic structure (Linck and Battey 2019), we did not filter for a

minimum MAF because we previously showed that changing MAF did not significantly impact

pairwise F_{ST} calculations in *C. americana* (Manthey, et al. 2021).

Phylogenomics. We estimated "gene trees" in non-overlapping windows (window size 50 kbp

and 100 kbp; datasets C and D in Table 1) using RAxML v8.2.12 (Stamatakis 2014) with the

GTRCAT model of sequence evolution. For each phylogeny, we used 100 rapid bootstraps to

assess support and determined a best-supported tree for each window. For a window to be

included, we required a minimum of 10 kbp in the alignment following filtering. This resulted in

19,639 phylogenies for the 50 kbp sliding windows and 9846 phylogenies for the 100 kbp sliding

windows. We summarized the best-supported trees using the sumtrees.py script, part of the

DendroPy Python package (Sukumaran and Holder 2010). Lastly, we used ASTRAL III v 5.6.3

(Zhang, et al. 2018) to calculate a species tree from all gene trees, using quartet frequencies as

a measure of local support (Sayyari and Mirarab 2016).

For each gene tree, we estimated the genealogical sorting index (GSI) (Cummings, et al.

2008) for (1) each of the seven populations, and (2) each of the two main lineages (i.e., north

and south shaded blue and red, respectively, in Fig. 1). The GSI statistic uses rooted gene trees

to quantify lineage sorting; the GSI quantifies exclusive ancestry of individuals in labelled groups

by dividing the minimum number of nodes to produce monophyly for a defined group divided by

the observed number of nodes necessary to connect all individuals of a group in a phylogeny.

The maximum GSI value is one and occurs when a defined group is monophyletic. We

calculated the GSI statistic for each sampled population and for each of the two main lineages

(Fig. 1).

For each individual across every gene tree, we calculated the root-to-tip distance (RTD)

using the R package ape (Paradis, et al. 2004). As a summary of these RTD distances, we

calculated the mean RTD per population relative to the maximum RTD value for each gene tree

(i.e., this relative statistic's range = 0 – 1). We then measured the relationships between

harmonic mean effective population sizes (mean per population) and relative RTD in regions

with low recombination and high recombination, as well as the shift in values between high and

low recombination regions. We also estimated relationships between these characteristics while

accounting for the evolutionary relationships of the samples using phylogenetic independent

contrasts (PICs) calculated in the R package ape (Paradis, et al. 2004).

Genetic structure. We estimated genetic structure using the program ADMIXTURE v1.3.0

(Alexander, et al. 2009). Here, we used genome-wide SNPs thinned to either a minimum of 50

kbp or 100 kbp distance between SNPs (datasets A and B in Table 1). We ran ADMIXTURE

with an assumed number of genetic clusters equal to two or seven (i.e., k = 2 and k = 7) based

on our knowledge of phylogeographic structure in this species from previous studies (Manthey,

et al. 2015, 2011a, b). To assess how genetic structure estimation varied across the genome,

we also ran ADMIXTURE for SNP datasets in non-overlapping sliding windows of 50 kbp and

100 kbp (datasets E and F in Table 1). We compared the genome-wide ADMIXTURE results to

those in each window by summing the differences in group membership for each individual. In

other words, this deviation value ranges from zero to nearly one, where a value of zero is

identical group assignment for all individuals with the same admixture coefficients for each

genetic cluster, and higher values indicate more dissimilar matrices.

We also estimated variation in genetic structure across the genome using LOSTRUCT

(Li and Ralph 2019). This method runs in three steps: (1) reducing the dimensionality of the data

using principal component analysis (PCA) for each genomic window, (2) find distances between

PCA maps, and (3) using multidimensional scaling (MDS) to display variation in PCA-based

genetic structure across the genome. We used LOSTRUCT in sliding windows of 50 kbp and

100 kbp (datasets E and F in Table 1). We used variation in MDS axis one (hereafter MDS1) to

describe variation in these PCA-based estimates of genetic structure across the genome.

Summary statistics. Across 50 kbp and 100 kbp sliding windows (datasets C and D in Table

1), we estimated population genomic summary statistics to summarize patterns of genetic

variation across the genome both within and among populations. We measured observed

heterozygosity per individual and estimated nucleotide diversity per population as measures of

genetic diversity. We calculated genetic differentiation (F_{ST}) and genetic divergence (D_{XY})

between geographically neighboring populations as measures of distinct evolutionary

trajectories among populations. We used the Reich et al. (2009) F_{ST} estimator because it is

unbiased when using small sample sizes, unlike some other common F_{ST} estimators (Willing, et

al. 2012). Lastly, we measured Tajima's D (Tajima 1989) as an estimator of whether populations

tend to evolve neutrally across the genome.

Population genomic correlations. We used the R package Hmisc (Harrell and Dupont 2020)

to estimate both Pearson and Spearman correlation coefficients between population genomic

summary statistics and characteristics of genomic architecture in sliding windows. We used the

R package corrplot (Wei, et al. 2017) to visualize correlations between all summary statistics.

Lastly, we used variance partitioning, implemented in the R package vegan (Oksanen, et al.

2007), to assess the proportion of variance in phylogeographic structure estimates explained by

genomic architecture.

Population demographic history. We used the program MSMC2 v1.1.0 (Schiffels and Durbin

2014) to estimate demographic history for each individual. For use in MSMC, we masked

regions of the genome not genotyped due to low coverage or low genotype quality scores.

Additionally, we did not include the sex chromosomes in demography calculations. We ran

MSMC for each individual allowing up to 20 iterations (default setting) and used up to 23

inferred distinct time segments because this setting worked well to reduce spurious or

inconsistent results in other songbirds that we have studied (Manthey, et al. 2022). We

performed 10 bootstraps for each demography estimate, using 1 Mbp bootstrapped segments of

the genome, to assess how demographic signal varies when subsetting parts of the genome.

Because the output of MSMC is interpreted relative to assumed mutation rates and generation

times, we used the Brown Creeper genome mutation rate estimate of 2.506×10^{-9} substitutions

per site per year (Manthey, et al. 2021). Because there are no published estimates of Brown

Creeper generation times, we used a proxy generation time of double the age of sexual maturity

(Nadachowska-Brzyska, et al. 2015) using maturity estimates from the Animal Aging and

Longevity Database (age of maturity = one year) (Tacutu, et al. 2017). With the MSMC

demographic model output, we estimated the harmonic mean population sizes for each

population by using the mean MSMC population size estimates for each individual in 1000-year

discrete time intervals over the past 200 kya.

Potential mutational load. Because we identified different relative rates of molecular evolution

in differently sized populations, we wanted to identify whether smaller populations were

accumulating a relatively greater potential mutational load. First, we used SnpEff (Cingolani, et

al. 2012) to annotate estimated functional effects of SNPs in coding regions based on a

previous functional annotation of the *C. americana* reference genome (Manthey, et al. 2021).

Next, as a relative estimate of potential mutational load, we measured the proportion of

polymorphisms in each population that changed the amino acid sequence (i.e., missense and

nonsense mutations). Specifically, we calculated the number of nonsynonymous polymorphisms

(P_N) and synonymous polymorphisms (P_S) and measured the ratio $\frac{P_N}{P_N + P_S}$ for each population in

genomic quintiles of recombination rate variation.

Code availability. All computer code used for analyses and figure creation (applicable for some figures) for this project is available here: github.com/jdmanthey/certhia_phylogeography. We used the following R packages for figure creation and file manipulation that were not cited in other parts of the methods section: Biostrings (Pagès, et al. 2017), ggplot2 (Wickham 2011), palettetown (Lucas 2016), phytools (Revell 2012), and RColorBrewer (Neuwirth 2014).

RESULTS

Phylogeographic structure and demography. We used genomic sequencing data from 21 ingroup individuals sequenced at ~18-29× genomic coverage to estimate genome-wide patterns of phylogeographic structure. We corroborated previous subgenomic scale work by identifying hierarchical and strong phylogeographic structure among sampled populations (Manthey, et al. 2015, 2011a, b). The deepest phylogenomic split separates northern from southern populations and there is additional support for distinctiveness of each of the regionally sampled populations (Fig. 1C). Genome-wide ADMIXTURE results were consistent across different thinning strategies (i.e., 50kbp and 100 kbp thinning; only 50 kbp results plotted) and showed patterns of genetic structure that aligned well with the species tree (Fig. 1C). Demographic history estimates for each individual were consistent within populations, but each population exhibited a distinct demographic history (Fig. 1D). In the Pacific and Central American localities, population sizes have fluctuated somewhat, but generally stay below an N_E of 200,000 over the past 100 ky (Fig. 1D). The Sierra Madre Oriental population showed evidence for a sharp decline in N_E over the past 50 ky, shifting from an N_E ~ 200,000 at about 50 kya to an N_E ~ 10,000 in the past ten kya (Fig. 1D). The Rocky Mountains population has remained relatively stable over the past 100 ky with an N_E ~ 200,000 (Fig. 1D). Lastly, the Eastern North America and Central Mexico populations have exhibited fluctuations and much larger N_E than all other populations over the past 100 ky (Fig. 1D). Harmonic mean effective population sizes over the past 200 ky ranged from ~85,000 (Central America South) to ~320,000 (Central Mexico) (Fig. 1D). Harmonic mean effective population size estimates over the past 200 ky are highly correlated with observed heterozygosity estimates for each individual ($r = 0.962$).

Phylogeographic structure variation across the genome. Pairwise values of genetic divergence and differentiation between geographically proximate populations varied widely across the genome (Fig. 2; Fig. S2; Fig. S3; Fig. S4, Fig. S5). Estimates between the two main

lineages (Rocky Mountain × Central Mexico comparison) were the highest, as expected; for all

other comparisons, genetic divergence (D_{XY}) had pairwise mean values between 0.0025 and

0.0050, and genetic differentiation (F_{ST}) had pairwise mean values between 0.15 and 0.25 (Fig.

2; Fig. S4). Notably, only two pairwise comparisons of populations—Rocky Mountain × Central

Mexico and Rocky Mountain × Eastern North America—showed considerable variation in F_{ST}

across the genome, while the other pairwise comparisons showed relatively consistent

estimates of F_{ST} across the genome (Fig. S2). Notably, while there was a negative relationship

between F_{ST} and D_{XY} when using the mean values across all pairwise comparisons, this

relationship also varied by pairwise population comparison (Fig. S5); the Rocky Mountain ×

Central Mexico showed a strong positive correlation between F_{ST} and D_{XY}, the Rocky Mountain

× Eastern North America showed a strong negative correlation, and the other pairwise

comparisons showed a weaker relationship between the two summary statistics (Fig. S5).

Genetic diversity in each population varied widely across the genome, with mean

nucleotide diversity values for each population between ~0.002 and 0.004 (Fig 1B; Fig. S4; Fig.

S6). Estimates of Tajima's D for each population appeared to be somewhat consistent across

the genome in three populations (Central Mexico, Eastern North America, and Rocky Mountain),

but highly variable across the genome in the other four populations (Fig. S7).

We also estimated phylogeographic structure across the genome using PCA,

ADMIXTURE, and the GSI metric from phylogenies; all estimates of phylogeographic structure

varied across the genome, with the outliers deviating the most from genome-wide patterns

clustering mostly on small chromosomes and the ends of large chromosomes (Fig. 2; Fig. S4).

The most extreme deviations from genome-wide genetic structure appear to cluster the Central

Mexico population with the northern lineage (Fig. S8), and in some cases also cluster the Sierra

Madre Oriental population with the northern lineage (Fig. S8; Fig. 3). As an example, we plot the

phylogeographic structure for one of these outlier windows in Fig. 3, demonstrating the

clustering of the Sierra Madre Oriental and Central Mexico populations with the northern

lineage.

Correlations between phylogeographic structure estimates and genomic architecture.

Almost all characteristics of the genome, population genomic summary statistics, and statistics

characterizing variation in phylogeographic structure were strongly correlated in sliding windows

across the genome (Fig. 4; Fig. S9). Notably, the relationship between population pairwise

genetic divergence (D_{XY}) and recombination rate variation varied with different population

comparisons (Fig. S5); D_{XY} was positively correlated in all the comparisons except the Rocky

Mountain × Central Mexico population, which exhibited a strong negative correlation between

D_{XY} and recombination rate (Fig. S5).

We found a strong correlation between different measures of phylogeographic structure,

including GSI at the lineage level, MDS1 of the PCA, and ADMIXTURE deviations (Fig. 4; Fig.

S9). Phylogeographic structure deviations (i.e., most different from genome-wide patterns) were

strongly positively correlated with recombination rate, GC content, and genetic diversity (Fig. 4;

Fig. S9). In contrast, TE content, Tajima's D, and GSI calculations at the population level had no

or weak correlations with all other statistics that we calculated (Fig. 4; Fig. S9). Correlations

were consistent between 50 kbp and 100 kbp estimates, with no correlations deviating more

than ~0.12 when estimated with different window sizes (Fig. S9).

Using variance partitioning, recombination rate variation most strongly explained

deviations in phylogeographic structure, but interactions of recombination rate, GC content,

CDS content, and TE content also explained some portions of the variance (Table 2). Most

notably, genomic architecture explained approximately 60% of the variance in both (1) the

MDS1 axis explaining differences in PCA-based estimates of genetic structure and (2)

ADMIXTURE results at the lineage level with an assumed k = 2 (Table 2). The relationship

between recombination rate and phylogeographic structure is exemplified when comparing the

genomic windows with the highest or lowest values of recombination rate; estimates of

phylogeographic structure and genetic diversity have little to no overlap in statistic distributions

in high recombination regions versus low recombination regions (Fig. 5). Additionally, when

comparing the genomic windows with relatively higher or lower recombination rates, all the

populations exhibit shifts in their relative RTD distributions (Fig. S10).

Because the effective population sizes for each locality varied substantially (Fig. 1; Fig.

S11), we looked to see if there was a relationship between relative RTD and population size in

regions with very different recombination rates. Using both the raw measurements and PICs, we

found a negative association between relative rates of evolution and harmonic mean effective

population sizes (Fig. 6). Notably, there was also a strong positive association between effective

population sizes and shifts in relative rates of evolution between high and low recombination

regions (Fig. 6). We also looked at the relationship between population sizes and potential

mutational load; we found no statistically significant associations but did find a slight, non-

significant, negative association between population sizes and potential mutational load in high

recombination regions (Fig. S12).

DISCUSSION

We investigated how variation in genomic architecture shapes local genomic phylogeographic structure in the Brown Creeper. We found that recombination rate and other characteristics of the genome explain 6-68% of the variance in phylogeographic structure inference using different metrics and genomic sliding window sizes (Table 2).

Genomic architecture shapes genomic diversity, divergence, and differentiation.

For most population genomic statistics, including measures of genetic diversity, divergence, and differentiation, we found significant associations with multiple measures of genomic characteristics including recombination rate and GC, gene, and repetitive DNA content (Fig. 4; Fig. S9). The strongest correlations were those with local recombination rate, so we focus our discussion here on the relationships between population genomics summary statistics and recombination rate.

In sliding windows across the genome, we found a strong negative correlation between genetic differentiation (F_{ST}) and local recombination rate (Fig. 4). Notably, this relationship was largely driven by population pairwise comparisons with strong genetic differentiation (Fig. S5); population comparisons with weak differentiation showed little relationship between genetic differentiation and recombination rate (Fig. S5). A negative relationship between local genomic differentiation and recombination rate has been identified in several other species, including birds, fruit flies, fishes, and monkeyflowers (Kulathinal, et al. 2008; Manthey, et al. 2021; Roesti, et al. 2013; Stankowski, et al. 2019; Vijay, et al. 2016). We may expect this relationship due to increased effects of linked selection and genetic drift in genomic regions with low recombination rates, particularly when gene flow is not strong enough to homogenize large genomic portions between differentiating taxa.

Genetic diversity—as measured by nucleotide diversity and observed heterozygosity— was positively associated with recombination rate across the genome (Fig. 4; Fig. S9). We may expect increased diversity in high recombination regions due to decreased effects of linked selection and genetic drift as well as a higher propensity for gene flow. This positive association between genetic diversity and recombination rate has been observed in multiple species of animals, fungi, and plants (Branca, et al. 2011; Cutter and Payseur 2013, 2003; Dutoit, et al. 2017b; Kraft, et al. 1998; Kulathinal, et al. 2008; Moreira, et al. 2022; Roesti, et al. 2013; Takahashi, et al. 2004; Thom, et al. 2021; Vijay, et al. 2016; Vijay, et al. 2017).

In our dataset, genetic divergence (D_{XY}) has a complicated relationship with

recombination rate; when using the mean genetic divergence across all population pairwise

comparisons measured, we identified a positive association between recombination rate and

genetic divergence (Fig. 4; Fig. S9). This relationship holds true for most population pairwise

comparisons except when we compare the two most divergent lineages (Central Mexico ×

Rocky Mountains) where we find a negative association between recombination rate and

genetic divergence (Fig. S5). We found positive, negative, and no association between genetic

divergence (D_{XY}) and differentiation (F_{ST}) across different population pairwise comparisons (Fig.

S5). These alternative relationships between genetic divergence and both recombination rate

and genetic differentiation may be interpreted in the context of the populations' divergence

scenario. In a model of population differentiation lacking gene flow, differentiation and

divergence may both positively scale with time in isolation, and both genetic differentiation and

divergence will exhibit a negative association with recombination rate variation (e.g., Manthey,

et al. 2021). In contrast, in a model of differentiation with gene flow, we may expect mixed

associations between genetic divergence and both genetic differentiation and recombination

rate variation because of the complex interactions between gene flow and selection against

gene flow in different genomic regions (Stankowski, et al. 2019). Based on these patterns, we

may intuit that the two main Brown Creeper lineages (red and blue in Fig. 1A) have diverged in

isolation (as previously suggested by Manthey, et al. 2021) while all the other population

comparisons examined here have differentiation scenarios including some amount of gene flow.

Phylogeographic structure shaped by genomic architecture

We used genetic clustering, ordination, and phylogenetic methods to assess phylogeographic

structure in sliding windows across the genome. We found that recombination rate variation was

the strongest predictor of phylogeographic structure variation across the genome (Table 2).

Previous work in insects, birds, and mammals has shown that phylogenomic signal

varies across the genome (Edelman, et al. 2019; Fontaine, et al. 2015; Li, et al. 2019; Martin, et

al. 2019; Thom, et al. 2021). Generally, these studies have found or suggested that the

background species tree is best represented by phylogenomic patterns exhibited in genomic

regions with low recombination, and that gene flow among taxa is more prevalent in genomic

regions with high recombination rate (Edelman, et al. 2019; Li, et al. 2019; Martin, et al. 2019;

Thom, et al. 2021). Our results in the Brown Creeper echo these findings, whereby we find

highest support for the genome-wide supported phylogenomic relationships in genomic regions

with low recombination rate (Fig. 2; Fig. S4). We may interpret these patterns in the context of

lineage sorting during the speciation process, whereby genomic regions with low introgression

exhibit quicker polymorphism fixation due to relatively increased effects of linked selection and

genetic drift and reduced homogenization among lineages due to gene flow.

Genomic studies investigating intraspecific population and phylogeographic structure

have more commonly used PCA-based methods (e.g., LOSTRUCT) to investigate variation in

genomic structure across the genome. In two species of woodpeckers, a small portion of the

variance in PCA-based phylogeographic structure was found to be associated with local

recombination rate (Moreira, et al. 2022). We found that variation in phylogeographic structure

across the genome of the Brown Creeper—inferred using both ordination and genetic clustering

methods—was strongly associated with recombination rate variation (Fig. 2; Fig. S4; Table 2).

We may infer that the same interactions between recombination rate variation and population

genomics processes shaping variance in phylogenomic signal are also shaping variance in

clustering- and ordination-based estimates of phylogeographic structure.

In contrast to generalizations about low recombination regions often reflecting a taxon's

background evolutionary history, in some scenarios we may expect regions of extremely low

recombination to greatly differ from genome-wide signatures of population genetic structure. In

several taxa, including sunflowers, flies, fishes, and birds, the greatest deviations from the

genome-wide pattern of intraspecific population structure (inferred with PCA) have been

identified in genomic regions with extremely low recombination rates that are often inferred to be

polymorphic inversions within populations or species (Hale, et al. 2021; Huang, et al. 2020; Li

and Ralph 2019; Mérot, et al. 2021; Perrier, et al. 2020; Shi, et al. 2021; Todesco, et al. 2020;

Whiting, et al. 2021). In these cases, if a taxon has relatively little population structure but also

exhibits polymorphic inversions, we may expect that the biggest deviations in population genetic

structure from genome-wide patterns to reflect these polymorphic inversions that exhibit

suppressed recombination.

Interaction of recombination rate, population demography, and rate of evolution

While we may expect that the interactions between recombination rate and population genomics

processes will impact the rate of evolution across the genome, we did not simply observe little

phylogeographic structure in genomic regions with high recombination rates and strong

phylogeographic structure in genomic regions with low recombination rates. Conversely, we

found strong—and differing—patterns of phylogeographic structure across the genome (Fig. 1;

Fig. 3; Fig. S8). Because these differing patterns of phylogeographic structure generally

separated the Central American populations and sometimes the Sierra Madre Oriental

population from all others, we hypothesized that populations with relatively smaller N_E exhibit

relatively higher rates of evolution in genomic regions with high recombination. Indeed, using

relative RTD measures as a relative rate of evolution for each population, we found that larger

populations exhibited reduced rates of molecular evolution in genomic windows with high

recombination rates (Fig. 6; Fig. S10) and that shifts in relative rates of molecular evolution in

high versus low recombination genomic regions were positively associated with effective

population sizes (Fig. 6; Fig. S10).

We suspect that molecular evolution in genomic regions with high recombination rates is

a product of complex interactions between population demography and the relative strengths of

linked selection and genetic drift. In the Brown Creeper genome, recombination rate variation is

positively associated with gene density (Fig. 2; Fig. 4) and we may interpret gene-dense

genomic regions as targets for natural selection. Indeed, in a previous study in the Brown

Creeper (Manthey, et al. 2021), we showed that the two main Brown Creeper lineages exhibited

relatively less neutral evolution in the gene-dense microchromosomes relative to the larger

macrochromosomes. In gene-dense and high recombination regions under the effects of

purifying selection, we may expect populations with relatively smaller N_E to accumulate

substitutions faster than populations with higher N_E (Lanfear, et al. 2014), consistent with the

trends we identified here in the Brown Creeper (Fig. 6; Fig. S11). Additionally, when we looked

at potential mutational load, we found a weak but nonsignificant negative association between

effective population size and mutational load accumulation in genomic regions with high

recombination rates (Fig. S12); this weak association also lends support to the idea that smaller

populations have relatively less efficient purifying selection than larger populations in high

recombination regions.

Implications for future phylogeographic studies

If phylogeographic structure may vary widely across the genome, what are realistic and best-

practice approaches for future phylogeographic work? Are studies that use few genes or

reduced representation genomic datasets doomed to be wrong from the start? We expect that

strong phylogeographic structure will be detected even with few genetic markers. For example,

more than ten years ago we used 20 genetic markers obtained with Sanger sequencing to study

the phylogeography of the Brown Creeper (Manthey, et al. 2011b) and found the same general

trends observed with our whole-genome dataset here. In contrast, any study where most of the

genetic markers are taken from genomic regions where the pattern differs from a taxon's true

evolutionary history (e.g., high recombination regions in the Brown Creeper case) may run into

problems inferring that true history. Additionally, in any taxa with weak population genetic

structure or widespread and rampant gene flow, a whole-genome method may be the only

approach to identify the few genomic regions differentiating taxa (e.g., Toews, et al. 2016).

Overall, we suggest using the most amount of genetic data feasible for future phylogeographic

studies given sources of biological material (e.g., quality of preserved tissue) and funding

available, as well as investigating the patterns and potential causes of population genetic

structure variation across the genome when possible.

CONCLUSIONS

We used whole genome sequencing to estimate how genomic architecture shapes variation in

phylogeographic structure across the genome of the Brown Creeper. We found that population

genetic structure—as measured using summary statistics, phylogenomic methods, and

population clustering methods—was strongly associated with regional genomic variation in

recombination rates. In low recombination regions, we recovered phylogeographic structure

concordant with genome-wide patterns. The most divergent phylogeographic patterns were in

high recombination regions, wherein populations with small effective population sizes were

distinct from all other populations due to relatively faster evolution than larger populations in

these high recombination regions. Because these high recombination regions are rich in coding

genes, we suspect that large populations have higher relative effects of purifying selection in

these regions, and overall slower relative evolution compared to smaller populations. Overall,

our results show that phylogeographic structure may vary widely across the genome, and that

effective population sizes of sampled populations and genomic architecture and their complex

interactions will impact regional genomic variation in phylogeographic structure.

ACKNOWLEDGEMENTS

We would like to thank the collection managers, curators, and contributors for generous tissue loans for all individuals used in this study from the following museums: Burke Museum of Natural History and Culture, and University of Kansas Biodiversity Institute. Sequencing was supported by Texas Tech University start-up funding to JDM and a generous donation from the Ferguson family to GMS and the Denver Museum of Nature and Science. Support was also provided by NSF awards #1953688 to JDM and #0814841 to GMS. The High-Performance Computing Center at TTU supported computational analyses.

DATA ACCESSIBILITY

All raw resequencing data are currently being uploaded to NCBI's SRA database. All code used for analysis on this project is available on GitHub: (github.com/jdmanthey/certhia_phylogeography).

AUTHOR CONTRIBUTIONS

JDM and GMS designed the study. JDM performed laboratory and bioinformatic work and wrote the first draft of the manuscript. JDM and GMS contributed to revising and improving the final draft of the manuscript.

 TABLES AND FIGURES

Figure 1. Sampling map, genetic diversity, phylogeographic structure, and demography. (A)
Sampling localities for this study. Colors for localities are consistent across plots. (B) Violin plots
of nucleotide diversity across all 50 kbp sliding windows. White circles indicate means. (C)
Consensus phylogeny using 19,639 gene trees from the 50 kbp windows. Support values are
labelled for all branches showing the highest support from all estimation methods (N = 4),
including consensus and ASTRAL phylogenies for both the 50 kbp and 100 kbp window
datasets. At tips of phylogeny are ADMIXTURE results from 19,602 genome-wide SNPs
(thinned every 50 kbp) for either two or seven assumed genetic clusters (k = 2 or k = 7,
respectively). (D) Demographic history estimated with MSMC2. Different individuals are
represented with differently colored lines and bootstraps are shown with thin lines. In (D), HM =
harmonic mean. Brown Creeper photo in (A) by JDM from the Chiricahua Mountains in southern
Arizona.

Figure 2. Genomic architecture and phylogeographic structure variation across the genome in
50kbp sliding windows. Gray points indicate window values and dark gray lines indicate means
across 20 windows (i.e., 1 Mbp windows). Blue indicates top 2.5% of outliers and orange
indicates bottom 2.5% of outliers for each statistic. Bottom outliers not shown for CDS, ADMIX.,
and GSI as these each have large proportions of values equal to zero. Abbreviations:
recombination rate (ρ), gene content (CDS), transposable element and repetitive DNA content
(TE), multidimensional scaling axis one from principal components analyses (-MDS1), mean
genealogical sorting index for the two main *Certhia* lineages (1 - GSI), ADMIXTURE deviation
relative to genome-wide analysis (ADMIX.), and mean genetic diversity across individuals as
measured using observed heterozygosity (H_O). MDS1 is plotted as negative and GSI is plotted
as (1 - GSI) so that deviations from expected phylogeographic structure for MDS1, GSI, and
ADMIXTURE are all represented as higher values.

591 Figure 3. Example phylogeographic structure in a window that deviates from genome-wide
592 patterns. This window had the 15th lowest MDS1 value of all windows from the principal
593 components analysis (PCA) across the genome. (A) PCA, (B) RAxML phylogeny, and (C)
594 ADMIXTURE results for a window on chromosome 4.

Figure 4. Correlation plot of genomic architecture, genomic diversity, and phylogeographic structure for 50 kbp windowed statistics. Pearson's and Spearman's r coefficients are plotted above and below the diagonal, respectively. Abbreviations shown in plot: recombination rate (ρ); GC content (GC%); gene content (CDS%); transposable element content (TE%); mean observed heterozygosity (H_O); mean nucleotide diversity (π); mean pairwise genetic divergence (D_{XY}); mean pairwise genetic differentiation (F_{ST}); genealogical sorting index at the population [1 - GSI (Pop.)] and lineage [1 - GSI (Lin.)] levels; negative multidimensional scaling axis one from principal components analyses (-MDS1); ADMIXTURE deviation relative to genome-wide analysis for two [ADMIX. (k = 2)] or seven [ADMIX. (k = 7)] assumed genetic clusters.

Figure 5. Violin plots of phylogeographic structure and genetic diversity variation in genomic
regions with very high or low recombination rates. 50 kbp windows with the highest or lowest
2.5% of recombination rate values are included. Higher values for ADMIXTURE deviations, 1 -
GSI, and -MDS1 indicate values deviating relatively more from the genome-wide patterns of
phylogeographic structure. For nucleotide diversity (π), D_{XY}, and F_{ST}, values for each window
are the mean across all populations (π) or population pairwise comparisons (D_{XY} and F_{ST}).

Figure 6. Relationship between effective population sizes and rate of evolution (measured by
relative root-to-tip distance for each population) in genomic regions that vary in recombination
rate. Extreme low and high recombination (rho) regions represent the quintiles of the genome
with the lowest or highest recombination rates (i.e., highest 20% and lowest 20% of windows).
Results for raw data are shown in (A-C) and for phylogenetic independent contrasts (PIC) in (D-
F). The right panels in each row (C + F) show the shift in relative rates of evolution between
high and low recombination genomic regions.

681 Table 1. Different datasets used in this study and their characteristics. For inclusion, all datasets
682 required sites to have a minimum quality score of 20, minimum of 20 genotype quality, a
683 minimum depth per individual for a site to be called of 6, and a max mean depth across
684 individuals of 50.

Set	Missing Allowed	OG[1]	Window size	# Windows	Min. Sep. per SNP[2]	# SNPs	# Sites	Analyses
A	0%	-	-	-	50 kbp	19,602	19,602	Genome-wide ADMIXTURE
B	0%	-	-	-	100 kbp	9,838	9,838	Genome-wide ADMIXTURE
C	20%	X	50 kbp	19,713	-	35,463,408	890,335,872	RAxML, GSI, H_O, F_{ST}, D_{XY}, Tajima's D
D	20%	X	100 kbp	9,862	-	35,463,408	890,335,872	RaxML, GSI, H_O, F_{ST}, D_{XY}, Tajima's D
E	0%		-	-	-	21,657,220	21,657,220	LOSTRUCT
F	0%		50 kbp	19,697	-	21,657,220	21,657,220	ADMIXTURE per window
G	0%		100 kbp	9,860	-	21,657,220	21,657,220	ADMIXTURE per window
H	0%		-	-	-	*	*	MSMC2

[1]OG = outgroup included

[2]Min. Sep. per SNP = minimum separation between SNPs (i.e., thinning)

*SNPs and sites not labeled for MSMC2 analyses because these values varied per individual. For each individual, no missing data was allowed for that particular individual's MSMC2 demographic analyses input files. The MSMC2 dataset did not include the Z chromosome.

688 Table 2. Variance partitioning where the explanatory variables are characteristics of genomic
689 architecture and response variables are different measures of population genetic structure.
690 Values indicative of variance proportion explained by each explanatory variable or their
691 interactions.

50 kbp windows						
Measure of Genetic Structure	ρ	CDS	TE	GC	Interactions	Total
MDS1	0.23	0.00	0.01	0.04	0.33	0.61
GSI (Lineages)	0.16	0.00	0.00	0.02	0.22	0.40
GSI (Populations)	0.02	0.00	0.00	0.01	0.03	0.06
ADMIXTURE (k = 2)	0.24	0.00	0.00	0.03	0.30	0.57
ADMIXTURE (k = 7)	0.19	0.00	0.00	0.01	0.18	0.38
Mean F_{ST}	0.13	0.00	0.01	0.01	0.15	0.30
100 kbp windows						
Measure of Genetic Structure	ρ	CDS	TE	GC	Interactions	Total
MDS1	0.22	0.01	0.01	0.03	0.41	0.68
GSI (Lineages)	0.16	0.00	0.00	0.01	0.23	0.40
GSI (Populations)	0.02	0.00	0.00	0.01	0.05	0.08
ADMIXTURE (k = 2)	0.25	0.00	0.00	0.02	0.35	0.62
ADMIXTURE (k = 7)	0.20	0.00	0.00	0.01	0.22	0.43
Mean F_{ST}	0.15	0.00	0.02	0.01	0.17	0.35

SUPPLEMENTAL TABLES AND FIGURES

Table S1. Sampling metadata and characteristics of sequencing amount and coverage when aligned to the reference genome. Samples from Utah, Morelos, Jalisco, and the outgroup were retrieved from the NCBI SRA and originally sequenced for a previous study by Manthey and colleagues (2021).

Museum #	Lineage	Locality	Lat.	Long.	Raw bp	Filtered bp	Coverage
UWBM103417	South	Copan, HN	14.866	-89.050	40,501,485,458	34,714,897,849	22.79
UWBM105618	South	Copan, HN	14.866	-89.050	30,149,924,172	26,420,616,801	18.55
UWBM93696	South	Copan, HN	14.866	-89.050	40,591,948,048	36,089,400,965	26.06
UWBM113447	South	Chiapas, MX	16.691	-92.606	37,741,499,908	33,063,229,986	23.32
UWBM113461	South	Chiapas, MX	16.691	-92.606	32,696,721,580	27,488,157,228	18.51
UWBM113462	South	Chiapas, MX	16.691	-92.606	48,519,014,074	42,841,474,607	27.09
UWBM106936	South	Morelos, MX	19.0873	-99.194	40,136,994,108	35,563,542,819	26.35
UWBM107890	South	Jalisco, MX	21.882	-103.865	43,601,103,228	38,563,138,625	26.14
UWBM107897	South	Jalisco, MX	21.882	-103.865	30,536,607,690	27,038,872,945	18.55
UWBM106768	South	Nuevo Leon, MX	24.872	-100.224	42,330,853,397	37,109,282,807	27.62
UWBM111691	South	Nuevo Leon, MX	24.872	-100.224	41,523,724,956	36,804,470,356	26.21
UWBM111708	South	Nuevo Leon, MX	24.872	-100.224	43,392,483,440	36,820,385,840	27.04
UWBM113198	North	California, US	36.245	-121.700	40,117,812,880	35,379,027,102	26.26
UWBM112794	North	California, US	36.245	-121.553	37,692,762,846	33,128,089,974	23.33
UWBM112795	North	California, US	36.245	-121.553	44,376,620,806	38,809,177,747	28.71
UWBM113162	North	Utah, US	37.317	-113.455	40,934,172,636	36,217,919,293	26.25
UWBM113168	North	Utah, US	37.317	-113.455	39,295,062,200	34,925,676,137	25.72
UWBM113167	North	Utah, US	37.377	-113.467	42,535,860,138	37,425,027,332	27.55
UWBM107061	North	West Virginia, US	38.289	-79.937	43,886,484,470	38,699,509,454	28.65
UWBM112055	North	West Virginia, US	38.289	-79.937	47,972,794,224	40,458,960,409	29.36
UWBM112054	North	West Virginia, US	38.641	-79.838	39,431,412,482	34,774,499,050	25.96
KU92846	Outgroup	England			21,417,629,506	19,195,682,239	13.58

708   Figure S1. Sequencing coverage aligned to the reference genome for each individual. The

709   vertical red lines indicate the mean coverage per individual.

731 Figure S2. Genetic differentiation (F_{ST}) variation across the genome in 100kbp sliding windows.

732 Gray points indicate window values and dark gray lines indicate means across ten windows

733 (i.e., 1 Mbp windows).

742 Figure S3. Genetic divergence (D_{XY}) variation across the genome in 100kbp sliding windows.

743 Gray points indicate window values and dark gray lines indicate means across ten windows

744 (i.e., 1 Mbp windows).

Figure S4. Genomic architecture and phylogeographic structure variation across the genome in 100kbp sliding windows. Gray points indicate window values and dark gray lines indicate means across ten windows (i.e., 1 Mbp windows). Blue indicates top 2.5% of outliers and orange indicates bottom 2.5% of outliers for each statistic. Bottom outliers not shown for CDS, ADMIX., and GSI as these each have large proportions of values equal to zero. Abbreviations: recombination rate (ρ), gene content (CDS), transposable element and repetitive DNA content (TE), multidimensional scaling axis one from principal components analyses (-MDS1), mean genealogical sorting index for the two main *Certhia* lineages (1 - GSI), ADMIXTURE deviation relative to genome-wide analysis (ADMIX.), and mean genetic diversity across individuals as measured using observed heterozygosity (H_O). MDS1 is plotted as negative and GSI is plotted as (1 - GSI) so that deviations from expected phylogeographic structure for MDS1, GSI, and ADMIXTURE are all represented as higher values.

765   Figure S5. Correlations between genetic divergence (D_{XY}), genetic differentiation (F_{ST}), and

766   recombination rate variation in 100 kbp sliding windows.

787 Figure S6. Nucleotide diversity variation across the genome in 100kbp sliding windows. Gray
788 points indicate window values and dark gray lines indicate means across ten windows (i.e., 1
789 Mbp windows).

798   Figure S7. Tajima's D variation across the genome in 100kbp sliding windows. Gray points

799   indicate window values and dark gray lines indicate means across ten windows (i.e., 1 Mbp

800   windows).

809   Figure S8. The 15 most extreme outlier principal components analysis (PCA) windows identified

810   from those windows with low MDS dimension one values.

Figure S9. Correlation plot of genomic architecture, genomic diversity, and phylogeographic
structure for (A) 50 kbp and (B) 100 kbp windowed statistics. Pearson's and Spearman's r
coefficients are plotted above and below the diagonal, respectively. In (C), the absolute value
difference between correlations in 50 kbp and 100 kbp windows. Abbreviations shown in plot:
recombination rate (ρ); GC content (GC%); gene content (CDS%); transposable element
content (TE%); mean observed heterozygosity (H_O); mean nucleotide diversity (π); mean
pairwise genetic divergence (D_{XY}); mean pairwise genetic differentiation (F_{ST}); genealogical
sorting index at the population [1 - GSI (Pop.)] and lineage [1 - GSI (Lin.)] levels; negative
multidimensional scaling axis one from principal components analyses (-MDS1); ADMIXTURE
deviation relative to genome-wide analysis for two [ADMIX. (k = 2)] or seven [ADMIX. (k = 7)]
assumed genetic clusters.

Figure S10. Violin plots of evolutionary rates (as measured by relative root-to-tip distances) for

each population in the upper and lower genomic quintiles of high or low recombination (rho)

values.

Figure S11. Demographic history estimated with MSMC2 for each individual. Thick lines indicate

the whole genome estimate and thin lines indicate results from ten bootstrap replicates. Sample

numbers correspond with those reported in Table S1.

Figure S12. Relationship between effective population sizes and potential mutational load in

genomic regions that vary in recombination rate. Extreme low and high recombination (rho)

regions represent the quintiles of the genome with the lowest or highest recombination rates

(i.e., highest 20% and lowest 20% of windows). Results for raw data are shown in (A-C) and for

phylogenetic independent contrasts (PIC) in (D-F). The right panels in each row (C + F) show

the shift in potential mutational load between high and low recombination genomic regions.

LITERATURE CITED

Alexander DH, Novembre J, Lange K 2009. Fast model-based estimation of ancestry in unrelated individuals. Genome Research 19: 1655-1664.

Allmon WD 1992. A causal analysis of stages in allopatric speciation. Oxford surveys in evolutionary biology 8: 219-219.

Branca A, et al. 2011. Whole-genome nucleotide diversity, recombination, and linkage disequilibrium in the model legume Medicago truncatula. Proceedings of the National Academy of Sciences, USA 108: E864-E870.

Bushnell B. 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner. In: Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, CA (US).

Cingolani P, et al. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly 6: 80-92.

Cruickshank TE, Hahn MW 2014. Reanalysis suggests that genomic islands of speciation are due to reduced diversity, not reduced gene flow. Molecular Ecology 23: 3133-3157.

Cummings MP, Neel MC, Shaw KL 2008. A genealogical approach to quantifying lineage divergence. Evolution 62: 2411-2422.

Cutter AD, Payseur BA 2013. Genomic signatures of selection at linked sites: unifying the disparity among species. Nature reviews genetics 14: 262-274.

Cutter AD, Payseur BA 2003. Selection at linked sites in the partial selfer *Caenorhabditis elegans*. Molecular Biology and Evolution 20: 665-673.

Danecek P, et al. 2011. The variant call format and VCFtools. Bioinformatics 27: 2156-2158.

Dutoit L, Burri R, Nater A, Mugal CF, Ellegren H 2017a. Genomic distribution and estimation of nucleotide diversity in natural populations: perspectives from the collared flycatcher (Ficedula albicollis) genome. Molecular Ecology Resources 17: 586-597.

Dutoit L, et al. 2017b. Covariation in levels of nucleotide diversity in homologous regions of the avian genome long after completion of lineage sorting. Proceedings of the Royal Society B: Biological Sciences 284: 20162756.

Edelman NB, et al. 2019. Genomic architecture and introgression shape a butterfly radiation. Science 366: 594-599.

Ellegren H 2010. Evolutionary stasis: the stable chromosomes of birds. Trends in Ecology & Evolution 25: 283-291.

Fontaine MC, et al. 2015. Extensive introgression in a malaria vector species complex revealed by phylogenomics. Science 347: 1258524.

Haenel Q, Laurentino TG, Roesti M, Berner D 2018. Meta-analysis of chromosome-scale crossover rate variation in eukaryotes and its significance to evolutionary genomics. Molecular Ecology 27: 2477-2497.

Hale MC, Campbell MA, McKinney GJ 2021. A candidate chromosome inversion in Arctic charr (*Salvelinus alpinus*) identified by population genetic analysis techniques. G3 11: jkab267.

Harrell FE, Dupont C. 2020. Hmisc: Harrell Miscellaneous. Version 4.3-1. R: R.

Hey J, Kliman RM 2002. Interactions between natural selection, recombination and gene density in the genes of Drosophila. Genetics 160: 595-608.

Hobolth A, Dutheil JY, Hawks J, Schierup MH, Mailund T 2011. Incomplete lineage sorting
patterns among human, chimpanzee, and orangutan suggest recent orangutan speciation
and widespread selection. Genome Research 21: 349-356.
Huang K, Andrew RL, Owens GL, Ostevik KL, Rieseberg LH 2020. Multiple chromosomal
inversions contribute to adaptive divergence of a dune sunflower ecotype. Molecular
Ecology 29: 2535-2549.
Kapusta A, Suh A 2017. Evolution of bird genomes—a transposon's-eye view. Annals of the
New York Academy of Sciences 1389: 164-185.
Kawakami T, et al. 2014. A high-density linkage map enables a second-generation collared
flycatcher genome assembly and reveals the patterns of avian recombination rate
variation and chromosomal evolution. Molecular Ecology 23: 4035-4058.
Koonin EV 2009. Evolution of genome architecture. The international journal of biochemistry &
cell biology 41: 298-306.
Kraft T, Säll T, Magnusson-Rading I, Nilsson N-O, Halldén C 1998. Positive correlation
between recombination rates and levels of genetic variation in natural populations of sea
beet (Beta vulgaris subsp. maritima). Genetics 150: 1239-1244.
Kulathinal RJ, Bennett SM, Fitzpatrick CL, Noor MA 2008. Fine-scale mapping of
recombination rate in *Drosophila* refines its correlation to diversity and divergence.
Proceedings of the National Academy of Sciences, USA 105: 10051-10056.
Lanfear R, Kokko H, Eyre-Walker A 2014. Population size and the rate of evolution. Trends in
Ecology & Evolution 29: 33-41.
Li G, Figueiró HV, Eizirik E, Murphy WJ 2019. Recombination-aware phylogenomics reveals
the structured genomic landscape of hybridizing cat species. Molecular Biology and
Evolution 36: 2111-2126.
Li H, Durbin R 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics 25: 1754-1760.
Li H, et al. 2009. The sequence alignment/map format and SAMtools. Bioinformatics 25: 2078-
2079.
Li H, Ralph P 2019. Local PCA shows how the effect of population structure differs along the
genome. Genetics 211: 289-304.
Linck E, Battey C 2019. Minor allele frequency thresholds strongly affect population structure
inference with genomic data sets. Molecular Ecology Resources 19: 639-647.
Louis M, et al. 2020. Selection on ancestral genetic variation fuels parallel ecotype formation in
bottlenose dolphins. hal-03020428.
Lucas T 2016. palettetown: Use Pokemon Inspired Colour Palettes. R package version 0.1.1.
https://CRAN.R-project.org/package=palettetown.
Manthey J, Bourgeois Y, Meheretu Y, Boissinot S 2022. Varied diversification patterns and
distinct demographic trajectories in Ethiopian montane forest bird (Aves: Passeriformes)
populations separated by the Great Rift Valley. doi: 10.1111/mec.16417. Molecular
Ecology.
Manthey JD, Klicka J, Spellman GM 2015. Chromosomal patterns of diversity and
differentiation in creepers: a next-gen phylogeographic investigation of *Certhia*
americana. Heredity 115: 165-172.
Manthey JD, Klicka J, Spellman GM 2011a. Cryptic diversity in a widespread North American
songbird: phylogeography of the Brown Creeper (Certhia americana). Molecular
Phylogenetics and Evolution 58: 502-512.

Manthey JD, Klicka J, Spellman GM 2021. The genomic signature of allopatric speciation in a songbird is shaped by genome architecture (Aves: *Certhia americana*). Genome Biology and Evolution 13: evab120.

Manthey JD, Klicka J, Spellman GM 2011b. Isolation-driven divergence: speciation in a widespread North American songbird (Aves: Certhiidae). Molecular Ecology 20: 4371-4384.

Martin SH, Davey JW, Salazar C, Jiggins CD 2019. Recombination rate variation shapes barriers to introgression across butterfly genomes. PLoS Biology 17: e2006288.

Mayr E, Mayr E, Mayr E, Mayr E. 1963. Animal species and evolution. In: Belknap Press of Harvard University Press Cambridge, Massachusetts.

McKenna A, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. Genome Research 20: 1297-1303.

Mérot C, et al. 2021. Locally adaptive inversions modulate genetic variation at different geographic scales in a seaweed fly. Molecular Biology and Evolution 38: 3953-3971.

Moreira LR, Klicka J, Smith BT 2022. Demography and linked selection interact to shape the genomic landscape of codistributed woodpeckers during the Ice Age. bioRxiv: DOI: https://doi.org/10.1101/2022.1102.1104.479011.

Nachman MW, Payseur BA 2012. Recombination rate variation and speciation: theoretical predictions and empirical results from rabbits and mice. Philosophical Transactions of the Royal Society B: Biological Sciences 367: 409-421.

Nadachowska-Brzyska K, Li C, Smeds L, Zhang G, Ellegren H 2015. Temporal dynamics of avian populations during Pleistocene revealed by whole-genome sequences. Current Biology 25: 1375-1380.

Neuwirth E 2014. RColorBrewer: ColorBrewer Palettes. R package version 1.1-2. https://CRAN.R-project.org/package=RColorBrewer.

Oksanen J, et al. 2007. The vegan package. Community ecology package 10: 631-637.

Pagès H, Aboyoun P, Gentleman R, DebRoy S 2017. Biostrings: Efficient manipulation of biological strings. R Package Version 2.0.

Paradis E, Claude J, Strimmer K 2004. APE: analyses of phylogenetics and evolution in R language. Bioinformatics 20: 289-290.

Perrier C, Rougemont Q, Charmantier A 2020. Demographic history and genomics of local adaptation in blue tit populations. Evolutionary applications 13: 1145-1165.

Poulin J, et al. 2020. Brown Creeper (Certhia americana), version 1.0. In Birds of the World. . In: Poole A, editor: Cornell Lab of Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.brncre.01.

Ravinet M, et al. 2018. The genomic landscape at a late stage of stickleback speciation: High genomic divergence interspersed by small localized regions of introgression. PLoS Genetics 14: e1007358.

Reich D, Thangaraj K, Patterson N, Price AL, Singh L 2009. Reconstructing Indian population history. Nature 461: 489.

Revell LJ 2012. phytools: an R package for phylogenetic comparative biology (and other things). Methods in Ecology and Evolution 3: 217-223.

Roesti M, Moser D, Berner D 2013. Recombination in the threespine stickleback genome—patterns and consequences. Molecular Ecology 22: 3014-3027.

Sayyari E, Mirarab S 2016. Fast coalescent-based computation of local branch support from quartet frequencies. Molecular Biology and Evolution 33: 1654-1668.

Schiffels S, Durbin R 2014. Inferring human population size and separation history from
multiple genome sequences. Nature Genetics 46: 919.
Shi Y, et al. 2021. Gene flow influences the genomic architecture of local adaptation in six
riverine fish species. bioRxiv: DOI: https://doi.org/10.1101/2021.1105.1118.444736.
Stamatakis A 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30: 1312-1313.
Stankowski S, et al. 2019. Widespread selection and gene flow shape the genomic landscape
during a radiation of monkeyflowers. PLoS Biology 17: e3000391.
Sukumaran J, Holder MT 2010. DendroPy: a Python library for phylogenetic computing.
Bioinformatics 26: 1569-1571.
Tacutu R, et al. 2017. Human Ageing Genomic Resources: new and updated databases. Nucleic
Acids Research 46: D1083-D1090.
Tajima F 1989. Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. Genetics 123: 585-595.
Takahashi A, Liu Y-H, Saitou N 2004. Genetic variation versus recombination rate in a
structured population of mice. Molecular Biology and Evolution 21: 404-409.
Thom G, et al. 2021. Genomic architecture controls spatial structuring in Amazonian birds.
bioRxiv: DOI: https://doi.org/10.1101/2021.1112.1101.470789.
Todesco M, et al. 2020. Massive haplotypes underlie ecotypic differentiation in sunflowers.
Nature 584: 602-607.
Toews DP, et al. 2016. Plumage genes and little else distinguish the genomes of hybridizing
warblers. Current Biology 26: 2313-2318.
Vijay N, et al. 2016. Evolution of heterogeneous genome differentiation across multiple contact
zones in a crow species complex. Nature Communications 7: 1-10.
Vijay N, et al. 2017. Genomewide patterns of variation in genetic diversity are shared among
populations, species and higher-order taxa. Molecular Ecology 26: 4284-4295.
Wei T, et al. 2017. Package 'corrplot'. Statistician 56: e24.
White MA, Ané C, Dewey CN, Larget BR, Payseur BA 2009. Fine-scale phylogenetic
discordance across the house mouse genome. PLoS Genetics 5: e1000729.
Whiting JR, et al. 2021. Drainage-structuring of ancestral variation and a common functional
pathway shape limited genomic convergence in natural high-and low-predation guppies.
PLoS Genetics 17: e1009566.
Wickham H 2011. ggplot2. Wiley Interdisciplinary Reviews: Computational Statistics 3: 180-
185.
Willing E-M, Dreyer C, Van Oosterhout C 2012. Estimates of genetic differentiation measured
by FST do not necessarily require large sample sizes when using many SNP markers.
PLoS One 7: e42649.
Zhang C, Rabiee M, Sayyari E, Mirarab S 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics 19: 153.
1035