Recombination rate variation shapes genomic variability of phylogeographic structure in a

widespread North American songbird (Aves: *Certhia americana*)

Joseph D. Manthey[1]*, Garth M. Spellman[2]

[1]Department of Biological Sciences, Texas Tech University. Lubbock, Texas, USA.

[2]Department of Zoology, Denver Museum of Nature & Science, Denver, Colorado, USA.

*Corresponding author: jdmanthey@gmail.com

26   **Running head:** Recombination rate variation shapes phylogeographic structure variation

ABSTRACT

The nonrandom distribution of chromosomal characteristics and functional elements—genomic architecture—impacts the relative strengths and impacts of population genetic processes across the genome. Due to this relationship, genomic architecture has the potential to shape variation in population genetic structure across the genome. Population genetic structure has been shown to vary across the genome in a variety of taxa, but this body of work has largely focused on pairwise population genomic comparisons between closely related taxa. Here, we used whole genome sequencing of seven phylogeographically structured populations of a North American songbird, the Brown Creeper (*Certhia americana*), to determine the impacts of genomic architecture on phylogeographic structure variation across the genome. Using multiple methods to infer phylogeographic structure—ordination, clustering, and phylogenetic methods— we found that recombination rate variation explained a large proportion of phylogeographic structure variation. Genomic regions with low recombination showed phylogeographic structure consistent with the genome-wide pattern. In regions with high recombination, we found strong phylogeographic structure, but with discordant patterns relative to the genome-wide pattern. In regions with high recombination rate, we found that populations with small effective population sizes evolve relatively more rapidly than larger populations, leading to discordant signatures of phylogeographic structure. These results suggest that the interplay between recombination rate variation and effective population sizes shape the relative impacts of linked selection and genetic drift in different parts of the genome. Overall, the combined interactions of population genetic processes, genomic architecture, and effective population sizes shape patterns of variability in phylogeographic structure across the genome of the Brown Creeper.

Keywords: phylogeography, population genomics, genetic architecture, demography

INTRODUCTION

The formation of population structure is implicit in the speciation process (Allmon, 1992; Mayr et al., 1963). Speciation is a continuum; early differentiating populations may have few genomic regions with evidence for genetic differentiation or barriers to gene flow (Ellegren et al., 2012; Nosil and Feder, 2012; Toews et al., 2016; Wu, 2001) while later in the speciation process two diverging populations may exhibit strong genetic differentiation across their entire genomes (Nosil and Feder, 2012; Ravinet et al., 2018; Wu, 2001). Across the genome, variation in local evolutionary history is shaped by the relative strengths of the population genomic processes of natural selection, gene flow, and genetic drift. For example, population genetic structure may differ in genomic regions with strong natural selection or differential patterns of gene flow relative to the rest of the genome (Louis et al., 2020; Whiting et al., 2021).

Variable impacts of selection, drift, and gene flow across the genome may be partially attributed to genomic architecture—the nonrandom distribution of chromosomal characteristics and functional elements (Koonin, 2009)—whereby variation in gene content, recombination rates, or other genomic characteristics impact the relative effects of population genomic processes. For example, we expect that genomic regions with lower recombination rates will be relatively depleted of genetic diversity and genetic differentiation will accumulate faster, due to relatively increased effects of linked selection (Cruickshank and Hahn, 2014; Haenel et al., 2018; Hey and Kliman, 2002; Nachman and Payseur, 2012). Indeed, speciation and population genomic studies have often found a negative relationship between recombination rates and genetic differentiation across the genome (Kulathinal et al., 2008; Manthey et al., 2021; Roesti et al., 2013; Stankowski et al., 2019; Vijay et al., 2016).

If genomic architecture impacts population genomic patterns within populations and between pairs of taxa, we may naturally extend this idea to phylogeographic patterns in taxa with multiple lineages at different stages of evolutionary distinctiveness. As such, we may

predict that estimates of phylogeographic structure will vary across the genome and be shaped

by several factors and their interactions: (1) population size variation of differentiating

populations shaping relative strengths of selection and drift, (2) time since divergence began,

(3) genomic architecture, and (4) quantity of interpopulation gene flow. Exemplar taxa to better

understand the impacts of genomic architecture on population genetic structure are species (or

species complexes) with multiple lineages on distinct evolutionary trajectories, populations with

different effective sizes, and genomes exhibiting heterogeneous genomic architecture.

The Brown Creeper (*Certhia americana*) is an excellent study organism to understand

the impacts of genomic architecture on phylogeographic structure; its genome is highly

heterogeneous in both structure and content, and it has seven distinct phylogeographic lineages

(Manthey et al., 2011a; Manthey et al., 2015). The genome of the Brown Creeper, as like other

birds, has chromosomes that (1) span two orders of magnitude in size, (2) vary in their gene

and repetitive element content, and (3) exhibit local effective recombination rates that can vary

by an order of magnitude or more (Dutoit et al., 2017; Ellegren, 2010; Kapusta and Suh, 2017;

Kawakami et al., 2014; Manthey et al., 2021). Additionally, the Brown Creeper has strong

phylogeographic structure, with two main lineages that have been diverging around one million

years (Fig. 1A), multiple phylogeographically structured clades within each of these two main

lineages, and populations with different effective sizes (Manthey et al., 2011a, b; Manthey et al.,

2015, 2021).

Using the Brown Creeper as our study taxon, we used whole-genome sequencing data

to assess how genome architecture influences phylogeographic structure; we used

phylogenetic, ordination, and population genetic clustering methods to characterize

phylogeographic structure across the genome. Our aims here were twofold. First, we looked to

decipher how genomic architecture and effective population sizes shape phylogeographic

structure variation across the genome. Second, because phylogeographic structure may be

estimated in many ways, we wanted to identify whether phylogeographic signal variation was

consistent using different methodologies.

Based on previous population genomic research, we hypothesized that genomic regions

with low recombination rates will have faster lineage sorting. Based on this hypothesis, we

would predict that phylogeographic estimates consistent with the species or population level

evolutionary history of the group will be concentrated in genomic regions with low

recombination. Additionally, we may predict a couple alternative patterns in high recombination

regions. We may identify mixed or weak phylogeographic structure in high recombination

regions. Alternatively, we may identify strong, but divergent patterns of phylogeographic

structure in high recombination regions linked with varied molecular evolutionary rates among

sampled populations. We may also predict an interaction between recombination rate and

population sizes; whereby, relatively smaller populations will undergo faster molecular evolution

in genomic regions characterized by stabilizing selection, such as in gene-dense, high

recombination genomic regions.

MATERIALS AND METHODS

Study organism, sampling, and sequencing. The Brown Creeper is a songbird found in

forested regions of the Americas; it has a widespread distribution ranging from Honduras in the

south to Alaska in the north (Fig. 1A) (Poulin et al., 2020). The Brown Creeper is aptly named; it

creeps up trees in search of its largely invertebrate diet and has dark coloration on its back and

sides (Fig. 1A inset) that makes for excellent camouflage on trees with dark bark. In previous

work using tens to thousands of genetic markers, we have identified phylogeographic structure

in the Brown Creeper, with up to seven distinct lineages (Manthey et al., 2011a, b; Manthey et

al., 2015). Here, we aimed to obtain population genomic sequencing data for seven populations

with three individuals sampled per population (Fig 1, Table S1). We downloaded previously

published sequencing data from two of these populations and one outgroup sample of *C.*

familiaris (Table S1) (Manthey et al., 2021). For the other five populations, we obtained

representative tissue samples from natural history museums and extracted genomic DNA using

QIAGEN DNeasy blood and tissue kits following manufacturer guidelines. DNA from extractions

was used to create standard Illumina sequencing libraries, followed by sequencing on an

Illumina NovaSeq6000 at the Texas Tech University Center for Biotechnology and Genomics

with the goal of obtaining ~15-30× coverage per individual.

Reference genome and genomic architecture. For the reference genome, we used the

chromosome-scale *Certhia americana* genome we assembled for a prior study (NCBI assembly

ASM1869719v1; Manthey et al. 2021). From data reported in this previous study (Manthey et al.

2021), we summarized mean effective recombination rates for each of the two main Brown

Creeper lineages, transposable element (TE) and gene density, and GC content across the

genome in 50kbp and 100 kbp non-overlapping sliding windows (Fig. 2). We used the mean

effective recombination rates of the two lineages because they were highly correlated (r =

0.829-0.861 across different correlation metrics; all p << 0.001), and we can assume relatively

conserved recombination rates over short evolutionary time frames (Singhal et al., 2015).

Briefly, these estimates were initially obtained with the following methods: effective

recombination rate was estimated using LDhat (McVean and Auton, 2007), TEs were annotated

with RepeatMasker v4.08 (Smit et al., 2015), and gene content was annotated with MAKER

v2.31.10 (Cantarel et al., 2008).

Sequencing data filtering and genotyping. We used the program bbduk (Bushnell, 2014) to

quality filter raw sequencing data. We then use BWA v0.7.17 (Li and Durbin, 2009) to align

filtered reads to the *Certhia americana* reference genome. Next, we used samtools v1.4.1 (Li et

al., 2009) to convert the BWA output SAM file to BAM format, followed by using the Genome

Analysis Toolkit (GATK) v4.1.0.0 (McKenna et al., 2010) to clean, sort, add read groups to, and

remove duplicates from BAM files. We used the samtools 'depth' command to measure

sequencing coverage across the reference genome for each individual (Fig. S1, Table S1).

We then used GATK in three steps to genotype all individuals: we used the

'HaplotypeCaller' function to call preliminary genotypes for each individual, built a database for

all samples using the 'GenomicsDBImport' function, and lastly used the 'GenotypeGVCFs'

function to group genotype all individuals together for both variant and invariant sites. We

filtered the genotype data using VCFtools v0.1.14 (Danecek et al., 2011) with the following

restrictions: (1) minimum site quality of 20, (2) minimum genotype quality of 20, (3) minimum

depth of coverage of 6, (4) maximum mean depth of coverage of 50, and (5) removal of indels.

For downstream analyses, we used various settings for (1) amount of missing data allowed, (2)

whether the outgroup was included, (3) inclusion or exclusion of invariant sites, and (4) thinning

between sites (see Table 1 for exact dataset characteristics used for all seven datasets used for

analyses). Although it has been suggested that variation in minor allele frequency (MAF) filtering

may impact estimates of genetic structure (Linck and Battey, 2019), we did not filter for a

minimum MAF because we previously showed that changing MAF did not significantly impact

pairwise F_{ST} calculations in *C. americana* (Manthey et al., 2021).

Phylogenomics. We estimated "gene trees" in non-overlapping windows (window size 50 kbp

and 100 kbp; datasets C and D in Table 1) using RaxML v8.2.12 (Stamatakis, 2014) with the

GTRCAT model of sequence evolution. For a window to be included, we required a minimum of

10 kbp in the alignment following filtering. This resulted in 19,639 phylogenies for the 50 kbp

sliding windows and 9846 phylogenies for the 100 kbp sliding windows. We summarized the

best-supported trees using the sumtrees.py script, part of the DendroPy Python package

(Sukumaran and Holder, 2010). Lastly, we used ASTRAL III v 5.6.3 (Zhang et al., 2018) to

calculate a species tree from all gene trees, using quartet frequencies as a measure of local

support (Sayyari and Mirarab, 2016).

For each gene tree, we estimated the genealogical sorting index (GSI) (Cummings et al.,

2008) for (1) each of the seven populations, and (2) each of the two main lineages (i.e., north

and south shaded blue and red, respectively, in Fig. 1). The GSI statistic uses rooted gene trees

to quantify lineage sorting; the GSI quantifies exclusive ancestry of individuals in labelled groups

by dividing the minimum number of nodes to produce monophyly for a defined group divided by

the observed number of nodes necessary to connect all individuals of a group in a phylogeny.

The maximum GSI value is one and occurs when a defined group is monophyletic. We

calculated the GSI statistic for each sampled population and for each of the two main lineages

(Fig. 1). We expect that each sampled population will show evidence of phylogenetic clustering,

regardless of position in the genome. As such, GSI at the population level may have little

variation tied to genomic architecture. In contrast, which populations group with which other

populations are likely to be influenced by rates of lineage sorting and are therefore likely

influenced by genomic architecture. As such, we may expect GSI measures at the lineage level

(i.e., two main clades) to be impacted by genomic architecture.

To obtain a relative rate of molecular evolution for each population per genomic window,

we calculated root-to-tip distances (RTD) in our phylogenies. For each individual across every

gene tree, we calculated the RTD using the R package ape (Paradis et al., 2004). As a

summary of these RTDs, we calculated the mean RTD per population relative to the maximum

RTD value for each gene tree (i.e., this relative statistic's range = 0 − 1). We then measured the

relationships between harmonic mean effective population sizes (mean per population) and

relative RTD in regions with low recombination and high recombination, as well as the shift in

values between high and low recombination regions. We also estimated relationships between

these characteristics while accounting for the evolutionary relationships of the samples using

phylogenetic independent contrasts (PICs) calculated in the R package ape (Paradis et al.,

2004).

Genetic structure. We estimated genetic structure using the program ADMIXTURE v1.3.0

(Alexander et al., 2009). Here, we used genome-wide SNPs thinned to either a minimum of 50

kbp or 100 kbp distance between SNPs (datasets A and B in Table 1). We ran ADMIXTURE

with an assumed number of genetic clusters equal to two or seven (i.e., $k = 2$ and $k = 7$) based

on our knowledge of phylogeographic structure in this species from previous studies (Manthey

et al., 2011a, b; Manthey et al., 2015). To assess how genetic structure estimation varied across

the genome, we also ran ADMIXTURE for SNP datasets in non-overlapping sliding windows of

50 kbp and 100 kbp (datasets E and F in Table 1). We compared the genome-wide

ADMIXTURE results to those in each window by summing the differences in group membership

for each individual. In other words, this deviation value ranges from zero to nearly one, where a

value of zero is identical group assignment for all individuals with the same admixture

coefficients for each genetic cluster, and higher values indicate more dissimilar matrices.

We also estimated variation in genetic structure across the genome using LOSTRUCT

(Li and Ralph, 2019). This method runs in three steps: (1) reducing the dimensionality of the

data using principal component analysis (PCA) for each genomic window, (2) find distances

between PCA maps, and (3) using multidimensional scaling (MDS) to display variation in PCA-

based genetic structure across the genome. We used LOSTRUCT in sliding windows of 50 kbp

and 100 kbp (datasets E and F in Table 1). We used variation in MDS axis one (hereafter

MDS1) to describe variation in these PCA-based estimates of genetic structure across the

genome.

We expect that genomic regions with low recombination will exhibit strong

phylogeographic structure similar to genome-wide patterns (e.g., Fig. 1). In contrast, we expect

that genomic regions with high recombination will either exhibit little genetic structure or

alternative patterns of genetic structure. As such, we expect that variation in ADMIXTURE and

PCA results will covary with genomic architecture.

Genetic diversity. Across 50 kbp and 100 kbp sliding windows (datasets C and D in Table 1),

we measured observed heterozygosity per individual as a measure of genetic diversity.

Population genomic correlations. We used the R package Hmisc (Harrell and Dupont, 2020)

to estimate both Pearson and Spearman correlation coefficients between population genomic

summary statistics and characteristics of genomic architecture in sliding windows. We used the

R package corrplot (Wei et al., 2017) to visualize correlations between all summary statistics.

Lastly, we used variance partitioning, implemented in the R package vegan (Oksanen et al.,

2007), to assess the proportion of variance in phylogeographic structure estimates explained by

genomic architecture.

Population demographic history. We used the program MSMC2 v1.1.0 (Schiffels and Durbin,

2014) to estimate demographic history for each individual. For use in MSMC, we masked

regions of the genome not genotyped due to low coverage or low genotype quality scores.

Additionally, we did not include the sex chromosomes in demography calculations. We ran

MSMC for each individual allowing up to 20 iterations (default setting) and used up to 23

inferred distinct time segments because this setting worked well to reduce spurious or

inconsistent results in other songbirds that we have studied (Manthey et al., 2022). We

performed 10 bootstraps for each demography estimate, using 1 Mbp bootstrapped segments of

the genome, to assess how demographic signal varies when subsetting parts of the genome.

Because the output of MSMC is interpreted relative to assumed mutation rates and generation

times, we used the Brown Creeper genome mutation rate estimate of 2.506×10^{-9} substitutions

per site per year (Manthey et al., 2021). Because there are no published estimates of Brown

Creeper generation times, we used a proxy generation time of double the age of sexual maturity

(Nadachowska-Brzyska et al., 2015) using maturity estimates from the Animal Aging and

Longevity Database (age of maturity = one year) (Tacutu et al., 2017). With the MSMC

demographic model output, we estimated the harmonic mean population sizes for each

population by using the mean MSMC population size estimates for each individual in 1000-year

discrete time intervals over the past 200 kya.

Code availability. All computer code used for analyses and figure creation (applicable for some

figures) for this project is available here: github.com/jdmanthey/certhia_phylogeography. We

used the following R packages for figure creation and file manipulation that were not cited in

other parts of the methods section: Biostrings (Pagès et al., 2017), ggplot2 (Wickham, 2011),

palettetown (Lucas, 2016), phytools (Revell, 2012), and RcolorBrewer (Neuwirth, 2014).

RESULTS

Genome-wide phylogeographic structure and demography. We used genomic sequencing

data from 21 ingroup individuals sequenced at ~18-29× genomic coverage to estimate genome-

wide patterns of phylogeographic structure. We corroborated previous genetic work (that used

few to thousands of genetic markers) by identifying hierarchical and strong phylogeographic

structure among sampled populations (Manthey et al., 2011a, b; Manthey et al., 2015). The

deepest phylogenomic split separates northern from southern populations and there is

additional support for distinctiveness of each of the regionally sampled populations (Fig. 1C).

Genome-wide ADMIXTURE results were consistent across different thinning strategies (i.e., 50

kbp and 100 kbp thinning; only 50 kbp results plotted) and showed patterns of genetic structure

that aligned well with the species tree (Fig. 1C). Demographic history estimates for each

individual were consistent within populations, but each population exhibited a distinct

demographic history (Fig. 1D). In the Pacific and Central American localities, population sizes

have fluctuated somewhat, but generally stay below an N_E of 200,000 over the past 100 ky (Fig.

1D). The Sierra Madre Oriental population showed evidence for a sharp decline in N_E over the

past 50 ky, shifting from an $N_E \sim$ 200,000 at about 50 kya to an $N_E \sim$ 10,000 in the past ten kya

(Fig. 1D). The Rocky Mountains population has remained relatively stable over the past 100 ky

with an $N_E \sim$ 200,000 (Fig. 1D). Lastly, the Eastern North America and Central Mexico

populations have exhibited fluctuations and much larger N_E than all other populations over the

past 100 ky (Fig. 1D). Harmonic mean effective population sizes over the past 200 ky ranged

from ~85,000 (Central America South) to ~320,000 (Central Mexico) (Fig. 1D). Harmonic mean

effective population size estimates over the past 200 ky are highly correlated with observed

heterozygosity estimates for each individual (r = 0.962).

Genetic diversity. Genetic diversity in each individual varied widely across the genome, with

mean observed heterozygosity values for each individual between ~0.00165 and 0.00463 (Fig

1B). Observed heterozygosity was strongly correlated across the genome in all pairwise

comparisons of individuals (mean r = 0.409, range = 0.189 – 0.853; all p << 0.001). Notably, the

Sierra Madre Oriental, Pacific, and Central American localities had more windows with no

heterozygosity than the other populations (Fig. 1B).

Phylogeographic structure variation across the genome. We estimated phylogeographic

structure across the genome using PCA, ADMIXTURE, and the GSI metric from phylogenies; all

estimates of phylogeographic structure varied across the genome, with the outliers deviating the

most from genome-wide patterns clustering mostly on small chromosomes and the ends of

large chromosomes (Fig. 2; Fig. S2). The most extreme deviations from genome-wide

phylogeographic structure appear to cluster the Central Mexico population with the northern

lineage (Fig. S3), and in some cases also cluster the Sierra Madre Oriental population with the

northern lineage (Fig. S3; Fig. 3). As an example, we plot the phylogeographic structure for one

of these outlier windows in Fig. 3, demonstrating the clustering of the Sierra Madre Oriental and

Central Mexico populations with the northern lineage.

Correlations between phylogeographic structure estimates and genomic architecture. We

found a strong correlation between different measures of phylogeographic structure, including

GSI at the lineage level, MDS1 of the PCA, and ADMIXTURE deviations (all r ≥ 0.58; p <<<

0.001; Fig. 4; Fig. S4). In contrast, GSI at the population level did not vary consistently with the

other metrics (all r between -0.21 and -0.03), likely because the population-level GSI usually

showed similar patterns across the genome (i.e., individuals generally clustered with individuals

sampled from the same population).

Phylogeographic structure patterns most different from genome-wide patterns were

strongly positively correlated with recombination rate, GC content, and genetic diversity (Fig. 4;

Fig. S4). In contrast, TE content had no or weak correlations with all other statistics that we

calculated (Fig. 4; Fig. S4). Correlations were consistent between 50 kbp and 100 kbp

estimates, with no correlations deviating more than ~0.08 when estimated with different window

sizes (Fig. S4).

Using variance partitioning, recombination rate variation most strongly explained

deviations in phylogeographic structure, but interactions of recombination rate, GC content,

CDS content, and TE content also explained some portions of the variance (Table 2). Most

notably, genomic architecture explained approximately 60% of the variance in both (1) the

MDS1 axis explaining differences in PCA-based estimates of phylogeographic structure and (2)

ADMIXTURE results at the lineage level with an assumed k = 2 (Table 2).

The relationship between recombination rate and phylogeographic structure is

exemplified when comparing the genomic windows with the highest or lowest values of

recombination rate; estimates of phylogeographic structure and genetic diversity have little to no

overlap in statistic distributions in high recombination regions versus low recombination regions

(Fig. 5). These patterns hold regardless of whether we are looking at autosomes or the Z

chromosome (Fig. 5).

Because the effective population sizes for each locality varied substantially (Fig. 1; Fig.

S5), we looked to see if there was a relationship between relative RTD and population size in

regions with very different recombination rates. Using both the raw measurements and PICs, we

found a negative association between relative rates of molecular evolution and harmonic mean

effective population sizes (Fig. 6; Fig. S6). Notably, there was also a strong positive association

between effective population sizes and shifts in relative rates of molecular evolution between

high and low recombination regions (Fig. 6).

DISCUSSION

Using whole-genome sequencing for seven populations of the Brown Creeper, we aimed to

decipher how genome architecture influences phylogeographic structure variation across the

genome. We found that recombination rate and other characteristics of the genome explain 6-

68% of the variance in phylogeographic structure inference using different metrics and genomic

sliding window sizes (Table 2).

Phylogeographic structure shaped by genomic architecture.

We used genetic clustering, ordination, and phylogenetic methods to assess phylogeographic

structure in genomic sliding windows. We found that recombination rate variation was the

strongest predictor of phylogeographic structure variation across the genome (Table 2).

Consistent with our expectation, phylogeographic signal best representing the bifurcating

evolutionary history of the Brown Creeper was identified in genomic regions with relatively low

recombination rate; this was consistent across all methods used (Fig. 2). Notably, the genomic

windows with the lowest recombination rates deviated little from the genome-wide patterns (Fig.

5).

Previous work in insects, birds, and mammals has shown that phylogenomic signal

varies across the genome (Edelman et al., 2019; Fontaine et al., 2015; Li et al., 2019; Martin et

al., 2019; Thom et al., 2024). Generally, these studies have found or suggested that the

background species tree is best represented by phylogenomic patterns exhibited in genomic

regions with low recombination, and that gene flow among taxa is more prevalent in genomic

regions with high recombination rate (Edelman et al., 2019; Li et al., 2019; Martin et al., 2019;

Thom et al., 2024). Our results in the Brown Creeper echo these findings *at a shallow*

evolutionary timescale, whereby we find highest support for the genome-wide supported

phylogeographic relationships in genomic regions with low recombination rate (Fig. 2; Fig. S2).

We may interpret these patterns in the context of lineage sorting during the speciation process,

whereby genomic regions with low introgression exhibit quicker polymorphism fixation due to

relatively increased effects of linked selection and genetic drift and reduced homogenization

among lineages due to gene flow.

Genomic studies investigating intraspecific population and phylogeographic structure

have more commonly used PCA-based methods (e.g., LOSTRUCT) to investigate variation in

genomic structure across the genome. In two species of woodpeckers, a small portion of the

variance in PCA-based phylogeographic structure was found to be associated with local

recombination rate (Moreira et al., 2023). We found that variation in phylogeographic structure

across the genome of the Brown Creeper—inferred using both ordination and genetic clustering

methods—was strongly associated with recombination rate variation (Fig. 2; Fig. S2; Table 2).

We may infer that the same interactions between recombination rate variation and population

genomics processes shaping variance in phylogenomic signal are also shaping variance in

clustering- and ordination-based estimates of phylogeographic structure.

In contrast to generalizations about low recombination regions often reflecting a taxon's

background evolutionary history, in some scenarios we may expect regions of extremely low

recombination to greatly differ from genome-wide signatures of population genetic structure. In

several taxa, including sunflowers, flies, fishes, and birds, the greatest deviations from the

genome-wide pattern of intraspecific population structure (inferred with PCA) have been

identified in genomic regions with extremely low recombination rates that are often inferred to be

polymorphic inversions within populations or species (Hale et al., 2021; Huang et al., 2020; Li

and Ralph, 2019; Mérot et al., 2021; Perrier et al., 2020; Shi et al., 2021; Todesco et al., 2020;

Whiting et al., 2021). In these cases, if a taxon has relatively little population structure but also

exhibits polymorphic inversions, we may expect the biggest deviations in population genetic

structure from genome-wide patterns to reflect these polymorphic inversions that exhibit

suppressed recombination. These cases present interesting patterns to think about when

interpreting genomic phylogeography data. Genomic regions with low recombination may best

represent a taxon's evolutionary history, excepting times when these regions are associated

with large structural variants.

Interaction of recombination rate, population demography, and rate of evolution.

While we may expect that the interactions between recombination rate and population genomics

processes will impact the rate of evolution across the genome, we did not simply observe little

phylogeographic structure in genomic regions with high recombination rates and strong

phylogeographic structure in genomic regions with low recombination rates. Conversely, we

found strong—and differing—patterns of phylogeographic structure across the genome (Fig. 1;

Fig. 3; Fig. S3). Because these differing patterns of phylogeographic structure generally

separated the Central American populations and sometimes the Sierra Madre Oriental

population from all others, we hypothesized that populations with relatively smaller N_E exhibit

relatively higher evolutionary rates in genomic regions with high recombination.

Using relative RTD measures as a relative rate of molecular evolution for each

population, we found that larger populations exhibited reduced rates of molecular evolution in

genomic windows with high recombination rates (Fig. 6; Fig. S6) and that shifts in relative rates

of molecular evolution in high versus low recombination genomic regions were positively

associated with effective population sizes (Fig. 6; Fig. S6). This is consistent with simulations by

Tigano and colleagues (2021) where the authors showed the interplay between recombination

rate and selection causes greater genomic variance in evolutionary rates in larger populations

relative to more consistent evolutionary rates across the genome in smaller populations.

We suspect that molecular evolution in genomic regions with high recombination rates is

a product of complex interactions between population demography and the relative strengths of

linked selection and genetic drift. In the Brown Creeper genome, recombination rate variation is

positively associated with gene density (Fig. 2; Fig. 4) and we may interpret gene-dense

genomic regions as targets for natural selection. Indeed, in a previous study in the Brown

Creeper (Manthey et al., 2021), we showed that the two main Brown Creeper lineages exhibited

relatively less neutral evolution in the gene-dense microchromosomes relative to the larger

macrochromosomes. In gene-dense and high recombination regions under the effects of

purifying selection, we may expect populations with relatively smaller N_E to accumulate

substitutions faster than populations with higher N_E (Lanfear et al., 2014), consistent with the

trends we identified here in the Brown Creeper (Fig. 6; Fig. S6).

Implications for future phylogeographic studies

If phylogeographic structure varies widely across the genome, what are realistic and best-

practice approaches for future phylogeographic work? Are studies that use few genes or

reduced representation genomic datasets doomed to be wrong from the start? We expect that

strong phylogeographic structure will be detected even with few genetic markers. For example,

more than ten years ago we used 20 genetic markers obtained with Sanger sequencing to study

the phylogeography of the Brown Creeper (Manthey et al., 2011b) and found the same general

phylogeographic trends observed here with our whole-genome dataset. In contrast, any study

where most of the genetic markers are taken from genomic regions where the pattern differs

from a taxon's true evolutionary history (e.g., high recombination regions in the Brown Creeper

case) may run into problems inferring that true history. Additionally, in any taxa with weak

population genetic structure or widespread and rampant gene flow, a whole-genome method

may be the only approach to identify the few genomic regions differentiating taxa (e.g., Toews et

al., 2016). Overall, we suggest using the most amount of genetic data feasible for future

phylogeographic studies given sources of biological material (e.g., quality of preserved tissue)

and funding available, as well as investigating the patterns and potential causes of population

genetic structure variation across the genome when possible.

CONCLUSIONS

We used whole genome sequencing to estimate how genomic architecture shapes variation in

phylogeographic structure across the genome of the Brown Creeper. We found that

phylogeographic structure—as measured using ordination, phylogenetic, and clustering

methods—was strongly associated with regional genomic variation in recombination rates. In

low recombination regions, we recovered phylogeographic structure concordant with genome-

wide patterns with all three types of methods. The most divergent phylogeographic patterns

were in high recombination regions; populations with small effective population sizes were

distinct from all other populations due to relatively faster evolution than larger populations in

these high recombination regions. Because these high recombination regions are rich in coding

genes, we hypothesize that large populations have higher relative effects of purifying selection

in these regions, and overall slower relative molecular evolutionary rates compared to smaller

populations. Overall, our results show that phylogeographic structure may vary widely across

the genome, and that effective population sizes of sampled populations and genomic

architecture and their interactions will impact regional genomic variation in phylogeographic

structure.

ACKNOWLEDGEMENTS

We would like to thank the collection managers, curators, and contributors for generous tissue

loans for all individuals used in this study from the following museums: Burke Museum of

Natural History and Culture, and University of Kansas Biodiversity Institute. Sequencing was

supported by Texas Tech University start-up funding to JDM and a generous donation from the

Ferguson family to GMS and the Denver Museum of Nature and Science. Support was also

provided by NSF awards #1953688 to JDM and #0814841 to GMS. The High-Performance

Computing Center at TTU supported computational analyses.

DATA ACCESSIBILITY

484　All raw resequencing data uploaded to NCBI's SRA database under BioProject PRJNA1068991.

485　All code used for analysis on this project is available on GitHub:

486　(github.com/jdmanthey/certhia_phylogeography).

AUTHOR CONTRIBUTIONS

JDM and GMS designed the study. JDM performed laboratory and bioinformatic work and wrote

the first draft of the manuscript. JDM and GMS contributed to revising and improving the final

draft of the manuscript.

DECLARATION OF INTEREST

The authors declare no conflicts of interest.

 TABLES AND FIGURES

Figure 1. Sampling map, genetic diversity, phylogeographic structure, and demography. (A)
Sampling localities for this study. Colors for localities are consistent across plots. (B) Violin plots
of observed heterozygosity across all 50 kbp sliding windows. White circles indicate means. (C)
Consensus phylogeny using 19,639 gene trees from the 50 kbp windows. Support values are
labelled for all branches showing the highest support from all estimation methods (N = 4),
including consensus and ASTRAL phylogenies for both the 50 kbp and 100 kbp window
datasets. At tips of phylogeny are ADMIXTURE results from 19,602 genome-wide SNPs
(thinned every 50 kbp) for either two or seven assumed genetic clusters (k = 2 or k = 7,
respectively). (D) Demographic history estimated with MSMC2. Different individuals are
represented with differently colored lines and bootstraps are shown with thin lines. In (D), HM =
harmonic mean. Brown Creeper photo in (A) by JDM from the Chiricahua Mountains in southern
Arizona.

Figure 2. Genomic architecture and phylogeographic structure variation across the genome in 50kbp sliding windows. Gray points indicate window values and dark gray lines indicate means across 20 windows (i.e., 1 Mbp windows). Blue indicates top 2.5% of outliers and orange indicates bottom 2.5% of outliers for each statistic. Bottom outliers not shown for CDS, ADMIX., and GSI as these each have large proportions of values equal to zero. Abbreviations: mean effective recombination rate of the two main lineages (ρ), gene content (CDS), transposable element and repetitive DNA content (TE), multidimensional scaling axis one from principal components analyses (-MDS1), mean genealogical sorting index for the two main *Certhia* lineages (1 - GSI), ADMIXTURE deviation relative to genome-wide analysis (ADMIX.), and mean genetic diversity across individuals as measured using observed heterozygosity (H_O). MDS1 is plotted as negative and GSI is plotted as (1 - GSI) so that deviations from expected phylogeographic structure for MDS1, GSI, and ADMIXTURE are all represented as higher values.

548 Figure 3. Example phylogeographic structure in a window that deviates from genome-wide
549 patterns. This window had the 15th lowest MDS1 value of all windows from the principal
550 components analysis (PCA) across the genome. (A) PCA, (B) RAxML phylogeny, and (C)
551 ADMIXTURE results for a window on chromosome 4.

Figure 4. Correlation plot of genomic architecture, genomic diversity, and phylogeographic structure for 50 kbp windowed statistics. Pearson's and Spearman's r coefficients are plotted above and below the diagonal, respectively. Abbreviations shown in plot: mean effective recombination rate of the two main lineages (ρ); GC content (GC%); gene content (CDS%); transposable element content (TE%); mean observed heterozygosity (H_O); genealogical sorting index at the population [1 - GSI (Pop.)] and lineage [1 - GSI (Lin.)] levels; negative multidimensional scaling axis one from principal components analyses (-MDS1); ADMIXTURE deviation relative to genome-wide analysis for two [ADMIX. (k = 2)] or seven [ADMIX. (k = 7)] assumed genetic clusters.

587 Figure 5. Violin plots of phylogeographic structure variation in genomic regions with very high or
588 low recombination rates, separated for autosomal (A) and Z-linked (Z) genomic regions. 50 kbp
589 windows with the highest or lowest 2.5% of recombination rate values are included. Higher
590 values for ADMIXTURE deviations, 1 - GSI, and -MDS1 indicate values deviating relatively
591 more from the genome-wide patterns of phylogeographic structure.
592

Figure 6. Relationship between effective population sizes and rate of evolution (measured by

relative root-to-tip distance for each population) in genomic regions that vary in recombination

rate. Extreme low and high recombination (rho) regions represent the quintiles of the genome

with the lowest or highest recombination rates (i.e., highest 20% and lowest 20% of windows).

Results for raw data are shown in (A-C) and for phylogenetic independent contrasts (PIC) in (D-

F). The right panels in each row (C + F) show the shift in relative rates of evolution between

high and low recombination genomic regions.

631 Table 1. Different datasets used in this study and their characteristics. For inclusion, all datasets
632 required sites to have a minimum quality score of 20, minimum of 20 genotype quality, a
633 minimum depth per individual for a site to be called of 6, and a max mean depth across
634 individuals of 50.

Set	Missing Allowed	OG[1]	Window size	# Windows	Min. Sep. per SNP[2]	# SNPs	# Sites	Analyses
A	0%		-	-	50 kbp	19,602	19,602	Genome-wide ADMIXTURE
B	0%		-	-	100 kbp	9,838	9,838	Genome-wide ADMIXTURE
C	20%	X	50 kbp	19,713	-	35,463,408	890,335,872	RAxML, GSI, H_O
D	20%	X	100 kbp	9,862	-	35,463,408	890,335,872	RaxML, GSI, H_O
E	0%		-	-	-	21,657,220	21,657,220	LOSTRUCT
F	0%		50 kbp	19,697	-	21,657,220	21,657,220	ADMIXTURE per window
G	0%		100 kbp	9,860	-	21,657,220	21,657,220	ADMIXTURE per window
H	0%		-	-	-	*	*	MSMC2

[1]OG = outgroup included

[2]Min. Sep. per SNP = minimum separation between SNPs (i.e., thinning)

*SNPs and sites not labeled for MSMC2 analyses because these values varied per individual. For each individual, no missing data was allowed for that particular individual's MSMC2 demographic analyses input files. The MSMC2 dataset did not include the Z chromosome.

638 Table 2. Variance partitioning where the explanatory variables are characteristics of genomic
639 architecture and response variables are different measures of population genetic structure.
640 Values indicative of variance proportion explained by each explanatory variable or their
641 interactions.

50 kbp windows						
Measure of Genetic Structure	ρ	CDS	TE	GC	Interactions	Total
MDS1	0.23	0.00	0.01	0.04	0.33	0.61
GSI (Lineages)	0.16	0.00	0.00	0.02	0.22	0.40
GSI (Populations)	0.02	0.00	0.00	0.01	0.03	0.06
ADMIXTURE (k = 2)	0.24	0.00	0.00	0.03	0.30	0.57
ADMIXTURE (k = 7)	0.19	0.00	0.00	0.01	0.18	0.38
100 kbp windows						
Measure of Genetic Structure	ρ	CDS	TE	GC	Interactions	Total
MDS1	0.22	0.01	0.01	0.03	0.41	0.68
GSI (Lineages)	0.16	0.00	0.00	0.01	0.23	0.40
GSI (Populations)	0.02	0.00	0.00	0.01	0.05	0.08
ADMIXTURE (k = 2)	0.25	0.00	0.00	0.02	0.35	0.62
ADMIXTURE (k = 7)	0.20	0.00	0.00	0.01	0.22	0.43

SUPPLEMENTAL TABLES AND FIGURES

Table S1. Sampling metadata and characteristics of sequencing amount and coverage when aligned to the reference genome. Samples from Utah, Morelos, Jalisco, and the outgroup were retrieved from the NCBI SRA and originally sequenced for a previous study by Manthey and colleagues (2021).

Museum #	Lineage	Locality	Lat.	Long.	Raw bp	Filtered bp	Coverage
UWBM103417	South	Copan, HN	14.866	-89.050	40,501,485,458	34,714,897,849	22.79
UWBM105618	South	Copan, HN	14.866	-89.050	30,149,924,172	26,420,616,801	18.55
UWBM93696	South	Copan, HN	14.866	-89.050	40,591,948,048	36,089,400,965	26.06
UWBM113447	South	Chiapas, MX	16.691	-92.606	37,741,499,908	33,063,229,986	23.32
UWBM113461	South	Chiapas, MX	16.691	-92.606	32,696,721,580	27,488,157,228	18.51
UWBM113462	South	Chiapas, MX	16.691	-92.606	48,519,014,074	42,841,474,607	27.09
UWBM106936	South	Morelos, MX	19.0873	-99.194	40,136,994,108	35,563,542,819	26.35
UWBM107890	South	Jalisco, MX	21.882	-103.865	43,601,103,228	38,563,138,625	26.14
UWBM107897	South	Jalisco, MX	21.882	-103.865	30,536,607,690	27,038,872,945	18.55
UWBM106768	South	Nuevo Leon, MX	24.872	-100.224	42,330,853,397	37,109,282,807	27.62
UWBM111691	South	Nuevo Leon, MX	24.872	-100.224	41,523,724,956	36,804,470,356	26.21
UWBM111708	South	Nuevo Leon, MX	24.872	-100.224	43,392,483,440	36,820,385,840	27.04
UWBM113198	North	California, US	36.245	-121.700	40,117,812,880	35,379,027,102	26.26
UWBM112794	North	California, US	36.245	-121.553	37,692,762,846	33,128,089,974	23.33
UWBM112795	North	California, US	36.245	-121.553	44,376,620,806	38,809,177,747	28.71
UWBM113162	North	Utah, US	37.317	-113.455	40,934,172,636	36,217,919,293	26.25
UWBM113168	North	Utah, US	37.317	-113.455	39,295,062,200	34,925,676,137	25.72
UWBM113167	North	Utah, US	37.377	-113.467	42,535,860,138	37,425,027,332	27.55
UWBM107061	North	West Virginia, US	38.289	-79.937	43,886,484,470	38,699,509,454	28.65
UWBM112055	North	West Virginia, US	38.289	-79.937	47,972,794,224	40,458,960,409	29.36
UWBM112054	North	West Virginia, US	38.641	-79.838	39,431,412,482	34,774,499,050	25.96
KU92846	Outgroup	England			21,417,629,506	19,195,682,239	13.58

659   Figure S1. Sequencing coverage aligned to the reference genome for each individual. The

660   vertical red lines indicate the mean coverage per individual.

Figure S2. Genomic architecture and phylogeographic structure variation across the genome in
100kbp sliding windows. Gray points indicate window values and dark gray lines indicate means
across ten windows (i.e., 1 Mbp windows). Blue indicates top 2.5% of outliers and orange
indicates bottom 2.5% of outliers for each statistic. Bottom outliers not shown for CDS, ADMIX.,
and GSI as these each have large proportions of values equal to zero. Abbreviations: mean
effective recombination rate of the two main lineages (ρ), gene content (CDS), transposable
element and repetitive DNA content (TE), multidimensional scaling axis one from principal
components analyses (-MDS1), mean genealogical sorting index for the two main *Certhia*
lineages (1 - GSI), ADMIXTURE deviation relative to genome-wide analysis (ADMIX.), and
mean genetic diversity across individuals as measured using observed heterozygosity (H_O).
MDS1 is plotted as negative and GSI is plotted as (1 - GSI) so that deviations from expected
phylogeographic structure for MDS1, GSI, and ADMIXTURE are all represented as higher
values.

Figure S3. The 15 most extreme outlier principal components analysis (PCA) windows identified

from those windows with low MDS dimension one values.

Figure S4. Correlation plot of genomic architecture, genomic diversity, and phylogeographic
structure for (A) 50 kbp and (B) 100 kbp windowed statistics. Pearson's and Spearman's r
coefficients are plotted above and below the diagonal, respectively. In (C), the absolute value
difference between correlations in 50 kbp and 100 kbp windows. Abbreviations shown in plot:
mean effective recombination rate of the two main lineages (ρ); GC content (GC%); gene
content (CDS%); transposable element content (TE%); mean observed heterozygosity (H_O);
genealogical sorting index at the population [1 - GSI (Pop.)] and lineage [1 - GSI (Lin.)] levels;
negative multidimensional scaling axis one from principal components analyses (-MDS1);
ADMIXTURE deviation relative to genome-wide analysis for two [ADMIX. (k = 2)] or seven
[ADMIX. (k = 7)] assumed genetic clusters.

Figure S5. Demographic history estimated with MSMC2 for each individual. Thick lines indicate

the whole genome estimate and thin lines indicate results from ten bootstrap replicates. Sample

numbers correspond with those reported in Table S1.

Figure S6. Violin plots of evolutionary rates (as measured by relative root-to-tip distances) for

each population in the upper and lower genomic quintiles of high or low recombination (rho)

values.

LITERATURE CITED

Alexander, D.H., Novembre, J., Lange, K., 2009. Fast model-based estimation of ancestry in
unrelated individuals. Genome Res. 19, 1655-1664.
Allmon, W.D., 1992. A causal analysis of stages in allopatric speciation. Oxford Surveys in
Evolutionary Biology 8, 219-219.
Bushnell, B., 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner. Ernest Orlando Lawrence
Berkeley National Laboratory, Berkeley, CA (US).
Cantarel, B.L., Korf, I., Robb, S.M., Parra, G., Ross, E., Moore, B., Holt, C., Alvarado, A.S.,
Yandell, M., 2008. MAKER: an easy-to-use annotation pipeline designed for emerging
model organism genomes. Genome Res. 18, 188-196.
Cruickshank, T.E., Hahn, M.W., 2014. Reanalysis suggests that genomic islands of speciation
are due to reduced diversity, not reduced gene flow. Mol. Ecol. 23, 3133-3157.
Cummings, M.P., Neel, M.C., Shaw, K.L., 2008. A genealogical approach to quantifying lineage
divergence. Evolution 62, 2411-2422.
Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E.,
Lunter, G., Marth, G.T., Sherry, S.T., 2011. The variant call format and VCFtools.
Bioinformatics 27, 2156-2158.
Dutoit, L., Burri, R., Nater, A., Mugal, C.F., Ellegren, H., 2017. Genomic distribution and
estimation of nucleotide diversity in natural populations: perspectives from the collared
flycatcher (Ficedula albicollis) genome. Mol. Ecol. Res. 17, 586-597.
Edelman, N.B., Frandsen, P.B., Miyagi, M., Clavijo, B., Davey, J., Dikow, R.B., García-
Accinelli, G., Van Belleghem, S.M., Patterson, N., Neafsey, D.E., 2019. Genomic
architecture and introgression shape a butterfly radiation. Science 366, 594-599.
Ellegren, H., 2010. Evolutionary stasis: the stable chromosomes of birds. Trends Ecol. Evol. 25,
283-291.
Ellegren, H., Smeds, L., Burri, R., Olason, P.I., Backström, N., Kawakami, T., Künstner, A.,
Mäkinen, H., Nadachowska-Brzyska, K., Qvarnström, A., 2012. The genomic landscape
of species divergence in Ficedula flycatchers. Nature 491, 756.
Fontaine, M.C., Pease, J.B., Steele, A., Waterhouse, R.M., Neafsey, D.E., Sharakhov, I.V., Jiang,
X., Hall, A.B., Catteruccia, F., Kakani, E., 2015. Extensive introgression in a malaria
vector species complex revealed by phylogenomics. Science 347, 1258524.
Haenel, Q., Laurentino, T.G., Roesti, M., Berner, D., 2018. Meta☐analysis of
chromosome☐scale crossover rate variation in eukaryotes and its significance to
evolutionary genomics. Mol. Ecol. 27, 2477-2497.
Hale, M.C., Campbell, M.A., McKinney, G.J., 2021. A candidate chromosome inversion in
Arctic charr (*Salvelinus alpinus*) identified by population genetic analysis techniques. G3
11, jkab267.
Harrell, F.E., Dupont, C., 2020. Hmisc: Harrell Miscellaneous. R, R.
Hey, J., Kliman, R.M., 2002. Interactions between natural selection, recombination and gene
density in the genes of Drosophila. Genetics 160, 595-608.
Huang, K., Andrew, R.L., Owens, G.L., Ostevik, K.L., Rieseberg, L.H., 2020. Multiple
chromosomal inversions contribute to adaptive divergence of a dune sunflower ecotype.
Mol. Ecol. 29, 2535-2549.

Kapusta, A., Suh, A., 2017. Evolution of bird genomes—a transposon's eye view. Ann. N. Y.
Acad. Sci. 1389, 164-185.

Kawakami, T., Smeds, L., Backström, N., Husby, A., Qvarnström, A., Mugal, C.F., Olason, P.,
Ellegren, H., 2014. A high density linkage map enables a second generation collared
flycatcher genome assembly and reveals the patterns of avian recombination rate
variation and chromosomal evolution. Mol. Ecol. 23, 4035-4058.

Koonin, E.V., 2009. Evolution of genome architecture. The International Journal of
Biochemistry & Cell Biology 41, 298-306.

Kulathinal, R.J., Bennett, S.M., Fitzpatrick, C.L., Noor, M.A., 2008. Fine-scale mapping of
recombination rate in *Drosophila* refines its correlation to diversity and divergence. Proc.
Natl. Acad. Sci. USA 105, 10051-10056.

Lanfear, R., Kokko, H., Eyre-Walker, A., 2014. Population size and the rate of evolution. Trends
Ecol. Evol. 29, 33-41.

Li, G., Figueiró, H.V., Eizirik, E., Murphy, W.J., 2019. Recombination-aware phylogenomics
reveals the structured genomic landscape of hybridizing cat species. Mol. Biol. Evol. 36,
2111-2126.

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows–Wheeler
transform. Bioinformatics 25, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., 2009. The sequence alignment/map format and SAMtools. Bioinformatics 25,
2078-2079.

Li, H., Ralph, P., 2019. Local PCA shows how the effect of population structure differs along the
genome. Genetics 211, 289-304.

Linck, E., Battey, C., 2019. Minor allele frequency thresholds strongly affect population
structure inference with genomic data sets. Mol. Ecol. Res. 19, 639-647.

Louis, M., Galimberti, M., Archer, F., Berrow, S., Brownlow, A., Fallon, R., Nykänen, M.,
O'brien, J., Robertson, K.M., Rosel, P.E., 2020. Selection on ancestral genetic variation
fuels parallel ecotype formation in bottlenose dolphins. hal-03020428.

Lucas, T., 2016. palettetown: Use Pokemon Inspired Colour Palettes. R package version 0.1.1.
https://CRAN.R-project.org/package=palettetown.

Manthey, J.D., Bourgeois, Y., Meheretu, Y., Boissinot, S., 2022. Varied diversification patterns
and distinct demographic trajectories in Ethiopian montane forest bird (Aves:
Passeriformes) populations separated by the Great Rift Valley. doi: 10.1111/mec.16417.
Mol. Ecol.

Manthey, J.D., Klicka, J., Spellman, G.M., 2011a. Cryptic diversity in a widespread North
American songbird: phylogeography of the Brown Creeper (Certhia americana). Mol.
Phylogen. Evol. 58, 502-512.

Manthey, J.D., Klicka, J., Spellman, G.M., 2011b. Isolation driven divergence: speciation in a
widespread North American songbird (Aves: Certhiidae). Mol. Ecol. 20, 4371-4384.

Manthey, J.D., Klicka, J., Spellman, G.M., 2015. Chromosomal patterns of diversity and
differentiation in creepers: a next-gen phylogeographic investigation of *Certhia*
americana. Heredity 115, 165-172.

Manthey, J.D., Klicka, J., Spellman, G.M., 2021. The genomic signature of allopatric speciation
in a songbird is shaped by genome architecture (Aves: *Certhia americana*). Genome Biol.
Evol. 13, evab120.

Martin, S.H., Davey, J.W., Salazar, C., Jiggins, C.D., 2019. Recombination rate variation shapes
barriers to introgression across butterfly genomes. PLoS Biol. 17, e2006288.
Mayr, E., Mayr, E., Mayr, E., Mayr, E., 1963. Animal species and evolution. Belknap Press of
Harvard University Press Cambridge, Massachusetts.
McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella,
K., Altshuler, D., Gabriel, S., Daly, M., 2010. The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. Genome
Res. 20, 1297-1303.
McVean, G., Auton, A., 2007. LDhat 2.1: A package for the population genetic analysis of
recombination. Department of Statistics, Oxford, OX1 3TG, UK.
Mérot, C., Berdan, E.L., Cayuela, H., Djambazian, H., Ferchaud, A.-L., Laporte, M.,
Normandeau, E., Ragoussis, J., Wellenreuther, M., Bernatchez, L., 2021. Locally
adaptive inversions modulate genetic variation at different geographic scales in a
seaweed fly. Mol. Biol. Evol. 38, 3953-3971.
Moreira, L.R., Klicka, J., Smith, B.T., 2023. Demography and linked selection interact to shape
the genomic landscape of codistributed woodpeckers during the Ice Age. Mol. Ecol.,
DOI: 10.1111/mec.16841.
Nachman, M.W., Payseur, B.A., 2012. Recombination rate variation and speciation: theoretical
predictions and empirical results from rabbits and mice. Philosophical Transactions of the
Royal Society B: Biological Sciences 367, 409-421.
Nadachowska-Brzyska, K., Li, C., Smeds, L., Zhang, G., Ellegren, H., 2015. Temporal dynamics
of avian populations during Pleistocene revealed by whole-genome sequences. Curr.
Biol. 25, 1375-1380.
Neuwirth, E., 2014. RColorBrewer: ColorBrewer Palettes. R package version 1.1-2.
https://CRAN.R-project.org/package=RColorBrewer.
Nosil, P., Feder, J.L., 2012. Genomic divergence during speciation: causes and consequences.
Philosophical Transactions of the Royal Society of London B: Biological Sciences 367,
332-342.
Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J., Suggests, M.,
2007. The vegan package. Community Ecology Package 10, 631-637.
Pagès, H., Aboyoun, P., Gentleman, R., DebRoy, S., 2017. Biostrings: Efficient manipulation of
biological strings. R Package Version 2.0.
Paradis, E., Claude, J., Strimmer, K., 2004. APE: analyses of phylogenetics and evolution in R
language. Bioinformatics 20, 289-290.
Perrier, C., Rougemont, Q., Charmantier, A., 2020. Demographic history and genomics of local
adaptation in blue tit populations. Evolutionary Applications 13, 1145-1165.
Poulin, J., D'Astous, É., Villard, M., Hejl, S., Newlon, K., McFadzen, M., Young, J., Ghalambor,
C., 2020. Brown Creeper (Certhia americana), version 1.0. In Birds of the World. . In:
Poole, A. (Ed.). Cornell Lab of Ornithology, Ithaca, NY, USA.
https://doi.org/10.2173/bow.brncre.01.
Ravinet, M., Yoshida, K., Shigenobu, S., Toyoda, A., Fujiyama, A., Kitano, J., 2018. The
genomic landscape at a late stage of stickleback speciation: High genomic divergence
interspersed by small localized regions of introgression. PLoS Genet. 14, e1007358.
Revell, L.J., 2012. phytools: an R package for phylogenetic comparative biology (and other
things). Methods Ecol. Evol. 3, 217-223.

Roesti, M., Moser, D., Berner, D., 2013. Recombination in the threespine stickleback genome—
patterns and consequences. Mol. Ecol. 22, 3014-3027.
Sayyari, E., Mirarab, S., 2016. Fast coalescent-based computation of local branch support from
quartet frequencies. Mol. Biol. Evol. 33, 1654-1668.
Schiffels, S., Durbin, R., 2014. Inferring human population size and separation history from
multiple genome sequences. Nat. Genet. 46, 919.
Shi, Y., Bouska, K.L., McKinney, G.J., Dokai, W., Bartels, A., McPhee, M.V., Larson, W.A.,
2021. Gene flow influences the genomic architecture of local adaptation in six riverine
fish species. BioRxiv, DOI: https://doi.org/10.1101/2021.1105.1118.444736.
Singhal, S., Leffler, E.M., Sannareddy, K., Turner, I., Venn, O., Hooper, D.M., Strand, A.I., Li,
Q., Raney, B., Balakrishnan, C.N., Griffith, S.C., McVean, G., Przeworski, M., 2015.
Stable recombination hotspots in birds. Science 350, 928-932.
Smit, A., Hubley, R., Green, P., 2015. RepeatMasker Open-4.0. 2013–2015. Institute for
Systems Biology. http://repeatmasker. org.
Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30, 1312-1313.
Stankowski, S., Chase, M.A., Fuiten, A.M., Rodrigues, M.F., Ralph, P.L., Streisfeld, M.A.,
2019. Widespread selection and gene flow shape the genomic landscape during a
radiation of monkeyflowers. PLoS Biol. 17, e3000391.
Sukumaran, J., Holder, M.T., 2010. DendroPy: a Python library for phylogenetic computing.
Bioinformatics 26, 1569-1571.
Tacutu, R., Thornton, D., Johnson, E., Budovsky, A., Barardo, D., Craig, T., Diana, E.,
Lehmann, G., Toren, D., Wang, J., 2017. Human Ageing Genomic Resources: new and
updated databases. Nucleic Acids Res. 46, D1083-D1090.
Thom, G., Moreira, L.R., Batista, R., Gehara, M., Aleixo, A., Smith, B.T., 2024. Genomic
architecture predicts tree topology, population structuring, and demographic history in
Amazonian birds. Genome Biol. Evol., evae002.
Tigano, A., Khan, R., Omer, A.D., Weisz, D., Dudchenko, O., Multani, A.S., Pathak, S.,
Behringer, R.R., Aiden, E.L., Fisher, H., 2021. Chromosome size affects sequence
divergence between species through the interplay of recombination and selection.
bioRxiv.
Todesco, M., Owens, G.L., Bercovich, N., Légaré, J.-S., Soudi, S., Burge, D.O., Huang, K.,
Ostevik, K.L., Drummond, E., Imerovski, I., 2020. Massive haplotypes underlie ecotypic
differentiation in sunflowers. Nature 584, 602-607.
Toews, D.P., Taylor, S.A., Vallender, R., Brelsford, A., Butcher, B.G., Messer, P.W., Lovette,
I.J., 2016. Plumage genes and little else distinguish the genomes of hybridizing warblers.
Curr. Biol. 26, 2313-2318.
Vijay, N., Bossu, C.M., Poelstra, J.W., Weissensteiner, M.H., Suh, A., Kryukov, A.P., Wolf,
J.B., 2016. Evolution of heterogeneous genome differentiation across multiple contact
zones in a crow species complex. Nat. Comm. 7, 1-10.
Wei, T., Simko, V., Levy, M., Xie, Y., Jin, Y., Zemla, J., 2017. Package 'corrplot'. Statistician
56, e24.
Whiting, J.R., Paris, J.R., van der Zee, M.J., Parsons, P.J., Weigel, D., Fraser, B.A., 2021.
Drainage-structuring of ancestral variation and a common functional pathway shape
limited genomic convergence in natural high-and low-predation guppies. PLoS Genet.
17, e1009566.

Wickham, H., 2011. ggplot2. Wiley Interdisciplinary Reviews: Computational Statistics 3, 180-
185.
Wu, C.I., 2001. The genic view of the process of speciation. J. Evol. Biol. 14, 851-865.
Zhang, C., Rabiee, M., Sayyari, E., Mirarab, S., 2018. ASTRAL-III: polynomial time species
tree reconstruction from partially resolved gene trees. BMC Bioinformatics 19, 153.
913