Antiretroviral Therapy Does Not Restore Brain Lipids During SIV Infection: Regional Analysis of Metabolic Homeostasis and Depletion

Cory J. White[1#], Andrew M. Gausepohl[1#], Herana Kamal Seneviratne[2,3], Namandjé N. Bumpus[2,4], Dionna W. Williams[1,2,4,5,6*]

[1]Department of Molecular and Comparative Pathobiology, [2]Department of Pharmacology and Molecular Sciences, [3]Department of Chemistry and Biochemistry, University of Maryland Baltimore County, Baltimore, Maryland 21250, [4]Department of Medicine, Division of Clinical Pharmacology, Johns Hopkins School of Medicine, Baltimore, Maryland 21205

[5]Department of Neuroscience, Johns Hopkins School of Medicine, Baltimore, Maryland 21205

[6]Department of Molecular Microbiology & Immunology, Johns Hopkins School of Public Health, Baltimore, Maryland 21205

[#]Authors contributed equally

*Address correspondence to**: Dionna W. Williams, Ph.D.,** Department of Molecular and Comparative Pathobiology, Johns Hopkins University School of Medicine, 733 N. Broadway St., Miller Research Building 849, Baltimore, MD 21205, Email: dwill201@jhmi.edu

ABSTRACT

Human immunodeficiency virus infection (HIV) continues to promote neurocognitive

impairment, mood disorders, and brain atrophy even in the modern era of viral suppression.

Brain lipids are vulnerable to HIV-associated energetic strain and contribute to HIV-associated

neurologic dysfunction due to alterations in lipid breakdown and structural lipid composition. HIV

neuropathology is region dependent, yet there has not been a comprehensive spatial evaluation

of brain lipids during infection that may impact neurologic function. To address this gap, we

evaluated brain lipid distribution using matrix laser desorption/ionization imaging mass

spectrometry (MALDI-IMS) across four brain regions (parietal cortex, midbrain, thalamus, and

temporal cortex), as well as kidney for a peripheral tissue control, in a virally suppressed simian

immunodeficiency virus (SIV)-infected rhesus macaque. We assessed lipids indicative of fat

breakdown (acylcarnitines) and critical structural lipids (phosphatidylcholines and

phosphatidylethanolamines) across fatty acid chain lengths and degrees of unsaturation.

Acylcarnitines with very long-chain, polyunsaturated fatty acids were more abundant across all

brain regions than shorter chain, saturated or monounsaturated species. We observed two

distinct brain lipid distribution patterns for acylcarnitines and phosphatidylcholines. However, no

clear expression patterns emerged for phosphatidylethanolamines. Surprisingly, acylcarnitines

were largely missing in kidney, while common phosphatidylcholines and

phosphatidylethanolamines were absent in kidney. Overall, our study provides substantial

evidence for persistent bioenergetic changes to the brain despite viral suppression, including

region-dependent mobilization of acylcarnitines for oxidation and disparities in the presence of

key phospholipids necessary for maintaining proper brain structure and function. These data

indicate that region-specific interventions to restore proper lipid metabolism are essential for

treating HIV neurologic disease in the era of ART.

INTRODUCTION

Human immunodeficiency virus (HIV) continues to pose a significant risk to neurologic health, despite increased life expectancy and quality of life afforded by antiretroviral therapy (ART)-mediated viral suppression (*1*). While the severity of neurologic disease has decreased in the ART era, virally suppressed people with HIV are living long enough to age with the virus and are more susceptible to neurocognitive impairment, mood disorders, and brain atrophy compared to uninfected individuals (*2, 3*). HIV induces considerable bioenergetic strain to the brain, which is a substantial driver of the neurologic dysfunction that persists despite viral suppression. Lipids are particularly vulnerable to this energetic strain, due to their high abundance and unique composition in the brain, which contributes substantially to neurologic decline when homeostasis of structural lipids are disrupted or the capacity for lipid utilization is decreased (*2, 3*). Therefore, it was not surprising when strong associations were found between the development of neurocognitive and mood disorders in people with HIV and disruptions in brain lipid metabolism (*4–18*). For example, post-mortem brain tissues from people with HIV that experienced cognitive decline were enriched in sphingomyelin containing very long-chain polyunsaturated fatty acids in parietal cortex, medial frontal gyrus, and cerebellum (*18*). In primary astrocyte cultures, brain cholesterol biosynthesis gene expression and brain cholesterol abundance were both decreased by the HIV viral protein, Tat (*19*). Further, Tat increased fatty oxidation gene expression in primary astrocytes (*20*). Another viral protein, gp120, had a similar effect in hippocampal neurons, where it increased sphingolipids containing very long-chain polyunsaturated fatty acids (*18*). These same very long-chain polyunsaturated fatty acid containing phospholipids were also enriched in the brain of transgenic rats that overexpressed regulatory and accessory HIV proteins (*21, 22*). These studies clearly demonstrate the contribution of lipid dysregulation to central nervous system dysfunction during HIV. However, previous *in vivo* and *ex vivo* studies focused exclusively on profiling total lipid abundance using brain homogenate, without consideration of the spatial differences inherent in the lipid profile

(*18*, *21*). This is a major limitation as it is well described that lipid distribution is not uniform

across brain regions (*23*, *24*), which likely occurs due to the region-specific expression of lipid

breakdown genes and regional differences in structural lipid composition (*25*, *26*). Similarly,

region-dependent neuropathological hallmarks of HIV are well recognized to occur, where

certain brain regions are preferentially vulnerable to HIV-induced damage and others remain

spared (*4–15*). For example, areas of the cortex and the basal ganglia are more prone to white

matter atrophy while areas of cortex and thalamus are prone to gray matter loss due to infection

(*4*, *15*). Despite strong evidence of region-dependent HIV neuropathology and sub-anatomic

heterogeneity inherit in lipid metabolism, there have been no studies to date that characterized

brain lipid distribution across multiple regions after infection.

Therefore, our goal was to characterize lipid distribution in brain utilizing a virally

suppressed model reflective of the modern landscape of HIV treatment to evaluate the direct

contribution of ART in restoring HIV-induced lipid alterations. We evaluated key lipids indicative

of fatty acid breakdown and structural membrane lipid homeostasis using a virally suppressed

SIV-infected rhesus macaque model using matrix assisted laser desorption/ionization imaging

mass spectrometry (MALDI-IMS). Of importance, this model allows for the direct evaluation of

the capacity for ART to restore HIV-induced brain lipid dysregulation in the absence of comorbid

disease or dietary and activity differences, which is a substantial confounding factor when

performing human studies. We also evaluated lipids in kidney to serve as a peripheral control

with known HIV pathology as altered lipid utilization is associated with kidney injury during HIV

(*27*). We first assessed acylcarnitines, fatty acid derivatives transiently generated during long-

chain fatty acid β-oxidation (**Fig. 1A**) (*23*). We also characterized phosphatidylcholines (**Fig. 1B**)

and phosphatidylethanolamines (**Fig. 1C**), prominent phospholipids that are integral in

comprising membranes necessary for maintaining cellular composition, integrity, and function

(*26*). Species from these lipid classes contain fatty acids with a wide range of chain lengths and

degrees of unsaturation that impact physical properties and signaling roles.

In this study, we determined that viral suppression afforded by ART was not sufficient to

restore brain lipid deposition to levels anticipated in the absence of HIV/SIV. Further, we

identified distinct lipid distribution patterns across diverse brain regions that varied according to

fatty acid chain length and degree of unsaturation. Very long chain acylcarnitines were more

abundant in midbrain, thalamus, and temporal cortex. Similar trends occurred for long-chain

saturated and monounsaturated phosphatidylcholines, which had the greatest enrichment in the

same three brain regions. Surprisingly, parietal cortex was depleted in both

phosphatidylcholines and phosphatidylethanolamines. Unexpectedly, we identified prominent

gaps in lipid enrichment across multiple lipid species in diverse brain regions, wherein lipids

were completely depleted from the brain tissue and essentially undetectable. Of note, this did

not occur throughout the entirety of the tissue section, and lipid enrichment often occurred in an

area directly adjacent to the region unpredictably devoid of lipids. These findings demonstrate

that lipid dysregulation continue to occur in people with HIV and suggest that altered abundance

of these lipids critical to normal brain function contribute to HIV CNS persistence in current the

ART era. Overall, this study demonstrates the need for targeted therapeutics to restore region-

specific changes in lipid metabolism linked to neurologic decline in people living with HIV.

MATERIALS AND METHODS

Animal Ethics Statement All procedures were performed in accordance with the *NIH's Guide*

for the Care and Use for Laboratory Animals, the *US Department of Agriculture Animal Welfare*

Act, and under the approval of the Johns Hopkins Medical School Animal Care and Use

Committee. The rhesus macaque used for this study was housed in a cage that was appropriate

for both size and weight in accordance with federal animal welfare regulation; 1 occupied a 9.0-

ft^2 stainless-steel perk bars cage. This animal was included in other studies that included blood

and cerebrospinal fluid (CSF) collections and ART injections performed between 8 and 12 a.m

(*28*). In those studies, the macaques was anesthetized with ketamine hydrochloride to induce

sedation while retaining physiological homeostasis for blood and CSF collections. The macaque

was monitored to ensure there were no signs of pain or distress. Criteria for human euthanasia

prior to planned endpoint if any of the following were observed: a) weight loss of greater than

15%, b) CD4+ T-lymphocytes count less than 5% baseline level, c) clinical signs of neurological

disease, d) intractable diarrhea, and opportunistic infection.

SIV-infected Rhesus Macaque Model

We used a well-established SIV-infected rhesus macaque (*Macaca mulatta*) model optimal for

studying brain pathology, immunological compromise, viral replication, and the impact of

therapeutics on HIV outcome (*28–43*). This study was conducted using historical samples

originally harvested from one juvenile 4-year-old, 6.1 kg SIV-infected, ART-treated rhesus

macaque male (*28*). The macaque wase inoculated intravenously using SIVmac251 as

previously described (*28*, *29*). Two weeks after SIV infection, the rhesus macaque was treated

with daily intraperitoneal injections of combined ARTs consisting of TFV (20 mg/kg), DTG (25

mg/kg), and FTC (50 mg/kg). ARTs were donated from Gilead, ViiV Healthcare, Bristol-Meyers

Squibb, Merck, Abbvie, Janssen, and Roche.

Quantitation of viral load in plasma and CSF

SIV was assessed in plasma and CSF as previously described (*28*, *44–47*). Briefly, RNA was

isolated from 140 µL of plasma or CSF using the QIAamp Viral RNA Mini kit (Qiagen) in

accordance with the manufacturer's protocol with the modification of on-column DNase

digestion (RNase free DNase kit; Qiagen) using RQ1 RNase free DNase (Promega) in the

enzyme mix. Quantification of SIV RNA was performed by reverse transcription polymerase

chain reaction (qRT-PCR) using the following primers and probes: FWD-(SGAG21) 5′-

GTCTGCGTCATCTGGTGCATTC-3′; REV-(SGAG22) 5′-

CACTAGGTGTCTCTGCACTATCTGTTTTG-3′, probe (pSGAG23) 5′-

(FAM)CTTCCTCAGTGTGTTTCACTTTCTCTTCTG-(BH1)-3′.

Necropsy and Harvest

At approximately 130 days post inoculation, the juvenile SIV-infected, ART-treated rhesus

macaque male was sacrificed in accordance with federal guidelines and institutional policies

(*28*). Necropsy protocols were performed in accordance with previous work routinely performed

within the Retrovirus Laboratory of the Department of Molecular and Comparative Pathobiology

in accordance with the American Veterinary Medical Association guidelines for euthanasia of

animals (*28*, *29*, *46*, *48*). Euthanasia occurred with an overdose of sodium pentobarbital while

under ketamine sedation (15- to 20- mg/kg intramuscular injection) before perfusion with

phosphate-buffered saline (PBS) (Gibco) to remove blood from tissues as described (*28*, *29*, *46*,

48). After harvest, the whole brain was placed in cold 2.5% agarose, and 4mm coronal sections

were obtained; fresh kidney was cut into four sections horizontally. Brain regions (parietal

cortex, temporal cortex, thalamus, and midbrain) and kidney were fresh frozen using liquid

nitrogen in optimum cutting temperature medium (OCT) (Sakura Finetek Inc, Torrance, CA) and

stored at -80°C until cryosectioning (*29*).

Materials & Chemicals for MALDI

MALDI matrix, a-cyano-4-hydroxycinnamic acid (CHCA) was obtained from MilliporeSigma (St.

Louis, MO). The CHCA matrix is widely suitable for lipids (*49*). All solvents and other chemicals

used were either reagent or high-performance liquid chromatography (HPLC) grade and

purchased from Fisher Scientific (Hampton, NH), unless otherwise specified.

Tissue Sections & Matrix Application

OCT-embedded brain and kidney were sectioned to 10 μm onto Superfrost Plus Microscope

Slides (Fisher Scientific, Pittsburgh, Pennsylvania) at -20°C using a Leica CM3050S cryostat

(Leica Biosystems, Buffalo Grove, IL). Matrix [CHCA, 10 mg/ml in ACN:H_2O (50:50, v/v)] was

applied to each slide using a TM-Sprayer (HTX Technologies, LLC, Chapel Hill, NC) at a flow

rate of 100 mL/min. The TM-Sprayer was operated at an air pressure (N_2) of 10 psi, spray

nozzle velocity of 1200 mm/min, track spacing at 2 mm, and spray nozzle temperature at 80°C.

Sixteen passes (matrix deposition cycles) were employed for homogeneous deposition of matrix

onto each slide.

MALDI Imaging

MALDI-MSI data were acquired using a LTQ Orbitrap XL (Thermo Fisher Scientific, Bremen,

Germany), equipped with a Fourier transform mass spectrometer (FTMS) with the MALDI ion

source fitted with a direct beam N_2 laser (l 5 337.7 nm). Data acquisition was performed using

the positive ion mode of the MS instrument at a mass range of m/z 100–1000 Da. Mass

resolution of the FTMS analyzer was 60,000. CHCA matrix was used at 10 mg/ml for ionization

of lipids and the laser energy of 7.8 mJ was used. The camera from the MALDI instrument was

used to define regions of interest and generate MALDI position files. Ion images had a spatial

resolution of 50 μm. Mass spectrometry data processing and analysis were done using Xcalibur

3.0 (ThermoFisher Scientific) and ion images were generated using ImageQuest 1.0.1 software

(Thermo Fisher Scientific, San Jose, CA).

Selection of Ion Images

Representative lipid ion images were selected for known m/z values of multiple species for the

following lipid classes: acylcarnitines, phosphatidylcholines, and phosphatidylethanolamines

using Xcalibur 3.0 (ThermoFisher Scientific) (*50*). Ion values were determined for H^+, K^+, and

Na^+ adducts from each neutral lipid using LIPID MAPS Lipidomics Gateway mass (m/z)

calculation tool for lipid classes. Representative images were selected upon determination of

uniform expression patterns from all three adducts.

RESULTS

Virally suppressed, SIV-infected macaque model

We utilized a virally suppressed rhesus macaque model because of the experimental control it facilitates that limit confounding factors inherent with postmortem human brain (28, 51–54). One juvenile male rhesus macaque was infected with SIVmac251 for 14 days, followed by 90 days of viral suppression. We showed that this model achieved peak viremia at 14 days post inoculation and accomplished complete viral suppression by day 42 post inoculation (28 days after ART initiation) (28). Upon necropsy, perfusion was performed, and the brain and kidney were harvested for lipid analysis via MALDI-IMS (18).

Determination of lipid spatial distribution in a virally suppressed, SIV-infected macaque

Brain lipids are necessary for maintaining structure, propagating action potentials, use as signaling molecules, and as a source of energy (3, 23, 55). Numerous neurocognitive disorders, mood disorders, and neurovirulent diseases, including HIV, are linked to dysregulated lipid metabolism (17, 56–58). HIV viral proteins decrease brain cholesterol and sphingolipid abundance and the expression of enzymes necessary for their biosynthesis (18, 19). HIV viral proteins also disrupted membrane lipid homeostasis when overexpressed in rats (21). However, previous studies failed to examine a component critical to lipid abundance and function: region-specific metabolic profiles necessary to meet energetic demands, supply structural needs, and maintain metabolic homeostasis.

To address this gap in knowledge, we evaluated brain lipid distribution by MALDI-IMS from HIV-relevant brain regions (midbrain, parietal cortex, thalamus, temporal cortex) (4, 28). We also performed MALDI-IMS on kidney as a point of reference for lipid distribution in peripheral tissues with known HIV-relevance as renal disorders remain prevalent even among virally suppressed people living with HIV (59). To examine lipid regional distribution, 10-μm coronal slices of parietal cortex, thalamus, temporal cortex, midbrain, and kidney were imaged

with a 50-µm resolution (**Fig. 1A; Fig. 1B**) using a m/z acquisition range of 100-1000 Da (**Fig**

1A; Fig. 1B). We focused on three main classes of lipids of key importance to lipid utilization

(acylcarnitines) and cellular structure/homeostasis (phosphatidylcholines and

phosphatidylethanolamines).

Acylcarnitines are abundant in brain, variably expressed in differing regions, and

minimally present in kidney

We first evaluated acylcarnitines, which have diverse chain lengths that range from two

to over twenty carbons with zero to over six degrees of unsaturation (*60*). There was a striking

pattern wherein very long-chain acylcarnitines were highly expressed in brain, but minimally

present in kidney, which was consistent across all imaged brain regions from our SIV-infected,

ART-treated rhesus macaque. This is evidenced by the very long chain acylcarnitine, C22:4,

that was highly expressed in brain but marginally detected in kidney (**Fig. 2**). Interestingly,

C18:3 was the only acylcarnitine species prominently observed in kidney (**Fig. 2**). Surprisingly,

this acylcarnitine was minimally represented in brain.

We determined that, while prominently represented, very long-chain acylcarnitines had a

high level of regional specificity across brain regions (**Fig. 2**). C20:0 and C22:3 were highly

expressed and homogenously distributed in parietal cortex. Yet, they were sparsely abundant in

midbrain, localizing only to the perimeter of the tissue (**Fig. 2**). These same acylcarnitines had a

distinct localization in thalamus where they were almost uniformly present with incomplete, but

prevalent, representation. Similar trends existed for C22:5, which was not present in parietal

cortex, but was abundant in both midbrain and thalamus. Interestingly, C22:5 had a unique

localization where it had the highest abundance in the innermost portion of tissue but was

absent at the outermost edges. C20:1 showed a similar trend in regional specificity where it was

homogenously present in parietal cortex but localized only to the central portions of midbrain

and thalamus (**Fig. 2**).

Phosphatidylcholines of longer-chain lengths are more prominent in brain than in kidney

We next evaluated phosphatidylcholines, focusing on species with common medium-,

long-, and very long-chain acyl groups (**Fig. 3**). Parietal cortex had the lowest enrichment of

analyzed phosphatidylcholines, where those of all chain lengths were minimally detected (**Fig.**

3A). Conversely, phosphatidylcholines were highly expressed and uniformly present in

midbrain. Thalamus and temporal cortex had high expression of medium-chain

phosphatidylcholines, but lower abundance of the longer chain lengths. Interestingly,

phosphatidylcholines had a comparable spatial distribution as some of the acylcarnitines in

thalamus, wherein the greatest expression was restricted to the uniquely patterned internal zone

(**Fig. 3A; Fig. 2**). The temporal cortex had the most unique and variable phosphatidylcholine

localization patterns of all examined brain regions. PC(32:0) and PC(32:1) were abundant in a

large, central portion of the temporal cortex, while PC(40:4) localized only to the top portion of

the tissue. In contrast, PC(44:6) was primarily only at the bottom portion of the temporal cortex

(**Fig. 3A**). Of interest, there was one notable exception that occurred uniformly across brain

regions. Very-long chain phosphatidylcholines were lowly expressed in brain (**Fig. 3A**). No clear

trend in phosphatidylcholines occurred in kidney, where some species were highly and

uniformly expressed, while others were present to a lower extent (**Fig. 3B**).

Phosphatidylethanolamines are heterogeneously present in brain and kidney

Phosphatidylethanolamines were the last lipid of investigation. Overall, the spatial

distributions of phosphatidylethanolamine species showed less consistent patterning than that

which occurred for acylcarnitines and phosphatidylcholines in both brain and kidney (**Fig. 4**).

Like phosphatidylcholines, smaller chain phosphatidylethanolamines were more prominent

across all brain regions and in kidney, while larger phosphatidylethanolamines were less

represented across both tissues (**Fig. 4**). Interestingly, phosphatidylethanolamines were not

visible in parietal cortex (**Fig. S1**).

DISCUSSION

Our study is among the first to evaluate the spatial distribution of major lipid classes related to lipid utilization (acylcarnitines) and membrane phospholipid homeostasis (phosphatidylcholines and phosphatidylethanolamines) in brain and kidney by MALDI-IMS in a virally suppressed non-human primate model of HIV infection. We identified distinct lipid distribution patterns lipid distribution for acylcarnitines and phosphatidylcholines, but no well-defined patterns for phosphatidylethanolamines. Acylcarnitines comprised of long-chain polyunsaturated fatty acids were more abundant across all brain regions compared to kidney. However, the distribution pattern within each brain region differed by species, where some species were present along the peripheries of regions and others were restricted to the in interior portions. Phosphatidylcholines containing common saturated and monounsaturated long-chain fatty acids were most abundant across brain regions and in kidney. Surprisingly, the parietal cortex was particularly vulnerable to lipid depletion and lacked both phosphatidylcholines and phosphatidylethanolamines. Our findings demonstrate distinct patterns of lipid distribution, including enrichment of very long chain polyunsaturated acylcarnitines across brain regions and limited phosphatidylcholines in parietal cortex. Further, they indicate that, despite viral suppression, ART is not sufficient to restore HIV-induced dysregulations in brain lipid abundance and composition, where some brain regions are at heightened risk for lipid depletion. This has profound implications for the persistence of HIV CNS disease in the era of ART, identifies a previously unrecognized marker of HIV neurologic dysfunction, and indicates that therapeutically targeting lipid metabolic processes in a region-specific manner will be integral in restoring appropriate neurologic function in people living with HIV. Our innovative approach in evaluating lipid distribution across multiple brain regions was imperative to understand HIV-induced changes in the brain lipidome and demonstrates the importance of spatially profiling lipids, rather than exclusively quantifying lipid concentrations in tissue homogenate (*23, 25*).

HIV viral proteins, which are expressed despite suppression of viral replication, greatly

contribute to neurologic dysfunction. These viral proteins contribute to neurotoxicity manifesting

as dendritic injuries (*61–63*), neurogenic deficits (*64*), and neuroendocrine defects (*65*).

Neurotoxic HIV proteins include gp120, gp41, Tat, Nef, Rev, and Vpr (*66, 67*). Viral proteins

alter brain bioenergetics in numerous ways, including Ca^{2+} dysregulation (*68–70*), modulated

fatty acid oxidation (*20*), and electron transport chain complex abnormalities (*71*). They also

disrupt metabolic coupling in the brain (*20*), elevate reactive oxygen species (*72, 73*), and

exacerbate neuroinflammation (*61, 74*). Previous studies showed that HIV viral proteins

increased fatty acid oxidation genes *in vitro* and disrupted the balance of membrane lipids (*20,*

21). Our findings in this study identify the high abundance of many acylcarnitines during viral

suppression. This is of major significance because acylcarnitines are fatty acid oxidation

intermediates, which is often contested to occur within brain (*23*). Our studies strongly implicate

that the HIV-infected brain undergoes a considerable amount of fatty acid oxidation and is

enriched in lipid species associated with countering HIV-associated neuroinflammation. Further,

while phosphatidylcholines containing polyunsaturated very long-chain fatty acids are less

prominent than phosphatidylcholines containing common saturated and unsaturated long-chain

fatty acids, they are still well represented across most brain regions. Previous work determined

brain tissue from people with HIV that experienced severe cognitive decline had increased very

long-chain fatty acids in sphingolipids, a class of structural membrane lipids, despite viral

suppression (*18*). It is possible that elevations in very long-chain, polyunsaturated fatty acids

contribute to, or are indicative of, cognitive decline. Determining the mechanisms by which

HIV/SIV and viral suppression impact these fatty acids is necessary to identify therapeutic

strategies to restore changes to brain lipid homeostasis that contribute to HIV-associated

cognitive decline.

Chronic inflammation is another major contributor to CNS dysfunction that persists in

people living with HIV, even with successful ART (*75–77*). Inflammation in the periphery,

perivascular space, and within the brain parenchyma contribute to CNS dysfunction during HIV

(*78–82*). Chronic inflammation promotes neuropathology, including accelerated brain aging,

increased risk of developing psychiatric disorders, and cognitive decline (*83*). In the short term,

the neuroimmune response is beneficial by promoting pathogen clearance, inducing

angiogenesis, and increasing wound healing (*84*). In addition to canonical cytokines and

chemokines, classes of specialized pro-resolving lipid mediators, including lipoxins, protectins,

resolvins, and meresins, also promote neuroinflammation (*85*). Our study identified high

abundance of the C22:5 fatty acid prominently found in specialized proresolving mediators

known to alleviate neuroinflammation in specific areas of midbrain, thalamus, and temporal

cortex (**Fig. 2**). This is important because previous studies determined that HIV viral proteins

increased the abundance of C22:5 acid in total lipid, phosphatidylethanolamine, and

triglycerides (*21*). Our findings indicate that lipid dysregulation in brain that contributes to

persistent neuroimmune activation during HIV/SIV-infection, even with adequate viral

suppression. It is likely that the high abundance of C22:5 exists to counter the chronic

neuroinflammation that occurs after HIV infection.

ART may also play a direct role in lipid dysregulation in brain as its adverse impact on

metabolism is well characterized. Systemically, it is well established that ART causes a mean

weight gain of 3-7 kg within the first year of initiating treatment (*86–89*). Further, specific ART

drugs or ART classes can have varying impacts on circulating lipid levels, blood pressure, and

risks for developing metabolic syndromes, including diabetes mellitus. For example, classical

non-nucleoside reverse transcriptase inhibitors and protease inhibitors were known for greatly

increasing the likelihood of developing dyslipidemia (*90*). Even though newer antiretrovirals do

not elevate circulating lipids to the same extent, tenofovir prodrugs, including tenofovir

alafenamide, increase cholesterol species (*91*). This occurs despite the known impact of HIV on

diminishing total cholesterol levels (*86, 92–95*). Conversely other ARTs, including raltegravir,

dolutegravir, and bictegravir, have either a beneficial or minimal impact on the cholesterol profile

after changing regimens from older protease inhibitors (*86*, *96*). Hypertension, another disorder

related to lipid dysregulation, is also more prevalent in individuals receiving ART. There is a

1.68-fold higher risk of developing hypertension in people living with HIV on ARTs in

comparison to those not on ART (*86*, *97*). However, it is not well characterized whether many of

these changes to metabolic parameters occur due to the antiretrovirals directly, or because of

the weight gain associated with many ARTs (*86*). With the clear role of ART in promoting

lipodystrophy and metabolic dysfunction systemically, it is likely that similar perturbations also

occur in brain.

Indices of kidney injury due to HIV are common as virally suppressed people living with

HIV are aging. However, ART also poses a direct insult to kidney physiology. For example,

tenofovir and its adenofir analog induced mitochondrial dysfunction and glycogen accumulation

in human kidney cells that may contribute to nephrotoxicity (*98*). While less is understood about

the impact ART on lipid metabolism in kidney, tenofovir decreased levels of low-density

lipoprotein and increased the rate of change in urine-liver-type fatty acid binding

protein/creatinine. These are known markers of elevated risk of hyperlipidemia and chronic

kidney disease respectively (*99*), demonstrating a precedence for ART promoting lipid

dysregulation in kidney. The contribution of ART to lipid perturbations in kidney has profound

implications for people with HIV as lipid metabolism is highly relevant to kidney health, as it is

the organ that preferentially oxidizes fatty acids used for energy. Further, lipid metabolism

dysfunction is a well-established risk factor for kidney disease (*27*). Despite the clear

importance and relevance of lipids in kidney function and normal physiology, we detected only a

single acylcarnitine species in kidney (C18:3) (**Fig. 2**). It is possible that the acylcarnitine

species we did not detect were exhausted to generate ATP, or that defective mitochondrial

metabolism occurred that prevented other acylcarnitines from being produced (*98*). Our findings

demonstrate that the kidney remains vulnerable to lipid dysfunction, even in the setting of ART-

mediated viral suppression, and HIV/SIV and ART may contribute to increased propensity for

kidney injury due to lipid dysregulation.

HIV poses a major risk to brain lipid metabolic homeostasis that contributes to

neurologic decline and brain pathology that continue to occur, despite viral suppression (*4–18*).

Characterizing region-dependent changes in lipid profiles is critical for determining region-

specific interventions that will restore proper brain function and metabolism disrupted due to HIV

CNS disease. We utilized advancements in lipid imaging by MALDI-IMS combined with our

virally suppressed SIV rhesus macaque model for an innovative approach to address this gap.

Here, we demonstrated that, despite viral suppression, lipids indicative of increased fatty acid

oxidation, dysregulated structural lipids, and activated neuroimmune response were prominent

in a SIV-infected non-human primate. These metabolic changes occurred in a region-dependent

manner, and it is imperative to fully understand HIV-induced changes to the brain lipidome that

may contribute to HIV-associated neurologic decline in order to develop therapeutics that target

spatially distinct areas of brain metabolic dysfunction.

Acknowledgements

We thank the members of the Johns Hopkins School of Medicine Retrovirus Laboratory in the Department of Molecular and Comparative Pathobiology for husbandry of non-human primate models, analysis of infection and viral suppression of the animal, necropsy and sample collection, and for veterinary care including Suzanne E. Queen, Erin N. Shirk, Brandon T. Bullock, Kelly A. Metcalf Pate, Lisa M. Mangus, Janice E. Clements, and Joseph L. Mankowski. We are also thankful for Gilead and ViiV for graciously providing antiretrovirals used by non-human primate models.

This study was supported by National Institutes of Health (NIH) grants R00 DA044838 and R01 DA052859 to Dionna W. Williams, U19AI113127 to Namandjé N. Bumpus, and K00 NS118713 Cory J. White. Also, this work was in part supported by pilot funding provided by parent funding under the JHU NIMH Center for Novel Therapeutics for HIV-associated Cognitive Disorders P30 MH075673 to Justin C. McArthur, and P01 AI131306 to Janice E. Clements.

Images in figures were created using BioRender.

Conflict of Interest

The authors report no conflicts of interest.

Author Contributions

D.W.W. and C.J.W. performed experiments, A.M.G. and C.J.W. analyzed data, D.W.W. and C.J.W. were responsible for conceptualization of the study design, H.K.S. and N.N.B. developed the methodologies, C.J.W. and D.W.W. wrote the original draft with review and editing from all authors.

FIGURE LEGENDS

Figure 1. MALDI MS Imaging of SIV-infected, ART-treated macaque tissues using CHCA matrix and positive mode of the LTQ Orbitrap XL instrument. Representative full scan spectra and images of total ion counts of SIV-infected, ART-treated rhesus macaque A) brain regions: (parietal cortex, midbrain, thalamus, and temporal cortex) and B) kidney. Red signifies the highest intensity and blue the lowest of each m/z. C) Acylcarnitine (red), phosphatidylcholine (blue), and phospatidylethanolamine (gray) structures. R = acyl (fatty chain) group.

Figure 2. Very long-chain acylcarnitines are prevalent in the brain, but not kidney, where spatial distribution varies by species composition. Representative MALDI-IMS of SIV-infected, ART-treated rhesus macaque brain regions [parietal cortex (P. ctx), midbrain, thalamus, and temporal cortex (T. ctx)] and kidney of A) very long-chain polyunsaturated AC C22:4 species, B) polyunsaturated C18:3 AC species, C) varying very long-chain un-, mono-, and polyunsaturated AC species demonstrating differences in spatial distribution. Red signifies the highest intensity and blue the lowest of each m/z.

Figure 3. Lower chain-length phosphatidylcholines are more abundant in both kidney and across brain regions. Representative MALDI-IMS of SIV-infected, ART-treated rhesus macaque brain regions of A) m/z values shared by representative PCs of similar chain length in A) the following brain regions: parietal cortex (P. ctx), thalamus, temporal cortex (T. ctx), and midbrain as well as B) in kidney. Red signifies the highest intensity and blue the lowest of each m/z.

Figure 4. Patterns in phosphatidylethanolamine spatial distribution across brain regions

or within a respective tissue are less clear. Representative MALDI-IMS of SIV-infected, ART-

treated rhesus macaque brain regions of A) m/z values shared by representative PEs of similar

chain length in A) the following brain regions: thalamus, midbrain, and temporal cortex (T. ctx)

as well as B) in kidney. Red signifies the highest abundance and blue the lowest of each m/z.

Figure S1. Phosphatidylethanolamines were not observed within parietal cortex.

MALDI-IMS of SIV-infected, ART-treated rhesus macaque parietal cortex of m/z values shared

by PEs in linked PowerPoint file. Red signifies the highest abundance and blue the lowest of

each m/z.

References

1. M. S. Espíndola, L. S. Soares, L. J. Galvão-Lima, F. A. Zambuzi, M. C. Cacemiro, V. S.
Brauer, F. G. Frantz, HIV infection: focus on the innate immune cells. *Immunol Res.* 64, 1118–
1132 (2016).

2. R. W. Mitchell, G. M. Hatch, Fatty acid transport into the brain Of fatty acid fables and lipid
tails. *Prostaglandins, Leukotrienes and Essential Fatty Acids.* 85, 293–302 (2011).

3. P. A. Watkins, J. A. Hamilton, A. Leaf, A. A. Spector, S. A. Moore, R. E. Anderson, H. W.
Moser, M. J. Noetzel, R. Katz, (2001), vol. 16 of *Journal of molecular neuroscience : MN*, pp.
87-92-discussion 151-7.

4. A. Gongvatana, B. C. Schweinsburg, M. J. Taylor, R. J. Theilmann, S. L. Letendre, O. M.
Alhassoon, J. Jacobus, S. P. Woods, T. L. Jernigan, R. J. Ellis, L. R. Frank, I. Grant, T. C.
Group, White matter tract injury and cognitive impairment in human immunodeficiency virus–
infected individuals. *J Neurovirol.* 15, 187–195 (2009).

5. M. A. Mohamed, P. B. Barker, R. L. Skolasky, O. A. Selnes, R. T. Moxley, M. G. Pomper, N.
C. Sacktor, Brain metabolism and cognitive impairment in HIV infection: a 3-T magnetic
resonance spectroscopy study. *Magnetic Resonance Imaging.* 28, 1251–1257 (2010).

6. K. J. Towgood, M. Pitkanen, R. Kulasegaram, A. Fradera, S. Soni, N. Sibtain, L. J. Reed, C.
Bradbeer, G. J. Barker, J. T. Dunn, F. Zelaya, M. D. Kopelman, Regional cerebral blood flow
and FDG uptake in asymptomatic HIV-1 men. *Human Brain Mapping.* 34, 2484–2493 (2012).

7. T. Ernst, C. S. Jiang, H. Nakama, S. Buchthal, L. Chang, Lower brain glutamate is associated
with cognitive deficits in HIV patients: A new mechanism for HIV-associated neurocognitive
disorder. *Journal of Magnetic Resonance Imaging.* 32, 1045–1053 (2010).

8. Å. B. Andersen, I. Law, K. S. Krabbe, H. Bruunsgaard, S. R. Ostrowski, H. Ullum, L.
Højgaard, A. Lebech, J. Gerstoft, A. Kjær, Cerebral FDG-PET scanning abnormalities in
optimally treated HIV patients. *J Neuroinflamm.* 7, 13–13 (2010).

9. H. N. Consortium, A. Gongvatana, J. Harezlak, S. Buchthal, E. Daar, G. Schifitto, T.
Campbell, M. Taylor, E. Singer, J. Algers, J. Zhong, M. Brown, D. McMahon, Y. T. So, D. Mi, R.
Heaton, K. Robertson, C. Yiannoutsos, R. A. Cohen, B. Navia, Progressive cerebral injury in the
setting of chronic HIV infection and antiretroviral therapy. *Journal of NeuroVirology.* 19, 209–218
(2013).

10. J. J. Weiss, R. Calvi, M. Naganawa, T. Toyonaga, S. F. Farhadian, M. Chintanaphol, J.
Chiarella, M.-Q. Zheng, J. Ropchan, Y. Huang, R. H. Pietrzak, R. E. Carson, S. Spudich,
Preliminary In Vivo Evidence of Reduced Synaptic Density in Human Immunodeficiency Virus
(HIV) Despite Antiretroviral Therapy. *Clin Infect Dis Official Publ Infect Dis Soc Am.* 73, 1404–
1411 (2021).

11. V. Samboju, C. L. Philippi, P. Chan, Y. Cobigo, J. L. K. Fletcher, M. Robb, J. Hellmuth, K.
Benjapornpong, N. Dumrongpisutikul, M. Pothisri, R. Paul, J. Ananworanich, S. Spudich, V.

Valcour, S. 010/RV254, R. protocol teams, Structural and functional brain imaging in acute HIV.
Neuroimage Clin. 20, 327–335 (2018).

12. C.-X. Li, J. G. Herndon, F. J. Novembre, X. Zhang, A Longitudinal Magnetization Transfer
Imaging Evaluation of Brain Injury in a Macaque Model of NeuroAIDS. *Aids Res Hum Retrov*.
31, 335–341 (2015).

13. C.-X. Li, X. Zhang, A. Komery, Y. Li, H. Mao, J. G. Herndon, F. J. Novembre, Longitudinal
cerebral metabolic changes in pig-tailed macaques infected with the neurovirulent virus
SIVsmmFGb. *J Neurovirol*. 20, 612–619 (2014).

14. C. Li, X. Zhang, A. Komery, Y. Li, F. J. Novembre, J. G. Herndon, Longitudinal diffusion
tensor imaging and perfusion MRI investigation in a macaque model of neuro-AIDS: A
preliminary study. *Neuroimage*. 58, 286–292 (2011).

15. D. Saylor, A. M. Dickens, N. Sacktor, N. Haughey, B. Slusher, M. Pletnikov, J. L.
Mankowski, A. Brown, D. J. Volsky, J. C. McArthur, HIV-associated neurocognitive disorder —
pathogenesis and prospects for treatment. *Nature Reviews Neurology*. 12, 234–248 (2016).

16. P. Deme, C. Rojas, B. S. Slusher, R. Rais, Z. Afghah, J. D. Geiger, N. J. Haughey,
Bioenergetic adaptations to HIV infection. Could modulation of energy substrate utilization
improve brain health in people living with HIV-1? *Exp Neurol*. 327, 113181 (2020).

17. B. Cotto, K. Natarajaseenivasan, D. Langford, HIV-1 infection alters energy metabolism in
the brain:contributions to HIV-associated neurocognitive disorders. *Progress in Neurobiology*,
101616 (2019).

18. N. J. Haughey, R. G. Cutler, A. Tamara, J. C. McArthur, D. L. Vargas, C. A. Pardo, J.
Turchan, A. Nath, M. P. Mattson, Perturbation of sphingolipid metabolism and ceramide
production in HIV-dementia. *Ann Neurol*. 55, 257–267 (2004).

19. B. Cotto, K. Natarajaseenivasan, K. Ferrero, L. Wesley, M. Sayre, D. Langford, Cocaine and
HIV-1 Tat disrupt cholesterol homeostasis in astrocytes: Implications for HIV-associated
neurocognitive disorders in cocaine user patients. *Glia*. 66, 889–902 (2018).

20. K. Natarajaseenivasan, B. Cotto, S. Shanmughapriya, A. A. Lombardi, P. K. Datta, M.
Madesh, J. W. Elrod, K. Khalili, D. Langford, Astrocytic metabolic switch is a novel etiology for
Cocaine and HIV-1 Tat-mediated neurotoxicity. *Cell Death Dis*. 9, 415 (2018).

21. A. Y. Taha, M. Basselin, E. Ramadan, H. R. Modi, S. I. Rapoport, Y. Cheon, Altered lipid
concentrations of liver, heart and plasma but not brain in HIV-1 transgenic rats. *Prostaglandins
Leukot Essent Fat Acids*. 87, 91–101 (2012).

22. M. Vigorito, K. P. Connaghan, S. L. Chang, The HIV-1 transgenic rat model of neuroHIV.
Brain Behav Immun. 48, 336–349 (2015).

23. C. J. White, J. Lee, J. Choi, T. Chu, S. Scafidi, M. J. Wolfgang, Determining the Bioenergetic
Capacity for Fatty Acid Oxidation in the Mammalian Nervous System. *Mol Cell Biol*. 40 (2020),
doi:10.1128/mcb.00037-20.

24. G. Sighinolfi, S. Clark, L. Blanc, D. Cota, B. Rhourri-Frih, Mass spectrometry imaging of
mice brain lipid profile changes over time under high fat diet. *Sci Rep-uk*. 11, 19664 (2021).

25. J. N. Jernberg, C. E. Bowman, M. J. Wolfgang, S. Scafidi, Developmental regulation and
localization of carnitine palmitoyltransferases (CPTs) in rat brain. *Journal of Neurochemistry*.
142, 407–419 (2017).

26. J. N. van der Veen, J. P. Kennelly, S. Wan, J. E. Vance, D. E. Vance, R. L. Jacobs, The
critical role of phosphatidylcholine and phosphatidylethanolamine metabolism in health and
disease. *Biochimica Et Biophysica Acta Bba - Biomembr*. 1859, 1558–1572 (2017).

27. H.-S. Jang, M. R. Noh, J. Kim, B. J. Padanilam, Defective Mitochondrial Fatty Acid Oxidation
and Lipotoxicity in Kidney Diseases. *Frontiers Medicine*. 7, 65 (2020).

28. C. M. Abreu, R. T. Veenhuis, C. R. Avalos, S. Graham, D. R. Parrilla, E. A. Ferreira, S. E.
Queen, E. N. Shirk, B. T. Bullock, M. Li, K. A. M. Pate, S. E. Beck, L. M. Mangus, J. L.
Mankowski, F. M. Gabhann, S. L. O'Connor, L. Gama, J. E. Clements, Myeloid and CD4 T Cells
Comprise the Latent Reservoir in Antiretroviral Therapy-Suppressed SIVmac251-Infected
Macaques. *Mbio*. 10, e01659-19 (2019).

29. A. M. Gorska, M. Donoso, S. Valdebenito, B. Prideaux, S. Queen, E. Scemes, J. Clements,
E. Eugenin, Human immunodeficiency virus-1/simian immunodeficiency virus infection induces
opening of pannexin-1 channels resulting in neuronal synaptic compromise: A novel therapeutic
opportunity to prevent NeuroHIV. *J Neurochem*. 158, 500–521 (2021).

30. J. E. Clements, J. L. Mankowski, L. Gama, M. C. Zink, The accelerated simian
immunodeficiency virus macaque model of human immunodeficiency virus-associated
neurological disease: From mechanism to treatment. *J Neurovirol*. 14, 309–317 (2008).

31. G. M. Shaw, M. E. Harper, B. H. Hahn, L. G. Epstein, D. C. Gajdusek, R. W. Price, B. A.
Navia, C. K. Petito, C. J. O'Hara, J. E. Groopman, E.-S. Cho, J. M. Oleske, F. Wong-Staal, R.
C. Gallo, HTLV-III Infection in Brains of Children and Adults with AIDS Encephalopathy.
Science. 227, 177–182 (1985).

32. J. Rappaport, D. J. Volsky, Role of the macrophage in HIV-associated neurocognitive
disorders and other comorbidities in patients on effective antiretroviral treatment. *J Neurovirol*.
21, 235–241 (2015).

33. K. Williams, S. Westmoreland, J. Greco, E. Ratai, M. Lentz, W.-K. Kim, R. A. Fuller, J. P.
Kim, P. Autissier, P. K. Sehgal, R. F. Schinazi, N. Bischofberger, M. Piatak, J. D. Lifson, E.
Masliah, R. G. González, Magnetic resonance spectroscopy reveals that activated monocytes
contribute to neuronal injury in SIV neuroAIDS. *J Clin Invest*. 115, 2534–2545 (2005).

34. V. Monceaux, L. Viollet, F. Petit, M. C. Cumont, G. R. Kaufmann, A. M. Aubertin, B. Hurtrel,
G. Silvestri, J. Estaquier, CD4 + CCR5 + T-Cell Dynamics during Simian Immunodeficiency
Virus Infection of Chinese Rhesus Macaques. *J Virol*. 81, 13865–13875 (2007).

35. J. B. Whitney, A. L. Hill, S. Sanisetty, P. Penaloza-MacMaster, J. Liu, M. Shetty, L.
Parenteau, C. Cabral, J. Shields, S. Blackmore, J. Y. Smith, A. L. Brinkman, L. E. Peter, S. I.
Mathew, K. M. Smith, E. N. Borducchi, D. I. S. Rosenbloom, M. G. Lewis, J. Hattersley, B. Li, J.

Hesselgesser, R. Geleziunas, M. L. Robb, J. H. Kim, N. L. Michael, D. H. Barouch, Rapid Seeding of the Viral Reservoir Prior to SIV Viremia in Rhesus Monkeys. *Nature*. 512, 74–77 (2014).

36. S. Tuyishime, L. H. Haut, R. K. Kurupati, J. M. Billingsley, D. Carnathan, S. Gangahara, T. M. Styles, Z. Xiang, Y. Li, M. Zopfs, Q. Liu, X. Zhou, M. G. Lewis, R. R. Amara, S. Bosinger, G. Silvestri, H. C. J. Ertl, Correlates of Protection Against SIVmac251 Infection in Rhesus Macaques Immunized With Chimpanzee-Derived Adenovirus Vectors. *Ebiomedicine*. 31, 25–35 (2018).

37. D. H. Barouch, G. Alter, T. Broge, C. Linde, M. E. Ackerman, E. P. Brown, E. N. Borducchi, K. M. Smith, J. P. Nkolola, J. Liu, J. Shields, L. Parenteau, J. B. Whitney, P. Abbink, D. M. Ng'ang'a, M. S. Seaman, C. L. Lavine, J. R. Perry, W. Li, A. D. Colantonio, M. G. Lewis, B. Chen, H. Wenschuh, U. Reimer, M. Piatak, J. D. Lifson, S. A. Handley, H. W. Virgin, M. Koutsoukos, C. Lorin, G. Voss, M. Weijtens, M. G. Pau, H. Schuitemaker, Protective efficacy of adenovirus/protein vaccines against SIV challenges in rhesus monkeys. *Science*. 349, 320–324 (2015).

38. D. H. Barouch, P. J. Klasse, J. Dufour, R. S. Veazey, J. P. Moore, Macaque studies of vaccine and microbicide combinations for preventing HIV-1 sexual transmission. *Proc National Acad Sci*. 109, 8694–8698 (2012).

39. M. Vaccari, B. F. Keele, S. E. Bosinger, M. N. Doster, Z.-M. Ma, J. Pollara, A. Hryniewicz, G. Ferrari, Y. Guan, D. N. Forthal, D. Venzon, C. Fenizia, T. Morgan, D. Montefiori, J. D. Lifson, C. J. Miller, G. Silvestri, M. Rosati, B. K. Felber, G. N. Pavlakis, J. Tartaglia, G. Franchini, Protection Afforded by an HIV Vaccine Candidate in Macaques Depends on the Dose of SIV mac251 at Challenge Exposure. *J Virol*. 87, 3538–3548 (2013).

40. L. Micci, X. Alvarez, R. I. Iriele, A. M. Ortiz, E. S. Ryan, C. S. McGary, C. Deleage, B. B. McAtee, T. He, C. Apetrei, K. Easley, S. Pahwa, R. G. Collman, C. A. Derdeyn, M. P. Davenport, J. D. Estes, G. Silvestri, A. A. Lackner, M. Paiardini, CD4 Depletion in SIV-Infected Macaques Results in Macrophage and Microglia Infection with Rapid Turnover of Infected Cells. *Plos Pathog*. 10, e1004467 (2014).

41. B. R. Magalis, D. J. Nolan, P. Autissier, T. H. Burdo, K. C. Williams, M. Salemi, Insights into the Impact of CD8 + Immune Modulation on Human Immunodeficiency Virus Evolutionary Dynamics in Distinct Anatomical Compartments by Using Simian Immunodeficiency Virus-Infected Macaque Models of AIDS Progression. *J Virol*. 91 (2017), doi:10.1128/jvi.01162-17.

42. J. Mallard, K. C. Williams, Chapter 4 Animal models of HIV-associated disease of the central nervous system. *Handb Clin Neurology*. 152, 41–53 (2018).

43. M. R. Lentz, S. V. Westmoreland, V. Lee, E. Ratai, E. F. Halpern, R. G. González, Metabolic markers of neuronal injury correlate with SIV CNS disease severity and inoculum in the macaque model of neuroAIDS. *Magnet Reson Med*. 59, 475–484 (2008).

44. A. Shen, M. C. Zink, J. L. Mankowski, K. Chadwick, J. B. Margolick, L. M. Carruth, M. Li, J. E. Clements, R. F. Siliciano, Resting CD4 + T Lymphocytes but Not Thymocytes Provide a Latent Viral Reservoir in a Simian Immunodeficiency Virus- Macaca nemestrina Model of

Human Immunodeficiency Virus Type 1-Infected Patients on Highly Active Antiretroviral
Therapy. *J Virol*. 77, 4938–4949 (2003).

45. L. Gama, E. N. Shirk, J. N. Russell, K. I. Carvalho, M. Li, S. E. Queen, J. Kalil, M. C. Zink, J.
E. Clements, E. G. Kallas, Expansion of a subset of CD14highCD16negCCR2low/neg
monocytes functionally similar to myeloid-derived suppressor cells during SIV and HIV infection.
J Leukocyte Biol. 91, 803–816 (2012).

46. L. Gama, C. M. Abreu, E. N. Shirk, S. L. Price, M. Li, G. M. Laird, K. A. M. Pate, S. W.
Wietgrefe, S. L. O'Connor, L. Pianowski, A. T. Haase, C. V. Lint, R. F. Siliciano, J. E. Clements,
Reactivation of simian immunodeficiency virus reservoirs in the brain of virally suppressed
macaques. *Aids*. 31, 5–14 (2017).

47. K. A. Meulendyke, M. V. Pletnikov, E. L. Engle, P. M. Tarwater, D. R. Graham, M. C. Zink,
Early Minocycline Treatment Prevents a Decrease in Striatal Dopamine in an SIV Model of HIV-
Associated Neurological Disease. *J Neuroimmune Pharm*. 7, 454–464 (2012).

48. M. C. Zink, K. Suryanarayana, J. L. Mankowski, A. Shen, M. Piatak, J. P. Spelman, D. L.
Carter, R. J. Adams, J. D. Lifson, J. E. Clements, High Viral Load in the Cerebrospinal Fluid and
Brain Correlates with Severity of Simian Immunodeficiency Virus Encephalitis. *J Virol*. 73,
10480–10488 (1999).

49. K. C. Pathmasiri, T. T. A. Nguyen, N. Khamidova, S. M. Cologna, Mass spectrometry-based
lipid analysis and imaging. *Curr Top Membr*. 88, 315–357 (2021).

50. E. Fahy, S. Subramaniam, R. C. Murphy, M. Nishijima, C. R. H. Raetz, T. Shimizu, F.
Spener, G. van Meer, M. J. O. Wakelam, E. A. Dennis, Update of the LIPID MAPS
comprehensive classification system for lipids 1. *J Lipid Res*. 50, S9–S14 (2009).

51. D. A. Lewis, The Human Brain Revisited: Opportunities and Challenges in Postmortem
Studies of Psychiatric Disorders. *Neuropsychopharmacol*. 26, 143–154 (2002).

52. J. Weickenmeier, M. Kurt, E. Ozkaya, R. de Rooij, T. C. Ovaert, R. L. Ehman, K. B. Pauly,
E. Kuhl, Brain stiffens post mortem. *J Mech Behav Biomed*. 84, 88–98 (2018).

53. A. C. Birdsill, D. G. Walker, L. Lue, L. I. Sue, T. G. Beach, Postmortem interval effect on
RNA and gene expression in human brain tissue. *Cell Tissue Bank*. 12, 311–318 (2011).

54. Z. ElHajj, A. Cachot, T. Müller, I. M. Riederer, B. M. Riederer, Effects of postmortem delays
on protein composition and oxidation. *Brain Res Bull*. 121, 98–104 (2016).

55. R. W. Mitchell, N. H. On, M. R. D. Bigio, D. W. Miller, G. M. Hatch, Fatty acid transport
protein expression in human brain and potential role in fatty acid transport across human brain
microvessel endothelial cells. *J Neurochem*. 117, 735–746 (2011).

56. A. T. Ahmed, S. MahmoudianDehkordi, S. Bhattacharyya, M. Arnold, D. Liu, D. Neavin, M.
A. Moseley, J. W. Thompson, L. S. J. Williams, G. Louie, M. K. Skime, L. Wang, P. Riva-Posse,
W. McDonald, W. V. Bobo, W. E. Craighead, R. Krishnan, R. M. Weinshilboum, B. W. Dunlop,
D. S. Millington, A. J. Rush, M. A. Frye, R. Kaddurah-Daouk, T. M. D. P. M. C. (MDPMC),

Acylcarnitine metabolomic profiles inform clinically-defined major depressive phenotypes. *J Affect Disorders.* 264, 90–97 (2020).

57. T. Clark-Taylor, B. E. Clark-Taylor, Is autism a disorder of fatty acid metabolism? Possible dysfunction of mitochondrial β-oxidation by long chain acyl-CoA dehydrogenase. *Medical Hypotheses*. 62, 970–975 (2004).

58. R. Barone, S. Alaimo, M. Messina, A. Pulvirenti, J. Bastin, M.-A. Group, A. Ferro, R. E. Frye, R. Rizzo, A Subset of Patients With Autism Spectrum Disorders Show a Distinctive Metabolic Profile by Dried Blood Spot Analyses. *Frontiers in Psychiatry*. 9, 1–11 (2018).

59. C. M. Wyatt, Kidney Disease and HIV Infection. *Top Antivir Medicine*. 25, 13–16 (2017).

60. C. C. C. R. de Carvalho, M. J. Caramujo, The Various Roles of Fatty Acids. *Mol J Synthetic Chem Nat Prod Chem*. 23, 2583 (2018).

61. S. Iskander, K. A. Walsh, R. R. Hammond, Human CNS cultures exposed to HIV-1 gp120 reproduce dendritic injuries of HIV-1-associated dementia. *J Neuroinflamm*. 1, 7 (2004).

62. S. M. Toggas, E. Masliah, E. M. Rockenstein, G. F. Rail, C. R. Abraham, L. Mucke, Central nervous system damage produced by expression of the HIV-1 coat protein gpl20 in transgenic mice. *Nature*. 367, 188–193 (1994).

63. B. O. Kim, Y. Liu, Y. Ruan, Z. C. Xu, L. Schantz, J. J. He, Neuropathologies in Transgenic Mice Expressing Human Immunodeficiency Virus Type 1 Tat Protein under the Regulation of the Astrocyte-Specific Glial Fibrillary Acidic Protein Promoter and Doxycycline. *Am J Pathology*. 162, 1693–1707 (2003).

64. R. Putatunda, Y. Zhang, F. Li, X.-F. Yang, M. F. Barbe, W. Hu, Adult neurogenic deficits in HIV-1 Tg26 transgenic mice. *J Neuroinflamm*. 15, 287 (2018).

65. M. F. Salahuddin, F. Mahdi, S. P. Sulochana, J. J. Paris, HIV-1 Tat Protein Promotes Neuroendocrine Dysfunction Concurrent with the Potentiation of Oxycodone's Psychomotor Effects in Female Mice. *Viruses*. 13, 813 (2021).

66. A. Nath, J. Geiger, Neurobiological aspects of human immunodeficiency virus infection:Neurotoxic mechanisms. *Prog Neurobiol*. 54, 19–33 (1998).

67. V. Repunte-Canonigo, C. Lefebvre, O. George, T. Kawamura, M. Morales, G. F. Koob, A. Califano, E. Masliah, P. P. Sanna, Gene expression changes consistent with neuroAIDS and impaired working memory in HIV-1 transgenic rats. *Mol Neurodegener*. 9, 26–26 (2014).

68. S. Fitting, P. E. Knapp, S. Zou, W. D. Marks, M. S. Bowers, H. I. Akbarali, K. F. Hauser, Interactive HIV-1 Tat and Morphine-Induced Synaptodendritic Injury Is Triggered through Focal Disruptions in Na+ Influx, Mitochondrial Instability, and Ca2+ Overload. *J Neurosci*. 34, 12850–12864 (2014).

69. S. Fitting, S. Zou, N. El-Hage1, M. Suzuki, J. J. Paris, C. J. Schier, J. W. Rodríguez, M. Rodriguez, P. E. Knapp, K. F. Hauser, Opiate Addiction Therapies and HIV-1 Tat: Interactive

Effects on Glial [Ca2+]i, Oxyradical and Neuroinflammatory Chemokine Production and
Correlative Neurotoxicity. *Curr Hiv Res*. 12, 424–434 (2014).

70. A. Ciardo, J. Meldolesi, Effects of the HIV-1 Envelope Glycoprotein gp120 in Cerebellar
Cultures. [Ca2+]i Increases in a Glial Cell Subpopulation. *Eur J Neurosci*. 5, 1711–1718 (1993).

71. L. M. Villeneuve, P. R. Purnell, K. L. Stauch, S. E. Callen, S. J. Buch, H. S. Fox, HIV-1
transgenic rats display mitochondrial abnormalities consistent with abnormal energy generation
and distribution. *Journal of NeuroVirology*, 1–11 (2016).

72. J. Steiner, N. Haughey, W. Li, A. Venkatesan, C. Anderson, R. Reid, T. Malpica, C.
Pocernich, D. A. Butterfield, A. Nath, Oxidative Stress and Therapeutic Approaches in HIV
Dementia. *Antioxid Redox Sign*. 8, 2089–2100 (2006).

73. L. Agrawal, J.-P. Louboutin, E. Marusich, B. A. S. Reyes, E. J. V. Bockstaele, D. S. Strayer,
Dopaminergic neurotoxicity of HIV-1 gp120: Reactive oxygen species as signaling
intermediates. *Brain Res*. 1306, 116–130 (2010).

74. G. J. Jones, N. L. Barsby, E. A. Cohen, J. Holden, K. Harris, P. Dickie, J. Jhamandas, C.
Power, HIV-1 Vpr Causes Neuronal Apoptosis and In Vivo Neurodegeneration. *J Neurosci*. 27,
3703–3711 (2007).

75. P. Duffau, A. Ozanne, F. Bonnet, E. Lazaro, C. Cazanave, P. Blanco, E. Rivière, A.
Desclaux, C. Hyernard, N. Gensous, I. Pellegrin, L. Wittkop, Multimorbidity, age-related
comorbidities and mortality. *Aids*. 32, 1651–1660 (2018).

76. A. Edén, T. D. Marcotte, R. K. Heaton, S. Nilsson, H. Zetterberg, D. Fuchs, D. Franklin, R.
W. Price, I. Grant, S. L. Letendre, M. Gisslén, Increased Intrathecal Immune Activation in Virally
Suppressed HIV-1 Infected Patients with Neurocognitive Impairment. *Plos One*. 11, e0157160
(2016).

77. A. Winston, S. Spudich, Cognitive disorders in people living with HIV. *Lancet Hiv*. 7, e504–
e513 (2020).

78. R. T. Veenhuis, D. W. Williams, E. N. Shirk, C. M. Abreu, E. A. Ferreira, J. M. Coughlin, T.
T. Brown, P. M. Maki, K. Anastos, J. W. Berman, J. E. Clements, L. H. Rubin, Higher circulating
intermediate monocytes are associated with cognitive function in women with HIV. *Jci Insight*
(2021), doi:10.1172/jci.insight.146215.

79. A. C. Vecchio, D. W. Williams, Y. Xu, D. Yu, D. Saylor, S. Lofgren, R. O'Toole, D. R.
Boulware, N. Nakasujja, G. Nakigozi, A. Kisakye, J. Batte, R. Mayanja, A. Anok, S. J. Reynolds,
T. C. Quinn, R. H. Gray, M. J. Wawer, N. Sacktor, L. H. Rubin, Sex-specific associations
between cerebrospinal fluid inflammatory marker levels and cognitive function in antiretroviral
treated people living with HIV in rural Uganda. *Brain Behav Immun*. 93, 111–118 (2021).

80. L. H. Rubin, Y. Xu, P. J. Norris, X. Wang, R. Dastgheyb, K. C. Fitzgerald, S. M. Keating, R.
C. Kaplan, P. M. Maki, K. Anastos, G. Springer, L. Benning, S. Kassaye, D. R. Gustafson, V. G.
Valcour, D. W. Williams, Early Inflammatory Signatures Predict Subsequent Cognition in Long-
Term Virally Suppressed Women With HIV. *Frontiers Integr Neurosci*. 14, 20 (2020).

81. A. R. Kamkwalala, X. Wang, P. M. Maki, D. W. Williams, V. G. Valcour, A. Damron, P. C.
Tien, K. M. Weber, M. H. Cohen, E. E. Sundermann, V. J. Meyer, D. M. Little, Y. Xu, L. H.
Rubin, Brief Report: Higher Peripheral Monocyte Activation Markers Are Associated With
Smaller Frontal and Temporal Cortical Volumes in Women With HIV. *J Acquir Immune Defic*
Syndromes 1999. 84, 54–59 (2020).

82. D. W. Williams, D. Byrd, L. H. Rubin, K. Anastos, S. Morgello, J. W. Berman, CCR2 on
CD14(+)CD16(+) monocytes is a biomarker of HIV-associated neurocognitive disorders.
Neurology Neuroimmunol Neuroinflammation. 1, e36 (2014).

83. M. Deleidi, M. Jäggle, G. Rubino, Immune aging, dysmetabolism, and inflammation in
neurological diseases. *Front Neurosci-switz*. 9, 172 (2015).

84. T. Shabab, R. Khanabdali, S. Z. Moghadamtousi, H. A. Kadir, G. Mohan, Neuroinflammation
pathways: a general review. *Int J Neurosci*. 127, 1–10 (2016).

85. S. G. Fard, D. Cameron-Smith, A. J. Sinclair, n – 3 Docosapentaenoic acid: the iceberg n –
3 fatty acid. *Curr Opin Clin Nutrition Metabolic Care*. 24, 134–138 (2020).

86. M. Sapuła, M. Suchacz, A. Załęski, A. Wiercińska-Drapało, Impact of Combined
Antiretroviral Therapy on Metabolic Syndrome Components in Adult People Living with HIV: A
Literature Review. *Viruses*. 14, 122 (2022).

87. K. Bourgi, C. A. Jenkins, P. F. Rebeiro, B. E. Shepherd, F. Palella, R. D. Moore, K. N.
Althoff, J. Gill, C. S. Rabkin, S. J. Gange, M. A. Horberg, J. Margolick, J. Li, C. Wong, A. Willig,
V. D. Lima, H. Crane, J. Thorne, M. Silverberg, G. Kirk, W. C. Mathews, T. R. Sterling, J. Lake,
J. R. Koethe, for the N. A. A. C. C. on R. and D. (NA-ACCORD), Weight gain among treatment-
naïve persons with HIV starting integrase inhibitors compared to non-nucleoside reverse
transcriptase inhibitors or protease inhibitors in a large observational cohort in the United States
and Canada. *J Int Aids Soc*. 23, e25484 (2020).

88. P. E. Sax, K. M. Erlandson, J. E. Lake, G. A. McComsey, C. Orkin, S. Esser, T. T. Brown, J.
K. Rockstroh, X. Wei, C. C. Carter, L. Zhong, D. M. Brainard, K. Melbourne, M. Das, H.-J.
Stellbrink, F. A. Post, L. Waters, J. R. Koethe, Weight Gain Following Initiation of Antiretroviral
Therapy: Risk Factors in Randomized Comparative Clinical Trials. *Clin Infect Dis*. 71, 1379–
1389 (2019).

89. H. Stires, J. LaMori, W. Chow, Z. Zalewski, A. Vidulich, M. Avina, C. Sloan, R. Hughes, H.
Hardy, Weight Gain and Related Comorbidities Following Antiretroviral Initiation in the 2000s: A
Systematic Literature Review. *Aids Res Hum Retrov*. 37, 834–841 (2021).

90. D. M. Byonanebye, R. S. Group, Incidence of dyslipidemia in people with HIV who are
treated with integrase inhibitors versus other antiretroviral agents. *Aids*. 35, 869–882 (2021).

91. P.-E. Plum, N. Maes, A.-S. Sauvage, F. Frippiat, C. Meuris, F. Uurlings, M. Lecomte, P.
Léonard, N. Paquot, K. Fombellida, D. Vaira, M. Moutschen, G. Darcis, Impact of switch from
tenofovir disoproxil fumarate-based regimens to tenofovir alafenamide-based regimens on lipid
profile, weight gain and cardiovascular risk score in people living with HIV. *Bmc Infect Dis*. 21,
910 (2021).

92. J. Lagoutte-Renosi, M. Flammang, C. Chirouze, G. Beck-Wirth, F. Bozon, A.-S. Brunel, M.-
C. Drobacheff-Thiebaut, A. Foltzer, L. Hustache-Mathieu, J. Kowalczyk, C. Michel, S. Davani, P.
Muret, Real-Life Impact on Lipid Profile of a Switch from Tenofovir Disoproxil Fumarate to
Tenofovir Alafenamide in HIV-Infected Patients. *Curr Hiv Res*. 19, 84–89 (2021).

93. W. El-Sadr, C. Mullin, A. Carr, C. Gibert, C. Rappoport, F. Visnegarwala, C. Grunfeld, S.
Raghavan, Effects of HIV disease on lipid, glucose and insulin levels: results from a large
antiretroviral-naïve cohort. *Hiv Med*. 6, 114–121 (2005).

94. N. T. Funderburg, N. N. Mehta, Lipid Abnormalities and Inflammation in HIV Inflection. *Curr
Hiv-aids Rep*. 13, 218–225 (2016).

95. S. D. Yacovo, M. Saumoy, J. L. Sánchez-Quesada, A. Navarro, D. Sviridov, M. Javaloyas,
R. Vila, A. Vernet, H. Low, J. Peñafiel, B. García, J. Ordoñez-Llanos, D. Podzamczer, Lipids,
biomarkers, and subclinical atherosclerosis in treatment-naive HIV patients starting or not
starting antiretroviral therapy: Comparison with a healthy control group in a 2-year prospective
study. *Plos One*. 15, e0237739 (2020).

96. M. Saumoy, J. L. Sanchez-Quesada, J. Ordoñez-Llanos, D. Podzamczer, Do All Integrase
Strand Transfer Inhibitors Have the Same Lipid Profile? Review of Randomised Controlled
Trials in Naïve and Switch Scenarios in HIV-Infected Patients. *J Clin Medicine*. 10, 3456 (2021).

97. C. U. Nduka, S. Stranges, A. M. Sarki, P. K. Kimani, O. A. Uthman, Evidence of increased
blood pressure and hypertension risk among people living with HIV on antiretroviral therapy: a
systematic review with meta-analysis. *J Hum Hypertens*. 30, 355–362 (2016).

98. X. Zhao, K. Sun, Z. Lan, W. Song, L. Cheng, W. Chi, J. Chen, Y. Huo, L. Xu, X. Liu, H.
Deng, J. A. Siegenthaler, L. Chen, Tenofovir and adefovir down-regulate mitochondrial
chaperone TRAP1 and succinate dehydrogenase subunit B to metabolically reprogram glucose
metabolism and induce nephrotoxicity. *Sci Rep-uk*. 7, 46344 (2017).

99. M. Ikeda, Y. Wakabayashi, K. Okamoto, S. Yanagimoto, S. Okugawa, K. Moriya, Changing
trends in lipid profile and biomarkers of renal function and bone metabolism before and after
switching from tenofovir disoproxil fumarate to tenofovir alafenamide: a prospective
observational study. *Aids Res Ther*. 18, 30 (2021).

Figure 1.

Figure 2.

Figure 3.

Figure 4.