Investigating the ability of astrocytes to drive neural network synchrony

Gregory Handy[1,2*], Alla Borisyuk[3*]

1 Departments of Neurobiology and Statistics, University of Chicago, Chicago, Illinois, USA
2 Grossman Center for Quantitative Biology and Human Behavior, University of Chicago, Chicago, Illinois, USA
3 Department of Mathematics, University of Utah, Salt Lake City, Utah, USA

*Corresponding authors: borisyuk@math.utah.edu, ghandy@uchicago.edu

Abstract

Recent experimental works have implicated astrocytes as a significant cell type underlying several neuronal processes in the mammalian brain, from encoding sensory information to neurological disorders. Despite this progress, it is still unclear how astrocytes are communicating with and driving their neuronal neighbors. While previous computational modeling works have helped propose mechanisms responsible for driving these interactions, they have primarily focused on interactions at the synaptic level, with microscale models of calcium dynamics and neurotransmitter diffusion. Since it is computationally infeasible to include the intricate microscale details in a network-scale model, little computational work has been done to understand how astrocytes may be influencing spiking patterns and synchronization of large networks. We overcome this issue by first developing an "effective" astrocyte that can be easily implemented to already established network frameworks. We do this by showing that the astrocyte proximity to a synapse makes synaptic transmission faster, weaker, and less reliable. Thus, our "effective" astrocytes can be incorporated by considering heterogeneous synaptic time constants, which are parametrized only by the degree of astrocytic proximity at that synapse. We then apply our framework to large networks of exponential integrate-and-fire neurons with various spatial structures. Depending on key parameters, such as the number of synapses ensheathed and the strength of this ensheathment, we show that astrocytes can push the network to a synchronous state and exhibit spatially correlated patterns.

Author summary

In many areas of the brain, glial cells called astrocytes wrap their processes around synapses – the points of contact between neurons. The number of wrapped synapses and the tightness of wrapping varies between brain areas and changes during some diseases, such as epilepsy. We investigate the effect that this synaptic ensheathment has on communication between neurons and the resulting collective dynamics of the neuronal network. We present a general, computationally-efficient way to include astrocytes in neuronal networks using an "effective astrocyte" representation derived from detailed microscopic scale models. The resulting hybrid networks allow us to emulate and observe the effect of ensheathment conditions corresponding to different brain areas and

disease states. In particular, we find that it makes the networks more likely to switch into a highly correlated regime, contrary to predictions from the traditional neurons-only view. These results open a new perspective on neural network dynamics, where our understanding of conditions for generating correlated brain activity (e.g., rhythms associated with various brain functions, epileptic seizures) needs to be reevaluated.

Introduction

Recent theoretical work on neural circuits has successfully begun linking network structure to their underlying dynamics [1]. In particular, it has been shown that the statistics of connectivity (e.g., spatial scales of recurrent connectivity) can be used to predict the statistics of spiking behavior (e.g, level of network correlation) [2–4]. As reconstruction methods such as electron microscopy start to provide considerable amounts of data pertaining to the connectome of extensive brain regions [5–7], such theoretical work can guide the analysis of these massive datasets, suggesting the most useful network statistics to examine first. In this work, we develop a framework that extends computational spiking networks to efficiently include astrocytes, a non-neuronal cell that is often included in such reconstructed volumes [8]. As a result, this method allows for improving the scope and accuracy of theoretical predictions linking network structure to network dynamics.

While the overwhelming majority of computational networks only account for conventional synaptic connections across excitatory and inhibitory populations of neurons, many synapses found *in vivo* are ensheathed, or tightly wrapped around, by astrocytes [9–11]. It is well-known that astrocytes are engaged in bidirectional communication with nearby neurons, with recent breakthrough results demonstrating that astrocytes reliably encode sensory information [12]. Such communication is thought to occur through multiple pathways, such as neurotransmitter removal from the synaptic cleft, altering of extracellular ion concentrations, and the release of neuroactive substances [13–15]. The tightness of the ensheathment modulates the amount of extracellular space available around the synaptic site, and, as a result, is likely one of the contributing factors determining the properties of the synaptic connections [16] and the overall dynamics of the network. However, it is unclear how one may explore these factors computationally, as the field currently lacks an efficient method to implement such micro-scale details into a large spiking network.

Further, the presence of ensheathed and non-ensheathed synapses in a network would require a higher degree of heterogeneity than considered previously, since modeling efforts have largely focused on networks with weak heterogeneities across key synaptic parameter values to none at all [17–20]. Here, synaptic parameters would need to be drawn from distinct distributions depending on the presence of an astrocyte. Interestingly, it has been experimentally shown that the number of synapses ensheathed by astrocytes differs dramatically between brain regions (e.g., 74% of cerebellar Purkinje cell synapses, 29% of the dendritic spines in the mouse visual cortex, and 60% of hippocampal synapses are ensheathed [9, 10, 21]). Further, disease states such as epilepsy and Alzheimer's disease are associated with astrocytes altered in a variety of ways (so called "reactive astrocytes") [22–27], including alteration of their proximity to the synapse. Taken together, synaptic ensheathment by astrocytes is a source of heterogeneity in synaptic connections that needs to be accounted for to make possible predictions as to how neuronal dynamics shift across brain regions and states. Interestingly, this concept has not been included in previous mathematical studies that have investigate the so-called "tripartite synapse" (pre- and post-synaptic terminals plus neighboring astrocyte, choosing to assume the astrocyte portion does not vary from synapse to synapse [28–31].

Despite the wealth of evidence for synaptic ensheathment, the role it might play in regulating brain activity is unknown. For example, enhanced synaptic ensheathment during epilepsy in hippocampal CA3-CA1 network might be contributing to the run-away excitability and to seizures, or, conversely, they might be playing a compensatory, protective role [32]. We use computational modeling at different spatial scales to differentiate between these two options, and, more generally, to take a step towards understanding the role of astrocyte ensheathment in altering network dynamics.

As mentioned above, astrocytes have many pathways of interacting with neurons. In this work we limit ourselves to only considering two such interaction pathways, namely, the physical barrier around the synapses presented by astrocyte wrapping and a rapid removal of neurotransmitter from the synaptic cleft by glutamate transporters on the surface of astrocyte. Building on our earlier work on detailed synaptic models [33,34], we show that the ensheathed synapse can be thought of as faster and weaker than the same synapse without tight wrapping by an astrocyte. This allows us to create an "effective astrocyte ensheathment" model, which can be included in established neuronal network frameworks with only minimal increases in computational cost. Namely, an "effectively ensheathed" synapse is endowed with parameter values (synaptic time constant and synaptic strength) derived from detailed studies of a synapse at the microscopic level. These synaptic properties are considered as parameterized only by the degree of astrocytic proximity at that synapse. This approach allows us to reevaluate several important examples of neuronal networks in which the dynamics had been well-studied and understood in the absence of astrocytes. We endow some of the individual synapses with "effective astrocyte ensheathment" by altering their synaptic parameters accordingly, and study the resulting changes in network dynamics.

We start in this work by analyzing simulations of the microscale DiRT (Diffusion with Recharging Traps) model for different levels of astrocyte protrusion into the synaptic cleft. Interestingly, this model finds a strong negative linear relationship between protrusion depth and the strength and time constant of the ensheathed synapse. In other words, astrocyte ensheathment leads to weaker and faster synaptic communication across neurons. We then utilize this linear relationship and implement this result into a series of large-scale spiking models similar to those studied extensively in [2]. This implementation transforms a homogeneous network, a network with uniform synaptic parameters across populations, into a heterogeneous network, where the parameter values for an individual synapse depend explicitly on its level of astrocyte ensheathment. We investigate the impacts this novel heterogeneity has on fundamental states of the network, such as its firing rate, pairwise correlation, and stability of its dynamic states. We demonstrate that, depending on key parameters, such as the number of synapses ensheathed and the strength of this ensheathment, astrocytes can push a non-spatial network from a balanced, asynchronous firing regime into a highly synchronous one. Consistently, we then show that when neuronal assemblies are embedded into our non-spatial network, astrocyte ensheathment can enhance correlations within populations. Finally, broadening our network architectures to include models with spatially structured connections, we find that these results readily extend, namely that the spatial correlation patterns sharpen (show stronger correlation in more spatially-restricted regions) in the presence of astrocytes.

Overall, our results indicate that our understanding of what dynamical modes can be expected from a particular network of neurons needs to be reevaluated in light of synaptic heterogeneity induced by astrocyte ensheathment. Specific ensheathment-induced parameter modification for individual synapses will need to be expanded as we learn more about the effects of the synaptic ensheathment from the detailed models. In the meantime, this framework can easily be applied to other dynamic networks in other contexts using our "effective ensheathment" approach.

Models

Microscale model of neurotransmitter diffusion

We use the diffusion with recharging traps (DiRT) model [33, 34] to investigate the number of neurotransmitters that interact with receptors found on the postsynaptic terminal vs. those that diffuse out of the synaptic cleft and interact with the ensheathing astrocyte. We consider a similar idealized synapse as this previous work, taking the cleft to be a two-dimensional domain

$$\Omega^{\text{cleft}} = [0, c_w] \times [0, c_h],$$

where c_w is the width of the cleft, and c_h is the height. N_{rec} postsynaptic receptors of equal size are located along the postsynaptic density

$$\partial\Omega^{\text{cleft}}_{\text{rec}} = \{(x, y)|y = 0, x \in [\text{psd}_1, \text{psd}_2] \subseteq [0, c_w]\}.$$

The boundary of the ensheathing astrocyte is taken to be

$$\partial\Omega_{\text{astro}} = \{(x, y)|x = c_w \cdot (\phi/2) \text{ and } x = c_w \cdot (1 - \phi/2)\},$$

where $\phi < 1$ denotes the fraction of the synaptic cleft blocked due to astrocyte protrusion (Fig 1A). As a result, stronger astrocyte ensheathment corresponds to larger values of ϕ. Note that we also allow for ϕ to be negative (i.e., there exists space between the astrocytes and the synaptic cleft), in which case the domain outside of the cleft is referred to as the extracellular space Ω^{extra}.

Fig 1. Model of astrocyte ensheathment. A: Schematic of astrocyte (purple) ensheathment of a synapse (blue). The astrocyte is protruding into the synaptic cleft by amount ϕ. B: Time course of receptor activation in a diffusion with recharging traps simulation for different depths of astrocyte protrusions (red: $\phi = 0$, blue: $\phi = 0.4$) of the neighboring astrocyte, with the half-max widths denoted (dashed lines). C: Strength of synapse, defined as the area under the curve of activation time course, and D: half-max width as a function of protrusion depth. Dashed lines show a piecewise linear fit to the simulated data.

The simulation begins with N_{NT} neurotransmitters being released at $(c_w/2, c_h)$. While in the cleft or in the extracellular space, the neurotransmitters diffuse according to

$$d\mathbf{X}_k(t) = \sqrt{2D}d\mathbf{W}_k(t), k = 1, ..., N_{\text{NT}} \text{ for } \mathbf{X}_k(t) \in \left(\Omega^{\text{cleft}} \cup \Omega^{\text{extra}}\right)$$

where $\mathbf{X}_k(t)$ denotes the location of the neurotransmitter, $\mathbf{W}_j(t)$ denotes independent Wiener processes, and D is the diffusion coefficient. The receptors along the

postsynaptic terminal are taken to be partially absorbing, meaning the probability of a particle being absorbed upon making contact is

$$\text{Probability of absorption} = \frac{K\sqrt{\pi}}{\sqrt{D}}\sqrt{\Delta t},$$

where Δt is the time step of the diffusion model and K is the absorption rate. After successfully absorbing a neurotransmitter, the receptor becomes activated and switches to a transitory reflecting state and is unable to bind additional molecules. The time in this transitory state is taken to be exponentially distributed with mean $\tau_r > 0$. After this 'recharge' time, the receptor switches back to its partially absorbing state. The astrocyte boundary is taken to be perfectly absorbing, with no recharge needed between captures. We define the synaptic time course in this framework as the number of active receptors over time (Fig 1B), and the strength of the synapse to be the area under this curve. All parameter corresponding to this microscale model can be found in Table 1.

Table 1. Default parameter value for the DiRT model (arbitrary units).

Parameter	Default value	Description
N_{NT}	1000	number of neurotransmitters released
N_{rec}	50	number of postsynaptic receptors
τ_r	0.1	mean receptor recharge time
K	1	absorption rate
D	1	diffusion coefficient
c_w	1	cleft width
c_h	0.1	cleft height
$[\text{psd}_1, \text{psd}_2]$	[0.25 0.75]	postsynaptic density location
ϕ	-1 to 0.95	fraction of the cleft blocked by protruding astrocyte
Δt	0.00001	diffusion time step

Exponential integrate-and-fire neuron

Throughout this paper, we examine the effects that astrocyte ensheathment has on the network dynamics of exponential integrate-and-fire (EIF) neurons (similar to those used in [2]),

$$C_m \frac{dV_j^a}{dt} = I_L^a(V_j^a) + \psi^a(V_j^a) + I_j^a(t),$$

where V_j^a denotes the membrane potential for neuron j in population a $(= e, i)$. The first two terms on the right-hand side correspond to the leak

$$I_L^a(V) = -g_L^a(V - E_L^a),$$

and spike-generating current

$$\psi^a(V) = g_L^a \Delta_T^a \exp\left[\frac{V - V_T^a}{\Delta_T^a}\right],$$

respectively. The third term represents the synaptic input currents, and is given by

$$C_m^{-1} I_j^a(t) = F_j^a(t) + R_j^a(t),$$

where $F_j^a(t)$ represents the feedforward inputs and $R_j^a(t)$ are the recurrent inputs. We consider recurrent inputs of the form

$$R_j(t) = \sum_{b=e,i} \sum_{k=1}^{N_b} \frac{J_{jk}^{ab}}{\sqrt{N}} \left(\eta_b^{jk} * S_k^b \right)(t), \tag{1}$$

where $S_k^b(t)$ is the spike train, $\eta_b^{jk}(t)$ represents the synaptic kinetics and takes the following form

$$\eta_b^{jk}(t) = \frac{1}{\tau_b^{jk}} e^{-t/\tau_b^{jk}} \mathcal{H}(t), \tag{2}$$

and $\mathcal{H}(t)$ is the Heaviside function.

Network connectivity and feedforward input

The $1/\sqrt{N}$ scaling of the synaptic weights J_{jk}^{ab} in Eq 1 highlights that we are in the regime of strong recurrent coupling, a key feature of the so-called tightly balanced network [35, 36]. In this regime, if the firing rates of the inhibitory neurons are kept fixed, the strong coupling across excitatory neurons leads to amplification and unstable dynamics [37]. However, if recurrent connections with inhibitory neurons are sufficiently strong and fast, inhibition can effectively track and balance recurrent excitation, leading to asynchronous firing across the network [2, 36–40]. Such asynchronous firing behavior is robust to parameter choices, and is observed across networks with different connectivity rules. The focus in this work is to investigate how the dynamics established by this strong coupling might be modulated by the ensheathment of synapses by astrocytes. To do this, we consider both a randomly-connected and a spatially extended recurrent layer defined by connectivity rules previously considered in this literature. Unless otherwise noted in the figure captions, the parameter used for each network can be found in Table 2.

Non-spatial model

In the case of the non-spatial model, for each postsynaptic neuron in population b, we randomly and uniformly chose K_{eb}^{out} excitatory and K_{ib}^{out} inhibitory presynaptic neurons in the network for it to be connected to (i.e., we fixed the number of outgoing connections). The random network receives feedforward inputs as smoothly varying input biases that can target specific subpopulations. Specifically, the feedforward input to each neuron was given by the sum of an input bias and a smoothly varying signal, $F_j^a(t)$

$$F_j^a(t) = \sqrt{N} m_a + \sigma_s s(t),$$

where $s(t)$ is smooth, unbiased Gaussian noise that is shared across neurons. We take its auto-covariance function to be

$$\text{cov}(s(t), s(t+\tau)) = \exp\left(-\frac{\tau}{2\tau_s^2} \right),$$

where τ_s sets the correlation timescale and σ_s scales the magnitude of the fluctuations. We consider two cases: 1) all neurons receive the same signal $s(t)$ (Fig 2A), and 2) half of the neurons receive realization $s_1(t)$ and the other half receive $s_2(t)$ (Fig 2B).

Table 2. Default parameter value for the non-spatial and spatial model (spatial model parameters in parentheses if different).

Parameter	Default value	Description
N_e	10,000 (40,000)	number of excitatory neurons
N_i	10,000	number of inhibitory neurons
N_F	N/A (5,625)	number of ffwd neurons
$\left[C_m^e/g_l^e, C_m^i/g_l^i\right]$	[15, 10] ms	membrane time constant
V_{th}	-10 mV	spike threshold
V_T	-50 mV	soft spike threshold
E_L	-60 mV	leak potential
V_{re}	-65 mV	reset threshold
$\left[\tau_{\text{ref}}^e, \tau_{\text{ref}}^i\right]$	[1.5, 0.5] ms	refractory time
$[\tau_e, \tau_i]$	[5, 4] ([6, 5]) ms	synaptic time constants
$\left[\Delta_T^e, \Delta_T^i\right]$	[2, 0.5] mV	EIF slope parameter
τ_s	40 (N/A) ms	noise correlation timescale
$[m_e, m_i]$	[0.015, 0.01] (N/A) mV/ms	static bias to ffwd input
σ_s	0.1 (N/A) mV/ms	noise fluctuations magnitude
r_F	N/A (5) Hz	firing rate of ffwd neurons
K_{ee}^{out}	2500 (2000)	number of out-going connections from E to E neurons
K_{ie}^{out}	2500 (500)	number of out-going connections from E to I neurons
K_{ei}^{out}	2500 (2000)	number of out-going connections from I to E neurons
K_{ii}^{out}	2500 (500)	number of out-going connections from I to I neurons
J_{ee}	12.5 (40) mV	synaptic strength from E to E neurons
J_{ie}	20 (120) mV	synaptic strength from E to I neurons
J_{ei}	50 (400) mV	synaptic strength from I to E neurons
J_{ii}	50 (400) mV	synaptic strength from I to I neurons

Spatial model

To model the spatial network, we arranged N_e excitatory and N_i inhibitory neurons on a uniform grid covering a two-dimensional square domain (Fig 2C). Similarly, the feedforward layer consisted of N_F excitatory neurons uniform grid covering a square that is parallel to the recurrent network. We then normalized each grid such that the domain lied in the unit square $\Gamma = [0, 1] \times [0, 1]$. Neurons were connected randomly and the probability that two neurons were connected depended on their distance measured periodically on Γ. We used the same algorithm presented in [2] to form the connections. Given a presynaptic neuron at coordinates $y = (y_1, y_2)$ in population b and a postsynaptic neuron at $x = (x_1, x_2)$ in population a, this algorithm creates a network

Fig 2. Network schematics. We consider networks of excitatory (red triangles) and inhibitory (blue circles) neurons with different connectivity rules. Individual synapses can be ensheathed by an astrocyte (purple stars). A: One population, non-spatial network where all neurons receive the same Gaussian noise $s(t)$. B: Two population, non-spatial network where the neurons are randomly split to either receive $s_1(t)$ or $s_2(t)$. C. Spatial network on a periodic domain, where the recurrent connections can be ensheathed by astrocytes.

where the expected number of synaptic contacts from b to a is given by

$$p_{ab}(x-y) = \frac{K_{ab}^{\mathrm{out}}}{N_a} g(x_1 - y_1; \alpha_b) g(x_2 - y_2; \alpha_b) \tag{3}$$

where $g(u; \alpha)$ is a wrapped Gaussian distribution [41]. The N_F feedforward neurons are taken to be Poisson-spiking, so that the feedforward input to the recurrent layer is

$$F_j^a(t) = \sum_{k=1}^{N_F} \frac{J_{jk}^{aF}}{\sqrt{N}} \left(\eta_e * S_k^F \right)(t).$$

Results

Modeling astrocyte ensheathment with an effective astrocyte

We start by investigating how absorption of excess neurotransmitter by astrocytes can shape synaptic time courses. After neurotransmitters are released into the synaptic cleft, they either interact with receptors on the postsynaptic terminal or are absorbed by neighboring astrocytes (e.g., via glutamate transporters). While it is reasonable to assume the depth that an astrocyte protrudes into a cleft impacts the number of neurotransmitters ending up in each destination, it is unclear whether this impact is enough to significantly alter the strength of the transmitted signal. To address this unknown, we make use of the DiRT model [33, 34] to simulate particles interacting with the receptors on this microscale domain (Fig 1A). We find that the protrusion of an astrocyte noticeably decreases the synaptic time course by effectively assisting the cleft in clearing out neurotransmitters (Fig 1B, red: $\phi = 0$, blue: $\phi = 0.4$). Further, by defining the strength of the synapse to be the area under the curve of these synaptic time courses, we find that protrusion also causes the synapse to become noticeably weaker. We explore this relationship between the amount of protrusion and synaptic strength in more detail by considering a range of protrusion values ($\phi \in [-1, 0.95]$). In the extreme case where the astrocyte sits significantly away from the cleft ($\phi < -0.2$), the synaptic strength rests at a constant value (Fig 1C). As the astrocyte approaches and then begins to fill in the cleft ($\phi \in [-0.2, 0.95]$), we see that the protrusion amount and the strength vary linearly, decreasing to zero as the astrocyte completely fills in the extracellular space. This narrative also largely holds true for the half-max width of the synaptic time course (Fig 1D).

Such microscale simulations are computationally intractable to conduct in a large network of spiking neurons, but they provide insight into how one might include

"effective" astrocytes in such a model. Namely, a synapse that is ensheathed by an astrocyte should be weaker and faster than an unsheathed synapse. As a result, an ensheathed synapse should have smaller values for J and τ, which correspond to the synapse's strength and time constant respectively (see Eqs 1 and 2).

We implement these results of the microscale model in the following way. First, we let $s_{en} \in (0,1)$ denote the strength of astrocyte ensheathment in the network, where stronger astrocyte ensheathment implies a greater amount of protrusion into the synaptic cleft (i.e., $s_{en} \sim \phi$). Second, for a synaptic connection from neuron k to neuron j, we let $\mathbf{1}_{jk \text{ ensheathed}}$ be an indicator function that equals 1 if the jk synapse is ensheathed and zero otherwise. We then choose to set this synapse's strength J_{jk}^{ab} and time constant τ_b^{jk} to be

$$
\begin{aligned}
J_{jk}^{ab} &= J^{ab} \cdot \left(1 - s_{en} \cdot \mathbf{1}_{jk \text{ ensheathed}}\right), \text{ and} \\
\tau_b^{jk} &= \tau_b \cdot \left(1 - s_{en} \cdot \mathbf{1}_{jk \text{ ensheathed}}\right).
\end{aligned}
$$

This model captures the key results from the DiRT simulations, namely the synaptic strength (J_{jk}^{ab}) and the half-max width of the synaptic time course ($\tau_b^{jk} \cdot \ln(2)$) are linearly decreasing functions of the ensheathment strength (Fig 1C,D). Lastly, the probability that an excitatory (inhibitory) synapse is ensheathed is given by p_{en}^e (p_{en}^i). The difference between this and previous network models that accounted for synaptic heterogeneity is that the synaptic strength J_{jk}^{ab} and synaptic time constant τ_b^{jk} of different synapses are highly variable across the network. Though this adjustment appears deceptively subtle, allowing each synapse to have its own synaptic parameters introduces a level of heterogeneity not seen in previous work.

Utilizing this readily implementable model of astrocyte-neuron interactions, the rest of paper investigates how this heterogeneity introduced by astrocyte ensheathment may modulate the network's ability to synchronize in several commonly used configurations introduced in Models (Fig 2). Specifically, we consider non-spatial networks receiving one or two smoothly varying input biases and a spatial network driven by a feedforward layer of Poisson-spiking neurons. Since recent experimental evidence suggests that glutamatergic (i.e., excitatory) neurons are preferentially approached by astrocytes [42], we initially focus on exploring the $p_{en}^e \times s_{en}$ parameter space. Building upon these initial results, we then perform a more in-depth parameter sweep that accounts for the ensheathment of both excitatory and inhibitory synapses.

Astrocyte ensheathment can break E-I balance, causing synchronous activity

We first consider a non-spatial recurrent layer with each neuron receiving the same smoothly varying signal $s(t)$ (Fig 2A). Surprisingly, despite each neuron receiving this same input signal, previous work has shown that such a network with homogeneous parameters can exhibit robust asynchronous firing, as long as the network is placed in strongly coupled, or tightly balanced, parameter regime [2,39]. Placing our EIF network into such a regime reproduces this result, namely a network that exhibits asynchronous spiking (Fig 3A). However, introducing astrocyte ensheathment onto the excitatory synapses broke E-I balance and caused the network to show strong levels of synchrony (Fig 3B and 3C). We emphasize that all three networks in Fig 3 share the same underlying connectivity and receive the same feedforward input. Therefore, the only differences between the networks used in Fig 3B and 3C are the specific synapses that were selected for ensheathment. Together, these results demonstrate that network activity can be driven to synchrony at different times based on the subtle interactions and loss of balance between internal excitatory and inhibitory activity and the external drive.

Fig 3. Astrocyte ensheathment can lead to synchronous firing across the network.
A: The time course of the input current (top), spiking times of sampled excitatory (red) and
inhibitory (blue) neurons from the one population, non-spatial network (middle), and average
firing rate or the default network without astrocytes showing asynchronous firing rates
(bottom). B and C: same as panel A but for two network with astrocytes realizations, with
both networks exhibiting synchronous activity. Ensheathment parameters for B and C:
$p_{en}^e = 0.7$, $p_{en}^i = 0$, $s_{en} = 0.5$.

To investigate what caused the balance state to be lost, we examine the normalized
shared current across 500 neurons (Fig 4A; solid: default network, dashed: realization 1
of the network with astrocyte ensheathment from Fig 3B). Initially, we observe that the
recurrent activity in both networks tracks and cancels the feedforward input, the crucial
characteristic of a balanced network (Fig 4A; gray lines). However, we see that this
breaks down at ~1.8 seconds. At this time, the feedforward input into the networks
significantly decreases (Fig 4A; black line), and the recurrent activity in the astrocyte
network is unable to successfully cancel it. As a result, the balanced state is lost and
the network transitions into synchronous firing.

To better understand the mechanisms supporting the asynchronously firing state
and, more importantly, what causes the loss of this state in the network with astrocyte
ensheathment, we take a closer look at two time windows during this simulation
(Fig 4B). During the first time window (I), the balanced state is maintained in both
networks. We note that the normalized feedforward current is large and positive, and by
necessity, the net shared/recurrent current is negative to compensate (Fig 4Bi). This
cancellation of the feedforward input is achieved via small fluctuations in the excitatory
currents that are then effectively tracked and balanced by the inhibitory currents
accordingly (Fig 4Bii and 4Biii).

However, during the second time window (II), the normalized feedforward current
drops, which must be met with an increase in the net shared current. This increase runs

Fig 4. Astrocyte ensheathment can break E-I balance. A: Normalized shared fluctuations in the feedforward (black) and recurrent (purple) synaptic inputs for two one population, non-spatial networks (solid: default; dashed: realization 1 from Fig 3B). Initially, both networks exhibit recurrent dynamics that effectively cancel the feedforward inputs, leading to weak shared fluctuations (gray), until the balance state is lost in the network with astrocyte ensheathment. The curves were computed by averaging the synaptic input currents to 500 neurons, convolving with a Gaussian-shaped kernel ($\sigma = 15$ ms), subtracting the mean and dividing by the neurons' rheobase. B: Normalized shared fluctuations in the currents zoomed in on a time window when the asynchronous state is maintained in both networks (left column) and for the time window when the asynchronous state is lost in the network with astrocyte ensheathment (right column). The following currents are displayed: i) feedforward, all recurrent, total, ii) recurrent excitatory, and iii) recurrent inhibitory currents in the default network (solid) and the network with astrocyte ensheathment (dashed). C: Percent of the ten EIF networks that led to synchronous dynamics in 5 seconds of simulation as a function of ensheathment parameters. D: The $\hat{s}_{en} = s_{en} \cdot p_{en}^e = 0.4$ isoline showing that a naïve mean field approximation would fail at capturing this phenomenon. Colorbar is the same as panel C. $p_{en}^i = 0$ for all panels.

the risk of leading to uncontrolled runaway excitation, and as a result, synchronous   275
firing. We see that the default network's inhibitory currents are able to continue   276
tracking and balancing the excitatory currents, preventing this from occurring.   277
Meanwhile, the network with astrocyte ensheathment struggles to make these fine tuned   278
adjustments, which leads to growing oscillations in both the excitatory and inhibitory   279
currents. Interestingly, we note that the second realization of the astrocyte   280
ensheathment network (Fig 3C) does not fall into synchrony at this point in the   281
simulation, and instead is able to maintain the balanced state during this drop in the   282

feedforward current (S1 Fig D, center column) due to random differences between realizations (i.e., the specific synapses chosen for astrocyte ensheathment). Nonetheless, it fails later on in the simulation when the feedforward input reaches a more extreme local minimum (S1 Fig D, right column).

To explain this loss of the asynchronous state, we recall that our model of astrocyte ensheathment leads to faster and weaker synapses. As noted previously, the balanced state relies on inhibition to be sufficiently fast relative to excitation [39, 40]. Here, astrocytes are specifically targeting excitatory synapses, so the net effect is known: a subset of excitatory synapses throughout the network now have faster synaptic dynamics. As these simulations then suggest, their inhibitory counterparts are unable to effectively track their activity and the network experiences runaway excitation as the balanced state is lost. While consistent with previous work, our results show that adjusting the synaptic kinetics in only a *subset* of neurons is sufficient for the entire network to enter into a state of synchronous firing.

We now explore how the parameters underlying astrocyte ensheathment (i.e., the probability (p_{en}^e) and strength (s_{en}) of ensheathment of excitatory synapses) determine the size of subset needed to lead to synchronization and whether a mean-field approximation can be used to average across this heterogeneity and capture this effect. We consider ten default EIF networks, each with a different input signal $s(t)$ and no astrocyte ensheathment. These ten networks all exhibited asynchronous spiking solutions for the full length of the trial (taken to be 5 seconds). We then ask whether or not these same networks synchronized in the presence of astrocyte ensheathment. Discretizing the parameter space $p_{en}^e \times s_{en} = [0.1, 1] \times [0.2, 0.8]$ into seventy parameter sets, we observe that all ten networks eventually exhibit synchronous behavior as p_{en}^e and s_{en} increase (Fig 4C; red dots in upper right corner). However, we also observe that there is not a sharp transition point from blue (all networks remain in the asynchronous state) to red (all networks become synchronous). This suggests that for intermediate values of p_{en}^e and s_{en}, random differences in the connectivity diagram and the locations of astrocyte ensheathment are enough to prime some networks to tend toward synchrony over others.

These results highlight the shortcomings that a naïve mean field theory would have in capturing the dynamics brought on by this type of synaptic heterogeneity. More specifically, one might be tempted to simply average out the effect of astrocyte ensheathment across the network and adjust the strength of all synapses to be

$$\hat{s}_{en} = s_{en} \cdot p_{en}^e + 0 \cdot (1 - p_{en}^e).$$ (4)

However, by considering the outcomes across the $\hat{s}_{en} = 0.4$ iso-line (Fig 4D), the heterogeneous networks may rarely exhibit synchrony (low p_{en}^e, high s_{en}) or frequently (high p_{en}^e, low s_{en}).

Astrocyte ensheathment can enhance correlations already present within a network

The previous section explored how astrocyte ensheathment of excitatory synapses affects the dynamics of a balanced network operating in the asynchronous regime. Now we examine a network with the same non-spatial connectivity, but the neurons are split into two subpopulations, where each subpopulation receives its own feedforward input (Fig 2B). As previously shown in [2], this change in feedforward inputs leads to a dramatic shift away from the asynchronous state, with positive correlations arising among neurons within the same population and negative correlations across populations. As a result, the correlation distribution becomes strongly binomial (Fig 5A,B; solid lines). While we expect the results of the previous section to carry through, namely the

loss of stability for the strong ensheathment of excitatory synapses, it is unclear how the ³³¹
heterogeneity introduced by more moderate levels of ensheathment will modulated these ³³²
new correlations inherently found in the network.

Fig 5. Astrocyte ensheathment can lead to greater correlations. A: Normalized
histogram of pairwise spike count correlations between 1000 neurons for the default two
population, non-spatial network (solid) and the same network with astrocytes (dashed), with
parameters $(p_{en}^e, s_{en}) = (0.8, 0.5)$. B: Same as panel A, except the pairwise spike count
correlations calculated within and across each populations. C: Mean pairwise spike count
correlations within the same population as a function of ensheathment parameters. Gray
regions denote areas where the network lost stability. D: Raster plots for the two populations
for the i) default network and ii) network with astrocytes $(p_{en}^e, s_{en}) = (0.8, 0.5)$. E: Mean spike
count correlations plotted against the excitatory firing rate for different astrocyte conditions
(the color of dots corresponds to $p_{en}^e \cdot s_{en}$ for that simulation). $p_{en}^i = 0$ for all panels.

³³³

Setting $p_{en}^e = 0.8$ and $s_{en} = 0.5$, we find that the binomial distribution of spike ³³⁴
count correlations present in the default network becomes more extreme, with an ³³⁵
increase in the positive correlation among the same population and a decrease in the ³³⁶
negative correlations across populations (Fig 5A,B; dashed lines). This trend is true in ³³⁷
general: as p_{en}^e and s_{en} increase, the correlations among a single population also ³³⁸
increase (Fig 5C), while the correlations across different populations decrease (figure not ³³⁹
shown). As anticipated, we also find that the network stability is loss for more extreme ³⁴⁰
values of astrocyte ensheathment (Fig 5C, gray boxes). ³⁴¹

To gain additional insight into how astrocytes are modulating the spiking behavior ³⁴²
of the network, we compare the raster plot of the default network to a network with ³⁴³
astrocytes, where both networks receive the same feedforward inputs (Fig 5D). As ³⁴⁴
expected from the presence of negative correlations across populations (Fig 5B), the two ³⁴⁵
populations show clear competitive dynamics. Interestingly, the network with astrocyte ³⁴⁶
ensheathment shows a much more subtle modulation of network dynamics than our ³⁴⁷
previous results. In fact, it is clear from these raster plots that key quantitative traits, ³⁴⁸
such as the times the dominate spiking populations switches, are largely conserved ³⁴⁹
across the two conditions. ³⁵⁰

One observable difference between the two networks is the population firing rate: both populations in the network with astrocyte ensheathment appears to have higher firing rates. Previous work [43] suggests that this difference in firing rates could be sufficient to explain the increase in correlations. Indeed, we find a strong positive relationship ($R^2 = 0.96$) between the spike count correlations and the excitatory firing rate, with higher firing rates arising with stronger astrocyte ensheathment (Fig 5E; color corresponding to $p_{en}^e \cdot s_{en}$). This strong relationship between firing rate, correlations and astrocyte ensheathment suggest that unlike the last network configuration, a mean-field model utilizing the parameter defined in Eq 4 could be used to capture the effect of astrocytes on the neuronal dynamics. See Discussion for additional considerations regarding a mean-field approach.

Spatial correlations are enhanced by astrocyte ensheathment

In the previous two sections, we showed that the ensheathment of excitatory synapses by astrocytes strongly modulated the underlying correlations across neurons in a non-spatial network. However, it is unclear the impact this ensheathment may have on networks with more realistic connectivity. To pursue this, we now implement our "effective" astrocytes in a spatial model (Fig 2C). Specifically, we arrange the recurrently connected neurons in a two dimensional grid, with a Gaussian connectivity profile (see Eq 3), the spread of which is given by parameter α_{rec}. Unlike the previous sections, the feedforward inputs are provided by neurons that are Poisson-spiking. These neurons also lie on a two-dimensional grid and make connections based on a Gaussian connectivity profile with spread parameter α_{ff}. We continue to consider astrocyte ensheathment of only recurrent excitatory synapses.

In agreement with previous work [2], we find that when the spatial profile of recurrent connections is less than the feedforward (i.e., $\alpha_{\mathrm{rec}} < \alpha_{\mathrm{ff}}$), the default network exists in a spatially asynchronous state, with small pairwise spike count correlations between neurons of any distance (Fig 6A, left and 6B, black). In fact, [2] proves that the asynchronous state requires $\alpha_{\mathrm{rec}} < \alpha_{\mathrm{ff}}$. However, by introducing astrocyte ensheathment to this model ($p_{en}^e = 0.9$ and $s_{en} = 0.5$), we find that the asynchronous state can be broken without changing the spatial profiles of connectivity. Specifically, we observe strong positive correlations arising between nearby neurons and negative correlations from those farther away (Fig 6B, orange), resulting in tight clusters of spiking neurons (Fig 6A, right). We note these tight clusters do not persist across time (S1 Video).

These results mirror our earlier results corresponding to the non-spatial network with a single population of neurons (Fig 3), and arise from similar mechanisms. Specifically, astrocyte ensheathment speeds up excitatory synapses, and as a result, the inhibitory synapses are unable to track and cancel recurrent excitation, leading to a loss of the asynchronous balance state. However, unlike the result for the non-spatial network, the network activity organizes in tight spatial patterns, as opposed the synchronous firing of the entire network.

In further agreement with [2], when the recurrent connection's spatial profile is larger than the feedforward projections (i.e., $\alpha_{\mathrm{rec}} > \alpha_{\mathrm{ff}}$), we find that the default network now exhibits spatial correlations (Fig 6C, black). In this case, astrocyte ensheathment of excitatory synapses does not qualitatively change this behavior, but instead simply enhances these spatial correlations at almost all distances (Fig 6C, orange), with the exception being the distances where spatial correlations are zero. This is true for a range of astrocyte ensheathment parameters: stronger astrocyte ensheathment, either through p_{en}^e or s_{en}, leads to stronger spatial correlations, while not affecting the spatial footprint of these correlations (S2 Fig). In this case, with the asynchronous state already broken due to the widths of the spatial projections, astrocyte ensheathment of

Fig 6. Astrocyte ensheathment can lead to greater spatial correlations. A: Spatial raster plots at three 20 msec time points for the default network (left) and network with astrocyte ensheathment (right). Parameters were chosen such that the default network exhibited no spatial correlations ($\alpha_{\mathrm{rec}} = 0.05$ and $\alpha_{\mathrm{ffwd}} = 0.1$). Ensheathment parameters are $p_{en}^{e} = 0.9$ and $s_{en} = 0.5$. B: Spike count correlation plots as a function of distance between neuron pairs corresponding to the parameters in panel A for the default (black) and network with astrocytes (orange). C: Same as panel B, except with a default network placed in the spatial correlation regime ($\alpha_{\mathrm{rec}} = 0.2$ and $\alpha_{\mathrm{ffwd}} = 0.1$). Here, the ensheathment parameters are $p_{en}^{e} = 0.8$ and $s_{en} = 0.5$). $p_{en}^{i} = 0$ for all panels.

excitatory synapses simply amplifies the inability of inhibitory neurons to effectively track and cancel recurrent excitation across all spatial scales.

Astrocyte ensheathment of both excitatory and inhibitory synapses can boost stability and within-population correlations

Thus far, we have explored the effects on network dynamics when excitatory synapses are targeted by astrocyte ensheathment. We found that across networks with different connectivity rules and feedforward input, the neuronal dynamics shifted towards a more correlated regime. However, despite evidence suggesting that excitatory synapses are preferentially targeted [42], it is important to note that inhibitory neurons can and are ensheathed, and that the level of ensheathment may vary by the particular brain region. For this reason, we now consider astrocyte ensheathment that can target both types of synapses, expanding our parameter search over the larger $p_{en}^{e} \times p_{en}^{i} \times s_{en}$ space to examine whether our previous results carry through.

For this, we return to the non-spatial model with two subpopulations that each receive its own feedforward input (Fig 2B). Recall that in this framework, the within

population correlations were positive and astrocyte ensheathment of excitatory synapses ₄₁₇
increased these correlations (Fig. 7A, top left). Across different values of p_{en}^i, we see ₄₁₈
that this relationship is not only intact, but amplified: As p_{en}^i increases, the within ₄₁₉
population correlations also increase. Further, we note the network's stability is more ₄₂₀
robust, with the higher levels of p_{en}^i (i.e., ≥ 0.4) lacking the unstable gray regions. ₄₂₁
Lastly, we investigated whether the relationship across correlations, excitatory firing ₄₂₂
rate, and average ensheathment level of excitatory synapses ($p_{en}^e \cdot s_{en}$) remains. Indeed, ₄₂₃
we again find a strong linear relationship between correlations and firing rate, and both ₄₂₄
metrics with a clear correspondence to the strength of astrocyte ensheathment (Fig 7B). ₄₂₅
In total, these results suggest that while the ensheathment of inhibitory neurons in ₄₂₆
addition to excitatory do not drastically alter our earlier results, stability and ₄₂₇
correlations are both reliably enhanced. As a result, this could be a key mechanism ₄₂₈
utilized by the brain to support high levels of correlation while preventing pathological ₄₂₉
states from arising. ₄₃₀

Fig 7. Investigation of neuronal dynamics when astrocyte ensheathment targets both excitatory and inhibitory synapses. A: Mean pairwise spike count correlations within the same population as a function of ensheathment parameters. Gray regions denote areas where the network lost stability. B: Mean spike count correlations plotted against the excitatory firing rate for different astrocyte conditions (the color of dots corresponds to $p_{en}^e \cdot s_{en}$ for that simulation).

Discussion ₄₃₁

Through microscale modeling of neurotransmitter diffusion, this work derives an efficient ₄₃₂
way to implement astrocytes ensheathment of synapses in large scale neuronal networks. ₄₃₃
Specifically, simulations on the microscale domain of the synaptic cleft suggested that ₄₃₄
ensheathed synapses are weaker and faster than their non-ensheathed counterparts. ₄₃₅
This investigation identified the synaptic strength and synaptic time constant as the key ₄₃₆
parameters to vary to denote the presence of an "effective" astrocyte. ₄₃₇

While we have chosen to consider a network of EIF neurons, such parameters are ₄₃₈
commonly found across different types of recurrently connected spiking networks (e.g., ₄₃₉
networks of current- and conductance-base neurons, with and without ₄₄₀
adaptation [37, 40, 44–46]), making the implementation presented here applicable to ₄₄₁
many network architectures and parameter regimes. We showed this specifically by ₄₄₂

examining the consequences of including these effective astrocytes in both non-spatial and spatial networks placed in the regime of strong recurrent coupling, which is also referred to as the "tight balanced" regime [36]. In this regime, we uncovered that astrocyte ensheathment can dramatically change firing dynamics, break the balance state, and cause synchronous behavior. While this parameter regime yields a type of inhibition-stabilized network (ISN) [47], meaning the fixed point is stabilized by feedback inhibition, it is unclear whether our results extend to other types of ISN balanced networks (e.g., the "loose balance" regime, also referred to as a stabilized supralinear network [36, 48]) and non-ISN parameter regimes. For example, we were able to explain the synchronous firing in our non-spatial network (Fig 3) as due to the astrocytes targeting and speeding up excitatory synapses, which the inhibitory currents were unable to track and cancel. In these other parameter regimes, this tight canceling does not occur, so it is unclear if similar synchronous firing would arise. Future work will push these lines of inquiry further, made easier in part by the straightforward implementation of our astrocyte framework.

The investigation of the balance state as a function of synaptic time courses was also completed recently in [40], but our work differs in a few notable areas. First, the adjustment of the synaptic time courses here arose from the implementation of astrocyte ensheathment around individual synapses. This resulted in a heterogeneous network, which behaves qualitatively differently than its homogeneous network counterpart found by a naïve mean field (see Fig 4). Second, we observed spatial patterns emerge in the $\alpha_{\text{rec}} < \alpha_{\text{ff}}$ regime when astrocyte ensheathment was present, showing that this condition, as derived in [2], is a necessary condition for the balance state, but not a sufficient one. Meanwhile [40] only considered homogeneous networks in the $\alpha_{\text{rec}} > \alpha_{\text{ff}}$ regime.

We also note that the heterogeneity introduced by astrocyte ensheathment considered here is significantly different than other models that consider heterogeneous parameters. For example, previous works have implemented heterogeneity in connection strengths by having the J_{jk}^{ab}'s be randomly distributed according to the normal distribution $\mathcal{N}(J, \sigma)$ [18–20], resulting in only small variations of parameters across the network. Here, the synaptic parameters corresponding to ensheathed and unsheathed synapses are pulled from entirely distinct distributions, specifically

$$
(J_{jk}^{ab}, \tau_b^{jk}) \quad = \begin{cases} (J^{ab}, \tau_b) & \text{if synapse } k \to j \text{ is unsheathed} \\ (J^{ab} \cdot (1 - s_{en}), \tau_b \cdot (1 - s_{en})) & \text{if synapse } k \to j \text{ is ensheathed} \end{cases}.
$$

In the future work, it would be important to extend this heterogeneity even further, either by considering distributions for the synaptic parameters as done in previous works, or by considering a distribution on the ensheathment parameter s_{en}. This would allow various degrees of astrocyte ensheathment across a single network and potentially across different populations (i.e., define s_{en}^{ab} as the strength of ensheathment from population b to a). In addition, since astrocytes segregate the space into non-overlapping control domains, which can also be disrupted in epilepsy [49]), it would also be interesting and important to perform a more thorough investigation that includes spatial inhomogeneities across all ensheathment parameters for the spatial network. Further, it may also be interesting to consider a time varying component to these ensheathment parameters. In fact, along with increasing the number of postsynaptic receptors [50], long-term potentiation induction has recently been shown to decrease the ensheathment level of astrocytes [51]. While these two mechanisms operate on a much slower timescale as considered in this work, as shown here and in our previous work [33], they would work together to strengthen and slow synapses. Including all of these additional forms of heterogeneities, spatially and temporally, would lead to a more robust understanding of how astrocytes are modulating network dynamics via ensheathment.

Additionally, we found that it is non-trivial to capture this novel form of network

heterogeneity in a mean-field approximation, complicating the extension of this ₄₉₂ framework to be included in rate-based models of average population dynamics. ₄₉₃ Specifically, when we considered astrocyte ensheathment of excitatory synapses, we ₄₉₄ showed that a naïve mean-field theory (see Eq 4) would fail to capture key qualitative ₄₉₅ dynamics of the full model (Fig 4D). On the other hand, when we allowed astrocytes to ₄₉₆ equally ensheathed both excitatory and inhibitory synapses, we found a tight ₄₉₇ relationship between correlations, firing rates, and astrocyte ensheathment (Fig 5E), ₄₉₈ suggesting that a mean-field approximation in this parameter regime could be possible. ₄₉₉ So while a mean-field approach could work in the latter case, it would be interesting ₅₀₀ future work to investigate how robust such an approximation would be in the presence ₅₀₁ of symmetry breaking changes in parameters, either in the probability and/or strength ₅₀₂ of ensheathment. ₅₀₃

Lastly, our simplified model of astrocyte-neuron interactions at the synaptic level ₅₀₄ only accounts for the ability of astrocytes to assist in the clearance of neurotransmitters ₅₀₅ from the synaptic cleft by acting as an effective buffer and transporter [13, 52], but ₅₀₆ other astrocyte-neuron communication pathways are likely to play a role in modulating ₅₀₇ neuronal dynamics. For example, neurotransmitters may also activate receptors found ₅₀₈ on astrocyte membranes (e.g., mGluR5), leading to the activation of calcium ₅₀₉ pathways [53, 54]. While the downstream effects of this pathway are still being ₅₁₀ investigated, evidence suggests that it may impact other aspects of neuronal signaling ₅₁₁ not considered in this work, such as the overall excitability of the network by either ₅₁₂ adjusting the resting potential of neurons [55] or by modifying neurotransmitter profile ₅₁₃ in the extracellular space through the direct release of gliotransmitters [15] (but ₅₁₄ see [56–58]). Here, we show the unanticipated result that this calcium pathway is not ₅₁₅ needed for astrocytes to strongly tune network dynamics. ₅₁₆

Astrocytes are also known to have a role in restoring the resting membrane potential ₅₁₇ of nearby neurons and preventing hyperexcitability through the removal of potassium ₅₁₈ via Kir4.1 channels [59]. It is therefore possible that increased ensheathment, as ₅₁₉ considered in this work, might result in improved potassium removal by bringing ₅₂₀ astrocyte Kir4.1 closer, which may counteract to some extent the tendency to ₅₂₁ synchronize. On the other hand, rapid reduction of extracellular potassium would also ₅₂₂ render glutamate removal more efficient [60, 61] making the synchronization that we ₅₂₃ observe here even more likely. Lastly, our model suggests that tight ensheathment by ₅₂₄ astrocytes would drastically decrease the amount of neurotransmitters that 'spillover' ₅₂₅ into the extrasynaptic space, which is in alignment with experimental ₅₂₆ observations [51, 62]. Since such spillover is known to allow for cross-talk between ₅₂₇ neurons, it is another mechanism that could set the excitability of the network [63, 64]. ₅₂₈ Future work should seek to connect these additional communication pathways to the ₅₂₉ ensheathment interaction investigated here to further advance our understanding of how ₅₃₀ astrocyte-neuron interactions modulate spiking dynamics across brain regions. ₅₃₁

Supporting information ₅₃₂

S1 Video. Video of spatial raster plots showing tight clusters of spiking ₅₃₃ **neurons in the network with astrocytes and asynchronous spiking in the** ₅₃₄ **default network.** Will be made available upon acceptance for publication. ₅₃₅

S1 Fig. Broken E-I balance in Realization 2. A: Normalized shared fluctuations in the feedforward (black) and recurrent (purple) synaptic inputs for two one population, non-spatial networks (solid: default; dashed: realization 2 from Fig 3C). B,C: Spiking times of sampled excitatory (red) and inhibitory (blue) neurons from the default network and network with astrocyte ensheathment, respectively. D: Normalized shared fluctuations in the currents zoomed in on time windows when the asynchronous state is maintained in both networks (left and middle columns) and for the time window when the asynchronous state is lost in the network with astrocyte ensheathment (right column). The following currents are displayed: i) feedforward, all recurrent, total, ii) recurrent excitatory, and iii) recurrent inhibitory currents in the default network (solid) and the network with astrocyte ensheathment (dashed). Same parameters as Fig 3-4.

S2 Fig. Spatial correlations as a function of ensheathment parameters. Spike count correlations as a function of distance for different probabilities of ensheathment (A: $p_{en}^{ae} = 0.9$, B: $p_{en}^{ae} = 0.85$, C: $p_{en}^{ae} = 0.8$) and ensheathment strengths ($s_{en} \in [0.35, 0.5]$), where the default network exhibits spatial correlations ($\alpha_{\text{rec}} = 0.2$ and $\alpha_{\text{ffwd}} = 0.1$).

Data availability

The code used to generate the main figures in the manuscript is available on the GitHub
database (https://github.com/gregoryhandy/Neuronal_Net_with_Astro_Ensheathment).

Acknowledgments

G.H. would like to acknowledge support from the Swartz Foundation Fellowship for 540
Theory in Neuroscience and the Burroughs Wellcome Fund's Career Award at the
Scientific Interface. A.B. would like to acknowledge support from National Science
Foundation award NSF-DMS-1853673 and the support and resources from the Center
for High Performance Computing at the University of Utah.

References

1. Ocker GK, Hu Y, Buice MA, Doiron B, Josić K, Rosenbaum R, et al. From the Statistics of Connectivity to the Statistics of Spike Times in Neuronal Networks. Current Opinion in Neurobiology. 2017;46:109–119. doi:10.1016/j.conb.2017.07.011.

2. Rosenbaum R, Smith MA, Kohn A, Rubin JE, Doiron B. The Spatial Structure of Correlated Neuronal Variability. Nature Neuroscience. 2017;20(1):107–114. doi:10.1038/nn.4433.

3. Hu Y, Trousdale J, Josić K, Shea-Brown E. Motif Statistics and Spike Correlations in Neuronal Networks. Journal of Statistical Mechanics: Theory and Experiment. 2013;2013(03):P03012. doi:10.1088/1742-5468/2013/03/P03012.

4. Trousdale J, Hu Y, Shea-Brown E, Josić K. Impact of Network Structure and Cellular Response on Spike Time Correlations. PLoS Computational Biology. 2012;8(3):e1002408. doi:10.1371/journal.pcbi.1002408.

5. Lee WCA, Bonin V, Reed M, Graham BJ, Hood G, Glattfelder K, et al. Anatomy and Function of an Excitatory Network in the Visual Cortex. Nature. 2016;532(7599):370–374. doi:10.1038/nature17192.

6. Kasthuri N, Hayworth KJ, Berger DR, Schalek RL, Conchello JA, Knowles-Barley S, et al. Saturated Reconstruction of a Volume of Neocortex. Cell. 2015;162(3):648–661. doi:10.1016/j.cell.2015.06.054.

7. Helmstaedter M, Briggman KL, Turaga SC, Jain V, Seung HS, Denk W. Connectomic Reconstruction of the Inner Plexiform Layer in the Mouse Retina. Nature. 2013;500(7461):168–174. doi:10.1038/nature12346.

8. Consortium M, Bae JA, Baptiste M, Bodor AL, Brittain D, Buchanan J, et al. Functional Connectomics Spanning Multiple Areas of Mouse Visual Cortex. bioRxiv. 2021; p. 2021.07.28.454025.

9. Špaček J. Three-Dimensional Analysis of Dendritic Spines. Anatomy and Embryology. 1985;171(2):245–252. doi:10.1007/BF00341419.

10. Ventura R, Harris KM. Three-Dimensional Relationships between Hippocampal Synapses and Astrocytes. The Journal of neuroscience : the official journal of the Society for Neuroscience. 1999;19(16):6897–906.

11. Pannasch U, Freche D, Dallérac G, Ghézali G, Escartin C, Ezan P, et al. Connexin 30 Sets Synaptic Strength by Controlling Astroglial Synapse Invasion. Nature Neuroscience. 2014;17(4):549–558. doi:10.1038/nn.3662.

12. Doron A, Rubin A, Benmelech-Chovav A, Benaim N, Carmi T, Refaeli R, et al. Hippocampal Astrocytes Encode Reward Location. Nature. 2022;doi:10.1038/s41586-022-05146-6.

13. Tzingounis AV, Wadiche JI. Glutamate Transporters: Confining Runaway Excitation by Shaping Synaptic Transmission. Nature reviews Neuroscience. 2007;8(12):935–47. doi:10.1038/nrn2274.

14. Anderson CM, Swanson RA. Astrocyte Glutamate Transport: Review of Properties, Regulation, and Physiological Functions. Glia. 2000;32(1):1–14. doi:10.1002/1098-1136(200010)32:1¡1::AID-GLIA10¿3.0.CO;2-W.

15. Covelo A, Araque A. Neuronal Activity Determines Distinct Gliotransmitter Release from a Single Astrocyte. eLife. 2018;7:1–19. doi:10.7554/eLife.32237.

16. Herde MK, Bohmbach K, Domingos C, Vana N, Komorowska-Müller JA, Passlick S, et al. Local Efficacy of Glutamate Uptake Decreases with Synapse Size. Cell Reports. 2020;32(12):108182. doi:10.1016/j.celrep.2020.108182.

17. Van Vreeswijk C, Abbott LF, Bard Ermentrout G. When Inhibition Not Excitation Synchronizes Neural Firing. Journal of Computational Neuroscience. 1994;1(4):313–321. doi:10.1007/BF00961879.

18. Rajan K, Abbott LF. Eigenvalue Spectra of Random Matrices for Neural Networks. Physical Review Letters. 2006;97(18):188104. doi:10.1103/PhysRevLett.97.188104.

19. Rajan K, Abbott LF, Sompolinsky H. Stimulus-Dependent Suppression of Chaos in Recurrent Neural Networks. Physical Review E. 2010;82(1):011903. doi:10.1103/PhysRevE.82.011903.

20. Mastrogiuseppe F, Ostojic S. Linking Connectivity, Dynamics, and Computations in Low-Rank Recurrent Neural Networks. Neuron. 2018;99(3):609–623.e29. doi:10.1016/j.neuron.2018.07.003.

21. Lippman JJ, Lordkipanidze T, Buell ME, Yoon SO, Dunaevsky A. Morphogenesis and Regulation of Bergmann Glial Processes during Purkinje Cell Dendritic Spine Ensheathment and Synaptogenesis. Glia. 2008;56(13):1463–1477. doi:10.1002/glia.20712.

22. Volman V, Bazhenov M, Sejnowski TJ. Computational Models of Neuron-Astrocyte Interaction in Epilepsy. Frontiers in Computational Neuroscience. 2012;6. doi:10.3389/fncom.2012.00058.

23. Coulter SC Douglas. Role of Astrocytes in Epilpesy. Cold Spring Harb Perspect Med. 2015;5(a022434):1–13. doi:10.1038/clpt.2011.183.

24. Heneka MT, Carson MJ, Khoury JE, Landreth GE, Brosseron F, Feinstein DL, et al. Neuroinflammation in Alzheimer's Disease. The Lancet Neurology. 2015;14(4):388–405. doi:10.1016/S1474-4422(15)70016-5.

25. Matias I, Morgado J, Gomes FCA. Astrocyte Heterogeneity: Impact to Brain Aging and Disease. Frontiers in Aging Neuroscience. 2019;11:59. doi:10.3389/fnagi.2019.00059.

26. Verhoog QP, Holtman L, Aronica E, van Vliet EA. Astrocytes as Guardians of Neuronal Excitability: Mechanisms Underlying Epileptogenesis. Frontiers in Neurology. 2020;11:591690. doi:10.3389/fneur.2020.591690.

27. Price BR, Johnson LA, Norris CM. Reactive Astrocytes: The Nexus of Pathological and Clinical Hallmarks of Alzheimer's Disease. Ageing Research Reviews. 2021;68:101335. doi:10.1016/j.arr.2021.101335.

28. Wade J. Self-Repair in a Bidirectionally Coupled Astrocyte-Neuron (AN) System Based on Retrograde Signaling. Frontiers in Computational Neuroscience. 2012;6(September):1–12. doi:10.3389/fncom.2012.00076.

29. Reato D, Cammarota M, Parra LC, Carmignoto G. Computational Model of Neuron-Astrocyte Interactions during Focal Seizure Generation. Frontiers in computational neuroscience. 2012;6(October):81. doi:10.3389/fncom.2012.00081.

30. Huguet G, Joglekar A, Messi LM, Buckalew R, Wong S, Terman D. Neuroprotective Role of Gap Junctions in a Neuron Astrocyte Network Model. Biophysical Journal. 2016;111(2):452–462. doi:10.1016/j.bpj.2016.05.051.

31. Garnier A, Vidal A, Benali H. A Theoretical Study on the Role of Astrocytic Activity in Neuronal Hyperexcitability by a Novel Neuron-Glia Mass Model. The Journal of Mathematical Neuroscience. 2016;6(1):10. doi:10.1186/s13408-016-0042-0.

32. Umpierre AD, West PJ, White JA, Wilcox KS. Conditional Knock-out of mGluR5 from Astrocytes during Epilepsy Development Impairs High-Frequency Glutamate Uptake. Journal of Neuroscience. 2019;39(4):727–742. doi:10.1523/JNEUROSCI.1148-18.2018.

33. Handy G, Lawley SD, Borisyuk A. Role of Trap Recharge Time on the Statistics of Captured Particles. Physical Review E. 2019;99(2):022420. doi:10.1103/PhysRevE.99.022420.

34. Handy G, Lawley SD, Borisyuk A. Receptor Recharge Time Drastically Reduces the Number of Captured Particles. PLOS Computational Biology. 2018;14(3):e1006015. doi:10.1371/journal.pcbi.1006015.

35. Van Vreeswijk C, Sompolinsky H. Chaos in Neuronal Networks with Balanced Excitatory and Inhibitory Activity. Science. 1996;doi:10.1126/science.274.5293.1724.

36. Ahmadian Y, Miller KD. What Is the Dynamical Regime of Cerebral Cortex? Neuron. 2021;109(21):3373–3391. doi:10.1016/j.neuron.2021.07.031.

37. Sanzeni A, Histed MH, Brunel N. Emergence of Irregular Activity in Networks of Strongly Coupled Conductance-Based Neurons. 2020; p. 1–37.

38. Amit DJ, Brunel N. Model of Global Spontaneous Activity and Local Structured Activity during Delay Periods in the Cerebral Cortex. Cerebral Cortex. 1997;7(3):237–252. doi:10.1093/cercor/7.3.237.

39. Renart A, De La Rocha J, Bartho P, Hollender L, Parga N, Reyes A, et al. The Asynchronous State in Cortical Circuits. Science. 2010;327(5965):587–590. doi:10.1126/science.1179850.

40. Huang C, Ruff DA, Pyle R, Rosenbaum R, Cohen MR, Doiron B. Circuit Models of Low-Dimensional Shared Variability in Cortical Networks. Neuron. 2019;101(2):337–348.e4. doi:10.1016/j.neuron.2018.11.034.

41. Rosenbaum R, Doiron B. Balanced Networks of Spiking Neurons with Spatially Dependent Recurrent Connections. Physical Review X. 2014;4(2):1–9. doi:10.1103/PhysRevX.4.021039.

42. Brunskine C, Passlick S, Henneberger C. Structural Heterogeneity of the GABAergic Tripartite Synapse. Cells. 2022;11(19):3150. doi:10.3390/cells11193150.

43. de la Rocha J, Doiron B, Shea-Brown E, Josić K, Reyes A. Correlation between Neural Spike Trains Increases with Firing Rate. Nature. 2007;448(7155):802–806. doi:10.1038/nature06028.

44. Li B, Routh BN, Johnston D, Seidemann E, Priebe NJ. Voltage-Gated Intrinsic Conductances Shape the Input-Output Relationship of Cortical Neurons in Behaving Primate V1. Neuron. 2020; p. 1–12. doi:10.1016/j.neuron.2020.04.001.

45. Baker C, Zhu V, Rosenbaum R. Nonlinear Stimulus Representations in Neural Circuits with Approximate Excitatory-Inhibitory Balance. PLOS Computational Biology. 2020;16(9):e1008192. doi:10.1371/journal.pcbi.1008192.

46. Veit J, Handy G, Mossing DP, Doiron B, Adesnik H. Cortical VIP Neurons Locally Control the Gain but Globally Control the Coherence of Gamma Band Rhythms. bioRxiv. 2021; p. 1–29.

47. Ozeki H, Finn IM, Schaffer ES, Miller KD, Ferster D. Inhibitory Stabilization of the Cortical Network Underlies Visual Surround Suppression. Neuron. 2009;62(4):578–592. doi:10.1016/j.neuron.2009.03.028.

48. Sanzeni A, Akitake B, Goldbach HC, Leedy CE, Brunel N, Histed MH. Inhibition Stabilization Is a Widespread Property of Cortical Networks. eLife. 2020;9:1–39. doi:10.7554/eLife.54875.

49. Oberheim NA, Tian GF, Han X, Peng W, Takano T, Ransom B, et al. Loss of Astrocytic Domain Organization in the Epileptic Brain. The Journal of Neuroscience. 2008;28(13):3264–3276. doi:10.1523/JNEUROSCI.4980-07.2008.

50. Maren S, Tocco G, Standley S, Baudry M, Thompson RF. Postsynaptic Factors in the Expression of Long-Term Potentiation (LTP): Increased Glutamate Receptor Binding Following LTP Induction in Vivo. Proceedings of the National Academy of Sciences. 1993;90(20):9654–9658. doi:10.1073/pnas.90.20.9654.

51. Henneberger C, Bard L, Panatier A, Reynolds JP, Kopach O, Medvedev NI, et al. LTP Induction Boosts Glutamate Spillover by Driving Withdrawal of Perisynaptic Astroglia. Neuron. 2020;108(5):919–936.e11. doi:10.1016/j.neuron.2020.08.030.

52. Clarke LE, Barres BA. Emerging Roles of Astrocytes in Neural Circuit Development. Nature Reviews Neuroscience. 2013;14(5):311–321. doi:10.1038/nrn3484.

53. Handy G, Taheri M, White JA, Borisyuk A. Mathematical Investigation of IP3-dependent Calcium Dynamics in Astrocytes. Journal of Computational Neuroscience. 2017;42(3):257–273. doi:10.1007/s10827-017-0640-1.

54. Taheri M, Handy G, Borisyuk A, White JA. Diversity of Evoked Astrocyte Ca2+ Dynamics Quantified through Experimental Measurements and Mathematical Modeling. Frontiers in Systems Neuroscience. 2017;11(23):1–19. doi:10.3389/fnsys.2017.00079.

55. Larsen BR, Assentoft M, Cotrina ML, Hua SZ, Nedergaard M, Kaila K, et al. Contributions of the Na+/K+-ATPase, NKCC1, and Kir4.1 to Hippocampal K+ Clearance and Volume Responses. GLIA. 2014;62(4):608–622. doi:10.1002/glia.22629.

56. Sloan SA, Barres BA. Looks Can Be Deceiving: Reconsidering the Evidence for Gliotransmission. Neuron. 2014;84(6):1112–1115. doi:10.1016/j.neuron.2014.12.003.

57. Fiacco TA, McCarthy KD. Multiple Lines of Evidence Indicate That Gliotransmission Does Not Occur under Physiological Conditions. The Journal of Neuroscience. 2018;38(1):3–13. doi:10.1523/JNEUROSCI.0016-17.2017.

58. Savtchouk I, Volterra A. Gliotransmission: Beyond Black-and-White. The Journal of Neuroscience. 2018;38(1):14–25. doi:10.1523/JNEUROSCI.0017-17.2017.

59. Nwaobi SE, Cuddapah VA, Patterson KC, Randolph AC, Olsen ML. The Role of Glial-Specific Kir4.1 in Normal and Pathological States of the CNS. Acta Neuropathologica. 2016;132(1):1–21. doi:10.1007/s00401-016-1553-1.

60. Barbour B, Brew H, Attwell D. Electrogenic Glutamate Uptake in Glial Cells Is Activated by Intracellular Potassium. Nature. 1988;335(6189):433–435. doi:10.1038/335433a0.

61. Kucheryavykh YV, Kucheryavykh LY, Nichols CG, Maldonado HM, Baksi K, Reichenbach A, et al. Downregulation of Kir4.1 Inward Rectifying Potassium Channel Subunits by RNAi Impairs Potassium Transfer and Glutamate Uptake by Cultured Cortical Astrocytes. Glia. 2007;55(3):274–281. doi:10.1002/glia.20455.

62. Piet R, Vargová L, Syková E, Poulain DA, Oliet SHR. Physiological Contribution of the Astrocytic Environment of Neurons to Intersynaptic Crosstalk. Proceedings of the National Academy of Sciences. 2004;101(7):2151–2155. doi:10.1073/pnas.0308408100.

63. Huang H, Bordey A. Glial Glutamate Transporters Limit Spillover Activation of Presynaptic NMDA Receptors and Influence Synaptic Inhibition of Purkinje Neurons. The Journal of Neuroscience. 2004;24(25):5659–5669. doi:10.1523/JNEUROSCI.1338-04.2004.

64. Murphy-Royal C, Dupuis J, Groc L, Oliet SHR. Astroglial Glutamate Transporters in the Brain: Regulating Neurotransmitter Homeostasis and Synaptic Transmission: Astroglial Glutamate Uptake. Journal of Neuroscience Research. 2017;95(11):2140–2151. doi:10.1002/jnr.24029.