CRISPR/Cas9 mediated gene editing in non-model nematode *Panagrolaimus* sp. PS1159

Viktoria Hellekes[1,2]*, Denise Claus[2], Johanna Seiler[1], Felix Illner[1], Philipp Schiffer[2] and
Michael Kroiher[1]

[1]Institute for Zoology, University of Cologne, Cologne, NRW, Germany

[2]worm~lab, Institute for Zoology, University of Cologne, Cologne, NRW, Germany

* Correspondence: v.hellekes@uni-koeln.de

Abstract

The phylum Nematoda consists of a huge diversity of species in a broad range of ecosystems and
habitats. Nematodes share a largely conserved Bauplan but major differences have been found in early
developmental processes. The development of the nematode model organism *Caenorhabditis elegans*
has been studied in great detail for decades. These efforts have provided the community with a large
number of protocols and methods. Unfortunately, many of these tools are not easily applicable in non-
Caenorhabditis nematodes. In recent years it has become clear that many crucial genes in the *C.*
elegans developmental toolkit are absent in other nematode species. It is thus necessary to study the
developmental program of other nematode species in detail to understand evolutionary conservation
and novelty in the phylum.

Panagrolaimus sp. PS1159 is a non-parasitic nematode exhibiting parthenogenetic reproduction and
we are establishing the species to comparatively study evolution, biodiversity and alternative
reproduction and survival strategies. Here, we demonstrate the first successful application of the
CRISPR/Cas9 system for genome editing in *Panagrolaimus* sp. PS1159 and its closely related
hermaphroditic species *Propanagrolaimus* sp. JU765 with both the non-homologous end joining and
the homology-directed repair mechanism. Using microinjections and modifying published protocols
from *C. elegans and P. pacificus* we induced mutations in the orthologue of *unc-22,* which resulted in
a visible uncoordinated twitching phenotype. We also compared the HDR efficiency following the
delivery of different repair templates. This work will expand the applicability for a wide range of non-
model nematodes from across the tree and facilitate functional analysis into the evolution of
parthenogenesis, changes in the developmental program of Nematoda, and cryptobiosis.

1   Introduction

Understanding the molecular machinery controlling cellular fate during development of an organism

is a major goal in developmental biology. Functional analysis of genes and how they are regulated aids

to understand how, for example, body shape is encoded in genomes (Levine and Davidson, 2005).

The phylum Nematoda comprises of an estimated 1 million species (Lambdshead and Boucher, 2003)

occupying a broad range of habitats including terrestrial, marine, and aquatic environments (Figure 1)

(Holterman et al., 2019). Many species have adopted a parasitic lifestyle, or have evolved survival

strategies to deal with extreme cold and desiccation (Blaxter and Koutsovoulos, 2015; Shannon et al.,

2005). Throughout the Nematoda, different modes of reproduction are realized. For example,

parthenogenesis, a form of asexual reproduction, occurs frequently. Despite their different biotopes

and lifestyles, all nematodes share a morphologically similarly structured and largely conserved

Bauplan (Poinar, 2011). In contrast to this conserved outcome of development, variations are found in

early development on the cellular level with divergent cleavage patterns (Dolinski et al., 2001; Schulze

and Schierenberg, 2009). This variability of embryogenesis shows that the patterns and structures

observed in a single species, such as the highly derived nematode model *Caenorhabditis elegans*, do

not allow a generalization across the phylum. A special case in this regard are parthenogenetic

nematodes, which initiate development without male input.

On the genetic level, many aspects of the molecular machinery for development have been studied and

are understood in the nematode model system. However, it has recently become clear that many

developmentally crucial *C. elegans* genes are absent in other nematode species. This includes for

example sex determination, axis formation and endo-mesoderm specific genes (Kraus et al., 2017;

Schiffer et al., 2013, 2018). It is thus necessary to study functions of genes and their regulations

throughout the diversity of the phylum. Some other nematode species have been studied to greater

detail, for example *Pristionchus pacificus,* a species from the same clade as *C. elegans* (Figure 1), but

we lack approachable and informative systems from across the tree. Parthenogenesis, for example, is

not present in any for the more than 50 *Caenorhabditis* species named to date, nor in species close to

Pristionchus.

In nematodes many species can easily be cultured in the laboratory, and the availability of more high-

quality genomes through recent advances in sequencing technology is enabling us to define precise

targets for gene knock-out studies. Unfortunately, many methods and protocols established in the

model organisms do not work out-of-the-box in non-*Caenorhabditis* nematodes and organisms with

less background on the genetic level. For example, RNAi has been restricted to a few nematodes (Félix,

Figure 1

Figure 1: A cladogram showing the position of the two main species in this study, *Panagrolaimus* sp. PS1159 and *Propanagrolaimus* sp. JU765, in relation to the nematode model organisms *C. elegans* and *P. pacificus*.

2008), as they are often insensitive to this method and important genes of the pathway are missing

(Koutsovoulos et al. in prep).

The CRISPR/Cas9 system (Wang et al., 2016) is a powerful experimental tool for gene-editing. The

simple design, its high efficiency, and low cost have made it the go-to-method in non-model organisms

in recent years (Dickinson et al., 2020). The most important and widely adapted CRISPR/Cas

application for gene editing is the type II system, which relies on one protein only (Makarova et al.,

2011). The *Streptococcus pyogenes* Cas9 requires two small RNAs to build a ribonucleoprotein

complex (RNP): the CRISPR RNA (crRNA) with a 20 bp guide sequence and its downstream

protospacer-adjacent motif (PAM; NGG) that determines the target, and the universal trans-activating

CRISPR RNA (tracrRNA). The crRNA binds to the DNA target by base-pair binding. Cas9 recognizes

the PAM sequence, leading to a cleavage of both strands 3 nucleotides away from the PAM (Doudna

and Charpentier, 2014), typically followed by either one of two cellular repair mechanisms. The Non-

homologous End Joining (NHEJ) repairs the double-strand break (DSB) by ligating the ends. This

mechanism sometimes results in the loss of nucleotides (Davis and Chen, 2013), a frame shift,

premature stop and therefore creating knock-out mutations. The homology directed repair (HDR) is a

more accurate system, in which user-provided DNA templates with homology arms serve as a guide

to precisely repair the DSB (Mali et al., 2013; Paix et al., 2017) generating knock-in mutations. Cas9

can be delivered to the target DNA in various ways, for example via expression plasmids, or as a

purified protein (Lino et al., 2018). It was shown that direct injection of the RNP complex into the

adult gonad is more efficient and less time consuming than injection of plasmids (Paix et al., 2015).

Over the past years, CRISPR/Cas9 was successfully applied in *C. elegans*. But the system has been

proven to be useful also in non-model nematodes, such as *C. briggsae* (Culp et al., 2020), *P.*

pacificus (Witte et al., 2015), the human- and rat-parasitic worms *S. stercoralis* and *S. ratti* (Gang et

al., 2017) Auanema (Adams et al., 2019) and *O. tipulae* (Vargas-Velazquez et al., 2019).

Here we report the successful and reproducible use of microinjection, and CRISPR/Cas9 in two new

nematode strains from clade IV (Figure 1), the parthenogenetic strain *Panagrolaimus* sp. PS1159 and

the hermaphroditic species *Propanagrolaimus* sp. JU765.

We also provide an optimization of previously published protocols using modified microinjection

needles, different target regions as well as temperature and concentration adjustment during the RNP

complex assembly. We also compare the HDR efficiency following the delivery of different repair

templates.

2   Results

2.1   Microinjection in *Panagrolaimus* sp. PS1159

In nematodes CRISPR/Cas9 components are delivered through injection into the adult gonad. We found the cuticle in *Panagrolaimus* to be more robust and harder to breach than in *C. elegans*. At the same time the gonad appeared to be less tolerant to the injury resulting by the needles used for *C. elegans*. Thus, for successful microinjection in *Panagrolaimus* sp. PS1159 we adapted the *C. elegans* microinjection protocol (Evans, 2006) accordingly (see supplementary video V1).

Panagrolaimus worms were also less likely to stick on the 2% agarose pads. We therefore used higher concentrations of approximately 2.5% agarose pads for good adhesion of the worms. To prepare injection needles using a puller (Sutter Instruments, P-2000), we use a 1-lined so called "bee stinger" program (Oesterle, 2018) as a basis for a 3-lined program (table 1), producing sharper needles with a short taper. With these modifications we were able to easily inject *C. elegans, P. pacificus*, PS1159, JU765, and even the very small nematode *D. coronatus* (data not shown) with a high survival rate.

2.2   Gene knock-out of *unc-22* in PS1159

We targeted the PS1159 orthologue of the *C. elegans unc-22* gene, which was previously successfully mutagenized by CRISPR/Cas9 in *C. elegans* (Kim et al., 2014), and the non-model nematodes *S. stercoralis,* and *S. ratti* (Gang et al., 2017). *unc-22* encodes Twitchin, a large protein, which is involved in the regulation of muscle contraction (Moerman et al., 1988). Mutation of *unc-22* in *C. elegans* results in motility defects and a twitching phenotype, which is easily detectable under the dissecting microscope.

2.2.1 Identification of the PS1159 *unc-22* ortholog

The Twitchin protein (UNC-22, isoform a) of *C. elegans* was downloaded from www.wormbase.org (WormBase ID CE33017) and blasted against the PS1159 genome using the TBLASTN tool. A potential orthologue with 62.9% sequence similarity was selected (PS1159_v2.g4256). Reciprocal BLASTN against the *C. elegans* genome confirmed the *C. elegans unc-22* gene as the best hit indicating the gene found is an *unc-22* orthologue. Target guide RNA sequences, preferably following the form G(N)G(NGG) were searched for in the exonic regions of the *unc-22* orthologue in our PS1159 genome assembly (Schiffer et al., 2019) using CRISPOR (Figure 2A; Concordet and Haeussler, 2018). Guide sequences with a GG motif at the 3' end have been observed to have a higher cleavage efficiency in *C. elegans* (Farboud and Meyer, 2015). Sequence alignment analysis of the chosen guide RNA revealed no significant off-target sequences in the entire genome. Two pairs of repair templates with

Figure 2

A PS1159 *unc-22* guide design

1000 bp

5'-TCGTCTTTCAACCATATATCCTCGA|ATACGGCCACCACCATCATGTTCAGGG-3'

5'-ATTCATCACCAACTTCTCGTCTTTCAACCATATATCTCTAGACGAATACGGCCACCACCATCATGTTCAGGGCGACG-3'

Homology arm   *Xba*I   Homology arm

B PS1159 *unc-22* RNP complex formation

tracrRNA

+

+ Cas9 →

crRNA

Cas9
tracrRNA
crRNA

C Screening for *unc-22* mutants

D Motility assay

WT   0 min

WT   15 min

unc-22   0 min

unc-22   15 min

E Survival of *unc-22* mutants in nicotine

Surviving nematodes

Time (min)

strong
weak
WT

Figure 2: **(A)** CRISPR target site for *Panagrolaimus* sp. PS1159 *unc-22*. Predicted gene structure. Black boxes = exons. Red arrows = CRISPR target regions. Zoom into part of exon 15 coding strand with target region (blue) and complementary PAM site (red). Vertical bar indicates the expected Cas9 cleavage site. Example of single-stranded oligodesoxyribonucleotide (ssODN) homology arms flanking the cleavage site on both sides and insertion of a *Xba*I cleavage site (dark yellow). Gene figure was designed by using the Exon-Intron Graphic Maker (version 4) **(B)** Ribonucleoprotein complex assembly. Cas9 protein, crRNA and trRNA are mixed and incubated **(C+E)** Strategy for screening in PS1159. RNP complexes are introduced into the syncytial part of the gonad of an adult worm by microinjection. F1 progeny is screened in 0.5% nicotine for twitching phenotypes. Comparative assay of survival in nicotine of wild type, strong and weak twitching worms over a time period of 10 minutes **(D)** Reduced motility of wild-type worms vs. *unc-22* F1 in the time course of 15 min; black line shows movement; scale bar = 1 mm.

homology arms of 35 and 98 nt flanking the predicted Cas9 cleavage site with a built-in mutation and

restriction enzyme *Xba*I site were produced (Figure 2A).

2.2.2 CRISPR-Cas9 targeting of PS1159 *unc-22* causes an uncoordinated phenotype

CRISPR/Cas9 components were assembled and introduced into the syncytial part of the gonad of

PS1159 following the protocol of Paix et al, 2017. After injections, all worms were recovered on a

single agar plate. After 12 days (generation time is 8-10 days in *Panagrolaimus*; Schiffer et al., 2019),

16 F1 animals showing reduced crawling activity and a twitching phenotype in water were transferred

into single drops of *Plectus* nematode growth medium (PNGG; supplementary Table S1). We were

able to isolate 4 different twitchers with varying intensity of their twitching behavior, most

likely indicating strong and weak phenotypes. The motility of strong twitchers was monitored and

compared to wild-type worms. The movement of twitching worms appeared uncoordinated. In a

comparative assay, wild type nematodes appeared to have moved a longer distance after 15 minutes,

indicating reduced motility in mutants (Figure 2D).

In *C. elegans*, wild-type worms stop moving and eventually die under exposure of nicotinic

acetylcholine receptor agonists, such as levamisole or nicotine. *C. elegans unc-22* mutants are resistant

to these poisons and twitching is enhanced by exposure (Lewis et al., 1980; Moerman et al., 1988). We

tested exposure of PS1159 wild-type and CRISPR-induced *unc-22* mutants to both, levamisole (1mM)

and 0.5% nicotine. Wild-type animals and both twitching phenotypes were analyzed (Figure 2C,E;

supplementary video V1). After 10 minutes all worms were rescued in M9. No wild-type animal, 9

strong and 5 weak twitchers survived the procedure. All future progeny of injected worms were

screened in water or, if confirmation was needed, additionally in 0.5% nicotine for 2-3 minutes.

2.3   Development of an optimal cultivation method after injections for PS1159

The transfer of single worms per plate, as has been done in *C. elegans* and *P. pacificus*, was not optimal

for *Panagrolaimus* sp. PS1159, as the worms frequently crawled out of the agar to the walls of the

plates and dried out. To find the optimal way of cultivating the worms after microinjections, different

approaches were taken. Injected worms were transferred either (1) into single PNGG drops on 12-well

plates in a moist chamber individually, (2) all individuals onto a single agar plate or (3) onto single

plates with each worm covered by a slice of agar. Although surviving the transfer into single PNGG

drops, worms could not be screened properly because of strong bacterial growth in the drops. When

worms were placed on a plate altogether, they remained on the plate but analysis of F1s from individual

P0s was impossible. In the latter approach, almost all worms remained under the agar pad and laid

eggs. Occasionally some worms still left and dried out, but the advantage of this method outweigh the

loss of single individuals. However, since the reproduction cycle of PS1159 is much longer, than in *C.*

elegans and *P. pacificus*, mothers were kept at 25°C for at least 5-7 days on the plates, allowing them

to lay more eggs.

2.4   Modification of the CRISPR mix preparation increases efficiency

During the first experiments, we were only able to isolate a few worms with a twitching phenotype

and injection results were thus unsatisfactory and not consistent. After the development of the optimal

cultivation strategy and the possibility to screen progeny of individual injected P0s, we aimed to

adapt the protocol of the CRISPR injection mix to yield a higher efficiency.

2.4.1 Using guide RNAs targeting different regions in the gene shows only minor variations

To examine if the genomic region of the double strand break affects the editing success, we used 4

targets in different exons (Figure 2A; Table 4). The total outcome of progeny with the desired

phenotype after injection of 60 µM crRNA varied between 1.14% and 2.63% (Table 2). When using a

modified crRNA from IDT (Alt-R CRISPR-Cas9 crRNA XT), which contains additional chemical

modifications providing an increased stability the efficiency rose dramatically to up to 16%.

2.4.2 Changes of temperature optimizes complex formation

In *Pristionchus pacificus* the CRISPR mix protocol differs from the one used in *C. elegans* (Paix et al.,

2017) by an additional pre-annealing step of the tracrRNA and crRNA (James Lightfoot, pers.

comment). We tested if the incubation temperature has an impact on the complex formation in PS1159.

We therefore injected the complex with the modified crRNA with 60 µM concentration. When pre-

annealing the crRNA and tracrRNA at 95° C for 5 minutes (Hiraga et al., 2021) at the beginning we

yielded 8.25% efficiency. When assembling the mix without the annealing step at 95° C, but with an

incubation step at 37° C for 10 minutes after adding the Cas9 we reached to 11.57% of twitching

progeny. When using both steps, the percentage of F1s with the phenotype increased up to 16.34 (Table

2).

2.4.3 Guide RNA concentration determines effect-size of gene editing

We then asked how the concentration of the crRNA affected the mutation frequency. We used

concentrations of 12,5 µM, 15 µM, 30 µM, 60 µM and 125 µM and discovered that by increasing up

to 60 µM improved the efficiency rate but did not change much when going up to 125 µM (Table 2).

After these findings we decided that our future experiments would be performed with that optimal

concentration of 60 µM.

2.4.4 The optimal time-window for screening is during the first 48 hours after injection

After establishing the enhanced protocol for the injection mix, we further investigated the optimal time

window for observing CRISPR induced knock-outs in PS1159. After microinjections, P0s were

isolated onto single agar plates with a slice of agar and transferred every 24 hours for a total of 6 days

(Figure 3A). We found that in the first 24 hours the average percentage of mutated progeny was 25%

and then dropped to 8.5% after 48 hours (Figure 3B). After 72 hours, there was still an average of 7.5%

and then 0% to 1.2% in the next 3 days. From these results we decided that the preferred time window

to screen for mutations after injections, is 48 hours post injection after considering the feasibility and

simplicity of maintenance of the progeny and achieving mutations arising from a higher number of

P0s.

2.5 Sequencing verifies *unc-22* mutation after double strand break within the target site

We aimed to molecularly characterize the mutations produced after knock-out and therefore amplified

the target region using primers located 328 bp upstream and 454 bp upstream of the PAM and

sequenced it after single worm lysis and PCR. Sanger sequencing of the PCR product confirmed

different types of mutations near the cleavage site (Figure 4A). To quantify and identify the genotypes

from the trace data, we used the ICE tool from Synthego (Conant et al., 2022; Figure 4B). To determine

the exact mutations, we cloned and subsequent sequenced each allele. The results were in accordance

with the prediction from the Synthego Analysis tool (Figure 4C).

In some cases, the amplification failed and only the WT sequences were observed even though a clear

phenotype was visible, suggesting the deletion of at least one primer site. We attempted to localize the

exact site of the deletion by increasing the PCR amplification region to up to 6000 bp and used different

primer combinations. We found an amplification at 6000 bp (WT) and 3000 bp (supplementary Figure

S1 A) of one mutant (T4) and at 3000 bp when using only reverse primers. We next sequenced this

region and found a deletion around the target and forward primer site and an inversion of the reserve

primer region (supplementary Figure S1 B).

2.6 Longer homology arms increase efficiency

To simplify the detection of knock-in success we used homology arms with a built-in insert containing

a recognition site for the restriction enzyme *Xba*I, which additionally delivers an in frame stop, thus

allowing us to easily check for gene editing in progeny with a phenotype via restriction digest (Figure

5A) after PCR. Restriction digest was successful and Sanger sequencing after DNA cloning confirmed

the desired insertion in one allele (Figure 5B). We designed longer homology arms (98nt) with an

Figure 3

Figure 3:   **(A)** Scheme of experimental workflow. Microinjection of adult P0 into gonad. Every 24 h individual worms are transferred onto fresh agar plates covered by a slice of agar **(B)** Number of wild-type (light grey) and twitching (dark grey) progeny per injected P0 in time intervals of 24 h for a total of 6 days

Figure 4

A PS1159 *unc-22* Sanger traces

B Distribution of Indels

C Sequences of each allele

Figure 4: **(A)** *Panagrolaimus* sp. PS1159 *unc-22* Sanger Sequencing traces after PCR of edited and wild-type (control) in the region around the guide sequence using the Synthego Performance Analysis, ICE Analysis. 2019. v3.0. Synthego. Black underlined region shows the guide sequence. Red underline represents the PAM site. Vertical dotted line shows cut site **(B)** Predicted distribution and percentage of Indels in the entire population of genomes using ICE **(C)** Sanger sequencing reveals deletions in single alleles after DNA cloning. Target region (blue) and complementary PAM site (red). Vertical bar indicates the expected (actual) Cas9 cleavage site.

Figure 5

A PS1159 *unc-22* restriction digest

B Sanger sequences of each allele

wild type   5'-TCGTCTTTCAACCATATATCCTCGA|ATACGGCCACCACCATCATGTTCAGGG-3'

unc-22 mutant

5'-TCGTCTTTCAACCATATATCTCTAGA|CGAATACGGCCACCACCATCATGTTCAGGG-3'   **+4 bp**

5'-TCGTCTTTCAACCATATATCCTCGA | ATACGGCCACCACCATCATGTTCAGGG-3'   **+0 bp**

5'-TCGTCTTTCAACCATATATCC----|-------CACCACCATCATGTTCAGGG-3'   **- 11 bp**

Figure 5: **(A)** *Panagrolaimus* sp. PS1159 *unc-22:* Representative examples of agarose gel electrophoresis after restriction digest with *Xba*I of knock-in experiments. Multiple band patterns of mutants with desired insertion. M=Marker Lambda PstI digest. C=positive control of approved knock-in positive F1. T1 - T5= twitchers 1-5 **(B)** Sanger sequencing revealing insertion in one allele after DNA cloning of mutant after positive restriction digest. Target region (blue) and complementary PAM site (red). Vertical bar indicates the expected (actual) Cas9 cleavage site. *Xba*I cleavage site in dark yellow.

additional incorporated phosphorothioate (PS) bond modification and observed the effect of knock-in

efficiency when compared to the results of injections with 35 nt arms flanking the target site. We found

that in ~10% of twitching progeny the desired knock-in has been generated with short homology arms

whereas using modified 98 nt resulted in a significantly higher outcome of knock-ins with 58% (Fishers

exact test p value = 0.00233; n=30).

2.7   The same protocol can be applied in *Propanagrolaimus* sp. JU765

Following our improved protocol for PS1159, we asked whether we could transfer the injection mixture

assembly and microinjection strategy to modify the genome of other non-model nematodes. We hence

aimed to induce mutations in the *Propanagrolaimus* sp. JU765 orthologue of *unc-22*. After successful

injections of 13 individuals, we screened the progeny after 7 days and recovered 30.7% with a twitching

phenotype (total F1=501). To verify that the phenotype arose from a CRISPR induced knock-out, we

amplified the target region with primers 437 bp upstream and 416 bp downstream of the PAM and

performed a T7EI assay indicating a mutation within the region (Figure 6A). We identified the exact

indels with Sanger Sequencing (Figure 6B). This result supports that we provide an optimal protocol

that might be useful in a variety of non-model species in the phylum Nematoda.

3   Discussion

CRISPR/Cas9 has become an indispensable tool for precise gene editing in different eukaryotes, and

has revolutionized genetics in the past years in many model organisms (Chang et al., 2013; Cho et al.,

2013; Cong et al., 2013; Miki et al., 2021). In this study, we demonstrate the successful and efficient

application of the CRISPR/Cas9 system in the nematodes *Panagrolaimus* sp. PS1159 and

Propanagrolaimus sp. JU765 by generating heritable *unc-22* mutants. We further optimized the

protocol for a higher cleavage efficiency.

Microinjection is a useful tool to introduce genetic material into an adult worm. In *C. elegans* this

method has long been used for different approaches, including RNAi and CRISPR/Cas9 (Conte Jr. et

al., 2015; Ghanta et al., 2021; Mello and Fire, 1995) and has successfully been used in some non-model

nematodes as well (Adams et al., 2019; Cinkornpumin and Hong, 2011; Ratnappan et al., 2016).

Because of differences in the cuticle and gonad, the published protocols for *C. elegans* have not been

working in PS1159. We therefore adapted existing protocols for the microinjection pads and needles.

With these changes it was possible to inject not only PS1159, but a handful other nematodes with high

survival rates.

Figure 6

A JU765 *unc-22* T7EI assay

B Sanger Sequences of mutants T1-3

wild type   5'- ATATCAAATGGGATCCTCCTCATGAICGACGGTGGAAGTCCAATTGATC-3'

unc-22
mutants

5'- ATATCAAATGGGATCCTCCTCATGA|----------AGTCCAATTGATC-3'   **-10 bp**

5'- ATATCAAATGGGATCCTCCTCATGA|NNNNNNNNNNNNNNNNNNNNNCGACGG-3'   **+20 bp**

5'- ATATCAAATGGGATCCTCCTC----|----GGTGGAAGTCCAATTGATC-3'   **-8 bp**

Figure 6: **(A)** *Propanagrolaimus* sp. JU765 *unc-22:* Representative examples of 3 individuals T1-T3. Agarose gel electrophoresis of T7EI assay after knock-out experiments. Multiple band patterns detect indels. M=Marker Lambda *PstI* digest. C=negative control. T1 - T3= twitchers 1-3 **(B)** Sanger sequencing of T1-T3 PCR product confirms indels in the target region. Target region (blue) and complementary PAM site (red). Vertical bar indicates the expected (actual) Cas9 cleavage site.

We generated knock-out mutations in the PS1159 and JU765 orthologues of the *C. elegans unc-22*

gene which resulted in a visible uncoordinated twitching phenotype with reduced motility increased

by nicotine exposure. This phenotype is consistent with what has been observed in *C. elegans* and *S.*

stercoralis. To determine if the phenotype was caused by mutations in the desired target, we recovered

and isolated single worms with a phenotype. We amplified, cloned, and sequenced the regions spanning

the predicted sites. We were able to identify indels showing that the genotypes were consistent with

the observed phenotypes. However, in some cases the amplification failed, and we only observed wild-

type sequences which was in line with previous results from knock-out experiments in the nematodes

Strongiloides and *Pristionchus* (Gang et al., 2017; Witte et al., 2015). We took a closer look and found

a deletion of the guide sequence and an inversion of the reserve primer site, removing the forward

primer site. It has been demonstrated in different organisms that repair of cleavage induced by Cas9

can lead to large deletions and genomic rearrangements (Adikusuma et al., 2018; Kosicki et al., 2018)

and should therefore be considered as a possible outcome. To create knock-in mutations of small

insertions we provided single-stranded homology arms on both sites. We were able to achieve even

higher knock-in efficiency when using longer homology arms with a protection of the donor DNA with

phosphorothioate modification. This result is in accordance with previous findings regarding optimized

HDR donors (Liang et al., 2017; Schubert et al., 2021).

We found that best results can be obtained when using a more stable, modified version of a crRNA

with an additional pre-annealing step with the tracrRNA before incubation with the Cas9 protein. It

has previously been reported that certain chemical alterations on terminal ends lead to increased

metabolic stability of guide RNA and subsequent higher CRISPR/Cas9 efficiency (Allen et al., 2021;

Hendel et al., 2015). Increasing the concentration of the crRNA resulted in an increase from ~1-2% to

up to 16%. Moreover, we observed the editing efficiency over a time window of one week for every

24 hours and our analysis indicates that the CRISPR effect in PS1159 performs at its best in the first

48 hours post injection, leading to even up to 25% cleavage efficiency. In *Pristionchus pacificus* it has

been reported that most mutations arose in eggs laid in the first 9 hours after injections (Witte et al.,

2015).

In summary, testing and adapting experimental conditions, such as needle size, temperature and

concentration of the RNP complex components during preparation as well as using modified reagents

is necessary for successfully applying microinjections and CRISPR/Cas9 in divergent nematode

species. Our work will facilitate functional analysis into the evolution of parthenogenesis, changes in

the developmental program of Nematoda, and cryptobiosis. The *unc-22* target might be a useful and

easily identifiable co-marker for future functional analysis, in particular for large insertions of

fluorescent markers.

4   Materials and methods

4.1   Nematodes and maintenance

All experiments have been performed using the parthenogenetic *Panagrolaimus* sp. PS1159 and the

hermaphroditic *Propanagrolaimus* sp. JU765, cultured on 55 mm diameter petri dishes at 20° C with

low-salt agar (Lahl et al., 2003) seeded with *Escherichia coli* strain OP50 as a food source (Brenner,

1974). After injections worms were kept at 25° C (20° C for experiments with older protocol) either

on single drops of 15 µl plectus nematode growth medium (PNGG, supplemental Table S1; 30 µl OP50

in 1000 µl PNGG) on 12-well plates in a moist chamber, or on single agar plates covered with a slice

of agar.

4.2   Selection of guide RNA targets

290   The *C. elegans* sequences of UNC-22 were downloaded from NCBI. TBLASTN and BLASTN tools

291   were used to identify orthologues for PS1159 and JU765 in WormBase ParaSite

292   (https://parasite.wormbase.org). We searched for potential guide RNAs and analyzed the putative off-

293   targets using a web-based software (CRISPOR, http://crispor.tefor.net; Concordet and Haeussler,

294   2018). A list of guide RNA's can be found on table 3.

4.3   Preparation of HDR donor

296   For knock-in of small insertions using the homology directed repair (HDR), ssODN (single-stranded

297   oligodeoxynucleotides) repair templates with 35 nt and 98 nt homology arms flanking the cleavage site

298   with an inserted *Xba*I site were designed using Geneious Pro 5.5.6 (Kearse et al, 2012).

4.4   Preparation of injection mix

300   1.25 µl tracrRNA (100 µM, catalog# 1073191, IDT) and x µM crRNA were mixed and incubated at

301   95°C for 5 min following incubation at RT for 5 min. 0.5 µl Cas9 (10 mg/ml, catalog# 10811059, IDT)

302   was added and incubated at 37°C for 10 min for RNP (ribonucleoprotein) complex formation. 1 µl

303   ssODN repair template (25 µM) was added and the mixture was filled up with TE buffer to 10 µl,

304   centrifuged at 13.000 x g for 5 min and kept on ice for injections.

4.5   Preparation of injection needles, agar pads and microinjection

2.5 % agarose pads were produced, and microinjections performed as described in Evans, 2006.
PS1159 and JU765 are monodelphic and the syncytial region of the gonadal arm looks similar as in *C.*
elegans. Borosilicate glass capillary (GB120F-10, 0.69x1.20x100 mm, Science products GmbH) were
pulled in a needle puller (P-2000, Sutter instruments) and loaded with the final injection mixture. The
needle was opened by gently tapping the tip against the edge of a broken piece of coverslip placed on
an agar pad and covered with oil. An Inverted DIC microscope (Zeiss IM 35) equipped with standard
Nomarski objectives (non-oil immersion, 6,3x and 40x) was used. A micromanipulator (Bachofer) with
a needle holder with a fine mobility in the x, y and z axes (Piezo manipulator, PM10, Bachofer) was
attached to the microscope and a pressure regulator (Pneumatic PicoPump PV 820) with a foot petal
connection to a vacuum pump (Vacuubrand ME2C). Pressure was adjusted between 15 and 20 psi for
injections. Worms were recovered in M9, transferred to a single agar plate, and covered with a small
piece of agar at 25°C (video of microinjection steps in supplemental video V1).

4.6   Screening for knock-out mutations

4.6.1 Single worm PCR and Sanger Sequencing
Individual worms were picked and transferred into tubes with 10 µl H2O or M9 buffer. Tubes were
placed at – 80° C for at least 1 h. 10 µl 2x worm lysis buffer (supplemental Table S2) was added,
incubated at 55° C for 3 hours and heated at 95° C for 10 minutes. We designed primers to amplify the
genomic DNA fragment containing the cut site. PCR products were resolved on a 0.8 % agarose gel.
Sanger sequencing was performed through Eurofins Scientific SE. Sequences were analyzed using the
ICE CRISPR analysis tool v3 (https://www.synthego.com; Conant et al., 2022) and Geneious Pro 5.5.6
(Kearse et al., 2012). List of primers used for this study can be found on table 4.

4.6.2 T7 Endonuclease I assay
10 µl single worm PCR product was used for T7 Endonuclease I assay according to the protocol from
New England Biolabs Inc. Samples were loaded onto a 1.4% agarose gel.

4.7   Screening for Knock-ins / HDR

4.7.1 Restriction digest
10 µl single worm PCR products of F2 individuals were mixed with 1 µl Tango buffer, 1 µl *Xba*I
enzyme and 13 µl H$_2$O and incubated at 37°C for 3 h. Samples were loaded in a 1.4% agarose gel.

4.8   Data analysis and editing

Gene structure diagrams were generated with the Exon-Intron Graphic Maker (Version 4,
www.wormweb.org). Plots were generated with R package ggplot2 (Wickham, 2016). Images and
Videos were edited with InkScape 1.2, Adobe Illustrator CS6 v. 16.0.0 and Avidemux 2.8.0.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

6   Author Contributions

VH conducted all major experiments, was involved in planning the experiments, analyzed the data,
wrote the manuscript. DC conducted some experiments and contributed to figure design. JS conducted
some experiments. FI contributed to one experiment. PS conceived the study, wrote the manuscript,
and supervised VH and DC. MK conceived the study, planned the experiments, analyzed data, edited
the manuscript, supervised VH, JS and FI and co-supervised DC.

7   Funding

VH is funded by a DFG ENP grant to PS (grant number: 434028868).

8   Acknowledgments

We thank James Lightfoot for sharing RNP complex components and protocols for pilot experiments.
This manuscript is dedicated to the memory of Einhard Schierenberg.

9   References

Adams, S., Pathak, P., Shao, H., Lok, J.B., Pires-daSilva, A., 2019. Liposome-based transfection
enhances RNAi and CRISPR-mediated mutagenesis in non-model nematode systems. Sci Rep 9, 483.
https://doi.org/10.1038/s41598-018-37036-1

Adikusuma, F., Piltz, S., Corbett, M.A., Turvey, M., McColl, S.R., Helbig, K.J., Beard, M.R., Hughes,
J., Pomerantz, R.T., Thomas, P.Q., 2018. Large deletions induced by Cas9 cleavage. Nature 560, E8–
E9. https://doi.org/10.1038/s41586-018-0380-z

Allen, D., Rosenberg, M., Hendel, A., 2021. Using Synthetically Engineered Guide RNAs to Enhance
CRISPR Genome Editing Systems in Mammalian Cells. Frontiers in Genome Editing 2.

Blaxter, M., Koutsovoulos, G., 2015. The evolution of parasitism in Nematoda. Parasitology 142, S26–
S39. https://doi.org/10.1017/S0031182014000791

Brenner, S., 1974. The genetics of Caenorhabditis elegans. Genetics 77, 71–94.
https://doi.org/10.1093/genetics/77.1.71

Chang, N., Sun, C., Gao, L., Zhu, D., Xu, X., Zhu, X., Xiong, J.-W., Xi, J.J., 2013. Genome editing
with RNA-guided Cas9 nuclease in zebrafish embryos. Cell Res 23, 465–472.
https://doi.org/10.1038/cr.2013.45

Cho, S.W., Lee, Jihyun, Carroll, D., Kim, J.-S., Lee, Junho, 2013. Heritable Gene Knockout in
Caenorhabditis elegans by Direct Injection of Cas9–sgRNA Ribonucleoproteins. Genetics 195, 1177–
1180. https://doi.org/10.1534/genetics.113.155853

Cinkornpumin, J.K., Hong, R.L., 2011. RNAi mediated gene knockdown and transgenesis by
microinjection in the necromenic Nematode Pristionchus pacificus. J Vis Exp e3270.
https://doi.org/10.3791/3270

Conant, D., Hsiau, T., Rossi, N., Oki, J., Maures, T., Waite, K., Yang, J., Joshi, S., Kelso, R., Holden,
K., Enzmann, B.L., Stoner, R., 2022. Inference of CRISPR Edits from Sanger Trace Data. CRISPR J
5, 123–130. https://doi.org/10.1089/crispr.2021.0113

Concordet, J.-P., Haeussler, M., 2018. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome
editing experiments and screens. Nucleic Acids Res 46, W242–W245.
https://doi.org/10.1093/nar/gky354

Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W., Marraffini,
L.A., Zhang, F., 2013. Multiplex genome engineering using CRISPR/Cas systems. Science 339, 819–
823. https://doi.org/10.1126/science.1231143

Conte Jr., D., MacNeil, L.T., Walhout, A.J.M., Mello, C.C., 2015. RNA Interference in Caenorhabditis
elegans. Current Protocols in Molecular Biology 109, 26.3.1-26.3.30.
https://doi.org/10.1002/0471142727.mb2603s109

Culp, E., Richman, C., Sharanya, D., Jhaveri, N., Van Den Berg, W., Gupta, B.P., 2020. Genome
editing in the nematode Caenorhabditis briggsae using the CRISPR/Cas9 system. Biology Methods
and Protocols 5. https://doi.org/10.1093/BIOMETHODS/BPAA003

Davis, A.J., Chen, D.J., 2013. DNA double strand break repair via non-homologous end-joining.
Translational cancer research 2, 130–143. https://doi.org/10.3978/J.ISSN.2218-676X.2013.04.02

Dickinson, M.H., Vosshall, L.B., Dow, J.A.T., 2020. Genome editing in non-model organisms opens
new horizons for comparative physiology. Journal of Experimental Biology 223, jeb221119.
https://doi.org/10.1242/jeb.221119

Dolinski, C., Baldwin, J.G., Thomas, W.K., 2001. Comparative survey of early embryogenesis of
secernentea (nematoda), with phylogenetic implications. Canadian Journal of Zoology 79, 82–94.
https://doi.org/10.1139/z00-179

Doudna, J.A., Charpentier, E., 2014. The new frontier of genome engineering with CRISPR-Cas9.
Science 346, 1258096. https://doi.org/10.1126/science.1258096

Evans, T.C., 2006. Transformation and microinjection, WormBook: The Online Review of C. elegans
Biology [Internet]. WormBook.

Farboud, B., Meyer, B.J., 2015. Dramatic enhancement of genome editing by CRISPR/Cas9 through
improved guide RNA design. Genetics 199, 959–971. https://doi.org/10.1534/genetics.115.175166

Félix, M.-A., 2008. RNA interference in nematodes and the chance that favored Sydney Brenner. J
Biol 7, 34. https://doi.org/10.1186/jbiol97

Gang, S.S., Castelletto, M.L., Bryant, A.S., Yang, E., Mancuso, N., Lopez, J.B., Pellegrini, M., Hallem,
E.A., 2017. Targeted mutagenesis in a human-parasitic nematode. PLoS Pathogens 13.
https://doi.org/10.1371/journal.ppat.1006675

Ghanta, K.S., Ishidate, T., Mello, C.C., 2021. Microinjection for precision genome editing in
Caenorhabditis elegans. STAR Protocols 2, 100748. https://doi.org/10.1016/j.xpro.2021.100748

Hendel, A., Bak, R.O., Clark, J.T., Kennedy, A.B., Ryan, D.E., Roy, S., Steinfeld, I., Lunstad, B.D.,
Kaiser, R.J., Wilkens, A.B., Bacchetta, R., Tsalenko, A., Dellinger, D., Bruhn, L., Porteus, M.H., 2015.
Chemically modified guide RNAs enhance CRISPR-Cas genome editing in human primary cells. Nat
Biotechnol 33, 985–989. https://doi.org/10.1038/nbt.3290

Hiraga, H., Ishita, Y., Chihara, T., Okumura, M., 2021. Efficient visual screening of CRISPR/Cas9
genome editing in the nematode Pristionchus pacificus. Development, Growth & Differentiation 63,
488–500. https://doi.org/10.1111/dgd.12761

Holterman, M., Schratzberger, M., Helder, J., 2019. Nematodes as evolutionary commuters between
marine, freshwater and terrestrial habitats, Biological Journal of the Linnean Society.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A.,
Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A., 2012. Geneious Basic:
an integrated and extendable desktop software platform for the organization and analysis of sequence
data. Bioinformatics 28, 1647–1649. https://doi.org/10.1093/bioinformatics/bts199

Kim, H., Ishidate, T., Ghanta, K.S., Seth, M., Conte, D., Jr., Shirayama, M., Mello, C.C., 2014. A Co-
CRISPR Strategy for Efficient Genome Editing in Caenorhabditis elegans. Genetics 197, 1069–1080.
https://doi.org/10.1534/genetics.114.166389

Kosicki, M., Tomberg, K., Bradley, A., 2018. Repair of double-strand breaks induced by CRISPR-
Cas9 leads to large deletions and complex rearrangements. Nat Biotechnol 36, 765–771.
https://doi.org/10.1038/nbt.4192

Kraus, C., Schiffer, P.H., Kagoshima, H., Hiraki, H., Vogt, T., Kroiher, M., Kohara, Y., Schierenberg,
E., 2017. Differences in the genetic control of early egg development and reproduction between C.
elegans and its parthenogenetic relative D. coronatus. EvoDevo 8. https://doi.org/10.1186/s13227-017-
0081-y

Lahl, V., Halama, C., Schierenberg, E., 2003. Comparative and experimental embryogenesis of
Plectidae (Nematoda). Dev Genes Evol 213, 18–27. https://doi.org/10.1007/s00427-002-0289-1

Lambdshead, P.J.D., Boucher, G., 2003. Marine nematode deep-sea biodiversity-hyperdiverse or
hype? Journal of Biogeography 30, 475–485.

Levine, M., Davidson, E.H., 2005. Gene regulatory networks for development. Proceedings of the
National Academy of Sciences 102, 4936–4942. https://doi.org/10.1073/pnas.0408031102

Lewis, J.A., Wu, C.H., Berg, H., Levine, J.H., 1980. The genetics of levamisole resistance in the
nematode Caenorhabditis elegans. Genetics 95, 905–928. https://doi.org/10.1093/genetics/95.4.905

Liang, X., Potter, J., Kumar, S., Ravinder, N., Chesnut, J.D., 2017. Enhanced CRISPR/Cas9-mediated
precise genome editing by improved design and delivery of gRNA, Cas9 nuclease, and donor DNA. J
Biotechnol 241, 136–146. https://doi.org/10.1016/j.jbiotec.2016.11.011

Lino, C.A., Harper, J.C., Carney, J.P., Timlin, J.A., 2018. Delivering CRISPR: a review of the
challenges   and   approaches.   Drug   Delivery   25,   1234–1234.
https://doi.org/10.1080/10717544.2018.1474964

Makarova, K.S., Haft, D.H., Barrangou, R., J Brouns, S.J., Charpentier, E., Horvath, P., Moineau, S.,
M Mojica, F.J., Wolf, Y.I., Yakunin, A.F., van der Oost, J., Koonin, E.V., 2011. Evolution and
classification of the CRISPR–Cas systems. https://doi.org/10.1038/nrmicro2577

Mali, P., Yang, L., Esvelt, K.M., Aach, J., Guell, M., DiCarlo, J.E., Norville, J.E., Church, G.M., 2013.
RNA-Guided Human Genome Engineering via Cas9. Science (New York, N.Y.) 339, 823–823.
https://doi.org/10.1126/SCIENCE.1232033

Mello, C., Fire, A., 1995. Chapter 19 DNA Transformation, in: Epstein, H.F., Shakes, D.C. (Eds.),
Methods in Cell Biology, Cuenorhubditis Elegans: Modern Biologcal Analysis of an Organism.
Academic Press, pp. 451–482. https://doi.org/10.1016/S0091-679X(08)61399-0

Miki, D., Zinta, G., Zhang, W., Peng, F., Feng, Z., Zhu, J.-K., 2021. CRISPR/Cas9-Based Genome
Editing Toolbox for Arabidopsis thaliana, in: Sanchez-Serrano, J.J., Salinas, J. (Eds.), Arabidopsis
Protocols, Methods in Molecular Biology. Springer US, New York, NY, pp. 121–146.
https://doi.org/10.1007/978-1-0716-0880-7_5

Moerman, D.G., Benian, G.M., Barstead, R.J., Schriefer, L.A., Waterston, R.H., 1988. Identification
and intracellular localization of the unc-22 gene product of Caenorhabditis elegans. Genes &
development 2, 93–105. https://doi.org/10.1101/GAD.2.1.93

Oesterle, A. 2018. Pipette Cookbook 2018. P-97 & P-1000 Micropipette Pullers [online]. Available
at: https://www.sutter.com/PDFs/pipette_cookbook.pdf

Paix, A., Folkmann, A., Rasoloson, D., Seydoux, G., 2015. High efficiency, homology-directed
genome editing in Caenorhabditis elegans using CRISPR-Cas9ribonucleoprotein complexes. Genetics
201, 47–54. https://doi.org/10.1534/GENETICS.115.179382/-/DC1

Paix, A., Folkmann, A., Seydoux, G., 2017. Precision genome editing using CRISPR-Cas9 and linear
repair   templates   in   C.   elegans.   Methods   (San   Diego,   Calif.)   121–122,   86–93.
https://doi.org/10.1016/J.YMETH.2017.03.023

Poinar, G.O., 2011. The evolutionary history of nematodes : as revealed in stone, amber and mummies.
Nematology Monographs and Perspectives.

Ratnappan, R., Vadnal, J., Keaney, M., Eleftherianos, I., O'Halloran, D., Hawdon, J.M., 2016. RNAi-
mediated gene knockdown by microinjection in the model entomopathogenic nematode
Heterorhabditis bacteriophora. Parasit Vectors 9, 160. https://doi.org/10.1186/s13071-016-1442-4

Schiffer, P.H., Danchin, E.G.J., Burnell, A.M., Creevey, C.J., Wong, S., Dix, I., O'Mahony, G.,
Culleton, B.A., Rancurel, C., Stier, G., Martínez-Salazar, E.A., Marconi, A., Trivedi, U., Kroiher, M.,
Thorne, M.A.S., Schierenberg, E., Wiehe, T., Blaxter, M., 2019. Signatures of the Evolution of
Parthenogenesis and Cryptobiosis in the Genomes of Panagrolaimid Nematodes. iScience 21, 587–
602. https://doi.org/10.1016/j.isci.2019.10.039

Schiffer, P.H., Kroiher, M., Kraus, C., Koutsovoulos, G.D., Kumar, S., Camps, J.I.R., Nsah, N.A.,
Stappert, D., Morris, K., Heger, P., Altmüller, J., Frommolt, P., Nürnberg, P., Thomas, K., Blaxter,
M.L., Schierenberg, E., 2013. The genome of Romanomermis culicivorax: revealing fundamental
changes in the core developmental genetic toolkit in Nematoda, BMC Genomics.

Schiffer, P.H., Polsky, A.L., Cole, A.G., Camps, J.I.R., Kroiher, M., Silver, D.H., Grishkevich, V.,
Anavy, L., Koutsovoulos, G., Hashimshony, T., Yanai, I., 2018. The gene regulatory program of
Acrobeloides nanus reveals conservation of phylum-specific expression. Proceedings of the National
Academy of Sciences of the United States of America 115, 4459–4464.
https://doi.org/10.1073/pnas.1720817115

Schubert, M.S., Thommandru, B., Woodley, J., Turk, R., Yan, S., Kurgan, G., McNeill, M.S., Rettig,
G.R., 2021. Optimized design parameters for CRISPR Cas9 and Cas12a homology-directed repair. Sci
Rep 11, 19482. https://doi.org/10.1038/s41598-021-98965-y

Schulze, J., Schierenberg, E., 2009. Embryogenesis of Romanomermis culicivorax: An alternative way
to construct a nematode. Developmental Biology 334, 10–21.
https://doi.org/10.1016/j.ydbio.2009.06.009

Shannon, A.J., Browne, J.A., Boyd, J., Fitzpatrick, D.A., Burnell, A.M., 2005. The anhydrobiotic
potential and molecular phylogenetics of species and strains of Panagrolaimus (Nematoda,
Panagrolaimidae). Journal of Experimental Biology 208, 2433–2445.
https://doi.org/10.1242/jeb.01629

Vargas-Velazquez, A.M., Besnard, F., Félix, M.-A., 2019. Necessity and Contingency in
Developmental Genetic Screens: EGF, Wnt, and Semaphorin Pathways in Vulval Induction of the
Nematode Oscheius tipulae. Genetics 211, 1315–1330. https://doi.org/10.1534/genetics.119.301970

Wang, H., La Russa, M., Qi, L.S., 2016. CRISPR/Cas9 in Genome Editing and Beyond. Annu Rev
Biochem 85, 227–264. https://doi.org/10.1146/annurev-biochem-060815-014607

Wickham H, 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. ISBN
978-3-319-24277-4, https://ggplot2.tidyverse.org.

Witte, H., Moreno, E., Rödelsperger, C., Kim, J., Kim, J.S., Streit, A., Sommer, R.J., 2015. Gene
inactivation using the CRISPR/Cas9 system in the nematode Pristionchus pacificus. Development
genes and evolution 225, 55–62. https://doi.org/10.1007/S00427-014-0486-8

10 FIGURE CAPTIONS

Figure 1: A cladogram showing the position of the two main species in this study, *Panagrolaimus* sp.
PS1159 and *Propanagrolaimus* sp. JU765, in relation to the nematode model organisms *C. elegans*
and *P. pacificus*.

Figure 2: (A) CRISPR target site for *Panagrolaimus* sp. PS1159 *unc-22*. Predicted gene structure.
Black boxes = exons. Red arrows = CRISPR target regions. Zoom into part of exon 15 coding strand
with target region (blue) and complementary PAM site (red). Vertical bar indicates the expected Cas9
cleavage site.   Example of single-stranded oligodesoxyribonucleotide (ssODN) homology arms
flanking the cleavage site on both sides and insertion of a *Xba*I cleavage site (dark yellow). Gene figure
was designed by using the Exon-Intron Graphic Maker (version 4) **(B)** Ribonucleoprotein complex
assembly. Cas9 protein, crRNA and tracrRNA are mixed and incubated **(C-E)** Strategy for screening
in PS1159. RNP complexes are introduced into the syncytial part of the gonad of an adult worm by
microinjection. Reduced motility of wild-type worms vs. *unc-22* F1 in the time course of 15 min; black
line shows movement; scale bar = 1 mm F1 progeny is screened in 0.5% nicotine for twitching
phenotypes.

Figure 3: (A) Scheme of experimental workflow. Microinjection of adult P0 into gonad. Every 24 h
individual worms are transferred onto fresh agar plates covered by a slice of agar **(B)** Number of wild-
type (light grey) and twitching (dark grey) progeny per injected P0 in time intervals of 24 h for 6 days

Figure 4: (A) *Panagrolaimus* sp. PS1159 *unc-22* Sanger Sequencing traces after PCR of edited and
wild-type (control) in the region around the guide sequence using the Synthego Performance Analysis,
ICE Analysis. 2019. v3.0. Synthego. Black underlined region shows the guide sequence. Red underline
represents the PAM site. Vertical dotted line shows cut site **(B)** Predicted distribution and percentage
of Indels in the entire population of genomes using ICE **(C)** Sanger sequencing reveals deletions in
single alleles after DNA cloning. Target region (blue) and complementary PAM site (red). Vertical bar
indicates the expected (actual) Cas9 cleavage site.

Figure 5: (A) *Panagrolaimus* sp. PS1159 *unc-22*: Representative examples of agarose gel
electrophoresis after restriction digest with *Xba*I of knock-in experiments. Multiple band patterns of
mutants with desired insertion. M=Marker Lambda *Pst*I digest. C=positive control of approved knock-
in positive F1. T1 - T5= twitchers 1-5 **(B)** Sanger sequencing revealing insertion in one allele after
DNA cloning of mutant after positive restriction digest. Target region (blue) and complementary PAM
site (red). Vertical bar indicates the expected (actual) Cas9 cleavage site. *Xba*I cleavage site in dark
yellow.

Figure 6: (A) *Propanagrolaimus* sp. JU765 *unc-22*: Representative examples of 3 individuals T1-T3.
Agarose gel electrophoresis of T7EI assay after knock-out experiments. Multiple band patterns detect
indels. M=Marker Lambda *Pst*I digest. C=negative control. T1 - T3= twitchers 1-3 **(B)** Sanger
sequencing of T1- T3 PCR product confirms indels in the target region. Target region (blue) and
complementary PAM site (red). Vertical bar indicates the expected (actual) Cas9 cleavage site.

11   Tables

Table 1: Needle puller P-2000 (A) One-line program "bee stinger needles" B) newly written 3-line
program for our microinjection needles

Program	Line	Heat	Filament	Velocity	Delay	Pull
A	1	350	4	40	200	0
	1	350	4	40	200	0
B	2	350	4	40	200	0
	3	450	4	60	130	100/150

Table 2: Summary of CRISPR/Cas9 experiments targeting the PS1159 *unc-22* gene

Guide	Concentration crRNA/µM tracrRNA (12,5 µM)	Temperature (*95° C and 37° C)	Surviving injected worms	P0 producing mutations	F1 animals scored	F1 animals with phenotype	%	Standard deviation
1			22	4	585	7	1.14	0.44
2	60	*	28	8	719	14	2.05	1.15
3			28	10	1493	41	2.63	0.44
	12.5	*	16	3	812	4	0.49	
4	60		9	3	407	5	1.2	
	125	95	30	5	1295	36	2.67	0.23
		95	17	3	275	5	1.82	
	12.5	37	17	11	860	59	6.86	
4 XT		*	21	14	862	83	9.63	
	60	95	13	8	419	35	8.25	
		37	12	7	337	39	11.57	
	15		18	12	1113	72	6.47	
	30	*	18	11	809	68	8.41	
4 XT	60		53	33	1663	255	16.34	5.13
	125	95	14	3	114	14	12.28	
		*	26	9	516	69	12.93	2.23

Table 3: List of gRNAs used for this study

Nematode	Exon	Sequence (5'-3')	Predicted efficiency (Doench)	Potential off-targets
JU765	16	GATCCTCCTCATGACGACGG	69	0
	7	GCACCAACCTAGTGCCTCGG	72	0
	12	GGTTAAAGACTGCCGCCACG	56	0
PS1159	15	GACTCACTAGTTTAATCAAG	67	0
	15	GATGGTGGTGGCCGTATTCG	50	9

Table 4: List of primers used for this study

Nematode	Name	Sequence (5'-3')
JU765	*unc-22* F1P	TCTGCAAAAAGATCTGACAGTGG
	unc-22 R1P	GATTGCATCAATTCTGCACTGG
	unc-22 F2	TGTGCCAAAATTGCATGAAGG
	unc-22 F1	GCGGCAATATCATGGAATCC
PS1159	*unc-22* R1	AGTTCTTTCAATCCTCGGAACC
	unc-22 SR2	GCTGTGTTCTGAATTTCCTGC