1   **NBS1 binds directly to TOPBP1 via disparate interactions between the NBS1**
2   **BRCT1 domain and the TOPBP1 BRCT1 and BRCT2 domains**
3

Oanh Huynh[1], Kenna Ruis[1], Katrina Montales[1], and W. Matthew Michael[1,2]

[1]Molecular and Computational Biology Section

Department of Biological Sciences

University of Southern California

Los Angeles, CA 90089

[2]Author for correspondence at: mattm@usc.edu

Abstract

The TOPBP1 and NBS1 proteins are key components of DNA repair and DNA-based

signaling systems. TOPBP1 is a multi-BRCT domain containing protein that plays

important roles in checkpoint signaling, DNA replication, and DNA repair. Likewise,

NBS1, which is a component of the MRE11-RAD50-NBS1 (MRN) complex, functions

in both checkpoint signaling and DNA repair. NBS1 also contains BRCT domains, and

previous works have shown that TOPBP1 and NBS1 interact with one another. This

interaction occurs efficiently under basal conditions and efficiency increases during

a DNA damage response (DDR). In this work we examine the basal interaction

between TOPBP1 and NBS1, and we show that the two proteins bind to one another

in a non-canonical manner. We report that NBS1 uses its BRCT1 domain to directly

interact with a central region of TOPBP1's BRCT1, and that it also binds to a C-

terminal region within TOPBP1's BRCT2 domain. Thus, NBS1 can interact with

TOPBP1 via two different modes of binding. For both of TOPBP1's BRCT1 and

BRCT2 domains we delineate short peptide sequences, less than 30 amino acids,

that are sufficient for binding to NBS1 and can confer binding upon transfer to a

heterologous protein, such as maltose binding protein (MBP). We also show that

NBS1 requires an intact phosphate-binding pocket (PBP) within its BRCT1 domain

to interact with TOPBP1, however the presence of phosphate is not required for

binding. These data expand our understanding of how BRCT-containing proteins

bind to one another by revealing a non-canonical mode of binding and by showing

that PBPs can contribute to binding in a phosphate-independent manner.

Key words

BRCT domain, TOPBP1, NBS1, MRN complex, protein-protein interaction, scaffold

protein

1. Introduction

Scaffold proteins are platforms for the assembly of multi-protein complexes and are key contributors to the cellular response to DNA damage. Scaffolds typically contain multiple domains that allow for either protein::protein or protein::nucleic acid interactions, and they function by recruiting multiple partners and bringing these disparate factors together in a manner that nucleates DNA-based signaling and repair complexes (Reinhardt and Yaffee, 2013; Wan et al, 2016; Day et al, 2021). One well-studied scaffold is the XRCC1 protein, an excision repair factor that binds to damaged chromatin via interaction with auto-modified PARP1, and in turn recruits DNA polymerase beta and DNA ligase III to facilitate repair of the lesion. The modular structure of XRCC1 explains its function as XRCC1 uses three distinct domains to bind damaged chromatin, bind DNA polymerase beta, and to bind DNA ligase III (London, 2020). Another well-studied scaffold protein is TOPBP1, which plays numerous roles in the cell including checkpoint signaling, DNA repair, DNA replication, and transcription (Day et al, 2021; Bagge et al, 2021; Ma et al, 2020; Wardlaw et al, 2013). TOPBP1's scaffold function has been most clearly demonstrated in yeast during the initiation of DNA replication, where the TOPBP1 ortholog DPB11 binds both SLD3 and SLD2 to allow activation of the replicative helicase (Costa and Diffley, 2022). TOPBP1 may also play a scaffolding role during DNA damage signaling. Using a cytological assay, recent work has suggested that TOPBP1 physically links the DNA signaling proteins 53BP1 and RAD9 together at sites of damage in G1 cells (Bigot et al, 2019). Other work has also promoted a scaffold function for TOPBP1, during RAD51 filament formation for DNA double-

strand break (DSB) repair. In this scenario, TOPBP1 binds both PLK kinase and

RAD51, thereby promoting RAD51 phosphorylation by PLK, a step required for

filament formation (Moudry et al, 2016).

For scaffolds to function they must contain sequence motifs that allow for

interaction with exogenous binding partners. One such motif that is prominent

amongst proteins involved in the DDR is the BRCA1 C-Terminal repeat, or BRCT

domain. BRCT domains are multi-faceted protein-protein interaction domains that

also have the capacity to bind DNA (Sheng et al, 2011), as well as the poly ADP

ribose (PAR) chains synthesized by PARP enzymes at sites of damage (Hou et al,

2019). One well-studied feature of BRCT domains is their ability to bind substrate

proteins in a manner regulated by substrate phosphorylation (Leung and Glover,

2011; Reinhardt and Yaffee, 2013; Day et al, 2021). A subset of BRCT domains

contain so-called phosphate-binding pockets (PBPs), whereby a triad of highly

conserved residues within the BRCT domain come together to form a phosphate-

binding surface that promotes interaction with phosphorylated substrates (Leung

and Glover, 2011; Reinhardt and Yaffee, 2013; Day et al, 2021). Another feature that

can distinguish different classes of BRCT domains is whether they appear as a single

module, or two or even three closely spaced modules (Sheng et al, 2011).

Interestingly, BRCT domains that contain PBPs are always found in the multi-

module configuration, and this is because BRCT::BRCT packing is considered

important for formation of the PBP (Leung and Glover, 2011; Reinhardt and Yaffee,

2013; Day et al, 2021).

One previously identified feature of BRCT domains is that they can bind to

one another. For example, the XRCC1 BRCT2 domain can form homodimers, as well

as heterodimers with the BRCT domain from DNA ligase III (London, 2020). This is

also true of TOPBP1, which contains nine BRCT domains. We have recently shown

that TOPBP1's BRCT1&2 domains can dimerize, and they can also bind to BRCT5

(Kim et al, 2020). Though not as well-studied as interactions involving BRCT

domains and phosphorylated substrates, homotypic and heterotypic BRCT::BRCT

interactions are likely to be just as important for an efficient DDR. In this work we

extend our understanding of BRCT::BRCT interactions by studying how TOPBP1

interacts with the NBS1 protein. We chose this interaction to study because recent

work from our group showed that NBS1, a component of the critical DNA repair

factor MRN, plays an important role in recruiting TOPBP1 to sites of damage

(Montales et al, 2022). Here, we report that the NBS1::TOPBP1 interaction is driven

by BRCT::BRCT interactions, whereby the BRCT1 domain from NBS1 binds directly

to the both the BRCT1 and BRCT2 domains of TOPBP1. Interestingly, NBS1 binds

BRCT1 differently than it does BRCT2, with the interaction with BRCT1 using a

similar mode of binding as XRCC1::DNA ligase III, and the interaction with BRCT2

occurring via what is likely to be a novel mode of BRCT::BRCT interaction. These

finding help fill a large knowledge gap in our understanding of how BRCT domain

containing proteins can interact with one another.

2. Material and Methods

2.1 Materials

2.1.1 Plasmids

The expression vector to produce GST-RAD9 Tail has previously been described

(Duursma et al., 2013). All other plasmids used in this study are described below, in

table format, and all are *Xenopus* proteins. Specific details about plasmid

construction are available upon request.

Plasmid	Name	Vector	AA coordinates
5Xmyc-BRCT0-2	pHG128	pCS2+MT	1-333
5Xmyc-BRCT0-2 KK155,249EE	pAK42	pCS2+MT	1-333
5Xmyc-BRCT4&5	pJA19	pCS2+MT	480-758
5Xmyc-BRCT0	pHG126	pCS2+MT	1-99
5Xmyc-BRCT1	pHG129	pCS2+MT	100-191
5Xmyc-BRCT2	pJA23	pCS2+MT	191-333
5Xmyc-NBS1	pKM107	pCS2+MT	1-763
5Xmyc-NBS1 1-400	pKM105	pCS2+MT	1-400
5Xmyc-NBS1 1-200	pKR34	pCS2+MT	1-200
5Xmyc-NBS1 100-400	pKR35	pCS2+MT	100-400
5Xmyc-NBS1 1-100	pOH4	pCS2+MT	1-100
5Xmyc-NBS1 100-200	pKR36	pCS2+MT	100-200
5Xmyc-NBS1 200-400	pKR37	pCS2+MT	200-400
5Xmyc-NBS1 393-763	pKM106	pCS2+MT	393-763
MBP-TOPBP1 1-333	pAK53	pUC57-MBP	1-333

MBP-TOPBP1 91-275	pAK56	pUC57-MBP	91-275
MBP-TOPBP1 91-196	pKR23	pUC57-MBP	91-196
MBP-TOPBP1 91-179	pKR24	pUC57-MBP	91-179
MBP-TOPBP1 91-156	pKR25	pUC57-MBP	91-156
MBP-TOPBP1 91-139	pKR26	pUC57-MBP	91-139
MBP-TOPBP1 128-179	pKR29	pUC57-MBP	128-179
MBP-TOPBP1 138-179	pKR30	pUC57-MBP	138-179
MBP-TOPBP1 128-156	pKR38	pUC57-MBP	128-156
MBP	pOH10	pUC57-MBP	-
MBP-TOPBP1 221-273	pOH6	pUC57-MBP	221-273
MBP-TOPBP1 231-273	pOH7	pUC57-MBP	231-273
MBP-TOPBP1 221-250	pOH8	pUC57-MBP	221-250
MBP-TOPBP1 197-231	pOH9	pUC57-MBP	197-231
MBP-TOPBP1 242-273	pOH11	pUC57-MBP	421-273
MBP-TOPBP1 250-273	pOH12	pUC57-MBP	250-273
MBP-TOPBP1 251 - 273 5Am5	pOH26	pUC57-MBP	251 - 273
MBP-TOPBP1 251 - 273 5Am6	pOH27	pUC57-MBP	251 - 273
MBP-TOPBP1 251 - 273 5Am7	pOH28	pUC57-MBP	251 - 273
MBP-TOPBP1 251 - 273 5Am8	pOH29	pUC57-MBP	251 - 273

2.1.2 Recombinant proteins

The recombinant proteins used in this study were GST, GST-NBS1 1-400, GST-NBS1

1-400 K158E, GST-NBS1 393-763, GST-RAD9Tail, GST-BRCT4&5, MBP, MBP-

BRCT1&2, and 6His-T7-BRCT0-2. Expression vectors for GST-RAD9Tail (Duursma et

al., 2013), GST-BRCT4&5, MBP-BRCT1&2, and 6His-T7-BRCT0-2 (Kim et al, 2020)

have been described. Expression vectors for GST-NBS1 1-400, GST-NBS1 1-400

K158E, and GST-NBS1 393-763 were made using pGEX4T-2 as the backbone. All

proteins were expressed in *E. coli* BL21(DE3) cells, at 37°C for 4 hours, and all

proteins were purified from the soluble fraction according to standard procedures

(detailed below).

2.1.3 Antibodies

We used the following commercially sourced antibodies in this work: Myc

(Millipore Sigma #M4439), MBP (New England Biolabs #E8032S), GST (Millipore

Sigma #05-782), and T7 (Millipore Sigma #69522). We also used our own antibody

against *Xenopus* TOPBP1, HU142, which has been described (Van Hatten et al.,

2002).

2.2 Methods

2.2.1 Xenopus egg extracts

Extracts were prepared as previously described (Van Hatten et al., 2002, Yan et al.,

2006).

2.2.2 IVTT production of proteins and determination of protein concentrations

IVTT reactions were performed using the SP6 TnT® Quick Coupled

Transcription/Translation System (Promega #L2080) according to the

manufacturer's instructions. For Bug-Bunny binding assays we used 20ul of the

IVTT reaction/sample.

2.2.3 Recombinant Protein Expression and Purification

Plasmids were transformed in BL21 (DE3) Competent *E. Coli* cells and grown in LB

media (GST fusion proteins) or LB media with 0.2% glucose (MBP fusion proteins)

with ampicillin. At an OD600 of 0.6-0.8, cells were induced with 0.3 mM IPTG for 4

hours at 37°C. Cells were then harvested by centrifugation (4000 rpm for 20 min)

and washed with phosphate-buffered saline (PBS). Cells were lysed by sonication in

GST column buffer (40% PBS, 60% TEN buffer: 10 mM Tris-Cl, pH 8.0, 1 mM EDTA,

pH 8.0, 100 mM NaCl) supplemented with cOmplete protease inhibitor cocktail

tablets (Roche #04693159001) and 10 mg lysozyme for GST fusion proteins or in

maltose column buffer (20 mM Tris-HCl, pH 7.4, 0.3 M NaCl, and 1 mM EDTA) with

1mM DTT and cOmplete protease inhibitor cocktail tablets for MBP fusion proteins.

Lysates were cleared by centrifugation (15,000 rcf for 20 min) and incubated with

glutathione sepharose (BioVision #6566) for GST fusion proteins or amylose resin

(New England Biolabs #E8021S) for MBP fusion proteins for 1 hour at 4°C. Beads

were poured over a column and washed with column buffer. GST fusion proteins

were eluted with 20 mM glutathione in 75 mM HEPES, pH 7.5, 150 mM NaCl, 5 mM

DTT, and 0.1% Triton-X. MBP fusion proteins were eluted with 10 mM maltose in

maltose column buffer. Proteins were dialyzed overnight into PBS.

2.2.4 Protein binding assays

Three kinds of protein binding assays were utilized. The first involved the addition

of recombinant proteins to XEE (Figures 1B-D). For these assays 25 ul of XEE was

supplemented with 5ug of GST-tagged recombinant protein. DSBs, in the form of

EcoRI-digested lambda DNA, were also optionally added (500 ng). After a 30-minute

incubation the samples were diluted into 400 ul of Binding Buffer (PBS + 0.1% NP-

40) and incubated with glutathione sepharose (BioVision, #6566). The beads were

washed four times with 500 ul of Binding Buffer and then eluted with 2X Sample

Buffer (Millipore Sigma #S3401-10VL). The total and bound fractions were then

analyzed via Western blotting using standard conditions. The second involved

incubation of IVTT proteins with recombinant GST or MBP fusion proteins purified

from *E. coli,* in what we refer to as the Bugs-Bunny system. For these assays 20 ul of

IVTT proteins were mixed with 5 ug of GST or MBP fusion protein and incubated for

60 minutes at 30°C. The mixtures were diluted into 400 ul of Binding Buffer and

then incubated with the appropriate resin (amylose for MBP, New England Biolabs

E8021S; glutathione sepharose for GST, BioVision #6566) for one hour at 4°C.

The beads were washed four times with 500 ul of Binding Buffer and then eluted

with 2X Sample Buffer. The input and bound fractions were then analyzed via

Western blotting using standard conditions. The third type of binding assay (done

once, in Figure 4F) was a variant of the pull-down where MBP recombinant proteins,

purified from *E. coli,* were mixed at 10 ug each with 100 ng of 6His-T7-BRCT0-2, also

purified from *E. coli.* The samples were then treated as described above for the

Bugs-Bunny system.

3. Results

3.1 The TOPBP1-NBS1 interaction can be recapitulated using E. coli- and IVTT-produced proteins.

Previous work from many different laboratories has shown that TOPBP1 and MRN act together to promote DDR signaling, and that NBS1 and TOPBP1 can bind to another (Morishima et al, 2007; Yoo et al, 2009; Duursma et al., 2013; Shiotani et al, 2013; Montales et al, 2022). The domain structures of the two proteins are shown in Figure 1A. As detailed above, TOPBP1 contains nine BRCT domains, named 0-8. Of these, BRCT domains 1,2,5, and 7 contain PBPs. NBS1 is less complex, with three protein-protein interaction motifs, two BRCT domains and one FHA domain. For NBS1 the sole PBP is located in BRCT1. One previous study analyzed the interaction between TOPBP1 and NBS1 in some detail (Yoo et al, 2009). Using XEEs, Yoo and colleagues found that TOPBP1 and NBS1 could co-precipitate under resting conditions, and that the interaction was enhanced when DDR signaling was activated. This work went on to show that TOPBP1 uses its BRCT0-2 region to bind NBS1, and that the relevant region within NBS1 mapped to the amino-terminal 400 amino acids of the protein. Finally, Yoo and colleagues showed that mutations in the PBPs from both TOPBP1 (BRCT1) and NBS1 (BRCT1) prevented efficient interaction. We have recently shown that MRN is important for recruitment of TOPBP1 to long, linearized dsDNA templates that mimic DSBs (Montales et al, 2022). Given these findings, we were motivated to analyze the TOPBP1-NBS1 interaction in more detail.

We first sought to repeat some of the earlier findings by Yoo and colleagues

(Yoo et al, 2009), that a fragment of NBS1 corresponding to amino acids 1-400 could

bind TOPBP1 after incubation in XEE. Accordingly, we expressed and purified from

E. coli two GST-NBS1 1-400 fusion proteins, one corresponding to the wild type

sequence and the other containing a K158E mutation that attenuates the PBP in

BRCT1 of NBS1 (Figure 1A). These proteins, along with GST alone, were added to

XEE in either the presence or absence of linearized 5kb dsDNA, which we have

previously shown will potently activate both ATR and ATM in our XEEs (Montales et

al, 2021; Ruis et al, 2022; Montales et al, 2022). The GST protein complexes were

then recovered back out of the XEE using glutathione-sepharose beads, and the

presence of endogenous TOPBP1 was detected via Western blotting. As displayed in

Figure 1B, TOPBP1 did not bind to GST alone, but it did bind well to the wild type

form of GST-NBS1 1-400. Binding was clearly evident in the sample lacking DSBs,

and was enhanced by the presence of DSBs. By contrast, binding to the K158E form

of GST-NBS1 1-400 was severely attenuated, under both -DSB and +DSB conditions

(Figure 1B). We next asked if there are additional binding sites for TOPBP1 within

NBS1, beyond those present in the 1-400 region. We found that this is not likely, as

TOPBP1 did not bind to a GST fusion containing NBS1 from residue 393 to the end

of the protein, amino acid 763 (Figure 1C). Lastly, we wanted to confirm that the

BRCT0-2 region of TOPBP1 could bind to NBS1 1-400, and for this a myc-tagged

TOPBP1 fragment corresponding to BRCT0-2 was produced by *in vitro* transcription

and translation (IVTT) and added to XEE along with GST-NBS1 1-400. DSBs were

optionally added, and we observed that myc-TOPBP1 BRCT0-2 could readily bind to

GST-NBS1 1-400, but not to GST alone, and that binding was modestly enhanced by

DSBs (Figure 1D). Taken together, the data in Figures 1B-D are all in excellent

agreement with prior work of Yoo et al (Yoo et al, 2009). We note that neither full-

length TOPBP1 nor the myc-tagged BRCT0-2 fragment can bind to GST alone

(Figures 1B,D; see also Kim et al, 2020), and thus moving forward we did not include

the GST alone control for every experiment.

Previous work, and data shown here, reveal that there is a constitutive, or

basal, interaction between TOPBP1 and NBS1 that occurs in XEE even in the absence

of DDR signaling (Figures 1B-D; Yoo et al, 2009). We wanted to study this basal

interaction further, and we first asked if we could recapitulate it using a protein-

binding assay that we nicknamed Bugs-Bunny (Figure 2A). In the Bugs-Bunny

system one binding partner is expressed and purified from *E. coli* (Bugs) and the

other is synthesized by IVTT in rabbit reticulocyte lysates (Bunny). After synthesis

of one binding partner in the IVTT reaction we simply add the purified form of the

other partner to the rabbit reticulocyte lysates. After incubation the samples are

diluted in binding buffer and protein complexes isolated using an affinity resin

corresponding to the tag on the *E. coli* produced protein (for this study the tag is

predominantly GST, but sometimes MBP). This system has several advantages for

the study of protein-protein interactions. One, the endogenous proteins present in

the reticulocyte lysates help to prevent non-specific binding of the target protein to

the resin. Two, as detailed below, the reticulocyte lysates can contain modifying

enzymes, such as casein kinase 2 (CK2), that are important for certain interactions.

Three, and most importantly, because synthesis of a target protein by IVTT is far

less labor intensive than purification of the target from bacterial or insect cells, one

can rapidly screen through a collection of mutants to analyze sequence

determinants required for the interaction. Lastly, in our experience, proteins are

much better behaved, in terms of proper folding and solubility, when made by IVTT

relative to other expression systems.

We first examined binding of GST-NBS1 1-400 to myc-tagged TOPBP1

BRCT0-2 in the Bugs-Bunny assay. As shown in Figure 2B, myc-BRCT0-2 could bind

to wild type GST-NBS1 1-400 under these assay conditions, and binding was greatly

reduced when the K158A form of NBS1 was used. These data show that the

NBS1::TOPBP1 interaction is readily observed using the Bugs-Bunny system, and

thus that XEE is not required for the interaction to occur. We also asked if the PBP

pockets within TOPBP1's BRCT1 and 2 domains are required for the interaction, by

testing the "EE" mutant, where the conserved lysines (K155 and K249) within the

PBPs of both BRCT1 and BRCT2 had been mutated to glutamic acid (Figure 1A). We

found that the EE mutant binds as efficiently as wild type to NBS1 (Figure 2B). Thus,

the NBS1::TOPBP1 interaction, as it occurs in the Bugs-Bunny system, requires the

NBS1 PBP, but not those of TOPBP1. We next assessed the specificity of the

interaction between NBS1 and TOPBP1's BRCT0-2 region by asking if NBS1 could

also bind to TOPBP1's BRCT4&5 domains. Binding assays were run with the

following partners: GST-NBS1 1-400 with myc-tagged TOPBP1 BRCT0-2 or 4&5

domains, and GST fused to the RAD9 Tail domain (RAD9Tail) with the same pair of

TOPBP1 proteins. We included RAD9Tail because we had previously shown that it

binds well to myc-BRCT0-2 in our Bugs-Bunny system (Kim et al, 2020). As shown

in Figure 2C, myc-BRCT0-2 could bind as efficiently to NBS1 as it did to the

RAD9Tail, and myc-BRCT4&5 failed to bind efficiently to either NBS1 or RAD9Tail.

We conclude that the Bugs-Bunny system faithfully recapitulates a specific

interaction between NBS1 and TOPBP1.

Because the NBS1::TOPBP1 interaction requires the NBS1 BRCT1 PBP, we

were curious if the interaction was dependent on phosphorylation. Previous work

from our group has shown that the interaction between TOPBP1 and the RAD9Tail,

which requires phosphorylation of RAD9Tail, can be recapitulated in the Bugs-

Bunny system and this is because the reticulocyte lysates used for IVTT contain CK2,

the kinase responsible for RAD9Tail phosphorylation (Kim et al, 2020). Myc-BRCT0-

2 was produced by IVTT and, prior to addition of the GST fusion proteins, the

samples were optionally treated with lambda phosphatase. As expected,

phosphatase treatment eliminated the RAD9Tail::TOPBP1 interaction, however the

NBS1::TOPBP1 interaction was refractory to phosphatase treatment (Figure 2D).

We conclude that, despite the requirement for the NBS1 BRCT1 PBP, the

NBS1::TOPBP1 interaction does not require the presence of phosphate on TOPBP1.

3.2 NBS1 can bind to both BRCT1 and BRCT2 of TOPBP1

To pursue these observations, we next asked if NBS1 required an intact BRCT0-2

region for binding to TOPBP1. To explore this, we subdivided TOPBP1's BRCT0-2

region into the individual BRCT domains and tested them for binding to NBS1.

Surprisingly, both BRCT1 and BRCT2 could bind to NBS1 (Figure 3A), whereas

BRCT0 could not. Efficient binding of either BRCT1 or BRCT2 to NBS1 was

dependent on NBS1 K158 (Figure 3B), demonstrating that the interactions are

specific and not the result of the isolated BRCT1 or BRCT2 domains becoming

"sticky" proteins upon separation from one another. In addition, while NBS1 readily

binds the isolated BRCT1 or BRCT2 domains, the same is not true of RAD9Tail, as it

could only interact with the contiguous BRCT0-2 region, and not the individual

BRCT1 or BRCT2 domains (Figure 3C). Taken together, the data presented in

Figures 1-3 show that the TOPBP1 BRCT0-2 region interacts with NBS1 in a

fundamentally different manner than it does with RAD9Tail. The RAD9Tail::TOPBP1

interaction requires an intact BRCT0-2 region and is dependent on phosphorylation

(summarized in Figure 3D), whereas the NBS1::TOPBP1 interaction requires neither

an intact BRCT0-2 region nor phosphorylation (Figure 3E).

NBS1 uses its BRCT1 domain to bind directly to TOPBP1.

Having determined that BRCT1 or BRCT2 from TOPBP1 are both capable of

binding to NBS1 we next flipped the problem on its head and asked which domains

within NBS1 1-400 could bind TOPBP1. The constructs used for this are

summarized in Figure 4A. We first wanted to confirm that full-length NBS1 could

bind TOPBP1's BRCT1&2 domains in the Bugs-Bunny system, and to do so we

employed an MBP-BRCT1&2 fusion protein purified from E. coli, together with myc-

tagged NBS1, made by IVTT. As shown in Figure 4B, full-length myc-NBS1 can bind

MBP-BRCT1&2, but not MBP alone. Given that full-length NBS1 is incapable of

binding to MBP alone, moving forward we did not include the MBP alone control in

all experiments. We also observed that MBP-BRCT1&2 could bind the NBS1 1-400

region, but not the 393-763 region (Figure 4C). These data show that despite

switching production of the TOPBP1 region from IVTT to E. coli, and vise-versa for

NBS1, the specificity of the interaction was not altered. We next used MBP-BRCT1&2

to delineate the region within NBS1 to which it binds. We found that myc-tagged

NBS1 1-400 bound well, as expected (Figure 4D). A smaller fragment, amino acids 1-

200 and comprised of the FHA domain plus BRCT1, also bound well to MBP-

BRCT1&2. Further delineation revealed that loss of the FHA domain did not alter

binding, as a fragment containing just the two BRCT domains could bind to

TOPBP1's BRCT1&2 domains (Figure 4D). Interestingly, when the two BRCT

domains were tested for binding in isolation, we saw that BRCT1 could bind well,

but BRCT2 showed no detectable binding. The data in Figure 4D show clearly that

NBS1 BRCT1 can interact with TOPBP1 BRCT1&2, and that NBS1 BRCT2 cannot, but

they left open the possibility that the FHA domain could bind TOPBP1 BRCT1&2. To

explore this we ran the experiment shown in Figure 4E. Unlike NBS1 BRCT1, the

FHA domain showed no detectable binding to TOPBP1. We conclude that it is

BRCT1 from NBS1 that mediates the interaction with TOPBP1's BRCT1 or BRCT2

domains.

Having defined the region of NBS1 that binds TOPBP1, we next asked if the

NBS1::TOPBP1 interaction is direct, or if it requires a bridging factor that might be

present in the reticulocyte lysates used for our binding assays. For this we

expressed and purified an MBP-NBS1 1-200 fusion protein from E. coli. We initially

attempted to purify MBP-NBS1 100-200, however we found that this fusion protein

was insoluble and difficult to express in E. coli. We also produced a 6His-T7-TOPBP1

BRCT0-2 recombinant protein, one that we have used in the past (Kim et al, 2020).

Thus, in this experiment, both proteins were expressed in E. coli and then purified,

thereby eliminating concerns about bridging factors. To block non-specific binding

to the resin the assays were performed in binding buffer supplemented with 1%

BSA. As shown in Figure 4F, 6His-T7-TOBP1 BRCT0-2 could readily bind MBP-NBS1

1-200, but not MBP alone. We conclude that the interaction between NBS1 and

TOPBP1 is direct and does not require any post-translational modifications on

either partner.

3.4 A 29 amino-acid sequence in the central region of TOPBP1 BRCT1 promotes

binding to NBS1 in a transferrable manner.

To learn more about the NBS1::TOPBP1 interaction we decided to identify the

sequences within the TOPBP1 BRCT1 and 2 domains that mediate the interaction.

To do so we engineered a system whereby MBP fusions proteins could be rapidly

produced by IVTT and then used in binding assays with GST-NBS1 1-400. Figure 5A

shows the constructs used for the delineation analysis. As shown in Figure 5B, we

found that all constructs that contained an intact BRCT1 domain would bind to

NBS1, showing that the NBS1-binding capacity of BRCT1 can be transferred to a

heterologous protein (MBP). Further delineation within BRCT1 revealed the

following results: we found that while a fragment corresponding to amino acids 91-

156 could bind NBS1, a slightly smaller fragment, amino acids 91-139, could not

bind (Figure 5C). Thus, the C-terminal border of the NBS1 binding region,

operationally defined, is amino acid 156. We then came in from the N-terminal side

and observed that a fragment running from amino acids 128 to 179 could bind

NBS1, but the fragment from amino acids 138-179 could not (Figure 5D). This

operationally establishes the N-terminal border at position 128. Finally, we asked if

a fragment corresponding to the two borders, 128 to 156, retained the capacity to

bind NBS1 and we found that it did (Figure 5E). These data show that, surprisingly,

an intact BRCT1 domain is dispensable for the NBS1 interaction, and that the ability

to bind NBS1 is encoded by a small, 29 amino-acid sequence centrally located within

the domain. We have named this sequence NBR1 for NBS-binding region 1 (Figure

5F). Interestingly, previous work has shown that the interaction between the BRCT2

domain of XRCC1 and the BRCT domain of DNA ligase III depends on a similar

sequence in XRCC1 as the TOPBP1 NBR1 (Taylor et al, 1998). As detailed in the

Discussion, these data suggest that the TOPBP1 BRCT1::NBS1 BRCT1 interaction is

mechanistically similar to the well-studied XRCC1 BRCT2::DNA ligase III BRCT

interaction.

3.5 A novel mode of BRCT::BRCT interaction for binding of NBS1 to TOPBP1 BRCT2.

We next examined BRCT2 and going into this we expected that its NBR would

be analogous to the BRCT1 NBR1, i.e. centrally located. This, however, was not the

case. Figure 6A shows the constructs used for the BRCT2 analysis, and the construct

that corresponds to the central region is 221-250. As shown in Figure 6B, this

construct could not bind to NBS1, and in the same experiment we see again that the

central region of BRCT1, amino acids 128-156, does bind to NBS1. In addition, we

observed that a fragment corresponding to the N-terminal half of BRCT2, amino

acids 197-231, also failed to bind (Figure 6B). This focused attention on the C-

terminal portion of BRCT2, and we found through a series of N-terminal truncations

that we could remove most of the N-terminal BRCT2 sequences and still retain

binding to NBS1 (Figures 6C&D). These data show that a small, 24-amino acid

region at the extreme C-terminus of BRCT2, running from amino acids 250 to 273, is

sufficient to confer binding to NBS1 (Figure 6E). We named this motif NBR2.

We next wanted to determine which amino acids within TOPBP1's NBR2 are

required for binding to NBS1. For this analysis we systematically mutated all but

two residues within NBR2 to alanine, and we did so in blocks of 5 or 6 amino acids

(Figure 6F). These mutations were introduced in the context of the full BRCT2

domain (amino acids 197-296) fused to MBP. As shown in Figures 6G, mutation of

the extreme C-terminal and N-terminal blocks did not disrupt binding to NBS1. By

contrast, mutation of the internal blocks did impact binding (Figure 6H), and one

mutant (aa 263-268) showed substantially reduced binding. These data show that

that the key residues mediating association of BRCT2 with NBS1 are amino acids

256-268.

3.6 Binding of TOPBP1 BRCT1 and BRCT2 to TOPBP1 BRCT 4&5 relies on the same

motifs that are needed to bind NBS1.

417   Previous work from our group had shown that TOPBP1 self-associates via an

array of BRCT::BRCT domain interactions (Kim et al, 2020). In this previous work

we showed that the BRCT 1&2 pair can bind isolated BRCT domains 1,2, and 5, and

that the BRCT4&5 pair can bind to isolated BRCT domains 1 and 2. Data shown here

demonstrate that NBS1 can bind to isolated BRCT domains 1 and 2 as well, and thus

it was of interest to see if the same determinants used by BRCT1 and BRCT2 to bind

NBS1 are also required for binding to BRCT4&5. As shown in Figure 7A, the MBP-

BRCT1 91-156 construct could bind to GST-BRCT4&5, but not to GST alone. The 91-

139 fragment failed to bind GST-BRCT4&5, while the minimally defined region for

NBS1 binding, 128-156, did bind. This is the same pattern of binding/not binding

that was observed for NBS1 (Figure 5), and thus we conclude that BRCT1 uses the

same motif, NBR1, to bind both NBS1 and TOPBP1 BRCT4&5. For BRCT2, we found

that the minimally defined region for NBS1 binding, amino acids 250-273, was not

sufficient to bind BRCT4&5 (Figures 7B); however, a slightly larger fragment

corresponding to amino acids 242-273 did bind. Thus, BRCT2 requires a handful of

additional N-terminal amino acids to bind BRCT4&5, relative to NBS1, but

nonetheless the determinants are largely overlapping (Figure 7C).

4. Discussion

BRCT domain containing proteins are major players in the cellular response to DNA damage and, as such, it is crucial that we understand how they function and that we fully describe all the different modes by which they interact with their binding partners. Two BRCT proteins that are at the heart of the DDR are TOPBP1 and NBS1. They are known to bind one another, and this interaction is conserved from frogs to man (Morishima et al, 2007; Yoo et al, 2009). Interestingly, in both frogs and humans, TOPBP1 and NBS1 constitutively bind to one another, and the interaction is then enhanced during the DDR (Morishima et al, 2007; Yoo et al, 2009). In this study we analyzed the constitutive interaction using the Bugs-Bunny assay system. It was important to ensure that the NBS1-TOPBP1 interaction could be faithfully recapitulated in Bugs-Bunny. Data in support of a specific interaction include the finding that the NBS1 K158E point mutation abrogates the interaction in Bugs-Bunny (Figure 2B), as it does in XEE (Figure 1B and Yoo et al, 2009). Furthermore, NBS1 does not bind TOPBP1's BRCT4&5 domains (Figure 2C), showing that it is not non-specifically interacting with just any BRCT domain. In addition, TOPBP1's BRCT1&2 domains are highly selective for the NBS1 BRCT1 domain, as binding to the NBS1 BRCT2 or FHA domains is undetectable (Figures 4D&E). Thus, Bugs-Bunny can reveal highly specific interactions, and is thereby suitable for a detailed analysis of the TOPBP1-NBS1 interaction.

Our data support the conclusion that the NBS1 BRCT1 domain can bind directly to both TOPBP1's BRCT1 and BRCT2 domains, and that these interactions

allow NBS1 to bind TOPBP1 in a non-regulated, constitutive manner. There are

several intriguing questions that emerge from this discovery. First, why would

evolution select for two distinct binding sites for NBS1 on TOPBP1? Both

interactions rely on NBS1 K158, and, therefore, it is not the case that NBS1 also has

two binding sites for TOPBP1, as NBS1 likely uses the same interface for both

interactions. It thus appears that NBS1 can occupy one or the other binding site on

TOPBP1, or perhaps two molecules of NBS1 can simultaneously engage TOPBP1 via

the two binding sites. It is also curious that K158, which is part of the NBS1 BRCT1

PBP, is required for binding, however phosphate is not (Figure 2D). One possibility

is that, under resting conditions, the interaction between NBS1 and TOPBP1 serves

to mask the NBS1 PBP, and thereby prevents spurious and/or inappropriate

interactions with phosphorylated proteins. We note that our data show no

requirement for the PBPs present in TOPBP1 BRCT1 or BRCT2 for the interaction

with NBS1 (Figure 2B). For this experiment we mutated the conserved lysine

residues within the two PBPs to glutamic acid, and in data not shown we confirmed

that this mutant cannot bind to the RAD9 Tail. We note that Yoo and colleagues

mutated the same lysine in BRCT1 (K155) that we did, but they also mutated the

neighboring lysine (K154) to generate a KK->AM mutant. They observed that

binding to NBS1 was attenuated by these mutations (Yoo et al, 2009). One

explanation for this discrepancy is that mutation of both lysines (K154 & K155) in

BRCT1 is needed to observe a defect in binding.

One important finding revealed by our work is the remarkable versatility

that TOPBP1's BRCT0-2 region displays in its interactions with various binding

partners. On the one hand, there is the well-studied, phosphate-dependent

interaction with RAD9Tail (and other phospho-substrates), and on the other hand,

as we have shown here, is a very different mode of interaction, involving phosphate-

free and direct binding to NBS1. There are important differences between the

phosphate-dependent and phosphate-free modes of binding. For the former, a

contiguous BRCT0-2 module is required for efficient binding. In the case of the

RAD9Tail, previous work has shown that it is BRCT1 that contacts phosphate

(Rappas et al, 2011), however BRCT1 alone is not sufficient for the interaction

(Figure 3C), and in data not shown we have observed that BRCT0 also plays a

positive role, as the BRCT0-2 region binds RAD9Tail more efficiently than does the

BRCT1&2 region. Thus, the BRCT0-2 continuum is needed for optimal binding in the

phosphate -dependent mode, and nothing could be further from the truth for the

phosphate-independent mode. All that is required for the phosphate-independent

mode are relatively small sequences within the BRCT domains. These sequences,

termed NBR1 and NBR2, are <30 amino acids and both can be transferred to

heterologous proteins such as MBP to allow binding to NBS1.

A critical question that arises from our studies is how NBR1 and NBR2

mediate the interaction with NBS1. NBR1 maps to the $\alpha1$-$\beta2$-$\beta3$ region of BRCT1

(Figures 8A&B). Importantly, previous work on the XRCC1 BRCT2::DNA ligase III

BRCT interaction had shown that a similar region is used by XRCC1 BRCT2 to bind

DNA ligase III (Taylor et al, 1998; Dulic et al, 2001; Cuneo et al, 2011; Hammel et al,

2021). Furthermore, just like TOPBP1 BRCT1, a short peptide sequence from the

analogous position in XRCC1 can mediate binding to DNA ligase III upon transfer to

a heterologous protein (Taylor et al, 1998). In Figure 8C this sequence is underlined

in black, it maps to the β2-β3 region of XRCC1, just as NBR1 maps to the β2-β3

region of TOPBP1's BRCT1. Extensive structural analysis has defined the interface

that XRCC1 uses to bind DNA ligase III, and it is largely produced by the C-terminal

portion of α1 extending into and beyond β2 (Cuneo et al, 2011; Hammel et al, 2021).

Mutagenesis experiments have shown that residues positioned in the C-terminal

region of XRCC1 BRCT2's α1 contribute to the binding (shown in red in Figure 8C),

even though they are not required in the context of the short peptide that confers

binding to heterologous proteins. Given this, it is likely that, within the context of

the properly folded domain, TOPBP1's BRCT1 also relies on a more substantial

interface than what is represented by NBR1 to bind to NBS1. Nonetheless, for both

XRCC1 BRCT2 and TOPBP1 BRCT1, the small transferrable regions retain enough of

the interface to allow binding. This has important implications for the design of

peptide modifiers that can potentially disrupt BRCT::BRCT interactions and thereby

impact DDR signaling, possibly in a therapeutic manner.

Recent data have examined how ECT2, the only other triplet BRCT containing

protein besides TOPBP1, binds to its partner CYK4 (Gómez-Cavazos et al, 2020). In

this scenario, ECT2 uses its BRCT1 domain to bind CYK4 that has been

phosphorylated by PLK – however, despite the requirement for phosphate, the PBP

within ECT2 is dispensable for binding. Instead, ECT2 uses a negatively charged

interface located in the central, β2-β3 region of BRCT1. Figure 8D shows an

alignment of the BRCT1 domains from TOPBP1 and ECT2 and shown in red coloring

are three basic amino acids in the ECT2 domain that mediate binding to CYK4. Note

that despite good overall conservation between these two BRCT domains the basic

residues are not found in TOPBP1, and thus it is unlikely that the analogous

interface in TOPBP1 BRCT2 is able to interact with phosphate. In summary, data

presented here together with previous work show that a binding interface located in

the central, α1-β2-β3 region of certain BRCT domains can mediate BRCT::BRCT

interactions, in the case of XRCC1 and TOPBP1, and phosphate-dependent

interactions in the case of ECT2.

It is presently unclear how NBR2 mediates its interaction with NBS1, as there

are no precedents (to our knowledge) for a BRCT domain to utilize its extreme C-

terminal region for interaction with a heterologous BRCT domain. NBR2 is well

conserved in human TOPBP1, with only one non-conservative substitution (Figure

8D). This 95.8% similarity is substantially higher than the overall similarity between

human and frog TOPBP1 (66%) and is also higher than the similarity between the

BRCT2 domains (79%), strongly suggesting a conservation of function. A similar but

less dramatic pattern is observed when comparing frog TOPBP1 to its Drosophila

ortholog, MUS-101. The NBR2 region shows 45.8% conservation (Figure 8E),

whereas conservation drops to 37.6% when comparing the entire BRCT2 domains.

Future studies will address the possibility that the NBR2 mode of BRCT::BRCT

interaction is found in other binding partners or if it is unique to TOPBP1::NBS1.

Author contribution

W.M.M. conceptualized the project, acquired the funding, and wrote the manuscript

together with O.H. O.H., K.R., and K.M. performed the experiments.

Conflict of interest statement

The authors declare no conflict of interest.

Acknowledgments

This work was supported by NIH grant R01GM12287. We thank Karlene Cimprich

for the gift of the GST-RAD9 Tail expression vector.

References

Bagge J, Oestergaard VH, Lisby M. Functions of TopBP1 in preserving genome integrity during mitosis. Semin Cell Dev Biol. 2021 May;113:57-64. doi: 10.1016/j.semcdb.2020.08.009.

Bigot N, Day M, Baldock RA, Watts FZ, Oliver AW, Pearl LH. Phosphorylation-mediated interactions with TOPBP1 couple 53BP1 and 9-1-1 to control the G1 DNA damage checkpoint. Elife. 2019 May 28;8:e44353. doi: 10.7554/eLife.44353.

Costa A, Diffley JFX. The Initiation of Eukaryotic DNA Replication. Annu Rev Biochem. 2022 Jun 21;91:107-131. doi: 10.1146/annurev-biochem-072321-110228.

Cuneo MJ, Gabel SA, Krahn JM, Ricker MA, London RE. The structural basis for partitioning of the XRCC1/DNA ligase III-α BRCT-mediated dimer complexes. Nucleic Acids Res. 2011 Sep 1;39(17):7816-27. doi: 10.1093/nar/gkr419.

Day M, Oliver AW, Pearl LH. Phosphorylation-dependent assembly of DNA damage response systems and the central roles of TOPBP1. DNA Repair (Amst). 2021 Dec;108:103232. doi: 10.1016/j.dnarep.2021.103232.

Dulic A, Bates PA, Zhang X, Martin SR, Freemont PS, Lindahl T, Barnes DE. BRCT domain interactions in the heterodimeric DNA repair protein XRCC1-DNA ligase III. Biochemistry. 2001 May 22;40(20):5906-13. doi: 10.1021/bi002701e.

Duursma AM, Driscoll R, Elias JE, Cimprich KA. A role for the MRN complex in ATR activation via TOPBP1 recruitment. Mol Cell. 2013 Apr 11;50(1):116-22. doi: 10.1016/j.molcel.2013.03.006.

Gómez-Cavazos JS, Lee KY, Lara-González P, Li Y, Desai A, Shiau AK, Oegema K. A Non-canonical BRCT-Phosphopeptide Recognition Mechanism Underlies RhoA Activation in Cytokinesis. Curr Biol. 2020 Aug 17;30(16):3101-3115.e11. doi: 10.1016/j.cub.2020.05.090.

Hammel M, Rashid I, Sverzhinsky A, Pourfarjam Y, Tsai MS, Ellenberger T, Pascal JM, Kim IK, Tainer JA, Tomkinson AE. An atypical BRCT-BRCT interaction with the XRCC1 scaffold protein compacts human DNA Ligase IIIα within a flexible DNA repair complex. Nucleic Acids Res. 2021 Jan 11;49(1):306-321. doi: 10.1093/nar/gkaa1188.

Hou WH, Chen SH, Yu X. Poly-ADP ribosylation in DNA damage response and cancer therapy. Mutat Res Rev Mutat Res. 2019 Apr-Jun;780:82-91. doi: 10.1016/j.mrrev.2017.09.004.

Kim A, Montales K, Ruis K, Senebandith H, Gasparyan H, Cowan Q, Michael WM.
Biochemical analysis of TOPBP1 oligomerization. DNA Repair (Amst). 2020
Dec;96:102973. doi: 10.1016/j.dnarep.2020.102973.

Leung CC, Glover JN. BRCT domains: easy as one, two, three. Cell Cycle. 2011 Aug
1;10(15):2461-70. doi: 10.4161/cc.10.15.16312.

London RE. XRCC1 - Strategies for coordinating and assembling a versatile DNA
damage response. DNA Repair (Amst). 2020 Sep;93:102917. doi:
10.1016/j.dnarep.2020.102917.

Ma M, Rodriguez A, Sugimoto K. Activation of ATR-related protein kinase upon DNA
damage recognition. Curr Genet. 2020 Apr;66(2):327-333. doi: 10.1007/s00294-
019-01039-w.

Montales K, Kim A, Ruis K, Michael WM. Structure-function analysis of TOPBP1's
role in ATR signaling using the DSB-mediated ATR activation in Xenopus egg
extracts (DMAX) system. Sci Rep. 2021 Jan 11;11(1):467. doi: 10.1038/s41598-020-
80626-1.

Montales K, Ruis K, Lindsay H, Michael WM. MRN-dependent and independent
pathways for recruitment of TOPBP1 to DNA double-strand breaks. PLoS One. 2022
Aug 2;17(8):e0271905. doi: 10.1371/journal.pone.0271905.

Morishima K, Sakamoto S, Kobayashi J, Izumi H, Suda T, Matsumoto Y, Tauchi H, Ide
H, Komatsu K, Matsuura S. TopBP1 associates with NBS1 and is involved in
homologous recombination repair. Biochem Biophys Res Commun. 2007 Nov
3;362(4):872-9. doi: 10.1016/j.bbrc.2007.08.086.

Moudry P, Watanabe K, Wolanin KM, Bartkova J, Wassing IE, Watanabe S, Strauss R,
Troelsgaard Pedersen R, Oestergaard VH, Lisby M, Andújar-Sánchez M, Maya-
Mendoza A, Esashi F, Lukas J, Bartek J. TOPBP1 regulates RAD51 phosphorylation
and chromatin loading and determines PARP inhibitor sensitivity. J Cell Biol. 2016
Feb 1;212(3):281-8. doi: 10.1083/jcb.201507042.

Rappas M, Oliver AW, Pearl LH. Structure and function of the Rad9-binding region of
the DNA-damage checkpoint adaptor TopBP1. Nucleic Acids Res. 2011
Jan;39(1):313-24. doi: 10.1093/nar/gkq743.

Reinhardt HC, Yaffe MB. Phospho-Ser/Thr-binding domains: navigating the cell
cycle and DNA damage response. Nat Rev Mol Cell Biol. 2013 Sep;14(9):563-80. doi:
10.1038/nrm3640.

Ruis K, Huynh O, Montales K, Barr NA, Michael WM. Delineation of a minimal
topoisomerase II binding protein 1 for regulated activation of ATR at DNA double-
strand breaks. J Biol Chem. 2022 Jul;298(7):101992. doi:10.1016/j.jbc.2022.101992.

Sheng ZZ, Zhao YQ, Huang JF. Functional Evolution of BRCT Domains from Binding
DNA to Protein. Evol Bioinform Online. 2011;7:87-97. doi: 10.4137/EBO.S7084.

Shiotani B, Nguyen HD, Håkansson P, Maréchal A, Tse A, Tahara H, Zou L. Two
distinct modes of ATR activation orchestrated by Rad17 and Nbs1. Cell Rep. 2013
May 30;3(5):1651-62. doi: 10.1016/j.celrep.2013.04.018.

Taylor RM, Wickstead B, Cronin S, Caldecott KW. Role of a BRCT domain in the
interaction of DNA ligase III-alpha with the DNA repair protein XRCC1. Curr Biol.
1998 Jul 16;8(15):877-80. doi: 10.1016/s0960-9822(07)00350-8.

Van Hatten RA, Tutter AV, Holway AH, Khederian AM, Walter JC, Michael WM. The
Xenopus Xmus101 protein is required for the recruitment of Cdc45 to origins of
DNA replication. J Cell Biol. 2002 Nov 25;159(4):541-7. doi:
10.1083/jcb.200207090.

Wan B, Hang LE, Zhao X. Multi-BRCT scaffolds use distinct strategies to support
genome maintenance. Cell Cycle. 2016 Oct;15(19):2561-2570. doi:
10.1080/15384101.2016.1218102.

Wardlaw CP, Carr AM, Oliver AW. TopBP1: A BRCT-scaffold protein functioning in
multiple cellular pathways. DNA Repair (Amst). 2014 Oct;22:165-74. doi:
10.1016/j.dnarep.2014.06.004.

Yan S, Lindsay HD, Michael WM. Direct requirement for Xmus101 in ATR-mediated
phosphorylation of Claspin bound Chk1 during checkpoint signaling. J Cell Biol.
2006 Apr 24;173(2):181-6. doi: 10.1083/jcb.200601076.

Yoo HY, Kumagai A, Shevchenko A, Shevchenko A, Dunphy WG. The Mre11-Rad50-
Nbs1 complex mediates activation of TopBP1 by ATM. Mol Biol Cell. 2009
May;20(9):2351-60. doi: 10.1091/mbc.e08-12-1190.

Figures and Legends

Figure 1. NBS1 binds to TOPBP1 in a manner dependent on K158 of NBS1.

A. Schematic representation of the proteins of interest. TOPBP1 contains 9 BRCT

domains, four of which contain PBPs. NBS1 contains two BRCT domains, one of

which harboring a PBP, as well as an N-terminal FHA domain. In this study we

mutated two of the TOPBP1 PBPs and the sole PBP in NBS1, as depicted by the K->E

symbols **B.** Either GST or GST-NBS1 recombinant proteins (5 ug) were added to

25ul of XEE as indicated. EcoRI-digested lambda DNA (referred to as "DSBs", 500 ng,

see Montales et al, 2020) was also included, as indicated. After a 30-minute

incubation the samples were diluted in binding buffer and incubated with

glutathione-sepharose beads. The beads were washed and bound proteins then

eluted and probed by Western blot for the presence of TOPBP1 and the GST

proteins, as indicated. A sample of the total extract, taken prior to dilution of the

samples, was also probed for TOPBP1 (panel "total"). The experiment shown is

representative of two independently performed biological replicates. **C.** Same as (B)

except GST-NBS1 393-763 was utilized. The experiment shown is representative of

two independently performed biological replicates. **D.** Either GST or GST-NBS1

recombinant proteins (5 ug) were added to 25ul of XEE as indicated. Myc-tagged

TOPBP1 BRCT0-2, produced by IVTT, was also added (2.5 ul of the IVTT reaction).

EcoRI-digested lambda DNA DSBs (500 ng) was also included, as indicated. The

samples were then processed as in (B). The experiment shown is representative of

two independently performed biological replicates.

Figure 2. The Bugs-Bunny system recapitulates a specific, phosphate-independent interaction between NBS1 and TOPBP1.

A. Overview of the Bugs-Bunny protein binding assay system. **B.** The indicated GST proteins, expressed and purified from *E. coli*, were incubated in IVTT reactions expressing the indicated myc-tagged proteins. The samples were then processed as described in (A). Samples of the total binding reaction, taken prior to dilution, were also probed for myc, showing that the myc-tagged proteins were present in equivalent amounts across all of the samples (lane "input"). The experiment shown is representative of two independently performed biological replicates. **C.** Same as (B). The experiment shown is representative of two independently performed biological replicates. **D.** Same as (B) except that lambda phosphatase (2ul) was optionally added, as indicated. Phosphatase was added after the IVTT reactions were finished and prior to addition of the GST proteins. The experiment shown is representative of two independently performed biological replicates.

Figure 3. NBS1 independently binds to both BRCT1 and BRCT2 from TOPBP1.

A-C. Bugs-Bunny binding assays, using the indicated proteins, were performed exactly as in Figure 2B. The experiments shown are representative of two independently performed biological replicates. **D-E.** Cartoons highlighted the different ways that RAD9Tail and NBS1 interact with TOPBP1's BRCT0-2 region. See text for details.

Figure 4. NBS1 uses its BRCT1 to bind directly to the TOPBP1 BRCT0-2 region.

A. Cartoons summarizing the domains located within the N-terminus of NBS1 and

the various truncations used in the binding assays. **B-E.** Bugs-Bunny binding assays,

using the indicated proteins, were performed exactly as in Figure 2B with exception

that MBP and not GST fusion proteins were utilized. The experiments shown are

representative of two independently performed biological replicates. **F.** As detailed

in the Methods section, the indicated purified, recombinant proteins were mixed

and then assessed for interaction by MBP pull-down. The experiment shown is

representative of two independently performed biological replicates.

Figure 5. Delineation of the NBS1 binding region within TOPBP1's BRCT1

domain.

A. Cartoons summarizing the various truncations used in the binding assays. These

MBP-fusion proteins were produced by IVTT and used in Bugs-Bunny binding

assays with recombinant GST-NBS1 1-400. **B-E.** Bugs-Bunny binding assays, using

the indicated proteins, were performed exactly as in Figure 2B. The MBP fusions

were detected with an antibody recognizing the MBP tag. The experiments shown

are representative of two independently performed biological replicates. **F.** The

sequence of TOPBP1's BRCT1 domain is shown. The red box highlights the

experimentally defined NBS1 binding region 1.

Figure 6. Delineation of the NBS1 binding region within TOPBP1's BRCT2

domain.

A. Cartoons summarizing the various truncations used in the binding assays. These

MBP-fusion proteins were produced by IVTT and used in Bugs-Bunny binding

assays with recombinant GST-NBS1 1-400. **B-D.** Bugs-Bunny binding assays, using

the indicated proteins, were performed exactly as in Figure 2B. The MBP fusions

were detected with an antibody recognizing the MBP tag. The experiments shown

are representative of two independently performed biological replicates. **E.** The

sequence of TOPBP1's BRCT2 domain is shown. The red box highlights the

experimentally defined NBS1 binding region 2. **F.** Cartoon showing the alanine

scanning mutants used to define the critical amino acids within NBR2 that are

needed to bind NBS1. **G-H.** Bugs-Bunny binding assays, using the indicated proteins,

were performed exactly as in Figure 2B. The experiments shown are representative

of two independently performed biological replicates.

Figure 7. TOPBP1's BRCT1 and BRCT2 use similar sequences to bind TOPBP1's

BRCT4&5 domains as for NBS1.

A-B. Bugs-Bunny binding assays, using the indicated proteins, were performed

exactly as in Figure 2B. The MBP constructs are the same as those depicted in

Figures 5A and 6A. The experiments shown are representative of two independently

performed biological replicates. **C.** The sequences for NBR1 and NBR2 are shown

and the relevant regions for binding to BRCT4&5 are shown.

Figure 8. A comparison of the TOPBP1::NBS1 interactions to previously

studied BRCT::BRCT interactions involving XRCC1, DNA ligase II, and ECT2.

A. The sequences for TOPBP1's BRCT1 and BRCT2 domains are shown. The

canonical BRCT domain secondary structures are also highlighted. Shown in red

lettering are NBR1 and NBR2. **B.** The tertiary structure of human TOPBP1's BRCT0-

2 region is shown at two different angles. The regions corresponding to NBR1 and

NBR2 are indicated and colored in beige. The images were generated using NCBI's

web-based "Structure" protein analysis tool. **C.** The sequence of XRCC1's BRCT1

domain is shown and the canonical BRCT domain secondary structures are

highlighted. Shown in red lettering are amino acids that mediate the interaction

with DNA ligase III. **D.** The BRCT1 domains from TOPBP1 and ECT2 are aligned.

Shown in red lettering are the ECT2 amino acids that mediate the interaction with

phosphorylated CYK4. **E.** The sequence of *Xenopus* TOPBP1's NBR2 region is shown

and compared to its human and *Drosophila* counterparts. Shown in red lettering are

conserved amino acids for the human and fly proteins.

 Figure 1

791   **Figure 2**

794   **Figure 3**

Figure 4

 Figure 5

Figure 6

806   **Figure 7**

 Figure 8

A

```
                β1              α1                    β2              β3              α2              β4          α3
BRCT1  91  DVTISCTSLDKETREDVHHYVQIMGGCVYRDLNVS-VTHLIAGEVGSKKYLVAA-LEKPILLPSWVKELWEKS  179
BRCT2 197  GCTICVTGLSSLDRKEVQRLTALHGGEYTGQLKMNESTHLIVQEAKGQKYECARKWNVHCISVQWFFDSIEKG  273
```

B

NBR1

NBR2

C

XRCC1 BRCT2

```
                β1                    α1                β2              β3          α2              β4          α3
545-GKHFFLYGEFPGDERRKLIRYVTAFNGELEDYMSDRVQFVITAQEWDPSFEEALMDNPSLAFVRPRWIYSCNEKQ-619
                                                        allows MBP to bind DNA ligase III
```
.... TOPBP1 NBR 1

D

BRCT1

```
                β1              α1                β2              β3          α2              β4          α3
TOPBP1 197  GCTICVTGLSSLDRKEVQRLTAL—HGGEYTGQLKMNESTHLIVQEAKGQKYECARKWNVHCISVQWFFDSIEKG  273
  ECT2 176  NLVLCFTGFRK—KEELVKLVNLVHHMGGTIRKDFSSKVTHLIANSTHGDKFRVAVSLGTPIMRADWIIQAWEKR  251
```
.... TOPBP1 NBR 1

E

```
HUMAN 262-YECAKRWNVHCVTTQWFFDSIEKG-284

FROG  251-YECARKWNVHCISVQWFFDSIEKG-273

FLY   271-YKAAIRYKKDVLLPEWIFDSCNRG-295
```