1 **NBS1 binds directly to TOPBP1 via disparate interactions between the NBS1**
2 **BRCT1 domain and the TOPBP1 BRCT1 and BRCT2 domains**
3

Oanh Huynh[1], Kenna Ruis[1], Katrina Montales[1], and W. Matthew Michael[1,2]

[1]Molecular and Computational Biology Section

Department of Biological Sciences

University of Southern California

Los Angeles, CA 90089

[2]Author for correspondence at: mattm@usc.edu

Abstract

The TOPBP1 and NBS1 proteins are key components of DNA repair and DNA-based

signaling systems. TOPBP1 is a multi-BRCT domain containing protein that plays

important roles in checkpoint signaling, DNA replication, and DNA repair. Likewise,

NBS1, which is a component of the MRE11-RAD50-NBS1 (MRN) complex, functions

in both checkpoint signaling and DNA repair. NBS1 also contains BRCT domains, and

previous works have shown that TOPBP1 and NBS1 interact with one another. In

this work we examine the interaction between TOPBP1 and NBS1 in detail. We

report that NBS1 uses its BRCT1 domain to interact with TOPBP1's BRCT1 domain

and, separately, with TOPBP1's BRCT2 domain. Thus, NBS1 can make two distinct

contacts with TOPBP1. We report that recombinant TOPBP1 and NBS1 proteins

bind one another in a purified system, showing that the interaction is direct and

does not require post-translational modifications. Surprisingly, we also report that

intact BRCT domains are not required for these interactions, as truncated versions

of the domains are sufficient to confer binding. For TOPBP1, we find that small 24-

29 amino acid sequences within BRCT1 or BRCT2 allow binding to NBS1, in a

transferrable manner. These data expand our knowledge of how the crucial DNA

damage response proteins TOPBP1 and NBS1 interact with one another and set the

stage for functional analysis of the two disparate binding sites for NBS1 on TOPBP1.

Key words

BRCT domain, TOPBP1, NBS1, MRN complex, protein-protein interaction, scaffold

protein

1. Introduction

The BRCA1 C-terminal (BRCT) domain is an interaction module that allows

the host protein to bind to various biological substrates such as other proteins,

nucleic acids, and poly (ADP-ribose) chains (reviewed by Gerloff et al, 2012; Leung

and Glover, 2011; Reinhardt and Yaffe, 2013; Wan et al, 2016). BRCT proteins are

major players in the DNA damage response (DDR), and it is therefore crucial that we

understand how they function and that we fully describe all the different modes by

which they interact with their binding partners. Regarding how BRCT domains bind

to other proteins, there are two general categories: interactions that require the

BRCT binding substrate to be phosphorylated, and those that do not. For phospho-

driven interactions, some BRCT domains contain a so-called phosphate-binding

pocket (PBP), which features a triad of conserved amino acid residues that interact

with the phosphate group on the substrate protein (reviewed by Leung and Glover,

2011; Reinhardt and Yaffe, 2013; Wan et al, 2016; Day et al, 2021). BRCT domains

that bind phospho-epitopes are typically found in pairs, where one domain within

the pair contains the PBP. While the PBP directly interacts with phosphate, both

BRCT domains within the pair are required for binding as the PBP forms via

intramolecular interactions between the two domains (reviewed by Leung and

Glover, 2011; Reinhardt and Yaffee, 2013; Day et al, 2021). Intramolecular

interactions between BRCT domains also allow binding of the host protein to

substrate in a phospho-independent manner, and both the phospho-dependent and

independent modes of binding have been observed within the same protein. The

yeast Rtt107 protein contains 6 BRCT domains, with the C-terminal two BRCTs

supplying phospho-dependent binding to histone H2A, and the four N-terminal

domains forming a compact, higher-order assembly that binds peptide ligands from

a number of proteins (Wan et al, 2019). The net result is that Rtt107 binds to

chromatin via the phospho-dependent mechanism, and then recruits other proteins

to chromatin via the phospho-independent mechanism.

BRCT domains also make intermolecular contacts to allow assembly of both

homo- and heterotypic multimers. A well-studied example of BRCT::BRCT

intermolecular interaction is found with the XRCC1 and DNA Ligase 3 (LIG3)

proteins (Taylor et al, 1998; Dulic et al, 2001; Beernink et al, 2005; Cuneo et al,

2011; London, 2015; London et al, 2020; Hammel et al, 2021) . Both proteins can

homodimerize via BRCT::BRCT interaction, and the two can also heterodimerize via

their BRCT domains. Recent work from our group was focused on the

Topoisomerase II Binding Protein 1 (TOPBP1) protein, which contains nine BRCT

domains, and we found that TOPBP1 forms oligomers and that the protein is capable

of a series of BRCT::BRCT interactions (Kim et al, 2020). More specifically, we found

that BRCT1&2 could bind to BRCT domains 1,2, and 5, and that BRCT4&5 could bind

BRCTs 1 and 2 but not 5. In this work we examine how TOPBP1 uses its BRCT

domains to bind a heterologous protein, by studying its interaction with the NBS1

protein.

NBS1 is part of a trimeric complex, termed MRN for MRE11-RAD50-NBS1,

that senses DNA double-strand breaks (DSBs) and then plays multiple roles in

ensuring efficient repair of the breaks (reviewed by McCarthy-Leo et al, 2022;

Reginato and Cejka, 2020; Syed and Tainer, 2018). These roles include holding the

two broken ends together, initiating DNA end resection, and activation of the DNA

damage checkpoint kinases ATM and ATR. MRN activates ATM directly (reviewed by

Paull, 2015), and ATR indirectly through its ability to recruit TOPBP1 to the DSB

(Montales et al, 2022). Recent work from our laboratory has shown that loss of MRN

attenuates ATR signaling at DSBs, and TOPBP1's occupancy at the DSB is also

reduced (Montales et al, 2022). TOPBP1 has a well-studied function in directly

interacting with ATR (and its binding partner ATRIP) at sites of damage to stimulate

ATR kinase activity (reviewed by Wardlaw et al, 2015), and thus our recent work

has revealed a pathway for ATR signaling that runs from the DNA end through MRN

to TOPBP1 and then ATR (Montales et al, 2022). One open and important question

for this pathway is: what is the molecular mechanism by which MRN recruits

TOPBP1 to DSBs?

In this study we focused on the NBS1 subunit as previous works have shown

that TOPBP1 and NBS1 bind one another, and that this interaction is conserved from

frogs to man (Morishima et al, 2007; Yoo et al, 2009). Interestingly, in both frogs and

humans, TOPBP1 and NBS1 constitutively bind to one another, and the interaction is

then enhanced during the DSB response (Morishima et al, 2007; Yoo et al, 2009).

Here, we study the constitutive interaction, and we find that NBS1 uses its BRCT1

domain to make direct contacts with TOPBP1's BRCT1 domain and, separately, the

BRCT2 domain. Our data provide a molecular explanation for how MRN recruits

TOPBP1 to DSBs, and they also expand our understanding of how BRCT::BRCT

interactions drive the cellular response to DNA damage.

2. Material and Methods

2.1 Materials

2.1.1 Plasmids

The expression vector to produce GST-RAD9 Tail has previously been

described (Duursma et al., 2013). All other plasmids used in this study are

described below, in table format, and all are *Xenopus* proteins. Specific details about

plasmid construction are available upon request.

Plasmid	Name	Vector	AA coordinates
5Xmyc-BRCT0-2	pHG128	pCS2+MT	1-333
5Xmyc-BRCT0-2 KK155,249EE	pAK42	pCS2+MT	1-333
5Xmyc-BRCT4&5	pJA19	pCS2+MT	480-758
5Xmyc-BRCT0	pHG126	pCS2+MT	1-99
5Xmyc-BRCT1	pHG129	pCS2+MT	100-191
5Xmyc-BRCT2	pJA23	pCS2+MT	191-333
5Xmyc-NBS1	pKM107	pCS2+MT	1-763
5Xmyc-NBS1 1-400	pKM105	pCS2+MT	1-400
5Xmyc-NBS1 1-200	pKR34	pCS2+MT	1-200
5Xmyc-NBS1 1-100	pOH4	pCS2+MT	1-100
5Xmyc-NBS1 100-200	pKR36	pCS2+MT	100-200
5Xmyc-NBS1 200-400	pKR37	pCS2+MT	200-400
5Xmyc-NBS1 393-763	pKM106	pCS2+MT	393-763
5Xmyc-NBS1 1-400 K158A	pOH34	pCS2+MT	1-400
5Xmyc-NBS1 1-175	pOH35	pCS2+MT	1-175

5Xmyc-NBS1 1-150	pOH36	pCS2+MT	1-150
5Xmyc-NBS1 128-400	pOH39	pCS2+MT	128-400
5Xmyc-NBS1 128-400	pOH40	pCS2+MT	150-400
MBP-TOPBP1 1-333	pAK53	pUC57-MBP	1-333
MBP-TOPBP1 91-275	pAK56	pUC57-MBP	91-275
MBP-TOPBP1 91-196	pKR23	pUC57-MBP	91-196
MBP-TOPBP1 91-179	pKR24	pUC57-MBP	91-179
MBP-TOPBP1 91-156	pKR25	pUC57-MBP	91-156
MBP-TOPBP1 91-139	pKR26	pUC57-MBP	91-139
MBP-TOPBP1 128-179	pKR29	pUC57-MBP	128-179
MBP-TOPBP1 138-179	pKR30	pUC57-MBP	138-179
MBP-TOPBP1 128-156	pKR38	pUC57-MBP	128-156
MBP	pOH10	pUC57-MBP	-
MBP-TOPBP1 221-273	pOH6	pUC57-MBP	221-273
MBP-TOPBP1 231-273	pOH7	pUC57-MBP	231-273
MBP-TOPBP1 221-250	pOH8	pUC57-MBP	221-250
MBP-TOPBP1 197-231	pOH9	pUC57-MBP	197-231
MBP-TOPBP1 242-273	pOH11	pUC57-MBP	421-273
MBP-TOPBP1 250-273	pOH12	pUC57-MBP	250-273

2.1.2 Recombinant proteins

The recombinant proteins used in this study were GST, GST-NBS1 1-400,

GST-NBS1 1-400 K158E, GST-NBS1 393-763, GST-RAD9Tail, GST-BRCT4&5, MBP,

MBP-BRCT1&2, and 6His-T7-BRCT0-2. Expression vectors for GST-RAD9Tail

(Duursma et al., 2013), GST-BRCT4&5, MBP-BRCT1&2, and 6His-T7-BRCT0-2 (Kim

et al, 2020) have been described. Expression vectors for GST-NBS1 1-400, GST-NBS1

1-400 K158E, and GST-NBS1 393-763 were made using pGEX4T-3 as the backbone.

All proteins were expressed in *E. coli* BL21(DE3) cells, at 37°C for 4 hours, and all

proteins were purified from the soluble fraction according to standard procedures

(detailed below).

2.1.3 Antibodies

We used the following commercially sourced antibodies in this work: Myc

(Millipore Sigma #M4439), MBP (New England Biolabs #E8032S), GST (Millipore

Sigma #05-782), and T7 (Millipore Sigma #69522). We also used our own antibody

against *Xenopus* TOPBP1, HU142, which has been described (Van Hatten et al.,

2002).

2.2 Methods

2.2.1 Xenopus egg extracts (XEE)

Extracts were prepared as previously described (Van Hatten et al., 2002, Yan

et al., 2006).

2.2.2 IVTT production of proteins and determination of protein concentrations

In-vitro transcription and translation (IVTT) reactions were performed using

the SP6 TnT® Quick Coupled Transcription/Translation System (Promega #L2080)

according to the manufacturer's instructions. For Bug-Bunny binding assays we

used 20µl of the IVTT reaction/sample.

2.2.3 Recombinant Protein Expression and Purification

Plasmids were transformed in BL21 (DE3) Competent *E. Coli* cells and grown

in LB media (GST fusion proteins) or LB media with 0.2% glucose (MBP fusion

proteins) with ampicillin. At an OD600 of 0.6-0.8, cells were induced with 0.3 mM

IPTG for 4 hours at 37℃. Cells were then harvested by centrifugation (4000 rpm for

20 min) and washed with phosphate-buffered saline (PBS). Cells were lysed by

sonication in GST column buffer (40% PBS, 60% TEN buffer: 10 mM Tris-Cl, pH 8.0,

1 mM EDTA, pH 8.0, 100 mM NaCl) supplemented with complete protease inhibitor

cocktail tablets (Roche #04693159001) and 10 mg lysozyme for GST fusion proteins

or in maltose column buffer (20 mM Tris-HCl, pH 7.4, 0.3 M NaCl, and 1 mM EDTA)

with 1mM DTT and complete protease inhibitor cocktail tablets for MBP fusion

proteins. Lysates were cleared by centrifugation (15,000 rcf for 20 min) and

incubated with glutathione sepharose (BioVision #6566) for GST fusion proteins or

amylose resin (New England Biolabs #E8021S) for MBP fusion proteins for 1 hour at

4℃. Beads were poured over a column and washed with column buffer. GST fusion

proteins were eluted with 20 mM glutathione in 75 mM HEPES, pH 7.5, 150 mM

NaCl, 5 mM DTT, and 0.1% Triton-X. MBP fusion proteins were eluted with 10 mM

maltose in maltose column buffer. Proteins were dialyzed overnight into PBS.

2.2.4 Protein binding assays

Three kinds of protein binding assays were utilized. The first involved the

addition of recombinant proteins to XEE (Figures 1B-D). For these assays 25 μl of

XEE was supplemented with 5 µg of GST-tagged recombinant protein. DSBs, in the

form of EcoRI-digested lambda DNA, were also optionally added (500 ng). After a

30-minute incubation the samples were diluted into 400 µl of Binding Buffer (PBS +

0.1% NP-40) and incubated with glutathione sepharose (BioVision, #6566). The

beads were washed four times with 500 µl of Binding Buffer and then eluted with 2X

Sample Buffer (Millipore Sigma #S3401-10VL). The total and bound fractions were

then analyzed via Western blotting using standard conditions. The second involved

incubation of IVTT proteins with recombinant GST or MBP fusion proteins purified

from *E. coli,* in what we refer to as the Bugs-Bunny system. For these assays 20 µl of

IVTT proteins were mixed with 5 µg of GST or MBP fusion protein and incubated for

60 minutes at 30°C. The mixtures were diluted into 400 µl of Binding Buffer and

then incubated with the appropriate resin (amylose for MBP, New England Biolabs

E8021S; glutathione sepharose for GST, BioVision #6566) for one hour at 4°C.

The beads were washed four times with 500 µl of Binding Buffer and then eluted

with 2X Sample Buffer. The input and bound fractions were then analyzed via

Western blotting using standard conditions. The third type of binding assay (done

once, in Figure 4F) was a variant of the pull-down where MBP recombinant proteins,

purified from *E. coli,* were mixed at 10 µg each with 100 ng of 6His-T7-BRCT0-2, also

purified from *E. coli.* The samples were then treated as described above for the

Bugs-Bunny system. ImageJ software was used to quantify the signal intensity of

bound and input samples, and the percent of bound to input was presented in

Figure 2C.

3. Results

3.1 The TOPBP1-NBS1 interaction can be recapitulated using E. coli- and IVTT-produced proteins.

Previous work from many different laboratories has shown that TOPBP1 and MRN act together to promote DDR signaling, and that NBS1 and TOPBP1 can bind to one another, although how this interaction occurs is not known (Morishima et al, 2007; Yoo et al, 2009; Duursma et al., 2013; Shiotani et al, 2013; Montales et al, 2022). The domain structures of the two proteins are shown in Figure 1A. As detailed above, TOPBP1 contains nine BRCT domains, named 0-8. NBS1 is less complex, with three protein-protein interaction motifs, two BRCT domains and one FHA domain. One previous study analyzed the interaction between TOPBP1 and NBS1 in some detail (Yoo et al, 2009). Using *Xenous* egg extracts, Yoo and colleagues found that TOPBP1 and NBS1 could co-precipitate under resting conditions, and that the interaction was enhanced when DDR signaling was activated. This work went on to show that TOPBP1 uses its BRCT0-2 region to bind NBS1, and that the relevant region within NBS1 mapped to the amino-terminal 400 amino acids of the protein. We have recently shown that MRN is important for recruitment of TOPBP1 in frog egg extracts to long, linearized dsDNA templates that mimic DSBs (Montales et al, 2022). Given these findings, we were motivated to analyze the TOPBP1-NBS1 interaction in more detail.

We first sought to repeat some of the earlier findings by Yoo and colleagues (Yoo et al, 2009), that a fragment of *Xenopus* NBS1 corresponding to amino acids 1-

400 could bind to *Xenopus* TOPBP1 after incubation in frog egg extracts.

Accordingly, we expressed and purified from *E. coli* a GST-NBS1 1-400 fusion

protein. This protein, along with GST alone, was added to egg extract in either the

presence or absence of linearized 5kb dsDNA, which we have previously shown will

potently activate both ATR and ATM in our extracts (Montales et al, 2021; Ruis et al,

2022; Montales et al, 2022). The GST protein complexes were then recovered back

out of the extract, using glutathione-sepharose beads, and the presence of

endogenous TOPBP1 was detected via Western blotting. As displayed in Figure 1B,

TOPBP1 did not bind to GST alone, but it did bind to GST-NBS1 1-400. Binding was

evident in the sample lacking DSBs and was modestly enhanced by the presence of

DSBs. We next asked if there are additional binding sites for TOPBP1 within NBS1,

beyond those present in the 1-400 region, and we found that this is not likely as

TOPBP1 did not bind to a GST fusion containing NBS1 from residue 393 to the end

of the protein, amino acid 763 (Figure 1C). Lastly, we wanted to confirm that the

BRCT0-2 region of TOPBP1 could bind to NBS1 1-400, and for this a myc-tagged

TOPBP1 fragment corresponding to BRCT0-2 was produced by *in vitro* transcription

and translation (IVTT) and added to egg extract along with GST-NBS1 1-400. DSBs

were optionally added, and we observed that myc-TOPBP1 BRCT0-2 could readily

bind to GST-NBS1 1-400, but not to GST alone, and that binding was modestly

enhanced by DSBs (Figure 1D). Taken together, the data in Figures 1A-D are all in

excellent agreement with the previous work (Yoo et al, 2009).

As shown here, and by others, a constitutive interaction between TOPBP1

and NBS1 occurs in egg extract even in the absence of DDR signaling (Figures 1B-D;

Yoo et al, 2009). We wanted to study this basal interaction further, and we first

asked if we could recapitulate it using a protein-binding assay that we nicknamed

Bugs-Bunny (Figure 2A). In the Bugs-Bunny system one binding partner is

expressed and purified from *E. coli* (Bugs) and the other is synthesized by IVTT in

rabbit reticulocyte lysates (Bunny). After synthesis of one binding partner by IVTT

we simply add the purified form of the other partner to the rabbit reticulocyte

lysates. After incubation the samples are diluted in binding buffer and protein

complexes isolated using an affinity resin corresponding to the tag on the *E. coli*

produced protein (for this study the tag is predominantly GST, but sometimes MBP).

This system has several advantages for the study of protein-protein interactions.

One, the endogenous proteins present in the reticulocyte lysates help to prevent

non-specific binding of the target protein to the resin. Two, as detailed below, the

reticulocyte lysates can contain modifying enzymes, such as casein kinase 2 (CK2),

that are important for certain interactions. Three, and most importantly, because

synthesis of a target protein by IVTT is far less labor intensive than purification of

the target from bacterial or insect cells, one can rapidly screen through a collection

of mutants to analyze sequence determinants required for the interaction. Lastly, in

our experience, proteins are much better behaved, in terms of proper folding and

solubility, when made by IVTT relative to other expression systems.

258   We first examined binding of GST-NBS1 1-400 to myc-tagged TOPBP1

259 BRCT0-2 in the Bugs-Bunny assay. These are both derivatives of the *Xenopus*

260 proteins, as are all the factors analyzed in this study. As shown in Figure 2B, myc-

261 BRCT0-2 could bind to GST-NBS1 1-400 under these assay conditions, but not to

262 GST alone. These data show that the NBS1::TOPBP1 interaction is readily observed

263 using the Bugs-Bunny system, and thus that egg extract is not required for the

264 interaction to occur. We next assessed the specificity of the interaction between

265 NBS1 and TOPBP1's BRCT0-2 region by asking if NBS1 could also bind to TOPBP1's

266 BRCT4&5 domains. Binding assays were run with the following partners: GST-NBS1

267 1-400 with myc-tagged TOPBP1 BRCT0-2 or 4&5 domains, and GST fused to the

268 RAD9 Tail domain (RAD9Tail) with the same pair of TOPBP1 proteins. We included

269 RAD9Tail because we had previously shown that it binds well to myc-BRCT0-2 in

270 our Bugs-Bunny system (Kim et al, 2020). As shown in Figure 2C, myc-BRCT0-2

271 could bind both NBS1 and the RAD9Tail, and myc-BRCT4&5 failed to bind efficiently

272 to either NBS1 or RAD9Tail. Quantification revealed that binding of myc-TOPBP1

273 BRCT0-2 to NBS1 was slightly more efficient than binding to RAD9Tail, and that

274 binding of myc-BRCT4&5 to either GST substrate was largely undetectable. We

275 conclude that the Bugs-Bunny system faithfully recapitulates a specific interaction

276 between NBS1 and TOPBP1.

278   Both BRCT1 and BRCT2 from TOPBP1, as well as BRCT1 from NBS1, contain

279 PBPs, suggesting that the TOPBP1::NBS1 interaction may be phospho-dependent.

280 Previous work from our group has shown that the interaction between TOPBP1 and

the RAD9Tail, which requires phosphorylation of RAD9Tail, can be recapitulated in

the Bugs-Bunny system and this is because the reticulocyte lysates used for IVTT

contain CK2, the kinase responsible for RAD9Tail phosphorylation (Kim et al, 2020).

Thus, we next asked if including phosphatase in the binding assay would prevent

the NBS1::TOPBP1 interaction, as it does the RAD9Tail::TOPBP1 interaction. Myc-

BRCT0-2 was produced by IVTT and, prior to addition of the GST fusion proteins,

the samples were optionally treated with lambda phosphatase. As expected,

phosphatase treatment eliminated the RAD9Tail::TOPBP1 interaction, however the

NBS1::TOPBP1 interaction was refractory to phosphatase treatment (Figure 2D).

We conclude that the NBS1::TOPBP1 interaction does not require the presence of

phosphate on either binding partner.

3.2 NBS1 can bind to both BRCT1 and BRCT2 of TOPBP1

To pursue these observations, we next asked if NBS1 required an intact

BRCT0-2 region to associate with TOPBP1. To explore this, we subdivided TOPBP1's

BRCT0-2 region into the individual BRCT domains and tested them for binding to

NBS1. Surprisingly, both BRCT1 and BRCT2 could bind to NBS1 (Figure 3A),

whereas BRCT0 could not. In addition, while NBS1 readily binds the isolated BRCT1

or BRCT2 domains, the same is not true of RAD9Tail, as it could only interact with

the contiguous BRCT0-2 region, and not the individual BRCT1 or BRCT2 domains

(Figure 3B). Taken together, the data presented in Figures 1-3 show that the

TOPBP1 BRCT0-2 region interacts with NBS1 in a fundamentally different manner

than it does with RAD9Tail. The RAD9Tail::TOPBP1 interaction requires an intact

BRCT0-2 region and is dependent on phosphorylation (summarized in Figure 3C),

whereas the NBS1::TOPBP1 interaction requires neither an intact BRCT0-2 region

nor phosphorylation (Figure 3D).

3.3 NBS1 uses its BRCT1 domain to bind TOPBP1.

309   We next examined the sequence determinants within NBS1 that allow

binding to TOPBP1's BRCT1&2 region. The constructs used for this are summarized

in Figure 4A. We first confirmed that full-length NBS1 could bind TOPBP1's

BRCT1&2 domains in the Bugs-Bunny system, and to do so we employed an MBP-

TOPBP1 BRCT1&2 fusion protein purified from *E. coli*, together with myc-tagged

NBS1, made by IVTT. As shown in Figure 4B, full-length myc-NBS1 can bind MBP-

TOPBP1 BRCT1&2, but not MBP alone. We also observed that MBP- TOPBP1

BRCT1&2 could bind the NBS1 1-400 region, but not the 393-763 region (Figure

4C). These data show that despite switching production of the TOPBP1 region from

IVTT to *E. coli*, and vise-versa for NBS1, the ability to interact was not altered. We

next used MBP-TOPBP1 BRCT1&2 to delineate the region within NBS1 to which it

binds. The N-terminal portion of NBS1 was subdivided into three fragments, 1-100

(FHA domain), 100-200 (BRCT1), and 200-400 (BRCT2 plus additional sequences,

see Figure 4A), and we observed that only the BRCT1 domain could bind to TOPBP1

BRCT1&2 (Figure 4D). Thus, unlike TOPBP1 where both BRCT1 and BRCT2 bind

NBS1, for NBS1 just BRCT1 and not BRCT2 binds to TOPBP1.

3.4 Direct interaction between NBS1 and TOPBP1.

An important question became is the NBS1::TOPBP1 interaction direct, or

does it require a bridging factor that might be present in the reticulocyte lysates

used for our binding assays? To answer this we expressed and purified an MBP-

NBS1 1-200 fusion protein from *E. coli* (Figure 5A). We attempted to purify MBP-

NBS1 100-200, however we found that this protein was insoluble and difficult to

express in *E. coli*. We also produced a 6His-T7-TOPBP1 BRCT0-2 recombinant

protein (Figure 5A), one that we have used in the past (Kim et al, 2020). Thus, in this

experiment, both proteins were expressed in *E. coli* and then purified, thereby

eliminating concerns about bridging factors. To block non-specific binding to the

resin the assays were performed in binding buffer supplemented with 1% BSA. As

shown in Figure 5B, 6His-T7-TOBP1 BRCT0-2 could readily bind MBP-NBS1 1-200,

but not MBP alone, and binding was not inhibited by inclusion of phosphatase. We

conclude that the interaction between NBS1 and TOPBP1 is direct and does not

require any post-translational modifications on either partner.

3.5 No requirement for phosphate-binding pockets in the TOPBP1::NBS1 interaction.

Our recent study (Kim et al, 2020), together with data shown here, have

demonstrated that TOPBP1's BRCT1&2 region interacts with several different BRCT

domains, including TOPBP1's BRCT1, BRCT2, and BRCT5 domains as well as BRCT1

from NBS1 (Figure 5C). One feature that all these domains have in common is a PBP.

Even though we have shown that phosphate *per se* is not needed for the

TOPBP1::NBS1 interaction (Figures 2D and 5B), it was still possible that the PBPs

played a role in the binding. To test this idea we produced mutant proteins where a

key and highly conserved lysine residue within the PBP was mutated. For TOPBP1

we mutated both K155 in BRCT1 and K249 in BRCT2 to glutamic acid, in the context

of a myc-tagged BRCT0-2 fragment. Previous work with human TOPBP1 has shown

that a similar K to E substitution in BRCT1 prevents binding to RAD9 (Rappas et al,

2011), however we observed that the KK155,249EE double mutant could still

readily bind to GST-NBS1 1-400 (Figure 5D). For NBS1, we initially made the same K

to E mutation at K158 in BRCT1 and observed that this mutant could not bind to

TOPBP1 (data not shown). Because this charge-reversal mutation might non-

specifically disrupt folding of the NBS1 domains we also tested a more subtle

mutation, substituting K158 with alanine. The K158A mutation is still expected to

inactivate the PBP, as has been shown for the BRCA1 PBP (Botuyan et al, 2004),

however we observed that it had no impact on binding to TOPBP1 (Figure 5E). Thus,

for both TOPBP1 and NBS1, an intact PBP is dispensable for their interaction.

3.6 Intact BRCT domains are not required for the TOPBP1::NBS1 interaction.

To learn more about the NBS1::TOPBP1 interaction we asked how creating

truncations of TOPBP1's BRCT0-2 region would impact binding to NBS1 1-400. A

series of deletion mutants were constructed as fusions to MBP and are shown in

Figure S2A. These MBP fusions were produced by IVTT and tested for binding

against GST-tagged NBS1 1-400. We found that MBP-BRCT0-2 (amino acids 1-333)

and MBP-BRCT1&2 (amino acids 91-275) could bind NBS1 (Figure S2B). We next

asked if the linker sequence between BRCTs 1 and 2 influenced binding of BRCT1,

by comparing a construct containing the linker (amino acids 91-196) to one that

does not (amino acids 91-179). As shown in Figure S2B, both proteins could bind

NBS1. Moving forward, we next asked if BRCT1 must be in an intact form to bind to

NBS1. Structural analysis of BRCT domains has shown that they are highly ordered,

using a series of alpha helices and beta sheets to fold into a unit with a conserved

topology (reviewed by Gerloff et al, 2012; Leung and Glover, 2013; Reinhardt and

Yaffe, 2013; Wan et al, 2016). Thus, our expectation was that by making deletions

within BRCT1 the domain would lose its ability to fold properly and binding to NBS1

would be lost. This, however, was not the case. The Conserved Domain sequence

analysis tool that is available on the NCBI Structure website

(www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi) defines *Xenopus* TOPBP1 BRCT1

as running from amino acid 108 to 179. As shown in Figure 6A&B, and we observed

that a fragment containing amino acids 128-179, and therefore missing the N-

terminal 20 amino acids of BRCT1, could bind well to NBS1. A slightly smaller piece,

corresponding to amino acids 138-179, could not bind NBS1 (Figure 6B). Intrigued

by these observations we next asked if loss of C-terminal sequences within

TOPBP1's BRCT1 would impact binding to NBS1. We found that truncation of the C-

terminal 23 amino acids of BRCT1, using a construct containing amino acids 91-156,

did not disrupt binding and, furthermore, that an even smaller fragment, amino

acids 128-156, could bind efficiently to NBS 1 1-400 (Figure 6C). Thus, an intact

BRCT1 domain is not required for its interaction with NBS1. To see if this is also true

of TOPBP1's BRCT2 we generated a series of deletion constructs (Figure S2C) and

tested them for binding to GST-NBS1 1-400. These results showed that a small

fragment of 23 amino acids, corresponding to residues 250-273, was sufficient for

binding to NBS1 (Figures S2D&E). We conclude that small sequences within both

BRCT1 and BRCT2 of TOPBP1 can confer binding to NBS1 onto a heterologous

protein, in this case MBP.

We next examined sequence determinants within the NBS1 BRCT domain

that allow binding to TOPBP1. For this we generated the series of deletion mutants

shown in Figure 6D, and we tested them for binding to *E. coli*-produced MBP-

TOPBP1 BRCT1&2. As expected, we found that the fragment corresponding to intact

FHA and BRCT1 domains (1-200) could bind efficiently to TOPBP1 (Figure 6E). We

also observed that a fragment where the carboxyl terminal 25 amino acids of BRCT1

were removed (1-175) retained the ability to bind TOPBP1. Thus, like TOPBP1's

BRCT1&2 domains, NBS1's BRCT1 domain can be truncated without attenuating the

interaction. We then asked if removing more of the carboxyl terminal sequences

impacted binding and found that neither a fragment corresponding to amino acids

1-150 nor one corresponding to 1-128 could bind TOPBP1. We also examined

amino-terminal deletions and found that removing the amino-terminal half of the

NBS1 BRCT1 domain did little to inhibit the interaction with TOPBP1 (Figure 6E).

These data show that, like TOPBP1's BRCT1 and BRCT2 domains, the NBS1 BRCT1

domain need not be in an intact and properly folded configuration to bind TOPBP1.

Data presented thus far show that, for all the relevant BRCT domains in this

study, binding persists even after truncation of the domain. These curious results

inspired us to ask if the interaction required that at least one partner contain an

intact BRCT domain. For example, if the NBS1 BRCT1 domain and the TOPBP1

BRCT1 domain are both in a truncated form, would the interaction be lost? As it

turned out, the answer is no. For this experiment we used an assay system where

both binding partners were produced by IVTT. This was because all the required

constructs for this experiment were available for protein production by IVTT, and

thus there was no need to remake them for expression in *E. coli*. Proteins were

produced by IVTT and then the lysates were mixed, to allow interaction. The

samples were then diluted in binding buffer and incubated with maltose beads,

followed by washing, elution, and Western blotting. As shown in Figure 6F, binding

was readily observed when the BRCT1 domains for both TOPBP1 and NBS1 were

intact (lane 1). Similarly, binding was observed when TOPBP1 BRCT1 was intact and

NBS1 BRCT1 was truncated (lane 2), and vise versa (lane 3). These results are all in

accordance with the data derived from the Bugs-Bunny assays shown in Figures 6B,

C, and E. When both binding partners where in a truncated form we also observed

efficient binding (lane 4), and binding was not observed when the NBS1 derivative

was truncated further (lane 5), consistent with data shown in Figure 6E. Based on

these data, we conclude that neither the TOPBP1 BRCT1 domain nor its counterpart

in NBS1 need to be in an intact and properly folded configuration for the interaction

to occur.

4. Discussion

In this study we examined how two crucial DDR proteins, TOPBP1 and NBS1, interact with one another. Prior to this work, it was known that TOPBP1 and NBS1 can form a complex, in a manner conserved from frogs to man (Morishima et al, 2007; Yoo et al, 2009). However, it was not known which domains within either protein control the interaction, and neither was it known if the interaction was direct or mediated by a bridging factor. Here, we show that the NBS1 BRCT1 domain binds both the BRCT1 and BRCT2 domain from TOPBP1. These are specific interactions, as revealed by our finding that NBS1 binds well to TOPBP1 BRCTs 1&2 but shows very little affinity for BRCTs 4&5 (Figure 2C). Likewise, TOPBP1 binds NBS1 BRCT1 efficiently but there is no detectable binding to either the FHA or BRCT2 domains from NBS1 (Figure 4D). Our data also show that the interaction between TOPBP1 and NBS1 is constitutive, and it is also direct and independent of any post translational modifications (Figure 5). Thus, at least a subset of MRN complexes contain TOPBP1 even under resting conditions. In our previous work we showed that immunodepletion of MRN from frog egg extracts with antibodies targeting NBS1 co-depletes all detectable MRE11 but does not influence the levels of TOPBP1 (Montales et al, 2022). This shows that the association of TOPBP1 with NBS1 in egg extract is not as strong as the association of NBS1 with other members of MRN, although it is possible that the antibodies used for depletion dislodge TOPBP1 from NBS1.

What might be the function of the TOPBP1::NBS1 interaction(s)? In addition,

a related question is why did TOPBP1 evolve two disparate binding sites for NBS1,

positioned closely together at the amino terminus of the protein? We propose that

one function of the interaction is to recruit TOPBP1 to sites of damage so that it can

activate the ATR kinase. Three previous studies, all performed with frog egg

extracts, have implicated MRN in recruitment of TOPBP1 to stalled replication forks

(Lee and Dunphy, 2013), to an ATR-activating DNA structure comprised of M13

ssDNA to which primers are annealed (Duursma et al, 2013), and to 5kb linear

dsDNAs that mimic DSBs (Montales et al, 2022). Thus, it is clear that MRN is

important for how TOPBP1 arrives at sites of damage, and we propose that at least

one of the interactions defined in this paper plays a role in TOPBP1 recruitment for

the purpose of ATR signaling. But why does have TOPBP1 harbor more than one site

for interaction with NBS1? One possibility is that TOPBP1 also collaborates with

MRN is the physical repair of a DSB. Indeed, a recent study using Repair-seq to order

DNA repair proteins into common pathways showed that TOPBP1 clusters with

NBS1, MRE11, and RAD50 in that all four show a common repair defect upon their

depletion (Hussmann et al, 2021). Thus, it may be that TOPBP1 interacts with NBS1

differently for ATR signaling versus DNA repair, and if so, this would explain the

multiple binding sites. Clearly, further work, combining mutagenesis with functional

assays, is needed to clarify the roles of the two NBS1 binding sites on TOPBP1.

Another important question raised by our studies is how are the BRCT

domains from NBS1 and TOPBP1 interacting with one another? There is precedent

for BRCT domains to form both homo- and heterodimers. The best studied example

of this is found with the XRCC1 and LIG3 proteins (reviewed most recently by

London, 2020). XRCC1 has two BRCT domains, with BRCT2 being the one that binds

to itself as well as to the single BRCT domain within LIG3, which also interacts with

itself. Our previous work has shown that TOPBP1's BRCT1 and 2 domains can

interact with one another, and we have shown here that they also bind to NBS1

BRCT1. Thus, the TOPBP1::NBS1 system is similar to XRCC1::LIG3 in that both

systems can form homo- and heterotypic interactions. Recent structural analysis of

a human XRCC1 BRCT1::LIG3 BRCT dimer revealed that the two BRCT domains bind

one another largely through electrostatic interactions between residues located in

the alpha helix 1 portion of the respective domains (Hammel et al, 2021). Thus, it

will be of interest in future work to determine if a similar molecular interface

controls the interactions between TOPBP1 and NBS1.

Work presented here also reveals the remarkable versatility that TOPBP1's

BRCT0-2 region displays in its interactions with various binding partners. On the

one hand, there is the well-studied, phosphate-dependent interaction with RAD9Tail

(and other phospho-substrates), and on the other hand, as we have shown here, is a

very different mode of interaction, involving phosphate-free and direct binding to

NBS1. There are important differences between the phosphate-dependent and

phosphate-free modes of binding. For the former, a contiguous BRCT0-2 module is

required for efficient binding. In the case of the RAD9Tail, previous work has shown

that it is BRCT1 that contacts phosphate (Rappas et al, 2011), however BRCT1 alone

is not sufficient for the interaction (Figure 3B), and in data not shown we have

observed that BRCT0 also plays a positive role, as the BRCT0-2 region binds

RAD9Tail more efficiently than does the BRCT1&2 region. Thus, the BRCT0-2

continuum is needed for optimal binding in the phosphate -dependent mode, while

the phosphate-free mode can occur with isolated domains.

Another unexpected finding reported in this work is that interaction between

the TOPBP1 and NBS1 BRCT domains can occur in the context of truncated, and thus

improperly folded, domains. As shown in Figures 6B&C, for TOPBP1's BRCT1, we

used MBP fusions to ask if truncation of the intact domain would impede binding

and found that it did not. Indeed, we were able to show that a small peptide of just

29 amino acids can confer the ability to bind NBS1 when grafted on to MBP.

Similarly, we found that an even smaller peptide from TOPBP1's BRCT2 domain has

the same properties (Figures S2D&E). For NBS1, we found that removal of large

portions of BRCT1 did not impair binding to TOPBP1 (Figure 6E). In Figure 7A the

amino acid sequences of the three BRCT domains studied here are shown. For

TOPBP1, the residues that allow biding to NBS1, in a transferrable manner, are

highlighted in red. Also highlighted in red are the residues within NBS1 that

represent the minimal region sufficient for binding to TOPBP1, as defined by

deletion analysis (Figure 6E). We do not yet know if this sequence is transferrable,

but this seems likely. Also shown is the canonical arrangement of beta sheets and

alpha helices that promote proper folding of BRCT domains. While the minimally

defined region within BRCT2 shows good overlap with the corresponding sequence

from NBS1, there is very poor overlap between the two TOPBP1 sequences, perhaps

suggesting that TOPBP1's BRCT1 uses a different binding mode to interact with

NBS1 than does its BRCT2 domain.

Given these results with truncations we went on to ask if binding could still

occur when both partners where in a truncated form. We found that it could as

truncated TOPBP1 BRCT1 clearly interacts with a truncated form of NBS1 BRCT1

(Figure 6F). This interaction is specific as the NBS1 1-175 truncation binds both

intact TOPBP1 BRCT1 (91-179) and the truncated form (128-179), but further

truncation of NBS1 to yield 1-150 yields a protein unable to bind either intact or

truncated TOPBP1 BRCT1 (Figure 6F). Thus, truncated TOPBP1 BRCT1 binds to

NBS1 with the same specificity as the intact form. While these results are difficult to

reconcile with the notion that proper folding of an intact BRCT domain is a

prerequisite for its ability to specifically interact with any given binding partner,

they are not without precedent. Previous work has shown that a truncated form of

XRCC1 retains the ability to bind to LIG3 (Taylor et al, 1998). Indeed, this work

showed that a small peptide of 20 amino acids can be transferred to MBP, and this

confers binding to LIG3. Furthermore, binding of the peptide to LIG3 was

stoichiometric and of equivalent efficiency as the full-length XRCC1 protein (Taylor

et al, 1998). The sequence of XRCC1 BRCT2 is shown in Figure 7B, and the LIG3-

binding peptide underlined in black. Interestingly, this peptide maps to a similar

position within its BRCT domain as does the peptide from TOPBP1's BRCT1 that

confers binding to NBS1, as both sequences are comprised of the residues that form

beta sheet 3 within an intact domain, along with some flanking sequences (compare

Figures 7A and 7B). Curiously, however, recent structural studies of the

XRCC1::LIG3 complex have defined electrostatic interactions between residues

largely located in alpha helix 1 as comprising the binding interface for the

XRCC1::LIG3 interaction (Hammel et al, 2021). Thus, it is currently unclear how the

XRCC1 20mer peptide maintains key properties of the interaction with LIG3 when

the structural studies suggest the binding interface resides elsewhere in the domain.

This question is relevant to the work presented here, given that small peptides from

TOPBP1 can also confer binding to NBS1.

563   Lastly, we note that the crystal structure of human TOPBP1 in complex with a

phospho-peptide from RAD9 has been solved (Day et al, 2018), and thus it was of

interest to visualize where in this structure the two NBS1-binding peptides reside.

As shown in Figure 7C, left panel, much of the NBS1 binding peptide from TOPBP1's

BRCT1 is solvent-exposed and overlaps with the region that binds RAD9 (in Figure

7C the NBS1-binding peptide is colored in green and the RAD9 peptide is colored in

yellow). For the NBS1-binding peptide from BRCT2 we see that it is also solvent-

exposed and thus, in principle, able to interact with heterologous proteins (Figure

7C, right panel). Having defined these two regions of TOPBP1 as able to interact

with NBS1, the next important steps will be to first map the amino acids that are

crucial for the interaction and to then test these mutants in functional assays for

ATR activation and DNA repair, and such experiments are now in progress.

Author contribution

W.M.M. conceptualized the project, acquired the funding, and wrote the manuscript

together with O.H. O.H., K.R., and K.M. performed the experiments.

Conflict of interest statement

The authors declare no conflict of interest.

Acknowledgments

This work was supported by NIH grant R01GM12287. We thank Karlene Cimprich

for the gift of the GST-RAD9 Tail expression vector.

References
587
588 Bagge J, Oestergaard VH, Lisby M. Functions of TopBP1 in preserving genome
589 integrity during mitosis. Semin Cell Dev Biol. 2021 May;113:57-64. doi:
590 10.1016/j.semcdb.2020.08.009.
591
592 Beernink PT, Hwang M, Ramirez M, Murphy MB, Doyle SA, Thelen MP. Specificity of
593 protein interactions mediated by BRCT domains of the XRCC1 DNA repair protein. J
594 Biol Chem. 2005 Aug 26;280(34):30206-13. doi: 10.1074/jbc.M502155200.
595
596 Bigot N, Day M, Baldock RA, Watts FZ, Oliver AW, Pearl LH. Phosphorylation-
597 mediated interactions with TOPBP1 couple 53BP1 and 9-1-1 to control the G1 DNA
598 damage checkpoint. Elife. 2019 May 28;8:e44353. doi: 10.7554/eLife.44353.
599
600 Botuyan MV, Nominé Y, Yu X, Juranic N, Macura S, Chen J, Mer G. Structural basis of
601 BACH1 phosphopeptide recognition by BRCA1 tandem BRCT domains. Structure.
602 2004 Jul;12(7):1137-46. doi: 10.1016/j.str.2004.06.002. PMID: 15242590; PMCID:
603 PMC3652423.
604
605 Costa A, Diffley JFX. The Initiation of Eukaryotic DNA Replication. Annu Rev
606 Biochem. 2022 Jun 21;91:107-131. doi: 10.1146/annurev-biochem-072321-110228.
607
608 Cuneo MJ, Gabel SA, Krahn JM, Ricker MA, London RE. The structural basis for
609 partitioning of the XRCC1/DNA ligase III-α BRCT-mediated dimer complexes.
610 Nucleic Acids Res. 2011 Sep 1;39(17):7816-27. doi: 10.1093/nar/gkr419.
611
612 Day M, Rappas M, Ptasinska K, Boos D, Oliver AW, Pearl LH. BRCT domains of the
613 DNA damage checkpoint proteins TOPBP1/Rad4 display distinct specificities for
614 phosphopeptide ligands. Elife. 2018 Oct 8;7:e39979. doi: 10.7554/eLife.39979.
615
616 Day M, Oliver AW, Pearl LH. Phosphorylation-dependent assembly of DNA damage
617 response systems and the central roles of TOPBP1. DNA Repair (Amst). 2021
618 Dec;108:103232. doi: 10.1016/j.dnarep.2021.103232.
619
620 Dulic A, Bates PA, Zhang X, Martin SR, Freemont PS, Lindahl T, Barnes DE. BRCT
621 domain interactions in the heterodimeric DNA repair protein XRCC1-DNA ligase III.
622 Biochemistry. 2001 May 22;40(20):5906-13. doi: 10.1021/bi002701e.
623
624 Duursma AM, Driscoll R, Elias JE, Cimprich KA. A role for the MRN complex in ATR
625 activation via TOPBP1 recruitment. Mol Cell. 2013 Apr 11;50(1):116-22. doi:
626 10.1016/j.molcel.2013.03.006.
627
628 Gerloff DL, Woods NT, Farago AA, Monteiro AN. BRCT domains: A little more than
629 kin, and less than kind. FEBS Lett. 2012 Aug 14;586(17):2711-6. doi:
630 10.1016/j.febslet.2012.05.005.
631

632 Hammel M, Rashid I, Sverzhinsky A, Pourfarjam Y, Tsai MS, Ellenberger T, Pascal JM,
633 Kim IK, Tainer JA, Tomkinson AE. An atypical BRCT-BRCT interaction with the
634 XRCC1 scaffold protein compacts human DNA Ligase IIIα within a flexible DNA
635 repair complex. Nucleic Acids Res. 2021 Jan 11;49(1):306-321. doi:
636 10.1093/nar/gkaa1188.
637
638 Hou WH, Chen SH, Yu X. Poly-ADP ribosylation in DNA damage response and cancer
639 therapy. Mutat Res Rev Mutat Res. 2019 Apr-Jun;780:82-91. doi:
640 10.1016/j.mrrev.2017.09.004.
641
642 Hussmann JA, Ling J, Ravisankar P, Yan J, Cirincione A, Xu A, Simpson D, Yang D,
643 Bothmer A, Cotta-Ramusino C, Weissman JS, Adamson B. Mapping the genetic
644 landscape of DNA double-strand break repair. Cell. 2021 Oct 28;184(22):5653-
645 5669.e25. doi: 10.1016/j.cell.2021.10.002.
646
647 Kim A, Montales K, Ruis K, Senebandith H, Gasparyan H, Cowan Q, Michael WM.
648 Biochemical analysis of TOPBP1 oligomerization. DNA Repair (Amst). 2020
649 Dec;96:102973. doi: 10.1016/j.dnarep.2020.102973.
650
651 Lee J, Dunphy WG. The Mre11-Rad50-Nbs1 (MRN) complex has a specific role in the
652 activation of Chk1 in response to stalled replication forks. Mol Biol Cell. 2013
653 May;24(9):1343-53. doi: 10.1091/mbc.E13-01-0025.
654
655 Leung CC, Glover JN. BRCT domains: easy as one, two, three. Cell Cycle. 2011 Aug
656 1;10(15):2461-70. doi: 10.4161/cc.10.15.16312.
657
658 London RE. The structural basis of XRCC1-mediated DNA repair. DNA Repair
659 (Amst). 2015 Jun;30:90-103. doi: 10.1016/j.dnarep.2015.02.005.
660
661 London RE. XRCC1 - Strategies for coordinating and assembling a versatile DNA
662 damage response. DNA Repair (Amst). 2020 Sep;93:102917. doi:
663 10.1016/j.dnarep.2020.102917.
664
665 Ma M, Rodriguez A, Sugimoto K. Activation of ATR-related protein kinase upon DNA
666 damage recognition. Curr Genet. 2020 Apr;66(2):327-333. doi: 10.1007/s00294-
667 019-01039-w.
668
669 McCarthy-Leo C, Darwiche F, Tainsky MA. DNA Repair Mechanisms, Protein
670 Interactions and Therapeutic Targeting of the MRN Complex. Cancers (Basel). 2022
671 Oct 27;14(21):5278. doi: 10.3390/cancers14215278.
672
673 Montales K, Kim A, Ruis K, Michael WM. Structure-function analysis of TOPBP1's
674 role in ATR signaling using the DSB-mediated ATR activation in Xenopus egg
675 extracts (DMAX) system. Sci Rep. 2021 Jan 11;11(1):467. doi: 10.1038/s41598-020-
676 80626-1.
677

678 Montales K, Ruis K, Lindsay H, Michael WM. MRN-dependent and independent
679 pathways for recruitment of TOPBP1 to DNA double-strand breaks. PLoS One. 2022
680 Aug 2;17(8):e0271905. doi: 10.1371/journal.pone.0271905.
681
682 Morishima K, Sakamoto S, Kobayashi J, Izumi H, Suda T, Matsumoto Y, Tauchi H, Ide
683 H, Komatsu K, Matsuura S. TopBP1 associates with NBS1 and is involved in
684 homologous recombination repair. Biochem Biophys Res Commun. 2007 Nov
685 3;362(4):872-9. doi: 10.1016/j.bbrc.2007.08.086.
686
687 Moudry P, Watanabe K, Wolanin KM, Bartkova J, Wassing IE, Watanabe S, Strauss R,
688 Troelsgaard Pedersen R, Oestergaard VH, Lisby M, Andújar-Sánchez M, Maya-
689 Mendoza A, Esashi F, Lukas J, Bartek J. TOPBP1 regulates RAD51 phosphorylation
690 and chromatin loading and determines PARP inhibitor sensitivity. J Cell Biol. 2016
691 Feb 1;212(3):281-8. doi: 10.1083/jcb.201507042.
692
693 Paull TT. Mechanisms of ATM Activation. Annu Rev Biochem. 2015;84:711-38. doi:
694 10.1146/annurev-biochem-060614-034335.
695
696 Rappas M, Oliver AW, Pearl LH. Structure and function of the Rad9-binding region of
697 the DNA-damage checkpoint adaptor TopBP1. Nucleic Acids Res. 2011
698 Jan;39(1):313-24. doi: 10.1093/nar/gkq743.
699
700 Reginato G, Cejka P. The MRE11 complex: A versatile toolkit for the repair of broken
701 DNA. DNA Repair (Amst). 2020 Jul-Aug;91-92:102869. doi:
702 10.1016/j.dnarep.2020.102869.
703
704 Reinhardt HC, Yaffe MB. Phospho-Ser/Thr-binding domains: navigating the cell
705 cycle and DNA damage response. Nat Rev Mol Cell Biol. 2013 Sep;14(9):563-80. doi:
706 10.1038/nrm3640.
707
708 Ruis K, Huynh O, Montales K, Barr NA, Michael WM. Delineation of a minimal
709 topoisomerase II binding protein 1 for regulated activation of ATR at DNA double-
710 strand breaks. J Biol Chem. 2022 Jul;298(7):101992. doi:10.1016/j.jbc.2022.101992.
711
712 Sheng ZZ, Zhao YQ, Huang JF. Functional Evolution of BRCT Domains from Binding
713 DNA to Protein. Evol Bioinform Online. 2011;7:87-97. doi: 10.4137/EBO.S7084.
714
715 Shiotani B, Nguyen HD, Håkansson P, Maréchal A, Tse A, Tahara H, Zou L. Two
716 distinct modes of ATR activation orchestrated by Rad17 and Nbs1. Cell Rep. 2013
717 May 30;3(5):1651-62. doi: 10.1016/j.celrep.2013.04.018.
718
719 Syed A, Tainer JA. The MRE11-RAD50-NBS1 Complex Conducts the Orchestration of
720 Damage Signaling and Outcomes to Stress in DNA Replication and Repair. Annu Rev
721 Biochem. 2018 Jun 20;87:263-294. doi: 10.1146/annurev-biochem-062917-012415.
722

723   Taylor RM, Wickstead B, Cronin S, Caldecott KW. Role of a BRCT domain in the
724   interaction of DNA ligase III-alpha with the DNA repair protein XRCC1. Curr Biol.
725   1998 Jul 16;8(15):877-80. doi: 10.1016/s0960-9822(07)00350-8.
726

727   Van Hatten RA, Tutter AV, Holway AH, Khederian AM, Walter JC, Michael WM. The
728   Xenopus Xmus101 protein is required for the recruitment of Cdc45 to origins of
729   DNA replication. J Cell Biol. 2002 Nov 25;159(4):541-7. doi:
730   10.1083/jcb.200207090.
731

732   Wan B, Hang LE, Zhao X. Multi-BRCT scaffolds use distinct strategies to support
733   genome maintenance. Cell Cycle. 2016 Oct;15(19):2561-2570. doi:
734   10.1080/15384101.2016.1218102.
735

736   Wan B, Wu J, Meng X, Lei M, Zhao X. Molecular Basis for Control of Diverse Genome
737   Stability Factors by the Multi-BRCT Scaffold Rtt107. Mol Cell. 2019 Jul 25;75(2):238-
738   251.e5. doi: 10.1016/j.molcel.2019.05.035.
739

740   Wardlaw CP, Carr AM, Oliver AW. TopBP1: A BRCT-scaffold protein functioning in
741   multiple cellular pathways. DNA Repair (Amst). 2014 Oct;22:165-74. doi:
742   10.1016/j.dnarep.2014.06.004.
743

744   Yan S, Lindsay HD, Michael WM. Direct requirement for Xmus101 in ATR-mediated
745   phosphorylation of Claspin bound Chk1 during checkpoint signaling. J Cell Biol.
746   2006 Apr 24;173(2):181-6. doi: 10.1083/jcb.200601076.
747

748   Yoo HY, Kumagai A, Shevchenko A, Shevchenko A, Dunphy WG. The Mre11-Rad50-
749   Nbs1 complex mediates activation of TopBP1 by ATM. Mol Biol Cell. 2009
750   May;20(9):2351-60. doi: 10.1091/mbc.e08-12-1190.

Main Figures and Legends

Figure 1. NBS1 and TOPBP1 interact in frog egg extracts.

A. Schematic representation of the proteins of interest. TOPBP1 contains 9 BRCT

domains while NBS1 contains two BRCT domains, as well as an N-terminal FHA

domain. **B.** A GST pull-down assay was performed, as described in the Methods, with

the indicated GST proteins. This was done in the presence of egg extract, with

optionally added DSBs, and the presence of endogenous TOPBP1 in the bound

fraction was detected by Western blot. The experiment shown is representative of

two independently performed biological replicates. **C.** Same as (B) except GST-NBS1

393-763 was included. The experiment shown is representative of two

762   independently performed biological replicates. **D.** Myc-tagged TOPBP1 BRCT0-2,

763   produced by IVTT, was added to XEE, and DSBs were optionally included, as

764   indicated. The indicated GST proteins were also included, and then the samples

765   were then processed as in (B). The experiment shown is representative of two

766   independently performed biological replicates.

Figure 2. The Bugs-Bunny system recapitulates a phosphate-independent

interaction between NBS1 and TOPBP1.

A. Overview of the Bugs-Bunny protein binding assay system. **B.** The indicated GST

proteins, expressed and purified from *E. coli*, were incubated in IVTT reactions

expressing myc-tagged TOPBP1 BRCT0-2. The samples were then processed for the

GST pull-down assay as described in the Methods. Samples of the total binding

reaction, taken prior to dilution, were also probed for myc, showing that the myc-

tagged proteins were present in equivalent amounts across all the samples (lane

"input"). The experiment shown is representative of two independently performed

biological replicates. We note that the image shown is derived from a larger gel, and

the irrelevant samples were removed. The white dashed line depicts where the gel

was spliced together. The full gel with all original samples is shown in Figure S1A.

C. A GST pull-down assay was performed, as in part A, with the indicated GST- and

myc-tagged proteins. The experiment shown is representative of three

independently performed biological replicates. Quantification of all three

experiments is shown in the graph. Please see Methods for how these experiments

were quantified. **D.** A GST pull-down assay was performed, as in part A, with the

indicated GST proteins and myc-tagged TOPBP1 BRCT0-2. In this experiment

lambda phosphatase was optionally added, as indicated. The experiment shown is

representative of two independently performed biological replicates.

Figure 3. NBS1 independently binds to both BRCT1 and BRCT2 from TOPBP1.

A. GST pull-down assays, using GST-NBS1 1-400 with myc-tagged versions of

TOPBP1's N-terminal BRCT domains, were performed exactly as in Figure 2B. **B.**

GST pull-down assays, using GST-NBS1 1-400 or GST-RAD9Tail together with myc-

tagged versions of TOPBP1's N-terminal BRCT domains, were performed exactly as

in Figure 2B. The experiments shown are representative of two independently

performed biological replicates. **C-D.** Cartoons highlighting the different ways that

RAD9Tail and NBS1 interact with TOPBP1's BRCT0-2 region. See text for details.

Figure 4. NBS1 uses its BRCT1 to bind TOPBP1.

A. Cartoons summarizing the domains located within the N-terminus of NBS1 and

the various truncations used in the binding assays. **B.** MBP pull-down assays were

performed using either MBP alone or MBP fused to TOPBP1's BRCT1&2 domains,

together with myc-tagged and full-length NBS1. The experiment shown is

representative of two independently performed biological replicates. **C.** MBP pull-

down assays were performed using MBP-TOPBP1 BRCT1&2, together with the

indicated myc-tagged forms of NBS1. The experiment shown is representative of

two independently performed biological replicates. **D.** MBP pull-down assays were

performed using either MBP alone or MBP fused to TOPBP1's BRCT1&2 domains,

together with the indicated myc-tagged forms of NBS1. The experiment shown is

representative of two independently performed biological replicates.

Figure 5. Direct interaction between TOPBP1 and NBS1 and the role of PBPs in the interaction.

A. Cartoons showing the recombinant proteins used in the binding assay displayed

in part B. **B.** All the proteins used in this binding assay were expressed and purified

from *E. coli*. Shown is an MBP pull-down assay with either MBP alone or MBP fused

to NBS1 1-200, together with 6His-T7 tagged TOPBP1 BRCT0-2. The binding

reaction was performed in the presence of a 1% BSA solution, to suppress non-

specific binding to the maltose resin. As in Figure 2D, phosphatase was optionally

included (as indicated by the "P"). The experiment shown is representative of two

independently performed biological replicates. **C.** A schematic summarizing the

disparate BRCT domains that are known to bind the TOPBP1 BRCT1&2 region. All of

these domains contain a PBP. **D.** GST pull-down assays, using GST-NBS1 1-400 with

myc-tagged TOPBP1 BRCT0-2, were performed exactly as in Figure 2B. Shown are

both the wild type (WT) and K155K249 to EE ("KK->EE") forms of TOPBP1 BRCT0-

2. The experiment shown is representative of two independently performed

biological replicates. We note that the image shown is derived from a larger gel, and

the irrelevant samples were removed. The white dashed line depicts where the gel

was spliced together. The full gel with all original samples is shown in Figure S1B.

E. An MBP pull-down assay was performed with MBP-TOPBP1 BRCT1&2 together

with either wild type ("WT") or the lysine 158 to alanine ("K158A") forms of myc-

tagged NBS1. 1-400. The experiment shown is representative of two independently

performed biological replicates.

Figure 6. Intact BRCT domains are dispensable for the TOPBP1::NBS1

interaction.

A. Cartoons summarizing the various TOPBP1 BRCT1 truncations used in the

binding assays. These MBP-fusion proteins were produced by IVTT and used in pull-

down assays with recombinant GST-NBS1 1-400. **B-C.** GST pull-down assays, using

GST-NBS1 1-400 together with the indicated MBP-tagged TOPBP1 derivatives. The

experiments shown are representative of two independently performed biological

replicates. **D.** Cartoons summarizing the various NBS1 truncations used in the

binding assays. **E.** MBP pull-down assays using MBP-TOPBP1 BRCT1&2 together

with the indicated myc-tagged NBS1 derivatives. The experiment shown is

representative of two independently performed biological replicates. **F.** MBP pull-

down assays were performed with the indicated proteins. In this experiment, both

the MBP-TOPBP1 derivatives and the myc-NBS1 derivatives were produced by

IVTT. The experiment shown is representative of two independently performed

biological replicates.

A

	β1	α1	β2	β3	α2	β4	α3

TOPBP1 BRCT1 (108-179) DVTISCTSLDKETREDVHHY**VQIMGGCVYRDLNVS-VTHLIAGEVG-SKKY**LVAA-LEKPILLPSWVKELWEKS
TOPBP1 BRCT2 (201-273) GCTICVTGLSSLDRKEVQRLTALHGGEYTGQLKMNESTHLIVQEAK-GQK**YECARKWNVHCISVQWFFDSIEKG**
NBS1 BRCT1 (109-182) EPLVVCSSCLDNSEKNSLNQNLIHLGGHVLNNWTEK-STHLVMT**SIKVTIKTICALICCKPIIKPDYFC**ELLRAI

B

XRCC1 BRCT2

	β1	α1	β2	β3	α2	β4	α3

545-GKHFFLYGEFPGDER**R**KLI**RY**VTA**F**NGELEDYMSDRVQFVITAQEWDPSFEEALMDNPSLAFVRPRWIYSCNEKQ-619

allows MBP to bind DNA ligase III

C

Yellow: RAD9 phospho-peptide; Blue: NBS1 binding region from BRCT2
Green: NBS1 binding region from BRCT1; Magenta: remainder of the BRCT0-2 domain

Figure 7. Summary of the BRCT domain delineation analyses.

A. The amino acid sequences of the indicated domains are displayed. Above the sequences are the positions of either beta sheets or alpha helices, as indicated. Highlighted in red for TOPBP1 are the small stretches of amino acids that confer binding to NBS1 when fused to MBP. Highlighted in red for NBS1 is the region of BRCT1 that binds TOPBP1, as defined by deletion analysis. **B.** The human XRCC1 BRCT2 sequence is shown. Underlined in black is the small region that confers binding to LIG3 when fused to MBP, as shown in Taylor et al, 1998, and highlighted in red are residues shown by Hammel et al, 2021 to be required for binding to LIG3. **C.** Two views of the crystal structure of the human TOPBP1 BRCT0-2 region in complex with a RAD9 phospho-peptide. The images were generated using Pymol structure view software and data set PDB ID: 6HM5.