External Validation and Further Exploration of Fall Prediction Models based on Questionnaires and Daily-life Trunk Accelerometry

Yuge Zhang[1], Roel H.A. Weijer[1,2], Kimberley S. van Schooten[3,4,] Sjoerd M. Bruijn[1,5,6*], and Mirjam Pijnappels[1]

[1] Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
[2] Department of Neurology, Leiden University Medical Center, Leiden, The Netherlands
[3] Neuroscience Research Australia, University of New South Wales, Sydney, Australia
[4] School of Population Health, University of New South Wales, Sydney, Australia
[5] Institute of Brain and Behavior Amsterdam, Amsterdam, The Netherlands
[6] Biomechanics Laboratory, Fujian Medical University, Quanzhou, Fujian, PR China

Corresponding Author:
Sjoerd M. Bruijn
Vrije Universiteit Amsterdam, De Boelelaan 1105, 1081 HV Amsterdam, Netherlands.
Email: s.m.bruijn@vu.nl
Telephone Number: +31 618844777

Abstract

Background: Ambulatory measurements of trunk accelerations can provide valuable insight into the amount and quality of daily life activities. Such information has been used to create models that aim to identify individuals at high risk of falls. However, external validation of such prediction models is lacking, yet crucial for clinical implementation. We externally validated three previously described fall prediction models (van Schooten et al., 2015a).

Methods: Complete questionnaires and one week of trunk acceleration data were obtained in 263 community-dwelling people (mean age 71.8 years, 68.1% female). To validate models, we first used the coefficients and optimal cut-offs from original cohort, then recalibrated the original models, as well as optimized parameters based on our new cohort.

Results: Among all participants, 39.9% experienced falls during 6-month follow-up. All models showed poor precision (0.20-0.49), poor sensitivity (0.32-0.58), and good specificity (0.45-0.89). Calibration of the original models had limited effect on model performance. Using coefficients and cut-offs optimized on the external cohort also had limited benefit. Last, the odds ratios in our cohort were different from those in the original cohort and indicated that gait characteristics, except for index of harmonicity ML, were not statistically significantly associated with falls.

Conclusions: Prediction of fall risk in our cohort was not as effective as in the original cohort. Recalibration as well as optimized model parameters resulted in limited increase in accuracy. Fall prediction models are highly specific to the cohort studied. This highlights the need for large representative cohorts, preferably with an external validation cohort.

Keywords: Wearable devices, Aged, Locomotion, Accidental falls, Verification

1. Introduction

Falls among older adults are a major cause of disability and death. Each year, one in
three older adults aged 65 years or over experience at least one fall (Morrison et al.
2013; Salvà et al. 2004). Therefore, early identification of fall risk is necessary for
timely intervention. Fall risk can be estimated by questionnaires, functional tests in a
clinical setting or in a biomechanical laboratory, but all these methods have limited
predictive value (Hamacher et al. 2011). Specifically, questionnaires have low
reliability due to the subjective bias of the participants or testers (Furnham 1986;
Malatesta et al. 2003; Winstein 1987). Clinical gait tests, such as Berg Balance scale,
Timed Up and Go and POMA, cannot quantify fall risk well, especially in frail older
populations (Barry et al. 2014; Boulgarides et al. 2003; Mancini & Horak 2010). Gait
tests in a biomechanics laboratory do not reflect exposure to environmental and task-
related hazards(Hausdorff et al. 2001; Maki 1997).

Inertial Measurement Units (IMU's) are wearable motion sensors, which can be
used in an everyday environment. Their validity and reliability have been shown in
several studies. For instance, Felius and colleagues studied the test-retest reliability of
107 gait features assessed with IMUs (Felius et al. 2022). They found the relative
minimal detectable change of these features ranged between 0.5 and 1.5 standard
deviation. Several other studies compared IMUs to gold-standard gait-analysis
technologies and suggested IMU-based gait metrics have sufficient accuracy and
reliability to be used in clinical setting(Berner et al. 2020; Betteridge et al. 2021;
Bezold et al. 2021; Zhong & Rau 2020).

Several studies have shown that daily-life gait quality characteristics, assessed
with a single IMU, are able to discriminate fallers from non-fallers. Examples of such
characteristics are local dynamic stability, variability of gait, frequency domain
parameters, and gait smoothness (Bourke et al. 2010; Brodie et al. 2017; Rispens et al.
2015; Weiss et al. 2013). However, only a few research teams focus on developing
fall risk prediction models based on daily-life activities in community environment
(Brodie et al. 2017; Rispens et al. 2015; van Schooten et al. 2016; Weiss et al. 2013).
Among these, three predictive models from van Schooten and colleagues (van
Schooten et al. 2015a) had moderate to good predictive ability for falls happening in
the next 6 months, and demonstrated that daily life accelerometry contributed
substantially to the identification of individuals at risk of falls (areas under curve
(AUC) of 0.68, 0.71 and 0.82 respectively). These models were logistic regression
models based on a relatively large (n=169) data set of the FARAO cohort, a project
concerning fall-risk assessment in older adults, with independent variables obtained
either from questionnaires, from daily-life accelerometry, or from a combination of
both.

However, to date, none of the fall-risk prediction models have been externally
validated, so the repeatability and generalizability cannot be guaranteed (Nouredanesh
et al. 2021), which hinders the implementation of a fall prediction model for
community-dwelling older adults. Shany and colleagues (Shany et al. 2015) also
pointed out that current studies of sensor-based fall risk estimation were overly
optimistic due to small sample sizes, flawed modelling and validation methods, (e.g.
inherent problems with overly cross-validation techniques, lack of external validation).
In this study, we aimed to assess the external validity of the FARAO fall prediction
models (van Schooten et al. 2015a) by using an external cohort of community-
dwelling older adults. Moreover, as the ambulatory assessments specifically quantify
the amount and quality of daily-life gait, we performed additional analyses that
focused on walk-falls only. In addition, we calibrated the FARAO prediction models.
Finally, we evaluated models based on the optimal model parameters of the new
cohort and compared the odds ratio for individual variables between original and new
cohorts.
54

2. Methods

2.1 Participants

The data in this study were available from the "Veilig in beweging blijven" (VIBE),
which translates to "Safely remaining active" cohort. Participants were included if 1)
they were 65 years of age or older, but not exceed 90 years 2) they had a Mini-Mental
State Examination (MMSE) $\geq$ 19 (Folstein et al. 1975) , 3) they were able to walk at
least 20 meters, if necessary with a walking aid, without dizziness or perceiving chest
pain or pressure, and 4) they were able to understand the Dutch language. The
protocol was approved by the ethics committee of the faculty of Behavioural and
Movement Sciences of the Vrije Universiteit Amsterdam (VCWE-2016-129). Among
441 persons who expressed interest, 283 were eligible to participate and agreed to a
12-month follow-up on falls. All participants provided signed informed consent
before participation.
68

2.2 Questionnaire related to fall risk

Descriptive characteristics such as age, weight, height, the use of a walking aid and 6-
month fall history were obtained from all participants. In addition, participants
independently filled out validated questionnaires and tests on fall-risk factors, i.e.
cognitive function (Mini Mental State Examination; MMSE), concern about falling
(16-item fall efficacy scale; FES-I (Yardley et al. 2005)), LASA fall-risk profile
(Longitudinal Aging Study Amsterdam) (Tromp et al. 2001) and Geriatric Depression

Scale (GDS-15 (Yesavage et al. 1982)). The LASA fall-risk profile comprised
questions concerning dizziness, independence in daily life, having pets, alcohol
consumption, education duration and required the measurement of hand grip strength,
which was quantified using a handgrip dynamometer (TKK 5401, Takei Scientific
Instruments, Tokyo, Japan). The GDS-15 is an effective scale for depressive
symptoms in older populations, where 5 negative items and 10 positive items indicate
symptoms of depression (Greenberg 2012). Missing ≤20% of items in GDS-15 forms
were imputed (Imai et al. 2014). Scores of 0-4 are considered normal, depending on
age, education, and complaints; 5-8 indicate mild depression; 9-11 indicate moderate
depression; and 12-15 indicate severe depression. Since the GDS-30 was used in the
original model, we had to convert our GDS-15 scores to GDS-30 scores, according to
a regression equation, GDS-30 scores = 1.95*GDS-15 scores + 1.57 (Zhang et al.
2023).
89

2.3 Assessment and analyses of daily-life gait characteristics

Conforming to the FARAO cohort study (van Schooten et al. 2015a), participants
were instructed to wear a tri-axial accelerometer (DynaPort MoveMonitor, McRoberts,
The Hague, the Netherlands) at home for 7 consecutive days, except during aquatic
activities such as showering. The sensor was worn on the trunk at the level of L5
using an elastic belt, whose three axes of orientation corresponded to vertical (VT),
anterior-posterior (AP) and medial-lateral (ML), ranged from −6g to +6g, with
sampling frequency set to 100 Hz. We omitted the first 6 hours of the measurements
from analysis, to discard any possible artefacts caused by transportation from the
research team to the participants' home. To have a valid measure of the amount of
daily activity, the accelerometer had to be worn for more than twelve hours each day
(van Schooten et al. 2015b). At the same time, the minimum valid days for daily
physical activities are 2 days except for lying for which it is 3 days (van Schooten et
al. 2015b), so only subjects who had accelerometer data with worn time ≥ 2 days
were used. The identification of non-wearing, locomotion, sitting, lying, shuffling,
and standing, were obtained from the manufacturer's validated algorithms (Dijkstra et
al. 2010). According to these episodes of daily activities, the duration of lying,
walking, and the number of strides per day were calculated as outcome measures.
The data analyses were performed in Matlab R2019a (Mathworks, Natwick, MA),
using the Gait Analysis toolbox (Human Movement Sciences 2022). Prior to
estimation of gait characteristics, data from each locomotion episode was realigned
with the vertical axis, which was estimated from the direction of the mean
acceleration signal (Moe-Nilssen 1998), and anteroposterior and mediolateral
directions, which were estimated by optimizing the left–right symmetry (Rispens et al.

2014). Locomotion episodes were then split into epochs of 10 seconds to avoid
possible sample size-related bias in outcome measures (Bruijn et al. 2009). Median
gait characteristics were calculated over epochs of each gait episode, identical to the
data length requirements of characteristics determined, i.e., logarithmic divergence
rate AP (Wolf et al. 1985), range VT, index of harmonicity ML (Lamoth et al. 2002)
and sample entropy VT (Richman & Moorman 2000). The mean over all episodes in
a week for these measures were calculated as outcome measure (van Schooten et al.
2016) . Detailed description and formulas of sensor-based gait measures can be seen
in the supplementary materials of (Felius et al. 2022).
123
124 2.4 Outcome: Falls and walk-falls
Although participants were monitored prospectively for falls over a 12-month period
following the baseline assessment by means of a fall diary and monthly telephone
contact, we only focused on the first 6 months of follow-up for consistency with the
original models. However, to explore if extended follow-up periods resulted in better
model performance, we also analyzed data obtained from the 12-month fall history
and 12-month prospective fall recordings; these results are reported in the
supplementary materials (see *Table S1-2* and *Fig. S1-2)*. A fall was defined as an
unintentional change in position resulting in coming to rest at a lower level or on the
ground (Gibson 1987). Participants were classified as being a faller based on the
number of prospectively monitored falls (≥1) they experienced over the 6-month
period. Participants who did not record falls in their fall diary for the complete 6
months but experienced at least one fall within the recorded follow-up period were
also classified as fallers.
When a fall happened, the activities and circumstances were also recorded,
including walking, climbing the stair, riding a bicycle, showering, etc. Because the
models were based on gait parameters, we also investigated how all models
performed for people that fell specifically during walking. Participants were classified
as walk-fallers if they had one record of a fall during walking.
143
144 2.5 Statistical analysis
Statistical analysis was performed in Matlab R2021a (MathWorks, Natick, MA) and
R (R core team, Vienne). Unpaired t-tests were used to compare outcomes between
fallers and non-fallers, and between walk-fallers and non-fallers, separately. The
FARAO predictive models were all originally based on stepwise forward logistic
regression analyses, with all continuous parameters transformed to z-scores as input.
So, we did the same when we did our logistic regression. Model 1 only included 6-
month history of falls and GDS-30, model 2 contained accelerometry-based daily life
gait quantity and quality characteristics, and model 3 combined variables from both

model 1 and model 2. Model performance was evaluated by area under the ROC
curve (AUC), accuracy, precision, sensitivity, and specificity. Usually, AUC=0.5-0.7
means poor discriminative ability, 0.7-0.8 means moderate discriminative ability, 0.8-
0.9 means good discriminative ability and 0.9-1.0 means excellent discriminative
ability (Hosmer Jr et al. 2013).

To validate the FARAO predictive models on our VIBE cohort data, we first
calculated model performance for our VIBE cohort when we used the mean/SDs of
the FARAO cohort to calculate the z-scores for our data, and used the FARAO model
coefficients, original cut-offs (0.279, 0.352 and 0.396). Either all falls or walking falls
were used as outcome.

Next, these models were recalibrated (Van Calster et al. 2019) for possible model
improvement. To this aim, the agreement between the predicted probability of the
models and observed binary outcomes was evaluated using a logistic regression, in
which the predicted probability was the independent variable and falling status the
dependent variable, see equation 1

Logit (Actual Values) = $\beta_0 + \beta_1$ * (predicted probability)   [1]
Where the slope (β_1) indicates the level of overfitting (<1) or underfitting (>1) and
the intercept (β_0) suggests an general overestimation (<0) or underestimation (>0)
(Van Calster et al. 2016). A slope of 1 and an intercept of 0 imply perfect model
calibration (Van Calster et al. 2019). If these recalibrated models perform well, it
suggests a simple scaling of the model would be enough to use the model in other
populations.

Next, to see whether performance could be improved, considering potential
differences in the sample measured, we fitted optimal logistic regression models for
our VIBE dataset given the input variables as specified by the three FARAO models.

Finally, we performed unpaired t-tests on the means and SD of outcome measures
of the VIBE data as compared to the published FARAO data, to see if potential
differences in populations could explain performance. Lastly, univariate logistic
regression was performed to identify the association between model variables and
prospective falls, and these resulting odds ratio were compared with those from the
FARAO cohort (van Schooten et al. 2015a).

3. Results

3.1 Descriptive Characteristics

283 participants filled in questionnaires and wore an accelerometer for one week at
baseline, but 5 participants' acceleration data could not be read. Among the remaining
278 participants, 15 participants were excluded due to missing more than 20% of the
GDS items (n=3, 1 person missed 5 items and 2 persons missed 15 items), incomplete

records within 6 months and no record of falls (n=3), accelerometer assessment
duration of less than 2 days (n=3), ≤ 50 episodes of overall measurement time (n=6).
Figure 1 presents a flowchart of the included fallers and non-fallers.
Descriptive characteristics of the participants are summarized in table 1. Of the
263 participants, 39.9% (n=105) experienced one or more falls during follow-up, 19%
(n=50) had fallen during walking. In addition, 37.6% of all participants, 43.8% of all
fallers, 50% of walk-fallers and 33.5% of non-fallers had a fall history in the past 6
months. The average number of prospective and past falls, GDS scores, FES-I and
LASA fall risk profile (score) were significantly different between fallers and non-
fallers, and between walk-fallers and non-fallers. Furthermore, walk-fallers had
significantly lower hand grip strength, and poorer independence in life, compared to
non-fallers.
Figure 2(A) shows the main activities when a fall happened, including walking
(39%), riding a bicycle (13%), movement transfer (12%), going up or down the stairs
(12%), doing sports (6%), standing, and sitting (in static positions) (4%), and other
activities (15%). Among these, hiking was classified as a sport in this study. Working
in the garden, getting dressed, etc. were classified as "Other". Furthermore, main
causes of falls during walking were shown in figure 2(B), which were trip/stumble
(34%), balance loss (31%), slip (17%), physiological discomfort (8%), being
distracted (2%) and other (8%). Trip/stumble often happened on uneven roads or
roads with obstacles, and balance loss was more likely to occur when there was
sudden disturbance, opposite walking height to the expected. Falling due to slips
usually occurred in the bathroom or on a slippery floor. In addition, physiological
discomfort included ankle sprain, sudden dizziness, drops in blood pressure, eye
injury and weakened muscle strength, etc. Falling caused by distraction included
chatting, looking at map, or doing other tasks when walking. Lastly, some unknown,
poorly classified reasons were divided into "Other".
218
3.2 Validation of predicting all falls and walk-falls based on
existing models
By using optimal cut-offs (0.279, 0.352 and 0.396), coefficients and mean (SD) (for
z-score) from the original cohort, we obtained model performance of the three
prediction models based on only questionnaires, only accelerometry, and combined
questionnaires and accelerometry. Table 2a shows the results of each prediction
model, with either all falls or walk-falls as outcome (Model performance for 12-
months follow-up on FARAO cohort can be seen in *Table S2a*). Model 1, with only
questionnaires, performed best for predicting all falls and walk-falls, with AUC (area
under the curve) of 0.62 and 0.68, respectively. AUC of model 2 in all-fall and walk-
fall groups was 0.50 and 0.49, respectively, and 0.53 and 0.59 in model 3. At the same

time, the precision and sensitivity in both groups were low, which meant the ability of
predicting a fall was low for all models.
232
3.3 Model recalibration
In our VIBE cohort, the actual fall incidence and walk-fall incidence in 6 months was
0.40 and 0.19. The recalibrated fall rate was the same as actual fall rate, but the
recalibrated walk-fall rate was 0.24 (see *Table S3*). Calibration intercepts of model 1
and model 3 in the group with all-fallers were significantly negative (-1.06 and -0.83,
respectively), as well as intercepts of all models with walk-fallers (-2.17, -1.14 and -
1.82, respectively). The slopes of model 1 and model 3 were significantly >1 in both
groups (2.22 and 1.40, 3.24 and 2.08, respectively), indicating underfitting. The
performance of the calibrated models is shown in table 2b. The AUC of all calibrated
models ranged from 0.51-0.68.
243
3.4 Validation of predicting all falls and walk-falls based on the external
cohort
Model performances did not improve as compared to the original FARAO model
when the models were fitted again (with the same variables as in the original FARAO
model), shown in table 2c (see ROC curves in *Fig. S3*) (12-months follow-up model
performance and ROC curves based on VIBE cohort can be seen in *Table S2b* and *Fig.*
S2, respectively). Specificity and sensitivity were still poor for these models (ranging
from 0.59-0.66).
252
3.5 Comparison of cohorts and individual odds ratios
To better understand differences in model performance between the original models
and models fitted on our external dataset, the comparison of descriptive statistics and
odds ratio of measures for the our external VIBE cohort and original FARAO cohort
(van Schooten et al. 2015a) are presented in table 3. There were small but significant
differences in age, and height, with the VIBE cohort being slightly younger and
shorter. Prevalence of fall history prior to 6 months in VIBE cohort was 37.6%, which
was slightly higher than 35.5% in the FARAO cohort. Cognitive function (MMSE
score) of VIBE cohort was 28.2, slightly higher than 27.7 in FARAO cohort. LASA
fall-risk profile score, percent of inability to use public transportation and use of
walking aid in VIBE cohort were lower than those in FARAO cohort. In agreement
with this, the participants of the VIBE cohort had a higher degree of living
independence. Number of strides walked, and walking duration were significantly
higher in the VIBE cohort than in the FARAO cohort. However, subjects in the VIBE
cohort walked with lower walking speed (0.6 m/s for VIBE, vs 0.9 m/s for FARAO).

The duration of lying was 0.9 hours less than FARAO cohort. All gait characteristics
differed significantly between the VIBE and FARAO cohorts.
For both the VIBE and the FARO cohorts, significant odds ratios for
questionnaire metrics 6-months history of falls, GDS-30, FES-I, and LASA profile
score were found. For the VIBE cohort, the only gait measure that was significantly
associated with falls was the index of harmonicity ML. However, the confidence
interval of the odds ratios for the other gait parameters overlapped with those in
FARAO cohort, except for the logarithmic divergence rate AP.
276

4. Discussion

We validated the predictive ability of three fall prediction models based on
questionnaires, accelerometry of daily life gait, and both, on the data from the VIBE
cohort of community-dwelling older adults. We found that the external validity of the
models developed based on the FARAO cohort was limited for predicting all falls as
well as when focusing on walk-falls. Subsequently, we explored why this may have
been so.
When comparing the population characteristics between the two cohorts, we
found that participants in the VIBE cohort were slightly younger, more often female,
a little shorter, had a higher level of cognition, had less fall risk, lived more
independently, use public transport more often and used less walking aids.
Participant's in both studies had a normal (i.e., low) degree of depressive symptoms
(Greenberg 2012). Both cohorts had the same definition of a faller in past that the
number of falls in follow-up 6 months was ≥1. The prevalence of falls in our study
was higher than in the original study (39.9% and 34.9%, respectively), with 14.5%
participants having two or more falls in the follow-up 6 months compared to 16.5% in
the FARAO cohort. Studies suggested that only one fall may be an ambiguous fall,
with no apparent cause (Nouredanesh et al. 2021), and people who were 'once only
fallers' were characteristically more closely related to non-fallers than 'recurrent
fallers'(Lord et al. 1994) or injury-prone fallers. Hence, the fall rate of the FARAO
cohort was lower than VIBE cohort if based on one fall, but higher than the VIBE
cohort if based on multiple falls.
The definition and record methods of falls were the same in both cohorts,
including fall calendars and monthly phone calls. However, this does not exclude that
there was a slight difference in interpreting the fall during the phone calls that may
have resulted in more reported falls on the VIBE cohort. Causes of falls were divided
into 7 categories and 39% of falls occurred during walking, similar to other studies
(Niino et al. 2000; Peel 2011; Talbot et al. 2005). However, it was not known if falls
in the FARAO cohort had a similar distribution of causes. Hence the different

proportion of falls during walking may have affected the results, but the models also
did not work well for the walking fallers. In post-hoc analysis, results did not change
based on the 12-month follow-up data available from the VIBE cohort (refer to *Table*
S2). In conclusion, there may have been differences in the classification of fallers that
influenced the accuracy of the models. However, since such differences are most
likely due to interpretation of the fall definition by the people who performed the
monthly phone calls, we are unable to identify such potential differences to test if
these affected the results.
Limited model performance in both groups was obtained after recalibration of the
model from the FARAO cohort. Calibrated slopes were significantly larger than 1 and
intercepts were significantly negative. This indicates underfitting and overestimation
of original models. Some studies recommend use dataset with more than 100 events
and 100 non-events to prevent the potential underfitting of models (Van Calster et al.
2016), and it is clear that the original FARAO cohort, with 169 subjects, does not
fulfill this criterium.
When we explored the optimal coefficients and optimal cut-offs for the VIBE
cohort, we found that model 3 performed best for both all falls and walk-falls. The
AUC of all models improved a little, but the precision and sensitivity remained
moderate. More importantly, the model which combined the gait quality
characteristics and questionnaires was better than either one alone – reinforcing the
conclusion of van Schooten et al.'s study that daily-life accelerometry has value for
the identification of people at risk for falls. Thus, while the same variable may yield a
reasonable fall prediction model, it appears that coefficients and cut-offs may be
rather specific to a given population sample (Punt et al. 2016).
To further study the predictors in the models, odds ratios of individual variables
were compared with the initial FARAO cohort. Studies have shown that fall history,
depressive symptoms, LASA fall-risk profile, fear of falling, and poor executive
function (as assessed by TMT-A) were associated with prospective falls (Holtzer et al.
2007; Pluijm et al. 2006; Tromp et al. 2001). Significant odds ratios of these measures
in the original study confirmed this (van Schooten et al. 2015a). However, of the gait
quality characteristics, only index of harmonicity ML had significant odds ratios in
VIBE cohort, while other gait quality measures have been shown to be related to
future falls in FARAO cohort. Specifically, lower harmonic ratio, reflecting lower
gait symmetry, was shown to be related to falls by (Doi et al. 2013); higher
logarithmic rate of divergence, reflecting lower local stability, was associated with
future falls, according to previous findings (Lockhart & Liu 2008; Rispens et al. 2015;
Toebes et al. 2012). This was most likely due to the differences in the VIBE cohort,
who were younger, and more active than participants in the FARAO cohort. Still,
these findings suggest that even though we could not successfully use the FARAO fall

prediction models for an external data set, there may be merit in the use of daily-life
IMU data.
Our study has some limitations. First, we included elderly subjects who did or not
use walking aids, but it is uncertain whether the use of a walking aid influenced the
extracted gait characteristics, and their relationship with fall risks, which is also a
limitation in the original FARAO models. Second, we were unable to pinpoint the
source of some large differences in gait parameters between our cohort. For instance,
we found a complete lack of overlap for the index of harmonicity ML in the mean
values and its standard deviation (0.4(0.2) and 2.0 (0.3)). However, this also
highlights the inherent difficulty with creating widely applicable fall-prediction
models; even though our inclusion criteria were largely similar, and we used similar
recruitment procedures, our results were substantially different from previous studies.
Perhaps, even larger studies (or pooling of datasets) can prevent these shortcomings.
358

5. Conclusion

In conclusion, prediction of fall risk in our VIBE cohort based on the FARAO models
was not as effective as models in FARAO cohort (van Schooten et al. 2015a).
Moreover, recalibration of the models suggested that the original models are
underfitting and have limited external validity. Despite several earlier reports
indicating the importance of daily life gait measures for predicting falls, such
measures alone were of limited value in our cohort. Moreover, the fact that optimal
coefficients and cut-off were different for our cohort, as well as the fact that we found
almost no significant ORs for the gait parameters included in the model, suggest that
this kind of fall prediction models are highly specific to the cohort studied.
Nonetheless, even with these optimal coefficients and cut-offs, model performance
was still poor, which highlights the need for large representative cohorts, preferably
with an external validation cohort.
372

Acknowledgments

Author contributions: Conceptualization, MP and SMB; methodology, RW, SMB,
KS and MP; formal analysis, YZ, SMB, MP, RW and KS; resources, MP and SMB;
data collection, RW; data processing, YZ; writing-review and editing, YZ, RW, SMB,
KS and MP; visualization, YZ; supervision, SMB and MP; project administration, MP.
All authors have read and agreed to the published version of the manuscript.
Funding: This work was supported by China Scholarship Council (CSC)
(202009110145) to YZ; Dutch Organization for Scientific Research (NWO) (no.

91714344, 016.Vidi.178.014) to MP, RW and SMB; and Human Frontier Science

Program Fellowship (LT001080/2017) to KS.

Conflicts of Interest: The authors declare no conflict of interest.

References

Barry E, Galvin R, Keogh C, Horgan F, and Fahey T. 2014. Is the Timed Up and Go test a useful
predictor of risk of falls in community dwelling older adults: a systematic review and meta-
analysis. *BMC Geriatrics* 14:14. 10.1186/1471-2318-14-14

Berner K, Cockcroft J, Morris LD, and Louw Q. 2020. Concurrent validity and within-session
reliability of gait kinematics measured using an inertial motion capture system with repeated
calibration. *Journal of Bodywork and Movement Therapies* 24:251-260.
https://doi.org/10.1016/j.jbmt.2020.06.008

Betteridge CMW, Natarajan P, Fonseka RD, Ho D, Mobbs R, and Choy WJ. 2021. Objective falls-risk
prediction using wearable technologies amongst patients with and without neurogenic gait
alterations: a narrative review of clinical feasibility. *mHealth* 7.

Bezold J, Krell-Roesch J, Eckert T, Jekauc D, and Woll A. 2021. Sensor-based fall risk assessment in
older adults with or without cognitive impairment: a systematic review. *Eur Rev Aging Phys*
Act 18:15. 10.1186/s11556-021-00266-w

Boulgarides LK, McGinty SM, Willett JA, and Barnes CW. 2003. Use of clinical and impairment-
based tests to predict falls by community-dwelling older adults. *Physical Therapy* 83:328-339.
10.1093/ptj/83.4.328

Bourke AK, Van De Ven P, Gamble M, O'Connor R, Murphy K, Bogan E, McQuade E, Finucane P,
ÓLaighin G, and Nelson J. 2010. Assessment of waist-worn tri-axial accelerometer based fall-
detection algorithms using continuous unsupervised activities. 2010 Annual International
Conference of the IEEE Engineering in Medicine and Biology: IEEE. p 2782-2785.

Brodie MA, Coppens MJ, Ejupi A, Gschwind YJ, Annegarn J, Schoene D, Wieching R, Lord SR, and
Delbaere K. 2017. Comparison between clinical gait and daily-life gait assessments of fall risk
in older people. *Geriatrics & Gerontology International* 17:2274-2282. 10.1111/ggi.12979

Bruijn SM, van Dieën JH, Meijer OG, and Beek PJ. 2009. Statistical precision and sensitivity of
measures of dynamic gait stability. *Journal of Neuroscience Methods* 178:327-333.
10.1016/j.jneumeth.2008.12.015

Dijkstra B, Kamsma Y, and Zijlstra W. 2010. Detection of gait and postures using a miniaturised
triaxial accelerometer-based system: accuracy in community-dwelling older adults. *Age and*
Ageing 39:259-262. 10.1016/j.apmr.2010.05.004

Doi T, Hirata S, Ono R, Tsutsumimoto K, Misu S, and Ando H. 2013. The harmonic ratio of trunk
acceleration predicts falling among older people: results of a 1-year prospective study. *Journal*
of neuroengineering and rehabilitation 10:7. 10.1186/1743-0003-10-7

Felius RAW, Geerars M, Bruijn SM, van Dieën JH, Wouda NC, and Punt M. 2022. Reliability of IMU-
Based Gait Assessment in Clinical Stroke Rehabilitation. *Sensors (Basel)* 22.
10.3390/s22030908

Folstein MF, Folstein SE, and McHugh PR. 1975. "Mini-mental state": a practical method for grading
the cognitive state of patients for the clinician. *Journal of Psychiatric Research* 12:189-198.

10.1016/0022-3956(75)90026-6

Furnham A. 1986. Response bias, social desirability and dissimulation. *Personality and individual*
differences 7:385-400. 10.1016/0191-8869(86)90014-0

Gibson MJ. 1987. The Prevention of Falls in Later Life : A Report of the Kellogg International Work
Group on the Prevention of Falls by the Elderly. *Dan Med Bull* 34:1-24.

Greenberg SA. 2012. The geriatric depression scale (GDS). *Best Practices in Nursing Care to Older*
Adults 4:1-2.

Hamacher D, Singh NB, Van Dieën JH, Heller MO, and Taylor WR. 2011. Kinematic measures for
assessing gait stability in elderly individuals: a systematic review. *Journal of The Royal*
Society Interface 8:1682-1698. 10.1098/rsif.2011.0416

Hausdorff JM, Rios DA, and Edelberg HK. 2001. Gait variability and fall risk in community-living
older adults: A 1-year prospective study. *Archives of physical medicine and rehabilitation*
82:1050-1056. 10.1053/apmr.2001.24893

Holtzer R, Friedman R, Lipton RB, Katz M, Xue X, and Verghese J. 2007. The relationship between
specific cognitive functions and falls in aging. *Neuropsychology* 21:540. 10.1037/0894-
4105.21.5.540

Hosmer Jr DW, Lemeshow S, and Sturdivant RX. 2013. *Applied logistic regression*: John Wiley &
Sons.

Human Movement Sciences VUA. 2022. VU-HMS/Gait-Analysis: v3.4.1.1. *Available at*
https://doi.org/10.5281/zenodo.7362138 (accessed nov 2022).

Imai H, Furukawa TA, Kasahara Y, Ishimoto Y, Kimura Y, Fukutomi E, Chen W-l, Tanaka M,
Sakamoto R, Wada T, Fujisawa M, Okumiya K, and Matsubayashi K. 2014. Ipsative
imputation for a 15-item Geriatric Depression Scale in community-dwelling elderly people.
Psychogeriatrics 14:182-187. 10.1111/psyg.12060

Lamoth CJC, Beek PJ, and Meijer OG. 2002. Pelvis–thorax coordination in the transverse plane during
gait. *Gait & Posture* 16:101-114. 10.1016/S0966-6362(01)00146-1

Lockhart TE, and Liu J. 2008. Differentiating fall-prone and healthy adults using local dynamic
stability. *Ergonomics* 51:1860-1872. 10.1080/00140130802567079

Lord SR, Ward JA, Williams P, and Anstey KJ. 1994. Physiological factors associated with falls in
older community-dwelling women. *Journal of the American Geriatrics Society* 42:1110-1117.
10.1111/j.1532-5415.1994.tb06218.x

Maki BE. 1997. Gait changes in older adults: predictors of falls or indicators of fear? *Journal of the*
American Geriatrics Society 45:313-320.

Malatesta D, Simar D, Dauvilliers Y, Candau R, Borrani F, Préfaut C, and Caillaud C. 2003. Energy
cost of walking and gait instability in healthy 65-and 80-yr-olds. *Journal of applied*
physiology 95:2248-2256. 10.1152/japplphysiol.01106.2002

Mancini M, and Horak FB. 2010. The relevance of clinical balance assessment tools to differentiate
balance deficits. *European journal of physical and rehabilitation medicine* 46:239-248.

Moe-Nilssen R. 1998. A new method for evaluating motor control in gait under real-life environmental
conditions. Part 1: The instrument. *Clin Biomech (Bristol, Avon)* 13:320-327. 10.1016/s0268-
0033(98)00089-8

Morrison A, Fan T, Sen SS, and Weisenfluh L. 2013. Epidemiology of falls and osteoporotic fractures:
a systematic review. *ClinicoEconomics and outcomes research : CEOR* 5:9-18.
10.2147/CEOR.S38721

Niino N, Tsuzuku S, Ando F, and Shimokata H. 2000. Frequencies and circumstances of falls in the
National Institute for Longevity Sciences, Longitudinal Study of Aging (NILS-LSA). *Journal*
of Epidemiology 10:90-94. 10.2188/jea.10.1sup_90

Nouredanesh M, Godfrey A, Howcroft J, Lemaire ED, and Tung J. 2021. Fall risk assessment in the
wild: A critical examination of wearable sensor use in free-living conditions. *Gait & Posture*
85:178-190. 10.1016/j.gaitpost.2020.04.010

Peel NM. 2011. Epidemiology of falls in older age. *Canadian Journal on Aging/La Revue canadienne*
du vieillissement 30:7-19. 10.1017/S071498081000070X

Pluijm SMF, Smit JH, Tromp EAM, Stel VS, Deeg DJH, Bouter LM, and Lips P. 2006. A risk profile
for identifying community-dwelling elderly with a high risk of recurrent falling: results of a 3-
year prospective study. *Osteoporosis international* 17:417-425. 10.1007/s00198-005-0002-0

Punt M, Bruijn SM, van Schooten KS, Pijnappels M, van de Port IG, Wittink H, and van Dieën JH.
2016. Characteristics of daily life gait in fall and non fall-prone stroke survivors and controls.
Journal of neuroengineering and rehabilitation 13:1-7. 10.1186/s12984-016-0176-z

Richman JS, and Moorman JR. 2000. Physiological time-series analysis using approximate entropy and
sample entropy. *American Journal of Physiology-Heart and Circulatory Physiology*
278:H2039-H2049. 10.1152/ajpheart.2000.278.6.H2039

Rispens SM, Pijnappels M, van Schooten KS, Beek PJ, Daffertshofer A, and van Dieën JH. 2014.
Consistency of gait characteristics as determined from acceleration data collected at different
trunk locations. *Gait & Posture* 40:187-192. 10.1016/j.gaitpost.2014.03.182

Rispens SM, van Schooten KS, Pijnappels M, Daffertshofer A, Beek PJ, and van Dieën JH. 2015.
Identification of fall risk predictors in daily life measurements:Gait characteristics' reliability
and association with self-reported fall history. *Neurorehabilitation and neural repair* 29:54-61.
10.1177/1545968314532031

Salvà A, Bolíbar I, Pera G, and Arias C. 2004. Incidence and consequences of falls among elderly
people living in the community. *Medicina Clínica* 122:172-176. 10.1016/S0025-
7753(04)74184-6

Shany T, Wang K, Liu Y, Lovell NH, and Redmond SJ. 2015. Review: Are we stumbling in our quest to
find the best predictor? Over-optimism in sensor-based models for predicting falls in older
adults. *Healthcare Technology Letters* 2:79-88. 10.1049/htl.2015.0019

Talbot LA, Musiol RJ, Witham EK, and Metter EJ. 2005. Falls in young, middle-aged and older
community dwelling adults: perceived cause, environmental factors and injury. *BMC public*
health 5:1-9. 10.1186/1471-2458-5-86

Toebes MJP, Hoozemans MJM, Furrer R, Dekker J, and van Dieën JH. 2012. Local dynamic stability
and variability of gait are associated with fall history in elderly subjects. *Gait & Posture*
36:527-531. 10.1016/j.gaitpost.2012.05.016

Tromp AM, Pluijm SMF, Smit JH, Deeg DJH, Bouter LM, and Lips P. 2001. Fall-risk screening test: A
prospective study on predictors for falls in community-dwelling elderly. *Journal of Clinical*
Epidemiology 54:837-844. 10.1016/S0895-4356(01)00349-3

Van Calster B, McLernon DJ, Van Smeden M, Wynants L, and Steyerberg EW. 2019. Calibration: the
Achilles heel of predictive analytics. *BMC medicine* 17:1-7. 10.1186/s12916-019-1466-7

Van Calster B, Nieboer D, Vergouwe Y, De Cock B, Pencina MJ, and Steyerberg EW. 2016. A
calibration hierarchy for risk models was defined: from utopia to empirical data. *Journal of*
Clinical Epidemiology 74:167-176. https://doi.org/10.1016/j.jclinepi.2015.12.005

van Schooten KS, Pijnappels M, Rispens SM, Elders PJM, Lips P, Daffertshofer A, Beek PJ, and van
Dieën JH. 2016. Daily-ilfe gait quality as predictor of falls in older people: a 1-year
prospective cohort Study. *PLOS ONE* 11:e0158623. 10.1371/journal.pone.0158623

van Schooten KS, Pijnappels M, Rispens SM, Elders PJM, Lips P, and van Dieën JH. 2015a.
Ambulatory fall-risk assessment: Amount and quality of daily-life gait predict falls in older
adults. *The Journals of Gerontology: Series A* 70:608-615. 10.1093/gerona/glu225

van Schooten KS, Rispens SM, Elders PJM, Lips P, van Dieën JH, and Pijnappels M. 2015b. Assessing
physical activity in older adults: Required days of trunk accelerometer measurements for
reliable estimation. *Journal of Aging and Physical Activity* 23:9-17. 10.1123/japa.2013-0103

Weiss A, Brozgol M, Dorfman M, Herman T, Shema S, Giladi N, and Hausdorff JM. 2013. Does the
evaluation of gait quality during daily life provide insight into fall risk? A novel approach
using 3-day accelerometer recordings. *Neurorehabil Neural Repair* 27:742-752.
10.1177/1545968313491004

Winstein N. 1987. Unrealistic optimism about susceptibility to health problems: Conclusions from a
community-wide simple. *J Behav Med* 10:481-500. 10.1007/BF00846146

Wolf A, Swift JB, Swinney HL, and Vastano JA. 1985. Determining Lyapunov exponents from a time
series. *Physica D: Nonlinear Phenomena* 16:285-317. 10.1016/0167-2789(85)90011-9

Yardley L, Beyer N, Hauer K, Kempen G, Piot-Ziegler C, and Todd C. 2005. Development and initial
validation of the Falls Efficacy Scale-International (FES-I). *Age and ageing* 34:614-619.
10.1093/ageing/afi196

Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, and Leirer VO. 1982. Development and
validation of a geriatric depression screening scale: A preliminary report. *Journal of
Psychiatric Research* 17:37-49. 10.1016/0022-3956(82)90033-4

Zhang Y, Hoozemans M, Pijnappels M, and Bruijn SM. 2023. A formula for calculating 30-item
Geriatric Depression Scale (GDS-30) scores from the 15-item version (GDS-15).
Experimental Gerontology 172:112077. https://doi.org/10.1016/j.exger.2022.112077

Zhong R, and Rau PP. 2020. Are cost-effective technologies feasible to measure gait in older adults? A
systematic review of evidence-based literature. *Arch Gerontol Geriatr* 87:103970.
10.1016/j.archger.2019.103970

551 **Table 1.** Descriptive characteristics (Mean (SD)) of the VIBE participants

Characteristics	All participants	All fallers	Walk-fallers	Non-fallers
Number	263	105	50	158
Age (years)	71.8 (5.7)	71.9 (5.4)	71.9 (6.2)	71.8 (5.9)
Gender (%female)	68.1 (179)	68.6 (72)	72 (36)	67.7 (107)
Weight (kg)	74.1 (13.5)	72.6 (12.9)	70.6 (12.4)	75.1 (13.8)
Height (cm)	169.1 (8.5)	168.6 (9.1)	166.4 (8.2)	169.5 (8.1)
Number of falls in follow-up 6 months	0.6 (1.02)	1.6 (1.04) [A]	2.0 (1.3) [B]	0 [A, B]
At least one fall in follow-up 6 months (%/n)	39.9 (105)	100 (105)	100 (50)	0
Two or more falls in follow-up 6 months (%/n)	14.5 (38)	36.2 (38)	56 (28)	0
Number of falls in past 6 months	0.7 (1.4)	1 (1.9) [A]	1.4 (2.5) [B]	0.5 (0.8) [A, B]
At least one fall in past 6 months (%/n)	37.6 (99)	43.8 (46)	50 (25)	33.5 (53)
Two or more falls in past 6 months (%/n)	13.7 (36)	20 (21)	30 (15)	9.5 (15)
Hand grip strength (kg)	58.8 (16.6)	57.5 (17.5)	53.6 (17.5) [b]	59.7 (15.9) [b]
Geriatric depression score 30, GDS (score)	4.4 (4.3)	5.5 (5.3) [A]	6.5 (6.3) [B]	3.7 (3.3) [A, B]
Fall efficacy scale, FES-I(score)	20.5 (5.1)	21.4 (6.5) [a]	21.5 (6) [b]	19.9 (3.9) [a, b]
Cognitive function (MMSE score)	28.2 (1.6)	28.3 (1.6)	28.5 (1.7)	28.2 (1.6)
LASA fall risk profile (score)	3.4 (3.3)	4.4 (3.8) [A]	5.4 (4.5) [B]	2.8 (2.8) [A, B]
Inability of using transport(%/n)[1]	0.4 (1)	1 (1)	0 (0)	0 (0)
Living independently (%/n)	99.2 (261)	98.1 (103)	96 (48) [b]	100 (158) [b]
Use of a walking aid (%/n)	6.8 (18)	8.6 (9)	10 (5)	5.7 (9)

Note: [1]Either public or own transport. [A] significant difference between all fallers and non-fallers with p<0.01; [a]

significant difference between all fallers and non-fallers with p<0.05; [B] significant difference between walking

fallers and non-fallers with p<0.01; [b] significant difference between walking fallers and non-fallers with p<0.05.

Table 2. Model performance of fallers in follow-up 6 months for a) models based on

the FARAO cohort, b) the calibrated FARAO model performance, c) models based on

VIBE cohort

a) model based on FARAO cohort	All falls			Walk-falls		
	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3
Accuracy	0.58	0.50	0.49	0.63	0.53	0.75
Precision	0.48	0.37	0.40	0.34	0.20	0.49
Sensitivity	0.50	0.36	0.55	0.58	0.32	0.34
Specificity	0.64	0.59	0.45	0.64	0.59	0.89
AUC	0.62	0.50	0.53	0.68	0.49	0.59
b) calibrated FARAO model performance	All falls			Walk-falls		
	Calibrated Model1	Calibrated Model2	Calibrated Model3	Calibrated Model1	Calibrated Model2	Calibrated Model3
Optimal cut-offs	0.36	0.39	0.45	0.23	0.24	0.30
Accuracy	0.59	0.48	0.63	0.71	0.57	0.75
Precision	0.49	0.42	0.59	0.42	0.27	0.47
Sensitivity	0.53	0.86	0.28	0.50	0.48	0.36
Specificity	0.63	0.22	0.87	0.78	0.59	0.87
AUC	0.62	0.52	0.58	0.68	0.51	0.62
c) model based on VIBE cohort	All falls			Walk-falls		
	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3
Optimal cut-offs	0.38	0.40	0.37	0.20	0.23	0.20
Accuracy	0.62	0.53	0.57	0.66	0.54	0.6
Precision	0.53	0.43	0.47	0.38	0.27	0.33
Sensitivity	0.56	0.53	0.58	0.66	0.54	0.62
Specificity	0.66	0.53	0.56	0.66	0.54	0.59
AUC	0.63	0.53	0.65	0.69	0.56	0.73

561 **Table 3.** Mean (SD) and odds ratios (OR) for VIBE cohort (n=263) and FARAO
562 cohort (n=169) (van Schooten et al. 2015a)

Measures	Prevalence or mean (SD) for VIBE cohort	Prospective Falls OR (95% CI)	Prevalence or mean (SD) for FARAO cohort	Prospective Falls OR (95% CI)
Demographic information				
Age (years)	71.8 (5.7) ***	1.02 (0.8-1.3)	75.4 (6.8)	1.09 (0.8-1.5)
Gender (% female)	68.1 **	1.04 (0.6-1.8)	52.1	1.08 (0.6-2.06)
Weight (kg)	74.1 (13.5)	0.8 (0.6-1.1)	74.1 (14.2)	1.2 (0.9-1.7)
Height (cm)	169.1 (8.5) *	0.9 (0.7-1.1)	170.8 (8.9)	1.2 (0.9-1.7)
Questionnaires				
6-months history of falls	37.6%	1.6 (1.2-2.2) **	35.5%	3.1 (1.6–6.1) **
Hand grip strength (kg)	58.8 (16.6)	0.9 (0.7-1.1)	58.4 (20.5)	0.97 (0.7–1.4)
Geriatric depression score, GDS-30	4.4 (4.3)	1.5 (1.2-2.0) **	4.6 (4.2)	1.6 (1.1–2.2) **
Fall efficacy scale, FES-I (score)	20.5 (5.1)	1.3 (1.1-1.7) *	20.8 (4.4)	1.1 (1.0–1.1) *
Cognitive function (MMSE score)	28.2 (1.6) **	1.1 (0.9-1.4)	27.7 (2.2)	0.9 (0.7-1.3)
LASA fall-risk profile (score)	3.4 (3.3) ***	1.7 (1.3-2.2) **	5.5 (4.7)	1.6 (1.1–2.2) **
Inability to use public transportation	0.4% ***	N/A	7.1%	2.8 (0.8–9.5)
Living independently	99.2% ***	N/A	88.1%	1.0 (0.5-2.1)
Use of walking aid	6.8% ***	1.6 (0.6-4.1)	17.2%	1.4 (0.6–3.2)
Daily activities (per day)				
Number of strides (N)	4023 (2053) ***	1.1 (0.8-1.4)	3060 (1527)	0.9 (0.7–1.3)
Duration of lying (h)	9.2 (4.1) *	1.0 (0.8-1.3)	10.1 (2.01)	0.8 (0.6-1.1)
Locomotor duration (h)	1.5 (0.7) ***	1.1 (0.9-1.4)	1.2 (0.5)	0.8 (0.7–1.4)
Gait characteristics				
Walking speed (m/s)	0.6 (0.1) ***	0.8 (0.6-1.1)	0.9 (0.2)	0.6 (0.4–1.0) *
Logarithmic divergence rate AP (/stride)	1.5 (0.3) ***	0.9 (0.7-1.2)	1.9 (0.3)	1.7 (1.2–2.4) **
Range VT (m/s^2)	9.1 (1.9) ***	0.9 (0.7-1.1)	11.8 (5.6)	0.4 (0.2-0.8) **
Index of harmonicity ML	0.4 (0.2) ***	1.3 (1.0-1.7) *	2.0 (0.3)	1.4 (1.0-2.0) *
Sample entropy VT	0.2 (0.03) ***	0.9 (0.7-1.1)	0.3 (0.1)	1.4 (1.0-2.0)

Note: AP = anteroposterior direction; ML = mediolateral direction; VT = vertical direction; N/A = not applicable

with no solution (Infinity); Continuous parameters are z-transformed. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Significance indicators in the column for mean VIBE data illustrated significant differences between the mean (SD)

of the VIBE and FARAO data, as tested using an unpaired t-test. Significance indicators in the OR columns

indicate significance of the OR.

Figure 1. Flowchart of participants included for analyses. Fall records were
considered incomplete when one or more months of fall records were missing.

573
574 **Figure 2.** Main activities of all falls (A) and causes of walk-falls (B)