Deleterious effects of thermal and water stresses on life history and physiology: a case study on woodlouse

Charlotte DEPEUX,[1, 2], Angèle BRANGER[1], Théo MOULIGNIER[1], Jérôme MOREAU[3], Jean-François LEMAÎTRE[2], François-Xavier DECHAUME-MONCHARMONT[4], Tiffany LAVERRE[1], Hélène PAUHLAC[1], Jean-Michel GAILLARD[2], Sophie BELTRAN-BECH[1]

[1] Université de Poitiers, Laboratoire Ecologie et Biologie des interactions EBI, UMR CNRS 7267, 3, rue Jacques Fort, TSA 51106 86073 POITIERS Cedex 9, France.

[2] Université Lyon 1, Laboratoire de Biométrie et Biologie Evolutive UMR CNRS 5558, 43 Boulevard du 11 novembre 1918, 69622 Villeurbanne cedex, France.

[3] Université de Bourgogne, Laboratoire Biogéosciences, UMR CNRS 6282, 6 Boulevard Gabriel, 21000 Dijon, France.

[4] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622, Villeurbanne, France

Corresponding author: sophie.beltran.bech@univ-poitiers.fr

Abstract

We tested independently the influences of increasing temperature and decreasing moisture on life history and physiological traits in the arthropod *Armadillidium vulgare.* Both increasing temperature and decreasing moisture led individual body mass and reproductive success to decrease. While immune cells density decreased and β-galactosidase activity increased with increasing temperature and decreasing moisture, suggesting a negative impact of these stresses on individual quality, increased temperature and decreased moisture affected differently the other biomarkers conjuring different underlying mechanisms depending on the stress applied. Our findings demonstrate overall a negative impact of high temperature and low moisture on woodlice welfare. Changing temperature or moisture had slightly different effects, illustrating the need of testing further the respective role of each of these key components of climate change on organisms to predict more reliably the future of our ecosystems.

Key words

Abiotic stresses, life history traits, physiological traits, arthropods, climate change

Introduction

The Intergovernmental Panel on Climate Change (IPCC) forecasts an average increase in temperature between +1.5°C and +4°C in 2100 (Masson-Delmotte et al., 2021). Not only will average temperatures and the frequency and intensity of precipitation change, but extreme events will increase in frequency. Although the link between global warming and drought is still highly debated and may not be direct (Trenberth et al., 2014), droughts due to a decrease in rainfall and an increase in evaporation are expected to take place in the coming decades (Dai, 2013), and should be much more intense than current droughts (Trenberth et al., 2014). As deterioration of environmental conditions are known to impact life history traits such as growth rate, reproductive success or longevity (e.g. Chen et al., 2019; Johnson and Jones, 2016; Khadioli et al., 2014), identifying the potential implications of climate change for organisms is a research question of paramount importance.

Terrestrial arthropods are ectotherms that are particularly sensitive to temperature and moisture changes (Lister and Garcia, 2018; Maron et al., 2015). Global warming constitutes a threat for them, as increasingly reported (Johnson and Jones, 2016). Thus, in Lepidoptera, too

high temperatures prevent hatching (Khadioli et al., 2014). In both Lepidoptera and Hymenoptera, survival decreases with increasing temperature (Abou-Shaara et al., 2012; Khadioli et al., 2014). In some Coleoptera, egg viability decreases and hatching time increases for viable eggs when they are exposed to drought (Johnson et al., 2010). When facing the costs of increased temperature and drought frequency and intensity on life history traits, arthropods display different responses to resist or tolerate such changes (Strachan et al., 2015). For example, some arthropods can migrate to refugia, others can implement physiological resistance tactics (e.g. resistant eggs) and/or dormancy, and others are able to alter their life cycle and/or development (Strachan et al., 2015; Verberk et al., 2008). In organisms with limited movement capacity, increased temperature and decreased moisture are expected to induce pronounced physiological stresses. Studying how these stresses affect both life history traits and physiological traits would allow us to anticipate the effect of global warming on organisms with limited movement capacity.

The common woodlice *Armadillidium vulgare* is a key soil decomposer naturally exposed to a wide range of environmental conditions (Souty-Grosset et al., 1988) that provides major ecosystem services (David and Handa, 2010), notably in agrosystems and grassland habitats where it is used as an ecological indicator. In the course of its evolutionary history, the common woodlice had to adapt to terrestrial life. Consequently, it is still very sensitive to moisture and temperature variations, which can induce water loss (Smigel and Gibbs, 2008) and have major consequences in terms of distribution, behaviour and survival (Hassall et al., 2018; Paris and Pitelka, 1962). Moreover, their movement capacity is low (i.e. several hundred meters at the best during the entire lifetime, Durand et al., 2019) to allow them to migrate so to avoid the stress imposed by the environment. Our knowledge and ability to measure woodlice life history traits and the availability of molecular and cellular biomarkers of individual quality (Depeux et al., 2020a) make this species a highly relevant experimental model to study the influence of both temperature and moisture on life history and physiological traits.

In this study, we tested independently the effects of increased temperature (experiment 1) and of decreased moisture (experiment 2) on a selected set of key life history (i.e. growth, reproductive success, and survival) and physiological (i.e. immune cell parameters (cell viability, cell density and cell size) and β-galactosidase activity) traits in woodlice. For the

experiment 1 (testing the effect of increased temperature), we compared individuals maintained at 20°C and 80% of moisture (i.e. the standard temperature and moisture laboratory conditions) to individuals exposed at 28°C (simulating a temperature increase of 8°C) still at 80% of moisture. For the experiment 2 (testing the effect of decreased moisture), we compared animals in standard conditions to individuals exposed at 50% of moisture (simulating a moisture loss of 30%) still at 20°C. We predicted that a rise in temperature and a loss in moisture should be stressful and should induce changes in life history and physiological traits.

MATERIALS & METHODS

Biological Material – Routine Breeding

All specimens of *A. vulgare* used in this study descend from individuals sampled in Denmark (Helsingör) in 1982. Since then, animals were reared under laboratory conditions under the natural photoperiod of Poitiers (France) (46°35'N; 0°20'E), at 20°C and about 80-85% of moisture, in plastic boxes (length × width × height: 26.5 × 13.5 × 7.5 cm) containing humid loam, and fed ad libitum with carrot slices and dried linden leaves. Controlled breeding, for the maintenance of the lineage over years, is performed in individual boxes (diameter x height: 9,8cm x 4,9cm), with reproductive pairs selected from their pedigree to avoid inbreeding. One month after mating, the offspring exit the female *marsupium* (i.e. female ventral pouch on which the eggs develop) (Suzuki and Ziegler, 2005)*.* We transferred these offspring a few days after birth into a bigger box (length × width × height: 26.5 × 13.5 × 7.5 cm) with loam and food. After 3 months, once sexual characters have appeared but before sexual maturity, we placed young males and females in separate boxes (length × width × height: 26.5 × 13.5 × 7.5 cm) in laboratory conditions described above, enabling us to obtain virgin adults. For the maintenance of the lineage, about 40 crosses have been performed following this protocol each year. Each of the 40 broods provides at least one breeder for the next generation. The animals used in the experiments of this study came from this controlled lineage.

Experimental Design

Experiment 1: effect of increased temperature on life history and physiological traits

The experiment 1 performed in January 2019 involved the comparison between two groups of animals aged of 7 months old maintained in different temperature conditions in two

climatic chambers (Memmert HPP 256L with LED Light module cold white 6500K for HPP260 (15%) and Interior IP68 socket (for temperature restriction)) during two months after classical conditions of maintenance:

(i)   The "Control Temperature" group (CT) of animals maintained in standard conditions (i.e. at 20°C and 80% of moisture) in one of our two climatic chambers.

(ii)   The "High Temperature" (HT) group of animals exposed at 28°C (simulating increased temperature by 8°C) and at 80% of moisture in the second climatic chamber.

Eight degrees (i.e. difference in temperature between the two groups) corresponds to a temperature increase close to daily variations observed in Poitiers during some summers, which could chronically induce stress. Moreover, we had observed the stressful effect of this temperature increase in a preliminary experiment in which we did not control the moisture variation (Depeux et al., 2019).

In each group, animals were kept with fed *ad libitum* in 3 boxes (length × width × height: 26.5 × 13.5 × 7.5 cm; classical laboratory density conditions) of 30 females and 3 boxes of 30 males from 15 different clutches (i.e. all compared conditions contained animals with the same overall genetic background (from 15 same clutches) to be comparable). For each condition, one box was used to monitor survival and growth (mass gain over time) of animals from the beginning to the end of the experiment, another was used to evaluate reproductive success and the last box served to quantify physiological traits (i.e. immune cell parameters (cell viability, cell density and cell size) and β-galactosidase activity) (see below).

Experiment 2: effect of a loss of moisture on life history and physiological traits

The experiment 2 performed in January 2021 involved the comparison between two groups of animals also aged of 7 months old maintained in different conditions in our two climatic chambers during two months after standard conditions of maintenance:

(i)   The "Control Moisture" (CM) group of animals maintained in standard conditions (i.e. at 20°C and 80% of moisture) in one of our two climatic chambers

(ii)   The "Loss of Moisture" (LM) group of animals exposed at 50% of moisture (simulating a moisture loss of 30%) and at 20°C in the second climatic chamber.

Similar to the experiment 1, in each group, animals were fed *ad libitum* in 3 boxes of 30

females and 3 boxes (length × width × height: 26.5 × 13.5 × 7.5 cm; standard laboratory density

conditions) of 30 males from 15 different clutches (i.e. all boxes to compare contained animals

with the same overall genetic background (from 15 same clutches)). For each condition, one

box was used to monitor survival and growth of animals from the beginning to the end of the

experiment, another was used to evaluate reproductive success and the last box served to

quantify physiological traits (see below).

In our two experiments, we aimed to compare individuals of the same age because age

negatively impacts both reproductive success (Depeux et al., 2020b) and physiological traits

(Depeux et al., 2020a). Having initially only two climatic chambers, we had to perform our

experiments 1 and 2 in different years (i.e. experiment 1 in 2019 and experiment 2 in 2021).

Thus, we systematically compared the effect of each stress against its own control condition

group (i.e. CT for HT and CM for LM). Moreover, we selected, at the beginning of each

experiment (1 and 2), individuals of the same size and we checked, at the end of the

experiments, potential statistical differences between the two control groups (CT and CM) on

measures of life history and physiological traits (Supp. File1). Although β-galactosidase activity

and cell density were higher in the CT group than in the CM group (Supp. File1), we observed

the same dynamics in these measures in the face of their stressful condition (HT and LM,

respectively) (see Results part). The body mass at day 14 was higher in the CM group than in

the CT group (Supp. File1). Whatever the differences observed between the two control

groups (CT group used in 2019 and CM group used in 2021), we compared the effect of each

stress (HT in 2019 and LM in 2021, respectively) against its own control group (CT group in

2019 and CM group in 2021, respectively) for testing the effect of each stress.

Life history traits

Survival and growth

One box of males and one box of females from each group (i.e. for the groups CT, HT, CM, LM)

were used to monitor and compare changes of survival and body mass over time. All

individuals in these boxes were monitored for 124 days (i.e. about 4 months). We sampled

individuals at 14, 28, 42, 69 and 124 days (i.e. 5 sampling points per box) and assessed

survivorship and change in body mass (in grams) of all surviving animals in each box (body

mass was measured with a precision balance 650g|1mg Sartorius™ BCE653-1S Entris™ II

Essential). Then, we compared these traits over time and between groups (CT vs. HT groups

and CM vs. LM groups) to test independently the effect of temperature and moisture changes

on these traits (see section on Statistical analyses). Due to their regular moults, individual

identification among the 30 animals sharing in given box cannot be performed, leading our

measures to be average survival and growth instead of individual trajectories.

Reproductive success

At the end of the exposure to different conditions, one box of males and one box of females

were collected from each group (i.e. for the groups CT, HT, CM, and LM). We formed 20

breeding pairs composed of one male and one female per group. Each breeding pair was

placed in a box, at 20°C, with food provided *ad libitum* and in a photoperiod of 16:8 (L/D)

stimulating the reproduction (Mocquard et al., 1989). We followed all these pairs for 5 months

during which each clutch produced was recorded. At the end of this period, the ability to

produce a clutch (i.e. the probability that one clutch or more is produced by a given pair) for

the 80 pairs (i.e. 40 pairs for experiment 1 and 40 pairs for experiment 2) was compared

independently between CT and HT groups and between CM and LM groups to test the effect

of temperature and moisture changes on breeding success. As we created groups from similar

clutches to have the same overall genetic background among boxes, we cannot exclude that

some crosses were composed of related individuals although we expect this event to be rare.

However, the probability of forming sibling pairs (8%) was similar among groups that were

exposed either at 20°C vs. 28°C or at 80% vs. 50% of moisture.

Physiological traits

At the end of the experimental treatments (i.e. after two months in our experimental

conditions), one box of males and one box of females were taken from each group (i.e. for the

groups CT, HT, CM, and LM) for measuring the level of our set of physiological traits (i.e.

immune cells parameters and β-galactosidase activity) developed in Depeux et al. (2020a).

These physiological traits were firstly described as senescence biomarkers because they allow

predicting the amount of cellular senescence in different organisms and are strongly age-

dependent in *A. vulgare* (i.e. older the individual, higher the decline of these biomarkers,

Depeux et al. 2020a). We performed these measures on each remaining animals (Table 1) and

we compared these metrics independently between CT and HT groups for experiment 1 and

between CM and LM groups for experiment 2 (see section Statistical analyses).

Table 1. Numbers of individuals on which we measured quality biomarkers.

Groups	CT (Control temperature)	HT (High temperature)	CM (Control Moisture)	LM (Loss of Moisture)
Numbers of females	13	12	9	15
Numbers of males	17	17	15	15

Immune cells

As immune cells are free-circulating, they can inform about a potential premature biological

aging. When an individual *A. vulgare* ages, its immune cells decrease in density and viability

while increasing in size (Depeux et al., 2020a). To measure these parameters, we collected 3µL

of haemolymph per individual and placed it in 15µL of MAS-EDTA (EDTA 9 mM, Trisodium

citrate 27 mM, NaCl 336 mM, Glucose 115 mM, pH 7, (Rodriguez et al., 1995)). We then added

6µL of Trypan Blue at 0.4% (Invitrogen) to discriminate live and dead cells. After, 10µL of this

solution was put in Invitrogen Coutness® counting slide and put in an automated Cell Counter

(Invitrogen) to quantify cell density (measured as the number of cells per mL of haemolymph),

viability (measured as the proportion of live cells) as well as cell size (in µm). These three

parameters of the immune cells are physiological traits that were found to be reliable

biomarkers of cellular senescence in *A. vulgare* (Depeux et al., 2020a).

β-galactosidase activity

The β-galactosidase activity is a physiological trait commonly used as a marker of cellular

senescence (Lee et al., 2006). Its indirect activity in regards to the process of cellular

senescence increases with age in *A. vulgare* (Depeux et al., 2020a). To measure this enzymatic

activity, we dissected and removed the nerve cord of each individual after having collected

haemolymph for assessing the immune parameters. We put individual nerve cords in 300µL

of Lyse Buffer 1X (CHAPS 5 mM, Citric acid 40 mM, Sodium Phosphate 40 mM, Benzamidine

0.5 mM, PMSF 0.25 mM, pH = 6) (Gary and Kindell, 2005). We centrifuged the sample at 15

000g for 30 minutes at 4°C and then we collected and kept the supernatant at -80°C. We

quantified the protein concentration thanks to the BCA Assay and we homogenized all

samples at the 0.1mg/mL protein concentration. Then, 100µL of these protein extracts were

added to 100µL of reactive 4-methylumbelliferyl-D-galactopyranoside (MUG) solution. The synthesis of the fluorescent 4-methylumbelliferone (4-MU), the result of the contact of MUG reactive with β-galactosidase, was measured using the multimode microplate reader Mithras LB940 133 HTS III, Berthold; excitation filter: 120 nm, emission filter 460 nm, for 120 minutes. We included two technical replicates for each sample to obtain the measures.

Statistical analyses

The effects of the stress condition (control vs. high temperature, or control vs. low moisture) on life history and physiological traits were tested using the following models. *(i)* Life history traits. For the survival data, Cox proportional hazard models were fitted with stress condition, sex and their interaction term as fixed variables, using 'survival' package (Therneau, 2022). For the growth data, the body mass was modelled with linear model with Gaussian distribution with stress condition, sex, time (in days) and their two-by-two interaction term as fixed variables. The female reproductive success was modelled as binary data (*presence of at least one clutch or absence of clutch)* using linear regression with a binomial distribution, with stress condition as fixed variables. *(ii)* Physiological traits. The cell density, cell size, cell viability and the β-galactosidase activity were modelled with a linear model with Gaussian distribution with stress condition, sex and their interaction term as fixed variables.

We proceeded to model selection starting with full (saturated) model. We ranked all nested models according to their AICc using 'MuMIn' package (Barton, 2022). We selected the most parsimonious models among the top ranked models (ΔAICc < 2) (Galipaud et al., 2017). The tables summarizing the model selection procedure was presented in Supp. File2. To represent the effect of the two environmental stresses in each variable, we presented our results with indices of size effect. The effect of each stress on each measure of life history traits and individual quality biomarkers were measured using the standardized slopes (Schielzeth et al 2010) and their SE calculated by rescaling the variable of the selected model. For survival data, the effect size was the hazard ratio, calculated as the exponential of the regression parameter (Collett 2003). When the selected model did not include the effect of the stress, we took the model with the variable stress as fixed factor to obtain a size effect as made in Depeux et al. 2020a. All statistical analyses and representations of results were performed using the software R 4.2.1 (R core Team 2022).

All datasets and source code are available as electronic supplementary materials on public

repository: https://gitlab.com/fxdm/armadillidium_stress

RESULTS

Life history traits

The survival was not impacted by an increased temperature (χ^2_1=2.16, P=0.14, Fig.1A, Supp.

File2 Table S2a, Supp. File3-1.A.1) although mortality risk was almost twice as lower at low

compared to high temperature. The hazard ratio was 1.78, with a 95%CI including 1 [0.81;

3.88]. By contrast, individuals exposed to a loss of moisture had a 2.5 times higher mortality

risk (χ^2_1=4.54, P=0.03, Fig. 1B, Supp. File2 Table S2b, Supp. File3-1.A.2). The hazard ratio was

2.69, with a 95%CI excluding 1 [1.03; 7.01]. As a result, 90% of individuals placed at control

moisture were still alive at the end of the follow-up, whereas only 75% of individuals placed

at 50% of moisture survived at the end of the experiment.

In both temperature and moisture stresses, body mass increased during the entire experiment

duration (Fig. 1C and 1D, Supp. File3-1.B.1 and 1.B.2), as expected in an indeterminate grower

as *A. vulgare*, but there was no detectable interaction between day and sex (Supp. File2 Table

S2c and Table S2d). For the temperature experiment, interactions between sex and stress

($F_{1,528}$=6.90, P=0.0088, Supp. File3), and between day and stress ($F_{1,528}$=14.6, P=0.00015, Supp.

File2 Table S2c) showed up, illustrating the impact of an increasing temperature on growth.

By contrast, in the moisture stress experiment, the body mass was not affected by the sex

($F_{1,522}$=0.35, P=0.55), the stress condition ($F_{1,522}$=0.31, P=0.58), or any interaction between the

variables (all P > 0.10).

The reproductive success markedly decreased in both experiments for the stressful condition:

a fourfold increase of reproductive failure with temperature stress (χ^2_1=5.02, p=0.025, Odd-

ratio=0.23, 95%CI=[0.057;0.83], Fig. 1E, Supp. File2 Table S2e, Supp. File3-1.C.1), and a fivefold

increase of reproductive failure with moisture stress (χ^2_1=5.38, p=0.02, Odd-ratio=0.20,

95%CI=[0.045;0.79], Fig. 1F, Supp. File2 Table S2f, Supp. File 3-1.C.2). In both cases, it

corresponds to halving the reproductive success in the stress groups (moisture stress: 55% in

the control group vs. 20% in the stressed group; temperature stress: 65% in the control group

vs. 30% in the stressed group).

Figure 1. **Effect of the two environmental stresses (Temperature (A, C and E) and Moisture (B, D and F) on Survival (A and B), Body mass (C and D) and Reproductive success (E and F).** Blue colour: control groups, orange colour: stress groups. CT: Control Temperature (20°C), HT: High Temperature (28°C), CM: Control Moisture (80%), LM: Loss of Moisture (50%). NS: No significant; * p<0.05.

Physiological traits

Immune cells

Immune cell viability was not affected by increasing temperature ($F_{1,56}$=0.92, p=0.34,

standardized slope β=-0.25; 95%CI=[-0.79;0.27], Fig. 2A, Supp. File2 Table S2g and Supp. File3-

2.A.1) but decreased during a loss of moisture ($F_{1,50}$=4.17, p=0.046, standardized slope β=0.55;

95%CI=[0.01;1.09], Fig. 2B, Supp. File2 Table S2h and Supp. File3-2.A.2).

Immune cell size decreased during the temperature stress ($F_{1,55}$=5.72, p=0.02, standardized

slope β=-0.60; 95%CI=[-0.1;-1.1], Fig. 2C, Supp. File2 Table S2i and Supp. File 3-2.B.1) but not

under the moisture stress ($F_{1,50}$=3.79, p=0.057, standardized slope β=-0.55; 95%CI=[-

1.07;0.02], Fig. 2D, Supp. File2 Table S2j and Supp. File3-2.B.2).

Immune cell density decreased during the temperature stress ($F_{1,56}$=38.2, P<0.001,

standardized slope β=-1.26; 95%CI=[-1.67;-0.85], Fig. 2E, Supp. File2 Table S2k and Supp. File3-

2.C.1)) and the moisture stress ($F_{1,50}$=7.64, p=0.008, standardized slope β=0.72;

95%CI=[0.19;1.25], Fig. 2F, Supp. File2 Table S2l and Supp. File3-2.C.2).

β-galactosidase activity

The *β*-galactosidase activity increased when the temperature increased ($F_{1,54}$=11.32,

P=0.0014, standardized slope β=0.82; 95%CI=[0.33;1.32], Fig.2G, Supp. File2 Table S2m, and

Supp. File3-2.D.1), but also when the moisture decreased ($F_{1,50}$=10.50, P=0.002, standardized

slope β=-0.83; 95%CI=[-1.31;-0.32], Fig. 2H, Supp. File2 Table S2n and Supp. File3-2.D.2).

Figure 2. Effect of the two environmental stresses (Temperature (A, C, E, and G) and Moisture (B, D, F and H) on immune cell viability (A and B), immune cell size (C and D), immune cell density (E and F) and β-galactosidase activity (G and H). Blue colour: control groups, orange colour: stress groups. CT: Control Temperature (20°C), HT: High Temperature (28°C), CM: Control Moisture (80%), LM: Loss of Moisture (50%). NS: No significant; * p<0.05, ** p<0.01, *** p<0.001.

DISCUSSION

Our results highlight that life history traits were negatively impacted by the two environmental stresses (temperature and moisture) considered in this study. Moreover, the detrimental effects of these stresses on our set of individual quality biomarkers are consistent with an overall premature ageing of stressed animals compared to unstressed ones. To briefly summarize, an increase in temperature negatively impacted the body mass trajectory over time and the reproductive success of individuals. A decrease in moisture resulted in a decrease of both survival and reproductive success. Concerning our physiological traits: (1) the density of immune cells decreased under both stresses, (2) immune cell size decreased under temperature stress, but was not impacted under moisture stress, (3) the viability of the cells decreased when the moisture decreased (but not with temperature increase) and (4) finally, the β-galactosidase activity increased for the two stressed groups. In this context, our results globally support marked negative effect of higher temperature and lower moisture on woodlice performance, with some minor differences between the two kinds of stresses in their effects on the different measured life history and physiological traits.

About the life history traits, if the increase of temperature had no detectable effect on survival in *A. vulgare*, contrary to what was previously found in arthropods studied so far such as *Antestiopsis thunbergii*, *Calliphora stygia* and *Margaritifera margaritifera* (Azrag et al., 2017; Hassall et al., 2017; Kelly et al., 2013), this stress led to a slowdown in woodlice growth, in line to what has been reported in three other isopods (Angilletta et al., 2004). In parallel, we observed that the increase in temperature led to a decrease in reproductive success, as previously observed in females of *Antestiopsis thunbergii* (Azrag et al., 2017). In *A. vulgare*, the animal size is positively correlated with fecundity (Durand et al., 2018; Lawlor, 1976), meaning that the slowdown in growth could explain, at least partly, the decrease in reproductive success for stressed animals compared to non-stressed ones. Concerning the moisture stress, if the loss of moisture had no detectable effect on woodlouse growth, it caused a decrease in both survival and reproductive success. These findings suggest a high cost of drought on individual fitness in *A. vulgare*.

About the physiological traits, the results of the temperature stress experiment showed that although cell density was negatively impacted by increased temperature, cell viability was not affected. Moreover, contrary to what was expected when individuals are senescent, cell size

decreased instead of increasing. This last result supports our previous finding that cells decreases in small when the temperature raises (without controlling moisture level, Depeux et al., 2019). That little cells came from a more important cell renewal in stressed animals might explain this pattern. On the other hand, the increase of β-galactosidase activity seems to indicate premature ageing (and thus a decrease in quality) in individuals exposed to increased temperature. Concerning the moisture stress, the individual quality biomarkers indicated a decrease in cell density and viability, associated with an increase in β-galactosidase activity, which and suggest an acceleration of biological ageing in the individuals exposed to a decrease in moisture (Depeux et al., 2020a).

We observed global negative effect of temperature increase in *A. vulgare* in our study, but our results seemed to show an even higher and clearer effect of loss of moisture on both life history traits and individual quality biomarkers. Although the woodlouse has become terrestrial for a long time, they are still dependent on and require a substantial water supply (Smigel and Gibbs, 2008). Thus, behaviours like aggregation allowing them to resist to desiccation have been set up and allow them to maintain the rate of moisture required for their survival (Broly et al., 2013; Smigel and Gibbs, 2008). This can explain the strong effect of loss of moisture in our study. In natural conditions, increase in temperature and loss of moisture generally positively covary, leading to even higher negative consequences on the woodlouse performance. Further work will be required to test the influence of more extreme and maybe more realistic conditions by simultaneously increasing temperature by 8°C decreasing moisture by 30% on our measured life history and physiological traits. A study in the wild comparing life history and physiological traits on *A. vulgare* collected across areas with different temperature and drought gradient would also allow a better assessment of the combined effects of these stresses.

Thanks to our experimental design that allowed us to test independently the stresses that the organisms are likely to face, we showed that thermal and water stress do not have the same impact on organisms. Although simulations based on mathematical models have predicted that both temperature and drought changes overall affect arthropods, experimental approaches (such as this work) are required to quantify reliably the effect of temperature changes (Johnson et al. 2010). Drought can have serious physiological consequences on invertebrates, such as protein denaturation and undesirable macromolecular interactions

(Sano et al., 1999; Tang and Pikal, 2005), or oxidative damage (Lopez-Martinez et al., 2008),

which are known to be involved in the process of cellular senescence (Gilca et al., 2007) and

thus in decrease in performance of stressed organisms. Due to the role of arthropods in

services to many ecosystems (e.g. biochemical balance of ecosystems, agriculture, pest

management...), and in the context of global warming, it is crucial to understand the effects

of temperature and moisture changes on these organisms (Santos et al., 2021). As

temperature increase is not the only environmental change expected to take place in the

coming years, it is of paramount importance to assess the impact of other stresses too.

Although many predictive models have been proposed so far, getting more accurate

information on the expected responses of organisms facing with different kinds of stress

would provide the required information to test these model predictions.

To conclude, *A. vulgare* is an important actor that delivers ecosystem services in many

ecosystems because it actively impacts soil fertility (Souty-Grosset and Faberi, 2018) and it is

also used as an ecological indicator in grassland habitat (Paoletti and Hassall, 1999; Souty-

Grosset et al., 2005). This detritivorous species promotes decomposition processes and

nutrient cycling on which agricultural productivity and sustainability depend (Bredon et al.,

2018; Paoletti and Hassall, 1999), and plays thereby a key role in ecosystem services (David

and Handa, 2010). Extending knowledge in the response of soil biodiversity facing current

global changes could promote sustainability by helping to the development of new tools and

strategies for more efficient management of soils and associated crops, through more

effective and targeted recolonisation and/or restoration of soil biodiversity. Also, to better

understand what the future of the animal communities in the current context of global

warming will be, it seems necessary to multiply the studies on models presenting particular

ecological requirements, like the woodlouse.

Supplementary files

Supp. File1 Comparison of the two control groups

Supp. File2 Model selection

Supp. File3 Graphical representations of results per sex

ACKNOWLEDGMENTS

This work was supported by the French ministry of Education, the 2015–2020 State-Region Planning contract and European Regional Development Fund (FEDER) and intramural funds from the Centre National de la Recherche Scientifique and the University of Poitiers and by a grant from the Agence Nationale de la Recherche (ANR-15-CE32-0002-01). We would like to thank Alexandra Lafitte for technical assistance.

BIBLIOGRAPHY

Abou-Shaara, H.F., Al-Ghamdi, A.A., and Mohamed, A.A. (2012). Tolerance of two honey bee races to various temperature and relative humidity gradients. Environ. Exp. Biol. *10*, 133–138.

Angilletta, M.J., Jr., Steury, T.D., and Sears, M.W. (2004). Temperature, growth rate, and body size in ectotherms: fitting pieces of a life-history puzzle. Integr. Comp. Biol. *44*, 498–509.

Azrag, A.G.A., Murungi, L.K., Tonnang, H.E.Z., Mwenda, D., and Babin, R. (2017). Temperature-dependent models of development and survival of an insect pest of African tropical highlands, the coffee antestia bug *Antestiopsis thunbergii* (Hemiptera: Pentatomidae). J. Therm. Biol. *70*, 27–36.

Bartoń, K. (2022). MuMIn: Multi-Model Inference. R package version 1.46.0, https://CRAN.R-project.org/package=MuMIn.

Bredon, M., Dittmer, J., Noël, C., Moumen, B., and Bouchon, D. (2018). Lignocellulose degradation at the holobiont level: teamwork in a keystone soil invertebrate. Microbiome *6*, 162.

Broly, P., Deneubourg, J.-L., and Devigne, C. (2013). Benefits of aggregation in woodlice: a factor in the terrestrialization process? Insectes Soc. *60*, 419–435.

Chen, C., Yang, H., Xue, F., and Xia, Q. (2019). Geographical variation in life-history traits suggests an environmental-dependent trade-off between juvenile growth rate and adult lifespan in a moth. Bull. Entomol. Res. *109*, 626–632.

Collett, D. (2003). *Modelling survival data in medical research. Chapman and Hall, London.*

Dai, A. (2013). Increasing drought under global warming in observations and models. Nat. Clim. Change *3*, 52–58.

David, J.-F., and Handa, I.T. (2010). The ecology of saprophagous macroarthropods (millipedes, woodlice) in the context of global change. Biol. Rev. *85*, 881-895.

Depeux, C., Samba-Louaka, A., Braquart-Varnier, C., Moreau, J., Lemaître, J.-F., Laverre, T., Pauhlac, H., Dechaume-Moncharmont, F.-X., Gaillard, J.-M., and Beltran-Bech, S. (2019). Impact of temperature and photoperiod on survival and biomarkers of senescence in common woodlouse. BioRxiv https://www.biorxiv.org/content/10.1101/433011v4.

Depeux, C., Samba-Louaka, A., Becking, T., Braquart-Varnier, C., Moreau, J., Lemaître, J.-F.,
Laverre, T., Paulhac, H., Dechaume-Moncharmont, F.-X., Gaillard, J.-M., and Beltran-Bech, S.
(2020a). The crustacean *Armadillidium vulgare* (Latreille, 1804) (Isopoda: Oniscoidea), a new
promising model for the study of cellular senescence. J. Crustac. Biol. *40*, 194–199.

Depeux, C., Lemaître, J., Moreau, J., Dechaume-Moncharmont, F., Laverre, T., Paulhac, H.,
Gaillard, J., and Beltran-Bech, S. (2020b). Reproductive senescence and parental effects in an
indeterminate grower. J. Evol. Biol. *33*, 1256-1264.

Durand, S., Loiseau, V., Prigot, C., Braquart-Varnier, C., and Beltran-Bech, S. (2018). Producing
offspring in *Armadillidium vulgare* : Effects of genetic diversity and inbreeding. Evol. Dev. *20*,
65–77.

Durand, S., Grandjean, F., Giraud, I., Cordaux, R., Beltran-Bech, S., and Bech, N. (2019). Fine-
scale population structure analysis in *Armadillidium vulgare* (Isopoda: Oniscidea) reveals
strong female philopatry. Acta Oecol. *101*, 103478.

Galipaud M, Gillingham MA, and Dechaume-Moncharmont F-X (2017) A farewell to the sum
of Akaike weights: the benefits of alternative metrics for variable importance estimations in
model selection. *Methods in Ecology and Evolution* 8:1668–1678.

Gary, R.K., and Kindell, S.M., 2005. Quantitative assay of senescence-associated β-
galactosidase activity in mammalian cell extracts. Anal. Biochem. *343,* 329–334.

Gilca, M., Stoian, I., Atanasiu, V., and Virgolici, B. (2007). The oxidative hypothesis of
senescence. J. Postgrad. Med. *53*, 207–213.

Hassall, C., Amaro, R., Ondina, P., Outeiro, A., Cordero-Rivera, A., and San Miguel, E. (2017).
Population-level variation in senescence suggests an important role for temperature in an
endangered mollusc. J. Zool. *301*, 32–40.

Hassall, M., Moss, A., Dixie, B., and Gilroy, J.J. (2018). Interspecific variation in responses to
microclimate by terrestrial isopods: implications in relation to climate change. ZooKeys *801*,
5–24.

Johnson, S.N., and Jones, T.H. (2016). Introduction to global climate change and terrestrial
invertebrates. In global climate change and terrestrial invertebrates, S.N. Johnson, and T.H.
Jones, eds. (Chichester, UK: John Wiley & Sons, Ltd), pp. 1–8.

Johnson, S.N., Gregory, P.J., McNicol, J.W., Oodally, Y., Zhang, X., and Murray, P.J. (2010).
Effects of soil conditions and drought on egg hatching and larval survival of the clover root
weevil (*Sitona lepidus*). Appl. Soil Ecol. *44*, 75–79.

Kelly, M.A., Zieba, A.P., Buttemer, W.A., and Hulbert, A.J. (2013). Effect of temperature on the
rate of ageing: an experimental study of the blowfly *Calliphora stygia*. PloS One *8*, e73781.

Khadioli, N., Tonnang, Z.E.H., Muchugu, E., Ong'amo, G., Achia, T., Kipchirchir, I., Kroschel, J.,
and Le Ru, B. (2014). Effect of temperature on the phenology of *Chilo partellus* (Swinhoe)
(Lepidoptera, Crambidae); simulation and visualization of the potential future distribution of

C. partellus in Africa under warmer temperatures through the development of life-table parameters. Bull. Entomol. Res. *104*, 809–822.

Lawlor, L.R. (1976). Parental investment and offspring fitness in the terrestrial isopod *Armadillidium vulgare* (Latreille) (Crustacea: Oniscidea). Evolution *30*, 775–785.

Lee, B.Y., Han, J.A., Im, J.S., Morrone, A., Johung, K., Goodwin, E.C., Kleijer, W.J., DiMaio, D., and Hwang, E.S. (2006). Senescence-associated β-galactosidase is lysosomal β-galactosidase. Aging Cell *5*, 187–195.

Lister, B.C., and Garcia, A. (2018). Climate-driven declines in arthropod abundance restructure a rainforest food web. Proc. Natl. Acad. Sci. *115*, E10397–E10406.

Lopez-Martinez, G., Elnitsky, M.A., Benoit, J.B., Lee, R.E., and Denlinger, D.L. (2008). High resistance to oxidative damage in the Antarctic midge *Belgica antarctica*, and developmentally linked expression of genes encoding superoxide dismutase, catalase and heat shock proteins. Insect Biochem. Mol. Biol. *38*, 796–804.

Maron, M., McAlpine, C.A., Watson, J.E.M., Maxwell, S., and Barnard, P. (2015). Climate-induced resource bottlenecks exacerbate species vulnerability: a review. Divers. Distrib. *21*, 731–743.

Masson-Delmotte, V., P. Zhai, A., and Pirani, S.L. (2021). IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.

Mocquard, J.P., Juchault, P., and Souty-Grosset, C. (1989). The role of environmental factors (temperature and photoperiod) in the reproduction of the terrestrial isopod *Armadillidium vulgare* (Latreille, 1804). Monogr. Monit. Zool. Ital. *4*, 455–475.

Paoletti, M.G., and Hassall, M. (1999). Woodlice (Isopoda: Oniscidea): their potential for assessing sustainability and use as bioindicators. Agric. Ecosyst. Environ. *74*, 157–165.

Paris, O.H., and Pitelka, F.A. (1962). Population characteristics of the terrestrial isopod *Armadillidium vulgare* in California grassland. Ecology *43*, 229.

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rodriguez, J., Boulo, V., Mialhe, E., and Bachere, E., 1995. Characterisation of shrimp haemocytes and plasma components by monoclonal antibodies. J. Cell Sci. *108,* 1043–1050.

Sano, F., Asakawa, N., Inoue, Y., and Sakurai, M. (1999). A dual role for intracellular trehalose in the resistance of yeast cells to water stress. Cryobiology *39*, 80–87.

Santos, J.C., de Almeida, W.R., and Fernandes, G.W. (2021). Arthropods: why it is so crucial to know their biodiversity? In measuring arthropod biodiversity, J.C. Santos, and G.W. Fernandes, eds. (Cham: Springer International Publishing), pp. 3–11.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients. Methods Ecol. Evol. *1*:103–113.

Smigel, J.T., and Gibbs, A.G. (2008). Conglobation in the pill bug, *Armadillidium vulgare*, as a
water conservation mechanism. J. Insect Sci. *8*.

Souty-Grosset, C., and Faberi, A. (2018). Effect of agricultural practices on terrestrial isopods:
a review. ZooKeys *801*, 63–96.

Souty-Grosset, C., Chentoufi, A., Mocquard, J.P., and Juchault, P. (1988). Seasonal
reproduction in the terrestrial isopod *Armadillidium vulgare* (Latreille): geographical
variability and genetic control of the response to photoperiod and temperature. Int. J.
Invertebr. Reprod. Dev. *14*, 131–151.

Souty-Grosset, C., Badenhausser, I., Reynolds, J.D., and Morel, A. (2005). Investigations
on the potential of woodlice as bioindicators of grassland habitat quality. Eur. J. Soil Biol. *41*,
109–116.

Strachan, S.R., Chester, E.T., and Robson, B.J. (2015). Freshwater invertebrate life history
strategies for surviving desiccation. Springer Sci. Rev. *3*, 57–75.

Suzuki, S., and Ziegler, A. (2005). Structural investigation of the female genitalia and sperm-
storage sites in the terrestrial isopod *Armadillidium vulgare* (Crustacea, Isopoda). Arthropod
Struct. Dev. *34*, 441–454.

Tang, X., and Pikal, M.J. (2005). Measurement of the kinetics of protein unfolding in viscous
systems and implications for protein stability in freeze-drying. Pharm. Res. *22*, 1176–1185.

Therneau, T. (2022). A Package for Survival Analysis in R. R package version 3.3-1,
https://CRAN.R-project.org/package=survival.

Trenberth, K.E., Dai, A., van der Schrier, G., Jones, P.D., Barichivich, J., Briffa, K.R., and
Sheffield, J. (2014). Global warming and changes in drought. Nat. Clim. Change *4*, 17–22.

Verberk, W.C.E.P., Siepel, H., and Esselink, H. (2008). Life-history strategies in freshwater
macroinvertebrates. Freshw. Biol. *53*, 1722–1738.