1

Targeting neuroinflammation by pharmacologic down-regulation of inflammatory pathways is neuroprotective in protein misfolding disorders.

Sydney J. Risen[1, 5], Sean W. Boland[1,5], Sadhana Sharma[2], Grace M. Weisman[1], Payton M. Shirley[1], Amanda S. Latham[1], Arielle J.D. Hay[1], Vincenzo S. Gilberto[2], Amelia D. Hines[1], Stephen Brindley[3], Jared M. Brown[3], Stephanie McGrath[4], Anushree Chatterjee[2], Prashant Nagpal[2], and Julie A. Moreno[1, 4, 5]*

[1]Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO, USA

[2]Sachi Bioworks Inc., Colorado Technology Center 685 S Arthur Avenue, Louisville, Colorado 80027, United States

[3]Department of Pharmaceutical Sciences, University of Colorado Anschutz Medical Campus, Aurora, Colorado 80045, United States

[4]Clinical Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO, USA

[5]Brain Research Center, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO, USA

*Corresponding author. E-mail: julie.moreno@colostate.edu

Abstract

36
Neuroinflammation plays a crucial role in the development of neurodegenerative protein
misfolding disorders. This category of progressive diseases includes, but is not limited to,
Alzheimer's disease, Parkinson's disease, and prion diseases. Shared pathogenesis involves
the accumulation of misfolded proteins, chronic neuroinflammation, and synaptic dysfunction,
ultimately leading to irreversible neuronal loss, measurable cognitive deficits, and death.
Presently, there are little to no effective treatments to halt the advancement of
neurodegenerative diseases. We hypothesized directly targeting neuroinflammation by
downregulating the transcription factor, NF-κB and the inflammasome protein, NLRP3, with the
brain-penetrant, non-toxic, SB_NI_112, would be neuroprotective. To achieve this, we used a
cocktail of RNA targeting therapeutics (SB_NI_112) shown to be brain-penetrant, non-toxic,
and targeting both NF-κB and NLRP3. We utilized a mouse-adapted prion strain as a model for
neurodegenerative diseases to assess aggregation of misfolded proteins, glial inflammation,
neuronal loss, cognitive deficits, and lifespan. Prion-diseased mice were treated either
intraperitoneally or intranasally with SB_NI_112. Behavioral and cognitive deficits were
significantly protected by this combination of NF-κB and NLRP3 down-regulators. Treatment
reduced glial inflammation, protected against neuronal loss, prevented spongiotic change,
rescued cognitive deficits, and significantly lengthened lifespan of prion-diseased mice. We
have identified a non-toxic, systemic pharmacologic that down-regulates NF-κB and NLRP3,
prevents neuronal death and slows the progression of neurodegenerative disease. Though
mouse models do not always predict human patient success, and the study was limited due to
sample size and number of dosing methods utilized, these findings serve as a proof of
principle for continued translation of the therapeutic SB_NI_112 for prion disease and other
neurodegenerative diseases. Based on success in a murine prion model, we will be continual
testing SB_NI_112 in a variety of neurodegenerative disease models, including Alzheimer's
Disease and Parkinson's Disease.

62
63 **Keywords**
64 Neuroprotection, Neuroinflammation, neurodegeneration, inflammasome, prion disease

65
Introduction
Chronic inflammation in the brain has emerged as a significant contributor to
neurodegenerative disease pathogenesis and progression. Alzheimer's disease (AD),
Parkinson's disease (PD), and prion diseases share many common characteristics;
accumulation and aggregation of misfolded proteins, glial activation, chronic
neuroinflammation, neuronal loss, cognitive deficits, and death[1–5]. These diseases can be
placed under the broader category of neurodegenerative protein misfolding disorders
(neurodegenerative diseases) and are associated with the presence and aggregation of
misfolded amyloid-β (Aβ) in AD, alpha synuclein (α-Syn) in PD, or the prion protein (PrP[Sc]) in
the brain, which lead to behavioral deficits and subsequent progressive neuronal death[6]. Many
disease modifying treatments have been directly targeted toward the aggregated proteins
associated with such disease, though none have been successful in stopping disease
progression[7–9]. Innate immune activation of microglia and astrocytes appears to be particularly
important in the context of neurodegenerative diseases[10–12]. Furthermore, new evidence
suggests that the compounded effect of misfolded proteins leads to further glial and astrocyte
activation[6,13–15]. Therefore, chronic neuroinflammation has been identified as a promising

therapeutic target for neurodegenerative diseases. However, the only current interventions
include anti-inflammatory steroids, but their effectiveness as neurotherapeutic
countermeasures has not yet been established physiologicaly[16,17].
During neurodegenerative disease-induced brain stress, there is significant crosstalk between
microglia, astrocytes, and neurons via critical inflammatory cell signaling events involving
initiation of the nuclear factor kappa B (NF-κB) pathway and subsequent NOD (nucleotide-
binding oligomerization domain), LRR (leucine rich repeat), and pyrin-containing 3 (NLRP3)
inflammasome formation[18–20]. NF-κB, when translocated into the nucleus, promotes
transcription of several inflammatory cytokines and chemokines, including NLRP3, pro-
interleukin-1β (pro-IL-1β), and pro-interleukin-18 (pro-IL-18), triggering a cascade of microglial
and astrocytic inflammation[21]. These reactive astrocytes are directly toxic to neurons and
endothelial cells during prion disease[22,23]. Many recent studies, including human patient, *in*
vitro, and *in vivo* animal models, have pointed to the assembly of the NLRP3 inflammasome in
microglia and astrocytes as a significant contributor to inflammatory-related amplification of
misfolded proteins in neurodegenerative diseases[24–31]. Formation of the NLRP3 inflammasome
is a critical step for activation of pro-inflammatory cytokines and perpetuation of the immune
response through microglial pyroptosis[32,33]. Pro-caspase-1 is activated by proximity-induced
autocatalytic activation upon recruitment to the NLRP3 inflammasome. Active caspase-1
cleaves pro-IL-1β and pro-IL-18 into their mature and biologically active forms, cytokines IL-1β
and IL-18[32,34]. In mouse models of APP/PS1 mutation and a tauopathy, NLRP3 inflammasome
assembly has been identified as a significant contributor to inflammatory-related amplification
of Aβ plaques and neurofibrillary tangles[24,25]. Similarly, the NLRP3 inflammasome promotes α-
Syn production in mouse models of PD[26,27]. Activation of NF-κB in prion disease augments
neuronal death by increasing the expression of pro-apoptotic genes and triggering the release
of cytochrome C, resulting in mitochondrial dysfunction and endoplasmic reticulum stress[35].
Additionally, the upregulation of inflammatory cytokines IL-1β and tumor necrosis factor (TNF),
formerly known as tumor necrosis factor alpha (TNFα), at sites of synaptic damage trigger the
production of reactive oxygen species including hydrogen peroxide and nitric oxide, leading to
excitotoxicity and neuronal death[36–38]. Beyond induction of inflammation and propagation of
misfolded proteins, it has been proposed that activation of the NLRP3 inflammasome during
NMPDs inhibits microglial autophagy, dampening the identification and degradation of
misfolded protein aggregates in AD, PD, and prion disease[39–41].
114
Based on the extensive evidence supporting central involvement of NLRP3 inflammasome
formation following NF-κB activation in the progression of neurodegenerative diseases, it is
important to develop potential therapeutics to inhibit NF-κB and NLRP3. Therapeutic
intervention to target and inhibit NLRP3 inflammasome formation arises as a potential three-hit
treatment against neurodegenerative diseases by: 1) reducing inflammatory signaling amongst
microglia and astrocytes[18–20], 2) preventing propagation of misfolded proteins[24–31], and 3)
upregulating microglia autophagy to accelerate removal of misfolded proteins[39–41]. No studies
to date have explored combined NF-κB and NLRP3 inhibition as a therapeutic intervention for
neurodegenerative diseases.
In the present study, we utilized non-toxic, brain-penetrant Nanoligomers designed to
downregulate NF-κB and NLRP3 by blocking their translation[42] (Figure 1A). Nanoligomers are
stable, non-toxic, and bioavailable, with the capability to cross the blood brain barrier (BBB)
through various routes of administration (Figure 1A)[43]. A previous study displayed effective

gene regulation *in vivo* with a similar Nanoligomer therapeutic aimed at down-regulation of NF-
κB and tumor necrosis factor receptor 1 (TNFR1) in an LPS-induced murine model of
neuroinflammation[42]. The combined NF-κB and NLRP3 down-regulator, SB_NI_112, displays
high target specificity and minimal off-targeting[42]. To study the impact of targeting NLRP3
inflammasome formation as a therapeutic for neurodegenerative diseases, we utilized a mouse
model of infectious prion disease in wild-type (C57BL/6) mice. A prion model allows for study
of a naturally occurring neurodegenerative diseases without the need for transgenic mice.
Prion diseases are rare neurodegenerative diseases that result from the native conformation of
the cellular prion protein (PrP^C) misfolding to the infectious form, PrP^{Sc} [44,45]. All prion diseases
display the common characteristics of neurodegenerative diseases, including the accumulation
and aggregation of a misfolded protein (PrP^{Sc}), glial activation, chronic inflammation, neuronal
loss, cognitive deficits, and death[42,43]. Previous literature has shown that therapeutics
developed for prion disease mice can be translated into other neurodegenerative diseases,
though the few compounds shown to reduce signs of prion disease in mouse models are
associated with several negative side effects[36]. In this study, found treatment with SB_NI_112
to down-regulate NF-κB and NLRP3 reduced glial inflammation, protected against neuronal
loss, prevented spongiotic change, rescued cognitive deficits, and significantly lengthened
lifespan of prion-diseased mice. These findings, if validated clinically, have important
implications for improving the quality of life for patients suffering with neurodegenerative
diseases.
148
Results & Discussion
Attempts to slow the progression of neurodegenerative diseases have proven to be difficult.
Many promising treatments are toxic or must be delivered directly to the brain, making
translation to human patients difficult[7,36,46,47]. Neurodegenerative diseases have highly
complex etiology, making the root cause, or multiple causes, difficult to identify and target.
Neuroinflammation has arisen as a promising target, as chronic stimulation of the brain's
immune system leads to increased protein misfolding and neuronal death[6,11,15]. The NF-κB
pathway and subsequent NLRP3 inflammasome formation have been identified as key
contributors to chronic neuroinflammation present in neurodegenerative diseases[30,31,35,37].
Here, we employed a mouse prion model to test the efficacy of a combined NF-κB and NLRP3
down-regulator for neurodegenerative disease treatment.
160
Wild-type, C57Bl6/J, (Jackson Laboratories) mice were infected intracerebrally with 0.1% prion
brain homogenates from Rocky Mountain Laboratories (RML) or normal brain homogenates
(NBH) derived from C57Bl6/J mice as the controls. SB_NI_112 treatment began at 10 weeks
post-inoculation (wpi) or approximately 40% of prion disease progression and continued until
the mice succumbed to the disease (Figure 1B). As the disease progressed and treatment
continued, behavioral assessments including novel object recognition, burrowing, and nesting,
were performed at 12 wpi to train and establish a baseline for each experimental cohort, then
continued throughout the study. All three behavioral assessments measure hippocampal
integrity, the brain region known to show the first clinical pathology including glial inflammation
and spongiotic change[48,49].
171
SB_NI_112 penetrates the blood-brain barrier and is systemically non-toxic.

With a focus on reducing inflammation in the brain, it was important to assess whether
SB_NI_112 effectively crosses the blood brain barrier (BBB). Biodistribution and toxicity
testing was completed in naïve C57BL/6 female and male mice. Penetration of SB_NI_112
across the blood-brain barrier was confirmed with ICP-MS and detected in the hippocampus
(76.1 nM +/- 44.3), cortex (49.9 nM +/- 11.4), cerebellum, (96.9 nM +/- 45.0) and brainstem
(96.6 nM +/- 32.4) (Figure 2A), which is significantly higher than the dissociation constant of
SB_NI_112[50].

Down-regulation of NF-κB and NLRP3 with SB_NI_112 leads to dampening of chronic
inflammation while preserving key immune pathways needed to respond to infectious
pathogens (63). Despite down-regulation of NF-κB and NLRP3, interferon gamma (IFN-γ)
expression was significantly higher in prion infected mice treated with SB_NI_112 (p = 0.048
(2-way t-test, Excel) or 0.0468 (One-way ANOVA, Graphpad Prism)), and not significantly
different from control NBH mice (Figure 2B). This shows the use of SB_NI_112 specifically
interferes with the NF-κB and NLRP3 pathway but does not alter expression of immune
signaling molecules essential for response to infections.

Both female and male naïve C57BL/6 mice treated with a single dose of SB_NI_112 at 500
mg/kg intraperitoneally (i.p.) had no weight loss 1 week after dosing (Figure 2C) and displayed
no morphologic toxicity in the colon (Figure 2C, F), kidney (Figure 2D, G), or liver (Figure 2E,
H) via hematoxylin and eosin staining. No immune cell infiltration was identified in these
organs and the parenchymal cells were intact. Based on previous studies completed with
similar therapeutics[42,43], we chose to dose mice conservatively at 10 mg/kg two times per
week. Prion-diseased mice were split into 3 treatment groups: vehicle i.p., SB_NI_112,
intranasal (i.n.), and SB_NI_112 i.p.

To confirm therapeutic SB_NI_112 successfully inhibits both NF-κB and NLRP3, we performed
western blots to analyze the ratio of active NF-κB (phosphorylated p65; p-p65) compared to
inactive form (p65) and NLRP3 protein expression. Interestingly, we identified a higher
success of inhibition with intraperitoneal delivery than intranasal. SB_NI_112 provided
significant reduction of p-p65 in prion disease mice compared to untreated counterparts
(Figure 3A, B, and D: p-value = 0.0333, Figure S3). Intraperitoneal treatment also significantly
reduced NLRP3 expression in prion-diseased mice compared to untreated counterparts
(Figure 3C, E; p-value = 0.0380, Figure S3). This difference in effectiveness is likely due to
inconsistency of intranasal delivery compared to the intraperitoneal method. The higher
success of SB_NI_112 dosed intraperitoneally to inhibit NF-κB and NLRP3 is supported by the
following behavioral and histological results obtained.

It is important to note that though many treatments for neurodegenerative diseases have been
promising in early research, many fail due to toxicity and low efficacy rates in animal models or
human trials. At this time, our research has identified no measurable toxicity or side effects
associated with Nanoligomer SB_NI_112 at single high doses of 500 mg/kg or repeated dosing
of 150 mg/kg. Additionally, the NF-κB and NLRP3 down-regulator is stable and bioavailable,
has confirmed ability to cross the BBB, and displays high target specificity with minimal off-
targeting. Our mechanism of action results (Figure 3) displaying superior inhibition of NF-κB
and NLRP3 via i.p. route of delivery compared to i.n., combined with behavioral and

histological data, support that an intraperitoneal dosing method is more successful at delivery
therapeutic SB_NI_112 to the brain, thus providing better protection against neuroinflammation
and prion disease progression.
222
A secondary study was completed at 150 mg/kg three times per week to explore the impact of
a higher dose. However, we saw no difference in RML prion associated behavioral
assessment scores and clinical signs of 10 mg/kg 2x per week and high dose 150 mg/kg 3x
per week treatment groups (Figure 3 and Supplemental Figure 1 respectively), indicating there
may be a threshold at which the treatment against neuroinflammation, specifically NF-κB and
NLRP3, is effective against RML prion disease. Because there is no indication to treat with a
higher dose, as it does not increase the therapeutic effect, we did not continue with additional
methods to assess disease progression in the high dose study.
231
Prion-induced behavioral and cognitive deficits occur significantly later in mice treated
with NF-κB and NLRP3 down-regulator. Down-regulation of NF-κB and NLRP3 significantly
protects against behavioral and cognitive changes associated with RML treated prion disease.
RML mouse-adapted prion strain preferentially seeds and accumulates in the
hippocampus[48,49]. Two common hippocampal behaviors affected by RML prion disease include
nesting and burrowing activity[51,52]. To assess nesting, mice were given napkins to nest in
overnight and scored between one (no nest formed) and five (fluffy built nest) weekly starting
at 16 wpi. Monitoring of nest formation was found to be significantly higher among i.p. treated
mice at 20 wpi (average score 4.5, p = 0.0001), 21 wpi (average score 3.2, p = 0.0007), and 22
wpi (average score, 2.5, p = 0.0216) (Figure 4A) compared to vehicle-treated prion-infected
(average score 1 at 20 wpi, and 0.5 at 21 and 22 wpi) nest formation, while i.n. treated mice
show a trend though no significant change in nesting throughout the disease. Similarly, a
significant increase in the ability of the mice to build proper nests at 19 wpi (average score 4.5,
p = 0.0039) and 21 wpi (average score 3.2, p = 0.0005) was shown with high dose i.p.
treatment with SB_NI_112 compared to vehicle treatments (average score 2.5, p = 0.0039 at
19 wpi and 2.4, p = 0.0004 at 21 wpi) (Supplemental Figure 1A).
248
Mice were given a plastic tube filled with food pellets within their home cages for 30 min to
burrow freely. Following this time, the food pellets left in the tubes were weighed and the
percentage burrowed was calculated. Prion-diseased mice are known to stop burrowing as the
disease progresses[28,51,52] but at 20 wpi, 21 wpi, and 22wpi, prion-diseased mice given
therapeutic SB_NI_112 (i.n. and i.p.) had a later onset of deficits in burrowing activity (Figure
4B), showing a significance of protection at 20 wpi (average percent burrowed 47, p = 0.0475
for i.n.; average percent burrowed 42, p = 0.0032 for i.p.), trending through 22 wpi. (Figure 4B)
A significant difference in the hippocampal burrowing assay was only indicated at 16 wpi
(average percent burrowed was 75%, p > 0.0365) when mice were treated with the high dose
SB_NI_112 compared to the vehicle (average percent burrowed was 51%) (Supplemental
Figure 1B).
260
Cognitively, mice with RML-induced prion disease have a significant decrease in the ability to
recognize a familiar object as the RML prion strain begins to aggregate within the
hippocampus, the brain region essential for memory and learning[48,51,52]. Both i.n. and i.p.
SB_NI_112 treatment protected prion-induced deficit in novel object recognition when

compared to vehicle-treated diseased mice (Figure 4C, D). The ratio of time the mouse spent
with the novel object to the familiar object (seen 24 hours beforehand) was calculated and
noted by the discrimination index. The closer the discrimination index (DI) is to a value of one
the more intact the hippocampal memory is, implying lower disease progression in the
animal[53]. At 20 wpi, only SB_NI_112 i.p. showed a significantly higher discrimination index
compared to untreated counterparts (mean discrimination index score of 0.4975, p = 0.0265),
with a trend of improvement with low dose i.n. SB_NI_112 (mean discrimination index score of
0.1974, p =0.4383) (Figure 4C). At 22 wpi, both i.p. (mean discrimination index score of
0.4491, p = 0.0008) and i.n. treated mice (mean discrimination index score of 0.4343, p =
0.0022) had a significant increase when compared to vehicle-treated diseased mice with a
discrimination index of -0.174 (i.e., preferring the familiar object) (Figure 4D). There was also a
trend towards improved novel object recognition in high dose treated mice compared to
untreated counterparts at 22 wpi (Supplemental Figure 1D).
278
Figure 5. Microglial inflammation is significantly decreased by NF-κB and NLRP3 down-
regulation with SB_NI_112 in prion diseased mice at 20 wpi. Representative images of the
hippocampus, thalamus, cortex, and cerebellum with Iba1+ cells, a marker of activated
microglia, with vehicle (A-D), SB_NI_112 i.n. (E-H), and SB_NI_112 i.p. (I-L) treatment.
Quantitative analysis of each brain region identifies a significantly lower number of Iba1+ cells
in the hippocampus (M), thalamus (N) cortex (O), and cerebellum (P) with both i.n. and i.p.
treatment. Arrows represent examples of positive cells. Box insets show zoomed-in image of
single microglia. Scale bar = 20μm, N=3-4. One-way ANOVA, error bars = SEM, *p > 0.05,
**p> 0.001, ***p > 0.0001.
288
Figure 6. Astrocytic inflammation is significantly decreased by NF-κB and NLRP3 down-
regulation with SB_NI_112 in prion disease mice at 20wpi. Representative images of the
hippocampus, thalamus, cortex, and cerebellum with S100®+ cells, a marker of activated
microglia, with vehicle (A-D), SB_NI_112 i.n. (E-H), and SB_NI_112 i.p. (I-L) treatment.
Quantitative analysis of each brain region identifies a significantly lower number of S100®+
cells in the thalamus (N) and a trend of a decrease of inflamed astrocytes in the hippocampus
(M) and cortex (O), and cerebellum (P) with treatment. Arrows represent examples of positive
cells. Box insets show zoomed-in image of single astrocytes. Scale bar = 20μm, N=3-4. One-
way ANOVA, error bars = SEM, * p< 0.05.
298
Down- regulation of NF-κB and NLRP3 with SB_NI_112 treatment reduces glial
inflammation.
Brain tissues were collected at 20 wpi and 24 wpi from low dose treatment groups. Formalin-
fixed tissues were processed and embedded into paraffin wax and sectioned at 5μm. Brains
were stained for both microglial and astrocytic inflammation (Figure 5-6). Microglia
inflammation was identified as Iba1+ cells (Figure 5), and S100β+ cells denote inflamed
astrocytes (Figure 6). Four main brain regions known to be affected in RML prion infection
were analyzed: the hippocampus, cortex, thalamus, and cerebellum. Representative images of
each brain region for all treatment groups are displayed for both microglial (Figure 5A-L) and
astrocytic gliosis (Figure 6A-L), and arrows point out examples of positively stained cells.
309

In the thalamic region, both microglial and astrocytic inflammation were significantly
suppressed by down-regulation of NF-κB and NLRP3 with SB_NI_112, both i.n. and i.p.
(Figure 5N and 6N respectively) at 20 wpi. Within the thalamic region, vehicle treated samples
had a mean of 775.4 Iba1+ microglia per mm^2, followed by SB_NI_112 i.p. treatment with a
mean of 168.6 Iba1+ microglia per mm^2 (p = 0.0115 compared to vehicle), and SB_NI_112 i.n.
treatment with a mean of 168.6 Iba1+ microglia per mm^2 (p = 0.0087 compared to vehicle). In
regard to astrogliosis in the thalamus, vehicle treated samples had a mean of 421 S100β+
astrocytes per mm^2, followed by SB_NI_112 i.p. treatment with a mean of 270 S100β+
astrocytes per mm^2, (p = 0.0613 compared to vehicle), and SB_NI_112 i.n. treatment with a
mean of 240.8 S100®+ astrocytes per mm^2, (p = 0.0282 compared to vehicle). Within the
hippocampal region, vehicle treated samples had a mean of 246 Iba1+ microglia per mm^2,
followed by SB_NI_112 i.n. treatment with a mean of 56.31 ba1+ microglia per mm^2 (p =
0.0019 compared to vehicle), and SB_NI_112 i.p. treatment with a mean of 36.75 Iba1+
microglia per mm^2 (p = 0.0010 compared to vehicle). In the cortex, vehicle-treated samples
had a mean of 203 Iba1+ microglia per mm^2, followed by SB_NI_112 i.n. treatment with a
mean of 33.22 Iba1+ microglia per mm^2 (p = 0.0014 compared to vehicle), and SB_NI_112 i.p.
treatment with a mean of 22.32 Iba1+ microglia per mm^2 (p = 0.0010 compared to vehicle).
Lastly, in the cerebellum vehicle-treated samples had a mean of 319.6 Iba1+ microglia per
mm^2, followed by SB_NI_112 i.n. treatment with a mean of 80.05 Iba1+ microglia per mm^2 (p =
0.0023 compared to vehicle), and SB_NI_112 i.p. treatment with a mean of 57.71 Iba1+
microglia per mm^2 (p = 0.0013 compared to vehicle). There was a trend towards astrocyte
reduction for both i.p. and i.n. treatment groups in the hippocampus (Figure 6M), Cortex
(Figure 6O), and cerebellum (Figure 5P).
333
Microglia are closely associated with plaques in multiple forms of human prion disease
including Creutzfeldt-Jakob disease, Gerstmann-Straussler-Scheinker, and kuru and it has
been shown that microglia play a role in neuronal cell death during prion disease[54,55].
Microglial activation appears before the manifestation of clinical signs of prion disease,
indicating microglial inflammation plays a role in the progression of prion disease. Furthermore,
the deposition of prion proteins alone does not directly predict regions of neurodegeneration[55].
In addition to microglial activation, astrogliosis is found early in prion disease pathology, and
reactive or inflamed astrocytes are known to be synaptotoxic in the pathogenesis of prion
disease[22,23,56,57]. Further, NF-κB and NLRP3 activation are implicated in the induction of
neurotoxic astrocytes by microglia. Global microglial ablation of NLRP3 ameliorates A1-like
astrocytes in vitro and in vivo, decreasing neuronal dysfunction. Our research supports this
claim by displaying decreased microglial and astrocytic inflammation down-regulating NLRP3
and NF-κB with brain-penetrant SB_NI_112 (Figure 5-6). Though a less significant difference
was seen in inflamed astrocytes (S100β+ cells) between treated and untreated groups, this
may be due to temporal differences in microglial and astrocyte activation throughout chronic
inflammation or a proposed cell-type specific activation of NLRP3 in microglia compared to
astrocytes[58,59].
351
Figure 7. Prion induced neuronal loss and spongiotic change is significantly decreased
by NF-κB and NLRP3 down-regulation with SB_NI_112 in prion diseased mice.
Representative images of the NeuN + cells (neuronal cell bodies) in the CA1 region of the
hippocampus (A-C) Significant decrease in the number of neurons lost at 20 wpi with treatment

via i.p. and i.n. SB_NI_112 (D). Scale bar = 20mm, N=3-4. One-way ANOVA *p > 0.05.
Representative images of the spongiotic change in hippocampus, thalamus, cortex, and
cerebellum with vehicle (E-H)), low dose SB_NI_112 i.n. (I-L), and low dose SB_NI_112 i.p.
(M-P). Pathological scoring of each brain region identifies a protection or decrease of
pathological score at all brain regions analyzed; + = mild spongiosis, ++ = moderate
spongiosis, +++ = severe spongiosis. N=3-4.
362
SB_NI_112 protects against prion-induced spongiosis and neuronal loss. Morphologic
spongiosis or vacuoles within the brain tissue is a hallmark of prion disease that increases as
the disease progresses[44,45]. To assess and quantify this change, we used pathological scoring
of the hippocampus, thalamus, cortex, and cerebellum following H&E staining. Scores were
given ranging from + (mild spongiosis) to +++ (high spongiosis) by three blinded researchers
Figure 6E-P. This revealed protection against spongiform change in both i.p. and i.n.
SB_NI_112 treatment of prion diseased mice when compared to vehicle-treated prion
diseased mice at 20 wpi. Representative images from each brain region with corresponding
average spongiosis score are displayed for RML + Vehicle, RML + SB_NI_112 i.n., and RML +
SB_NI_112 i.p. Overall, down regulation of NF-κB and NLRP3 with SB_NI_112 protects
against hallmark spongiosis in prion disease. Beyond spongiosis, the loss of neurons within
the hippocampus following RML prion infection is well established[48,49]. Therefore, we
quantified neuronal loss in SB_NI_112 treatment groups to untreated counterparts. Neuronal
numbers within the CA1 region of the hippocampus (Figure 7A-C) were significantly higher
with down-regulation of NF-⊦B and NLRP3 by low dose i.n. and i.p. delivered SB_NI_112
compared to the vehicle (Figure 7D, p>0.05). Critically, SB_NI_112 prevents the neuronal loss
and spongiform change induced by RML prion infection. As neuronal loss is the direct cause of
behavioral and cognitive deficits, and ultimate fatality due to prion disease, it is imperative
potential therapeutics show success in this metric[49].
382
Figure 8. Prion diseased mice treated with NF-κB and NLRP3 down-regulator had
significantly lower clinical scores and increased lifespan, independent of PrP^{Sc}
accumulation. Clinical scores of mice treated with i.n. or i.p. SB_NI_112 at 22 and 23 wpi
were significantly lower than vehicle treated counterparts, continued significance at 24 wpi with
i.p. SB_NI_112 (A). N=9 for prion positive i.p. and i.n. groups, N=9-12 for all other groups.
One-way ANOVA, error bars = SEM, * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001.
Mice treated with low dose SB_NI_112 i.p. lived significantly longer than untreated
counterparts (B) Survival curve, ***** (p <0.000001). Total PrP and PrP^{Sc} levels detected after
proteinase K (PK) (indicates not added, + indicates added) digestion at 20 wpi measured by
immunoblot (C). Total PrP relative to β-actin was significantly different among SB_NI_112 i.p.
treated and untreated groups (D). N=4 for all groups. One-way ANOVA, error bars = SEM, * p
< 0.05. No significant difference was seen in total PrP^{Sc} relative to PrP^{C} (E). Total PrP^{Sc}
relative to PrP^{C} was equivalent in prion-infected mice treated with vehicle and SB_NI_112 i.n.
and i.p. treatment (E). N=4 for all groups. One-way ANOVA, error bars = SEM. All
immunoblots were performed on frontal cortex lysates.
398
Treatment with SB_NI_112 improved clinical scores and survival in mice despite
accumulation of PrP^{Sc}. At 22 wpi, vehicle-treated mice begin to show a significant increase in
clinical signs of prion disease (Figure 8A). From 22 wpi to 24 wpi i.p. SB_NI_112 treated

diseased mice have significantly (p<0.0001 for all weeks) lower clinical scores than vehicle-
treated diseased mice, and SB_NI_112 i.n. treatments displayed significantly lower scores at
22 wpi (p <0.0001) and 23 wpi (p = 0.0197). Untreated Prion clinical signs included tail rigidity,
hyperactivity, ataxia, extensor reflex (clasping), tremors, righting reflex, kyphosis, and poor
grooming[28,48,52]. Each sign was rated on a scale from 0-5. All clinical sign scores were
combined for a total score. SB_NI_112 i.p. treated mice had clinical scores with a mean lower
than 2 at 22 wpi, 3 at 23 wpi, and 5 at 24 wpi, compared to clinical score mean of 6+ points
higher in untreated counterparts (Fig 8A). Mice receiving i.n. treatments averaged scores of 3
at 22 wpi and 7 at 23 wpi (Fig 8A). Of note, clinical scores in high dose treated mice were
significantly lower (p = 0.0120) at 22 wpi, with a trend towards lower scores at 23 and 24 wpi
(Supplemental Figures 1E). See supporting videos for visual of clasping (supporting videos 1-
3), righting reflex (supporting videos 4-6), and walking (supporting videos 7-9) in Control
(NBH), RML + vehicle, and RML + Sb_NI_112 i.p. at 22 wpi. Mice were considered terminal
and euthanized after reaching a total score of 9 or above. Additionally, mice with a weight loss
greater than 10% were considered terminal and euthanized. The date of euthanasia was
documented for survival analysis (Figure 8B).
418
The lifespan of prion disease mice was significantly increased with i.p. SB_NI_112
Nanoligomer treatment (p <0.000001) compared to vehicle-treated mice (Figure 8B). No
change or protection of i.n. delivered SB_NI_112 was seen in survival of prion-diseased mice.
Human sporadic Creutzfeldt-Jakob disease has a mean survival of 7-8 months after
diagnosis[60]. On average, i.p treatment with SB_NI_112 significantly increased prion-diseased
mouse lifespan. Untreated mice had a mean survival of 162 days, i.p. SB_NI_112 treated 174
days, and i.n. SB_NI_112 163 days. Of note, the first mouse without SB_NI_112 treatment
reached a clinical score indicative of sacrifice 15 days before the first mouse receiving i.p.
treatment. 15 mouse days equate to approximately 6 months in human years, a near doubling
of expected survival time after diagnosis[61]. Importantly, there was no toxic pathology induced
by the down regulation of NF-ⱡB and NLRP3 by SB_NI_112 (Figure 2A).
430
To determine if SB_NI_112 modulates the accumulation of the misfolded prion protein, PrP^{Sc},
immunoblotting was performed on cortical brain homogenates for levels of the cellular PrP^{C}
and proteinase K-resistant PrP^{Sc} (Fig 8C, Figure S4). Total PrP relative to β-actin was
significantly different among treated and untreated groups (Figure 8D). No significant
difference was seen in total PrP^{Sc} relative to PrP^{C} (Figure 8E, Figure S4). Similarly, other
studies examining potential prion therapeutics have shown similar results in regard to total
PrP^{Sc}[48,49]. This indicates dampening chronic inflammatory signaling protects against prion-
associated neurodegeneration independent of PrP^{Sc} accumulation. SB_NI_112, down-
regulator of NF-κB and NLRP3, shows efficacy in slowing the disease clinically, behaviorally,
and pathologically while increasing predicted survival after diagnosis, and use as a
combination therapy for individuals diagnosed with prion disease.
442
NLRP3 inflammasome involvement in progression of prion disease. The level of influence
the NLRP3 inflammasome has on prion disease progression is still debated within the
literature. The NLRP3 inflammasome was found to be involved in lipopolysaccharide (LPS)-
primed microglia after exposure to a synthetic neurotoxic prion fragment in vitro[29,30]. Previous
studies highlight the importance of the NLRP3 inflammasome in the production of IL-1β and IL-

18 in systemic inflammation and neurodegenerative disease, including prion disease[21,24–26,62].
An *in vitro* study in BV2 microglia and co-cultured glia reported that aggregated prion protein
was involved in both priming and activation of the NLRP3 inflammasome[29].
451
Regarding cytokine production in our study, at 16wpi, IL-28 is significantly reduced in i.p.
treated (average 30.1 pg/mg total protein, p <0.05) mice compared to untreated counterparts
(average 43.4 pg/mg total protein) (Figure S2). Interestingly, an *in vivo* study found levels of IL-
1β at end-stage prion disease were not affected by the absence of NLRP3 and that NLRP3
knockout mice succumbed to RML prion in a similar time frame to wild-type counterparts[63].
Conversely, our *in vivo* research supports the involvement of the NLRP3 inflammation in prion
disease progression, as downregulation of NLRP3 and upstream signaling molecule NF-κB
decreased glial inflammation, protected against neuronal loss, prevented spongiotic change,
rescued cognitive deficits, and significantly lengthened lifespan of prion-diseased mice.
Additionally, inflammatory cytokines IL-2 (Figure S2 B), IL-1β (Figure S2 C), and IL-1α (Figure
S2 D), trended lower in prion-diseased mice treated with i.p. SB_NI_112, like NBH controls,
when compared to untreated counterparts. Principal component analysis revealed trending
differences between treated and untreated prion-diseased mice (p = 0.094) (Figure S2 E).
Based on conflicting data, there are likely additional inflammatory pathways upregulated by
PrP^{Sc} in conjunction with the known direct degenerative impact PrP^{Sc} aggregates have on
neurons[64].
468
Limitations of our study include a low sample size at each time point. Higher significance may
have been achieved with a larger sample size, particularly in regard to later behavioral testing
timepoints. Moreover, adding more behavioral time points (weekly vs. biweekly) could display
more detailed changes in behavior as prion disease progresses. Additionally, there may be
other routes of dosing that provide stronger protection against prion disease progression,
including intravenous, subcutaneous, or oral routes. Finally, because the therapeutic showed
high safety efficacy, it would be beneficial to test a variety of difference dose concentrations
and frequencies.
477
Conclusions and Future Prospects
Down-regulation of NF-κB and NLRP3 in a mouse-adapted RML prion model slows disease
progression by reducing glial inflammation, delaying development of behavioral deficits and
clinical signs, preventing loss of neurons, and extending lifespan. Because treatments
successful in prion disease are anticipated to work in other neurodegenerative diseases due to
common pathogenesis, these findings, if validated clinically, have important implications for
patients diagnosed with prion disease and those suffering from more widespread
neurodegenerative diseases like AD and PD.
486
Though all mice eventually succumbed to disease in this study, PrP^{Sc} misfolding and
accumulation in prion disease, in comparison to other misfolded proteins in neurodegenerative
diseases (Aβ in AD and α-Syn in PD), is substantially more rapid and aggressive[65].
Additionally, properly functioning microglia can phagocytose and degrade both Aβ and α-Syn.
It is only when microglia lose the ability to degrade misfolded proteins and the load of Aβ or α-
Syn becomes too high that aggregation and progression of disease state occurs in AD and PD
respectively[24–27]. Recent research supports chronic inflammation and NLRP3 inflammasome

stimulation decreases microglial autophagy, leading to increased Aβ or α-Syn production and
aggregation in the brain[39–41]. Furthermore, inflammation in other glial cells, namely astrocytes,
leaves them unable to properly perform metabolic, structural, homeostatic, and neuroprotective
tasks[65]. Our study revealed down-regulation of NF-κB and NLRP3 decreases astrocyte
inflammation in addition to microglial inflammation.
499
Specifically in prion disease, there have been few small molecules, antibodies and other
compounds that show promise in treatment of disease pathogenesis. Therapeutic
oligonucleotides for the prion protein or PrP-lowering antisense oligonucleotides (ASO) have
proven to lower the toxic load of the disease-causing misfolded prion protein increasing
survival of infected mice. Critically this was irrespective of the strain or type of prion the mice
were exposed to but, only successful with the highly invasive intracerebroventricular injection,
a critical limitation of this treatment[66]. The use of antibodies as an immunotherapy and small
molecules against the prion protein has been studied for years and in rodents the progression
and acceleration of the toxic PrP^{Sc} is reduced, but unfortunately the protection of these do not
translate well to patients[47,67–72]. The development of human specific antibodies is showing
some efficacy in patients however the need for the ability to give the immunotherapy to
patients earlier, prior to severe disease states, seems to be essential[73].
512
Nanoligomers present benefits compared to other small-molecules, antibodies, and antisense
oligonucleotide (ASO) platforms. First, their efficient biodistribution and facile delivery across
the BBB and reaching various sections of the brain without using invasive administration
methods such as Intracerebroventricular injection used in other ASO based treatments.[66,74]
Second, lack of any immunogenic response, any adverse systemic effect or accumulation
makes these molecules preferable to other small molecules. This study focusing on a Prion
infection model, showed that Nanoligomers directed at the inflammasome demonstrate
neuroprotection, improved animal survival rates, and enhanced cognitive function. This
underscores the technology's safety and broader potential applications. Another significant
advantage of the Nanoligomers discovery engine lies in its capability to selectively target
specific gene(s) for either upregulation or downregulation of expression as needed. Notably,
recent research demonstrated that a low dosage of 5 mg/kg of Nanoligomer significantly
reduced inflammation in the mouse hippocampus during LPS-induced neuroinflammation.[42]
526
Limitations of our study include a low sample size at each individual time point. Higher
significance may have been achieved with a larger sample size, particularly regarding
behavioral testing timepoints for the later more diseased timepoints. Moreover, adding more
behavioral time points (weekly vs. biweekly) could display more detailed changes in behavior
as prion disease progresses. Additionally, there may be other routes of dosing that provide
stronger protection against prion disease progression, including intravenous, subcutaneous, or
oral routes. Finally, because the therapeutic showed high safety profile, it would be beneficial
to test a higher dose concentration and more often frequency in dosing. In summary, targeting
the NLRP3 inflammasome via down-regulation of NF-κB and NLRP3 with SB_NI_112 has
promising therapeutic potential for neurodegenerative diseases. SB_NI_112 reduces
inflammatory signaling amongst microglia and astrocytes, which is highly neuroprotective. At
the least, targeting chronic neuroinflammation in neurodegenerative diseases is an essential

component in the "cocktail" of multi-target therapeutic interventions that are likely necessary for
successful modification and halting of disease.
541
542 **Methods**
543
Mice and Brain homogenates. C57Bl/6 (Jackson Laboratory) mice were intracranially
inoculated with 30µl of 0.1% 22L or Rocky Mountain Laboratories (RML) strains of mouse-
adapted prions, or normal brain homogenate (NBH), in 1% Pen-Strep (1:1 penicillin-
streptomycin). Mice were monitored for weight loss and clinical signs of prion disease and
euthanized after showing signs of terminal illness. 20% brain homogenates in phosphate-
buffered saline (PBS) were made using beads and a tissue homogenizer (Benchmark Bead
Blaster 24) and stored at -80C. Brain homogenates were aliquoted and treated with UV light
for 30 minutes to sterilize before being used for cell culture.
552
Nanoligomer Design and Synthesis. The Nanoligomer design, sequences, and synthesis
has been described in our previous work (59). Nanoligomers were designed and synthesized
(Sachi Bioworks Inc.) according to published methods[42,43,50,75,76]. The Nanoligomer is
composed of a nanobiohybrid molecule based on antisense peptide nucleic acid
(PNA)[42,43,75,77,78] conjugated to nanoparticle[1,2,5] for improved delivery. The PNA moiety was
chosen to be 17 bases long (to optimize solubility and specificity) antisense to the start codon
regions of nfkb1 (N terminus- -C terminus, sequence: AGTGGTACCGTCTGCTA) and nlrp3 (N
terminus- -C terminus, sequence: CTTCTACTGCTCACAGG) within the Mouse genome
(GCF_000001635.27).[1] Potential PNA sequences were screened for solubility, self-
complementing sequences, and off-targeting within the mouse genome (GCF_000001635.27).
The PNA portion of the Nanoligomer was synthesized on Vantage peptide synthesizer
(AAPPTec, LLC) with solid-phase Fmoc chemistry. Fmoc-PNA monomers were obtained from
PolyOrg Inc., with A, C, and G monomers protected with Bhoc groups. Following synthesis, the
peptides were conjugated with gold nanoparticles and purified via size-exclusion filtration.
Conjugation and concentration of the purified solution was monitored through UV-Vis (Thermo
Scientific, NanoDrop) measurement of absorbance at 260 nm (for detection of PNA) and 400
nm (for quantification of nanoparticles). Nanoligomers were stored at 4 °C until further use.
Biodistribution and Toxicity. Six female and six male C57Bl/6 (Jackson Laboratory) mice, 8-
10 weeks of age, were injected with a single dose of 500 mg/kg intraperitoneal or intranasal
SB_NI_112. Mice were monitored for 2 weeks for safety/tox study and dosing. Weight was
monitored daily, along with any changes in behavior/well-being. At the end of the study, mice
were euthanized, and tissues harvested for histopathological scoring, blood concentration of
any inflammatory markers, albumin content in serum. Using three-fold lower concentration
than the maximum tolerated dose (MTD) (150 mg/kg), eight female and eight male C57Bl/6
mice were dosed for 2 weeks with 2 doses/week. Mice were euthanized and harvested to
measure distribution of Nanoligomer through different tissues using ICP-MS.
579
Nanoligomer Administration. At 10 weeks post RML prion or NBH inoculation, mice were
treated two times per week either intraperitoneally (i.p.) or intranasally (i.n.) with 10 mg/kg of
SB_NI_112 diluted in sterile phosphate buffered saline (PBS). For high dose study, mice were
treated three times per week with 150 mg/kg of SB_NI_112 i.p. Mice receiving treatment i.n.
were anesthetized prior to administration and laid in an intranasal apparatus that controlled

isoflurane throughout the intranasal procedure.
586
Cognitive assay: Novel Object Recognition. Mice were tested in a rectangular arena (28cm
x 43cm) and habituated by exploring the empty arena for ten minutes. During day one of
testing, mice were placed in the arena with two identical objects (constructed with Legos) and
video monitored for 5 minutes of exploration. On day two of testing, one of the known objects
was replaced by a novel object, and the 5-minute exploration was filmed using a 1080P FHD
Mini Video Camera. All objects and the arena were cleansed thoroughly with 70% ethanol
between trials to ensure the absence of olfactory cues. Time exploring both the known and
novel object was calculated by blinded observers. Discrimination index (DI) calculated
following Absolute vs Relative Analysis protocol[53]. All objects were tested prior to use to
ensure the mice would explore them.
597
Hippocampal Behavioral assays: Burrowing and Nesting. Briefly, mice were placed in a
large cage with a PVC tube (2.33" x 5.75") full of food pellets, as describe[28,48,51]. The
percentage of burrowing activity is calculated from the difference in the weight of pellets in the
tube before and after 2 hours. To test nesting, we placed three fresh napkins in each cage.
After 24 hours, cages were examined for nesting activity. Nests were scored on a scale of 0-5,
where 0 represents no nesting activity, and 5 represents a high-quality nest.
604
Clinical Scoring of mice. Eight key clinical signs were monitored in mice daily beginning at 20
or 21 wpi. Clinical signs included tail rigidity, hyperactivity, ataxia, extensor reflex, tremors,
righting reflex, kyphosis, and poor grooming. Each sign was rated on a scale from 0-5. All
clinical sign scores were combined for a total score. Mice considered terminal and euthanized
after reaching a total score of 9 or above. Additionally, mice with a weight loss greater than
10% were considered terminal and euthanized.
611
Immunohistochemistry. Paraffin-embedded brains were sectioned at 4 µm and stained with
NeuN antibody (1:250; Cell Signaling) for CA1 neuronal counts. Astrogliosis was detected with
S100β antibody (1:400; Abcam ab41548) and Iba1 antibody (1:400; Abcam, ab5076) was used
for microglial counts. Nonspecific binding was blocked before primary antibodies with 10%
Horse Serum (Sigma Aldrich). A biotinylated secondary antibody (Vector Labs) was used, and
stain was visualized with diaminobenzidine reagent. All images were taken with CellSens
Dimension Desktop 3.1 and counted with CellSens (Iba1 and S100β+) or manually (NeuN).
619
Immunoblotting. Cell lysates were isolated using the protein lysis buffer (50mM Tris, 150mM
NaCl, 2mM EDTA, 1mM MgCl2, 100mM NaF, 10% glycerol, 1% Triton X-100, 1% Na
deoxycholate, 0.1% SDS and 125mM sucrose) supplemented with Phos-STOP and protease
inhibitors (Roche). A BCA Protein Assay kit (Thermo Scientific) was used to quantify protein
concentration of lysates, and 250 mg or 500mg protein was digested with 20 µg/ml proteinase
K (PK) (Roche) for PrP^{Sc} blots for 1 hour at 37C. This digestion allows all other proteins to be
digested so PrP^{Sc} can be detected, therefore loading control antibodies cannot be used for
these blots. Digestion was terminated with 2mM PMSF and lysates were spun at 40,000 x g for
1 hour at 4C before being loaded on a gel. For PrP^{C} blots, 20 mg of samples was used.
Samples were run using 4-20% acrylamide SDS page gels (BioRad) and then transferred onto
0.45µm PVDF blotting paper (MilliPore). Primary antibody Bar-224 (Cayman Chemical

Company) was used at 1:1,000 dilution for PrP^{Sc} blots and 1:5,000 dilution for PrP^{C} blots.
HRP-conjugated secondary antibodies were used at a concentration of 1:5,000 (Vector
Laboratories). For PrP^{C} blots, loading control GAPDH was ran at a 1:5,000 dilution (MilliPore),
with HRP-conjugated anti-mouse secondary at 1:5,000 dilution (Southern Biotech). The protein
antibody complex was visualized using SuperSignal West Pico PLUS Chemiluminescent
Substrate (Thermo Scientific) and visualized with the BioRad ChemiDoc MP.
637
Quantification of Cytokine & Chemokine in mouse tissues. We used previously published
protocols (58-59). Briefly, dissected brain tissues were flash-frozen and homogenized using a
mortar and pestle to form a homogenate solution in tissue cell lysis buffer (Thermo Fisher,
EPX-99999-000). Samples were then centrifuged at 16,000xg for 10 min at 4°C and
supernatant was transferred to a fresh tube. Bio-Rad's DC Protein Assay Kit was used to
determine protein content and all samples were diluted to 10 mg protein/mL. Quantification of
cytokines/ chemokine was performed with Cytokine & Chemokine 36-Plex Mouse ProcartaPlex
Panel 1A (EPXR360-26092-901), analyzed on a Luminex MAGPIX xMAP instrument, and
quantified using xPONENT software. For standard curves, eight four-fold dilutions of protein
standards were used.
648
ICP-MS. Briefly, tissue samples were homogenized in a TissueLyser bead mill (Qiagen,
Germantown, MD) at 30 Hz for 3 min with the addition of 1 μl deionized water per 1 mg organ
tissue. Volumes of homogenates ranging from 10 to 130 μl were then digested in 500 μl of
aqua regia (3:1 hydrochloric acid to nitric acid) for 4 hours at 100 °C. Pellets were
resuspended in water and analyzed with a NexION 2000B single quadropole ICP-MS
(PerkinElmer, Waltham, MA). A Meinhard nebulizer was used with a cyclonic glass spray
chamber for the introduction of the sample. A nickel sample and skimmer cone were used with
an aluminum hyperskimmer cone. The ICP-MS was optimized daily with a calibration solution
of 1 ppb In, Ce, Be, U, and Pb. Au was the analyte monitored. Data was collected using the
sample acquisition module in Syngistix software (version 2.3). Data was analyzed using
Microsoft Excel by converting measured Au concentrations to pg/mg organ via organ
homogenate volume used. A seven-point linear curve with a 1000 ppm Au standard solution
(Ricca Chemical Co, Arlington, TX) was generated at a concentration range of 0- 250 ppb.
Indium was spiked into each sample at a concentration of 5 ppb. Linearity of the standard
curve was defined at an R^2 value of greater than 0.995. TraceMetal grade 70% HNO$_{3-}$ was
purchased from ThermoFisher Scientific (Waltham, MA). All H$_2$O used was from a Milli-Q
system (Millipore, Burlington, MA).
666
Statistical Analysis. All data is presented as mean +/- SEM unless otherwise specified. A
ROUT (Q = 10%) outlier test was performed on all datato identify any potential outliers, which
were removed from the data set. Difference between experimental groups were analyzed
using a One-way ANOVA or Two-way ANOVA. Statistical analysis was completed using Prism.
Significance is denoted throughout the manuscript as *= p $\leq$ 0.05, **= p $\leq$ 0.01, ***= p $\leq$ 0.001
and ****= p $\leq$ 0.001.
673
674 **Ethics Approval.** Mice were euthanized by deeply anaesthetizing with isoflurane followed by
675 decapitation. All mice were bred and maintained at Lab Animal Resources, accredited by the
676 Association for Assessment and Accreditation of Lab Animal Care International, in accordance

677   with protocols approved by the Institutional Animal Care and Use Committee at Colorado State

678   University.

Figure 1. Experimental Design. A) Schematic demonstrating the ability of Nanoligomers to cross the BBB, as well as the mechanism of action (translational inhibition) through which Nanoligomers repress NFKB and NLRP3 expression. **B)** Wild-type C57BL/6 mice were inoculated with RML mouse-adapted prion or normal brain homogenate (NBH). Incubation period is 10 weeks. Prion diseased mice were treated with NF-⊦B and NLRP3 down-regulator SB_NI_112 or vehicle via intraperitoneal (i.p.) or intranasal (i.n.) routes of exposure, twice a week at 10 mg/kg starting at 10 weeks post inoculation (wpi). Mice were monitored for changes in cognition and behavior throughout the disease progression. At 20 wpi brain tissue was dissected for analysis of disease progression including glial inflammation, spongiotic change, and neuronal loss. Created with BioRender.com

Figure 2. NF-⊦B and NLRP3 down-regulator SB_NI_112 penetrates the brain and is systemically non-toxic. Levels of SB_NI_112 in brain regions following dosing at 150 mg/kg i.p. three times per week for two weeks (A). Expression of IFN-© measured with ELISA (B). N=2 for NBH + Vehicle, N=4 for RML + Vehicle and RML + SB_NI_112 i.p. 10 mg/kg. One-way ANOVA, error bars = SEM, *p < 0.05. Weights of mice following a single dose of 500 mg/kg SB_NI_112 i.p. (C). Representative images at 20x of colon, kidney, and liver for female (D-F) and male (G-I) C57BL/6 mice one week after 500 mg/kg dose.

Figure 3. Intraperitoneally delivered SB_NI_112 significantly reduces phosphorylated p65 and NLRP3 protein expression in prion-diseased mice. Western blot images for phosphorylated-p65 (p-p65;A), p65 (B), and NLRP3 (C). Quantification of p-p65/p65 protein expression normalized to GAPDH (D). Quantification of NLRP3 protein expression normalized to GAPDH (E). N=3 for all groups. One-way ANOVA, error bars = SEM, * p< 0.05.

Figure 4. Behavioral and cognitive deficits are protected by NF-⊦B and NLRP3 down-regulation with SB_NI_112 in prion diseased mice. A significant increase in the ability of the mice to build proper nests at 20, 21, and 22 wpi with SB_NI_112 i.p. compared to vehicle treatments. Two-way ANOVA, error bars = SEM, * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001. In the hippocampal burrowing assay, i.p. and i.n. SB_NI_112 treated mice showed significantly better ability to burrow, along with high dose SB_NI_112 i.p. at 16 weeks, compared to untreated counterparts (B). Two-way ANOVA, error bars = SEM, * p< 0.05, ** p< 0.01. Only i.p. SB_NI_112 showed a significant difference in discrimination index at 20 wpi, higher discrimination index trends with i.n. SB_NI_112 (C). Both i.p. and i.n. SB_NI_112 treated mice had significantly higher scores in spatial novel object recognition at 22 wpi (D). N=9-12. One-way ANOVA with pos-hoc analysis, error bars = SEM, * p< 0.05, ** p< 0.01, *** p< 0.001, **** p< 0.0001.

Abbreviations

AD – Alzheimer's Disease

PD – Parkinson's Disease

NF-⊦B – Nuclear factor kappa B

NLRP3 - NOD, LRR, and pyrin-containing 3

wpi – weeks post inoculation

Author Information

Corresponding Author

Julie A. Moreno – *Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University; 1680 Campus Delivery. Fort Collins, CO, USA*; E-mail: julie.moreno@colostate.edu

Authors

Sydney J. Risen – *Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University; 1680 Campus Delivery. Fort Collins, CO, USA*

Sean W. Boland – *Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University; 1680 Campus Delivery. Fort Collins, CO, USA*

Sadhana Sharma – *Sachi Bio., Colorado Technology Center, 685 S Arthur Avenue, Louisville, Colorado 80027, United States*

Grace M. Weisman – *Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University; 1680 Campus Delivery. Fort Collins, CO, USA*

Payton M. Shirley – *Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University; 1680 Campus Delivery. Fort Collins, CO, USA*

Amanda S. Latham – *Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University; 1680 Campus Delivery. Fort Collins, CO, USA*

Arielle J.D. Hay – *Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University; 1680 Campus Delivery. Fort Collins, CO, USA*

Vincenzo S. Gilberto – *Sachi Bio., Colorado Technology Center, 685 S Arthur Avenue, Louisville, Colorado 80027, United States*

Amelia D. Hines – *Department of Environmental and Radiological Health Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University; 1680 Campus Delivery. Fort Collins, CO, USA*

Stephen Brindley – *Department of Pharmaceutical Sciences, University of Colorado Anschutz Medical Campus, Aurora, Colorado 80045, United States.*

Jared M. Brown – *Department of Pharmaceutical Sciences, University of Colorado Anschutz Medical Campus, Aurora, Colorado 80045, United States.*

Stephanie McGrath – *Clinical Sciences, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO, USA*

Anushree Chatterjee – *Sachi Bio., Colorado Technology Center, 685 S Arthur Avenue, Louisville, Colorado 80027, United States*

Prashant Nagpal – *Sachi Bio., Colorado Technology Center, 685 S Arthur Avenue, Louisville, Colorado 80027, United States*

JAM, SJR, SB, SM – *Brain Research Center, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO, USA*

Author Contributions
Conceptualization: JAM, PN, AC, SM
Methodology: SJR, SWB, SS, GMW, PS, ADH, AJDH, VG
Funding acquisition: JAM, PN and AC
Supervision: JAM, PN, AC
Writing – original draft: SJR, JAM
Writing – review & editing: JAM, PN, AC, SM, SJR

Conflict of Interest
The authors declare the following competing financial interest(s): SS, AC, and PN are employed by Sachi Bioworks where this technology was developed. AC and PN are the founders of Sachi Bioworks, and PN has filed a patent on this technology.

Funding Sources
NASA SBIR 80NSSC22CA116, Sachi Bio, and CVMBS Murphy Turney Fund at Colorado State University

Acknowledgments
The authors thank Colorado State University Lab Animal Resources for their animal husbandry.

Supporting Information

Supporting information contains behavioral and cognitive deficit data analysis at 150 mg/kg 2x/week dosing of SB_NI_112 (Figure S1). Additionally, data for ELISA analysis of IL-28, IL2, IL-1a, and IL-1β (Figure S2). Lastly, videos displaying clinical signs (clasping, righting reflex, and walking) for control and treatment groups at 22 weeks post inoculation (wpi) are available in Supplement Videos 1-9.

References

(1) Gavilán, M. P.; Revilla, E.; Pintado, C.; Castaño, A.; Vizuete, M. L.; Moreno-González, I.; Baglietto-Vargas, D.; Sánchez-Varo, R.; Vitorica, J.; Gutiérrez, A.; Ruano, D. Molecular and Cellular Characterization of the Age-Related Neuroinflammatory Processes Occurring in Normal Rat Hippocampus: Potential Relation with the Loss of Somatostatin GABAergic Neurons. *J. Neurochem.* **2007**, *103* (3), 984–996. https://doi.org/10.1111/j.1471-4159.2007.04787.x.

(2) Jimenez, S.; Baglietto-Vargas, D.; Caballero, C.; Moreno-Gonzalez, I.; Torres, M.; Sanchez-Varo, R.; Ruano, D.; Vizuete, M.; Gutierrez, A.; Vitorica, J. Inflammatory Response in the Hippocampus of PS1M146L/APP751SL Mouse Model of Alzheimer's Disease: Age-Dependent Switch in the Microglial Phenotype from Alternative to Classic. *J. Neurosci.* **2008**, *28* (45), 11650–11661. https://doi.org/10.1523/JNEUROSCI.3024-08.2008.

(3) Zhao, J.; Bi, W.; Xiao, S.; Lan, X.; Cheng, X.; Zhang, J.; Lu, D.; Wei, W.; Wang, Y.; Li, H.; Fu, Y.; Zhu, L. Neuroinflammation Induced by Lipopolysaccharide Causes Cognitive Impairment in Mice. *Sci. Rep.* **2019**, *9* (1), 5790. https://doi.org/10.1038/s41598-019-42286-8.

(4) Guan, Y.; Han, F. Key Mechanisms and Potential Targets of the NLRP3 Inflammasome in Neurodegenerative Diseases. *Front. Integr. Neurosci.* **2020**, *14*, 37. https://doi.org/10.3389/fnint.2020.00037.

(5) Holbrook, J. A.; Jarosz-Griffiths, H. H.; Caseley, E.; Lara-Reyna, S.; Poulter, J. A.; Williams-Gray, C. H.; Peckham, D.; McDermott, M. F. Neurodegenerative Disease and the NLRP3 Inflammasome. *Front. Pharmacol.* **2021**, *12*.

(6) González-Sanmiguel, J.; Schuh, C. M. A. P.; Muñoz-Montesino, C.; Contreras-Kallens, P.; Aguayo, L. G.; Aguayo, S. Complex Interaction between Resident Microbiota and Misfolded Proteins: Role in Neuroinflammation and Neurodegeneration. *Cells* **2020**, *9* (11), 2476. https://doi.org/10.3390/cells9112476.

(7) Yiannopoulou, K. G.; Anastasiou, A. I.; Zachariou, V.; Pelidou, S.-H. Reasons for Failed Trials of Disease-Modifying Treatments for Alzheimer Disease and Their Contribution in Recent Research. *Biomedicines* **2019**, *7* (4), 97. https://doi.org/10.3390/biomedicines7040097.

(8) Amirian, R.; Badrbani, M. A.; Derakhshankhah, H.; Izadi, Z.; Shahbazi, M.-A. Targeted Protein Degradation for the Treatment of Parkinson's Disease: Advances and Future Perspective. *Biomed. Pharmacother. Biomedecine Pharmacother.* **2023**, *166*, 115408. https://doi.org/10.1016/j.biopha.2023.115408.

(9) Relaño-Ginés, A.; Lehmann, S.; Crozet, C. Cell-Based Therapy against Prion Diseases. *Curr. Opin. Pharmacol.* **2019**, *44*, 8–14. https://doi.org/10.1016/j.coph.2018.11.001.

(10)   Kannarkat, G. T.; Boss, J. M.; Tansey, M. G. The Role of Innate and Adaptive Immunity
in Parkinson's Disease. *J. Park. Dis.* **2013**, *3* (4), 493–514. https://doi.org/10.3233/JPD-
130250.
(11)   Guzman-Martinez, L.; Maccioni, R. B.; Andrade, V.; Navarrete, L. P.; Pastor, M. G.;
Ramos-Escobar, N. Neuroinflammation as a Common Feature of Neurodegenerative
Disorders. *Front. Pharmacol.* **2019**, *10*, 1008. https://doi.org/10.3389/fphar.2019.01008.
(12)   Mabbott, N. A.; Bradford, B. M.; Pal, R.; Young, R.; Donaldson, D. S. The Effects of
Immune System Modulation on Prion Disease Susceptibility and Pathogenesis. *Int. J. Mol.*
Sci. **2020**, *21* (19), 7299. https://doi.org/10.3390/ijms21197299.
(13)   Stancu, I.-C.; Cremers, N.; Vanrusselt, H.; Couturier, J.; Vanoosthuyse, A.; Kessels, S.;
Lodder, C.; Brône, B.; Huaux, F.; Octave, J.-N.; Terwel, D.; Dewachter, I. Aggregated Tau
Activates NLRP3-ASC Inflammasome Exacerbating Exogenously Seeded and Non-
Exogenously Seeded Tau Pathology in Vivo. *Acta Neuropathol. (Berl.)* **2019**, *137* (4), 599–
617. https://doi.org/10.1007/s00401-018-01957-y.
(14)   Gustot, A.; Gallea, J. I.; Sarroukh, R.; Celej, M. S.; Ruysschaert, J.-M.; Raussens, V.
Amyloid Fibrils Are the Molecular Trigger of Inflammation in Parkinson's Disease. *Biochem.*
J. **2015**, *471* (3), 323–333. https://doi.org/10.1042/BJ20150617.
(15)   Carroll, J. A.; Chesebro, B. Neuroinflammation, Microglia, and Cell-Association during
Prion Disease. *Viruses* **2019**, *11* (1), 65. https://doi.org/10.3390/v11010065.
(16)   Akwa, Y. Steroids and Alzheimer's Disease: Changes Associated with Pathology and
Therapeutic Potential. *Int. J. Mol. Sci.* **2020**, *21* (13), 4812.
https://doi.org/10.3390/ijms21134812.
(17)   Ming Wang, J.; Irwin, R. W.; Liu, L.; Chen, S.; Brinton, R. D. Regeneration in a
Degenerating Brain: Potential of Allopregnanolone as a Neuroregenerative Agent. *Curr.*
Alzheimer Res. **2007**, *4* (5), 510–517. https://doi.org/10.2174/156720507783018262.
(18)   Hanslik, K. L.; Ulland, T. K. The Role of Microglia and the Nlrp3 Inflammasome in
Alzheimer's Disease. *Front. Neurol.* **2020**, *11*, 570711.
https://doi.org/10.3389/fneur.2020.570711.
(19)   Cheng, J.; Liao, Y.; Dong, Y.; Hu, H.; Yang, N.; Kong, X.; Li, S.; Li, X.; Guo, J.; Qin, L.;
Yu, J.; Ma, C.; Li, J.; Li, M.; Tang, B.; Yuan, Z. Microglial Autophagy Defect Causes
Parkinson Disease-like Symptoms by Accelerating Inflammasome Activation in Mice.
Autophagy 16 (12), 2193–2205. https://doi.org/10.1080/15548627.2020.1719723.
(20)   Kim, T.-K.; Lee, I.; Cho, J.-H.; Canine, B.; Keller, A.; Price, N. D.; Hwang, D.; Carlson,
G.; Hood, L. Core Transcriptional Regulatory Circuits in Prion Diseases. *Mol. Brain* **2020**,
13 (1), 10. https://doi.org/10.1186/s13041-020-0551-3.
(21)   Liu, T.; Zhang, L.; Joo, D.; Sun, S.-C. NF-κB Signaling in Inflammation. *Signal*
Transduct. Target. Ther. **2017**, *2*, 17023-. https://doi.org/10.1038/sigtrans.2017.23.
(22)   Kushwaha, R.; Sinha, A.; Makarava, N.; Molesworth, K.; Baskakov, I. V. Non-Cell
Autonomous Astrocyte-Mediated Neuronal Toxicity in Prion Diseases. *Acta Neuropathol.*
Commun. **2021**, *9* (1), 22. https://doi.org/10.1186/s40478-021-01123-8.
(23)   Kushwaha, R.; Li, Y.; Makarava, N.; Pandit, N. P.; Molesworth, K.; Birukov, K. G.;
Baskakov, I. V. Reactive Astrocytes Associated with Prion Disease Impair the Blood Brain
Barrier. *Neurobiol. Dis.* **2023**, *185*, 106264. https://doi.org/10.1016/j.nbd.2023.106264.
(24)   Venegas, C.; Kumar, S.; Franklin, B. S.; Dierkes, T.; Brinkschulte, R.; Tejera, D.; Vieira-
Saecker, A.; Schwartz, S.; Santarelli, F.; Kummer, M. P.; Griep, A.; Gelpi, E.; Beilharz, M.;
Riedel, D.; Golenbock, D. T.; Geyer, M.; Walter, J.; Latz, E.; Heneka, M. T. Microglia-

Derived ASC Specks Cross-Seed Amyloid-β in Alzheimer's Disease. *Nature* **2017**, *552*
(7685), 355–361. https://doi.org/10.1038/nature25158.

(25)   Ising, C.; Venegas, C.; Zhang, S.; Scheiblich, H.; Schmidt, S. V.; Vieira-Saecker, A.;
Schwartz, S.; Albasset, S.; McManus, R. M.; Tejera, D.; Griep, A.; Santarelli, F.; Brosseron,
F.; Opitz, S.; Stunden, J.; Merten, M.; Kayed, R.; Golenbock, D. T.; Blum, D.; Latz, E.;
Buée, L.; Heneka, M. T. NLRP3 Inflammasome Activation Drives Tau Pathology. *Nature*
2019, *575* (7784), 669–673. https://doi.org/10.1038/s41586-019-1769-z.

(26)   Scheiblich, H.; Bousset, L.; Schwartz, S.; Griep, A.; Latz, E.; Melki, R.; Heneka, M. T.
Microglial NLRP3 Inflammasome Activation upon TLR2 and TLR5 Ligation by Distinct α-
Synuclein Assemblies. *J. Immunol. Baltim. Md 1950* **2021**, *207* (8), 2143–2154.
https://doi.org/10.4049/jimmunol.2100035.

(27)   Kim, T.-K.; Bae, E.-J.; Jung, B. C.; Choi, M.; Shin, S. J.; Park, S. J.; Kim, J. T.; Jung, M.
K.; Ulusoy, A.; Song, M.-Y.; Lee, J. S.; Lee, H.-J.; Di Monte, D. A.; Lee, S.-J. Inflammation
Promotes Synucleinopathy Propagation. *Exp. Mol. Med.* **2022**, *54* (12), 2148–2161.
https://doi.org/10.1038/s12276-022-00895-w.

(28)   Hay, A. J. D.; Latham, A. S.; Mumford, G.; Hines, A. D.; Risen, S.; Gordon, E.;
Siebenaler, C.; Gilberto, V. S.; Zabel, M. D.; Moreno, J. A. Intranasally Delivered
Mesenchymal Stromal Cells Decrease Glial Inflammation Early in Prion Disease. *Front.*
Neurosci. **2023**, *17*, 1158408. https://doi.org/10.3389/fnins.2023.1158408.

(29)   Hay, A. J. D.; Murphy, T. J.; Popichak, K. A.; Zabel, M. D.; Moreno, J. A. Adipose-
Derived Mesenchymal Stromal Cells Decrease Prion-Induced Glial Inflammation in Vitro.
Sci. Rep. **2022**, *12*, 22567. https://doi.org/10.1038/s41598-022-26628-7.

(30)   Shi, F.; Yang, L.; Kouadir, M.; Yang, Y.; Wang, J.; Zhou, X.; Yin, X.; Zhao, D. The
NALP3 Inflammasome Is Involved in Neurotoxic Prion Peptide-Induced Microglial
Activation. *J. Neuroinflammation* **2012**, *9*, 73. https://doi.org/10.1186/1742-2094-9-73.

(31)   Hafner-Bratkovič, I.; Benčina, M.; Fitzgerald, K. A.; Golenbock, D.; Jerala, R. NLRP3
Inflammasome Activation in Macrophage Cell Lines by Prion Protein Fibrils as the Source
of IL-1β and Neuronal Toxicity. *Cell. Mol. Life Sci. CMLS* **2012**, *69* (24), 4215–4228.
https://doi.org/10.1007/s00018-012-1140-0.

(32)   Kelley, N.; Jeltema, D.; Duan, Y.; He, Y. The NLRP3 Inflammasome: An Overview of
Mechanisms of Activation and Regulation. *Int. J. Mol. Sci.* **2019**, *20* (13), 3328.
https://doi.org/10.3390/ijms20133328.

(33)   Shi, J.; Zhao, Y.; Wang, K.; Shi, X.; Wang, Y.; Huang, H.; Zhuang, Y.; Cai, T.; Wang, F.;
Shao, F. Cleavage of GSDMD by Inflammatory Caspases Determines Pyroptotic Cell
Death. *Nature* **2015**, *526* (7575), 660–665. https://doi.org/10.1038/nature15514.

(34)   Martinon, F.; Burns, K.; Tschopp, J. The Inflammasome: A Molecular Platform
Triggering Activation of Inflammatory Caspases and Processing of proIL-Beta. *Mol. Cell*
2002, *10* (2), 417–426. https://doi.org/10.1016/s1097-2765(02)00599-3.

(35)   Bacot, S. M.; Lenz, P.; Frazier-Jessen, M. R.; Feldman, G. M. Activation by Prion
Peptide PrP106-126 Induces a NF-kappaB-Driven Proinflammatory Response in Human
Monocyte-Derived Dendritic Cells. *J. Leukoc. Biol.* **2003**, *74* (1), 118–125.
https://doi.org/10.1189/jlb.1102521.

(36)   Prasad, K. N.; Bondy, S. C. Oxidative and Inflammatory Events in Prion Diseases: Can
They Be Therapeutic Targets? *Curr. Aging Sci.* **2019**, *11* (4), 216–225.
https://doi.org/10.2174/1874609812666190111100205.

(37)   Wu, G.-R.; Mu, T.-C.; Gao, Z.-X.; Wang, J.; Sy, M.-S.; Li, C.-Y. Prion Protein Is Required for Tumor Necrosis Factor α (TNFα)-Triggered Nuclear Factor κB (NF-κB) Signaling and Cytokine Production. *J. Biol. Chem.* **2017**, *292* (46), 18747–18759. https://doi.org/10.1074/jbc.M117.787283.

(38)   Bourgognon, J.-M.; Spiers, J. G.; Robinson, S. W.; Scheiblich, H.; Glynn, P.; Ortori, C.; Bradley, S. J.; Tobin, A. B.; Steinert, J. R. Inhibition of Neuroinflammatory Nitric Oxide Signaling Suppresses Glycation and Prevents Neuronal Dysfunction in Mouse Prion Disease. *Proc. Natl. Acad. Sci. U. S. A.* **2021**, *118* (10), e2009579118. https://doi.org/10.1073/pnas.2009579118.

(39)   Lv, Q.-K.; Tao, K.-X.; Wang, X.-B.; Yao, X.-Y.; Pang, M.-Z.; Liu, J.-Y.; Wang, F.; Liu, C.-F. Role of α-Synuclein in Microglia: Autophagy and Phagocytosis Balance Neuroinflammation in Parkinson's Disease. *Inflamm. Res.* **2023**, *72* (3), 443–462. https://doi.org/10.1007/s00011-022-01676-x.

(40)   Choi, I.; Wang, M.; Yoo, S.; Xu, P.; Seegobin, S. P.; Li, X.; Han, X.; Wang, Q.; Peng, J.; Zhang, B.; Yue, Z. Autophagy Enables Microglia to Engage Amyloid Plaques and Prevents Microglial Senescence. *Nat. Cell Biol.* **2023**, *25* (7), 963–974. https://doi.org/10.1038/s41556-023-01158-0.

(41)   Lai, M.; Yao, H.; Shah, S. Z. A.; Wu, W.; Wang, D.; Zhao, Y.; Wang, L.; Zhou, X.; Zhao, D.; Yang, L. The NLRP3-Caspase 1 Inflammasome Negatively Regulates Autophagy via TLR4-TRIF in Prion Peptide-Infected Microglia. *Front. Aging Neurosci.* **2018**, *10*, 116. https://doi.org/10.3389/fnagi.2018.00116.

(42)   Sharma, S.; Borski, C.; Hanson, J.; Garcia, M. A.; Link, C. D.; Hoeffer, C.; Chatterjee, A.; Nagpal, P. Identifying an Optimal Neuroinflammation Treatment Using a Nanoligomer Discovery Engine. *ACS Chem. Neurosci.* **2022**, *13* (23), 3247–3256. https://doi.org/10.1021/acschemneuro.2c00365.

(43)   Courtney, C. M.; Sharma, S.; Fallgren, C.; Weil, M. M.; Chatterjee, A.; Nagpal, P. Reversing Radiation-Induced Immunosuppression Using a New Therapeutic Modality. *Life Sci. Space Res.* **2022**, *35*, 127–139. https://doi.org/10.1016/j.lssr.2022.05.002.

(44)   Kovacs, G. G.; Budka, H. Molecular Pathology of Human Prion Diseases. *Int. J. Mol. Sci.* **2009**, *10* (3), 976–999. https://doi.org/10.3390/ijms10030976.

(45)   Prusiner, S. B. Prions. *Proc. Natl. Acad. Sci. U. S. A.* **1998**, *95* (23), 13363–13383. https://doi.org/10.1073/pnas.95.23.13363.

(46)   Mari, Z.; Mestre, T. A. The Disease Modification Conundrum in Parkinson's Disease: Failures and Hopes. *Front. Aging Neurosci.* **2022**, *14*, 810860. https://doi.org/10.3389/fnagi.2022.810860.

(47)   Geschwind, M. D.; Kuo, A. L.; Wong, K. S.; Haman, A.; Devereux, G.; Raudabaugh, B. J.; Johnson, D. Y.; Torres-Chae, C. C.; Finley, R.; Garcia, P.; Thai, J. N.; Cheng, H. Q.; Neuhaus, J. M.; Forner, S. A.; Duncan, J. L.; Possin, K. L.; Dearmond, S. J.; Prusiner, S. B.; Miller, B. L. Quinacrine Treatment Trial for Sporadic Creutzfeldt-Jakob Disease. *Neurology* **2013**, *81* (23), 2015–2023. https://doi.org/10.1212/WNL.0b013e3182a9f3b4.

(48)   Moreno, J. A.; Halliday, M.; Molloy, C.; Radford, H.; Verity, N.; Axten, J. M.; Ortori, C. A.; Willis, A. E.; Fischer, P. M.; Barrett, D. A.; Mallucci, G. R. Oral Treatment Targeting the Unfolded Protein Response Prevents Neurodegeneration and Clinical Disease in Prion-Infected Mice. *Sci. Transl. Med.* **2013**, *5* (206), 206ra138-206ra138. https://doi.org/10.1126/scitranslmed.3006767.

(49)   Moreno, J. A.; Radford, H.; Peretti, D.; Steinert, J. R.; Verity, N.; Martin, M. G.; Halliday, M.; Morgan, J.; Dinsdale, D.; Ortori, C. A.; Barrett, D. A.; Tsaytler, P.; Bertolotti, A.; Willis, A. E.; Bushell, M.; Mallucci, G. R. Sustained Translational Repression by eIF2α-P Mediates Prion Neurodegeneration. *Nature* **2012**, *485* (7399), 507–511. https://doi.org/10.1038/nature11058.

(50)   McCollum, C. R.; Courtney, C. M.; O'Connor, N. J.; Aunins, T. R.; Jordan, T. X.; Rogers, K. L.; Brindley, S.; Brown, J. M.; Nagpal, P.; Chatterjee, A. Safety and Biodistribution of Nanoligomers Targeting the SARS-CoV-2 Genome for the Treatment of COVID-19. *ACS Biomater. Sci. Eng.* **2023**, *9* (3), 1656–1671. https://doi.org/10.1021/acsbiomaterials.2c00669.

(51)   Deacon, R. M.; Raley, J. M.; Perry, V. H.; Rawlins, J. N. Burrowing into Prion Disease. *Neuroreport* **2001**, *12* (9), 2053–2057. https://doi.org/10.1097/00001756-200107030-00052.

(52)   Mallucci, G. R.; White, M. D.; Farmer, M.; Dickinson, A.; Khatun, H.; Powell, A. D.; Brandner, S.; Jefferys, J. G. R.; Collinge, J. Targeting Cellular Prion Protein Reverses Early Cognitive Deficits and Neurophysiological Dysfunction in Prion-Infected Mice. *Neuron* **2007**, *53* (3), 325–335. https://doi.org/10.1016/j.neuron.2007.01.005.

(53)   Lueptow, L. M. Novel Object Recognition Test for the Investigation of Learning and Memory in Mice. *J. Vis. Exp. JoVE* **2017**, No. 126, 55718. https://doi.org/10.3791/55718.

(54)   de Melo, A. S. L. F.; Lima, J. L. D.; Malta, M. C. S.; Marroquim, N. F.; Moreira, Á. R.; de Almeida Ladeia, I.; Dos Santos Cardoso, F.; Gonçalves, D. B.; Dutra, B. G.; Dos Santos, J. C. C. The Role of Microglia in Prion Diseases and Possible Therapeutic Targets: A Literature Review. *Prion* **2021**, *15* (1), 191–206. https://doi.org/10.1080/19336896.2021.1991771.

(55)   Aguzzi, A.; Zhu, C. Microglia in Prion Diseases. *J. Clin. Invest.* **2017**, *127* (9), 3230–3239. https://doi.org/10.1172/JCI90605.

(56)   Alibhai, J.; Blanco, R. A.; Barria, M. A.; Piccardo, P.; Caughey, B.; Perry, V. H.; Freeman, T. C.; Manson, J. C. Distribution of Misfolded Prion Protein Seeding Activity Alone Does Not Predict Regions of Neurodegeneration. *PLoS Biol.* **2016**, *14* (11), e1002579. https://doi.org/10.1371/journal.pbio.1002579.

(57)   Tahir, W.; Thapa, S.; Schatzl, H. M. Astrocyte in Prion Disease: A Double-Edged Sword. *Neural Regen. Res.* **2022**, *17* (8), 1659–1665. https://doi.org/10.4103/1673-5374.332202.

(58)   Li, S.; Fang, Y.; Zhang, Y.; Song, M.; Zhang, X.; Ding, X.; Yao, H.; Chen, M.; Sun, Y.; Ding, J.; Wang, Q.; Lu, M.; Wu, G.; Hu, G. Microglial NLRP3 Inflammasome Activates Neurotoxic Astrocytes in Depression-like Mice. *Cell Rep.* **2022**, *41* (4), 111532. https://doi.org/10.1016/j.celrep.2022.111532.

(59)   Latham, A. S.; Geer, C. E.; Ackart, D. F.; Anderson, I. K.; Vittoria, K. M.; Podell, B. K.; Basaraba, R. J.; Moreno, J. A. Gliosis, Misfolded Protein Aggregation, and Neuronal Loss in a Guinea Pig Model of Pulmonary Tuberculosis. *Front. Neurosci.* **2023**, *17*, 1157652. https://doi.org/10.3389/fnins.2023.1157652.

(60)   Salehi, P.; Clark, M.; Pinzon, J.; Patil, A. Sporadic Creutzfeldt-Jakob Disease. *Am. J. Emerg. Med.* **2022**, *52*, 267.e1-267.e3. https://doi.org/10.1016/j.ajem.2021.07.038.

(61)   Dutta, S.; Sengupta, P. Men and Mice: Relating Their Ages. *Life Sci.* **2016**, *152*, 244–248. https://doi.org/10.1016/j.lfs.2015.10.025.

(62) Agostini, L.; Martinon, F.; Burns, K.; McDermott, M. F.; Hawkins, P. N.; Tschopp, J.
NALP3 Forms an IL-1beta-Processing Inflammasome with Increased Activity in Muckle-
Wells Autoinflammatory Disorder. *Immunity* **2004**, *20* (3), 319–325.
https://doi.org/10.1016/s1074-7613(04)00046-9.
(63) Nuvolone, M.; Sorce, S.; Schwarz, P.; Aguzzi, A. Prion Pathogenesis in the Absence of
NLRP3/ASC Inflammasomes. *PloS One* **2015**, *10* (2), e0117208.
https://doi.org/10.1371/journal.pone.0117208.
(64) Soto, C.; Satani, N. The Intricate Mechanisms of Neurodegeneration in Prion Diseases.
Trends Mol. Med. **2011**, *17* (1), 14–24. https://doi.org/10.1016/j.molmed.2010.09.001.
(65) Giovannoni, F.; Quintana, F. J. The Role of Astrocytes in CNS Inflammation. *Trends
Immunol.* **2020**, *41* (9), 805–819. https://doi.org/10.1016/j.it.2020.07.007.
(66) Minikel, E. V.; Zhao, H. T.; Le, J.; O'Moore, J.; Pitstick, R.; Graffam, S.; Carlson, G. A.;
Kavanaugh, M. P.; Kriz, J.; Kim, J. B.; Ma, J.; Wille, H.; Aiken, J.; McKenzie, D.; Doh-ura,
K.; Beck, M.; O'Keefe, R.; Stathopoulos, J.; Caron, T.; Schreiber, S. L.; Carroll, J. B.;
Kordasiewicz, H. B.; Cabin, D. E.; Vallabh, S. M. Prion Protein Lowering Is a Disease-
Modifying Therapy across Prion Disease Stages, Strains and Endpoints. *Nucleic Acids
Res.* **2020**, *48* (19), 10615–10631. https://doi.org/10.1093/nar/gkaa616.
(67) Rovis, T. L.; Legname, G. Prion Protein-Specific Antibodies-Development, Modes of
Action and Therapeutics Application. *Viruses* **2014**, *6* (10), 3719–3737.
https://doi.org/10.3390/v6103719.
(68) Gayrard, V.; Picard-Hagen, N.; Viguié, C.; Laroute, V.; Andréoletti, O.; Toutain, P.-L. A
Possible Pharmacological Explanation for Quinacrine Failure to Treat Prion Diseases:
Pharmacokinetic Investigations in a Ovine Model of Scrapie. *Br. J. Pharmacol.* **2005**, *144*
(3), 386–393. https://doi.org/10.1038/sj.bjp.0706072.
(69) Collins, S. J.; Lewis, V.; Brazier, M.; Hill, A. F.; Fletcher, A.; Masters, C. L. Quinacrine
Does Not Prolong Survival in a Murine Creutzfeldt-Jakob Disease Model. *Ann. Neurol.*
2002, *52* (4), 503–506. https://doi.org/10.1002/ana.10336.
(70) Ghaemmaghami, S.; Ahn, M.; Lessard, P.; Giles, K.; Legname, G.; DeArmond, S. J.;
Prusiner, S. B. Continuous Quinacrine Treatment Results in the Formation of Drug-
Resistant Prions. *PLOS Pathog.* **2009**, *5* (11), e1000673.
https://doi.org/10.1371/journal.ppat.1000673.
(71) Collinge, J.; Gorham, M.; Hudson, F.; Kennedy, A.; Keogh, G.; Pal, S.; Rossor, M.;
Rudge, P.; Siddique, D.; Spyer, M.; Thomas, D.; Walker, S.; Webb, T.; Wroe, S.;
Darbyshire, J. Safety and Efficacy of Quinacrine in Human Prion Disease (PRION-1
Study): A Patient-Preference Trial. *Lancet Neurol.* **2009**, *8* (4), 334–344.
https://doi.org/10.1016/S1474-4422(09)70049-3.
(72) Haïk, S.; Brandel, J. P.; Salomon, D.; Sazdovitch, V.; Delasnerie-Lauprêtre, N.;
Laplanche, J. L.; Faucheux, B. A.; Soubrié, C.; Boher, E.; Belorgey, C.; Hauw, J. J.;
Alpérovitch, A. Compassionate Use of Quinacrine in Creutzfeldt-Jakob Disease Fails to
Show Significant Effects. *Neurology* **2004**, *63* (12), 2413–2415.
https://doi.org/10.1212/01.wnl.0000148596.15681.4d.
(73) Mead, S.; Khalili-Shirazi, A.; Potter, C.; Mok, T.; Nihat, A.; Hyare, H.; Canning, S.;
Schmidt, C.; Campbell, T.; Darwent, L.; Muirhead, N.; Ebsworth, N.; Hextall, P.; Wakeling,
M.; Linehan, J.; Libri, V.; Williams, B.; Jaunmuktane, Z.; Brandner, S.; Rudge, P.; Collinge,
J. Prion Protein Monoclonal Antibody (PRN100) Therapy for Creutzfeldt–Jakob Disease:

Evaluation of a First-in-Human Treatment Programme. *Lancet Neurol.* **2022**, *21* (4), 342–354. https://doi.org/10.1016/S1474-4422(22)00082-5.

(74)   Friberg, K. N.; Hung, G.; Wancewicz, E.; Giles, K.; Black, C.; Freier, S.; Bennett, F.; DeArmond, S. J.; Freyman, Y.; Lessard, P.; Ghaemmaghami, S.; Prusiner, S. B. Intracerebral Infusion of Antisense Oligonucleotides Into Prion-Infected Mice. *Mol. Ther. - Nucleic Acids* **2012**, *1*. https://doi.org/10.1038/mtna.2011.6.

(75)   McDonald, J. T.; Enguita, F. J.; Taylor, D.; Griffin, R. J.; Priebe, W.; Emmett, M. R.; Sajadi, M. M.; Harris, A. D.; Clement, J.; Dybas, J. M.; Aykin-Burns, N.; Guarnieri, J. W.; Singh, L. N.; Grabham, P.; Baylin, S. B.; Yousey, A.; Pearson, A. N.; Corry, P. M.; Saravia-Butler, A.; Aunins, T. R.; Sharma, S.; Nagpal, P.; Meydan, C.; Foox, J.; Mozsary, C.; Cerqueira, B.; Zaksas, V.; Singh, U.; Wurtele, E. S.; Costes, S. V.; Davanzo, G. G.; Galeano, D.; Paccanaro, A.; Meinig, S. L.; Hagan, R. S.; Bowman, N. M.; UNC COVID-19 Pathobiology Consortium; Wolfgang, M. C.; Altinok, S.; Sapoval, N.; Treangen, T. J.; Moraes-Vieira, P. M.; Vanderburg, C.; Wallace, D. C.; Schisler, J. C.; Mason, C. E.; Chatterjee, A.; Meller, R.; Beheshti, A. Role of miR-2392 in Driving SARS-CoV-2 Infection. *Cell Rep.* **2021**, *37* (3), 109839. https://doi.org/10.1016/j.celrep.2021.109839.

(76)   McCollum, C. R.; Courtney, C. M.; O'Connor, N. J.; Aunins, T. R.; Ding, Y.; Jordan, T. X.; Rogers, K. L.; Brindley, S.; Brown, J. M.; Nagpal, P.; Chatterjee, A. Nanoligomers Targeting Human miRNA for the Treatment of Severe COVID-19 Are Safe and Nontoxic in Mice. *ACS Biomater. Sci. Eng.* **2022**, *8* (7), 3087–3106. https://doi.org/10.1021/acsbiomaterials.2c00510.

(77)   Aunins, T. R.; Erickson, K. E.; Chatterjee, A. Transcriptome-Based Design of Antisense Inhibitors Potentiates Carbapenem Efficacy in CRE Escherichia Coli. *Proc. Natl. Acad. Sci. U. S. A.* **2020**, *117* (48), 30699–30709. https://doi.org/10.1073/pnas.1922187117.

(78)   Larsen, H. J.; Bentin, T.; Nielsen, P. E. Antisense Properties of Peptide Nucleic Acid. *Biochim. Biophys. Acta* **1999**, *1489* (1), 159–166. https://doi.org/10.1016/s0167-4781(99)00145-1.

Figure 1

Figure 2

Figure 3

A
75 kDa —   p-p65
35 kDa —   GAPDH

B
75 kDa —   p65
35 kDa —   GAPDH

C
100 kDa —
80 kDa —   NLRP3
35 kDa —   GAPDH

Legend:
- NBH + Vehicle (green)
- RML + Vehicle (red)
- RML + SB_NI_112 i.n. (magenta)
- RML + SB_NI_112 i.p. (blue)

D — p-p65/p65 protein expression (normalized to GAPDH)
*

E — NLRP3 protien expression (normalized to GAPDH)
* *

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supporting Figure 1