Identification of potential human pancreatic *α*-amylase inhibitors from natural products by molecular docking, MM/GBSA calculations, MD simulations, and ADMET analysis

Santosh Basnet†, Madhav Prasad Ghimire††, Tika Ram Lamichhane††, Rajendra Adhikari†††, Achyut Adhikari†∗

† Central Department of Chemistry, Tribhuvan University, Kirtipur-44613, Kathmandu, Nepal

†† Central Department of Physics, Tribhuvan University, Kirtipur-44613, Kathmandu, Nepal

††† Department of Physics, Kathmandu University, Dhulikhel, Nepal

∗ Corresponding author: achyutraj05@gmail.com

Abstract

Human pancreatic *α*-amylase (HPA), which works as a catalyst for carbohydrate hydrolysis, is one of the viable targets to control type 2 diabetes. The inhibition of *α*-amylase lowers blood glucose levels and helps to alleviate hyperglycemia complications. Herein, we systematically screened the potential HPA inhibitors from an in-house library of natural products by molecular modeling. The modeling encompasses molecular docking, MM/GBSA binding energy calculations, MD simulations, and ADMET analysis. This research identified newboulaside B, newboulaside A, quercetin-3-O-*β*-glucoside, and sasastilboside A as the top four potential HPA inhibitors from the library of natural products, whose Glide docking scores and MM/GBSA binding energies range from -9.191 to -11.366 kcal/mol and -19.38 to -50.29 kcal/mol, respectively. Based on the simulation, among them, newboulaside B was found as the best HPA inhibitor. Throughout the simulation, with the deviation of 3Å (acarbose = 3Å), it interacted with ASP356, ASP300, ASP197, THR163, ARG161, ASP147, ALA106, and GLN63 via hydrogen bonding. Additionally, the comprehensive ADMET analysis revealed that it has good pharmacokinetic properties having not acutely toxic,

moderately bioavailable, and non-inhibitor nature toward cytochrome P450. All the results suggest that newboulaside B might be a promising candidate for drug discovery against type 2 diabetes.

Keywords: Type 2 Diabetes, Human Pancreatic α-Amylase, Natural Products, Molecular Modeling, Newboulaside B

1. Introduction

Type 2 diabetes, associated with hyperglycemia, is a long-term endocrine metabolic disorder. It is most prevalent and caused by aberrant insulin activity. It causes various chronic complications. Despite medical improvements, it is still one of the leading causes of death in adults [1,2].

Delaying the digestion of carbohydrates ameliorates hyperglycemia impacts. Multiple enzymes, including human pancreatic α-amylase (HPA) and α-glucosidase, are involved in the hydrolysis of carbohydrates. According to the study, in the HPA, catalytic residues (ASP197, GLU233, and ASP300) are found in an eight-stranded parallel beta-barrel in domain A (1-99, 169-404). In carbohydrate hydrolysis, ASP197 acts as a nucleophile, and GLU233 and ASP300 may act as acid/base catalysts. Therefore, one of the proven scientific methods to manage type 2 diabetes is the inhibition of HPA by ligand, which lowers postprandial blood glucose levels [3–5].

A wide variety of oral anti-diabetic medications, including acarbose and miglitol, were used by doctors to treat type 2 diabetes. The undesirable, serious side effects, however, are becoming apparent. Therefore, the need for efficient, affordable, and low-toxic anti-diabetic medications is urgent [6–9].

To treat diabetes, ancient people relied on medications from natural sources [10]. Therefore, natural products derived from these sources may be an appealing strategy for controlling type 2 diabetes [11]. The research is ramping up to explore a potent HPA inhibitor. The *in vitro* study shows that different natural products exhibit α-amylase inhibition activity. There have been several *in vitro* studies on natural products up to this moment, but an efficient HPA inhibitor is still lacking. Therefore, to identify the potential HPA inhibitor from natural products, structure-based molecular modeling was employed. The modeling comprises molecular docking, MM/GBSA binding energy calculations, molecular dynamics

simulations, and absorption, distribution, metabolism, excretion, and toxicity (ADMET) analysis.

This computational study proposed newboulaside B as the potential HPA inhibitor from natural products. The study also suggests a possible binding mode of the inhibitor with HPA.

2. Material and methods

2.1 Curation and preparation of ligands

Based on *in vitro* data on α-amylase, an in-house library of 39 ligands (including acarbose as the standard ligand) was created to screen a potential HPA inhibitor. Supplementary data contains information about the ligands (**Table S1**). The ligand structures were sketched in Maestro's 2D sketcher and converted to a low-energy 3D state using Schrödinger's LigPrep module [12]. The OPLS4 force field was employed to generate a low-energy state structure [13]. Additionally, at pH 7.0 ± 2.0, Epik produced tautomeric and ionization states and generated a maximum of 32 conformers per ligand.

2.2 Protein preparation and target validation

The structure of HPA was obtained from the RCSB protein data repository and prepared using Schrödinger's Protein Preparation Wizard module (**Table S2**) [14,15]. It was preprocessed with Prime, which assigned bond ordering, added hydrogen atoms, and filled missing loops and side chains; Epik was then run at pH 7.0 ± 2.0 [16]. Additionally, after removing hetero groups, it was optimized by optimizing hydrogen bonding and minimized by employing the OPLS4 force field [17]. The prepared protein quality and its ligand binding region were assessed and validated by the SiteMap application [18]. By incorporating catalytic residues (ASP197, GLU233, and ASP300) receptor grid was created. Further, the cognate ligand was re-docked to validate the target.

2.3 Molecular docking and MM/GBSA binding energy calculation

Glide, in extra precision mode (XP), was used in the molecular docking of natural products

with HPA [19]. Schrödinger's prime module was employed to calculate the MM/GBSA binding energy of protein and ligand complexes. The implicit VSGB solvation model [20] and OPLS4 force field were employed to calculate the binding energy.

2.4 Molecular dynamics simulation

A common technique in drug discovery is rigid receptor-based molecular docking. However, pitfalls exist in it. Therefore, to study the molecular dynamics behavior of the protein-ligand complex, the 200 ns molecular dynamics simulation was performed using the Academic Desmond 2019 GPU version [21]. The molecular system was solvated using the explicit solvation model (SPC) under orthorhombic periodic boundary conditions with a buffer region of 10 Å. The default OPLS_2005 force field was used [22]. Na^+ and Cl^- ions were used in the system builder to neutralize the system. The system was made more relaxed by doing a short-duration simulation and minimization. The Noose-Hoover thermostat method and Martyna-Tobias-Klien barostat method were employed in the NPT ensemble to maintain the temperature and pressure at 300 K and 1.01325 bar, respectively. A multiple-time step RESPA integrator was used in the dynamics study of bonded, near, and far bonded interactions.

2.5 ADMET analysis

QikProp of Schrödinger was used to calculate the pharmacokinetic characteristics of the best ligand [23]. Additionally, using admetSAR, ProTox-II, pkCSM, and the SwissADME web server, the hit's ADMET properties were analyzed [24–27].

3. Results and discussion

3.1 Protein preparation and target validation

Before performing molecular docking, human pancreatic α-amylase (HPA) was prepared and validated. The catalytic region's validity was confirmed using SiteMap. It predicted SiteScore, Dscore, volume, and balance of the most likely ligand-binding (catalytic) area to

be 0.985, 1.010, 341.285, and 0.751, respectively, which meet the almost threshold. TRP58, TRP59, TYR62, GLN63, HIE101, TYR151, LEU162, LEU165, ARG195, ASP197, ALA198, LYS200, HIS201, GLU233, VAL234, ILE235, HIE299, ASP300, and HIE305 residues were found in this area, confirming the catalytic region as the ligand-binding region (**Table S3, Fig S1**). Further, the target, RMSD of less than 2 Å, was validated by re-docking the cognate ligand (**Fig S1**). As a result, the SiteMap calculation, and cognate ligand re-docking all support the target protein's feasibility for further modeling.

3.2 Molecular docking and MM/GMBSA binding energy calculation

In this study, 38 molecules and acarbose were docked with HPA using Glide XP (extra precision mode). Then, MM/GBSA was employed to predict the binding energy between the protein and ligand complex. Based on the docking pose and score, the molecular docking showed that newboulaside B, newboulaside A, quercetin-3-O-β-glucoside, and sasastilboside A are the best molecules (**Table 1**). Among them, according to the literature, newboulaside B, newboulaside A, and sasastilboside A are all new molecules [28,29]. The MM/GBSA approach, among them, showed that newboulaside B is the best binding molecule.

Table 1. Glide molecular docking and MM/GBSA binding energy of ligands

SN	Compound	docking score (kcal/mol)	glide gscore (kcal/mol)	MM/GBSA dG Bind (kcal/mol)
1	acarbose	-12.581	-12.832	-59.77
2	newboulaside B	-11.366	-11.514	-50.29
3	newboulaside A	-10.533	-10.68	-19.38
4	quercetin-3-O-β-glucoside	-10.038	-10.067	-19.65
5	sasastilboside A	-9.191	-9.192	-36.58
6	ligularoside A	-8.910	-8.91	-23.11
7	mangiferin	-8.895	-9.274	-33.56

8	daucosterol	-8.411	-8.411	-22.33
9	kaempferol-3-O-*β*-glucoside	-8.031	-8.06	-28.07
10	rhododendron	-7.436	-7.436	-26.13
11	valoneic acid dilactone	-7.48	-7.687	-39.38
12	DDBT	-7.174	-7.275	-20.65
13	myricetin	-7.095	-7.132	-34.03
14	peruviose E	-6.894	-6.894	-23.57
15	peruviose B	-6.888	-6.888	-26.28
16	peruviose D	-6.881	-6.881	-44.62
17	chrysoeriol	-6.655	-6.695	-10.06
18	*α*-viniferin	-6.464	-6.464	-44.5
19	1-acetoxypinoresinol	-5.451	-5.451	-29.28
20	emodin	-5.23	-5.323	-17.72
21	momilactone B	-5.129	-5.129	-6.6
22	*β*-sitosterol	-4.972	-4.972	-28.93
23	3′-geranylchalconaringenin	-4.863	-5.362	-12.56
24	diosgenin	-4.786	-4.786	-33.8
25	momilactone A	-4.772	-4.772	-29.36
26	lupenone	-4.658	-4.658	-33.24
27	rostratazine B	-4.582	-4.582	-19.18
28	ursolic acid	-4.42	-4.426	-22.29
29	norchelerythrine	-4.414	-4.415	-36.82
30	cycloeucalenone	-4.314	-4.314	-6.33
31	skimmianine	-4.151	-4.164	-39.66
32	sesamine	-4.147	-4.147	-31.28

33	gedunin	-4.057	-4.057	-17.86
34	chaylbemide B	-4.038	-4.038	-8.13
35	oleanolic acid	-3.866	-3.872	-5.88
36	dehydrodieugenol B	-3.623	-3.633	-40.32
37	trans-fagaramide	-3.616	-3.616	-5.16
38	chaylbemide C	-3.452	-3.452	-11.23
39	chaylbemide A	-2.716	-2.716	-16.86

Key: DDBT -   2-(4-(3,5-dihydroxyphenoxy)-3,5-dihydroxyphenoxy) benzene-1,3,5-triol

The newboulaside B docking score and MM/GBSA binding energy with HPA were predicted to be -11.366 kcal/mol (acarbose = -12.581 kcal/mol) and -50.29 kcal/mol (acarbose = -59.77 kcal/mol), respectively. It interacted with active site residues (ASP197 and ASP300) as well as additional binding site residues (ASN105, ALA106, ASP147, THR163, and ARG195) via hydrogen bonding, according to the molecular docking (**Fig 1a, Fig 1b**).

Fig 1. Docked poses and molecular interactions of newboulaside B in the binding site of HPA: (a) 3D molecular surface diagrams, and (b) 2D ligand interactions of newboulaside B with HPA. **Glide XP molecular docking contributing term in the HPA and newboulaside B complex:** (c) HBOND (hydrogen bond) between HPA and newboulaside B (pink interaction) and (d) PhobEn (hydrophobic enclosure reward) of HPA (CPK gray), and PhobEn of newboulaside B (green ball and stick). The hydrophobic/hydrophilic map is represented as gray/brown patches.

The binding of HPA by newboulaside B was aided more by hydrogen bonds (-6.93) and hydrophobic enclosure reward (-0.25) than in the binding of HPA by acarbose (**Fig 1c, Fig 1d**). In comparison to other parameters, in the binding, the hydrogen bond contribution was found significant. So, we speculate that inhibition of HPA by newboulaside B might be due to the formation of hydrogen bonding with active site residues ASP197 and ASP300.

When the ligand changes, it undergoes structural remodeling, resulting in a change in binding affinity. The docking score and MM/GBSA binding energy of the HPA newboulaside A complex were predicted to be -10.533 kcal/mol and -19.38 kcal/mol, respectively. ASN53, TRP59, GLN63, ASP197, GLU233, and HIE305 interacted with newboulaside A via hydrogen bonding (**Fig 2a, Fig 2b**). TRP59, THR163, GLU233, and HIE305 formed hydrogen bonds with quercetin-3-O-β-glucoside, while TRP59 interacted via pi-pi interaction, with a docking score of -10.038 kcal/mol, and the MM/GBSA binding energy of -19.65 kcal/mol, respectively (**Fig 2c, Fig 2d**). Similarly, THR163, ARG195, GLU233, HIE299, and ASP300 interacted with sasastilboside A via hydrogen bond, with a docking score of -9.191 kcal/mol and the MM/GBSA binding energy of -36.58 kcal/mol, respectively (**Fig 2e, Fig 2f**).

Fig 2. Docked poses and molecular interactions of HPA with inhibitors: 3D molecular surface diagrams and 2D ligand interactions of newboulaside A (a) and (b), quercetin-3-O-β-glucoside (c) and (d), and sasastilboside A (e) and (f), respectively. The hydrogen bonds are represented as pink lines.

3.3 Molecular dynamics simulation

Molecular docking predicts the binding mode between a protein and a ligand. However, it is riddled with flaws and unable to foretell the stability and dynamic behavior of the molecule during the binding process. Understanding these by experimental means is also tedious. However, by simulating molecular activity in a computational setting, we can forecast experimentally challenging molecular behavior. Therefore, the top four HPA ligand (acarbose, newboulaside B, newboulaside A, quercetin-3-O-β-glucoside, and sasastilboside A) complexes from molecular docking were subjected to the 200 ns of 1000 frame molecular dynamics simulation to confirm the hit.

Monitoring of protein secondary structure elements (SSE) revealed that the majority of

strands and helices contributed 100% throughout the simulation and accounted for 33.84 percent of the total composition; the time versus residue index plot also revealed that strands near the binding pocket (ASP197, GLU233, and ASP300) contributed almost time, indicating the structure of HPA's stability. However, due to the loop, there may be a little more fluctuation around residues 130 and 370 (**Fig S2**).

The molecular dynamics simulation anticipated that newboulaside B would be remarkably stable with HPA. The backbone of HPA in the newboulaside B complex was deviated by less than 2 Å. After a sharp initial deviation to 1.75 Å, it nearly stabilized. Throughout the simulation, newboulaside B deviated around 1.5 Å (acarbose = 1.5 Å) in its alignment with itself and approximately 3 Å (acarbose = 3 Å) in its alignment with the protein (**Fig 3a**).

Fig 3. RMSD and RMSF analysis of MD simulation trajectory: The RMSD plot obtained for HPA (a) acarbose and newboulaside B complex, and (b) newboulaside A, quercetin-3-O-β-glucoside, and sasastilboside A complex. The RMSF plot was obtained for HPA (c) acarbose complex, and (d) newboulaside B complex. The strand, helix, and loop region of HPA are represented as blue, red, and white, respectively.

In the interaction region, the C-alpha residues of the HPA newboulaside B complex fluctuated about 1 Å, but in the HPA acarbose complex, after residue 300, they fluctuated largely (**Fig 3c, Fig 3d**). Throughout the simulation, newboulaside B fluctuated less than 1 Å (acarbose = 1.2 Å) in its alignment with itself and less than 1.8 Å (acarbose = 2 Å) in its alignment with protein (**Fig S3a, Fig S3b**). The stability of other compounds, such as newboulaside A, quercetin-3-O-β-glucoside, and sasastilboside A, was predicted to be much less than that of acarbose and newboulaside B. The observed RMSD of their fitting with protein was found more than 3.5 Å (**Fig 3b**). As a result, among top compounds, acarbose, and newboulaside B's stability with HPA may be comparable, as indicated by the aforementioned comparative deviations and fluctuations.

A minimum of 10 protein-ligand interactions, of which at least 5 were significant, were observed between HPA and acarbose as well as HPA and newboulaside B throughout the simulation (**Fig 4a, Fig 4c**). Acarbose interacted with ASP300, ASP197, and THR163 via hydrogen bonds over 70% of the simulated duration (**Fig 4b**). Whereas, newboulaside B interacted with residues (ASP197, ASP147, and ALA106) over 90% of the simulation period via hydrogen bonding. Further, catalytic residue (ASP300) also interacted via multiple hydrogen bonding (**Fig 4d**).

Fig 4. Analysis of inhibitor's molecular interaction and type of contact with HPA after MD simulations: Normalized stacked bar chart of HPA binding site residues interacting with (a) acarbose and (c) newboulaside B via hydrogen bonds, hydrophobic and ionic interactions, and water bridges. Detailed schematic interaction of (b) acarbose and (d) newboulaside B atoms with the binding site residues of HPA. Interactions happening more than 30% of the simulation times are shown.

The timeline charts also showed that at least three bonds in the HPA complex of acarbose and newboulaside B are almost unbroken and interacted significantly throughout the simulation. THR163, ASP197, and ASP300 interacted via more or less uninterrupted bonding with acarbose, while TRP59 and GLU233 interacted via partial bonding (**Fig S4a**). Similarly, newboulaside B interacted via nearly continuous bonding with ALA106, ASP147, and ASP197, and partial bonding with GLN63, ARG161, THR163, ASP300, and ASP356 (**Fig S4b**). These interactions might help to prevent them from diffusing out from the ligand-binding pocket. This evidence substantiates the hypothesis that newboulaside B inhibits HPA. However, newboulaside A, quercetin-3-O-β-glucoside, and sasastilboside A demonstrated fewer interactions with active site residues than newboulaside B, according to the protein-ligand interaction analysis (**Fig S5a, Fig S5c, Fig S5e**). Further, the HPA complexes of these three were more unstable than the complex of newboulaside B, according to the ligand-

protein interaction diagram of MD simulation (**Fig S5b, Fig S5d, Fig S5f**). Overall, they exhibited significantly poorer interactions with HPA than acarbose and newboulaside B, as evidenced by the high deviations and weak interactions.

The outcome of the HPA newboulaside B complex's molecular docking and MM/GBSA binding energy is supported by the molecular dynamics simulation, too. Thus, based on comparative stability, binding affinity, and interactions with acarbose, newboulaside B is expected to be effective in inhibiting HPA.

Existing knowledge also reinforces our computational analysis. *Newbouldia laevis* (P. Beauv.), which is used in African traditional medicine to treat a variety of illnesses such as sickle cell, diarrhea, inflammation, aches, tumors, skin problems, hypertension, and diabetes, is reported to be the natural source of newboulaside B [30]. Ethanolic extracts and aqueous root extracts of the plant exhibited anti-diabetic activity in streptozocin-induced diabetic Wistar rats and alloxan-induced diabetic rats, respectively [31,32]. Ethanolic extracts and aqueous leaf extracts of the plant reported antioxidant activity in diabetic rats and insulin-inducing activity [33,34]. The methanolic extract of the plant suppressed induced diabetes by 60.2 percent [35].

Two new caffeic acid glycosides (newboulaside A and newboulaside B) isolated from ethanolic extracts of the *Newbouldia laevis*'s leaves inhibited α-amylase with IC_{50} values of 4.95 and 4.44 µg/mL, respectively (acarbose IC_{50} = 4.05 µg/mL) [28]. The aforementioned computational data also supports the binding nature of these compounds to HPA.

3.4 ADMET analysis

The drug should be highly selective and have as few side effects as possible. Due to effectiveness and unavoidable side effects, the therapeutic candidate has a high attrition rate at the late stage. ADMET problems arise due to the complexity of the biological system. The *in silico* study can help to reduce the risk of a drug candidate failure. As a result, as post-

processing, identified hit (newboulaside B) drug-likeness and ADMET properties were analyzed comprehensively.

The ethanolic extract of *Newbouldia laevis* was proven to be safe in terms of acute toxicity [36]. The assessment of the ethanolic extract of the plant revealed a low toxicity profile [37]. This finding supports the traditional uses of the plant against diabetes. However, research on the pure molecule was lacking. As a result, several models, including QikProp, admetSAR, ProTox-II, pkCSM, and SwissADME, were utilized to analyze the hit's pharmacokinetic properties (**Table S4 - Table S8**).

The QikProp computed lipophilicity (QPlogPo/w) of the hit -2.995 (-2.0 to 6.5), which indicates that the hit is more hydrophilic; this low lipophilicity may decrease absorption and distribution due to the lipoid nature of the cell membrane. Similarly, it predicted that hit is aqueous soluble with solubility value -2.52(-6.5 to 0.5) [38,39]. The water solubility of the hit is due to the hydroxyl group, which contributes to absorption. The hit is comparable to acarbose, according to molecular modeling; however, it defies recognized drug-likeness, it is partly due to its large molecular size.

The SwissADME projected that the hit would have poor intestinal absorption. Similarly, it predicted the hit as less skin permeable with LogKp value of -12.08 cm/s (acarbose = -16.30 cm/s). The pkCSM predicted CNS permeability (logPS) -5.747 (logPS<-3), which indicates that the hit is impermeable to CNS. The strong polarity, high molecular weight, and poor lipophilicity of this compound may be the cause of its limited absorption. QikProp's low human serum albumin binding value of the hit (-2.173) indicates the unbound molecule can easily reach the target [40]. As a result of the large molecular size, the hit may be moderately bioavailable on average.

The ProTox-II model projected the hit toxicity would be class V (slightly toxic), with an LD_{50} value of 5000 mg/kg, and also predicted that it is inactive for carcinogenicity, mutagenicity,

and cytotoxicity. Additionally, it was evaluated with negative mitochondrial, carcinogenicity, and nephrotoxicity by the admetSAR, which indicates it is not acutely toxic.

The hit was predicted as negative skin sensitization by the admetSAR, which reveals that the molecule would not show a cutaneous inflammatory reaction. It also predicted that the hit would not inhibit CYP 2D6, CYP 1A2, CYP 2C19, CYP 2C9, and CYP 3A4. This property helps to metabolize xenobiotics and reduce the adverse consequences of drug-drug interactions. Similarly, the admetSAR projected that the hit would be the non-inhibitor of OCT2, OATP2B1, and MATE1. Non-inhibition of OCT2, OATP2B1, and MATE1 help to distribute and excrete the second drug and reduces drug-drug interaction detrimental. Overall ADMET profile of newboulaside B might be good. However, further wet-lab research is necessary to support pharmacokinetics predictions.

4. Conclusions

The results of the current study imply that the computational approach effectively explores the potent human pancreatic α-amylase (HPA) inhibitor. Herein, the computational workflow, with almost similar binding affinity and stability to acarbose, identifies newboulaside B as the potential molecule for HPA inhibition. The study also shows a good ADMET profile for the molecule and reinforces prior reports. To further validate the outcome, wet-lab testing must be performed. The data presented here can be helpful for drug discovery against type 2 diabetes.

Data Accessibility

The data are provided in the electronic supplementary material.

Author's Contribution

Santosh Basnet: Conceptualization, Methodology, Investigation, Data Curation, Formal Analysis, Writing-original Draft, Funding Acquisition, Software, Validation; **Madhav Prasad Ghimire:** Resources for the Simulation, Technical Support in the Simulation,

Writing Reviewing and Editing, Formal Assessment; **Tika Ram Lamichhane:** Technical Support in the Simulation, Writing Reviewing and Editing, Formal Assessment; **Rajendra Adhikari:** Resources for the Simulation, Technical Support in the Simulation, Writing Reviewing and Editing, Formal Assessment; **Achyut Adhikari:** Conceptualization, Supervision, Writing Reviewing and Editing, Funding Acquisition, Software, Project Administration and Formal Assessment

Compliance with ethics requirement

There are no animal models used in this research.

Competing interest

The authors declare that there are no competing interests.

Funding

This work was supported by the University Grants Commission, Bhaktapur, Nepal, as part of a thesis grant fellowship award (MRS-77/78-S&T-36).

Acknowledgments

The author acknowledges Schrödinger Inc. for the software license support for the thesis research work. Similarly, the author acknowledges the Central Department of Physics, Tribhuvan University for providing the supercomputing facility supported by the Alexander von Humboldt Foundation, and IFW-Dresden, Germany. S. Basnet expresses gratitude to Schrödinger Inc. for providing training and fellowship in computational antibody engineering, rational drug design, and high-throughput virtual screening. R. Adhikari acknowledges NVIDIA for GPU grants. A. Adhikari gratefully acknowledges the Higher Education Reform Project (HERP) of Tribhuvan University for the startup grant.

Supplementary Information

Table S1-Table S8 and Fig S1-Fig S5's data are provided in the supplementary material.

References

1. Farooq MU, Mumtaz MW, Mukhtar H, Rashid U, Akhtar MT, Raza SA, Nadeem M. UHPLC-QTOF-MS/MS based phytochemical characterization and anti-hyperglycemic prospective of hydro-ethanolic leaf extract of *Butea monosperma*. Sci Rep. 2020; 10: 1-14. https://doi.org/10.1038/s41598-020-60076-5

2. American Diabetes Association. Classification, and diagnosis of diabetes: standards of medical care in diabetes-2019. Am Diabetes Assoc. 2019; 42. https://doi.org/10.2337/dc19-S002

3. Ponnusamy S, Ravindran R, Zinjarde S, Bhargava S, Ravi Kumar A. Evaluation of traditional Indian antidiabetic medicinal plants for human pancreatic amylase inhibitory effect in vitro. Evid Based Complementary Altern Med. 2010; 2011. https://doi.org/10.1155/2011/515647

4. Brayer GD, Luo Y, Withers SG. The structure of human pancreatic α-amylase at 1.8 Å resolution and comparisons with related enzymes. Protein Sci. 1995; 4: 1730–1742. https://doi.org/10.1002/pro.5560040908

5. Brayer GD, Sidhu G, Maurus R, Rydberg EH, Braun C, Wang Y, Nguyen NT, Overall CM, Withers SG. Subsite mapping of the human pancreatic *α*-amylase active site through structural, kinetic, and mutagenesis techniques. Biochem. 2000; 39: 4778-4791. https://doi.org/10.1021/bi9921182

6. Krentz AJ, Bailey CJ. Oral antidiabetic agents. Drugs. 2005; 65: 385–411 https://doi.org/10.2165/00003495-200565030-00005

7. American Diabetes Association. Standards of medical care in diabetes-2018 abridged for primary care providers. Clin Diabetes. 2018; 36: 14-37. https://doi.org/10.2337%2Fcd17-0119

8. Lipscombe L, Booth G, Butalia S, Dasgupta K, Eurich DT, Goldenberg R, Khan N, MacCallum L, Shah BR, Simpson S. Pharmacologic glycemic management of type 2 diabetes in adults. Can J Diabetes. 2018; 42. https://doi.org/10.1016/j.jcjd.2017.10.034

9. Akash MSH, Shen Q, Rehman K, Chen S. Interleukin-1 receptor antagonist: a new therapy for type 2 diabetes mellitus. J Pharm Sci. 2012; 101: 1647–1658. https://doi.org/10.1002/jps.23057

10. Shapiro K, Gong WC. Natural products used for diabetes. J Am Pharm Assoc. 2002; 42: 217-226. https://www.sciencedirect.com/science/article/abs/pii/S1086580215302989

11. Tundis R, Loizzo MR, Menichini F. Natural products as α-amylase and α-glucosidase inhibitors and their hypoglycaemic potential in the treatment of diabetes: an update. Mini-Rev Med Chem. 2010; 10: 315–331.
https://doi.org/10.2174/138955710791331007

12. Schrödinger Release 2021-4: LigPrep, Schrödinger, LLC, New York, NY, 2021.

13. Lu C, Wu C, Ghoreishi D, Chen W, Wang L, Damm W, Ross GA, Dahlgren MK, Russell E, Von Bargen CD, Abel R. OPLS4: Improving force field accuracy on challenging regimes of chemical space. J Chem Theory Comput. 2021; 17: 4291–4300. https://doi.org/10.1021/acs.jctc.1c00302

14. Maurus R, Begum A, Williams LK, Fredriksen JR, Zhang R, Withers SG, Brayer GD. Alternative catalytic anions differentially modulate human α-amylase activity and specificity. Biochem. 2008; 47: 3332–3344. https://doi.org/10.1021/bi701652t

15. Schrödinger Release 2021-4: Protein Preparation Wizard, Schrödinger, LLC, New York, NY, 2021.

16. Schrödinger Release 2021-4: Prime, Schrödinger, LLC, New York, NY, 2021.

17. Madhavi Sastry G, Adzhigirey M, Day T, Annabhimoju R, Sherman W. Protein and ligand preparation: parameters, protocols, and influence on virtual screening enrichments. J Comput Aided Mol Des. 2013; 27: 221–234. https://doi.org/10.1007/s10822-013-9644-8

18. Halgren TA. Identifying and characterizing binding sites and assessing druggability. J Chem Inf Model. 2009; 49: 377–389. https://doi.org/10.1021/ci800324m

19. Friesner RA, Murphy RB, Repasky MP, Frye LL, Greenwood JR, Halgren TA, Sanschagrin PC, Mainz DT. Extra precision glide: docking and scoring incorporating a model of hydrophobic enclosure for protein−ligand complexes. J Med Chem. 2006; 49: 6177–6196. https://doi.org/10.1021/jm051256o

20. Li J, Abel R, Zhu K, Cao Y, Zhao S, Friesner RA. The VSGB 2.0 model: a next-generation energy model for high-resolution protein structure modeling. Proteins: Str Fun Bioinform. 2011; 79: 2794–2812. https://doi.org/10.1002/prot.23106

21. Bowers KJ, Chow DE, Xu H, Dror RO, Eastwood MP, Gregersen BA, Klepeis JL, Kolossvary I, Moraes MA, Sacerdoti FD, Salmon JK. Scalable algorithms for molecular dynamics simulations on commodity clusters. SC '06: Proceedings Of The 2006 ACM/IEEE Conference On Supercomputing. 2006; 43–43. https://doi.org/10.1109/SC.2006.54

22. Banks JL, Beard HS, Cao Y, Cho AE, Damm W, Farid R, Felts AK, Halgren TA, Mainz DT, Maple JR, Murphy R. Integrated modeling program, applied chemical theory (impact). J Comput Chem. 2005; 26: 1752–1780. https://doi.org/10.1002/jcc.20292

23. Schrödinger Release 2021-4: QikProp, Schrödinger, LLC, New York, NY, 2021.

24. Yang H, Lou C, Sun L, Li J, Cai Y, Wang Z, Li W, Liu G, Tang Y. AdmetSAR 2.0: web-service for prediction and optimization of chemical ADMET properties. Bioinform. 2019; 35: 1067–1069. https://academic.oup.com/bioinformatics/article/35/6/1067/5085368

25. Banerjee P, Eckert AO, Schrey AK, Preissner R. ProTox-II: a webserver for the prediction of toxicity of chemicals. Nucleic Acids Res. 2018; 46. https://academic.oup.com/nar/article/46/W1/W257/4990033

26. Pires DEV, Blundell TL, Ascher DB. pkCSM: predicting small-molecule pharmacokinetic and toxicity properties using graph-based signatures. J Med Chem. 2015; 58: 4066–4072. https://doi.org/10.1021/acs.jmedchem.5b00104

27. Daina A, Michielin O, Zoete V. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. Sci Rep. 2017; 7: 42717. https://www.nature.com/articles/srep42717

28. Mbagwu IS, Akah PA, Ajaghaku DL, Ike JC, Okoye FBC. Newboulasides A and B, two new caffeic acid glycosides from *Newbouldia laevis* with α-amylase inhibitory activity. Nat Prod Res. 2020; 36: 1–9. https://doi.org/10.1080/14786419.2020.1799362

29. Cuc NT, Cuong NT, Anh LT, Yen DT, Tai BH, Thu Trang D, Yen PH, Kiem PV, Nam NH, Minh CV, Nhiem NX. Dihydrostilbene glycosides from *Camellia sasanqua* and their α-glucosidase and α-amylase inhibitory activities. Nat Prod Res. 2020; 1–7. https://doi.org/10.1080/14786419.2020.1756802

30. Okagu IU, Ndefo JC, Agbo MO. Trado-medical uses, chemical constituents and biological activities of *Newbouldia laevis* (Bignoniaceae): a review. Pharm Sci. 2021; 28: 51-59. https://doi.org/10.34172/PS.2021.29

31. Tanko Y. Anti-diabetic effect of ethanolic flower extract of *Newbouldia laevis* (Bignoniaceae) on blood glucose level of Streptozotocin-induced diabetic Wistar rats. Res J Med Sci. 2008; 2: 57–60.

32. Okonkwo P, Okoye Z. Hypoglycaemic effects of the aqueous extract of *Newbouldia laevis* root in rats. Int J Biol Chem Sci. 3: 998-1004. https://doi.org/10.4314/ijbcs.v3i5.51078

33. Anaduaka EG, Ogugua VN, Agu CV, Okonkwo CC. Ethanol extracts of *Newbouldia laevis* stem and leaves modulate serum liver marker enzymes and antioxidant enzymes activities in diabetic rats. Afr J Biotechnol. 2014; 13: 2265-2275. https://doi.org/10.4314/ajb.v13i22

34. Igbokwe UV, Eze ED, Adams MD, Atsukwei D. *Newbouldia laevis* leaf facilitates insulin secretion, glucose uptake, and pancreatic function in alloxan-induced hyperglycemia in Wistar. Int J Pharm Sci Res. 2018; 9: 5079-5085. http://dx.doi.org/10.13040/IJPSR.0975-8232.9(12).5079-85

35. JA B, S S, IU A. Antidiabetic and hypolipidemic effects of methanol leaf extract of *Newbouldia laevis* in alloxan-induced diabetic rats. Adv Res Gastroenterol Hepatol. 2019; 13: 56-62. http://dx.doi.org/10.19080/ARGH.2019.13.555863

36. Owolabi OJ, Amaechina FC, Okoro M. Effect of ethanol leaf extract of *Newboulda Laevis* on blood glucose levels of diabetic rats. Trop J Pharm Res. 2011; 10: 249-254. https://doi.org/10.4314/tjpr.v10i3.12

37. Kolawole OT, Akanji MA, Akiibinu MO. Toxicological assessment of ethanolic extract of the leaves of *Newbouldia laevis* (P. Beauv). Am J Med Sci. 2013; 3: 74-80. doi:10.5923/j.ajmms.20130304.04

38. Choudhary MI, Shaikh M, tul-Wahab A, ur-Rahman A. In silico identification of potential inhibitors of key SARS-CoV-2 3CL hydrolase (Mpro) via molecular docking, MMGBSA predictive binding energy calculations, and molecular dynamics simulation. PLOS ONE. 2020; 15. https://doi.org/10.1371/journal.pone.0235030

39. Jorgensen WL, Duffy EM. Prediction of drug solubility from structure. Adv Drug Del Rev. 2002; 54: 355–366. https://doi.org/10.1016/S0169-409X(02)00008-X

40. Sun L, Yang H, Li J, Wang T, Li W, Liu G, Tang Y. In silico prediction of compounds binding to human plasma proteins by QSAR models. ChemMedChem. 2018; 13: 572–581. https://doi.org/10.1002/cmdc.201700582

Supplementary Captions

Table S1. Ligand library of natural products

Table S2. Human pancreatic *α*-amylase

Table S3. SiteMap result

Table S4. ADMET properties of the acarbose and newboulaside B by QikProp

Table S5. ADMET properties of newboulaside B and acarbose by admetSAR

Table S6. Toxicity profile of newboulaside B and acarbose by ProTox-II

Table S7. ADMET properties of newboulaside B (1) and acarbose (2) by pkCSM

Table S8. ADMET properties of newboulaside B and acarbose by SwissADME

Fig S1. Pictorial representation of HPA with cognate ligand (green) and re-docked cognate ligand (white). The yellow and blue patches represent the SiteMap application view.

Fig S2. Analysis of secondary structure element (SSE) of HPA after MD simulation: The strand, helix, and loop region of HPA are represented as blue, red, and white, respectively.

Fig S3. RMSF of inhibitor's atoms after MD simulation: (a) acarbose and (b) newboulaside B

Fig S4. Timeline representation of protein-ligand contact of MD simulation trajectory: (a) HPA acarbose complex (b) HPA newboulaside B complex

Fig S5. Analysis of inhibitor's molecular interaction and type of contact with HPA after MD simulations: Normalized stacked bar chart of HPA binding site residues

interacting with (a) newboulaside A, (c) quercetin-3-O-*β*-glucoside, and (e) sasastilboside A via hydrogen bonds, hydrophobic and ionic interactions and water bridges. Detailed schematic interaction of (b) newboulaside A, (d) quercetin-3-O-*β*-glucoside, and (f) sasastilboside A atoms with the binding site residues of HPA. Interactions happening more than 30% of the simulation times are shown.

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure