# Allelic variations and gene cluster modularity act as non-linear bottlenecks for cholera emergence

Mario López-Pérez[a,b#], Deepak Balasubramanian[a#], Cole Crist[a], Trudy-Ann Grant[a], Jose M. Haro-Moreno[a,b], Asier Zaragoza-Solas[a,b], Salvador Almagro-Moreno[a*]

[a]Burnett School of Biomedical Sciences, College of Medicine, University of Central Florida, Orlando, Florida, USA; [b]Evolutionary Genomics Group, División de Microbiología, Universidad Miguel Hernández, Alicante, Spain.

\# Contributed equally

*Address correspondence to: Salvador Almagro-Moreno, samoreno@ucf.edu

Running title: Evolutionary bottlenecks of pathogen emergence

**Keywords**:

Pathogen emergence / *Vibrio cholerae* / modularity / population genomics / allelic variation / environmental reservoirs

**ABSTRACT**

Cholera, an acute secretory diarrhea, is caused by strains from a phylogenetically confined group within the *Vibrio cholerae* species, the pandemic cholera group (PCG). To date, the molecular and evolutionary factors that enable the isolated emergence of toxigenic *V. cholerae* from environmental populations remain mostly enigmatic. Comprehensive analyses of over 1,100 *V. cholerae* genomes, including novel environmental isolates from this study, reveal that the species consists of four major clades and several minor ones. PCG belongs to a large clade located within a lineage shared with environmental strains, the pandemic cholera lineage. This hierarchical classification provided us with a framework to unravel the eco-evolutionary dynamics of the genetic determinants associated with the emergence of toxigenic *V. cholerae*. Our analyses indicate that this phenomenon is largely dependent on the acquisition of unique modular gene clusters and allelic variations that confer a competitive advantage during intestinal colonization. We determined that certain PCG-associated alleles are essential for successful colonization whereas others provide a non-linear competitive advantage, acting as a critical bottleneck that elucidates the isolated emergence of PCG. For instance, toxigenic strains encoding non-PCG alleles of a) *tcpF* or b) a sextuple allelic exchange mutant for genes *tcpA*, *toxT*, *VC0176*, *VC1791*, *rfbT* and *ompU*, lose their ability to colonize the intestine. Interestingly, these alleles do not play a role in the colonization of model environmental reservoirs. Our study uncovers the evolutionary roots of toxigenic *V. cholerae* and offers a tractable approach for investigating the emergence of pathogenic clones within an environmental population.

**SIGNIFICANCE**

The underlying factors that lead to specific strains within a species to emerge as human pathogens remain mostly enigmatic. Toxigenic clones of the cholera agent, *Vibrio cholerae*, are encompassed within one phylogenomic clade, the pandemic cholera group (PCG). Here, we investigate the molecular and evolutionary factors that explain the confined nature of this group. Our analyses determined that the emergence of PCG is largely dependent on the acquisition of unique modular gene clusters and allelic variations that confer a competitive advantage during intestinal colonization. These allelic variations act as a critical bottleneck that elucidates the isolated emergence of PCG and provides a tractable blueprint for the study of the emergence of pathogenic clones within an environmental population.

## INTRODUCTION

The emergence of human pathogens is one of the most pressing public health concerns of modern times, with new outbreaks of varying magnitude occurring regularly around the world[1–4]. Pathogen emergence is a complex and multifactorial phenomenon that culminates with a microorganism acquiring the ability to colonize and harm the human host, cause outbreaks, and persist within the environment[5]. To date, the specific factors and evolutionary constraints that govern the emergence of bacterial pathogens from environmental populations remain mostly enigmatic[4]. Elucidating how the interplay between the numerous molecular and environmental drivers lead to this complex phenomenon is critical to develop surveillance platforms to predict emergent events and potential sources of outbreaks[4]. Furthermore, in addition to the profound public health implications, understanding pathogen emergence addresses one of the central questions of microbial ecology: what are the evolutionary processes within a bacterial population that underlie colonization and adaptation to a new environment?[6].

The Vibrionaceae is a highly diverse family of aquatic bacteria that encompass several species representing distinct paradigms of how pathogens can emerge from environmental populations[7,8]. *Vibrio cholerae* is the most widely studied member of the Vibrionaceae and has been a model organism for numerous molecular, ecological, and epidemiological studies[9–12]. Toxigenic *V. cholerae* is the causative agent of the severe diarrheal disease cholera, a scourge that affects millions of people each year and is responsible for over 100,000 deaths annually[9]. The first six pandemics of cholera are thought to have been caused by the Classical biotype *V. cholerae* O1, a strain that is now considered extinct[9,13–15]. The seventh and current pandemic is caused by the El Tor biotype of *V. cholerae* O1 and started in Indonesia in 1961[16,17]. The disease typically affects nations with limited access to clean water and poor sanitation and can be lethal if not properly treated[9,14]. Interestingly, only a limited number of *V. cholerae* strains can cause pandemic cholera[18–20]. Specifically, toxigenic strains of *V. cholerae* are confined to a phylogenetic group, what we term the pandemic cholera group

(PCG)[19,21–23]. Even though the mechanisms of cholera pathogenesis in humans have

been extensively studied over the last several decades, the evolutionary events and

molecular constraints that explain why only members from PCG can cause pandemic

cholera remains unknown[24–29].

Some of the major general events that are essential for the emergence of

toxigenic *V. cholerae* from environmental populations include the acquisition of the

lipopolysaccharide (LPS) O1 antigen cluster and mobile genetic elements (MGEs) such

as CTXΦ phage, Vibrio pathogenicity Island-1 (VPI-1) and Vibrio pathogenicity Island-2

(VPI-2)[15]. Among several other differences with Classical, El Tor strains have also

acquired two unique pathogenicity islands (PAIs): *Vibrio* Seventh Pandemic (VSP) I

and II. CTXΦ encodes the genes for the cholera toxin, which is responsible for the

severity of the profuse diarrhea associated with cholera[30]. The toxin coregulated pilus

(TCP), an essential colonization factor, is encoded within VPI-1[31,32], which also

encodes several major virulence regulators such as ToxT and TcpP[32,33]. VPI-2 encodes

the genes for sialic acid utilization, which confers a competitive advantage in the gut[34].

VSP-I codes for a regulator that plays a role in intestinal colonization and chemotaxis[35]

and VSP-II is associated with environmental survival and fitness of the bacterium[36–38]. It

was recently found that VSP-II and VPI-2 encode systems that prevent the uptake of

foreign genetic material[39]. These MGEs are always found in the PCG but are not

exclusive to them as they are also encoded in some environmental strains of *V.*

*cholerae*[40–42]. However, to date, their abundance, specific distribution, and dynamics

within the *V. cholerae* species are not well understood.

Horizontal acquisition of key virulence genes is a critical step in the emergence

of toxigenic clones of *V. cholerae*, however, it is not sufficient to explain why the ability

to cause cholera is limited to strains belonging to the PCG. This phylogenetically

confined distribution together with the presence of CTXΦ and the PAIs in some

environmental strains strongly indicate that other barriers and requirements exist that

stringently limit their emergence. We recently investigated the genomic prerequisites

that must be present in a population before a pathogenic clone can emerge from an

environmental gene pool[22]. We determined that the environmental ancestor of the PCG

had a particular genomic background containing a specific set of alleles of core genes

that conferred preadaptations to virulence and enhanced its pathogenic potential, what

we term virulence adaptive polymorphisms (VAPs)[22]. Interestingly, our data shows that

VAPs are present in environmental populations and confer virulence-associated traits

to strains encoding them. Overall, our results indicate that there are core genomic

factors beyond the acquisition of MGEs that are critical for the emergence of toxigenic

*V. cholerae*. To date, the unique core genes, and allelic variations in the genomic

background of toxigenic *V. cholerae* that explain their narrow distribution remain

enigmatic.

In this study, we investigated the evolutionary constraints that govern the

emergence of bacterial pathogens from environmental populations using toxigenic *V.*

*cholerae* as a model system. To do this, first, we isolated environmental *V. cholerae*

strains from the Indian River Lagoon (IRL), a highly diverse and expansive estuary

located in Eastern Florida. Genome comparisons of these strains against *V. cholerae*

PCG unveil a phylogeny in which strains form cluster-like groups. We extended our

analyses to over 1,100 publicly available genomes and found that *V. cholerae* strains

are divided into four major clusters and numerous other minor clades. We found that

PCG belong to one large cluster (C24) and is located within one lineage shared with

other environmental strains, the pandemic cholera lineage (PCL). Subsequently, we

examined the genetic determinants associated with the emergence of toxigenic *V.*

*cholerae* through multi-level genomic analyses by investigating **a)** the distribution and

abundance of the O1 LPS cluster and main virulence associated MGEs, **b)** the

evolutionary dynamics of the gene clusters that comprise these MGEs, and **c)** the

allelic diversity within the population. We found that these elements are made up of

several modules sparsely distributed among the different phylogenomic groups and

include small gene clusters and allelic variations. We determined that these allelic

variations confer a competitive advantage during intestinal colonization, acting as a

critical bottleneck that elucidates the isolated emergence of PCG, but do not play a role

in the colonization of model environmental reservoirs. Overall, our study provides a

multi-level scenario that serves as a blueprint for the study of the evolution of

facultative bacterial pathogens.

**RESULTS AND DISCUSSION**

**Isolation of environmental *Vibrio cholerae* from the Indian River Lagoon.** *V.*

*cholerae* is a natural inhabitant of aquatic ecosystems where it can be found as free-

living or attached to abiotic or biotic surfaces[43,44]. We recently established two sampling

sites along the eastern coast of Florida where we isolated pathogenic strains of *Vibrio*

*vulnificus*[45]. We sampled those two sites, Feller's House (FH) and Shepard Park (SP),

to investigate the potential presence of *V. cholerae* strains (**Fig. 1A**). Using the

isolation, enrichment and selection methods described below, we identified a total of

178 potential *V. cholerae* isolates from the different fractions at both locations during

our sampling events (42 FH and 136 SP). To confirm those isolates, we performed a

PCR screening for the species-specific gene *ompW*[46]. Only 65 isolates were confirmed

as *bona fide V. cholerae*, all of them isolated from SP (**Fig. 1A**). *V. cholerae* isolates

from SP were collected from all the fractions that we examined (sediment,

cyanobacteria, particle-associated and water) except for oysters. Subsequently, we

performed PCR-based fingerprinting of the strains to reduce redundancies and

minimize clonality. We used a combination of PCR-based fingerprinting analyses,

VCR-PCR and ERIC-PCR[47]. Based on fingerprinting patterns and cluster analysis

performed using BIONUMERICS v8.0 we generated a dendrogram to group these

observed patterns. Using this resulting dataset, we selected 35 isolates with unique

fingerprinting patterns for whole genome sequencing (**Fig. S1 and Table S1**).

**Phylogenomic characterization and clustering of *V. cholerae* strains.** The 35 *V. cholerae* strains that we sequenced were isolated from a diverse set of fractions: 6 from sediment, 10 from cyanobacteria, 9 from the free-living water fraction and 10 from the particle-associated one (**Table S1**). We examined the relationship of these strains with *V. cholerae* PCG by analyzing the average nucleotide identity (ANI) in a pairwise comparison against *V. cholerae* N16961, a reference O1 strain from the El Tor biotype (**Fig. S2**) [48]. The minimum values obtained were ca. 97.5, always within the established cut-off for strains of the same species. However, up to nine environmental strains showed a greater divergence with the PCG (ANI >98.2%) than with the rest of the strains (**Fig. S2**). Interestingly, phylogenomic analysis based on whole-genome single nucleotide polymorphism (SNP) (230,824 SNPs identified in 3.19 Mb of core genome) show that the majority of the IRL strains form two major clusters, one of them including the toxigenic strain of *V. cholerae* (**Fig. 1B**). This evolutionary relationship shares similarity with that of *V. vulnificus*, in which the pathogenic strains capable of causing septicemia in humans belong to one clearly defined cluster[49]. This prompted us to further characterize the *V. cholerae* species to investigate whether the evolution of the toxigenic strains follow a similar cluster-based pattern, as this would shed important light on the emergence and evolution of the pathogen.

First, we collected the *V. cholerae* genomes available in public databases (based on NCBI classification accessed in July 2020; see Materials and Methods) together with our environmental isolates. Naturally, there is a significant bias in the databases towards *V. cholerae* O1 and O139 strains, which are mostly clonal. Therefore, in order to obtain an unbiased phylogeny, we dereplicated all these genomes to an identity of ANI >99%. From a total of 1,127 initial genomes, we obtained 227 non-clonal reference *V. cholerae* clades (**Table S2**). To maximize the consistency of our analyses, we built both a SNP-based phylogeny and genome clustering using ANI to determine the groups that constitute the *V. cholerae* species. The application of a divergence threshold of less than 97% in ANI results in a single cluster indicating that

the boundary of intra-population sequence diversity in *V. cholerae* species is 97%.

However, clustering using a threshold of 98% of ANI shows that the species

encompasses 34 clusters that include *V. cholerae* strains (**Table S3**). Next, we used a

total of 458,798 SNPs from a core genome size of 2.46 Mb to build the species

phylogeny using non-clonal reference clades and IRL strains (**Fig. 1C**). The species

phylogeny topology based on SNPs or ANI clustering are virtually identical, with some

minor exceptions that might be due to certain strains having a convergent history.

Interestingly, both approximations reveal that four major clusters account for 69.2% of

the non-clonal clades (157 out of a total of 227) and 31 of the 35 IRL isolates from this

study (**Table S3**). All IRL isolates are grouped together with previously isolated strains

except for strains IRLE0050 and IRLE0074 which formed a separate cluster (**Table**

**S3**). Cluster 24 (C24) is the largest of the four main clusters containing 68 non-clonal

clades including PCG. On the other hand, most IRL strains belong to C5 (22 of the 35)

(**Fig. 1C**). We did not find a pattern that associate the phylogeny of the IRL isolates and

their fraction of isolation. The C24 phylogeny also reveals that PCG groups with nine

other non-clonal clades, including IRLE0178 from this study, fall into a separate branch

in the phylogenomic tree, what we have termed the Pandemic Cholera Lineage (PCL)

(**Fig. 1C**). This provides us with a novel hierarchical approach based on three levels to

decipher the evolutionary history of toxigenic strains of *V. cholerae* (from highest to

lowest): a) the C24 within all *V. cholerae* strains, b) the PCL within C24 non-clonal

clades and c) the clonal *V. cholerae* toxigenic strains within the PCL lineage.

**Lipopolysaccharide (LPS) cluster**.   The LPS plays numerous roles in the

pathogenesis of toxigenic *V. cholerae* (e.g., endotoxic activities, immunological

responses, etc.[50–52]) and is critical to determine the antigenic properties and

classification of the bacterium[52]. Specifically, given that only O1 strains, together with

the extinct O139, can cause pandemic cholera, we examined its LPS cluster to

determine its distribution, evolutionary history, and allelic diversity within the *V.*

*cholerae* species (**Fig. 2**). The O1 gene cluster is 29.7 Kb long and encompasses

approximately 26 ORFs with potential or demonstrated function (some are truncated

preventing an exact number). The *rfa* gene cluster (LPS core synthesis) is encoded at

the 5' end of the island (**Fig. 2A** blue arrows) and is widely distributed across all clades,

with the notable exception of RfaF (ADP-heptose--LPS heptosyltransferase 2,

VC_0235) and RfaL (O-antigen ligase, VC_0237) present in ca. 35 non-clonal clades.

Nonetheless, the genes in the *rfa* cluster exhibit the highest divergence within the LPS

cluster (median identity 86-88%) and accumulate numerous synonymous

polymorphisms (**Fig. 2A**). This indicates that, like in *V. vulnificus*, the *rfa* cluster

appears to be a preferred site to act as hot spots for the recombination of this island

due to the conservation and synteny of these genes[49]. The distribution of the central

region of the LPS cluster is much more restricted and is only present in 17 *V. cholerae*

non-clonal clades, however, those that encode it share 100% identity with the region of

PCG. This region contains the *rfb* cluster which is associated with antigen-O

synthesis[53] (**Fig. 2A,** green arrows). Interestingly, only 5 out the 17 clades that encode

this region belong to the PCL, indicating that there is no direct correlation between the

phylogeny and its presence in PCG. The LPS cluster of PCG encodes four

transposases, which likely increase the variability within the island (**Fig. 2A and 2B**).

These IS elements are widely distributed among *V. cholerae* strains providing insight

into how small gene clusters can be introduced in the island. For instance, even though

strains VC22 (C24) and 2479-86 (C27) encode the same version of the PCG LPS

cluster (identity >99%), they do not encode two of these transposases (**Fig. 2B**).

Interestingly, these 2 transposases are inserted within *rfbT* in PCG, resulting in a

truncated, non-functional version of the gene (**Fig. 2B and 2C**). *rfbT* encodes a

methyltransferase that is responsible for seroconversion between *V. cholerae* O1

Ogawa and Inaba serotypes, suggesting a major role for these IS sequences in the

emergence of PCG[54]. Furthermore, several strains from C23 encode the same version

of the hypervariable region as VC22 and 2479-86 (identity 99%) but the *rfa* cluster is

different (**Fig. 2B**). The first three genes of the island encode different alleles of *rfaF*,

*rfaG* and *rfaL*, and the identity of the remaining three is 95% (**Fig. 2B**). The *rfa* cluster

pattern of these strains is similar to that of strain A325 (C30) (**Fig. 2B**). In addition, in

this strain, the two transposases that truncate the *rfb*T gene in the PCG are inverted

maintaining an intact version of the gene (**Fig. 2C**). Overall, we found a remarkable

degree of conservation of the LPS cluster, which is present in almost twenty clades

encoding a PCG-like version. This is particularly striking as LPS hypervariability is a

universal bacterial characteristic, suggesting some strong evolutionary pressure

maintaining this version of the island widespread in the population. It appears that the

island is a) exchanged by homologous recombination leading to the whole gene cluster

being conserved and b) the presence of IS elements increases the variability by

truncating genes such as *rfbT*.

**Abundance and distribution of virulence-associated mobile genetic elements in**

***V. cholerae* species.** There are several horizontally acquired MGEs that are directly

associated with the emergence of toxigenic clones of *V. cholerae*. After we established

the phylogeny and clustering of the *V. cholerae* species, we investigated the

distribution, evolutionary dynamics, and allelic diversity of these elements within the

population.

*1. CTXΦ phage.* The genes for cholera toxin (CT) are encoded within the CTXΦ

phage[30]. The production of CT in the intestine is responsible for the severity of the

profuse diarrhea associated with cholera. CT leads to the secretion of electrolytes and

nutrients such as fatty acids into the lumen of the small intestine[9,55,56]. Due to the

lysogenic nature of CTXΦ, the insertion and deletion dynamics of this phage enables

recombination of genes, which leads to diversity within the toxigenic strains ultimately

serving as an opportunity to increase their pathogenic potential[30,57,58]. Inquiry into the

distribution and composition of this element within *V. cholerae* shows that there is no

environmental strain that recapitulates all the phage genes of PCG. At most, five

clades and two clusters encode 10 of the 14 genes that make up the CTXΦ phage of

*V. cholerae* N16961: three from C24 (represented by strains AAS91, PivertUAT4Aug

and V060002) and two from C17 (3541-04 and 3528-08) (**Fig. 3A**). Four of these

strains encode the two subunits of the cholera toxin (CtxAB). However, genes coding

for the duplicated transcriptional regulator RstR (VC_1455 and VC_1464) are only

found in one strain: AAS91 (**Fig. 3A**, brown shade). This strain belongs to the PCL and

encodes the complete version of CTXΦ except for VC_1460 that encodes a minor coat

protein pIII related to host interaction as part of the infection process (**Fig. 3B**)[59].

Instead, a transposase has been inserted in this locus in strain AAS91, followed by a

APH(3')-II family aminoglycoside O-phosphotransferase (**Fig. 3B**). In addition, AAS91

has a third copy of the triplet (RstA-RstB-RstR) instead of the two copies present in

PCG strains (**Fig. 3B**). The second most complete version of CTXΦ in *V. cholerae* is

found in strains V060002 from PCL and 3528-08 (C17). Both strains have lost the gene

coding for RstC and encode a different allele of the transcriptional regulator RstR (**Fig.**

**3B**).

Genomic comparison of *V. cholerae* strains reveal that the phage variation

dynamics are of the "additive" kind, where small gene cassettes are integrated into the

3' part of the gene coding for RtxA[60]. All the CTXΦ variants have conserved intergenic

regions that separate and mark the insertion of these gene cassettes (*att* sites, marked

as black arrows). These types of insertions have been reported previously in *V.*

*cholerae* O139[61]. Interestingly, some strains such as 3528-08 (C17) can have up to five

insertions (**Fig. 3C**). The first two of the five cassettes are identical, and each cassette

has a copy of the two subunits of CT. The other three have lost the toxin genes and

those coding for RstB, RstR and RstA, and acquired a replication endonuclease (**Fig.**

**3C**). The presence/absence of these cassettes is not correlated with the phylogeny

and, when present, they have high identity indicating a fast turnover of these elements

(**Fig. 3B**). Furthermore, these cassettes can be integrated into other genome locations,

as in the case of the strain PivertUAT4Aug (**Fig. 3B**). In addition, using the sequence of

*att* sites as reference we also found a collection of cassettes in the same order

(including the one containing the two subunits of CT) in the close relative species *Vibrio*

*mimicus*, suggesting another potential donor of CTXΦ to non-toxigenic *V. cholerae* (**Fig**

**S3**).

*2. Vibrio Pathogenicity Island-1 (VPI-1).* The toxin coregulated pilus (TCP), an

essential colonization factor, is encoded within VPI-1[31,32]. TCP mediates microcolony

formation, which is crucial for intestinal colonization, and acts as the receptor of

CTXΦ[33]. VPI-1 also encodes two critical virulence regulators: ToxT and TcpP[32,33]. VPI-

1 can be transferred, via generalized transduction, between strains of *V. cholerae* and

can also form circular intermediates[62,63]. The variable region of the island is located

between the CDS VC_0821 (hypothetical protein) and VC_0846 (integrase) (**Fig. 4A**).

The most conserved version of this region can be found in nine clades, four of them

within C24 (2 from PCL) and five from C17 (**Fig. 4B**), including strain IRLE0081

isolated in this study (**Fig. 1C**). The average identity of the proteins was highly

conserved compared to PCG (*ca*. >97%), however, we found some notable exceptions

(**Fig. 4A**). First, among the proteins that are part of the TCP operon, we found high

divergence rates in the master regulator ToxT (VC_0838) (**Fig. 4C**) and the toxin-

coregulated pilus major pilin, TcpA (VC_0828) (**Fig. 4D**) with a median protein identity

*ca*. 81% (**Fig. 4**). In addition, we found that the PCG allele of the colonization factor

TcpF (VC_0837) (**Fig. 4E**) was only present in three strains (one PCL and two C17). In

the rest, despite the conservation of synteny and similarity of the whole TCP operon,

the identity of this protein was less than 40%, encoding a different allele of this poorly

understood secreted protein[64–66] (**Fig. 4B**). Finally, our genomic comparisons reveal

that the gene coding for ToxR-activated gene A, TagA (VC_0820) was absent from the

C17 genomes, while being retained in three of the four C24 genomes (**Fig. 4B**).

Overall, despite widespread conservation in the synteny and size of the clusters of genes that comprise VPI-1 in the *V. cholerae* species, the existence of unique major allelic variations in a limited number of genes might act as a bottleneck in the emergence of PCG.

*3. Vibrio Pathogenicity Island-2 (VPI-2).* VPI-2 is a large PAI that encodes the genes for sialic acid utilization[34]. The capacity to utilize sialic acid as a carbon and energy source confers *V. cholerae* a competitive advantage in the mucus-rich environment of the gut, where sialic acid availability is extensive[67]. Sialic acid catabolism also mediates a chemotactic response towards mucin and several environmental reservoirs of *V. cholerae*[68]. VPI-2 is excisable and form a circular intermediate[63,69]. Interestingly, during the excision event, there is crosstalk between VPI-2 and VPI-1[70]. Unlike other MGEs, the differential abundance of the gene clusters within VPI-2 suggests that the island is made up of different modules. Also, all but one of the IS elements of the island are widespread among several clades in diverse locations, which favours the transfer of the different modules. The canonical VPI-2 from O1 strains encodes three major modules, from 5' to 3': a) a restriction modification system (RM), b) the Nan-Nag cluster, and c) a Mu-phage like region (**Fig. 4F**). At the 3' end of VPI-2 there is also a small group of genes encoding hypothetical proteins. The Nan-Nag cluster is present in 88 clades, whereas the distribution of the RM system is more limited, being encoded by only 7 clades (**Fig. 4F**). The Mu-phage like module is present in nine clades, however, there is no relationship between specific modules and species phylogeny. Like VPI-1, no clade other than PCG encode a PCG-like version of VPI-2. Furthermore, strains that contain the three individual modules sometimes encode them in different locations within the genome. For instance, strain PivertUAT4Aug (C24), the only one with the three major modules, encode the first two modules (Nan-Nag and RM) within a version of VPI-2, whereas the Mu-phage region is encoded in a different location of the genome (**Fig. 4G**). Similar to *rfb*T in the LPS gene

cluster, VC_1788 and VC_1791 in this strain comprise two parts of a gene coding for

the tape measure protein of phage Mu that have been truncated by the insertion of two

transposases in the PCG, annotated as VC_1789 and VC_1790 (**Fig. 4F**). Although the

synteny in the modules was conserved, and with high similarity (**Fig. 4G**), we found a

high divergence in the VC_1768 gene coding for a restriction endonuclease S subunit

(HsdS) between PivertUAT4Aug and the PCG-like version of VPI-2 with a protein

identity of 64% (**Fig. 4G**). Another instance of this modular distribution is exemplified by

strain N2824 (C16), which possesses the first two modules but not the Mu-phage.

Furthermore, the strain has lost the genes encoding the HsdS and HsdM proteins from

the RM cluster involved in DNA sequence recognition and methylation, respectively

(**Fig. 4G**). Strain 3541-04 (C17) encodes a different cluster of genes at the insertion

site of the Nan-Nag cluster and RM system, with the Nan-Nag cluster being inserted in

another chromosomal region (**Fig. 4G**). Overall, VPI-2 in PCG must be understood

based on its unique modular arrangement not by the individual units that constitute it

and the possibility of allelic variability in *hsdS* playing a unique role in the pandemic

group. Its four modules are associated with IS elements that are widely distributed

among *V. cholerae* environmental strains and scattered throughout the genome. This

leads to the convergence of all modules in the same genome being extremely low

which contributes to the rarity of PCG.

*4. Vibrio Seventh Pandemic Island I (VSP-I).* The role of VSP-I in *V. cholerae*

pathogenesis is not as clearly defined as other MGEs. VSP-I has been suggested to

play an environmental role related to chemotaxis[71] and encodes a regulator involved in

intestinal colonization (VC_0177)[35]. We found homologs of the eleven genes that

comprise this island in ten *V. cholerae* clades (**Fig. 5A**). Although four of them are

found in strains isolated from this study, IRLE0062, IRLE0049, IRLE0042 and

IRLE0077, that belonged to the same clade. The greatest sequence divergence

corresponds to its integrase (VC_0183) and an XRE family transcriptional regulator

(VC_0176) encoded within the island (**Fig. 5A**). Interestingly, only two strains encode a

PCG-like VSP-I, V060002 (C24), and VcCHNf4 (C5). Surprisingly, V060002 encodes

two identical copies of VSP-I in close proximity to each other with the LPS cluster

located between them (**Fig. 5B**). The landscape of evolutionary possibilities such as

modularity is much more limited for VSP-I than for the other MGEs due to its small size.

It is also the rarest of all the islands with one transcriptional regulator, VC_0176,

exhibiting high allelic diversity and very limited distribution. The rarity of the island and

diversity of the regulator suggests that this is a critical bottleneck in the emergence of

the Seventh Pandemic El Tor strains.

*5. Vibrio Seventh Pandemic Island II (VSP-II).* The presence of VSP-II in clinical

and environmental strains might be associated with environmental survival and fitness

of the bacterium[36–38]. A group of *V. cholerae* strains that caused an outbreak in Florida

associated with oyster consumption encoded a novel bacteriocin and a pyocin within

the VSP-II element, proteins typically associated with bacterial competition[38]. Our

analyses indicate that, similar to VPI-2, VSP-II is highly modular with several clusters of

genes that have a tRNA-Met insertion point. The direct repetition of the end of the

tRNA (highlighted as purple triangles) marks the insertion points of each module, which

in most cases match the presence of IS elements (**Fig. 5C and 5D**). Overall, this island

serves as another example of an "additive" island with three small modules. Module 1

comprises genes VC_0490 to VC_0498, five of which are of viral origin but lack

essential proteins for packaging, indicating that it is a defective prophage. Module 2

encompasses VC_0502 to VC_0510, however, the lack of possible annotation of its

genes precludes from assigning a specific function to this module. Nonetheless,

VC_0502 and VC_0503 from this module are only present in strains from C24 (**Fig.**

**5C**). Module 3 is comprised of four genes (VC_0512 to VC_0515): VC_0512 and

VC_0514 encode methyl-accepting chemotaxis proteins, VC_0513 a transcriptional

regulator and VC_0515 a predicted signal transduction protein. These four genes are

only present in C24 strains and according to their annotation could be associated with environmental adaptations (**Fig. 5D**). Interestingly, there are no identical versions of VSP-II outside of PCG. The closest versions are encoded by strains N2723, MZO-3, and V060002 belonging to C24, which are the only ones that possess a similar PCG-like modularity and synteny of VSP-II (**Fig. 5D**). N2723 has lost genes coding for two hypothetical proteins (VC_0496 and VC_0509) and a transcriptional regulator (VC_0497) and VSP-II from strain MZO-3 lost a transposase (VC_0501) in Module 1 and gained another in Module 3 (**Fig. 5D**). In the rest of *V. cholerae* strains, VSP-II only maintains some of these modules or clusters of genes, with the addition of other types of clusters being common. Finally, our analyses did not reveal any gene that exhibits high rates of divergence within the island, suggesting allelic diversity might not act as a major constraint in this element.

**Allelic variations confer a competitive advantage during intestinal colonization**. We identified a set of genes encoded within the elements described above that are dispersed in the phylogeny of the species and possess unique variations in PCG. As stated above, we hypothesized that they might act as bottlenecks for the emergence of toxigenic clones of *V. cholerae* and shed light on the evolution of PCG. In order to test this, we constructed several isogenic mutants where we exchanged the wild-type PCG allele for an environmental allele as specified in **Table 1**. Specifically, we mutated the genes encoding:

a) **RfbT** (VC_0255). A FkbM family methyltransferase located in the LPS cluster associated with Inaba-to-Ogawa serotype conversion[54]. The PCG allele is truncated by transposases and was replaced by the allele of *V. cholerae* A325, which has the complete version of the gene.

b) **TcpA** (VC_0828). The TCP major pilin subunit is encoded within VPI-1 and has an average protein identity of 81%. We exchanged the PCG *tcpA* allele for the one encoded by the IRL isolate *V. cholerae* IRL0081.

**c) ToxT** (VC_0838). The master virulence regulator ToxT is also encoded within VPI-1 and has an average identity of 81%. We exchanged the PCG *toxT* allele for the one encoded by the IRL isolate *V. cholerae* IRL0081.

**d) TcpF** (VC_0837). The secreted factor TcpF is also encoded within VPI-1 with only three strains encoding the PCG-allele. The rest had an average identity less than 40%. We exchanged the PCG *tcpF* allele for the one encoded by the IRL isolate *V. cholerae* IRL0081.

**e) VC_1791.** A tape measure protein in the Mu phage-like module encoded within VPI-2. The PCG allele is truncated by the insertion of two transposases and was replaced by the allele of *V. cholerae* strain PivertUAT4Aug, which has the complete version of the gene.

**f) HsdS** (VC_1768). This restriction endonuclease S subunit encoded within VPI-2 has the lowest median identity of the entire island (64%). We exchanged the PCG *hsdS* allele for the one encoded by the *V. cholerae* strain PivertUAT4Aug.

**g) VC_0176.** An XRE family transcriptional regulator encoded within VSP-1 that has the lowest median identity of the entire island (85%). We exchanged the VC0176 allele from the IRL isolate *V. cholerae* IRL0062.

We performed competitive assays using the infant mouse model of infection to determine whether the PCG-associated allelic variations of these genes confer a fitness advantage to toxigenic *V. cholerae.* Our results indicate that the PCG allele of *tcpF* is essential during the infectious process, with the mutant strain *tcpF*$^{IRL0081}$ exhibiting loss of intestinal colonization, similar to the $\Delta toxR$ strain used as negative control, which is avirulent (**Fig. 6A**). On the other hand, the allelic specificity of HsdS does not appear to play a direct role in the emergence of PCG as the competitive index (CI) of the *hsdS*$^{PIVERT}$ mutant is close to 1 (**Fig. 6A**). The mutant strains for the other 5 genes exhibit a varying degree of fitness loss during intestinal colonization with *tcpA*$^{IRL0081}$ showing the largest decrease (~3-logs), followed by *toxT*$^{IRL0081}$ (~1-log), *VC1791*$^{PIVERT}$ (~1-log), *rfbT*$^{A325}$ (~7-fold), and *VC0176*$^{IRL0062}$ (~5-fold) (**Fig. 6A**).

We recently determined that allelic variations in core genes, in the form of

virulence adaptive polymorphisms (VAPs), confer critical preadaptations for the

emergence of pathogenic traits[22]. For instance, toxigenic strains encoding

environmental alleles of the major outer membrane porin OmpU from the strain exhibit

bile sensitivity and reduced resistance to host antimicrobials[22,72]. Furthermore, those

strains (e.g., $ompU^{GBE1114}$) show a colonization defect in the infant mouse model (~1-

log) (**Fig. 6A**)[22]. In order to unveil the potential relationship among these allelic

variations (MGEs and core genome), we constructed a set of mutant strains combining

these alleles (**Fig. 6B**). We excluded those that showed no defect in colonization

($hsdS^{PIVERT}$) or led to complete loss of colonization ($tcpF^{IRL0081}$). We constructed the

mutants on the background of $VC0176^{IRL0062}$, as it was the one that showed the lowest

effect in the CI (**Fig 6A**) and exchanged the alleles in ascending order based on their

effect on the CI. Specifically, first we constructed a double mutant by introducing

$rfbT^{A325}$ that exhibits a ~5-fold decrease in its competitive index. Next, in the double

mutant background, we constructed a triple mutant by introducing $VC1791^{PIVERT}$ (~9-

fold). Subsequently, we constructed a quadruple mutant by introducing $ompU^{GBE1114}$

(~2.2-log) and a quintuple by introducing $toxT^{IRL0081}$ in the previous background (~2.9-

log). Finally, we generated a sextuple mutant encoding the previous 5 environmental

alleles and $tcpA^{IRL0081}$ which is fully defective for intestinal colonization. Overall, there is

a variable and non-linear effect between the alleles, with the largest effect being that of

the introduction of $ompU^{GBE1114}$ and $tcpA^{IRL0081}$ (**Fig. 6B**). Phenotypic analyses indicate

that the complete loss of colonization exhibited by $tcpF^{IRL0081}$ and the sextuple mutant is

not due to a growth defect, loss of motility, or changes in their ability to form biofilm

(**Fig S4**). Our results demonstrate that allelic variations in MGEs, LPS cluster and core

genome act as an additive yet stringent bottleneck that explicates the confined nature

of toxigenic *V. cholerae* isolates and the PCG.

**PCG alleles do not influence colonization of model environmental reservoirs.** Colonization of new niches or hosts is often associated with fitness trade-offs that make those adaptations beneficial within one context while detrimental on another[73,74]. In order to determine potential environmental fitness trade-offs associated with the PCG alleles, we examined the colonization dynamics of the two isogenic mutant strains that exhibit complete loss of intestinal colonization, *tcpF*[IRL0081] and the sextuple, during colonization of three model environmental reservoirs: crustaceans, cyanobacteria and molluscs.

**a) Crustaceans**. We developed a model of infection using the brine shrimp *Artemia salina* to examine the interactions of *V. cholerae* with crustaceans (**Figs 7A, 7B and S5**). First, we examined the colonization dynamics of *V. cholerae* C6706 at different concentrations and time points (**Fig S5**). We determined that infection with $10^7$ CFU/mL for 48 hours leads to consistent colonization of *A. salina* nauplii (**Fig S5**). Surprisingly, even though we expected *V. cholerae* to colonize the chitinaceous surface of *A. salina*, the bacterium primarily colonizes the gut space of the nauplii (**Fig 7B**). Next, we performed co-infection studies using a similar approach as those in the infant mouse. Our competition assays demonstrate that mutant strains encoding *tcpF*[IRL0081] or the sextuple mutant colonize *A. salina* at similar levels as the WT strain with a CI ~1 (**Fig 7C**).

**b) Cyanobacteria**. We used *Microcystis aeruginosa* as a model system to investigate the interactions of *V. cholerae* with cyanobacteria as the bacterium has been found associated with this species between cholera outbreaks in endemic areas[68,75]. Competition assays demonstrate that, similar to *A. salina*, both mutants colonize the mucilaginous sheath of *M. aeruginosa* comparably to the WT strain.

**c) Mollusks**. In order to examine the interaction of *V. cholerae* with molluscs, we extracted the mucus-containing gills of Eastern oysters (*Crassostrea virginica*) and performed competition assays using their homogenates (see materials and methods). Similarly, the sextuple mutant and *tcpF*[IRL0081] exhibit a CI in oyster gill homogenates

analogous to the WT toxigenic *V. cholerae* isolates (**Fig 7C**). Overall, our results

indicate that no significant fitness advantage or trade-off is associated with the

acquisition of PCG alleles in these model systems.

**CONCLUSIONS**

In this study, we took a systematic approach towards unraveling the evolutionary

history of the enigmatic Pandemic cholera group (PCG), which, to date, remains the

only clade within the *V. cholerae* species capable of causing the severe diarrheal

disease cholera. Phylogenomic analyses of the species reveal a cluster-based

evolution of *V. cholerae* with several hierarchical levels leading to the emergence of the

PCG. Our investigations on the abundance and distribution of the virulence-associated

genes and MGEs suggest the existence of several layers in the evolutionary dynamics

and emergence of *V. cholerae* (**Fig. 8**).

Our analyses indicate that **a)** strains from *V. cholerae* PCG have acquired the

gene clusters encoding the major virulence genes in a modular fashion, **b)** those MGEs

are more widespread than previously thought within the *V. cholerae* species, **c)** a

select set of genes dispersed in the phylogeny of the species have a very limited

distribution, and **d)** unique allelic variations in PCG act as stringent bottlenecks for

emergence. Pangenome analyses of the *V. cholerae* species reveal that, besides the

acquisition and modularity of the main MGEs, the absence or presence of other genes

does not appear to play a major role in the emergence and evolution of the PCG

(**Supp. text and Fig. S6A**). Furthermore, the phylogeny of *V. cholerae* shows that we

can only trace the evolutionary history of the LPS cluster (**Supp. text and Fig. S6B**).

This island-like element is exchanged by homologous recombination and appears to

have slower replacement dynamics than the MGEs. On the other hand, the MGEs are

additive genomic islands composed of several modules bordered by IS elements.

These modules have a rapid turnover and remain associated with each clade for a

short time. In addition, these modules have the potential to be inserted into other

regions of the genome. This mosaic-like degree of molecular sophistication explains

the absence of a complete set of PCG-like elements in any other *V. cholerae* clade.

Indicating that, in order to analyze the evolutionary history of these mobile elements, it

is necessary to examine the specific modules and individual alleles of some genes

within them rather than investigating presence or absence of the elements (**Fig. 8**).

To examine whether these intra-modular allelic variations act as an evolutionary

constraint that govern the emergence of toxigenic *V. cholerae*, we constructed isogenic

mutant strains where we exchanged the PCG-associated allele for an environmental

one and investigated their fitness during intestinal colonization. We determined that the

PCG-associated alleles lead to a competitive advantage during intestinal colonization.

Surprisingly, we uncovered a differential effect between alleles of genes encoded

within MGEs, the LPS, and core genome. Specifically, we found that some PCG-

associated alleles are essential for successful colonization, such as *tcpF*, whereas

others provide non-linear competitive advantages during infection that ultimately act as

a critical bottleneck that clarifies the isolated emergence of PCG. Future research will

elucidate the specific role and relative importance of these alleles and their potential

interactions. Elucidating this will open novel research avenues and provide essential

insights into the factors and network of genes associated with the emergence of

virulence traits in bacteria. Finally, it appears that none of the three environmental

reservoirs we examined in this study exert the ecological pressures associated with the

selection of the PCG alleles as the mutants exhibit a similar competitive index as the

wild-type strain. Given that these variations originate from the environment it is possible

that a combination of abiotic and biotic factors might select for the PCG alleles in the

aquatic ecosystem. Uncovering the environmental drivers fostering the selection of

factors associated with pathogen emergence will provide critical insights towards the

development of surveillance networks. Overall, our investigations provide unique

insights and perspectives to elucidate the barriers associated with the emergence of

human pathogens and can be extrapolated to the study of niche colonization and other

complex phenomena.

**MATERIALS AND METHODS**

**Sampling sites**. We collected samples at two environmentally distinctive locations

along the IRL (Eastern Florida, USA) in three sampling events. For each sampling

campaign, we obtained samples from several fractions: three associated with the water

bodies (water filtered through 20µm, 5µm, and 0.22µm) and one from the sediment.

One of the sampling locations, Fellers House Field Station (N28˚54'25.315";

W80˚49'15.017"), is located within the federally-protected Canaveral National

Seashore. The second sampling site, Shepard Park, is located in the Port St. Lucie

area (N27˚11'48.864"; W80˚15'33.172") that due to urbanization and agricultural

expansion, experiences nutrient over-enrichment leading to excessive macroalgal

bloom (**Fig. 1A**)[45].

**Isolation of *V. cholerae*. a)**. <u>*Water*</u>. Isolation of *V. cholerae* from water samples

collected at the sampling locations was performed using a modified protocol from Huq

*et al* [76]. Briefly, water samples (~3L) were collected in triplicate from each site and

stored cold until arrival at the laboratory. To investigate the prevalence of *V. cholerae* in

various fractions of the aquatic column and to separate particle-associated and free-

living bacteria, 500 mL of collected water was filtered successively through 20µm, 5µm

and 0.2µm filters (Sterlitech) using a vacuum filtration system. The respective

membrane filters harboring potential *vibrio* spp. were then suspended in 25ml

phosphate buffered saline (PBS) and vortexed vigorously to release the contents. Each

resuspended sample was then inoculated (1:10 v/v) in alkaline peptone water (APW)

and incubated at 37°C for 12-14 hrs. This was repeated for each replicate. Enriched

cultures were subjected to growth on a series of selective agar specific for the isolation

of *V. cholerae.* First, enriched cultures were serially diluted 10-fold and plated on

CHROMagar Vibrio (CaV; CHROMagar), a chromogenic media on which *V. cholerae*

forms turquoise blue colonies. Subsequently, turquoise blue colonies from CaV were

transferred onto Thiosulfate Citrate Bile Salts Sucrose (TCBS; Sigma) agar plates.

Colonies that appeared turquoise blue on CaV followed by yellow on TCBS were

preliminarily identified to be *V. cholerae*. **b)** _Sediment_. Isolation of *V. cholerae* was

done using established methods[45]. Briefly, sediment samples were collected in

triplicates from each sampling site using universal corer. 25 g of the sediment samples

from each replicate was weighed and suspended in 25 ml of PBS (Gibco). Samples

were homogenized, enriched, and studied for the presence of *V. cholerae* as described

above. **c)** _Cyanobacteria_. Cyanobacteria floccules were pelleted, supernatant removed

and fresh APW added to homogenize the sample. Samples were enriched and studied

for the presence of *V. cholerae* as described above.

**Strains and growth conditions.** *V. cholerae* strains were routinely grown on Luria

Bertani (LB) agar at 37 ºC for approximately 16hrs, unless otherwise stated. For routine

liquid cultures, isolated colonies selected from agar plates were grown aerobically at

37ºC in LB broth for ~16hrs. Tryptone broth (tryptone 10 g/L, NaCl 5 g/L), artificial sea

water and M9 minimal media (Fisher) supplemented with 0.1% glycerol were used for

biofilm formation, studies using *Artemia* and oyster gill homogenates, respectively.

Growth media was supplemented with the following antibiotics and reagents, as

needed: polymyxin B 50 U/mL, kanamycin 45 µg/mL, streptomycin 1000 µg/mL,

gentamycin 15 µg/mL, X-gal 40 µg/mL.

**Clonal analysis of *V. cholerae* isolates.** To confirm isolates of *V. cholerae,* presence

of the species-specific gene *ompW* was determined by PCR[46]. Genomic DNA from the

isolates was extracted using the Gentra Puregene Yeast/Bact Kit (Qiagen) and used as

template for PCRs. To identify the clonal populations of *V.* cholerae in the confirmed

dataset, DNA fingerprinting patterns were examined using a combination of *Vibrio*

*cholerae* Repeats-PCR (VCR-PCR) and Enterobacterial Repetitive Intergenic

Consensus Sequence (ERIC)-PCR as described by Teh *et al* [47]. Genomic DNA was

used to perform VCR- and ERIC-PCR and resulting PCR products were

electrophoresed and imaged using UVP ChemStudio (AnalytikJena) to observe

fingerprinting patterns. Cluster analysis of DNA patterns was performed using

BioNumerics v8.0 (Applied Maths, Inc.) and dendrogram generated based on the

Unweighted Pair Group Method with the Arithmetic Mean (UPGMA) method[47,77,78] (**Fig.**

**S1).** Based on resulting DNA fingerprinting patterns, a total of 35 independent clones

were identified and selected for whole genome sequencing (WGS) (**Table S1**).

Genomic DNA of each clone was submitted to Microbial Genome Sequencing Center

(MiGS) for WGS using the Illumina NextSeq 2000 platform.

**Genome assembly, gene prediction, and annotation.** Reads were trimmed using

Trimmomatic v0.36[79] and assembled de novo with SPAdes version 3.11.1[80]. Coding

DNA sequences (CDS) from the assembled contigs were predicted using Prodigal

version 2.6.3 using *-a output.proteins -d output.genes -c -p meta* parameters[81]. Then,

tRNAs were obtained using tRNAscan-SE version 1.4[82] together with ssu-align version

0.1.1, and rRNA genes using meta-rna[83]. Predicted proteins were compared against

the National Center for Biotechnology Information nonredundant database (NCBI nr)

using DIAMOND[84]. In addition, for functional annotation we used HMMscan version

3.1b2[85] for the comparison against COG v2003 (update 2014)[86] and TIGRFAM v15.0

(September 2014)[87] databases.

**Recovery of *V. cholerae* genomes.** A total of 1,127 *V. cholerae* genomes were

downloaded from RefSeq (accessed July 2021) and subjected to a dereplication step

with drep[88] at 99 % nucleotide identity. Briefly, we force dRep to sort and select as

seeds for the clustering genomes with better score. For instance, chromosomes in a

single contig, followed by higher N50 and L50 values, and lastly by the completeness

and contamination values calculated with checkm[89]. This resulted in 227 dereplicated

reference *V. cholerae* genomes.

**Phylogeny of *V. cholerae* dereplicated genomes.** Single nucleotide polymorphisms

(SNPs) among *V. cholerae* dereplicated genomes and the PCG reference genome

*Vibrio cholerae* El Tor N16961 (RefSeq accession number GCF_900205735.1) were

calculated with PhaME[90] with default options, resulting in a core genome length of

2,468,623 bp and 458,798 polymorphic positions considered. Then, a maximum

likelihood phylogenetic tree of SNPs was constructed using iqtree v 1.6.12[91] using the

ultrafast bootstrap approach (5000 replicates) and the best fitted model GTR+F+R4.

**Abundance and distribution of pathogenicity-related islands.** To determine

virulence-associated mobile genetic elements and LPS cluster distribution and degree

of prevalence of proteins within the studied *V. cholerae* genomes, we extracted these

genomic regions from PCG reference genome (*V. cholerae* El Tor N16961) based on

the manual annotation found in the NCBI Genbank.  Then, proteins were extracted and

used as a reference in a BLASTP search against the proteins from the rest of the *V.*

*cholerae* species genomes[92,93]. We kept all proteins that matched against the

pathogenic protein database with at least 70% identity and the alignment covered at

least 70% of both proteins.

**Construction of mutant strains.** *V. cholerae* constructs were all generated in the El

Tor strain C6706 background following previously published methods[22]. Briefly, the

environmental alleles of interest were synthesized commercially (Gene Universal)

including 500 bp flanking regions from *V. cholerae* C6706 to facilitate recombination.

The synthesized products were digested and ligated into the *V. cholerae* suicide vector

pKAS154, electroporated into *Escherichia coli* S17λ-*pir* and selected on kanamycin

plates. Following sequence verification, allelic exchange was carried out in *V. cholerae*

C6706 using the appropriate antibiotic selection, as described[94]. Positive clones were

first screened by colony PCR using flanking primers and confirmed by sequencing the

exchange locus from purified genomic DNA. Fluorescent tagging of the strains was

performed using the mini-Tn7 system that integrates at the *att* site between the *glmS*

(*VC0487*) and *VC0488* genes in *V. cholerae*[95]. Briefly, triparental mating was

performed using LB-grown overnight cultures of the donor *E. coli* S17$\lambda$-*pir* cells

harboring pUC18T-mini-Tn7T-GFP or RFP expressed from a strong constitutive

promoter, with the recipient *V. cholerae* cells and helper *E. coli* encoding the

transposase gene on a suicide vector. The mating mixture were selected twice on LB-

Pb-Gm plates and followed by two rounds of selection on LB-Sm plates, and

subsequently confirmed by sequencing the Tn7 insertion site and fluorescence

microscopy.

**Infant mouse colonization assays.** Intestinal colonization competition assays in

three-to-five-day-old infant mice were performed with LB-grown *lacZ*[+] cells of the WT

and allelic exchange mutants competed against an isogenic WT strain harboring a *lacZ*

deletion, essentially as described[22,96,97]. The colonization efficiencies of the different

strains in the small intestine harvested post infection are represented as competitive

indices. Briefly, strains were inoculated from fresh LB plates into LB-Sm broth,

incubated for 12 hours at 37 °C with aeration, and diluted 1:1000 fold in 1x PBS (pH

7.4, Gibco). The test (*lacZ*[+]) and control (*lacZ*) strains were mixed 1:1 to obtain ~$10^6$

CFU/mL. To monitor successful inoculation, Evan's blue dye (0.01% final, Sigma) was

added to the cell mixture. Fifty microliters of the bacterial mixture, corresponding to

~$10^5$ CFUs, were then used to intragastrically inoculate 3-5-day-old infant mice and

maintained at 30 °C post-infection. The exact input CFU numbers were determined by

plating dilutions of the inoculum on LB-Sm-X-Gal plates. Approximately 22 hours post

infection, the mice were euthanized, their small intestines harvested and transferred

into 4 mL LB-10% glycerol stored on ice. The samples were then processed for CFU

determination by homogenizing the tissues, serially diluting in 1x PBS (pH 7.4) and

plating appropriate dilutions on LB-Sm-X-Gal plates. *In vitro* competition experiments

were performed in parallel by inoculating 50 μL of the input CFU in LB-Sm broth and

incubating for 22 hours at 37 °C with aeration and enumerating CFUs as above.

Results are presented as the competitive index (CI), which is the ratio of the intestinal

*lacZ*$^+$ CFU (blue colonies) to *lacZ* CFU (white colonies) normalized to the input CFU

ratio and *in vitro* competition CFU ratio.

**Colonization of environmental hosts**. Bacterial inocula for infections of the

environmental hosts were prepared essentially as described for the mice colonization

assays but used at different dilutions and without the Evan's blue dye. Coinfections

were performed with GFP-tagged *lacZ*-positive test strains competing against RFP-

tagged wild type strain harboring a *lacZ* deletion. **a)** *Crustaceans*. We modified

previously established protocols as follows[98], eggs of *A. salina* were hatched in artificial

sea water supplemented with 2% sodium chloride (ASW), incubated at room

temperature and aeration under a 12-hour light-dark cycle for 48 hours. Infections were

performed in 200 μL ASW containing 20-30 individual crustaceans and the 10$^7$

CFU/mL of bacteria. After 48 hours incubation at room temperature, samples were

washed twice in ASW to remove unattached bacteria and either homogenized for

determining CIs (as above) or used directly for microscopy. **b)** *Cyanobacteria*. We

modified previously established protocols as follows[99], *Microcystis aeruginosa* (UTEX

LB 2385) were incubated in flasks containing BG-11 media (Gibco) under a 12-hour

light-dark cycle with aeration (100 rpm) for about 14 days. *M. aeruginosa* cells were

concentrated by centrifugation at 7500 rpm for 10 minutes, resuspended in BG-11 and

co-cultured with 10$^7$ CFU/mL *V. cholerae* cells at a final OD700 of 0.3-0.4 in a 200 μL

reaction volume. Samples were harvested at 24 hours and CIs were determined on LB-

Sm-X-Gal plates, as above. **c)** *Mollusks*. Mucus-containing gills of the Eastern oyster

*Crassostrea virginica* were harvested in 1x PBS pH 7.4 (1 g/mL), homogenized and

autoclaved[68]. M9 minimal media was supplemented with 0.1% sterile gill homogenate

and inoculated with 50 µL of $10^7$ CFU/mL of bacterial inoculum. Subsequently, cultures

were incubated for 24 hours at 37 °C under aerated conditions. CIs were determined by

plating appropriate dilutions on LB-Sm-X-Gal plates.

**Phenotypic assays**. **a)** Growth on LB. Overnight cultures were diluted to ~$10^5$ CFU/ml

in fresh LB and 200 µL aliquots in flat-bottom 96-well plates (CytoOne) were monitored

at OD595 nm, 37 °C for 24 hours in a Tecan Sunrise plate reader. **b)** Biofilm formation.

These assays were performed as previously described with minor modifications[100].

Briefly, overnight cultures in tryptone broth were diluted 1:500 in fresh broth and 200 µL

aliquots in 96-well polystyrene plates (CytoOne) were incubated at 37 °C for 24 hours.

After washing to remove unattached cells, biofilm cells were stained with 0.1% crystal

violet and quantified by measuring OD595 after elution with 50% glacial acetic acid. **c)**

Motility. These assays were performed in soft agar plates (LB-0.3% agar) as previously

described[68]. Diameter was measured for 12 hours every two hours.

**Pangenome analysis.** To evaluate the pangenome of *V. cholerae*, in a first approach

dereplicated sequences were grouped in clusters at 98% identity using cd-hit[101],

resulting in 34 clusters, four of them (C5, C11, C16, and C24) comprising 188 out of

262 genomes. Then, pangenomes were calculated for each cluster individually using

PPanGGOLiN[102] with default parameters.  A second pangenome step with the protein

families of clusters C5, C11 y C16 was performed using PPanGGOLiN to simplify the

comparison among the PCG reference genome (*V. cholerae* El Tor N16961), C24 and

the new superpangenome C5-C11-C16. Ortholog proteins among the three groups (El

Tor, C24 and C5-C11-C16 were calculated using orthoMCL[103] (> 30 % amino acid

identity, > 70 % alignment). A Venn diagram showing proteins shared and not shared

among groups was performed in R using the ggVennDiagram package

(https://github.com/gaospecial/ggVennDiagram).

**Determination of Gain and Loss of genes in the pandemic cholera lineage (PCL).**
In order to understand the flow of genes among pandemic *V. cholerae* and the closest genomes from cluster C24, those belonging to the PCL, gene presence and absence was tested using the gain loss mapping engine GLOOME[104]. The expectations and probabilities of both gain and loss events were estimated using stochastic mapping. The phyletic pattern of presence/absence of orthologous genes was then classified using hmmscan[85] against the Pfam database v35.0[105], using an evalue of $< 1e^{-5}$ as a threshold for protein annotation.

**Microdiversity analysis.** To identify which genes in each genome were under positive selection, we calculated the nonsynonymous (d$N$) and synonymous (d$S$) substitutions between ortholog proteins found in both the studied *V. cholerae* genomes and the reference PCG strain, we used the orthologr package in R[49], a pipeline that identifies orthologous protein pairs, performs codon alignment and finally computes the final dN/dS values. We considered a gene under positive selection if the dN/dS ratio > 1.

**Data availability**
The genomes have been deposited as BioSample SAMN38476510 to SAMN38476544 under BioProject PRJNA1035537.

**ACKNOWLEDGEMENTS**

We thank Zac Carver for his technical assistance. This work was supported by a National Science Foundation (NSF) CAREER award (#2045671) and a Burroughs Wellcome Fund Investigator in the Pathogenesis of Infectious Disease award (#1021977) to SAM. MLP was supported by a "FLEX3GEN" PID2020-118052GB-I00 (cofounded with FEDER funds) from the Spanish Ministerio de Economía, Industria y Competitividad.

**COMPETING INTERESTS**

The authors declare no competing interests.

**AUTHOR CONTRIBUTIONS**

SAM conceived the study. DB and TAG collected samples, isolated *V. cholerae* and sequenced genomes. MLP, AZS and JHM performed bioinformatic analyses. DB and CC constructed mutants and performed host colonization assays. The manuscript was written by MLP, DB and SAM. All authors read and approved the final version.

**FIGURE LEGENDS**

**Fig. 1. Phylogenomic and population structure of *Vibrio cholerae*. (A)** Map of Florida indicating the locations where the environmental *V. cholerae* strains were isolated (Shepard Park and Feller´s House). **(B)** Whole-genome SNP-based phylogenetic tree showing the relationships among the *V. cholerae* genomes obtained in this study and a representative strain from the Pandemic Cholera Group (highlighted with a red arrow). **(C)** Phylogenomic tree of dereplicated *V. cholerae* strains available in public databases and strains from this study. The tree was reconstructed from single nucleotide polymorphisms of the core genome. Branches of members belonging to the four main clusters are colored in green (C5), violet (C11), blue (C16) and red (C24). The colored circles represent the source of isolation of the environmental strains collected in this study. Grey circles located at the branch nodes represent the bootstrap values.

**Fig. 2. Distribution and evolutionary dynamics of the O1 Lipopolysaccharide (LPS) cluster in the *V. cholerae* species. (A)** The area outlined in gray represents the number of homologous proteins found in the dereplicated *V. cholerae* genomes. The white circles with dotted lines indicate the median protein identity. **(B)** Genomic

comparison of PCG-like LPS clusters identified in *V. cholerae* environmental genomes.

**(C)** Schematic representation of the variable zone of the LPS containing the *rfb*T gene

and the IS elements inserted in the PCG allele. IS elements are highlighted in yellow.

**Fig. 3. Integron-like dynamics govern genetic variation of CTXΦ phage. (A)**

Distribution and structure of the CTXΦ phage in the *V. cholerae* species. The area

outlined in gray represents the number of homologous proteins found in the

dereplicated *V. cholerae* genomes. The white circles with dotted lines indicate the

median protein identity. Red rectangles on the genes indicate their exclusive presence

in the PCL. The black arrows indicate the genomic position where the cassettes have

been inserted (*att* site). **(B)** Genomic comparison of similar CTXΦ phage found in

dereplicated *V. cholerae* genomes. **(C)** Schematic representation CTXΦ phage found

in *V. cholerae* 3528-08, both the numbers and the rectangles mark the different

clusters acquired. Clusters of genes with a known function follow the same color

pattern. IS elements are highlighted in yellow.

**Fig. 4. Modularity and allelic variations shape the evolutionary dynamics of**

**Vibrio pathogenicity islands 1 and 2.** Structures of **(A)** VPI-1 and **(F)** VPI-2. The area

outlined in gray represents the number of homologous proteins found in the

dereplicated *V. cholerae* genomes. The white circles with dotted lines indicate the

median protein identity. **(B)** Genomic comparison of homologous VPI-1 in dereplicated

*V. cholerae* genomes denote a reduced number of allelic variations in some PCG

genes. Sequence alignment of the protein sequences of **(C)** ToxT **(D)** TcpA and **(E)**

TcpF from PCG and environmental strain IRLE0081 (C17). **(G)** Genomic comparison of

VPI-2 variants encoded in dereplicated *V. cholerae* genomes highlight pervasive

modularity of the island. Clusters of genes with a known function follow the same color

pattern. IS elements are highlighted in yellow.

**Fig. 5. Distribution and evolutionary dynamics of the Vibrio seventh pandemic islands I and II.** Structures of **(A)** VSP-I and **(C)** VSP-II. The gray area represents the number of homologous proteins found in the dereplicated *V. cholerae* genomes. The white circles with dotted lines indicate the median protein identity. Red rectangles on the genes indicate their exclusive presence in the PCL. Pink triangles indicate the presence of tRNA-direct repeat genomic islands. **(B)** Genomic comparison of homologous VSP-I across *V. cholerae* genomes. **(D)** Genomic comparison of VSP-II variants encoded in dereplicated *V. cholerae* genomes highlight pervasive modularity of the island. IS elements are highlighted in yellow.

**Fig. 6**. **Allelic variations act as fitness bottleneck during intestinal colonization.** Isogenic *V. cholerae* mutant strains were orogastrically inoculated in 3-5-day-old CD-1 mice to perform competition assays. Strains were monitored for colonization of the small intestine. The relative colonization data is represented as competitive indices (CI) in comparison to a co-infected *V. cholerae* C6706 $\Delta lacZ$ strain. **(A)** Isogenic strains encoding individual non-PCG alleles exhibit different competitive indices during intestinal colonization. **(B)** Combination mutants of the non-PCG alleles demonstrate an additive decrease in colonization efficiency. n = 6 per strain. *p*-values indicate significance compared to the WT strain, unless otherwise specified, and are calculated by 2way ANOVA in Šidák's multiple comparisons test; *** <0.0001, **** < 0.00001.

**Fig 7. PCG alleles do not influence colonization of model environmental reservoirs. (A)** Image of *Artemia salina* uninfected with *V. cholerae*. **(B)** RFP-tagged *V. cholerae* C6706 cells are ingested by *A. salina* colonizing and proliferating to fill the gut space. **(C)** Competitive indices of *tcpF*[IRL0081] and the sextuple mutant in three model environmental reservoirs: Crustaceans (*A. salina*), cyanobacteria (*Microcystis aeruginosa*) and mollusks (gill homogenates of the Eastern oyster, *Crassostrea*

*virginica*). The mutants do not exhibit any significant differences in CI during

colonization of these reservoirs. n = 6. *p*-values indicate significance compared to the

WT strain and are calculated by 2way ANOVA.

**Fig. 8. Model of emergence of the Pandemic Cholera Group in** ***V. cholerae.*** The

evolutionary history of PCG suggests that the emergence of this group has not been a

linear process and contains a series of major bottlenecks that explains its rarity. **(A)** A

specific genomic background containing preadaptations to virulence in the core

genome (red arrows) is essential for the emergence of virulence traits in *V. cholerae.*

Green arrows: strains not encoding these preadaptations. **(B)** The acquisition of certain

gene clusters is critical (e.g., CTXΦ, VPI-1, etc.) together with (**C**) the presence of

specific modules within these gene clusters and **(D**) certain unique allelic variations

within them. **(E)** The correct combination of all the elements of this complex mosaic

provides the foundation for the emergence of pathogenic clones within an

environmental population of *V. cholerae* (red arrow).

**Table 1. List of strains used in this study**

| Strain | Description | Source |
|---|---|---|
| *V. cholerae* C6706 | Wild-type, clinical *V. cholerae* O1 El Tor strain C6706; $Sm^R$ | Lab collection |
| *E. coli* S17 λ pir | Host for cloning vector | Lab collection |
| *ompU*[GBE1114] | *V. cholerae* C6706 encoding *ompU* allele from environmental strain *V. cholerae* GBE1114 | [22] |
| *rfbT*[A325] | *V. cholerae* C6706 encoding *rfbT* allele from environmental strain *V. cholerae* A325 | This study |
| *tcpA*[IRLE0081] | *V. cholerae* C6706 encoding *tcpA* allele from environmental strain *V. cholerae* IRLE0081 | This study |
| *toxT*[IRLE0081] | *V. cholerae* C6706 encoding *toxT* allele from environmental strain *V. cholerae* IRLE0081 | This study |

| | | |
|---|---|---|
| $tcpF^{IRLE0081}$ | *V. cholerae* C6706 encoding *tcpF* allele from environmental strain *V. cholerae* IRLE0081 | This study |
| $hsdS^{Pivert}$ | *V. cholerae* C6706 encoding *hsdS* allele from environmental strain *V. cholerae* PivertUAT4Aug | This study |
| VC1791$^{Pivert}$ | *V. cholerae* C6706 encoding tape measure (VC1791) allele from environmental strain *V. cholerae* PivertUAT4Aug | This study |
| VC0176 $^{IRLE0062}$ | *V. cholerae* C6706 encoding VC0176 allele from environmental strain *V. cholerae* IRLE0062 | This study |
| VC0176 $^{IRLE0062}$:: $rfbT^{A325}$ | Double mutant; $rfbT^{A325}$ environmental allele introduced into the *V. cholerae* C6706 encoding $VC0176^{IRL0062}$ | This study |
| Double mutant:: VC1791$^{Pivert}$ | Triple mutant; $VC1791^{PivertUAT4Aug}$ environmental allele introduced into the *V. cholerae* C6706 double mutant | This study |
| Triple mutant::$ompU^{GBE1114}$ | Quadruple mutant; $ompU^{GBE1114}$ environmental allele introduced into the *V. cholerae* C6706 triple mutant | This study |
| Quadruple mutant::$toxT^{IRLE0081}$ | Quintuple mutant; $toxT^{IRLE0081}$ environmental allele introduced into the *V. cholerae* C6706 quadruple mutant | This study |
| Quintuple mutant::$tcpA^{IRLE0081}$ | Sextuple mutant; $tcpA^{IRLE0081}$ environmental allele introduced into the *V. cholerae* C6706 quintuple mutant | This study |

**SUPPLEMENTARY MATERIAL**

**Table S1.** List of *Vibrio cholerae* collected in this study, together with genomic features

and metadata.

**Table S2.** List of dereplicated (ANI>99%) *Vibrio cholerae* reference strains from the

NCBI

**Table S3.** List of dereplicated (ANI>98%) *Vibrio cholerae* reference including IRL

strains

**Table S4**. Shared family gene clusters in the pangenomes of the three hierarchical

levels

**Table S5**. Summary of the gene gain/loss of the branching event 3 from the pandemic

cholera lineage.

**Table S6**. Genes of the PCG under positive selection (dN/dS >1) compared to more

than 60 non-clonal *V. cholerae* clades.

**Figure S1**. **Dendrogram representing the relatedness of *V. cholerae* isolates**. The

dendrogram is based on PCR fingerprints generated by VCR-PCR. Groups of similarity

were established using the UPGMA method in BioNumerics v8.0. The red circles

indicate clonal isolates that were selected for whole genome sequencing.

**Figure S2. Pairwise comparison among the *V. cholerae* genomes.** The pairwise

comparison was obtained using the average nucleotide identity (ANI) from strains

isolated in this study and the Pandemic Cholera Group (*V. cholerae* El Tor N16961).

**Figure S3**. **Genomic comparison of CTXΦ phage**. Comparison between the CTXΦ

phage genomic island encoded by Pandemic Cholera Group strains (*V. cholerae* El Tor

N16961) and *Vibrio mimicus* strain 2011V-1073.

**Fig S4. Phenotypes of the *tcpF* IRL0081 and sextuple mutants**. Isogenic *V. cholerae*

C6706 strains encdoing either the *tcpF* IRL0081 allele or the sextuple alleles were

monitored for differences in **(A)** growth patterns in LB broth, **(B)** motility in 0.3% agar

and **(C)** biofilm formation. n=6. *p*-values were determined using one-way ANOVA.

**Fig. S5. Artemia model of *V. cholerae* infection. (A)** CFU/ml of *V. cholerae* C6706

recovered post-infection at different time points and inoculum concentrations

normalized to the input. Infection threshold (orange line) was determined by visualizing

colonization using fluorescent microscopy. **(B)** RFP-tagged *V. cholerae* C6706 cannot

colonize *A. salina* with an inoculum of $10^6$ CFU/ml after 48 hours and fluorescence

cannot be detected. **(C)** Inocula of $10^7$ CFU/ml lead to consistent colonization of *A.*

*salina* by RFP-tagged *V. cholerae* C6706 after 48 hours and red fluorescence can be

detected.

**Fig. S6. Evolutionary trajectories and genome dynamics of *V. cholerae* PCG. (A)**

*Pangenome analysis*. The Venn diagram shows the numbers of unique and shared

gene families between the pandemic cholera group (purple), pandemic cholera lineage

(blue), Cluster 24 (light blue) and the rest of clades (grey). **(B)** *PCL evolution*. The tree

designates the evolutionary relationships of the different clades within the PCL

highlighting the loss and gain of genes through the lineage. Black numbers indicate an

overall gain and grey an overall loss. The colored circles represent the specific element

as well as the percentage of completeness in each of the clades in reference to PCG.

LPS, O1 lipopolysaccharide cluster; VPI-1, Vibrio pathogenicity island-1; VPI-2, Vibrio

pathogenicity island-2; VSP-1, Vibrio seventh pandemic island-1; VSP-2, Vibrio

seventh pandemic island-2. **(C)** *Microdiversity analysis.* PCG genes under positive

selection compared to the rest of the clades. X-axis indicates the number of clades in

which d$N$/d$S$>1 and the Y-axis shows the average of the d$N$/d$S$ value.

**REFERENCES**

1. Alexander, K. A. *et al.* What factors might have led to the emergence of Ebola in
West Africa? *PLOS Negl Trop Dis* **9**, e0003652 (2015).

2. Faye, O. *et al.* Molecular Evolution of Zika Virus during Its Emergence in the 20th
Century. *PLOS Negl Trop Dis* **8**, e2636 (2014).

3. Wellington, E. M. *et al.* The role of the natural environment in the emergence of
antibiotic resistance in Gram-negative bacteria. *The Lancet Infectious Diseases* **13**,
155–165 (2013).

4. Balasubramanian, D., López-Pérez, M., Grant, T.-A., Ogbunugafor, C. B. & Almagro-
Moreno, S. Molecular mechanisms and drivers of pathogen emergence. *Trends*
*Microbiol* **30(9)**, 898–911 (2022).

5. Jones, K. E. *et al.* Global trends in emerging infectious diseases. *Nature* **451**, 990–
993 (2008).

6. Koskella, B. & Vos, M. Adaptation in Natural Microbial Populations. *Annu. Rev.*
*Ecol., Evol., Syst.* **46**, 1–20 (2014).

7. Thompson, F. L., Iida, T. & Swings, J. Biodiversity of vibrios. *Microbiology and*
*molecular biology reviews : MMBR* **68**, 403–431 (2004).

8. Reen, F. J., Almagro-Moreno, S., Ussery, D. & Boyd, E. F. The genomic code:
inferring Vibrionaceae niche specialization. *Nature reviews. Microbiology* **4**, 697–704
(2006).

9. Kanungo, S., Azman, A. S., Ramamurthy, T., Deen, J. & Dutta, S. Cholera. *Lancet*
**399**, 1429–1440 (2022).

10. Matson, J. S., Withey, J. H. & DiRita, V. J. Regulatory networks controlling Vibrio
cholerae virulence gene expression. *Infection and immunity* **75**, 5542–5549 (2007).

11. Almagro-Moreno, S., Pruss, K. & Taylor, R. K. Intestinal colonization dynamics of
Vibrio cholerae. *PLoS pathogens* **11**, e1004787 (2015).

12. Sakib, S. N., Reddi, G. & Almagro-Moreno, S. Environmental role of pathogenic
traits in Vibrio cholerae. *Journal of bacteriology* **200**, 2123 (2018).

13. Ali, M., Nelson, A. R., Lopez, A. L. & Sack, D. A. Updated Global Burden of Cholera
in Endemic Countries. *PLOS Negl Trop Dis* **9**, e0003832 (2015).

14. Clemens, J. D., Nair, G. B., Ahmed, T., Qadri, F. & Holmgren, J. Cholera. *The*
*Lancet* **390**, 1539–1549 (2017).

15. Balasubramanian, D., Murcia, S., Ogbunugafor, C. B., Gavilan, R. & Almagro-
Moreno, S. Cholera dynamics: lessons from an epidemic. *Journal of medical*
*microbiology* **20**, 106 (2021).

16. Mutreja, A. *et al.* Evidence for several waves of global transmission in the seventh
cholera pandemic. *Nature* **477**, 462–465 (2011).

17. Islam, M. T., Alam, M. & Boucher, Y. Emergence, ecology and dispersal of the
pandemic generating Vibrio cholerae lineage. *International microbiology : the official*
*journal of the Spanish Society for Microbiology* **20**, 106–115 (2017).

18. Chatterjee, S. *et al.* Incidence, virulence factors, and clonality among clinical strains
of non-O1, non-O139 Vibrio cholerae isolates from hospitalized diarrheal patients in
Kolkata, India. *Journal of clinical microbiology* **47**, 1087–1095 (2009).

19. Chun, J. *et al.* Comparative genomics reveals mechanism for short-term and long-
term clonal transitions in pandemic Vibrio cholerae. *Proceedings of the National*
*Academy of Sciences* **106**, 15442–15447 (2009).

20. Dziejman, M. *et al.* Comparative genomic analysis of Vibrio cholerae: genes that
correlate with cholera endemic and pandemic disease. *Proceedings of the National*
*Academy of Sciences of the United States of America* **99**, 1556–1561 (2002).

21. Boucher, Y. Sustained local diversity of Vibrio cholerae O1 biotypes in a previously
cholera-Free country. *mBio* **7**, e00570-16 (2016).

22. Shapiro, B. J., Levade, I., Kovacikova, G., Taylor, R. K. & Almagro-Moreno, S.
Origins of pandemic Vibrio cholerae from environmental gene pools. *Nature*
*Microbiology* **2**, 16240 (2017).

23. Balasubramanian, D., López-Pérez, M. & Almagro-Moreno, S. Pandemic dynamics
and cholera emergence. in *Vibrio spp. Infections* (eds. Almagro-Moreno, S. & Pukatzki,
S.) vol. 1404 127–147 (Springer Nature, 2023).

24. DiRita, V. J. & Mekalanos, J. J. Periplasmic interaction between two membrane
regulatory proteins, ToxR and ToxS, results in signal transduction and transcriptional
activation. *Cell* **64**, 29–37 (1991).

25. DiRita, V. J., Parsot, C., Jander, G. & Mekalanos, J. J. Regulatory cascade controls
virulence in Vibrio cholerae. *Proceedings of the National Academy of Sciences of the*
*United States of America* **88**, 5403–5407 (1991).

26. Finkelstein, R. A. & LoSpalluto, J. J. Pathogenesis of experimental cholera.
Preparation and isolation of choleragen and choleragenoid. *The Journal of*
*experimental medicine* **130**, 185–202 (1969).

27. Miller, V. L., Taylor, R. K. & Mekalanos, J. J. Cholera toxin transcriptional activator
toxR is a transmembrane DNA binding protein. *Cell* **48**, 271–279 (1987).

28. Skorupski, K. & Taylor, R. K. A new level in the Vibrio cholerae ToxR virulence
cascade: AphA is required for transcriptional activation of the tcpPH operon. *Molecular*
*microbiology* **31**, 763–771 (1999).

29. Taylor, R. K., Manoil, C. & Mekalanos, J. J. Broad-host-range vectors for delivery of
TnphoA: use in genetic analysis of secreted virulence determinants of Vibrio cholerae.
*The Journal of Bacteriology* **171**, 1870–1878 (1989).

30. Waldor, M. K. & Mekalanos, J. J. Lysogenic conversion by a filamentous phage
encoding cholera toxin. *Science* **272**, 1910–1914 (1996).

31. Taylor, R. K., Miller, V. L., Furlong, D. B. & Mekalanos, J. J. Use of phoA gene
fusions to identify a pilus colonization factor coordinately regulated with cholera toxin.
*Proc National Acad Sci* **84**, 2833–2837 (1987).

32. Karaolis, D. K. *et al.* A Vibrio cholerae pathogenicity island associated with
epidemic and pandemic strains. *Proceedings of the National Academy of Sciences of*
*the United States of America* **95**, 3134–3139 (1998).

33. Faruque, S. M. & Mekalanos, J. J. Pathogenicity islands and phages in Vibrio
cholerae evolution. *Trends in microbiology* **11**, 505–510 (2003).

34. Jermyn, W. S. & Boyd, E. F. Characterization of a novel Vibrio pathogenicity island
(VPI-2) encoding neuraminidase (nanH) among toxigenic Vibrio cholerae isolates.
*Microbiology* **148**, 3681–3693 (2002).

35. Davies, B. W., Bogard, R. W., Young, T. S. & Mekalanos, J. J. Coordinated
regulation of accessory genetic elements produces cyclic di-nucleotides for V. cholerae
virulence. *Cell* **149**, 358–370 (2012).

36. O'Shea, Y. A. *et al.* The Vibrio seventh pandemic island-II is a 26.9 kb genomic
island present in Vibrio cholerae El Tor and O139 serogroup isolates that shows
homology to a 43.4 kb genomic island in V. vulnificus. *Microbiology* **150**, 4053–4063
(2004).

37. Taviani, E. *et al.* Discovery of novel Vibrio cholerae VSP-II genomic islands using
comparative genomic analysis. *FEMS Microbiology Letters* **308**, 130–137 (2010).

38. Haley, B. J. *et al.* Genomic and Phenotypic Characterization of Vibrio cholerae
Non-O1 Isolates from a US Gulf Coast Cholera Outbreak. *PLoS ONE* **9**, e86264
(2014).

39. Jaskólska, M., Adams, D. W. & Blokesch, M. Two defence systems eliminate
plasmids from seventh pandemic Vibrio cholerae. *Nature* **604**, 323–329 (2022).

40. Rivera, I. N., Chun, J., Huq, A., Sack, R. B. & Colwell, R. R. Genotypes associated
with virulence in environmental isolates of Vibrio cholerae. *Applied and environmental*
*microbiology* **67**, 2421–2429 (2001).

41. Faruque, S. M. *et al.* Genetic diversity and virulence potential of environmental
Vibrio cholerae population in a cholera-endemic area. *Proceedings of the National*
*Academy of Sciences of the United States of America* **101**, 2123–2128 (2004).

42. Gennari, M., Ghidini, V. & Lleo, M. M. Virulence genes and pathogenicity islands in
environmental Vibrio strains non-pathogenic to humans. *FEMS microbiology ecology*
**82**, 563–573 (2012).

43. Almagro-Moreno, S. & Taylor, R. K. Cholera: Environmental Reservoirs and Impact
on Disease Transmission. *Microbiology Spectrum* **1**, 149–165 (2013).

44. Lutz, C., Erken, M., Noorian, P., Sun, S. & McDougald, D. Environmental reservoirs
and mechanisms of persistence of Vibrio cholerae. *Frontiers in Microbiology* **4**, 375
(2013).

45. López-Pérez, M. *et al.* Ecological diversification reveals routes of pathogen
emergence in endemic Vibrio vulnificus populations. *Proc National Acad Sci* **118**,
e2103470118 (2021).

46. Nandi, B. *et al.* Rapid Method for Species-Specific Identification of Vibrio cholerae
Using Primers Targeted to the Gene of Outer Membrane Protein OmpW. *J Clin*
*Microbiol* **38**, 4145–4151 (2000).

47. Teh, C. S. J., Thong, K. L., Osawa, R. & Chua, K. H. Comparative PCR-based
fingerprinting of Vibrio cholerae isolated in Malaysia. *J Gen Appl Microbiol* **57**, 19–26
(2011).

48. Heidelberg, J. F. *et al.* DNA sequence of both chromosomes of the cholera
pathogen Vibrio cholerae. *Nature* **406**, 477–483 (2000).

49. López-Pérez, M. *et al.* Evolutionary model of cluster divergence of the emergent
marine pathogen Vibrio vulnificus: From genotype to ecotype. *mBio* **10**, 697 (2019).

50. Chatterjee, S. N. & Chaudhuri, K. Lipopolysaccharides of Vibrio cholerae I. Physical
and chemical characterization. *Biochimica Et Biophysica Acta Bba - Mol Basis Dis*
**1639**, 65–79 (2003).

51. Chatterjee, S. N. & Chaudhuri, K. Lipopolysaccharides of Vibrio cholerae II.
Genetics of biosynthesis. *Biochimica Et Biophysica Acta Bba - Mol Basis Dis* **1690**, 93–
109 (2004).

52. Chatterjee, S. N. & Chaudhuri, K. Lipopolysaccharides of Vibrio cholerae: III.
Biological functions. *Biochimica Et Biophysica Acta Bba - Mol Basis Dis* **1762**, 1–16
(2006).

53. Schnaitman, C. A. & Klena, J. D. Genetics of lipopolysaccharide biosynthesis in
enteric bacteria. *Microbiol Rev* **57**, 655–682 (1993).

54. Liang, W. *et al.* Sequence polymorphisms of rfbT among the Vibrio cholerae O1
strains in the Ogawa and Inaba serotype shifts. *Bmc Microbiol* **13**, 173–173 (2013).

55. Harris, J. B., LaRocque, R. C., Qadri, F., Ryan, E. T. & Calderwood, S. B. Cholera.
*Lancet* **379**, 2466–2476 (2012).

56. Rivera-Chávez, F. & Mekalanos, J. J. Cholera toxin promotes pathogen acquisition
of host-derived nutrients. *Nature* **572**, 244–248 (2019).

57. Faruque, S. M. & Mekalanos, J. J. Phage-bacterial interactions in the evolution of
toxigenic Vibrio cholerae. *Virulence* **3**, 556–565 (2014).

58. Mukhopadhyay, A. K., Chakraborty, S., Takeda, Y., Nair, G. B. & Berg, D. E.
Characterization of VPI pathogenicity island and CTXphi prophage in environmental
strains of Vibrio cholerae. *The Journal of Bacteriology* **183**, 4737–4746 (2001).

59. Houot, L. *et al.* Electrostatic interactions between the CTX phage minor coat protein
and the bacterial host receptor TolA drive the pathogenic conversion of Vibrio cholerae.
*J Biol Chem* **292**, 13584–13598 (2017).

60. Rodriguez-Valera, F., Martin-Cuadrado, A.-B. & López-Pérez, M. Flexible genomic
islands as drivers of genome evolution. *Curr Opin Microbiol* **31**, 154–160 (2016).

61. Dorman, M. J. *et al.* High quality reference genomes for toxigenic and non-toxigenic
Vibrio cholerae serogroup O139. *Sci Rep-uk* **9**, 5865 (2019).

62. O'Shea, Y. A. & Boyd, E. F. Mobilization of the Vibrio pathogenicity island between
Vibrio cholerae isolates mediated by CP-T1 generalized transduction. *FEMS
Microbiology Letters* **214**, 153–157 (2002).

63. Murphy, R. A. & Boyd, E. F. Three pathogenicity islands of Vibrio cholerae can
excise from the chromosome and form circular intermediates. *Journal of bacteriology*
**190**, 636–647 (2008).

64. Kirn, T. J., Bose, N. & Taylor, R. K. Secretion of a soluble colonization factor by the
TCP type 4 pilus biogenesis pathway in Vibrio cholerae. *Molecular microbiology* **49**,
81–92 (2003).

65. Kirn, T. J. & Taylor, R. K. TcpF is a soluble colonization factor and protective
antigen secreted by El Tor and classical O1 and O139 Vibrio cholerae serogroups.
*Infection and immunity* **73**, 4461–4470 (2005).

66. Megli, C. J. *et al.* Crystal structure of the Vibrio cholerae colonization factor TcpF
and identification of a functional immunogenic site. *Journal of molecular biology* **409**,
146–158 (2011).

67. Almagro-Moreno, S. & Boyd, E. F. Sialic acid catabolism confers a competitive
advantage to pathogenic Vibrio cholerae in the mouse intestine. *Infection and immunity*
**77**, 3807–3816 (2009).

68. Reddi, G., Pruss, K., Taylor, R. K., Cottingham, K. L. & Almagro-Moreno, S.
Catabolism of mucus components influences motility of Vibrio cholerae in the presence
of environmental reservoirs. *PLoS ONE* **13**, e0201383 (2018).

69. Almagro-Moreno, S., Napolitano, M. G. & Boyd, E. F. Excision dynamics of Vibrio
pathogenicity island-2 from Vibrio cholerae: role of a recombination directionality factor
VefA. *BMC microbiology* **10**, 306 (2010).

70. Carpenter, M. R., Rozovsky, S., Boyd, E. F. & DiRita, V. J. Pathogenicity Island
Cross Talk Mediated by Recombination Directionality Factors Facilitates Excision from
the Chromosome. *The Journal of Bacteriology* **198**, 766–776 (2016).

71. Grim, C. J. *et al.* Occurrence of the Vibrio cholerae Seventh Pandemic VSP-I Island
and a New Variant. *OMICS: A Journal of Integrative Biology* **14**, 1–7 (2010).

72. Grant, T.-A., López-Pérez, M., Haro-Moreno, J. M. & Almagro-Moreno, S. Allelic
diversity uncovers protein domains contributing to the emergence of antimicrobial
resistance. *Plos Genet* **19**, e1010490 (2023).

73. Ferenci, T. Trade-off Mechanisms Shaping the Diversity of Bacteria. *Trends
Microbiol.* **24**, 209–223 (2016).

74. Sheppard, S. K. Strain wars and the evolution of opportunistic pathogens. *Curr.
Opin. Microbiol.* **67**, 102138 (2022).

75. Chaturvedi, P., Agrawal, M. K. & Bagchi, S. N. Microcystin-producing and non-
producing cyanobacterial blooms collected from the Central India harbor potentially
pathogenic Vibrio cholerae. *Ecotoxicology and Environmental Safety* **115**, 67–74
(2015).

76. Huq, A. *et al.* Detection, Isolation, and Identification of Vibrio cholerae from the
Environment. *Curr Protoc Microbiol* **26**, 6A.5.1-6A.5.51 (2012).

77. Naha, A. *et al.* Development and Evaluation of a PCR Assay for Tracking the
Emergence and Dissemination of Haitian Variant ctxB in Vibrio cholerae O1 Strains
Isolated from Kolkata, India. *J Clin Microbiol* **50**, 1733–1736 (2012).

78. Zou, W. *et al.* Evaluation of Pulsed-Field Gel Electrophoresis Profiles for
Identification of Salmonella Serotypes. *J Clin Microbiol* **48**, 3122–3126 (2010).

79. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

80. Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its
Applications to Single-Cell Sequencing. *Journal of Computational Biology* **19**, 455–477
(2012).

81. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11**, 1–11 (2010).

82. Lowe, T. M. & Eddy, S. R. tRNAscan-SE: a program for improved detection of
transfer RNA genes in genomic sequence. *Nucleic acids research* **25**, 955–964 (1997).

83. Huang, Y., Gilna, P. & Li, W. Identification of ribosomal RNA genes in metagenomic
fragments. *Bioinformatics* **25**, 1338–1340 (2009).

84. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
DIAMOND. *Nature methods* **12**, 59–60 (2015).

85. Eddy, S. R. Accelerated Profile HMM Searches. *PLOS Computational Biology* **7**,
e1002195 (2011).

86. Tatusov, R. L. *et al.* The COG database: new developments in phylogenetic
classification of proteins from complete genomes. *Nucleic acids research* **29**, 22–28
(2001).

87. Haft, D. H. *et al.* TIGRFAMs: a protein family resource for the functional
identification of proteins. *Nucleic acids research* **29**, 41–43 (2001).

88. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and
accurate genomic comparisons that enables improved genome recovery from
metagenomes through de-replication. *Isme J* **11**, 2864–2868 (2017).

89. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W.
CheckM: assessing the quality of microbial genomes recovered from isolates, single
cells, and metagenomes. *Genome Research* **25**, 1043–1055 (2015).

90. Shakya, M. *et al.* Standardized phylogenetic and molecular evolutionary analysis
applied to species across the microbial tree of life. *Sci Rep-uk* **10**, 1723 (2020).

91. Nguyen, L.-T., Schmidt, H. A., Haeseler, A. von & Minh, B. Q. IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol
Biol Evol* **32**, 268–274 (2015).

92. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local
alignment search tool. *J Mol Biol* **215**, 403–410 (1990).

93. Altschul, S. Gapped BLAST and PSI-BLAST: a new generation of protein database
search programs. *Nucleic acids research* **25**, 3389–3402 (1997).

94. Skorupski, K. & Taylor, R. K. Positive selection vectors for allelic exchange. *Gene*
**169**, 47–52 (1996).

95. Choi, K.-H. & Schweizer, H. P. mini-Tn7 insertion in bacteria with single attTn7
sites: example Pseudomonas aeruginosa. *Nat. Protoc.* **1**, 153–161 (2006).

96. Millet, Y. A. *et al.* Insights into Vibrio cholerae intestinal colonization from
monitoring fluorescently labeled bacteria. *PLoS pathogens* **10**, e1004405 (2014).

97. Matson, J. S. Vibrio Cholerae, Methods and Protocols. *Methods Mol. Biol.* **1839**,
147–152 (2018).

98. Lee, M.-N., Kim, S.-K., Li, X.-H. & Lee, J.-H. Bacterial virulence analysis using brine
shrimp as an infection model in relation to the importance of quorum sensing and
proteases. *J Gen Appl Microbiol* **60**, 169–174 (2014).

99. Islam, M. S. *et al.* Role of cyanobacteria in the persistence of Vibrio choleraeO139
in saline microcosms. *Canadian journal of microbiology* **50**, 127–131 (2004).

100. O'Toole, G. A. Microtiter Dish Biofilm Formation Assay. *Journal of visualized*
*experiments : JoVE* (2011) doi:10.3791/2437.

101. Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the
next-generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

102. Gautreau, G. *et al.* PPanGGOLiN: Depicting microbial diversity via a partitioned
pangenome graph. *Plos Comput Biol* **16**, e1007732 (2020).

103. Li, L., Stoeckert, C. J. & Roos, D. S. OrthoMCL: identification of ortholog groups
for eukaryotic genomes. *Genome Research* **13**, 2178–2189 (2003).

104. Cohen, O., Ashkenazy, H., Belinky, F., Huchon, D. & Pupko, T. GLOOME: gain
loss mapping engine. *Bioinformatics* **26**, 2914–2915 (2010).

105. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res*
**49**, gkaa913- (2020).

1218

1219

**A**

Transcriptional regulator VC0176

Integrase VC0183

**B**

20000 (C24)
98.4% 98.7%
VcCHNf4 (C05)
99.8%
VSP-I PCG
93.6%
IRLE0062_S59 (C05)
99.9%
LPS Cluster PCG (99.9% identity)
99.9%
V060002 (C24)

**C**

VC0502
VC0503
VC0512
VC0513
VC0514
VC0515

**D**

TMA 21 (C24)
97.8%
H1 (C19)
98.3% 93.5%
VSP-II PCG
99.9% 99.1% 99.7%
N2723 (C24)
93.6% 97.9% 98.8% 99.2% 97.9%
MZO-3 (C24)
tRNA-direct repeat
tRNA-Met

environmental adaptations
viral origin
IS element

*Artemia salina* uninfected

*Artemia salina* infected
with *Vibrio cholerae*

**C**

Crustaceans
(*Artemia salina*)

Cyanobacteria
(*Microcystis aeruginosa*)

Mollusks
(*Crassostrea virginica*)