1

# Studies on the *Escherichia coli* ExbD Transmembrane Domain,

# Residue L132, and an Inhibitory Cyclic Peptide.

6      Bimal Jana[1,3], Dale Kopp[1,4], Mingchao Xie[1,5], Hema Vakharia-Rao[2,6] and Kathleen Postle[1,2]*

9    RUNNING TITLE:
10      ExbD homodimerizes through its transmembrane domain.
11
12
13    KEYWORDS:
14      ExbD, TonB, energy transduction, disulfide bond,
15      membrane protein, protonmotive force, *Escherichia coli*
16
17

18    [1]Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University

19    Park, PA, 16802, USA

20    [2]School of Molecular Biosciences, Washington State University, Pullman, WA 99164

21    *Address correspondence to Kathleen Postle at postle@psu.edu

22
23    Current addresses:
24
25    [3]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA 02114, USA
26
27    [4] Bristol Myers Squib, Bothell, WA 98021
28
29    [5]AstraZeneca, Waltham, MA, USA
30
[6] Bangalore 560016 India

**ABSTRACT**

The TonB system of *Escherichia coli* uses the cytoplasmic membrane protonmotive force (PMF) to

energize active transport of nutrients across the otherwise unenergized outer membrane.  Because it

overcomes limitations for nutrient diffusion through outer membrane size-limiting porins, it provides a

growth advantage and is widespread among Gram-negative bacteria.  It consists of three known

cytoplasmic membrane proteins, TonB, ExbB and ExbD that energize a variety of customized TonB-

dependent transporters in the outer membrane.  The sole ExbD transmembrane domain is proposed to

consist of residues 23-43 (Kampfenkel and Braun, 1992, J. Bacteriol. 174:5485-7).  Here we showed

that the charge and location of residue Asp25 were essential for activity of the TonB system, thus

identifying it as the only PMF-responsive element in the TonB system. The proposed boundaries of the

transmembrane domain α-helix were revised to consist of residues 23-39, with residues 40-43 initiating

the subsequent disordered region required for signal transduction (Kopp and Postle, 2020, J. Bacteriol.

202, e00687-19).  Trapping of disulfide-linked ExbD homodimers through T42C or V43C  prevented

TonB system activity that was restored by addition of the reducing agent dithiothreitol, indicating a

requirement for motion.  In *vivo* photo-cross-linking experiments suggested that motion was rotation of

ExbD transmembrane domains. Inactivity of ExbD L132Q, the first ExbD mutant identified, was likely

due to steric hindrance. A conserved and defined site of *in vivo* ExbD interaction with TonB was

identified.  Exogenous addition of a cyclic peptide based on that site inhibited ExbD-TonB interaction

while concomitantly decreasing iron transport efficiency.  This suggested that a novel antimicrobial

strategy against ESKAPE and other Gram-negative pathogens could be developed by targeting ExbD

protein-protein interactions.

## INTRODUCTION

Gram-negative bacteria are characterized by their distinctive outer membranes that serve as intrinsic barriers to environmental agents, antibiotics, and degradative enzymes. To overcome the concomitant nutrient limitations imposed by the outer membrane, the TonB system energizes transport across the outer membrane of a wide array of scarce, important nutrients such as iron-siderophores, heme, zinc, copper, cobalamin, nickel, maltodextrin, chitin oligosaccharides, and sucrose [for reviews see (1-8)]. The TonB system is also an essential component for pathogenesis by certain Gram-negative bacteria (9).

In *Escherichia coli* K12 the TonB-dependent transport ligands are iron-siderophores and cobalamin. The TonB system consists of integral cytoplasmic membrane proteins ExbB and ExbD, which harvest the protonmotive force (PMF) of the cytoplasmic membrane, and transmit it to the integral cytoplasmic membrane protein TonB, which then transmits mechanical energy to customized β-barrels in the outer membrane known as TonB-dependent transporters (TBDTs) (Fig.1). In the absence of PMF or TonB, ligands bind to TBDTs but do not cross the outer membrane or enter the periplasm. Because the TBDTs greatly outnumber TonB and compete for contact with it, it is likely that TonB engages in an energy transduction cycle involving transient interactions with individual TBDTs (10-12).

Structurally, the TBDT β-barrels consist of 22 strands, with a lumen that is occluded by an amino-terminal globular domain, called the cork or hatch. *In vivo*, the periplasmically exposed surface of the TBDT appears to signal that ligand is bound via a conformational change that allows for productive interaction with TonB (13-18). The cork domain moves *in vivo* during contact with TonB (18).

Among the three integral cytoplasmic membrane proteins in the TonB system, there are five transmembrane domains, any one of which could theoretically respond to the PMF of the cytoplasmic membrane. TonB and ExbD are both anchored by a single transmembrane domain with the majority of each protein occupying the periplasmic space (Fig.1) [(19-21). ExbB has three transmembrane domains,

with an N-out, C-in topology and primarily occupies the cytoplasm (22).  TonB and ExbD both depend

on ExbB for their proteolytic stability suggesting that ExbB constitutes a scaffolding for their assembly

[(23-25); unpublished results].  Consistent with that idea, the cellular ratio of TonB:ExbD:ExbB is 1:2:7

(11).  The TonB transmembrane domain remains homodimerized throughout the energy transduction

cycle, thus making the ratio 2:4:14 (12).  Our current model suggests that, as TonB initiates an energy

transduction cycle, its previously monomeric periplasmic domain becomes a homodimer, that can then

interact with ExbD homodimerized through its periplasmic domain; this transforms into a TonB-ExbD

heterodimer that subsequently configures TonB to energize active transport of nutrients through a TBDT

(Fig. 2).  In this model the proposed energy-dependent step is recycling of ExbD homodimers to an

energized state.

The sole potentially PMF-responsive residue in the TonB transmembrane domain, His20, can be

fully functionally replaced by a non-protonatable Asparagine residue, thus removing it from

consideration as part of the mechanism of energy transduction (26). Likewise, all of the potentially

PMF-responsive residues in the three ExbB transmembrane domains are excluded because non-

protonatable substitutions are active (27).  This leaves Asp25 in the ExbD transmembrane domain as the

sole actor that could potentially mediate PMF-dependence in the TonB system.

Because both active and inactive TonBs can bind to TBDTs that are both ligand-bound and

ligand-free both *in vivo* and *in vitro*, energy transduction cannot be explained by simple binding (14, 28-

34).  Only *in vivo* full-length active TonB leads to active transport of ligands through a TBDT,

suggesting that there must be a difference in the residues through which the active and inactive TonBs

bind TBDTs.  That difference appears to be manifested by the ability of TonB to interact correctly with

PMF-activated ExbD such that it achieves an "energized" conformation.

10-residue deletion scanning suggests that all ExbD residues are functionally important except

for a few amino acids at the cytoplasmic amino terminus (35). ExbD makes multiple *in vivo* contacts

during TonB-dependent energy transduction, including formaldehyde cross-linked ExbD-ExbD

homodimers, and ExbD-ExbB and ExbD-TonB heterodimers. Consistent with the formaldehyde cross-

linking studies, the ExbD periplasmic domain forms a disulfide cross-linked ExbD-ExbD homodimer

and ExbD-TonB heterodimer *in vivo*, with nearly the same ExbD residues involved in both complexes

(36). In contrast, TonB is essentially impervious to missense mutations, with H20 in the transmembrane

domain playing only a structural role [(26, 37-41); Potter and Postle, unpublished results]. Thus, TonB

appears to be a puppet with ExbD as the puppet-master.

The role of ExbD can be recapitulated in spheroplasts, which retain their PMF, as well as all of

the elements of the cytoplasm that the TonB system may require. The conformation of ExbD is PMF-

dependent and it is only by interaction with ExbD that TonB conformations respond to PMF (42).

There have been several indications as to the importance of ExbD Asp 25. ExbD D25N

inactivates ExbD (and thus the entire TonB system) and is dominant negative, consistent with being part

of a multiprotein complex (43). ExbD D25N mimics the effect of PMF collapse that prevents the ExbD-

TonB interaction characterized by *in vivo* formaldehyde cross-linking (29). In the proposed TonB

energy transduction cycle, ExbD D25N stalls the ExbD-TonB periplasmic domain interaction at Stage

II, the energy-<u>in</u>dependent stage, such that it cannot proceed to the energy-dependent ExbD-TonB

interaction [Fig. 2; (29, 36, 42-44)]. Like wild-type ExbD, ExbD D25N formaldehyde-cross-links

normally with ExbB monomer and forms homodimers *in vivo* (29).

Along with ExbD, MotB and TolQ participate in three paralogous cytoplasmic membrane

systems in *E. coli* K12. Each of the systems uses PMF to energize events at the outer membrane and has

two cytoplasmic membrane components in common: one protein with multiple transmembrane domains,

occupying primarily the cytoplasm (ExbB, TolQ, and MotA), and one protein anchored by a single

transmembrane domain, occupying primarily the periplasm (ExbD, TolR, and MotB) (45-48). TolR and

MotB carry an essential Asp residue in their single transmembrane domain at positions 23 and 32

respectively (47, 49), suggesting that ExbD Asp 25 might be equally important.

Here we present a detailed analysis of the ExbD transmembrane domain indicating the pivotal

role both charge and location of Asp25 play in TonB system response to cytoplasmic membrane PMF.

We redefine the boundaries of the transmembrane domain and identify an subsequent adjacent segment

that initiates a previously characterized disordered domain.  We show that the ExbD transmembrane

domains homodimerize,  appear to rotate relative to one another, and that, if dynamic movement is

prevented by disulfide trapping, ExbD ceases to function. These results suggest that the snapshots of

ExbB/D and MotA/B provided by recent cryo-EM structures do not yet capture the mechanism by which

they utilize the PMF (50-52).  We also exclude the overall importance of ExbD L132, first identified as

an inactive L132Q missense mutation, showing that several different amino acids can functionally

substitute at that position. Further, we identify an ExbD-specific cyclic peptide that, when added to *E*.

*coli* exogenously, prevents binding of ExbD to its protein partners and inhibits the TonB system.

## RESULTS

*The negative charge and location of Asp25 in the ExbD transmembrane domain are indispensable for*

*TonB system function*

The ExbD transmembrane domain is predicted to consist of residues 23-43 (20). The similarity of the ExbD transmembrane domain to those of paralogues TolR and MotB suggested that ExbD D25 (D23 in TolR and D32 in MotB) might be important (48, 49). To investigate that possibility, Cys substitutions were engineered individually at all predicted ExbD transmembrane domain residues from 23 to 43 on plasmid pKP999 under a propionate-inducible promoter [(29); see Fig. 3A for the primary amino acid sequence]. Individual Cys substitution mutants were expressed at chromosomal levels in strain RA1045 (ΔexbD, ΔtolQR), where the paralogues TolQR were also removed and could no longer serve as a source of limited but detectable TonB system activity (28, 53-55). All Cys-substituted ExbD mutant proteins were proteolytically stable and, excluding the slower migration of ExbD D25C and ExbD P39C, migrated at the same position as wild type ExbD in immunoblots (Fig. 4, lower panel). The slower migration of D25C appeared to reflect the loss of the Asp residue since D25N and D25A substitutions also decrease ExbD mobility in sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis (data not shown). A case where ExbD migrates more slowly because it appears to be chemically modified is considered below. Except for ExbD D25C, which was insensitive (tolerant) to all agents tested, the Cys substitutions of residues 23-43 supported sensitivity to colicin B, Ia, M and bacteriophage φ80 (Table 1). These TonB-system assays are the most sensitive, detecting the activity of as little as one TonB per bacterial cell (56).

To further test the importance of Asp25 in ExbD, all 19 other amino acids were substituted at that position. The conservative replacement ExbD D25E retained 5-10% wildtype activity in iron transport assays, which are the most discriminative of the TonB system assays, while the remaining 18

substitutions at Asp25 were completely insensitive in colicin and phage sensitivity assays, indicating

that they supported no TonB system activity (56).

Relocation of His20 in the TonB transmembrane domain from position 20 to a higher turn of the

spiral at position 24 supports detectable but decreased colicin and phage sensitivity (26).  To test the

importance of its position within the ExbD transmembrane domain, Asp25 was relocated to all other

positions in the ExbD transmembrane domain from position 23 to 43 on a plasmid template where

Asp25 was mutated to Ala.  In contrast to the situation with TonB His20, all the ExbD Asp25 relocation

mutants were completely insensitive to colicin and phage.  Taken together these results indicated that the

negative charge and location of ExbD Asp25 within the transmembrane domain were essential for its

activity.

*Sequence comparisons with ExbD paralogues TolR and MotB*

We had previously used the TOPCONS program to predict ExbB, TolQ, MotA and PomA

transmembrane domain boundaries (57). In that study, TOPCONS provided a prediction that was

congruent with a number of conserved residues within their transmembrane domains and led to a

redefinition of boundaries for the last two TMDs of each protein (27).  Here TOPCONS analysis of the

individual protein sequences was less helpful, potentially due to divergence in the function of these

proteins compared to ExbB and its paralogues.

The invariant and essential Asp residue found in ExbD, TolR, MotB and PomB transmembrane

domains did not align well using TOPCONS: the majority of homologies between the paralogous

proteins ExbD and TolR were missed; congruence between ExbD and MotB Cys substitutions that

formed high levels of disulfide-linked complexes were not optimized; and homologies between MotB

and PomB, both of which play similar roles in flagellar rotation, were not optimized (Fig 3A).  Because

ExbD appears to interact with the last two transmembrane domains of ExbB, it seemed likely that

TolR/MotB/PomB proteins should interact similarly with TolQ/MotA/PomA (27). This appeared to be

the case since the inconsistencies generated by the TOPCONS analysis were resolved when the

transmembrane domains were aligned using the invariant Asp residues (Fig. 3B). Notably, the ExbD

Cys substitutions that homodimerized most efficiently (described below) were now aligned with the

MotB substitutions that homodimerized most efficiently.

*ExbD homodimerizes through residues 23-63* in vivo.

We had previously identified residues in the region of residues 92-121 through which ExbD

periplasmic domains homodimerized *in vivo* (36). To study the ability of the ExbD transmembrane

domain to support homodimerization in the absence of its periplasmic domain, the DNA encoding ExbD

residues V23-P63 was cloned into the TOXCAT vector, pccKan (58). The cloned region included the

predicted ExbD transmembrane domain [(residues 23-43); (20)] and a subsequent unstructured domain

involved in signal transduction [(residues 44-63) (59, 60)].

As part of the TOXCAT fusion protein, if residues 23-63 homodimerized they would mediate

concomitant homodimerization of the amino terminal, cytoplasmically localized ToxR domain, thus

activating the chloramphenicol acetyl transferase gene (CAT) and supporting increased resistance to

chloramphenicol (CAM) compared to controls where the transmembrane domain does not

homodimerize (GlpA compared to GlpA G83I in the TOXCAT vector). Activity of carboxy terminally

fused periplasmic maltose binding protein (MBP) would serve as a control to demonstrate that the

inserted ExbD residues had translocated MBP to the periplasmic space.

To test homodimerization by ExbD residues 23-63, plasmid vector pccKan, plasmid pccGlpA,

plasmid pccGlpA G83I and pKP523 [pccExbD (*toxR-exbD$_{23-63}$-malE*)] were transformed into strain

NT326 [*araD*139, Δ(*argF-lac*)$_{U169}$, Δ*malE444, recA*1, *thi*, *strR* (61)] and KP1448 (NT326, *exbB*::Tn*10*).

Resistance to various concentrations of CAM, from 34 $\mu$g/ml to 1.0 mg/ml was measured by growth on

plates.  In NT326, both GlpA-TOXCAT and ExbD$_{23-63}$-TOXCAT supported strong growth on 125 $\mu$g/ml

CAM and poor growth at 500 $\mu$g/ml.  The GlpAG83I-TOXCAT mutant control supported poor growth

at 125 $\mu$g/ml and no growth at 500 $\mu$g/ml (Table 2).  In KP1448, where competition by wild-type ExbD

transmembrane domains was eliminated, ExbD$_{23-63}$-TOXCAT and GlpA-TOXCAT again supported

equally robust growth on 0.125 $\mu$g/ml CAM. Again, there was little growth of the negative control,

GlpA G83I-TOXCAT, at that concentration (Table 2).  The three TOX-CAT constructs containing

GlpA, GlpA G83I, or ExbD$_{23-63}$ residues equally supported growth with maltose as sole carbon source

and had equivalent levels of protein present on anti-MBP immunoblots (data not shown), indicating that

they were equally expressed and exported to the cytoplasmic membrane. We concluded that ExbD

residues 23-63, which included the predicted transmembrane domain, had an intrinsic ability to form

homodimers *in vivo* in both the presence and absence of ExbB.

*Identification of a face through which ExbD transmembrane domains homodimerize*

To determine if homodimers formed through the predicted ExbD transmembrane domain,

disulfide-mediated homodimerization of ExbD Cys-substitutions at residues 23-43 expressed at

chromosomal levels was evaluated using iodine (I$_2$) catalysis, a technique previously used to study

homodimerization of TolR and MotB transmembrane domains (46, 47).  In Fig. 4A, several of the Cys

substitutions resulted in bands on non-reducing anti-ExbD immunoblots that, at ~ 31 kDa, were twice

the mass of monomeric ExbD (~15 kDa) (45).  The ~ 31 kDa complex was absent from an ExbD

deletion strain (RA1045, $\Delta$*exbD, $\Delta$tolQR*) and from wild-type (Cys-less) ExbD samples (WT and

pExbD), indicating that it represented a disulfide-linked ExbD homodimer.  The efficiency of dimer

formation varied with the position of the Cys residue.  The most abundant dimers were formed by

residues I24C, M27C, F34C, with a periodicity of ~3-4 amino acids (and a gap at 30-31C), consistent

with an α-helical arrangement of each ExbD transmembrane domain and suggesting that they

homodimerize through one face (Fig. 4B).

Similar results are seen for paralogues MotB and TolQ (46, 47). TolR, MotB, and as we

demonstrated above, ExbD, all carry a functionally important Asp residue in their single transmembrane

domain at positions 23, 32, and 25 respectively (47, 49). Further similarity was demonstrated by ExbD

I24C, which existed entirely in homodimeric form, with ExbD D25C existing mostly in monomeric

form in this assay (Fig. 4A). A similar relationship was seen with corresponding MotB transmembrane

domain Cys substitutions at A31C and D32C (46), suggesting that the transmembrane domains

homodimerize similarly. In contrast to a predicted α-helical configuration, each of the ExbD Cys

substitutions from L40C- V43C formed homodimers with little to no monomer. It was not clear if the $I_2$

oxidant or ExbB was needed to form any of the complexes.

Although all the ExbD mutants were expressed at near chromosomal levels as determined by

reducing SDS polyacrylamide gel electrophoresis (Fig. 4A, lower), there was differential antibody

detection of both monomers and homodimers from the same samples immunoblotted under non-

reducing conditions (Fig. 4A, upper). For example, V26C monomer was detected at just slightly less

than wild-type monomer ExbD levels on the reducing immunoblot (Fig. 4A, lower) but was much more

strongly detected compared to the wild-type ExbD on the non-reducing immunoblot (Fig. 4A, upper)

and did not form homodimers. In contrast, ExbD P39C appeared to be undetectable on the non-reducing

immunoblot and was present at wild-type chromosomal levels on the reducing immunoblot. The

differential antibody detection suggested that there was structural variation among ExbD Cys

substitutions under non-reducing conditions, even in the presence of SDS.

*ExbD transmembrane domain Cys substitutions 23-39 require oxidant for homodimerization whereas*

*Cys residues at 40-43 do not.*

To determine if oxidant was required for transmembrane domain homodimerization, the

experiment in Fig. 4A was repeated but without the use of $I_2$ (Fig. 5).  Near the cytoplasmic side of the

predicted transmembrane domain, I24C spontaneously formed only modest levels of homodimers in

contrast to the high level I24C formed when $I_2$ was present, thus placing it within the transmembrane

domain.  In contrast, ExbD F23C appeared to be on the transmembrane domain boundary because it

formed similarly low levels of homodimers with and without $I_2$ (compare Figs. 4A and 5).   Consistent

with an apparent transmembrane domain location, none of the residues from D25C through P39C

spontaneously formed significant levels of homodimers in the absence of $I_2$.

In contrast, residues from ExbD L40C- V43C each spontaneously formed high levels of

homodimers in the absence of $I_2$, suggesting they were not part of the transmembrane α-helix and likely

constituted a region of intrinsic disorder (compare Figs. 4A and 5).

*Trapping ExbD homodimers through T42C or V43C largely inactivates the TonB system.*

Because ExbD T42C and V43C resulted in almost 100% homodimer formation in the absence of

$I_2$ oxidant in Fig. 5, it was possible to determine whether dynamic movement of those residues was

required for TonB system activity.  We assayed the ability of plasmids expressing each variant to

support TonB-dependent iron transport with or without the reducing agent dithiothreitol (DTT) and

immunoblotted those same cultures to determine relative steady state levels of monomer and

homodimer.  As seen previously, both ExbD T42C and V43C were nearly 100% homodimerized in the

absence of DTT (Fig. 6A, upper panel).  All ExbD Cys substitutions were expressed at chromosomally

encoded levels (Fig. 6A, lower panel).

In Fig. 6B, DTT had an inhibitory effect, decreasing the initial transport rate of the wild-type

control to 77% of the rate seen in the absence of DTT.  We compensated for the fact that DTT was

somewhat inhibitory to wild-type transport by calculating the mutant transport rates as percentiles of the

wild-type transport rates.  The effect of DTT on the previously trapped inactive homodimers of

ExbDT42C and ExbDV43C, was to release them as monomers, thus restoring activity (Fig 6A, upper

panel).  Transport supported by trapped ExbD V43C was restored from 9% of wild-type -DTT to 89% of

wild-type +DTT (Fig. 6C; Table 3). Similarly, DTT restored transport supported by ExbD T42C from

23% of wild-type -DTT to 77% of wild-type +DTT.  These results indicated that inactivity of the

disulfide-linked homodimers was due to restriction of required conformational changes rather than the

inability of Cys to functionally replace Val or Thr.  The results were not contradictory with the spot titer

data from Table 1 where strains expressing ExbD T42C or V43C were sensitive to colicins and phage

due to the presence of residual monomer.   The highly sensitive colicin spot titers detect activity from as

little as 1 molecule of active TonB (56, 62).

The results were consistent with the existence of an energy transduction cycle where, if ExbD is

trapped in a homodimeric configuration such as Stage I or II, it cannot progress through the cycle (Fig.

2).  Residues L40-V43 are contiguous with a region of intrinsic disorder from 44-63 that dynamically

interacts with a second ExbD, with ExbB and with TonB *in vivo* (35, 59, 60).

*ExbD T42C appears to undergo increased modification in the absence of ExbB.*

Data presented in Fig. 6A (lower panel) suggested that monomeric ExbD T42C might be

chemically modified, because while a majority of ExbD T42C was found migrating at the same position

as wild-type ExbD on immunoblots of reducing SDS polyacrylamide gels, a smaller amount appeared to

migrate at a slightly higher apparent mass.  To determine the parameters that affected the potential

modification, the migration of ExbD T42C was evaluated on immunoblots of strains lacking ExbD, both

ExbB and ExbD, or both ExbD and TonB (Fig. 7).  As seen in Fig. 6A, both unmodified and modified

ExbD T42C were again observed, with ~ 2/3 in the unmodified form (Fig. 7, lane 6). The absence of

ExbB (and paralogue TolQ) converted all the ExbD T42C into the higher mass form (Fig. 7, lane 7).  In

the absence of TonB, the relative proportion of the two forms was reversed relative to the wild-type

situation, with ~2/3 of the ExbD T42C found at the position of the higher mass form (Fig. 7, lane 8).   It

is not clear whether the modification has any *in vivo* relevance for the mechanism of TonB-dependent

energy transduction. At the very least it provides evidence of *in vivo* interactions between ExbD and

ExbB. It is included here for future researchers.  Apparent chemical modification of full-length wild-

type ExbD also occurs when an inhibitory ExbD peptide (residues 44-63) is expressed endogenously,

resulting in the appearance of a higher mass ExbD species (63).

*Transmembrane domain Cys residues require ExbB for homodimerization whereas those of the*

*potentially disordered domain do not.*

327             Because the ExbD$_{23-63}$-TOXCAT fusion protein homodimerized in the absence of ExbB, we

tested the role of ExbB in ExbD homodimerization through its transmembrane domain by expressing

each of the Cys-substituted proteins in RA1017 (*ΔexbBD, ΔtolQRA*).  Unexpectedly, in light of the

TOXCAT results, the absence of ExbB prevented the previously detected I$_2$-dependent disulfide-linked

dimerization through ExbD substitutions F23C-P39C, even though I$_2$ was present (compare Figs. 4 and

8).  This finding notably contrasts with MotB transmembrane domain homodimer formation through

engineered Cys substitutions, which does not depend on MotA (46).

334             In contrast, even without ExbB, residues contiguous to the transmembrane domain, L40C, A41C

and V43C, formed homodimers, with V43C again achieving nearly 100% dimer formation (Fig. 8). The

results suggested that because two ExbD molecules were in sufficient proximity to homodimerize

through an adjacent region of likely intrinsic disorder, it seemed likely that the transmembrane domains

also interacted, but in a way not captured by disulfide cross-linking in the presence of I$_2$.  In addition,

these results suggested that the source of ExbB-independent homodimerization by the ExbD$_{23-63}$-

TOXCAT fusion protein resided in the disordered domain.

In the absence of ExbB, ExbD T42C formed essentially no homodimers, suggesting either that

the Cys substitution was the site of chemical modification observed in Fig. 7 or that without ExbB, two

T42C residues were no longer in proximity for disulfide formation.  The latter explanation was excluded

because A41C and V43C on either side of T42C formed disulfide-linked homodimers (Fig. 8). We

therefore conclude that chemical modification occurred at ExbD T42C in the absence of ExbB.

In vivo *photo-cross-linking suggests that ExbD transmembrane domains rotate relative to each other*

*and relative to ExbB*.

Disulfide linkage of ExbD Cys substitutions trapped transmembrane domains in a static

homodimeric state where the most abundant cross-linking occurred between residues on an apparently

helical face (Fig. 4A). Nonetheless, there was minor cross-linking that occurred through two ExbD

D25C residues on an opposing face of that helix, consistent with the idea that the transmembrane

domain helices might rotate relative to one another.  Because the homodimerized transmembrane

domains of TolR, which can functionally substitute for ExbD in the TonB system, rotate relative to one

another *in vivo* (47), we wanted to determine if that was also true for ExbD.  Rather than disulfide cross-

linking, we used *in vivo* photo-cross-linking by the reagent *p*-benzoyl-l-phenylalanine (pBpa), which

inserted efficiently at each amber codon for T21, V23, I24, D25, M27, V29, I32, D34, M35, and A37

(Fig. 9A, lower panel; monomeric ExbD).

In the presence of ultraviolet light, the ExbD pBpa monomers were pulled into three higher mass

complexes on immunoblots: ExbD homodimers and two ExbD-ExbB complexes with different apparent

masses (termed "upper" and "lower") characterized previously [Fig. 9A, upper panel (59)].  Each of the

ExbD pBpa substitutions formed homodimers and to different extents. Abundant homodimers were

formed by I24pBpa and F34pBpa, two sites that, as Cys substitutions, were part of the proposed α-

helical dimerization face (compare Figs. 4 and 9A). In addition, equally abundant homodimers were

formed by D25pBpa and V29pBpa, two residues on the opposite side of that face (Fig. 9 A and B).

Taken together, the results indicated that each of the transmembrane domain pBpa substitutions had

access to a second, homodimerized ExbD, most likely through its transmembrane domain.  To achieve

contacts through residues on opposite faces of the α-helix, the transmembrane domain must have moved

relative to a second ExbD, possibly by rotation.  Because all the transmembrane domain ExbD pBpa

substitutions formed complexes with ExbB (upper), they may have rotated relative to ExbB as well. Half

of the pBpa-substituted ExbDs also formed complexes with ExbB (lower) (Fig. 9A).

*ExbD L132 plays only a structural role in the TonB system.*

The Braun lab isolated the first inactive missense mutation in the ExbD periplasmic domain,

L132Q (43).  It stalls TonB at Stage I of the energy transduction cycle and, unlike wild-type ExbD, does

not formaldehyde cross-link to TonB *in vivo* [Fig. 2; (42)].  Deletion of the last 10 residues of ExbD

(Δ132-141) render it inactive, possibly due to the loss of L132 (35).  To determine if L132 was an

essential residue like D25 in the ExbD transmembrane domain, we engineered substitutions of all 19

amino acid residues at position L132 in pKP999.  The relative activity of the ExbD variants was

assessed by ability to complement the RA1045 Δ*exbD* mutation in a growth assay in M9 minimal

medium, where iron is limiting and only the TonB+ strains grow.  As shown in Table 4, L132 is not an

essential residue since it could be substituted with Cys, Val, Ile, Met, Phe, and Trp and still retain full

activity.  It seems likely that the Gln substitution in L132Q inactivated the TonB system by preventing

ExbD interaction with TonB through steric hindrance.

*A cyclic peptide derived from ExbD inhibits TonB system activity and prevents formaldehyde-specific*

*TonB-ExbB and TonB-ExbD cross-linked complexes* in vivo

The TonB system is a virulence factor for many Gram-negative bacteria (9). The periplasmic

domains of two key proteins in the system, ExbD and TonB, and their required interaction for

subsequent active transport through outer membrane TBDTs make it an appealing target for novel

antibiotic development. In general, antibiotics that inhibit periplasmic activities need possess only the

hydrophilic chemical characteristics that allow diffusion through outer membrane porins to access their

targets, a distinct advantage over those that must additionally cross the hydrophobic cytoplasmic

membrane (64).

Periplasmic ExbD residues 44-63 are disordered and thus a likely site for interaction with other

proteins (60, 65). We began to identify inhibitory regions of ExbD using endogenous secretion of

promising small peptides into the periplasmic space. Secretion of a disordered region peptide (residues

44-63) inhibits TonB-dependent iron transport by inhibiting an essential ExbD-TonB interaction, thus

establishing a proof of principle (63).

In this work we targeted interactions between ExbD and TonB periplasmic domains using

exogenously added peptides. Cyclic rather than linear peptides were chosen to decrease the likelihood

of proteolytic degradation by periplasmic proteases. None of the peptides contained cleavage sites for

the outer membrane protein, OmpT (66). Inhibitory cyclic peptides were designed based on three

regions of ExbD that are essential for TonB system activity *in vivo* [Fig. 10; (35, 36, 44)]: Region 1,

residues 44-63 from the ExbD periplasmic disordered region adjacent to the transmembrane domain, an

essential motif required for the ExbD-TonB PMF-dependent interaction (59); Region 2, residues 92-121

that are required for *in vivo* formaldehyde cross-linking of all known ExbD complexes (ExbD-ExbD,

ExbD-TonB, and ExbD-ExbB) (35); and Region 3, the region surrounding ExbD residue, L132, where a

single substitution, L132Q, completely abolishes activity (43).

The cyclic peptides from Region 1 of ExbD were the motifs V-K-V-N-L-P and V-K-V-N-L-P-A,

where the latter contained a carboxy terminal alanine to provide more conformational flexibility. For

Region 3, the sequence of ExbD residues 129-134 (K-I-G-L$_{132}$-V-G) was chosen along with the

corresponding mutant peptide representing ExbD L132Q (K-I-G-Q$_{132}$-V-G).

Because the larger ExbD Region 2 (residues 92-121) lacks specifically identified targets and has

extensive conservation of residues among most of the Gram-negative ESKAPE pathogens responsible

for some of the most deadly acquired hospital infections [(Fig. 10), (67)], studies of protein-protein

interactions were used to further guide the choice of residues.  The sequence I-F-F-R-A-D (residues 102-

107) was deemed likely to be a site of ExbD-ExbD interaction because it contains a concentration of

hydrophobic residues (IFFA), which are commonly found at protein-interaction interfaces (68).  In

addition, three of the top five "hotspot" amino acid preferences which contribute the most binding

energy in a protein interaction, were present within the semi-conserved sequence region shown in Fig.

10  [Ile, Arg, and Asp (69)]. Asp 207 was excluded from the cyclic peptide to facilitate transit through

porins in the outer membrane which have a size limit of ~ 600 da (70).

*--Cyc-IFFRA inhibits TonB-dependent activity*.  To eliminate potential efflux of peptides out of

the periplasm, strain SH03 (W3110, *tolC*::Tn*10*) was used (71). Initial rates of TonB-dependent $^{55}$Fe-

ferrichrome transport were determined in the presence of cyclic (cyc) peptides cyc-VKVNLP and cyc-

VKVNLPA from Region 1; cyc-IFFRA from Region 2; cyc-KIGLVG, and cyc-KIGQVG from Region

3 (Fig. 11).  In the presence of cyc-IFFRA, the $^{55}$Fe-ferrichrome initial transport rate was reduced to

~30% of the SH03 + DMSO control, indicating that it interfered with TonB system activity.  Cyc-

VKVNLP, cyc-VKVNLPA, cyc-KIGLVG, and cyc-KIGQVG had no effect on $^{55}$Fe- ferrichrome

transport.  These results suggested that either the peptides were not inhibitory or that they did not

efficiently enter the cells. In any case the L132Q mutation results were uninterpretable since the Region

3 peptide encoding the wild-type version had no detectable effect.

*--Cyc-IFFRA inhibits formation of TonB-ExbD and, surprisingly, TonB-ExbB complexes* in vivo.

Stage III of the TonB energy transduction cycle is characterized by the PMF-dependent formation of

formaldehyde cross-linked ExbD-TonB complexes (Fig. 2).  Cys substitutions K97C and T109C on

either side of the Region 2 peptide-derived sequence IFFRA capture multiple interactions with Cys

substitutions in the periplasmic TonB carboxy terminus *in vivo*, and formation of those disulfides is

significantly reduced or nonexistent in the PMF-unresponsive mutant, ExbD D25N (44).  Taken together

these results suggested that the inhibitory effect of cyc-IFFRA could be due to inhibition of Stage III of

the energy transduction cycle.  To test that hypothesis, strain SH03 was treated with formaldehyde in the

presence of the five cyclic peptides evaluated in Fig. 11 and processed for immunoblotting with both

anti-ExbD and anti-TonB antibodies (Fig. 12).

Consistent with its inhibition of iron transport in Fig. 11, the Region 2 cyc-IFFRA peptide

prevented the Stage III TonB-ExbD complex detected with both anti-TonB antibody (Fig. 12A) and

anti-ExbD antibody (Fig. 12B).  It also unexpectedly decreased formation of TonB-ExbB complexes

compared to the control (SH03 + DMSO), but not TonB-Lpp, TonB-OmpA or TonB-FepA complexes.

Formaldehyde cross-linking of TonB with Lpp and OmpA appear to be off-pathway contacts made by

unenergized TonB since a double *lpp, ompA* mutant has only slightly reduced iron transport (14). The

FepA complex likewise represents an inactive TonB-FepA complex which occurs with unenergized

TonB (72).  The cyc-IFFRA peptide had no effect on ExbD homodimer or ExbD-ExbB heterodimer

formation.

Inhibition of the TonB-ExbB complex by a cyclic peptide corresponding to a distal periplasmic

region of ExbD suggested that this region of ExbD played a role in assembly or conformation of the

TonB-ExbB interaction *in vivo*.  While monomeric formaldehyde is a probe for close contacts (73), the

absence of a formaldehyde-cross-linked complex only means that the formaldehyde-cross-linkable

residues have moved out of correct apposition--due either to dissolution of the complex or to a change in

conformation of one or more proteins within the complex.

The cyc-VKVNLPA peptide from Region 1 appeared to formaldehyde cross-link to ExbD

monomer (labeled "?") and disrupt ExbD homodimer cross-linking (Fig. 12B), consistent with our

identification of ExbD homodimerization through Region 1 residues in full-length ExbD (59).

Monomeric formaldehyde inserts a $CH_2$ group between two reactive residues (K, Y, H, C $\ggg$ R, W)

(74). It seems likely that cyc-VKVNLPA formaldehyde cross-linked to ExbD monomer through its Lys

residue. None of the other peptides contained strongly cross-linkable residues, which explains why

binding to ExbD monomer was not detected with the inhibitory cyc-IFFRA peptide from Region 2. If

cyc-VKVNLPA did cross-link to ExbD monomer, it means that it was able to penetrate the outer

membrane, even though it was the largest of the peptides. This result suggested how sensitive

translocation across the outer membrane, presumably via porin proteins, was to overall conformation

because the smaller version of the same peptide that also contained a cross-linkable Lys residue

apparently did not enter the periplasm. Alternatively, it did, but could not bind to ExbD. Because cyc-

VKVNLPA did not affect the iron transport rate in Fig. 11, it was not a candidate to be a TonB system

inhibitor. Cyc-IFFRA, however, showed promise.

**DISCUSSION**

Outer membrane active transporters of Gram-negative bacteria, including ESKAPE pathogens,

are powered by the TonB-system (75), which is one of three systems in *E. coli* K12 that each use two

similar cytoplasmic membrane proteins and the PMF to power cell envelope events:  ExbB and ExbD of

the TonB system energize active transport across the outer membrane (76); TolQ and TolR power

proper outer membrane invagination (77); MotA and MotB act as stators for flagellar rotation (78).   Of

the three systems, the TonB system is the least complex so far with ExbB/D and TonB located in the

cytoplasmic membrane and a variety of TBDTs in the outer membrane (Fig. 1), although there is

evidence for additional mystery proteins (79, 80).  What they all have in common are residue similarities

in the last two transmembrane domains of ExbB/TolQ/MotA and an invariant Asp residue in

approximately the same position in the single transmembrane domain of ExbD/TolR/MotB.  Other than

similarities among their transmembrane domains, the remaining residues most likely account for their

disparate functions. The overall mechanism of TonB-mediated energy transduction remains a mystery.

Results from the also-mysterious Mot and Tol systems provide informative comparisons.

*In vivo* studies of the paralogue MotB transmembrane domain demonstrate the importance of

Asp32 (D25 of ExbD; D23 of TolR) to flagellar rotation, where out of 15 substitutions only the

conservative replacement D32E shows a slight amount of activity (49).  MotB (308 residues) forms

homodimers through a face on its single α-helical transmembrane domain (46, 81).  MotB also binds to

peptidoglycan through residues ~ 209-227, a logical function for a flagellar rotation stator (82, 83).  The

need for a "plug" (MotB residues 52-65, C-terminal to the transmembrane domain) is explained if

unassembled MotA/B complexes leak protons until they are assembled in the flagellar motor (84).

*In vivo* studies on paralogue TolR (142 residues in length) show that two TolR transmembrane

domains homodimerize in a pattern similar to the MotB transmembrane domains (Fig. 3B). Like MotB,

TolR has a conserved Asp residue in its transmembrane domain that is inactivated by the D23R

substitution, while the conservative D23E substitution retains lower but detectable activity (48). ExbB/D

and TolQ/R are the most closely related paralogues and can in fact substitute (~ 10%) for one another in

their respective interactions with either non-cognate TonB or Tol-Pal proteins (45, 53-55, 85). Cys

substitution Y117C in the periplasmic carboxy terminus traps TolR such that it cannot undergo needed

conformational transformations (86). Similarly, disulfide-linked homodimers through the corresponding

residue in ExbD (Y112C) are inactive (36). Both results speak to the dynamic nature of their respective

periplasmic domain interactions.

Unlike the MotB transmembrane domains that apparently retain a static relationship and thereby

support binding of its periplasmic domain to peptidoglycan, homodimeric TolR transmembrane domains

appear to rotate relative to one another. Cys substitutions on opposite sides of the transmembrane

domain helix [L22 and V24 (corresponding to ExbD I24 and V26, Fig. 3B),] trap TolR as disulfide-

linked homodimers that inactivate the Tol system. Addition of the reducing agent DTT reduces the

disulfide bond and restores activity (47). This requirement for movement raises a question whether the

TolR periplasmic domain binds to peptidoglycan *in vivo*. Consistent with that, TolR is less than half the

length of MotB, seemingly making it less likely to access the peptidoglycan. In addition, TolR also does

not have an identifiable plug. In a contrasting view, TolR is hypothesized to reach the peptidoglycan as

a disordered protein (87). Cryptic peptidoglycan binding is observed for a truncated TolR periplasmic

domain, but it is not clear how the binding affinity compares to that of canonical OmpA peptidoglycan

binding (88).

*The ExbD transmembrane domain*. At 141 residues, ExbD is essentially the same length as

TolR (45). Braun et al. were the first to show that ExbD D25N is inactive (43). Our *in vivo* studies

described here on the predicted ExbD transmembrane domain (residues 23-43) were late to the party but

were somewhat more thorough from a mutagenic point of view. They first of all indicated that, like the

MotB and TolR transmembrane domains, the ExbD transmembrane domain homdimerized in an α-

helical configuration.

All residues in the ExbD transmembrane domain except Asp25 could be substituted with Cys

and retain function in colicin and phage sensitivity assays. Replacement by none of the remaining 19

amino acids at residue 25 except the conservative replacement D25E supported any activity, thus

aligning well with the importance of the transmembrane domain Asp in MotB and TolR (48, 49).  For

the first time, alternative positions for Asp25 within the transmembrane domain were assessed, showing

that ExbD Asp25 did not function at any other position within the transmembrane domain. Because

participation of the TonB and ExbB transmembrane domains in PMF-responsiveness is ruled out, it

appears that in the TonB system PMF is engaged solely through ExbD Asp25 (26, 27).  Consistent with

that, progression to Stage III in the energy transduction cycle is prevented either by the protonophore

CCCP (carbonyl cyanide *m*-chlorophenyl hydrazone) that collapses PMF or by the unresponsive *exbD*

D25N mutation [Fig. 2; (29, 42)].

*Two different domains in the ExbD amino terminal region*. Residues 23-43 were originally

proposed to constitute the ExbD transmembrane domain (20). Our results suggested instead that the

functional boundaries of the ExbD transmembrane domain could be revised to extend only from ExbD

F23 to P39.  A set of three ExbD Cys substitutions in this region appeared to represent an α-helical

configuration, requiring both oxidant and the presence of ExbB to form disulfide-linked homodimers

through those residues.  This revision somewhat positions the ExbD transmembrane domain between

two prolines at residues 22 and 39, that are conserved in TolR but not MotB (Fig. 3).

In contrast, each sequential residue from L40C through V43C homodimerized strongly without

the requirement for either added oxidant or ExbB, suggesting that they constituted an intrinsically

disordered domain.  [ExbD T42C did not homodimerize in the absence of ExbB because it appeared to

be chemically modified.]  Thus, ExbD residues L40-V43 appeared to initiate the previously identified

intrinsically disordered region of ExbD that terminates with residue 63, homodimerizes, and contains the

conserved motif V45-V47-L49-P50 that is required for signal transduction of PMF to the TonB

periplasmic domain (59, 60).

The $ExbD_{23-63}$-TOXCAT fusion protein, where none of the residues was Cys-substituted,

homodimerized either through ExbD transmembrane domain residues (residues 23-39) or an apparently

disordered region (residues 40-63), or both.  That result ruled out a requirement for participation of the

Dsb system in $ExbD_{23-63}$ homodimerization.  The $ExbD_{23-63}$-TOXCAT fusion protein homodimerization

did not require ExbB, indicating an intrinsic ability to homodimerize that was consistent with previous

studies identifying homodimerization through residues 44-56 using *in vivo* photo-cross-linking (59) and

disulfide-linked homodimerization of ExbD through Cys substitutions in its periplasmic domain

(residues 92-121) that also does not require ExbB *in vivo* (36).  It thus appeared that both disordered and

ordered regions of the ExbD periplasmic domain homodimerized independently of ExbB and it is likely

that the observed homodimerization of $ExbD_{23-63}$-TOXCAT both in the presence and absence of ExbB

was due to the presence of the disordered region and not the transmembrane domain.  If the entire

periplasmic domain of ExbD homodimerizes independently of ExbB, perhaps it might initiate formation

of ExbD transmembrane domain homodimers rather than vice versa.

*Required motion of the ExbD transmembrane domains*. Because L40C-V43C homodimerized in

the absence of ExbB, it suggested that the ExbD residues 23C-39C were likely also homodimerized due

to their proximity in the absence of ExbB, but without formation of disulfide-linked homodimers.

Combined, these results suggested that the ExbD transmembrane domains had the capacity to

homodimerize in two different configurations *in vivo* which could be consistent with a requirement for

motion.  The importance of motion was confirmed when spontaneously disulfide-linked Cys

substitutions at positions T42 and V43 inactivated TonB-dependent transport and reduction of the

disulfides reactivated it. These results mirrored similar approaches showing that TolR transmembrane

domains rotate relative to one another (47).

Results from *in vivo* photo-cross-linking studies suggested that the motion could be rotation

because sites on all faces of the ExbD transmembrane domain participated in formation of homodimers.

Consistent with that idea, *in vivo* photo-cross-linking also showed that all sites investigated in the ExbD

transmembrane domain formed photo-cross-linked complexes with ExbB.

Thus, two different techniques have now identified the likely rotation of ExbD/TolR

transmembrane domain homodimers relative to one another *in vivo*. By assessing disulfide-linked

homodimers, Celia et al. also suggested that the ExbD transmembrane helices rotated with respect to one

another (89).

*Results in the context of recent subcomplex cryo-EM structures.* The recent cryo-EM structures

of ExbB/D and MotA/B from several species show them to be asymmetric subcomplexes of 5

ExbB/MotA and 2 ExbD/MotB (51, 52, 89). The congruence of the proposed structures is striking,

especially since it is seen across different species and with the GldL/M proteins involved in gliding

motility of *Flavobacterium johnsoniae* (90). Authors of the MotA/B cryo-EM papers suggest that

MotA rotates around MotB homodimer, held static by interaction with peptidoglycan (51, 52). Hennell-

James et al. suggests that the GldL pentamer rotates relative to the GldM homodimer (like MotB) (90).

It has been suggested that these admittedly beautiful structures provide the key to the mechanism of the

protein complexes that all use PMF for energy (91) and that ExbB/D follows the same mechanism as

MotA/B (51).

An argument can be made for the validity of the MotAB structure because the first

transmembrane domain of MotA, which has four rather than three transmembrane domains like ExbB

and TolQ, stands in for their respective TonB/TolA transmembrane domains, thus obviating concerns

that TonB is not part of the ExbB/D cryo-EM structures and providing reassurance for the other

structures that, because they all exhibit the same 5:2 asymmetric architecture, they accurately reflect the

*in vivo* state of the complexes. Nonetheless, there is still a need to understand how and where

TonB/TolA fit into ExbB,TolQ/ExbD,TolR complexes and to address the cellular ratios of 7

ExbB/TolQ:4 ExbD/TolR:1TonB/TolA (11, 48, 92).

More remains to be discovered about the mechanism of these proteins. As Deme et al. states "No

ion permeation across the bilayer was seen in any simulation, supporting the idea that the complexes

currently seen will require rearrangement for activity." (51). Celia et al. likewise did not see a channel

in ExbB5/D2 through which a proton could be translocated and suggest that structural plasticity is likely

important (89). In support of a general need to incorporate *in vivo* results into interpretations based on

structural information, recent data from Nilaweera et al. demonstrate the importance of studying the

TonB system and its outer membrane TBDTs in their native cellular environment compared to *in vitro*

studies (93).

Evidence for the *in vivo* existence of ExbB tetramers and MotA tetramers (as opposed to

pentamers) is equally persuasive, and has the advantage that all required proteins, known and unknown,

are present in whole cells (79-81). For example, ExbB formaldehyde cross-links as a dimer of

homodimers *in vivo* (80). In another example, ExbD photo-cross-links to TonB *in vivo* through residues

that boundary its transmembrane domain suggesting that the ExbD and TonB transmembrane domains

are at some point adjacent (63), a result seemingly at odds with the structural data where the lumen of an

ExbB pentamer cannot accommodate additional transmembrane domains beyond the two of ExbD.

Heterodimerization of TonB-ExbD transmembrane domains would be consistent with Stage II in the

proposed energy transduction cycle (Fig. 2).

While *in vivo* studies of the TonB system have admittedly not always been interpreted correctly

[see (39) subsequently corrected in (94)], it is also worth noting that structural studies have failed to

correctly represent *in vivo* conformations of TonB carboxy terminal homodimers (41) and important

interactions between the TonB carboxy terminus and TBDTs FhuA and BtuB proteins (40). Structural

studies have also missed interactions between the TonB carboxy terminal amphipathic helix and TBDT

FepA (18). Additionally, Cys substitutions that formed ExbD disulfide-linked homodimers cannot be

modeled on the ExbD NMR structure without significant distortion (36). It seems possible that elements

of the model based on *in vivo* data summarized in Fig. 2 could provide important insights that have not

otherwise been gained by structural analysis. Also, considering the parable of the blind men and the

elephant, could there be *in vivo* equilibria among the several proposed ratios of TonB:ExbB:ExbD (11,

25, 80, 95, 96)?

Given the complexity and dynamic nature of the differing protein-protein interactions that occur

in each system, a true understanding of their mechanisms awaits *in vivo* data and *in vitro* structural

models that are significantly more congruent. Currently, much of the *in vivo* data for all three PMF-

dependent systems remains to be rationally incorporated into models arising from structural analyses.

*ExbD L132 is a non-essential residue*. Braun et al. isolated a spontaneous ExbD L132Q mutant

that was inactive, the first to demonstrate the importance of ExbD in all TonB system activities (43). In

this work, the construction and expression of all possible amino acid substitutions at *exbD* codon 132

showed that L132 is not a mechanistically important residue, even though L132Q prevents

formaldehyde crosslinking to TonB and stalls the energy transduction cycle at Stage I [Fig. 2; (42)].

Because other structurally diverse substitutions such as L132C and L132W were fully active, the L132Q

mutation likely caused structural distortion that prevented productive ExbD-TonB interaction.

*ExbD as a target for a novel antibiotic*. The TonB system has been recognized as a potential

target for novel antibiotics for many years because bacteria need sufficient iron to mount infection in

their hosts, which have evolved to sequester it from them (97-103). There appears to be an escalating

battle between bacterial pathogen and host, starting with the deliberately low levels of iron in body

fluids that can be thwarted by the specialized high-affinity iron acquisition systems such as the TonB

system that bacteria use to retrieve it (9).

TonB protein undergoes three different protein-protein interactions during an energy

transduction cycle: TonB-TonB, TonB-ExbD, and TonB-TBDT (Fig. 2). Clues that these could

constitute targets for intervention arise because overexpression of TonB exhibits a dominant negative

gene dosage effect. Proteolytic fragments of TonB that are produced when the level of TonB exceeds

the level of ExbB needed to proteolytically stabilize it can competitively inhibit energy transduction

(104, 105). An exogenously added peptide derived from a conserved consensus sequence known as the

"TonB box" in TBDTs showed promise (106) but was not examined further.

Shotgun and rational design approaches have also been used to search for compounds that could

inhibit the TonB system. High-throughput screening of small molecule libraries have identified

compounds that inhibit TonB system activity (107, 108). However, none of the compounds was shown

to directly target the TonB system and potential modes of inhibition were not identified.

Compounds that inhibit individual TonB-TBDT interactions might not be therapeutically optimal

since TonB interacts differently with the several different TBDTs of *E. coli* K12, which could

potentially diminish or eliminate the inhibition (41, 109). The essential TonB-ExbD interaction

appeared to be a better target since each protein and their interaction are required for energy transduction

through all TBDTs (Fig. 2).

We were expecting to discover that a peptide that is inhibitory when supplied endogenously

[residues 44-63; (63)] would also be inhibitory if supplied exogenously as a cyclic peptide (residues 45-

51), but that was not the case. Possibly it had no effect because it was too short or possibly because an

insufficient amount was able to cross the outer membrane and reach the periplasm. Instead, the cyclic

peptide cyc-IFFRA, derived from ExbD periplasmic domain residues 102-106 in a region that interacted

with the periplasmic domain of TonB prevented the formation of the essential ExbD-TonB complex *in*

*vivo* and inhibited TonB system activity. Because the peptide was cyclic and lacked an OmpT

recognition site, it appeared to have largely survived proteolysis by envelope proteases.  It would be

interesting to survey the effect of this peptide on ESKAPE or other Gram-negative pathogens. The use

of customized cyclic peptides to inhibit the essential ExbD-TonB interaction in Gram-negative bacteria

could serve as a prototype for future antibiotic development.

## MATERIALS AND METHODS

*Strains and plasmids*. Plasmids used in this work are listed in Table 5. Transmembrane domain codons 23-43 of *exbD* were engineered individually to Cys codons by site-directed mutagenesis PCR on plasmid pKP999 which encodes wild type ExbD expressed from a propionate-inducible promoter (29). Open reading frames of all *exbD* mutants were sequenced at the Penn State Genomics Core Facility, University Park, PA to rule out unintended base changes. The various individual Cys- substituted mutants were transformed to strains RA1045 [(*ΔexbD, ΔtolQR*); (54)], RA1017 [(*ΔexbBD, ΔtolQRA*); (110)], and KP1509 [(*ΔexbD, ΔtolQR, ΔtonB*); (29)] as described in the figure legends. All three strains are derivatives of W3110 (111). For studies of ExbD cyclic peptides, strain SH03 (W3110, *tolC*::Tn*10*) and KP1477 [W3110, *ΔtonB* (112)] were used.

*Media and culture conditions*. Luria-Bertani (LB), tryptone (T), and M9 minimal salts media were prepared as described previously (113, 114). Except for plasmid-less strains, all liquid cultures, agar plates, and T-top agar were supplemented with 35 $\mu$g of chloramphenicol/ml and plasmid-specific levels of propionate were added during the assay for the expression of ExbD cysteine substituted proteins at chromosomal levels as determined by immunoblot. M9 salts were supplemented with 1.0% glycerol (wt/vol), 0.4 mg of thiamine/ml, 1 mM $MgSO_4$, 0.5 mM $CaCl_2$, 0.2% Casamino Acids (wt/vol), 40 mg of tryptophan/ml, and 1.85 $\mu$M $FeCl_3$. Cultures were grown at 37°C with continuous aeration.

*Sensitivity to colicins and bacteriophage*. To measure the relative activity of ExbD mutants, strains expressing ExbD mutants at chromosomal levels were grown to mid-logarithmic phase in T-broth and plated in T-top agar on T-agar plates containing 35 $\mu$g of chloramphenicol/ml. The appropriate concentration of propionate inducer required for chromosomal level expression of each mutant was determined by immunoblotting and was present in growth media throughout the assay. 5-fold and 10-

fold dilutions of colicin or bacteriophage, respectively, were spotted on the inoculated T-top layer. The

zones of clearing were recorded after 18 h incubation at 37°C (56, 115).

*Iodine mediated and spontaneous disulfide cross-linking*. Cells grown overnight in LB were subcultured

1:100 into T-broth with various amounts of Na-propionate to achieve chromosomal levels of expression

(from 0.0-1.0 mM and in the case of ExbD V29C, glucose at 0.005% was added to repress expression).

At mid-exponential phase, 0.8 $A_{550}$-ml of cells were pelleted, washed and suspended in 100 $\mu$l of cross-

linking buffer (20mM sodium phosphate, pH 7.4, 150mM NaCl). Then cells were treated with 0.2 mM

iodine ($I_2$) from a 20 mM stock solution in ethanol and kept on ice for 10 min. Unreacted free Cys

residues were blocked with 20 mM *N*-ethylmaleimide (NEM) for 5 min on ice (46). Cells were then

precipitated on ice with equal volume of 20% TCA. Pellets were suspended in SDS gel sample buffer

supplemented with 50mM iodoacetamide and without β-mercaptoethanol (β-ME) (116). To check the

level of protein expression, 0.2 $A_{550}$-ml of mid-exponential phase cells from the experimental samples

were harvested immediately on ice with equal volumes of 20% TCA and solubilized in SDS gel sample

buffer containing β-ME. A kinetic study of dimer formation showed completion of iodine-mediated

disulfide cross-linking within 15 sec of incubation on ice (data not shown). Both iodine-treated non-

reduced and β-ME-treated reduced samples were resolved on 15% SDS-polyacrylamide gels and

immunoblotted with ExbD-specific polyclonal antibody (11). Equal loads were confirmed by Coomassie

staining of the PVDF membrane used for the immunoblot. For spontaneous disulfide cross-linking,

iodine and NEM were omitted.

In vivo *photo-cross-linking*. Plasmid pEVOL encodes an orthogonal tRNA synthetase and a

corresponding orthogonal tRNACUA that recognizes amber (UAG) codons (117). Plasmids with amber

substitutions in *exbD* were engineered, co-expressed with pEVOL in RA1045 (W3110, Δ*exbD*, Δ*tolQR*)

and processed as previously described (17). At $A_{550}$ of ~0.2, sub-cultures in supplemented M9 minimal

media were treated with 0.004% arabinose to induce expression of the tRNA synthetase/tRNA pair from

pEVOL, 0.8 mM of the photo-cross-linkable amino acid p-Benzoyl-L-phenylalanine (pBpa; Bachem

AG, www.bachem.com) and levels of propionate required for ExbD protein expression at near-

chromosomal levels.  2.0 $A_{550}$-ml aliquots were harvested at $A_{550} \sim 0.5$, pelleted at room temperature and

suspended in 2.0 ml supplemented M9 media (as described above with 1.85 μM $FeCl_3$ instead of 37 μM,

and without antibiotics, pBpa, and propionate). Equal volumes of each sample were transferred into

single wells of each of two 24-well plates (NEST Cell Culture Plate, cat: 702001). One plate was

irradiated with 365-nm light for 30 min on ice and the second plate was simultaneously incubated in the

dark for 30 min on ice. Samples were precipitated with an equal volume of 20% TCA as described

previously and processed for immunoblot analysis with ExbD-specific polyclonal antibodies and TonB-

specific monoclonal antibodies (11, 118).  Equal loads were confirmed by Coomassie staining of the

PVDF membrane used for the immunoblot.

*Initial rates of [$^{55}$Fe]-ferrichrome transport*. The initial rates of iron transport were determined as

described previously (56, 115).  Mid-exponential-phase subcultures were pelleted and suspended in iron

transport assay media, and the initial rates of [$^{55}$Fe]-ferrichrome transport were determined. Equal

proportions of the suspended cells were precipitated with an equal volume of 20% TCA to confirm the

chromosomal level expression of ExbD mutants by immunoblot. Precipitates were solubilized in SDS

gel sample buffer (116), electrophoresed on a 15% SDS polyacrylamide gel and immunobotted with

anti-ExbD polyclonal antibody (11).  Inhibitory ExbD-based cyclic peptides were purchased from

GenScript USA, Inc., Piscataway, New Jersey.  Assay of equal numbers of cells in each sample was

confirmed by Coomassie staining of the PVDF membrane used for the immunoblots.

*In-vivo formaldehyde cross-linking*

Cultures were grown overnight to saturation in LB broth at 37°C with aeration. The saturated cultures

were subsequently subcultured 1:100 in supplemented M9 minimal media. At mid-exponential phase,

volumes were adjusted to harvest 0.5 $A_{550}$-ml of cells. Bacteria were suspended in sodium phosphate

buffer (pH 6.8) and treated for 15 min at room temperature with 1% monomeric formaldehyde (Electron

Microscopy Sciences Cat #15710) as previously described (29). Following formaldehyde treatment,

bacteria were pelleted and suspended in SDS gel sample buffer (116), with equal cell numbers loaded on

SDS 13% polyacrylamide gels. ExbD complexes and TonB complexes were detected by

immunoblotting with ExbD-specific polyclonal antibodies and TonB-specific monoclonal antibodies,

respectively (11, 118). Equal loads were confirmed by Coomassie staining of the PVDF membrane used

for the immunoblot.

**Acknowledgements**

We thank Ann Ollis and Christine Dubowy for construction and assay of the Cys relocation mutants. We

thank Yan Shipelskiy for construction and assay of ExbD D25X mutants. We thank Garrick Treaster,

Anne Ollis, and Elise Zhao for construction and characterization of mutations at ExbD residue L132.

We thank Erica Johnson for construction of several amber mutations in *exbD*. We thank William Russ

and Donald Engelman for TOXCAT vectors and *E. coli* NT326 cells, and Kevin Bertrand and Stuart

Huntley for the gift of SH03. We thank Ray Larsen for editorial feedback on the manuscript. Support

from NIAID R21 AI113622, NIGMS R01 GM112710, and Violet S. Postle is gratefully acknowledged.

**REFERENCES**

1.  **Cornelissen CN, Hollander A.** 2011. TonB-Dependent transporters expressed by *Neisseria gonorrhoeae*. Front Microbiol **2:**117.

2.  **Noinaj N, Buchanan SK, Cornelissen CN.** 2012. The transferrin-iron import system from pathogenic *Neisseria* species. Mol Microbiol **86:**246-257.

3.  **Krewulak KD, Vogel HJ.** 2011. TonB or not TonB: is that the question? Biochemistry and cell biology **89:**87-97.

4.  **Kuehl CJ, Crosa JH.** 2010. The TonB energy transduction systems in *Vibrio* species. Future Microbiology **5:**1403-1412.

5.  **Blanvillain S, Meyer D, Boulanger A, Lautier M, Guynet C, Denance N, Vasse J, Lauber E, Arlat M.** 2007. Plant carbohydrate scavenging through TonB-Dependent receptors: A feature shared by phytopathogenic and aquatic bacteria. PLoS ONE **2:**e224.

6.  **Cescau S, Cwerman H, Letoffe S, Delepelaire P, Wandersman C, Biville F.** 2007. Heme acquisition by hemophores. Biometals **20:**603-613.

7.  **Lohmiller S, Hantke K, Patzer SI, Braun V.** 2008. TonB-dependent maltose transport by *Caulobacter crescentus*. Microbiology **154:**1748-1754.

8.  **Schauer K, Rodionov DA, de Reuse H.** 2008. New substrates for TonB-dependent transport: do we only see the 'tip of the iceberg'? Trends Biochem Sci **33:**330-338.

9.  **Sheldon JR, Laakso HA, Heinrichs DE.** 2016. Iron acquisition strategies of bacterial pathogens. Microbiol Spectr **4**.

10. **Kadner RJ, Heller KJ.** 1995. Mutual inhibition of cobalamin and siderophore uptake systems suggests their competition for TonB function. J Bacteriol **177:**4829-4835.

11.    **Higgs PI, Larsen RA, Postle K.** 2002. Quantitation of known components of the *Escherichia*

*coli* TonB-dependent energy transduction system: TonB, ExbB, ExbD, and FepA. Mol Microbiol

**44:**271-281.

12.    **Gresock MG, Kastead KA, Postle K.** 2015. From homodimer to heterodimer and back:

Elucidating the TonB energy transduction cycle. J Bacteriol **197:**3433-3445.

13.    **Larsen RA, Thomas MG, Postle K.** 1999. Protonmotive force, ExbB and ligand-bound FepA

drive conformational changes in TonB. Mol Microbiol **31:**1809-1824.

14.    **Higgs PI, Letain TE, Merriam KK, Burke NS, Park H, Kang C, Postle K.** 2002. TonB

interacts with nonreceptor proteins in the outer membrane of *Escherichia coli*. J Bacteriol

**184:**1640-1648.

15.    **Cadieux N, Kadner RJ.** 1999. Site-directed disulfide bonding reveals an interaction site

between energy-coupling protein TonB and BtuB, the outer membrane cobalamin transporter.

Proc Natl Acad Sc U. S. A. **96:**10673-10678.

16.    **Ogierman M, Braun V.** 2003. Interactions between the outer membrane ferric citrate

transporter FecA and TonB: studies of the FecA TonB box. J Bacteriol **185:**1870-1885.

17.    **Gresock MG, Postle K.** 2017. Going outside the TonB Box: Identification of novel FepA-TonB

interactions *in vivo*. J Bacteriol **199:**e00649-00616.

18.    **Postle K, Kho K, Gresock MG, Ghosh J, Larsen RA.** 2021. Identification of new *in vivo*

TonB-FepA rendevous sites. bioRxiv. doi: https://doi.org/10.1101/2021.12.08.471779

19.    **Plastow GS, Holland IB.** 1979. Identification of an *Escherichia coli* inner membrane

polypeptide specified by a lambda-*tonB* transducing phage. Biochem. Biophys Res Commun

**90:**1007-1014.

20.    **Kampfenkel K, Braun V.** 1992. Membrane topology of the *Escherichia coli* ExbD protein. J

Bacteriol **174:**5485-5487.

21.     **Roof SK, Allard JD, Bertrand KP, Postle K.** 1991. Analysis of *Escherichia coli* TonB

821           membrane topology by use of PhoA fusions. J Bacteriol **173:**5554-5557.

822     22.     **Kampfenkel K, Braun V.** 1993. Topology of the ExbB protein in the cytoplasmic membrane of

823           *Escherichia coli*. J Biol Chem **268:**6050-6057.

824     23.     **Fischer E, Günter K, Braun V.** 1989. Involvement of ExbB and TonB in transport across the

825           outer membrane of *Escherichia coli*: phenotypic complementation of *exb* mutants by

826           overexpressed *tonB* and physical stabilization of TonB by ExbB. J Bacteriol **171:**5127-5134.

827     24.     **Karlsson M, Hannavy K, Higgins CF.** 1993. ExbB acts as a chaperone-like protein to stabilize

828           TonB in the cytoplasm. Mol Microbiol **8:**389-396.

829     25.     **Pramanik A, Zhang F, Schwarz H, Schreiber F, Braun V.** 2010. ExbB protein in the

830           cytoplasmic membrane of *Escherichia coli* forms a stable oligomer. Biochemistry **49:**8721-8728.

831     26.     **Swayne C, Postle K.** 2011. Taking the *Escherichia coli* TonB transmembrane domain

832           "Offline"? Non-protonatable Asn substitutes fully for TonB His20. J Bacteriol **193:**3693-3701.

833     27.     **Baker KR, Postle K.** 2013. Mutations in *Escherichia coli* ExbB transmembrane domains

834           identify scaffolding and signal transduction functions and exclude participation in a proton

835           pathway. J Bacteriol **195:**2898-2911.

836     28.     **Skare JT, Ahmer BMM, Seachord CL, Darveau RP, Postle K.** 1993. Energy transduction

837           between membranes - TonB, a cytoplasmic membrane protein, can be chemically cross-linked *in*

838           *vivo* to the outer membrane receptor FepA. J Biol Chem **268:**16302-16308.

839     29.     **Ollis AA, Manning M, Held KG, Postle K.** 2009. Cytoplasmic membrane protonmotive force

840           energizes periplasmic interactions between ExbD and TonB. Mol Microbiol **73:**466-481.

841     30.     **Pawelek PD, Croteau N, Ng-Thow-Hing C, Khursigara CM, Moiseeva N, Allaire M,**

842           **Coulton JW.** 2006. Structure of TonB in complex with FhuA, *E. coli* outer membrane receptor.

843           Science  **312:**1399-1402.

31.   **Shultis DD, Purdy MD, Banchs CN, Wiener MC.** 2006. Outer membrane active transport:

structure of the BtuB:TonB complex. Science **312:**1396-1399.

32.   **Khursigara CM, De Crescenzo G, Pawelek PD, Coulton JW.** 2004. Enhanced binding of

TonB to a ligand-loaded outer membrane receptor. Role of the oligomeric state of TonB in

formation of a functional FhuA-TonB complex. J Biol Chem **279:**7405-7412.

33.   **Khursigara CM, De Crescenzo G, Pawelek PD, Coulton JW.** 2005. Kinetic analyses reveal

multiple steps in forming TonB-FhuA complexes from *Escherichia coli*. Biochemistry **44:**3441-

3453.

34.   **Freed DM, Lukasik SM, Sikora A, Mokdad A, Cafiso DS.** 2013. Monomeric TonB and the

Ton Box are required for the formation of a high-affinity transporter-TonB complex.

Biochemistry **52:**2638-2648.

35.   **Ollis AA, Kumar A, Postle K.** 2012. The ExbD periplasmic domain contains distinct functional

regions for two stages in TonB energization. J Bacteriol **194:**3069-3077.

36.   **Ollis AA, Postle K.** 2011. The same periplasmic ExbD residues mediate *in vivo* interactions

between ExbD homodimers and ExbD-TonB heterodimers. J Bacteriol **193:**6852-6863.

37.   **Skare JT, Roof SK, Postle K.** 1989. A mutation in the amino terminus of a hybrid TrpC-TonB

protein relieves overproduction lethality and results in cytoplasmic accumulation. J Bacteriol

**171:**4442-4447.

38.   **Larsen RA, Wood GE, Postle K.** 1993. The conserved proline-rich motif is not essential for

energy transduction by *Escherichia coli* TonB protein. Mol Microbiol **10:**943-953.

39.   **Larsen RA, Letain TE, Postle K.** 2003. *In vivo* evidence of TonB shuttling between the

cytoplasmic and outer membrane in *Escherichia coli*. Mol Microbiol **49:**211-218.

40.   **Vakharia-Rao H, Kastead KA, Savenkova MI, Bulathsinghala CM, Postle K.** 2007. Deletion

and substitution analysis of the *Escherichia coli* TonB Q160 region. J Bacteriol **189:**4662-4670.

41.    **Postle K, Kastead KA, Gresock MG, Ghosh J, Swayne CD.** 2010. The TonB dimeric crystal

structures do not exist *in vivo*. mBio **1:**e00307-00310.

42.    **Ollis AA, Postle K.** 2012. ExbD mutants define initial stages in TonB energization. J Mol Biol

**415:**237-247.

43.    **Braun V, Gaisser S, Herrmann C, Kampfenkel K, Killman H, Traub I.** 1996. Energy-

coupled transport across the outer membrane of *Escherichia coli*:  ExbB binds ExbD and TonB

*in vitro*, and leucine 132 in the periplasmic region and aspartate 25 in the transmembrane region

are important for ExbD activity. J Bacteriol **178:**2836-2845.

44.    **Ollis AA, Postle K.** 2012. Identification of functionally important TonB-ExbD periplasmic

domain interactions *in vivo*. J Bacteriol **194:**3078-3087.

45.    **Eick-Helmerich K, Braun V.** 1989. Import of biopolymers into *Escherichia coli*: nucleotide

sequences of the *exbB* and *exbD* genes are homologous to those of the *tolQ* and *tolR* genes,

respectively. J Bacteriol **171:**5117-5126.

46.    **Braun TF, Blair DF.** 2001. Targeted disulfide cross-linking of the MotB protein of *Escherichia*

*coli*: evidence for two H(+) channels in the stator complex. Biochemistry **40:**13051-13059.

47.    **Zhang XY, Goemaere EL, Thome R, Gavioli M, Cascales E, Lloubes R.** 2009. Mapping the

Interactions between *Escherichia coli* Tol Subunits: Rotation of the TolR Transmembrane Helix.

J Biol Chem **284:**4275-4282.

48.    **Cascales E, Lloubes R, Sturgis JN.** 2001. The TolQ-TolR proteins energize TolA and share

homologies with the flagellar motor proteins MotA-MotB. Mol Microbiol **42:**795-807.

49.    **Zhou J, Sharp LL, Tang HL, Lloyd SA, Billings S, Braun TF, Blair DF.** 1998. Function of

protonatable residues in the flagellar motor of *Escherichia coli*: a critical role for Asp 32 of

MotB. J Bacteriol **180:**2729-2735.

50.    **Celia H, Noinaj N, Buchanan SK.** 2020. Structure and stoichiometry of the Ton molecular

motor. Int J Mol Sci **21**.

51.    **Deme JC, Johnson S, Vickery O, Aron A, Monkhouse H, Griffiths T, James RH, Berks BC,**

**Coulton JW, Stansfeld PJ, Lea SM.** 2020. Structures of the stator complex that drives rotation

of the bacterial flagellum. Nat Microbiol **5:**1553-1564.

52.    **Santiveri M, Roa-Eguiara A, Kuhne C, Wadhwa N, Hu H, Berg HC, Erhardt M, Taylor**

**NMI.** 2020. Structure and function of stator units of the bacterial flagellar motor. Cell **183:**244-

257 e216.

53.    **Braun V.** 1989. The structurally related *exbB* and *tolQ* genes are interchangeable in conferring

*tonB*-dependent colicin, bacteriophage, and albomycin sensitivity. J Bacteriol **171:**6387-6390.

54.    **Brinkman KK, Larsen RA.** 2008. Interactions of the energy transducer TonB with noncognate

energy-harvesting complexes. J Bacteriol **190:**421-427.

55.    **Lloubes R, Goemaere E, Zhang X, Cascales E, Duche D.** 2012. Energetics of colicin import

revealed by genetic cross-complementation between the Tol and Ton systems. Biochem Soc

Trans **40:**1480-1485.

56.    **Larsen RA, Chen GJ, Postle K.** 2003. Performance of standard phenotypic assays for TonB

activity, as evaluated by varying the level of functional, wild-type TonB. J Bacteriol **185:**4699-

4706.

57.    **Bernsel A, Viklund H, Hennerdal A, Elofsson A.** 2009. TOPCONS: consensus prediction of

membrane protein topology. Nuc Acids Res **37:**W465-468.

58.    **Russ WP, Engelman DM.** 1999. TOXCAT: a measure of transmembrane helix association in a

biological membrane. Proc Natl Acad Sci U S A **96:**863-868.

59.    **Kopp DR, Postle K.** 2020. The intrinsically disordered region of ExbD is required for signal

transduction. J Bacteriol **202:**e00687-00619.

60.    **Garcia-Herrero A, Peacock RS, Howard SP, Vogel HJ.** 2007. The solution structure of the

periplasmic domain of the TonB system ExbD protein reveals an unexpected structural

homology with siderophore-binding proteins. Mol Microbiol **66:**872-889.

61. **Treptow NA, Shuman HA.** 1985. Genetic evidence for substrate and periplasmic-binding-

protein recognition by the MalF and MalG proteins, cytoplasmic membrane components of the

*Escherichia coli* maltose transport system. J Bacteriol **163:**654-660.

62. **Postle K, Larsen R.** 2004. The TonB, ExbB, and ExbD Proteins, p. 96-112. *In* Crosa JH, Mey

AR, Payne SM (ed.), Iron transport in Bacteria. ASM Press, Washington D.C.

63. **Kopp DR, Postle K.** 2020. An ExbD disordered domain peptide inhibits TonB system activity.

bioRxiv doi: https://doi.org/10.1101/2020.01.05.895219.

64. **Silver LL.** 2011. Challenges of antibacterial discovery. Clin Microbiol Rev **24:**71-109.

65. **Dyson HJ, Wright PE.** 2005. Intrinsically unstructured proteins and their functions. Nat Rev

Mol Cell Biol **6:**197-208.

66. **McCarter JD, Stephens D, Shoemaker K, Rosenberg S, Kirsch JF, Georgiou G.** 2004.

Substrate specificity of the *Escherichia coli* outer membrane protease OmpT. J Bacteriol

**186:**5919-5925.

67. **Chakravorty S, Shipelskiy Y, Kumar A, Majumdar A, Yang T, Nairn BL, Newton SM,**

**Klebba PE.** 2019. Universal fluorescent sensors of high-affinity iron transport, applied to

ESKAPE pathogens. J Biol Chem **294:**4682-4692.

68. **Young L, Jernigan RL, Covell DG.** 1994. A role for surface hydrophobicity in protein-protein

recognition. Protein Sci **3:**717-729.

69. **Bogan AA, Thorn KS.** 1998. Anatomy of hot spots in protein interfaces. J Mol Biol **280:**1-9.

70. **Nikaido H.** 2003. Molecular basis of bacterial outer membrane permeability revisited. Microbiol

Mol Biol Rev **67:**593-656.

71. **Hinchliffe P, Symmons MF, Hughes C, Koronakis V.** 2013. Structure and operation of

bacterial tripartite pumps. Annu Rev Microbiol **67:**221-242.

72. **Ghosh J, Postle K.** 2005. Disulphide trapping of an *in vivo* energy-dependent conformation of

*Escherichia coli* TonB protein. Mol Microbiol **55:**276-288.

73.   **Toews J, Rogalski JC, Clark TJ, Kast J.** 2008. Mass spectrometric identification of

formaldehyde-induced peptide modifications under in vivo protein cross-linking conditions. Anal

Chim Acta **618:**168-183.

74.   **Means GE, Feeney RE.** 1971. Chemical modification of proteins. Holden-Day, Inc., San

Francisco, Calif.

75.   **Klebba PE, Newton SMC, Six DA, Kumar A, Yang T, Nairn BL, Munger C, Chakravorty**

**S.** 2021. Iron acquisition systems of Gram-negative bacterial pathogens define TonB-dependent

pathways to novel antibiotics. Chem Rev **121:**5193-5239.

76.   **Braun V.** 2006. Energy transfer between biological membranes. ACS Chem Biol **1:**352-354.

77.   **Gerding MA, Ogata Y, Pecora ND, Niki H, de Boer PA.** 2007. The trans-envelope Tol-Pal

complex is part of the cell division machinery and required for proper outer-membrane

invagination during cell constriction in *E. coli.* Mol Microbiol **63:**1008-1025.

78.   **Kojima S, Blair DF.** 2001. Conformational change in the stator of the bacterial flagellar motor.

Biochemistry **40:**13041-13050.

79.   **Bulathsinghala CM, Jana B, Baker KR, Postle K.** 2013. ExbB cytoplasmic loop deletions

cause immediate, proton motive force-independent growth arrest. J Bacteriol **195:**4580-4591.

80.   **Jana B, Manning M, Postle K.** 2011. Mutations in the ExbB cytoplasmic carboxy terminus

prevent energy-dependent interaction between the TonB and ExbD periplasmic domains. J

Bacteriol **193:**5649-5657.

81.   **Braun TF, Al-Mawsawi LQ, Kojima S, Blair DF.** 2004. Arrangement of core membrane

segments in the MotA/MotB proton-channel complex of *Escherichia coli*. Biochemistry **43:**35-

45.

82.   **Koebnik R.** 1995. Proposal for a peptidoglycan-associating alpha-helical motif in the C-terminal

regions of some bacterial cell-surface proteins. Mol Microbiol **16:**1269-1270.

967 83. **Roujeinikova A.** 2008. Crystal structure of the cell wall anchor domain of MotB, a stator

968   component of the bacterial flagellar motor: implications for peptidoglycan recognition. Proc Natl

969   Acad Sci U S A **105:**10348-10353.

970 84. **Hosking ER, Vogt C, Bakker EP, Manson MD.** 2006. The *Escherichia coli* MotAB proton

971   channel unplugged. J Mol Biol **364:**921-937.

972 85. **Braun V, Herrmann C.** 1993. Evolutionary relationship of uptake systems for biopolymers in

973   *Escherichia coli*: cross-complementation between the TonB-ExbB-ExbD and the TolA-TolQ-

974   TolR proteins. Mol Microbiol **8:**261-268.

975 86. **Goemaere EL, Devert A, Lloubes R, Cascales E.** 2007. Movements of the TolR C-terminal

976   domain depend on TolQR ionizable key residues and regulate activity of the Tol complex. J Biol

977   Chem **282:**17749-17757.

978 87. **Szczepaniak J, Press C, Kleanthous C.** 2020. The multifarious roles of Tol-Pal in Gram-

979   negative bacteria. FEMS Microbiol Rev **44:**490-506.

980 88. **Wojdyla JA, Cutts E, Kaminska R, Papadakos G, Hopper JTS, Stansfeld PJ, Staunton D,**

981   **Robinson CV, Kleanthous C.** 2015. Structure and function of the *Escherichia coli* Tol-Pal

982   stator protein TolR. J Biol Chem **290:**26675-26687.

983 89. **Celia H, Botos I, Ni X, Fox T, De Val N, Lloubes R, Jiang J, Buchanan SK.** 2019. Cryo-EM

984   structure of the bacterial Ton motor subcomplex ExbB-ExbD provides information on structure

985   and stoichiometry. Commun Biol **2:**358.

986 90. **Hennell James R, Deme JC, Kjr A, Alcock F, Silale A, Lauber F, Johnson S, Berks BC, Lea**

987   **SM.** 2021. Structure and mechanism of the proton-driven motor that powers type 9 secretion and

988   gliding motility. Nat Microbiol **6:**221-233.

989 91. **Ratliff AC, Buchanan SK, Celia H.** 2021. Ton motor complexes. Curr Opin Struct Biol **67:**95-

990   100.

92.    **Guihard G, Boulanger P, Benedetti H, Lloubes R, Besnard M, Letellier L.** 1994. Colicin A

and the Tol proteins involved in its translocation are preferentially located in the contact sites

between the inner and outer membranes of *Escherichia coli* cells. J Biol Chem **269:**5874-5880.

93.    **Nilaweera TD, Nyenhuis DA, Cafiso DS.** 2021. Structural intermediates observed only in intact

*Escherichia coli* indicate a mechanism for TonB-dependent transport. Elife **10:**e68548.

94.    **Gresock MG, Savenkova MI, Larsen RA, Ollis AA, Postle K.** 2011. Death of the TonB

shuttle hypothesis. Front Microbiol **2:**206.

95.    **Pramanik A, Hauf W, Hoffmann J, Cernescu M, Brutschy B, Braun V.** 2011. Oligomeric

structure of ExbB and ExbB-ExbD isolated from *Escherichia coli* as revealed by LILBID-mass

spectrometry. Biochemistry **50:8950-6**

96.    **Sverzhinsky A, Chung JW, Deme JC, Fabre L, Levey KT, Plesa M, Carter DM,**
1002
1003    **Lypaczewski P, Coulton JW.** 2015. Membrane protein complex ExbB4-ExbD1-TonB1 from
1004
1005    *Escherichia coli* demonstrates conformational plasticity. J Bacteriol **197:**1873-85.
1006
97.    **Bullen JJ, Ward CG.** 1988. Iron and infection. Br Med J  **296:**1539.

98.    **Bullen JJ.** 1981. The significance of iron in infection. Rev Infect Dis **3:**1127-1138.

99.    **Bullen JJ, Rogers HJ, Griffiths E.** 1978. Role of iron in bacterial infection. Curr Top Microbiol

Immunol **80:**1-35.

100.    **Parrow NL, Fleming RE, Minnick MF.** 2013. Sequestration and scavenging of iron in

infection. Infection and immunity **81:**3503-3514.

101.    **Drakesmith H, Prentice AM.** 2012. Hepcidin and the iron-infection axis. Science **338:**768-772.

102.    **Bullen JJ, Griffiths E.** 1999. Iron and Infection, 2nd ed. John Wiley & Sons, Chichester, West

Sussex.

103.    **Sparling PF.** 1998. Iron and infection. Clin Infect Dis **27:**1367-1368.

104.    **Mann BJ, Holroyd CD, Bradbeer C, Kadner RJ.** 1986. Reduced activity of TonB-dependent

functions in strains of *Escherichia coli*. FEMS *Lett.* **33:**255-260.

105.    **Howard SP, Herrmann C, Stratilo CW, Braun V.** 2001. *In vivo* synthesis of the periplasmic

1020           domain of TonB inhibits transport through the FecA and FhuA iron siderophore transporters of

1021           *Escherichia coli*. J Bacteriol **183:**5885-5895.

1022    106.    **Tuckman M, Osburne MS.** 1992. *In vivo* inhibition of TonB-dependent processes by a TonB

1023           box consensus pentapeptide. J Bacteriol **174:**320-323.

1024    107.    **Nairn BL, Eliasson OS, Hyder DR, Long NJ, Majumdar A, Chakravorty S, McDonald P,**

1025           **Roy A, Newton SM, Klebba PE.** 2017. Fluorescence high-throughput screening for inhibitors

1026           of TonB action. J Bacteriol **199**(10):e00889-16

1027    108.    **Yep A, McQuade T, Kirchhoff P, Larsen M, Mobley HL.** 2014. Inhibitors of TonB function

1028           identified by a high-throughput screen for inhibitors of iron acquisition in uropathogenic

1029           *Escherichia coli* CFT073. mBio **5:**e01089-01013.

1030    109.    **Ghosh J, Postle K.** 2004. Evidence for dynamic clustering of carboxy-terminal aromatic amino

1031           acids in TonB-dependent energy transduction. Mol Microbiol **51:**203-213.

1032    110.    **Larsen RA, Deckert GE, Kastead KA, Devanathan S, Keller KL, Postle K.** 2007. His20

1033           provides the sole functionally significant side chain in the essential TonB transmembrane

1034           domain. J Bacteriol **189:**2825-2833.

1035    111.    **Hill CW, Harnish BW.** 1981. Inversions between ribosomal RNA genes of *Escherichia coli*.

1036           Proc Natl Acad Sci.USA **78:**7069-7072.

1037    112.    **Devanathan S, Postle K.** 2007. Studies on colicin B translocation: FepA is gated by TonB. Mol

1038           Microbiol **65:**441-453.

1039    113.    **Shedlovsky A, Brenner S.** 1963. A chemical basis for the host-induced modification of T-even

1040           bacteriophages. Proc Natl Acad Sci U S A **50:**300-305.

1041    114.    **Miller JH.** 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory Press, Cold

1042           Spring Harbor, N. Y.

1043    115.    **Postle K.** 2007. TonB system, *in vivo* assays and characterization. Meth Enzymol **422:**245-269.

1044    116.    **Laemmli UK.** 1970. Cleavage of structural proteins during the assembly of the head of

1045            bacteriophage T4. Nature **227:**680-685.

1046    117.    **Young TS, Ahmad I, Yin JA, Schultz PG.** 2010. An enhanced system for unnatural amino acid

1047            mutagenesis in *E. coli.* J Mol Biol **395:**361-374.

1048    118.    **Larsen RA, Myers PS, Skare JT, Seachord CL, Darveau RP, Postle K.** 1996. Identification

1049            of TonB homologs in the family *Enterobacteriaceae* and evidence for conservation of TonB-

1050            dependent energy transduction complexes. J Bacteriol **178:**1363-1373.

1051    119.    **Ahmer BMM, Thomas MG, Larsen RA, Postle K.** 1995. Characterization of the *exbBD*

1052            operon of *Escherichia coli* and the role of ExbB and ExbD in TonB function and stability. J

1053            Bacteriol **177:**4742-4747.

1054    120.    **Buchanan SK, Smith BS, Venkatramani L, Xia D, Esser L, Palnitkar M, Chakraborty R,**

1055            **van der Helm D, Deisenhofer J.** 1999. Crystal structure of the outer membrane active

1056            transporter FepA from *Escherichia coli*. Nat. Struct. Biol. **6:**56-63.

1057    121.    **Yakushi T, Kojima M, Homma M.** 2004. Isolation of Vibrio alginolyticus sodium-driven

1058            flagellar motor complex composed of PomA and PomB solubilized by sucrose monocaprate.

1059            Microbiology **150:**911-920.

1060    122.    **Higgs PI, Myers PS, Postle K.** 1998. Interactions in the TonB-dependent energy transduction

1061            complex: ExbB and ExbD form homomultimers. J Bacteriol **180:**6031-6038.

**TABLES**

**Table 1.  Phenotypes of strains producing Cys-substituted ExbD proteins**

| ExbD | ø80 | colicin B | colicin Ia | colicin M |
|------|-----|-----------|------------|-----------|
| W3110 | 8, 9, 8 | 6, 6, 6 | 5, 5, 5 | 4, 4, 4 |
| pExbD | 8, 8, 8 | 5, 5, 5 | 5, 5, 5 | 4, 4, 4 |
| F23C | 6, 7, 6 | 4, 4, 5 | 5, 5, 4 | 4, 4, 4 |
| I24C | 6, 5, 6 | 4, 4, 4 | 5, 5, 5 | 3, 3, 3 |
| D25C | T, T, T | T, T, T | T, T, T | T, T, T |
| V26C | 7, 7, 7 | 5, 5 ,5 | 5, 5, 5 | 4, 4, 4 |
| M27C | 7, 7, 7 | 5, 5 ,5 | 5, 5, 5 | 4, 4, 4 |
| L28C | 7, 7, 7 | 4, 4, 4 | 5, 5, 6 | 4, 4, 4 |
| V29C | 8, 7, 7 | 5, 5, 5 | 5, 5, 5 | 3, 3, 3 |
| L30C | 6, 6, 6 | 4, 5, 4 | 5, 5, 5 | 3, 3, 4 |
| L31C | 6, 6, 6 | 5, 5, 5 | 5, 5, 5 | 3, 4, 4 |
| I32C | 6, 6, 6 | 5, 6, 5 | 5, 5, 5 | 4, 4, 4 |
| I33C | 7, 7, 7 | 4, 4, 4 | 5, 5, 5 | 3, 3, 3 |
| F34C | 8, 7, 7 | 5, 5, 5 | 4, 4, 4 | 3, 3, 3 |
| M35C | 7, 7, 7 | 4, 4, 5 | 5, 5, 4 | 3, 3, 3 |
| V36C | 8, 7, 7 | 5, 5 ,5 | 4, 5, 4 | 3, 3, 3 |
| A37C | 8, 8, 8 | 5, 5, 5 | 5, 4, 5 | 4, 4, 4 |
| A38C | 7, 7, 7 | 5, 5, 5 | 4, 4, 4 | 3, 3, 4 |
| P39C | 7, 7, 7 | 4, 3 ,4 | 4, 4, 4 | 3, 3, 3 |
| L40C | 8, 8, 8 | 5, 5, 5 | 5, 5, 5 | 4, 4, 4 |
| A41C | 8, 7, 7 | 5, 5, 5 | 4, 5, 4 | 3, 3, 3 |
| T42C | 7, 7 ,7 | 5, 5, 5 | 5, 5, 5 | 3, 3, 3 |
| V43C | 8, 7, 7 | 5, 5, 5 | 5, 4, 5 | 3, 3, 3 |

Parent plasmid pExbD is pKP999.  Phenotypes of ExbD mutants were evaluated in RA1045

(ΔexbD, ΔtolQR) using spot titer assays (56). Cultures with the strains expressing ExbD mutated

proteins at chromosomal levels were plated on T-plates and spotted with fivefold serial dilutions

of colicins and 10-fold serial dilutions of bacteriophage Φ80. Values were recorded as the

reciprocal of the highest dilution at which clearing of the lawn was evident after 18 h of

incubation at 37°C. 'T' indicates tolerance (insensitivity).

**Table 2. ExbD residues 23-63 form homodimers independently of ExbB**

| Plasmid | [a]34 | 62.5 | 125 | 250 | 500 | 1000 |
|---|---|---|---|---|---|---|
| [b]WT/pccKan vector | - | - | - | - | - | - |
| WT/pccGlpA | [c]n.d. | 4+ | 4+ | 4+ | 1+ | - |
| WT/pccGlpA G83I | n.d. | 2+ | +/- | - | - | - |
| WT/pccExbD$_{23-63}$ | n.d. | 4+ | 3+ | 2+ | +/- | - |
| | | | | | | |
| [d]exbBD/pccKan vector | - | - | - | - | - | - |
| exbBD/pccGlpA | 4+ | 4+ | 4+ | 3+ | 1+ | - |
| exbBD/pccGlpA G83I | 3+ | 2+ | 1+ | | | |
| exbBD/pccExbD$_{23-63}$ | 4+ | 4+ | 4+ | 3+ | 1+ | |

[a]plate concentration of chloramphenicol (µg/ml)

[b]strain NT326 is wild-type for *exbB*: [*araD*139, $\Delta$(*argF-lac*)$_{U169}$, $\Delta malE444$, *recA*1, *thi*, *strR* (61)]

[c]growth; n.d. = not determined

[d]strain NT326, made *exbB*::Tn*10* by P1vir transduction (KP1448). The Tn*10* insertion in *exbB* is polar on downstream *exbD* expression (119).

**Table 3.  ExbD T42C and V43C disulfide-linked homodimers are impaired in TonB system-mediated iron transport.**

| | % rate relative to pExbD without DTT | % rate relative to pExbD with DTT |
|---|:---:|:---:|
| **RA1045** | 0.15% | |
| **W3110** | 121% | |
| **pExbD** | 100% | 100% |
| **pExbD L40C** | 101.9% | |
| **pExbD A41C** | 103.6% | |
| **pExbD T42C** | 22.9% | 77% |
| **pExbD V43C** | 9.2% | 89% |

RA1045 is [($\Delta exbD$, $\Delta tolQR$); (54)]; W3110 is wild-type *E. coli* K12 (111); pExbD is pKP999 (29).

Percentiles of transport rates were calculated by slopes derived from linear regression analysis of the individual data points, as described in (72). See Fig. 6 B and C.

# Table 4  ExbD L132 is not an essential residue

| [a]Position 132 a.a. | [b]Active? | [c]Plasmid | [d][propionate] in mM |
|---|---|---|---|
| L | Y | pKP999 | 0.3 |
| A | - | pKP1325 | 0.3 |
| C | Y | pKP1332 | 0.5 |
| Q | - | pKP1333 | 0.5 |
| S | - | pKP1409 | 0.5 |
| V | Y | pKP1449 | 0.2 |
| D | - | pKP1455 | 0.5 |
| T | - | pKP1699 | 0.5 |
| K | - | pKP1698 | 1.0 |
| R | - | pKP1712 | 0.3 |
| H | - | pKP1701 | 0.3 |
| E | - | pKP1702 | 0.5 |
| N | - | pKP1689 | 0.5 |
| G | - | pKP1745 | 0.4 |
| P | - | pKP1703 | 0.3 |
| I | Y | pKP1697 | 0.5 |
| M | Y | pKP1696 | 0.3 |
| F | Y | pKP1713 | 0.3 |
| W | Y | pKP1700 | 0.5 |
| Y | - | pKP1744 | 0.3 |

[a]Single letter code for the amino acid substituted at ExbD residue 132.

[b]Activity of the ExbD variant as determined by iron-dependent growth in RA1045 (*ΔexbD,*

   *ΔtolQR*): Y = activity was same as wild-type *exbD* in pKP999; (-) = inactive, no growth

[c]Plasmid encoding each variant

[d]Concentration of propionate inducer used for chromosomal level of expression (mM)

## Table 5  Plasmids used in this study

| Plasmid | Relevant phenotype | Source or reference |
|---------|--------------------|--------------------|
| pccKan | TOXCAT system vector | (58) |
| pccGlpA | GlpA-TOXCAT fusion | (58) |
| pccGlpAG83I | GlpA G83I-TOXCAT fusion | (58) |
| pKP523 | ExbD$_{23-63}$-TOXCAT fusion | (58) |
| pKP568 | TonB(C18G) | (72) |
| pKP945 | TonB(C18G, A150C) | (29) |
| pKP999 | pPro24 expressing wild-type ExbD | (29) |
| pKP1000 | ExbD(A92C) | (29) |
| pKP1064 | ExbD(D25N) | (29) |
| pKP1394 | ExbD(F23C) | Present study |
| pKP1264 | ExbD(I24C) | Present study |
| pKP1265 | ExbD(D25C) | Present study |
| pKP1284 | ExbD(V26C) | Present study |
| pKP1285 | ExbD(M27C) | Present study |
| pKP1286 | ExbD(L28C) | Present study |
| pKP1395 | ExbD(V29C) | Present study |
| pKP1396 | ExbD(L30C) | Present study |
| pKP1405 | ExbD(L31C) | Present study |
| pKP1397 | ExbD(I32C) | Present study |
| pKP1406 | ExbD(I33C) | Present study |
| pKP1389 | ExbD(F34C) | Present study |
| pKP1386 | ExbD(M35C) | Present study |
| pKP1399 | ExbD(V36C) | Present study |
| pKP1388 | ExbD(A37C) | Present study |
| pKP1390 | ExbD(A38C) | Present study |
| pKP1387 | ExbD(P39C) | Present study |
| pKP1398 | ExbD(L40C) | Present study |
| pKP1391 | ExbD(A41C) | Present study |
| pKP1392 | ExbD(T42C) | Present study |
| pKP1393 | ExbD(V43C) | Present study |
| pKP1412 | ExbD(D25N, L40C) | Present study |
| pKP1413 | ExbD(D25N, A41C) | Present study |
| pKP1445 | ExbD(D25N, T42C) | Present study |
| pKP1406 | ExbD(D25N, V43C) | Present study |
| pKP1191 | ExbD(D25A) | Present study |
| pKP1276 | ExbD(D25A, F23D) | Present study |

| pKP1273 | ExbD(D25A, I24D) | Present study |
| pKP1201 | ExbD(D25A, V26D) | Present study |
| pKP1213 | ExbD(D25A, M27D) | Present study |
| pKP1229 | ExbD(D25A, L28D) | Present study |
| pKP1202 | ExbD(D25A, V29D) | Present study |
| pKP1413 | ExbD(D25A, L30D) | Present study |
| pKP1272 | ExbD(D25A, L31D) | Present study |
| pKP1277 | ExbD(D25A, I32D) | Present study |
| pKP1287 | ExbD(D25A, I33D) | Present study |
| pKP1288 | ExbD(D25A, F34D) | Present study |
| pKP1322 | ExbD(D25A, M35D) | Present study |
| pKP1351 | ExbD(D25A, V36D) | Present study |
| pKP1359 | ExbD(D25A, A37D) | Present study |
| pKP1349 | ExbD(D25A, A38D) | Present study |
| pKP1360 | ExbD(D25A, P39D) | Present study |
| pKP1214 | ExbD(D25A ,L40D) | Present study |
| pKP1215 | ExbD(D25A, A41D) | Present study |
| pKP1199 | ExbD(D25A, T42D) | Present study |
| pKP1197 | ExbD(D25A, V43D) | Present study |
| pKP2234 | ExbD(T21am) | Present study |
| pKP2235 | ExbD(F23am) | Present study |
| pKP2064 | ExbD(I24am) | Present study |
| pKP2236 | ExbD(D25am) | Present study |
| pKP2229 | ExbD(M27am) | Present study |
| pKP2224 | ExbD(V29am) | Present study |
| pKP2237 | ExbD(I32am) | Present study |
| pKP2065 | ExbD(F34am) | Present study |
| pKP2225 | ExbD(M35am) | Present study |
| pKP2226 | ExbD(A37am) | Present study |

# FIGURES

**Figure 1. The TonB system of *Escherichia coli* K12.**  The TonB-dependent transporter,

FepA, is shown in the outer membrane. At its extreme amino terminus, the TonB box, a

site of interaction with TonB, is shown protruding into the periplasm. The topologies of

the cytoplasmic membrane proteins TonB, ExbB and ExbD are shown in the cytoplasmic

membrane. Cellular ratios of the three proteins are indicated in circles in each protein. The

protonmotive force gradient of the cytoplasmic membrane (H+) is shown.  The crystal

structure of FepA was solved by Buchanan et al. (120).

**Figure 2. Model for the TonB energy transduction cycle and the role of ExbD based on *in vivo* data [adapted from (12)].** Full length TonB (purple, upper left corner) and ExbD (blue, upper right corner) each have a single transmembrane domain signal anchor in the cytoplasmic membrane with the bulk of the residues occupying the periplasmic space. The TonB transmembrane domain remains homodimerized throughout the cycle (12). The ExbD transmembrane domain also homodimerizes (as shown in this work), but it is not known if this persists throughout the cycle. Interactions between the periplasmic carboxy-terminal domains of $TonB_2$ and $ExbD_2$ homodimers are shown. Purple circles/triangles are TonB carboxy termini. Blue circles/triangles are ExbD carboxy termini. ExbB tetramers ($ExbB_4$), shown to exist *in vivo*, independently stabilize both TonB and ExbD (23, 80). For simplicity, $ExbB_4$ is not shown in this model. The TonB-

dependent transporter, FepA, is represented by the green square. In Stage I, TonB

periplasmic domains form obligatory homodimers through residues in and near its single

carboxy terminal amphipathic helix (residues 199-216), mediated by H20 in the TonB

TMD, which plays a structural role (12, 26, 41, 72).  ExbD periplasmic domains also

homodimerize (36).  In Stage I there is no evidence that TonB and ExbD homodimers

interact, yet both are proteolytically stabilized by ExbB, suggesting that they might exist

in separate ExbB tetramers. Such a distribution also helps to account for cellular ratios of

14 ExbB: 4ExbD: 2TonB.  In Stage II TonB and ExbD periplasmic domain homodimers

are brought into close contact by movement of the ExbB tetramers such that ExbD

protects TonB in spheroplasts from degradation by exogenously added protease (36, 42).

In Stage III, the previously homodimeric TonBs and ExbD periplasmic domains convert

into a dimer of TonB-ExbD heterodimers in spheroplasts that become sensitive to

degradation by exogenously added protease (36). Energized ExbD correctly configures

TonB for productive interaction with FepA. Stage IV is binding of a monomeric

periplasmic domain of TonB to FepA, such that the siderophore enterochelin is actively

transported across the outer membrane (18).  After a transport event, H20, a key structural

residue in the transmembrane domain, is somehow required for re-establishment of TonB

periplasmic domain homodimers in Stage I (12, 26, 41).  The green circles represent that

all stages of the TonB energy transduction cycle involve the participation of the essential

carboxy terminal amphipathic helix (residues 199-216) in TonB (18). In this model, the

cellular ratios of 14 ExbB: 4ExbD: 2TonB leads to an excess of 2 ExbDs that are not

associated with TonB (11).  Because TonB functions as an obligate homodimer through its

transmembrane domain (12), we speculate that $ExbD_2$ is "de-energized" after Stage IV

(empty blue triangles) and is separately recycled.  In this model, $ExbD_2$ moves in and out

of interactions with TonB$_2$ escorted by ExbB$_4$ tetramers. A separate pool of recycled

ExbB$_4$-ExbD$_2$ is hypothesized to exist to replenish Stage I ExbB$_4$-ExbD$_2$ homodimers. See

(12) for a full explanation of the experimental basis for the model.

**Figure 3. The most meaningful sequence comparisons among ExbD paralogues are based on the conserved Asp residue in the transmembrane domains.**

## A. Alignment based on TOPCONS

## B. Alignment based on conserved Asp residue 25 in ExbD

A slightly expanded number of possible transmembrane domain residues are considered here to encompass the widest range of options. Alignment based on the conserved Asp residue in B. has been previously proposed (48).

Residues conserved between ExbD and TolR are indicated in bold underline in A and B.

Residues conserved between MotB and PomB are in upper case in both A and B.

Residues conserved in all four proteins in B. are indicated by an asterisk *.

Residues which, as Cys substitutions, cross-link strongly with oxidative catalysis ($I_2$ or cu-*o*-phenanthroline) are highlighted in magenta (46, 47). Cys substitutions in PomB have not been systematically analyzed for cross-links with $I_2$ catalysis (121).
Residues not evaluated as Cys substitutions, but that would be interesting to examine in the context of the results from this study, are in gray (46, 47, 121).

**A.**

$\alpha$ ExbD

**B.**

ExbD TMD 23-39

▲ I$_2$ catalyzed homodimers

**Figure 4. Oxidized Cys substitutions in the ExbD transmembrane domain define a helical homodimerization face. A.** ExbD Cys substitutions in RA1045 ($\Delta exbD$, $\Delta tolQR$) were subjected to I$_2$ catalysis as described in Materials and Methods. Divided samples were processed for immunoblotting with anti-ExbD antiserum on a non-reducing SDS

13% polyacrylamide gels. Strains and ExbD variants are indicated at the top of the immunoblot. The position of ExbD monomers and homodimers are indicated on the right with mass markers on the left. WT = W3110; pExbD = pKP999. The corresponding levels of ExbD variants resolved on a reducing SDS 13% polyacrylamide gel are indicated in the lower immunoblot. Equal numbers of cells were loaded in each lane. Two different immunoblots are shown with the partition occurring between ExbD I32C and I33C. **B.** Helical wheel diagram of residues 23-39 of the ExbD transmembrane domain. The most abundant complexes, formed by ExbD I24C, M27C and F34C, are labeled with red triangles and form a face on the helix. Residues 40-43 are not included in the helical wheel analysis since they represent a likely disordered region (see Fig. 5 results).

**Figure 5.  Spontaneous homodimer formation through ExbD L40C, A41C, T42C, and V43C suggests they are separate from the helical transmembrane domain.**

Divided samples of ExbD Cys substitutions in RA1045 (Δ*exbD*, Δ*tolQR*) prepared in the absence of $I_2$ were processed for immunoblotting with anti-ExbD antiserum on non-reducing SDS 13% polyacrylamide gels as described in Materials and Methods (upper). Strains and ExbD variants are indicated at the top of the immunoblot. The position of ExbD monomers and homodimers are indicated on the right with mass markers on the left. WT = W3110; pExbD = pKP999.  Two different immunoblots are shown with the partition occurring between ExbD I32C and I33C. The corresponding levels of ExbD variants resolved on a reducing SDS 13% polyacrylamide gel are indicated in the lower immunoblot.  Equal numbers of cells were loaded in each lane.

**Figure 6: DTT treatment releases ExbD disulfide-trapped through residues T42C and V43C to support TonB-dependent iron transport. A.** Immunoblot of a non-reducing SDS 13% polyacrylamide gel with anti-ExbD antiserum of the samples evaluated for iron transport. Identity of strains and ExbD variants is indicated at the top. Presence (+) or absence (-) of 2 mM DTT during the logarithmic phase of growth is indicated. The lower panel is an immunoblot of a reducing SDS 13% polyacrylamide gel to determine relative steady state levels of the same samples. Positions of ExbD monomer or dimer are indicated on the right. Equal numbers of cells were loaded in each lane. **B.** Initial rates of iron transport for RA1045 ($\Delta exbD$, $\Delta tolQR$) cultures with ExbD plasmids, both with and

without DTT.  Left panel evaluates plasmids expressing ExbD T42C; right panel evaluates

plasmids expressing ExbD V43C.  See Table 3 for calculation and comparison of initial

rates.

**Figure 7. Apparent *in vivo* modification of ExbDT42C occurs most prominently in the absence of ExbB**. Wild-type plasmid-encoded ExbD (pKP999) or ExbDT42C (pKP1392) were grown as described in Materials and Methods and immunoblotted under reducing conditions with anti-ExbD antibodies (upper panel) or anti-ExbB antibodies (lower panel). The positions of normal and modified ExbD and of ExbB are indicated on the right. Strain genotypes are listed at the top of the immunoblot; WT = W3110. RA1045 is (Δ*exbD*, Δ*tolQR*); RA1017 is (Δ*exbBD*, Δ*tolQRA*); KP1509 is (Δ*exbD*, Δ*tolQR*, Δ*tonB*). Lane 1, RA1045; lane 2, W3110; lane 3, RA1045/pKP999; lane 4, RA1017/pKP999; lane 5, KP1509/pKP999; lane 6, RA1045/pKP1392; lane 7, RA1017/pKP1392; lane 8,

KP1509/pKP1392. Equal numbers of cells were loaded in each lane. The immunoblots shown for each antibody are from the same exposure with non-relevant lanes removed (indicated by the white line). Amounts of propionate inducer added were: lane 1, none; lane 2, none, lane 3, 0.25 mM; lane 4, 3.0 mM, lane 5, none; lane 6, 0.018 mM; lane 7, 0.02 mM; lane 8, none.

**Figure 8. ExbB is required for homodimerization of full-length ExbD through its transmembrane domain.** ExbD Cys substitutions in RA1017 (*ΔexbBD, ΔtolQRA*) were subjected to $I_2$ catalysis as described in Materials and Methods. Divided samples were processed for immunoblotting with anti-ExbD antiserum on non-reducing SDS 13% polyacrylamide gels as described in Materials and Methods (upper). Strains and ExbD variants are indicated at the top of the immunoblot. The position of ExbD monomers and homodimers are indicated on the right with mass markers on the left. WT = W3110. Two different immunoblots are shown with the partition occurring between ExbD I33C and F34C. ExbB T42C did not homodimerize, likely because it was chemically modified in the absence of ExbB (see Fig. 7). The corresponding steady state levels of ExbD variants resolved on a reducing SDS 13% polyacrylamide gel are indicated in the lower immunoblot. Equal numbers of cells were loaded in each lane.

A.

B.

ExbD TMD 23-39

▲ I$_2$ catalyzed homodimers   ✻ Photocrosslinked homodimers

**Figure 9.** *In vivo* **photo-cross-linking suggests that the ExbD helices rotate with respect to one another.** **A.** ExbD with pBpa incorporated at various codons in the transmembrane domain was subjected to *in vivo* photo-cross-linking as described in Materials and Methods, resolved on an SDS 13% polyacrylamide gel and immunoblotted with anti-ExbD antisera. Strains and substituted ExbD residues are indicated at the top of the upper immunoblot. Strain labeled ExbD- is RA1045 (ΔexbD, ΔtolQR); pExbD is pKP999; W3110 is wild-type *E. coli*. The position of ExbD monomers, homodimers, and two different previously identified ExbD-ExbB homodimers (59), are indicated on the left with mass markers on the right. Prior to treatment of the samples in A. with ultraviolet light, the level of steady state monomeric ExbD expression and incorporation of pBpA were determined by processing the same volume of sample from A. for immunoblotting with anti-ExbD antiserum on reducing SDS 13% polyacrylamide gels shown in the lower immunoblot. Equal numbers of cells were loaded in each lane. **B.** Helical wheel diagram of residues 23-39 of the ExbD transmembrane domain. The most abundant homodimers formed by *in vivo* photo-cross-linking occurred through ExbD I24pBpa, D25pBpa, V29pBpa, and F34pBpa and are labeled with asterisks. The most abundant disulfide-linked homodimers, formed by ExbD I24C, M27C and F34C, are labeled with red triangles as in Fig. 4. Residues 40-43 are not included in the helical wheel analysis since they represent a likely disordered region (see Fig. 5 results).

**Figure 10: ExbD sequences chosen as cyclic peptide templates are semi-conserved among several ESKAPE pathogens.** Based on the *E. coli* K12 primary amino acid sequence, the ExbD cytoplasmic domain is highlighted in green, the transmembrane domain is highlighted in blue, and the periplasmic domain is highlighted in yellow. The *E. coli K12 exbD* DNA sequence is identified by the gray highlight in the sequence alignment. The selected ExbD regions from which synthetic cyclic peptides were designed based on *E. coli* K12 sequences are boxed in red. The asterisks (*) below the alignment

identify complete residue conservation; hyphens ( : ) identify conserved residue properties; periods ( . ) identify conserved similar residue properties. The orange bar labeled **(1)** above the alignment identifies the ExbD disordered domain in *E. coli* K12—residues 40-43 from this study plus residues 44-63 from (60). The orange bar labeled **(2)** above the alignment identifies the ExbD residues 92-121 in *E. coli* K12 that are necessary for all known ExbD formaldehyde cross-linked complexes (35). The blue star labeled **(3)** above the alignment identifies the L132 residue shown in this work to be relatively unimportant. The accession numbers are *P. aeruginosa* (AAF04084.1); *A. baumannii* (AOX72224.1); *E. cloacae* (ESS57504.1); *K. pneumoniae* (CDO12443.1); *S. dysenteriae* (ABB63085.1); *S. typhimurium* (TKE83773.1); *E. coli K12* (AAA69172.1). Alignment was generated using Clustal-Omega (1.2.4) multiple sequence alignment (http://www.clustal.org/omega).

**Figure 11: Cyclic peptide IFFRA inhibits TonB system activity.** The initial rate of [55]Fe-ferrichrome transport by strain SH03 (W3110, *tolC*::Tn*10*) in the presence of dimethyl sulfoxide (DMSO) or synthetic cyclic peptides in DMSO is shown. Cyclic peptide cyc-VKVNLP is based on ExbD residues 45-50; cyc-VKVNLPA is based on ExbD residues 45-51; cyc-IFFRA is based on ExbD residues 102-106, cyc-KIGLVG is based on ExbD residues 129-134, and cyc-KIGQVG is based on ExbD residues 129-134

with an L-to-Q substitution (see Fig. 10). Strains were grown until mid-exponential phase at which point an $A_{550}$ of 1.4 cells were harvested and suspended in assay medium as previously described (115). DMSO or synthetic peptides (500 µM) suspended in DMSO were added, and samples were incubated at 30˚C while shaking for five minutes prior to measuring the initial $^{55}$Fe-ferrichrome transport rate. The $^{55}$Fe-ferrichrome transport rate was recorded as the percent of the SH03 transport rate. Δ*tonB* refers to W3110 with the *tonB* gene deleted [KP1477; (112)]. The error bars indicate mean ± SEM for two biological replicates, each containing three technical replicates.

**Figure 12: IFFRA inhibits TonB-ExbD and TonB-ExbB complex formation.**

Formaldehyde cross-linking of SH03 (W3110, *tolC*::Tn*10*) in the presence of dimethyl

sulfoxide (DMSO) or synthetic cyclic peptides in DMSO is shown. Cyclic peptide cyc-

VKVNLP is based on ExbD residues 45-50; cyc-VKVNLPA is based on ExbD residues

45-51; cyc-IFFRA is based on ExbD residues 102-106, cyc-KIGLVG is based on ExbD

residues 129-134, and cyc-KIGQVG is based on ExbD residues 129-134 with an L-to-Q

substitution (see Fig. 10). Strains were grown to mid-exponential phase, at which point

cells were harvested and suspended in 100 mM sodium phosphate buffer (pH 6.8). DMSO

or the synthetic cyclic peptides (500 µM in DMSO) were added five minutes prior to

formaldehyde cross-linking as previously described (122). Proteins from equivalent

numbers of cells from the same cultures were analyzed on immunoblots of SDS 13%

polyacrylamide gels with (A) anti-TonB monoclonal antibodies and (B) anti-ExbD polyclonal antibodies. Identity of samples are indicated at the top of both immunoblots. Δ*exbD* refers to KP1045; Δ*tonB* refers to KP1477. (**A.**) Positions of the previously characterized TonB formaldehyde cross-linked complexes are indicated on the right: TonB-FepA, TonB-OmpA, TonB-ExbB, PMF-dependent TonB-ExbD, and the TonB-LPP complex (14, 28, 29). The position of TonB monomer is also indicated. Mass markers are indicated on the left. (**B.**) Positions of the previously characterized ExbD formaldehyde cross-linked complexes are indicated on the right: PMF-dependent ExbD-TonB complex, ExbD-ExbB complex, and ExbD-ExbD homodimer (29). The position of ExbD monomer is also indicated. The position of a potential complex between monomeric ExbD and the cyc-VKVNLPA peptide is indicated by "?". Mass markers are indicated on the left. Red text identifies the position of TonB-ExbB and ExbD-TonB complexes that are inhibited in the presence of cyc-IFFRA. Beneath A and B are shorter exposures of the immunoblots showing relative levels of monomers.