1   **Gliding towards an understanding of the origin of flight in bats**
2

3   **Abstract**
4
5

6   Bats are the only mammals capable of powered flight and have correspondingly

specialized body plans, particularly in their limb morphology. The origin of bat flight is still not

fully understood due to an uninformative fossil record but, from the perspective of a functional

transition, it is widely hypothesized that bats evolved from gliding ancestors. Here, we test

predictions of the gliding-to-flying hypothesis of the origin of bat flight by using phylogenetic

comparative methods to model the evolution of forelimb and hind limb traits on a dataset

spanning four extinct bats and 231 extant mammals with diverse locomotor modes. Our results

reveal that gliders exhibit adaptive trait optima (1) toward relatively elongate forelimbs that are

intermediate between those of bats and non-gliding arborealists, and (2) toward relatively

narrower but not longer hind limbs that are intermediate between those of non-gliders and bats.

We propose a hypothetical adaptive landscape based on limb length and width optimal trends

derived from our modeling analyses. Our results support the potential for glider-like postcranial

morphology to precede a bat-like morphology adapted to powered-flight, setting a foundation for

future developmental, biomechanical, and evolutionary research to test this idea.

Introduction

Powered flight has evolved at least four times over the course of organismal evolution,

consistently serving as a gateway to ecomorphological and species diversification (Heard and

Hauser 1995; Hunter 1998; Jetz et al. 2012; Nicholson et al. 2014). As the only mammals

capable of powered flight, bats have correspondingly specialized body plans that include highly

derived limbs. These specialized morphologies are understood to be the result of adaptations for

the demands of flight; the skeletal elements of bat forelimbs are elongated to support flight

membranes and increase aerodynamic efficiency, whereas bat hind limbs are not only specialized

for hanging and catching prey in flight (Norberg 1990; Gunnell and Simmons 2005; Giannini

2012), but also contribute to flight itself (Cheney et al. 2014).

The evolutionary origin of flight in bats remains one of the biggest mysteries in

mammalian evolution due to an uninformative fossil record linking bats to their non-flying

ancestors. Darwin (1859) posited the origin of bat flight as one of the biggest challenges to the

theory of evolution. Nevertheless, bat flight is widely hypothesized to be the result of an

ancestral "trees-down" transition from non-gliding tree-dweller (hereafter, 'arborealist') to glider

to flyer (Norberg 1985B; Bishop 2008; Simmons et al. 2008), although this idea has not been

tested using phylogenetic comparative methods. It should be noted that all phylogenomic

evidence shows that bats are not closely related to gliders (Upham et al., 2019; Álvarez-Carretero

et al., 2022); bats are nested within Laurasiatheria where neighboring clades are predominantly

adapted to terrestrial locomotion. Laurasiatheria is far from the placental gliders, including

dermopterans and rodents, located in the Eucarchontoglires superorder. The alternative "ground

up" hypothesis (Padian 1982; Caple et al. 1983; Adams and Shaw 2013) holds that flight evolved

from running and jumping vertebrates without a gliding intermediate step; bipedal running is a

key evolutionary step in how birds are thought to have evolved flight (Long et al. 2003;

Kurochkin and Bogdanovich 2010).

Following the arborealist-to-glider-to-flier hypothesis, the ancestors of bats –which

originated about 70-60 million years ago (Teeling et al. 2000; Upham et al. 2019; Álvarez-

Carretero et al. 2022)– are hypothesized to have evolved more elongated limbs, followed by

patagia (wing membranes) on the armwing to facilitate gliding, and finally membranes in the

handwing to enable powered flight. The armwing consists of stylopodal (i.e., humerus) and

zeugopodal (i.e., ulna and radius) bones, while the handwing consists of autopodal bones (i.e.,

the metacarpals and phalanges). The evolution of the handwing is thought to be the key

innovation that catalyzed bats' adaptive radiation (Vaughan 1970; Sears et al. 2006; Amador et

al. 2019), resulting in the evolution of over 1,500 extant species (Simmons and Cirranello

2022B).

Previous work has examined many biological features associated with the evolution of

bat flight, such as sensory adaptations (e.g., echolocation or vision), aerodynamics (e.g., aerofoil

and flight performance), ecology (e.g., diet, habitat, roosting), and development (e.g., patagia

development and digit elongation) (Bishop 2008; Cooper and Tabin 2008; Giannini 2012;

Anthwal et al., 2023; Feigin et al., 2023). Morphological approaches to understanding the origin

of bat flight further include analyses of skeletal dimensions in the context of development

(Adams 1992, 2008; Sears et al. 2006; Nojiri et al. 2023), biomechanics (Swartz et al. 1992;

Swartz 1997; Swartz and Middleton 2008), and functional morphology (Vaughan 1970). These

studies often focus on aspects of the bat forelimb (e.g., Amador et al. 2019) or hind limb (e.g.,

Riskin et al. 2005), but seldom both in tandem in a phylogenetic context (but see Swartz and

Middleton 2008 and Nojiri et al. 2023). Here, we test the gliding-to-flying hypothesis of the

origin of bat flight by modeling the evolution of forelimb and hind limb traits on a dataset

spanning four extinct bats and 231 extant mammals with diverse locomotor modes including

ground-dwellers, arborealists, gliders, and flyers.

Following the arborealist-to-glider-to-bat hypothesis, we test two predictions regarding

the evolution of limb morphologies using an Ornstein-Uhlenbeck evolutionary model-fitting

approach. First, we predict that selection will lead the forelimb skeletal traits of arborealists,

gliders, and bats towards three separate adaptive trait optima/attractors (i.e., θ parameters of

fitted OU models, which are commonly interpreted as the locations of adaptive peaks in

morphospace; hereafter "optima"), where the glider forelimb optimum will be intermediate in

value to those of arborealists and bats. If supported, this demonstrates that glider morphology

may fall on an intermediate adaptive peak between bat and arborealist morphologies.

Specifically, we predict wider and longer skeletal element optima values progressing from

arborealists to gliders to bats. Relatively longer bones (especially in the handwing) and larger-

diameter bones (especially in combination with thinner, highly mineralized walls), may give bats

enough resistance to torsion during powered flight (Swartz et al. 1992) while still increasing their

patagia area. Glider optima with intermediate forelimb length and width optima between

arborealist and bat optima may represent an intermediate adaptive zone between the anatomical

adaptations required for tree-dwelling (Cartmill 1985) and those required for powered flight

(Norberg 1990). Support for this prediction would suggest selective pressures for flight

influenced the evolution of bat forelimb skeletal elements from glider-like forelimbs.

Second, we predict that bat hind limb optima will exhibit a distinct trait optimum (i.e.,

adaptive peak) separate from ground-dwelling, arboreal, and gliding mammals. Previous work

has considered bat hind limbs to be "reduced" in size (e.g., Farnum et al. 2008; Simmons et al.

2008; Louzada et al. 2019), whereas gliders exhibit relatively lengthened hindlimbs (Grossnickle

et al. 2020; Linden et al. 2023). Therefore, the relative lengths and widths of the hindlimb bones

are unclear between these aerial groups as well as in relation to arborealists. Here, we aim to

identify where bat hind limb adaptive optima lie relative to other mammals. Specifically, we

predict narrower and longer skeletal element optima values progressing from arborealists to bats

and gliders. We predict similar but distinct hindlimb optima between gliders and bats because

relatively longer and lower-diameter hind limb bones in bats may decrease weight while still

generating lift by increasing their plagiopatagium surface area, as their hind limbs are

incorporated into this membrane (Giannini 2012). Similarly, relatively longer and wider

hindlimb bones in gliders would increase the surface area for lift during gliding. Support for this

prediction would suggest that shared selective pressures for less-quadrupedal motion (i.e., as

gliding and flying both incorporate hind limbs into the wing membrane) influenced the evolution

of bat hind limb skeletal elements from ancestors with glider-like hind limbs.

Methods

Morphological Data

We compiled linear measurements of 14 forelimb and 15 hind limb traits (Fig. 1) from

231 extant mammals (1–2 specimens per species) across 21 orders (78% of mammalian orders)

and 73 families (47% of mammalian families). Of these, 26 species were bats. Linear

measurements were obtained from specimens held at the Burke Museum of Natural History and

Culture or from previously published work (Chen and Wilson 2015; Grossnickle et al. 2020;

Weaver and Grossnickle 2020; Pevsner et al. 2022). All specimens measured for this study were

fully mature, determined by the closure of exoccipital–basioccipital and basisphenoid–

basioccipital sutures on the cranium and ossification of all vertebrae and limb bones. Specimens

were a mixture of male, female, and unknown sexes. We size-corrected all linear measurements

by calculating log-shape ratios of each trait measurement (Mosimann 1970) by $\log_{10}$-

transforming the mean trait value divided by the cube root of body mass prior to running all

statistical analyses. Unfortunately, most specimens lacked accompanying body mass data;

therefore we compiled species-level body mass data from the PanTHERIA database (Jones et al.

2009) and the *Handbook of the Mammals of the World* book series (Wilson and Mittermeier

2009).

We also collected the 29 measurements for four extinct bats (*Onychonycteris finneyi,*

Icaronycteris index, Hassianycteris messelensis, and Tachypteron franzeni) using ImageJ

(Schneider et al. 2012) on published full-body images that include scale bars (Storch et al. 2002;

Gunnell and Simmons 2005; Simmons et al. 2008). We estimated the body masses of extinct bats

using a linear regression between the body mass and geometric mean of all linear measurements

from the extant bats of our dataset and used those body mass estimations to size-correct our

fossil data again using log-shape ratios. We included fossil bats in our morphological dataset

because, although extinct bats have a high degree of morphological similarity to extant bats

(Norberg 1985B; Bishop 2008; Simmons et al. 2008), the fossil bat data may alter model

estimations of the adaptive zone between bats and gliders in an informative way. For example,

the inclusion of fossils may make the bat-glider trait optima more similar or more disparate by

adding more phylogenetic history information and thus inform the understanding of the adaptive

landscape underlying the evolution of mammalian flight.

To visualize the distribution of locomotor groups in the forelimb and hind limb

morphospaces, we performed principal component analyses (PCA) of the morphometric data

using the *princomp* function in the *R* package *stats* (R Core Team 2021).

Ecological Data

We categorized species into four locomotor groups: ground dwellers ($n = 117$),

arborealists ($n = 71$), gliders ($n = 16$), and flyers ($n = 27$) following previous work (Chen and

Wilson 2015; Grossnickle et al. 2020; Weaver and Grossnickle 2020; Pevsner et al. 2022). The

ground-dwelling mammals included diverse ambulatory, cursorial, saltatorial, and fossorial taxa.

This 'catch-all' category was intentional as we are interested in investigating the arboreal to

gliding to flying specialization trends only. Our arborealist category was also broad in that it

includes both fully arboreal (tree-dwelling) and scansorial taxa. Gliders included tree-dwelling

mammals with derived morphologies (i.e., patagia and elongated limbs) for gliding locomotion

(Grossnickle et al. 2020). Flyers included bats only.

Phylogenetic Comparative Analyses

We used generalized evolutionary models (Hansen 1997; Butler and King 2004) to

examine the influence of locomotor modes on the evolution of forelimb and hind limb

morphologies in our extant-only and extant+extinct datasets.

We first tested the hypothesis that locomotor modes influenced the evolution of the

forelimb and hind limb using multivariate generalized evolutionary models on our extant datasets

(Butler & King, 2004; Hansen, 1997) in the R package mvMORPH (Clavel et al., 2015). To

reduce the dimensions of our data, we conducted a principal component analysis (PCA) on our

14-trait forelimb dataset and 15-trait hind limb dataset. We retained the first three principal

components (PCs) of our forelimb dataset (91.8% of the forelimb variation) and hind limb

dataset (85.3% of hind limb variation). We then fit six multivariate models to the scores of these

three PCs. We fit a multivariate Brownian motion model (mvBM), which models trait variance

in a random walk over time; a single-regime multivariate Ornstein-Uhlenbeck model (mvOU1),

in which scores of each PC evolve towards a single trait optimum (θ parameter of OU models),

giving a single mean optimum for all locomotor groups; and multivariate OU models (mvOUM)

with a given number of optima corresponding to selective regimes (based on locomotor modes).

To test our hypotheses on overall limb evolution, we fit a four-peak OU model (mvOUM$_{loc4}$) for

distinct optima among ground-dweller, arborealist, glider, and flyer regimes. To test alternative

hypotheses, we fit three three-peak models (mvOUM$_{loc3a-c}$) with distinct combinations of the four

locomotor groups. The mvOUM$_{loc3a}$ model tests for distinct optima among ground-dweller,

arborealist, and a combined glider-flyer regimes. The mvOUM$_{loc3b}$ model tests for distinct

optima among ground-dweller, combined arborealist-glider, and flyer regimes. The mvOUM$_{loc3c}$

model tests for distinct optima among a combined ground-dweller-arborealist, glider, and flyer

regimes.

To account for uncertainty in phylogenetic topology and the ancestral character states

(Fig. 1), we used the *make.simmaps* function in the *R* package *phytools* (Revell 2012) to simulate

10 stochastic character maps across 1,000 tree topologies randomly drawn from the posterior

distribution of "completed" trees from Upham et al. (2019), resulting in 10,000 character maps

of locomotory regimes. We randomly sampled 250 trees and fit all six models to these trees. We

assessed relative model support using small sample corrected Akaike weights (AICcW). Models

with ΔAICc scores less than two were considered supported and interpreted as evidence of

locomotor groups influencing species' adaptive zones. We acknowledge that using a subset of

PCs instead of the full dataset could bias our results (D. C. Adams & Collyer, 2018; Uyeda et al.,

2015). However, our results remained consistent when we increased the number of PCs in our

model-fitting analyses to the first four or five PCs. Further, we used simulations to assess

whether we had adequate power to accurately distinguish between complex mvOU models from

Brownian motion (Boettiger et al., 2012).

If the mvOUM$_{loc4}$ model is best-supported, we interpreted this as each locomotor regime

having its own distinct adaptive optimum. This result alone does not provide evidence for the

evolutionary origin of bats. However, if gliders are located at an optimum between arborealists

and bats (i.e., if arborealist < glider < bat optima or arborealist > glider > bat optima), there is

support for an evolutionary "pathway" between the two locomotor modes relative to an ancestral

arborealist or ground-dwelling state (i.e., one could theoretically bridge adaptive peaks via these

pathways). Therefore, if the optima gradually increase/decrease from arborealists to gliders to

bats, our hypothesis that bats evolved from gliding ancestors is supported.

If the mvOUM$_{loc3a}$ model is best-supported, we interpret this as gliders+flyers having a

shared adaptive optimum and arborealists and ground-dwellers having their own distinct

evolutionary optima. This model provides the clearest support for flyers evolving from gliding

ancestors because lineages of both groups are evolving towards the same shared adaptive optima.

Therefore, if this model best fit the data, our hypothesis is supported.

If the mvOUM$_{loc3b}$ model is best-supported, we interpret this as arborealists+gliders

having a shared adaptive optimum and flyers and ground-dwellers having their own distinct

evolutionary optima. There is phylogenetic justification for this grouping because all gliders

evolved from ancestrally arboreal groups (e.g., gliding squirrels evolved from tree squirrels; see

tree in Fig. 1). These groups are similar to the 4-group model in that if arborealists+gliders have

an optimum between those of ground-dwellers and bats, there is support for an evolutionary

"pathway" between the two locomotor modes relative to an ancestral ground-dwelling state.

Therefore, if the optima gradually increase/decrease from ground-dwellers to arborealists+gliders

to bats, our hypothesis that bats evolved from gliding ancestors is supported.

Finally, if mvOUM$_{loc3c}$ model is best-supported, we interpret this as ground-

dwellers+arborealists having a shared adaptive optimum and flyers and gliders having their own

distinct evolutionary optima. As with the mvOUM$_{loc4}$ and mvOUM$_{loc3b}$ models, if gliders have an

optimum between those of ground-dwellers+arborealists and bats, there is support for an

adaptive "pathway" between the two locomotor modes relative to an ancestral arborealist or

ground-dwelling state, providing evidence for our hypothesis that bats evolved from gliding

ancestors.

We also examined the influence of locomotor modes on the evolution of each individual

forelimb and hind limb trait in our extant-only and extant+extinct datasets using univariate

evolutionary models. For each of the 29 traits, we fit a Brownian motion model (BM), a single-

optimum Ornstein-Uhlenbeck model (OU1), and four multiple-peak OU (OUM) models. The

OUM models included OUM$_{loc4}$, OUM$_{loc3a}$, OUM$_{loc3b}$, and OUM$_{loc3c}$, featuring the same

locomotor regimes as the multivariate mvMORPH models. All models were fit using the *R*

package *OUwie* (Beaulieu et al. 2012). We applied a single selection parameter (α) and rate

parameter (σ^2) across all selective regimes while allowing optima (θ) to vary between regimes

(Hansen 1997; Butler and King 2004). For traits in which an OUM model was the best-supported

model, we computed 95% confidence intervals using parametric bootstrapping of the model

parameters with the *OUwie.boot* function in the *R* package *OUwie* (Beaulieu et al. 2012). We

randomly sampled 100 of our character maps and performed 10 bootstrap replicates per

stochastic character map sampled.

We repeated the above multivariate and univariate model fitting approaches on a merged

dataset that includes extant species and the four fossil species. All four models were fit across 50

stochastically mapped trees to account for uncertainty in the ancestral character states. We

manually placed the four extinct bats on the phylogeny as a polytomy with extant bats at 69.8

MYA.

Results

Forelimb morphospace and multivariate model-fitting

PC1 explains 81.5% of the forelimb variation and separates flyers (both extant and

extinct) from the remaining three locomotor modes (Fig. 2a; Table S1a). PC1 also slightly

separates gliders from arborealists and ground-dwellers. This axis describes the relative lengths

of the armwing (humerus, radius, ulna) and the handwing (metacarpals and phalanges). PC1

varies across relatively shorter (-PC1) to elongate (+PC1) bones, revealing a trend within the

dataset in the elongation of multiple forelimb bone elements from arborealists to gliders to flyers.

PC2 explains 7.8% of the morphological variation and primarily describes the relative widths of

the long bones and hands. This variation of wide to narrow bones mostly occurs within

locomotor groups, especially within the diverse ground-dweller group. PC3 explains 2.9% of the

morphological variation with +PC3 dominated by metacarpal length, ulna olecranon length, and

scapula height and -PC3 dominated by intermediate phalanx length and humerus distal width.

Fossil bats overlap with extant bats in morphospace, which is consistent with the lack of

transitional forms in the bat fossil record.

The mvOUM$_{loc3b}$ model, which assigns arborealists and gliders to the same regime, was

the best-fitting model for the extant multivariate forelimb dataset (PCs 1-3) (AICcW $= 0.99$,

Table 1a). Distinct optima were supported for these three locomotor groups, with PC1 and PC2

optima scores increasing in value from ground-dweller to arborealist+glider to flyer hind limbs

(Fig. 2c). Repeating the multivariate model-fitting with the addition of the four extinct bat

species yielded a mvOUM$_{loc4}$ model as the best fit (mAICcW = 0.99; Table S6a). However, the

optima exhibited unrealistic magnitudes (Table S6b).

Forelimb univariate model-fitting

The OUM$_{loc4}$ model was the best-fitting model for six of the 14 forelimb traits

(AICcW = 0.51-1.00), OUM$_{loc3b}$ was the best-fitting model for five traits (AICcW = 0.36-0.65),

OUM$_{loc3c}$ was the best fitting model for metacarpal length (AICcW = 0.63), and OUM$_{loc3a}$ was

the best-fitting model for ulna olecranon length (AICcW = 0.58; Table S2a). The OU1 model

was the best fit for the proximal phalanx width of the third manual digit (AICcW = 0.51).

Armwing (i.e., humerus and radius) length traits have optima (from the best-fitting

OUM$_{loc4}$ models) that gradually increase from ground-dwellers to arborealists to gliders to flyers;

parametric bootstrapping revealed little to no overlap in the 95% confidence intervals of optima

between most locomotor groups (Fig. 3b-c). Optima of armwing widths (from the best-fitting

OUM$_{loc3b}$ models) indicated that bats exhibit relatively wider humeri at the distal end, proximal

end, and mid-shaft compared to all other locomotor groups (Fig. 3g-i; Table S3a).

Bats exhibit relatively higher optima for handwing length traits (i.e., longer third manual

metacarpal, proximal phalanx, and intermediate phalanx) than all other locomotor groups (Fig.

3d-f; Table S3a). For each, glider optima were intermediate between ground-dweller and

arborealist optima. For handwing width traits, bats exhibit relatively higher OUM$_{loc3b}$ optima for

metacarpal widths compared to all other locomotor groups (Fig. 3j), but no differences in

proximal phalanx width as evidenced by OU1 support for this trait.

An adaptive optimum for a bat-glider combined group in the OUM$_{loc3a}$ models was

supported for ulna olecranon length only (Table S2a). In total, however, glider optima (in

combination with arborealists) were between the 95% confidence intervals of ground-dweller

and bat optima for all traits except scapula length/width, humerus proximal width, and

metacarpal width (Fig. 3a, 3g, 3j; Table S3a). For these last trait optima, glider+arborealist

optima were statistically the same as that of ground-dwellers while bats exhibited higher optima

than all other locomotor groups (Fig. 3a, 3g, 3j; Table S3a).

Adding fossil bats into forelimb model-fitting analyses yielded overall slightly more

support for OUM_{loc3a} models compared to extant-only modeling; scapula height and humerus

proximal width demonstrate support for a bat-glider combined group (Table S4a). OUM_{loc4} and

OUM_{loc3b} models were still widely supported as the best-fitting model for six traits and five

traits, respectively; OU1 was still the best-fitting model for proximal phalanx width (Table S4a).

Overall, the length and width optima values for the forelimb were very similar between

extant+extinct and extant only-modeling (Table S5a); however, we were unable to examine the

trends in finer detail because forelimb bootstrapping failed with the addition of fossil species.

Hind limb morphospace and multivariate model-fitting

PC1 explains 47.5% of hind limb variation and more positive PC1 scores describe the

relative elongation (+PC1) of the fibula, metatarsal, and pedal proximal and intermediate

phalanges (Fig. 2b; Table S1b). PC1 slightly separates gliders and bats, with gliders falling

towards +PC1 and bats falling towards -PC1, indicating glider hind limb autopodia are overall

more elongated than bats'. Arborealists and ground-dwellers are spread across PC1. PC2

explains 17.8% of the variation, where +PC2 is associated with relatively lengthened pedal

proximal phalanges, intermediate phalanges, and ischium and -PC2 with relatively narrower

pedal proximal and intermediate phalanges. PC2 loosely separated ground-dwellers, arborealists,

and the glider-flyer group. PC3 explains 11.3% of the morphological variation with +PC3

dominated by ilium length, femur length, and fibula length and -PC3 dominated by metatarsal

width, intermediate phalanx width, and proximal phalanx width. These loadings show that

variation between the four locomotor groups mainly occurs throughout the hind limb autopodia

region.

305   The mvOUM$_{loc4}$ model was the best-fitting model for the extant multivariate hind limb

dataset (PCs 1-3) (AICcW = 0.92, Table 1b). Distinct optima were supported for the four

locomotor regimes, with PC1 and PC2 optima scores decreasing in value from ground-dweller to

arborealist to glider to flyer hind limbs, albeit with overlap (Fig. 2d). Adding in the four extinct

bat species yielded again the mvOUM$_{loc4}$ model as the best fit, and very little change in the

optima values (Table S6).

Hind limb univariate model-fitting

313   The OUM$_{loc4}$ model was the best-fitting model for three of 15 traits, all of which were

hind limb long bone lengths (AICcW = 0.33-0.60; Table S2b). Specifically, gliders have

relatively longer tibia and femoral optima compared to bats and the other locomotor groups (Fig.

4b-c, Table S3b). Fibula length was the third hind limb trait supported by OUM$_{loc4}$, but fibulae

are rudimentary in bats. The OUM$_{loc3a}$ model best fit three traits (AICcW = 0.36-0.52), all of

which were widths: femur shaft width, metatarsal width, and pedal proximal phalanx width

(Table S3b). Parametric bootstrapping of these OUM$_{loc3a}$-supported models revealed that the

glider-bat group had relatively narrower femur shafts, metatarsals, and pedal proximal phalanges

optima than the other locomotor groups (Fig. 4f, i, j; Table S3b). The OUM$_{loc3b}$ model was the

best-fitting model for four of the 15 traits (AICcW = 0.44-0.71): ilium, ischium, metatarsal, and

intermediate proximal phalanx length (Table S2b). Bats had significantly smaller relative ilium

and ischium length optima than other locomotor groups. Gliders+arborealists had metatarsal and

intermediate proximal phalanx optima intermediate between bats and ground-dwellers (Fig. 4d-e;

Table S3b). The OUM_{loc3c} model best fit three traits (AICcW = 0.42-0.48): pelvis length, femur

distal width, and tibia proximal width (Table S2b). In the latter two traits, gliders exhibited

intermediate width optima between bats (which were smaller) and ground-dwellers+arborealists

(where were larger). For pelvis length, like ilium and ischium length, there was no clear trend in

the direction of the optima values across locomotor groups (Table S3b).

Adding fossil bats into hind limb model-fitting analyses yielded only one difference

compared to extant-only modeling: pelvis length was now best fit with a OUM_{loc3b} instead of

OUM_{loc3c} model (Table S4b). The length and width optima for the hind limb were very similar

between extant+extinct and extant-only modeling (Table S5b). However, we were unable to

examine the extent of this consistency because hind limb bootstrapping failed with the addition

of fossil species.

Discussion

Forelimb trait evolution

We found evidence that mammalian glider forelimb morphologies are evolving toward

adaptive peaks that are intermediate between those of arborealists and bats. Our multivariate

forelimb results and univariate length trait results provide support for the hypothesis that

selective pressures for flight influenced the evolution of bat forelimbs from glider-like forelimbs.

Using multivariate modeling, we found a three optima model with distinct selective regimes for

ground-dwellers, arborealists+gliders, and bats as the best fit to the first three PCs of our data

(Table 1a). The arborealist+glider optima are intermediate between those of bats and ground-

dwellers (Fig. 2c). PC1 is mostly explained by the variation in the forelimb bone lengths and

PC2 by their widths, creating a coarse picture that forelimb bones likely become relatively longer

and possibly wider from ground-dwellers to arborealists+gliders to bats. Our univariate model

results are consistent with these findings, revealing that glider forelimb optima are intermediate

between bat and arborealist optima for length traits (Fig. 3a-f), but only partially for width traits

(Fig. 3g-j) across both the armwing and handwing (See Table S7 for a breakdown of whether

each univariate model supports our hypothesis).

Bats exhibit trait optima indicating evolution of relatively longer forelimb bones

compared to all other locomotor groups (Fig. 2a, Fig. 3a-f) particularly in the handwing, which

exhibited rapid evolution as a necessary adaptation for bats (Vaughan 1970; Sears et al. 2006;

Amador et al. 2019). Increased bone lengths may enhance the surface area of the patagium for

greater lift (Norberg 1985b). Bats also displayed evidence of evolution toward relatively wider

bones in the armwing compared to all other locomotor groups (Fig. 3g-j) consistent with findings

that bat humeri and radii exhibit significantly larger diameters compared to nonvolant mammals

(Swartz and Middleton 2008). These relatively wider proximal forelimb bones combined with

thinner yet highly mineralized walls provide resistance to bending and torsion during flapping

(Swartz et al. 1992; Papadimitrious et al. 1996). Furthermore, the transition from relatively more

robust proximal ends of the humerus to relatively less robust distal ends may also reflect greater

muscle mass closer to the center of rotation of the wing joints, which would greatly decrease

flight inertial power and facilitate economical flight (Norberg 1985A; Swartz 1997). In the

handwing, we found bats exhibit relatively thicker third metacarpals (Fig. 3j) that corresponded

with its relatively increased length (Fig. 3d) compared to nonvolant mammals. In contrast, the

proximal phalanx length of the third manual digit was relatively longer compared to nonvolant

mammals whereas its width was best supported with an OU1 model, consistent with previous

work showing that the proximal phalanx diameter is not significantly smaller in bats compared to

nonvolant mammals (Swartz and Middleton 2008). As a result, bats exhibit relatively thick bone

cortices to combat greater bending moments associated with this relatively increased elongation

(Swartz et al. 1992; Papadimitriou et al. 1996; Swartz 1997). Increased cortical thickness along

with the reduction of mineralization of the handwing work together to resist bending failure

while simultaneously maintaining flexibility during flying (Papadimitriou et al. 1996).

In gliders, we found evidence that forelimb bone length optima are largely intermediate

between ground-dwellers and bats. Gliders tend to exhibit relatively greater length optima in

most forelimb bones compared to ground-dwelling and arboreal mammals (Thorington and

Heaney 1981; Grossnickle et al. 2020; Linden et al. 2023) but do not reach the lengths of bat

forelimb bones. Nevertheless, like bats, the longer bones expand the surface area of the patagium

(Thorington and Heaney 1981) and enable further adjustment of their patagium during flight,

leading to more efficient aerodynamics, increased agility, and maneuverability (Thorington et al.

1998; Zhao et al. 2019). Forelimb bone widths in gliders, while clearly intermediate between

those of ground-dwellers and bats for some traits (Fig. 3h, i) share an optimum with arborealists

(shown by the OUM_{loc3b} support for these traits; Fig. 3g-j) and overlap with ground-dwelling

optima mammals for other traits (Fig. 3g, j). Together, these results suggest gliders may maintain

relative widths while continuing to elongate forelimb bones, supporting previous findings that

the evolutionary elongation of glider forelimbs is not also accompanied by slimming of the bones

(Grossnickle et al. 2020).

 Hind limb trait evolution

We found mixed evidence that mammal glider hind limb morphologies are intermediate

to those of arborealists and bats. Using our multivariate modeling approach, we found distinct

optima for ground-dwellers, arborealists, gliders, and bats that best fit the first three PCs of our

hind limb data (Table 1b). The trait loadings indicate that the autopodial bones dictate the

distribution of PC scores in PCs 1 and 2 (Table S1b). However, there is much overlap among the

optima of ground-dwellers, arborealists, and gliders in morphospace (Fig. 2d).

The univariate models uncovered further patterns: 1) width optima of hind limb bones

tend to be narrower in bats and gliders compared to arboreal mammals (Fig. 4f-j), and 2) there

was no pattern from arboreal to glider to bat in the hind limb length optima. Univariate modeling

revealed that the widths of hind limb bones tend not to differ between bats and gliders (Fig. 4f-j)

because the widths of hind limb bones in bats and gliders exhibited either a single optimum (i.e.,

OUM_{loc3a}) separate from ground-dwelling and arboreal mammals (Fig. 4 f, i, j) or an overlap in

confidence intervals of optima between bats and gliders (Fig. 4 g, h). This shared bat-glider

adaptive zone suggests that bats and gliders exhibit narrower hind limb optima, especially in the

autopod region (Fig. 4f-j). In contrast, univariate modeling revealed distinct optima in long bone

lengths of the hindlimb among the four locomotor regimes, in which gliders exhibited relatively

longer femoral and tibial optimal lengths compared to bats, arborealists, and ground-dwellers

(Fig. 4b-c). Unlike bat forelimbs, bat long bones in the hindlimb (i.e., the femur and tibia) are not

the longest compared to the other locomotor regimes; in fact, the confidence intervals of optimal

lengths largely overlap between bats and arborealists (Table S3b). Furthermore, bats exhibit

relatively shorter metatarsals but relatively longer pedal intermediate phalanges compared to all

other locomotor regimes, whereas these autopodial lengths do not differ between gliders and

arborealists (Fig. 4d-e). The different evolutionary patterns of hindlimb bones demonstrate the

unique locomotor diversity among the four regimes. Although gliders and bats are both aerial,

these behaviors require different functional morphologies. In gliders, relatively longer hindlimbs

are needed to increase the surface area of the patagium in tandem with relatively longer

forelimbs to generate lift during gliding (Thorington and Heaney 1981). In contrast, in bats, the

relative lengthening of the forelimb supports the majority of the wing membranes whereas the

hindlimb, specifically the ankle, serves as an anchor to the wing membrane. Bat hindlimbs are

still used for flight by helping modulate lift, drag, or pitch (Cheney et al. 2014).

Overall, gliders appear to be evolving relatively lengthened hindlimb bones while bats

are evolving more gracile hindlimb bones. This finding, while in opposition to our predictions,

may point to the fact that modern gliders are completely distinct evolutionarily and potentially

even morphologically from the gliding ancestors of bats. These results may highlight the pitfalls

of modeling modern gliders as proxies for ancestral ones instead of invalidating our hypothesis

that bats evolved from ancestral gliders.

Incorporating fossil bats

The morphospace positions of fossil bats in the forelimb PCA confirm extinct bats'

morphological similarity to extant bats in the degree of forelimb specialization (Fig. 2a). Amador

et al. (2019) modeled aerodynamic traits instead of skeletal traits and found that the Eocene bat

Onchonycteris finneyi occupies an area of aerofoil morphospace between extant gliders and bats.

In our morphospace, *O. finneyi* is the closest fossil bat to gliders across PC1 (Fig. S1a),

congruent with Amador et al.'s (2019) findings.

Fossil bats occupy the same region of hind limb morphospace as extant bats, but three of

four species overlap clearly with the morphospace of non-bats, a higher proportion compared to

that of extant bats (Fig. 2b). *Tachypteron franzeni* most clearly overlaps with extant bats and is

also the only fossil we included from an extant bat family, Emballonuridae (Storch et al. 2002).

Simmons et al. (2008) suggested that, based on its intermediate limb traits, *Onychonycteris*

finneyi was likely more proficient at climbing than extant bats and may have employed

quadrupedal locomotion and under-branch hanging. Interestingly, *O. finneyi* and *Desmodus*

rotundus (the common vampire bat) are almost indistinguishable in the hind limb morphospace

(Fig. S1b). Vampire bats are known for their adept quadrupedal locomotion and climbing (e.g.,

Riskin et al. 2005), so *O. finneyi'*s similar morphology may lend support to more walking-

capable early bats. However, the linear measurements we used here may not reflect the complex

adaptations required for these behaviors (e.g., ninety-degree joint rotations for under-branch

hanging, Riskin et al. 2005), so the placement of these species in morphospace leaves many

possibilities open for ancestral locomotor behaviors.

Including fossil bats in our model-fitting analyses as the closest morphologies we have to

bat-glider intermediaries did not significantly alter estimations of the adaptive zone between bats

and gliders; the inclusion of fossils appeared to make the bat-glider trait optima marginally more

similar for the forelimbs and had no change for the hind limbs (Table S7). Fossil addition to the

multivariate model-fitting analysis shifted support from mvOUM$_{loc3b}$ to mvOUM$_{loc4}$, suggesting

that the glider optimum is potentially distancing itself from arborealists' and towards bats'

(Table S6a). Unfortunately, we can only speculate about this result because the optima resist

comparison to extant-only mvOUM$_{loc3b}$ optima due to their extremely negative values, which

represent unrealistic morphologies (Table S6b). Fossil addition to univariate forelimb model-

fitting analyses yielded overall slightly more support for $\text{OUM}_{\text{loc3a}}$ models compared to extant-

only modeling (Table S4a). Fossil addition to hind limb model-fitting analyses yielded overall no

change in support for the $\text{mvOUM}_{\text{loc4}}$ model or $\text{OUM}_{\text{loc3a}}$ models compared to extant-only

modeling (Table S6, S4b). Regardless, overall the length and width trends are the same as above

between locomotor groups across the fore- and hind limb, though we lack bootstrap support to

gain more insight into whether any adaptive optima confidence intervals overlap (Table S5).

Limb trait optima and the adaptive landscape

Our results provide insight on the potential adaptive landscape connecting arboreal,

gliding, and flying locomotion, which is represented in Figure 5 with the z-axis describing the

qualitative summation of optima from the univariate model results. Overall, we found that glider

forelimb optima fall intermediate between those of arborealists and bats in their lengths but only

partially in their widths. Except for long bone lengths, glider hind limb optima are intermediate

between ground-dweller/arborealist and bat optima or even overlap with bat optima in their

lengths and widths. Together, these results emphasize that selection on key length/width traits

may be "pulling" glider traits towards a flying adaptive zone yet that certain glider traits (e.g.,

femur and tibia lengths) are derived specifically for gliding morphology. Bishop (2008) proposed

that the high degree of convergence among gliders suggests functional constraints on the

evolution of powered flight from gliding; however, more recent work found that gliding lineages

display incomplete convergence (species retain unique morphologies even while evolving some

similarities; Leal et al. 2002; Stayton 2006). For gliders, there is little evidence that gliding is

under strong selective pressures toward a single-optimum gliding morphology (Grossnickle et al.

2020). The clustering of bat species in forelimb morphospace (Fig. 2a) may represent a narrow

adaptive peak within which forelimb morphologies are highly conserved. In contrast, the broad

glider morphospace may represent a gradually inclined selection plane, with gliders still far from

a peak (Grossnickle et al. 2020). This broad adaptive slope could represent the foothill of a steep

bat peak such that gliders are very slowly evolving towards a bat peak (Fig. 5a). The

evolutionarily oldest extant glider lineages, dermopterans (colugos) and anomalurids (scaly-

tailed squirrels), have evolved farthest from ancestral arborealists (Grossnickle et al. 2020;

although some extinct gliders may have experienced relatively greater change from ancestral

arborealists, e.g., Meng et al. 2017; Luo et al. 2017). Here, the dermopteran species

Cynocephalus volans and *Galeopteras variegatus* are the gliders closest to the bat forelimb

morphospace (Fig. 3a; Fig. S1a), suggesting a trajectory in morphospace from gliding to flying.

Further, colugos are known as the "mitten gliders" because of their interdigital membranes,

which, along with their common carpal morphology with bats (Simmons and Geisler 1998),

reflects hypotheses that interdigital webbing was key to evolving powered flight (Anderson and

Ruxton 2020 but see Gardner and Dececchi 2022 for an alternate view).

Our results also suggest that both bats and gliders exhibit broad adaptive optima in the

hind limb due to gliders sharing an optimum or overlapping optima with bats, or being

intermediate between non-bats and bats for many traits (Fig. 2d, Fig. 4). These broad adaptive

optima could represent overlapping peaks for gliders and bats separated only by a shallow fitness

valley driven by the gliders' elongated femur and tibia lengths (Fig. 5b). The hypothetical

adaptive landscape is therefore valuable because the transition from quadrupedalism to hind limb

suspension has been hypothesized to enable forelimbs to evolve more freely since they would no

longer be weight-bearing structures (Bishop 2008). Indeed, there is both quadrupedal suspension

and hind limb suspension in dermopterans, resulting in a tradeoff in the ability to walk

quadrupedally upright (Fujiwara et al. 2011; Granatosky 2018). In the PC1 and PC2 hind limb

morphospace, *G. variegatus* is the only glider that overlaps with the bat morphologies (Fig. 2b;

Fig. S1b), reflecting a hind limb adaptive optimum potentially more easily attained than that of

bats' specialized forelimbs (Fig. 5).

Our modeling approach offers a first step towards support for the hypothesis that

selection may more strongly "pull" certain limb traits towards a bat adaptive zone. In particular,

gliders' intermediate forelimb length and overlapping hind limb length/width optima between

that of arborealists and bats could be the traits driving lineages across fitness valleys due to their

greater effect in, for example, the functionality of the gliding/flying apparatus. However, to

complete the tentative adaptive landscape presented here (Fig. 5), these results could be

strengthened by more comprehensive coverage of the lengths/widths across more mammal

species and biomechanical studies tying these traits to specific functions and selective pressures.

Conclusions

Our findings complement earlier work investigating the evolution of bat powered flight

by testing the selective regimes acting on bone dimensions across the mammalian forelimb and

hind limb. Our results reveal that the relatively lengthened forelimbs in gliders are intermediate

to those of bats and arborealists, whereas bats and gliders exhibit similarly relatively narrower

hind limbs compared to arborealists. Thus, solely based on morphometrics, the traditional highly

specialized view of glider/bat forelimbs and the "reduced" view of bat hind limbs may represent

more malleable adaptive zones than previously thought. There is a need for future studies to test

the biomechanical consequences of such trait optima alongside more synergistic studies taking

into account bat and gliders' complex physiology and development to test these hypotheses

regarding the evolution of mammalian powered flight.

530
531 **References**
532

Adams, D. C., and M. L. Collyer. 2018. Multivariate phylogenetic comparative methods:

evaluations, comparisons, and recommendations. Syst. Biol. 67:14–31.

Adams, R. A. 1992. Comparative skeletogenesis of the forearm of the little brown bat (*Myotis*

lucifugus) and the norway rat (*Rattus norvegicus*). J. Morphol. 214:251–260.

Adams, R. A. 2008. Morphogenesis in bat wings: linking development, evolution and ecology.

Cells Tissues Organs 187:13–23.

Adams, R. A., and J. B. Shaw. 2013. Time's arrow in the evolutionary development of bat flight.

Pp. 21–46 *in* R. A. Adams and S. C. Pedersen, eds. Bat Evolution, Ecology, and

Conservation. Springer New York, New York, NY.

Álvarez-Carretero, S., A. U. Tamuri, M. Battini, F. F. Nascimento, E. Carlisle, R. J. Asher, Z.

Yang, P. C. J. Donoghue, and M. dos Reis. 2022. A species-level timeline of mammal

evolution integrating phylogenomic data. Nature 602:263–267.

Amador, L. I., N. B. Simmons, and N. P. Giannini. 2019. Aerodynamic reconstruction of the

primitive fossil bat *Onychonycteris finneyi* (Mammalia: Chiroptera). Biol. Lett.

15:20180857.

Anderson, S. C., and G. D. Ruxton. 2020. The evolution of flight in bats: a novel hypothesis.

Mammal Rev. 50:426–439.

Anthwal, N., D. J. Urban, A. Sadier, R. Takenaka, S. Spiro, N. Simmons, R. R. Behringer, C. J.

Cretekos, J. J. Rasweiler, and K. E. Sears. 2023. Insights into the formation and

diversification of a novel chiropteran wing membrane from embryonic development.

BMC Biology 21:101.

Beaulieu, J. M., D.-C. Jhwueng, C. Boettiger, and B. C. O'Meara. 2012. Modeling stabilizing

selection: Expanding the Ornstein-Uhlenbeck model of adaptive evolution. Evolution

66:2369–2383.

Bishop, K. L. 2008. The evolution of flight in bats: narrowing the field of plausible hypotheses.

Q. Rev. Biol. 83:153–169.

Boettiger, C., G. Coop, and P. Ralph. 2012. Is your phylogeny informative? Measuring the

power of comparative methods. Evolution 66:2240–2251.

Butler, M. A., and A. A. King. 2004. Phylogenetic comparative analysis: a modeling approach

for adaptive evolution. Am. Nat. 164:683–695.

Caple, G., R. P. Balda, and W. R. Willis. 1983. The physics of leaping animals and the evolution

of preflight. Am. Nat. 121:455–476.

Cartmill, M. 1985. Chapter 5. Climbing. Pp. 73–88 *in* M. Hildebrand, D. M. Bramble, K. F.

Liem, and D. B. Wake, eds. Functional Vertebrate Morphology. Harvard University

Press.

Chen, M., and G. P. Wilson. 2015. A multivariate approach to infer locomotor modes in

Mesozoic mammals. Paleobiology 41:280–312.

Cheney, J. A., N. Konow, K. M. Middleton, K. S. Breuer, T. J. Roberts, E. L. Giblin, and S. M.

Swartz. 2014. Membrane muscle function in the compliant wings of bats. Bioinspir.

Biomim. 9:025007. IOP Publishing.

Clavel, J., G. Escarguel, and G. Merceron. 2015. mvMORPH:anR package for fitting

multivariate evolutionary models to morphometric data. Methods Ecol Evol 6:1311–

1319.

Cooper, K. L., and C. J. Tabin. 2008. Understanding of bat wing evolution takes flight. Genes

Dev. 22:121–124.

Darwin, C. 1859. On the origin of species. 1st ed. John Murray, London.

Farnum, C. E., M. Tinsley, and J. W. Hermanson. 2008. Forelimb versus hindlimb skeletal

development in the Big Brown Bat, *Eptesicus fuscus:* functional divergence is reflected

in chondrocytic performance in autopodial growth plates. Cells Tissues Organs 187:35–

47.

Feigin, C. Y., J. A. Moreno, R. Ramos, S. A. Mereby, A. Alivisatos, W. Wang, R. van

Amerongen, J. Camacho, J. J. Rasweiler, R. R. Behringer, B. Ostrow, M. V. Plikus, and

R. Mallarino. 2023. Convergent deployment of ancestral functions during the evolution

of mammalian flight membranes. Science Advances 9:eade7511. American Association

for the Advancement of Science.

Fujiwara, S., H. Endo, and J. R. Hutchinson. 2011. Topsy-turvy locomotion: biomechanical

specializations of the elbow in suspended quadrupeds reflect inverted gravitational

constraints: Elbow mechanics in suspended quadrupeds. J. Anat. 219:176–191.

Gardner, N. M., and T. A. Dececchi. 2022. Flight and echolocation evolved once in Chiroptera:

comments on 'The evolution of flight in bats: a novel hypothesis.' Mammal Rev. 52:284–

290.

Giannini, N. P. 2012. Toward an integrative theory on the origin of bat flight. Pp. 353–384 *in* G.

F. Gunnell and N. B. Simmons, eds. Evolutionary History of Bats. Cambridge University

Press, Cambridge.

Granatosky, M. C. 2018. Forelimb and hindlimb loading patterns during quadrupedal locomotion

in the large flying fox (*Pteropus vampyrus*) and common vampire bat (*Desmodus*

rotundus). J. Zool. 305:63–72.

Grossnickle, D. M., M. Chen, J. G. A. Wauer, S. K. Pevsner, L. N. Weaver, Q.-J. Meng, D. Liu,

Y.-G. Zhang, and Z.-X. Luo. 2020. Incomplete convergence of gliding mammal

skeletons. Evolution 74:2662–2680.

Gunnell, G. F., and N. B. Simmons. 2005. Fossil Evidence and the Origin of Bats. J. Mamm.

Evol. 12:209–246.

Hallgrímsson, B., K. Willmore, and B. K. Hall. 2002. Canalization, developmental stability, and

morphological integration in primate limbs. Am. J. Phys. Anthropol. Suppl 35:131–158.

Hansen, T. F. 1997. Stabilizing selection and the comparative analysis of adaptation. Evolution

51:1341–1351.

Heard, S. B., and D. L. Hauser. 1995. Key evolutionary innovations and their ecological

mechanisms. Hist. Biol. 10:151–173.

Hildebrand, M. 1985. Chapter 6. Digging of Quadrupeds. Pp. 89–109 *in* M. Hildebrand, D. M.

Bramble, K. F. Liem, and D. B. Wake, eds. Functional Vertebrate Morphology. Harvard

University Press.

Hunter, J. P. 1998. Key innovations and the ecology of macroevolution. Trends Ecol. Evol.

13:31–36.

Jetz, W., G. H. Thomas, J. B. Joy, K. Hartmann, and A. O. Mooers. 2012. The global diversity of

birds in space and time. Nature 491:444–448.

Jones, K. E., J. Bielby, M. Cardillo, S. A. Fritz, J. O'Dell, C. D. L. Orme, K. Safi, W. Sechrest,

E. H. Boakes, C. Carbone, C. Connolly, M. J. Cutts, J. K. Foster, R. Grenyer, M. Habib,

C. A. Plaster, S. A. Price, E. A. Rigby, J. Rist, A. Teacher, O. R. P. Bininda-Emonds, J.

L. Gittleman, G. M. Mace, and A. Purvis. 2009. PanTHERIA: a species-level database of

life history, ecology, and geography of extant and recently extinct mammals. Ecology 90:2648–2648.

Kurochkin, E. N., and I. A. Bogdanovich. 2010. Origin of feathered flight. Paleontol. J. 44:1570–1588.

Leal, M., A. K. Knox, and J. B. Losos. 2002. Lack of Convergence in Aquatic Anolis Lizards. Evolution 56:785–791.

Linden, T. J., A. E. Burtner, J. Rickman, A. McFeely, S. E. Santana, and C. J. Law. 2023. Scaling patterns of body plans differ among squirrel ecotypes. PeerJ 11:e14800. PeerJ Inc.

Long, C. A., G. P. Zhang, T. F. George, and C. F. Long. 2003. Physical theory, origin of flight, and a synthesis proposed for birds. J. Theor. Biol. 224:9–26.

Luo, Z.-X., Meng, Q.-J., Grossnickle, D. M., Liu, D., Zhang, Y.-G., Neander, A. I., and Q, Ji. 2017. New evidence for mammaliaform ear evolution and feeding adaptation in a Jurassic ecosystem. Nature. 548: 326-329 (doi:10.1038/nature23483).

Louzada, N. S. V., M. R. Nogueira, and L. M. Pessôa. 2019. Comparative morphology and scaling of the femur in yangochiropteran bats. J. Anat. joa.12996.

Meng, Q.-J., Grossnickle, D. M., Liu, D., Zhang, Y.-G., Neander, A. I., Ji, Q., and Z.-X. Luo. 2017. New gliding mammaliaforms from the Jurassic. Nature. 548: 291-296. (doi:10.1038/nature23476).

Mosimann, J. E. 1970. Size Allometry: Size and Shape Variables with Characterizations of the Lognormal and Generalized Gamma Distributions. J. Am. Stat. Assoc. 65:930–945.

Nicholson, D. B., A. J. Ross, and P. J. Mayhew. 2014. Fossil evidence for key innovations in the evolution of insect diversity. Proc. R. Soc. B Biol. Sci. 281:20141823. Royal Society.

Nojiri, T., I. Werneburg, V. T. Tu, D. Fukui, M. Takechi, S. Iseki, T. Furutera, and D. Koyabu.

2023. Timing of organogenesis underscores the evolution of neonatal life histories and

powered flight in bats. Proc. R. Soc. B Biol. Sci. 290:20221928. Royal Society.

Norberg, U. M. 1985. Chapter 8. Flying, Gliding, and Soaring. Pp. 148–152 *in* M. Hildebrand,

D. M. Bramble, K. F. Liem, and D. B. Wake, eds. Functional Vertebrate Morphology.

Harvard University Press.

Norberg, U. M. 1985. Evolution of Vertebrate Flight: An Aerodynamic Model for the Transition

from Gliding to Active Flight. Am. Nat. 126:303–327. [University of Chicago Press,

American Society of Naturalists].

Norberg, U. M. 1990. Vertebrate Flight. Springer Berlin Heidelberg, Berlin, Heidelberg.

Padian, K. 1982. Running, Leaping, Lifting Off. The Sciences 22:10–15.

Papadimitriou, H. M., S. M. Swartz, and T. H. Kunz. 1996. Ontogenetic and anatomic variation

in mineralization of the wing skeleton of the Mexican free-tailed bat, Tadarida

brasiliensis. Journal of Zoology 240:411–426.

Pevsner, S. K., D. M. Grossnickle, and Z.-X. Luo. 2022. The functional diversity of marsupial

limbs is influenced by both ecology and developmental constraint. Biol. J. Linn. Soc.

135:569–585.

R Core Team. 2021. R: A language and environment for statistical computing. Foundation for

Statistical Computing, Vienna, Austria.

Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other

things). Methods Ecol. Evol. 3:217–223.

Rickman, J., A. E. Burtner, T. J. Linden, S. E. Santana, and C. J. Law. 2023. Size And

Locomotor Ecology Have Differing Effects on the External and Internal Morphologies of

Squirrel (Rodentia: Sciuridae) Limb Bones. Integrative Organismal Biology 5:obad017.

Riskin, D. K., J. E. A. Bertram, and J. W. Hermanson. 2005. Testing the hindlimb-strength

hypothesis: non-aerial locomotion by Chiroptera is not constrained by the dimensions of

the femur or tibia. J. Exp. Biol. 208:1309–1319.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of

image analysis. Nat. Methods 9:671–675.

Sears, K. E., R. R. Behringer, J. J. Rasweiler, and L. A. Niswander. 2006. Development of bat

flight: Morphologic and molecular evolution of bat wing digits. Proc. Natl. Acad. Sci.

103:6581–6586.

Simmons, N. B., and A. L. Cirranello. 2022B. Bat Species of the World: A taxonomic and

geographic database.

Simmons, N. B., and J. H. Geisler. 1998. Phylogenetic relationships of Icaronycteris,

Archaeonycteris, Hassianycteris, and Palaeochiropteryx to extant bat lineages, with

comments on the evolution of echolocation and foraging strategies in Microchiroptera.

Bulletin of the AMNH ; no. 235. Relationships of Eocene bats. [New York] : American

Museum of Natural History.

Simmons, N. B., K. L. Seymour, J. Habersetzer, and G. F. Gunnell. 2008. Primitive Early

Eocene bat from Wyoming and the evolution of flight and echolocation. Nature 451:818–

821.

Smith, J. D. 1977. Comments on Flight and the Evolution of Bats. Pp. 427–437 *in* M. K. Hecht,

P. C. Goody, and B. M. Hecht, eds. Major Patterns in Vertebrate Evolution. Springer US,

Boston, MA.

Stayton, C. T. 2006. Testing hypotheses of convergence with multivariate data: morphological

and functional convergence among herbivorous lizards. Evolution 60:824–841.

Storch, G., B. Sigé, and J. Habersetzer. 2002. Tachypteron franzeni n. gen., n. sp., earliest

emballonurid bat from the Middle Eocene of Messel (Mammalia, Chiroptera). Paläontol.

Z. 76:189–199.

Swartz, S. M., M. B. Bennett, and D. R. Carrier. 1992. Wing bone stresses in free flying bats and

the evolution of skeletal design for flight. Nature 359:726–729.

Swartz, S. M. 1997. Allometric patterning in the limb skeleton of bats: Implications for the

mechanics and energetics of powered flight. J Morphol 234:277–294.

Swartz, S. M., and K. M. Middleton. 2008. Biomechanics of the bat limb skeleton: scaling,

material properties and mechanics. Cells Tissues Organs 187:59–84.

Teeling, E. C., M. Scally, D. J. Kao, M. L. Romagnoli, M. S. Springer, and M. J. Stanhope.

2000. Molecular evidence regarding the origin of echolocation and ‾ight in bats. 403:5.

Thorington, R. W., and L. R. Heaney. 1981. Body proportions and gliding adaptations of flying

squirrels (Petauristinae). J. Mammal. 62:101–114.

Thorington, R. W., K. Darrow, and C. G. Anderson. 1998. Wing Tip Anatomy and

Aerodynamics in Flying Squirrels. Journal of Mammalogy 79:245–250. [American

Society of Mammalogists, Oxford University Press].

.

Thorington, R. W., and E. M. Santana. 2007. How to make a flying squirrel: *Glaucomys* anatomy

in phylogenetic perspective. J. Mammal. 88:882–896.

Upham, N. S., J. A. Esselstyn, and W. Jetz. 2019. Inferring the mammal tree: Species-level sets

of phylogenies for questions in ecology, evolution, and conservation. PLoS Biol.

17:e3000494.

Uyeda, J. C., D. S. Caetano, and M. W. Pennell. 2015. Comparative analysis of principal

components can be misleading. Syst. Biol. 64:677–689.

Vaughan, T. A. 1970. Vaughan, T. A. (1970). The skeletal system. In Biology of Bats 1, ed. W.

A. Wimsatt. New York: Academic Press. pp. 98–139. Pp. 98–139 *in* Biology of Bats.

New York: Academic Press.

Weaver, L. N., and D. M. Grossnickle. 2020. Functional diversity of small-mammal postcrania is

linked to both substrate preference and body size. Curr. Zool. 66:539–553.

Wilson, and R. A. Mittermeier. 2009. Handbook of the Mammals of the World. Wilson&R. A.

Mittermeier, Eds. Lynx Edicions.

Young, N. M., and B. Hallgrímsson. 2005. Serial homology and the evolution of mammalian

limb covariation structure. Evolution 59:2691–2704.

Zhao, F., W. Wang, J. Zhang, J. Wyrwa, and F. Sun. 2019. Aerodynamic Characteristics and

Pitching Adjusting Mechanism of the Flying Squirrel With Deployed Patagium. IEEE

Access 7:185554–185564.

Table 1: AICcW scores from multivariate model-fitting analyses (via mvMORPH).
Comparisons of the fits of multivariate models to the first three principal components of the a)
forelimb dataset and b) hind limb dataset.
732

a) forelimb	AIC	ΔAICc	mAICcW	b) hind limb	AIC	ΔAICc	mAICcW
mvBM1	-94.42	225.50	0.00	mvBM1	-134.64	230.05	0.00
mvOU1	-248.73	71.20	0.00	mvOU1	-330.72	33.97	0.00
mvOUM$_{loc3a}$	-306.82	13.10	0.00	mvOUM$_{loc3a}$	-358.41	6.28	0.04
mvOUM$_{loc3b}$	**-319.93**	**0.00**	**0.99**	mvOUM$_{loc3b}$	-353.51	11.17	0.00
mvOUM$_{loc3c}$	-308.77	11.16	0.00	mvOUM$_{loc3c}$	-358.10	6.59	0.03
mvOUM$_{loc4}$	-302.35	17.57	0.00	**mvOUM$_{loc4}$**	**-364.69**	**0.00**	**0.92**

734
Figure 1. Morphological measurements and phylogeny. We conducted analyses based on this a)
phylogeny (Upham et al. 2019) across b) 14 forelimb linear measurements for each specimen of
scapula height and length (sh, sl), humerus length (hl), distal width (hdw), proximal width (hpw),
and mid-shaft width (hsw), radius length (rl), ulna length and olecranon length (ul, uol), and third
manual digit metacarpal length and width (mcl, mcw), proximal phalanx length and width (ppl,
ppw), and intermediate phalanx length (ipl) and we collected c) 15 hind limb measurements
consisted of pelvis, ilium, and ischium length (pel, il, isl); femur length (fl), distal width (fdw),
and midshaft width (fsw); tibia length (tl), proximal length (tpw), and midshaft width (tmw);
fibula length (fbl), and third pedal digit metatarsal length and width (mtl, mtw), proximal
phalanx length and width (ppxl, ppxw), and intermediate phalanx length (ipxl). We measured
only the third digit in this dataset (and not, for example, the fifth even though that might be of
biological interest) in order to build on previous datasets in the literature to generate a large
enough dataset for evolutionary modeling.
748
749

750
Figure 2. Forelimb and hind limb PCAs. Morphospaces consist of PC1 and PC2 of 14
forelimb traits (left panels) and 15 hind limb traits (right panels). The extant+extinct
morphospace is shown for the a) forelimbs and b) hind limbs. The extant-only morphospace of c)
the mvOUM$_{loc3b}$ optima (large icons) overlaid on the forelimb morphospace and the d) the
mvOUM$_{loc4}$ optima (large icons) overlaid on the hind limb morphospace. Cartoons of a e)
quadrupedal mammal skeleton (*Rattus norvegicus*) versus a f) bat skeleton (*Pipistrellus abramus*
are shown where dark gray shaded bones on the cartoon skeletons correspond to those measured
to generate the morphospace.
759

760
761 **Figure 3. Forelimb trait optima from model-fitting of univariate traits.** Forelimb OUM
762 optima (θ) as a size-corrected trait value (log-shape ratio) with 95% confidence intervals (Tables
763 S3a) for the best-fitting models (ΔAIC = 0) for 10 traits. The AICcW of the model is on top of
764 each panel (Table S2a).
765
766

767
768 **Figure 4. Hind limb trait optima from model-fitting of univariate traits.** Hind limb OUM
769 optima (θ) as a size-corrected trait value (log-shape ratio) with 95% confidence intervals (Tables
770 S3b) for the best-fitting models (ΔAIC = 0) for 10 traits. The AICcW of the model is on top of
771 each panel (Table S2b).
772

774
Figure 5. Hypothetical adaptive landscapes of forelimb and hind limb skeletal relative length and width evolution. Hypothetical fitness peaks are shown for arborealists, gliders, and bats for a broad summary of our conclusions from the skeletal trait optima results discussed. The "length" and "width" in the figure refer to the lengths and widths of individual limb elements. For the a) forelimb, we depict arborealists on a relatively steep adaptive peak at relatively lower length/width optima, gliders on a broad, shallow peak/incline at relatively higher length but not width optima, and bats on a very steep peak at relatively higher length and width optima. For the b) hind limb, we depict arborealists again on a fairly steep adaptive peak at relatively higher width but not length optima, bats on a shallow steep peak at relatively higher length and lower width optima, and gliders on a peak similar to and overlapping with bats' at relatively higher length but similar width optima.