A Genome Scale Transcriptional Regulatory Model of the Human Placenta

Alison Paquette, PhD[1,2+]; Kylia Ahuna[3]; Yeon Mi Hwang[4]; Jocelynn Pearl, PhD[5]; Hanna Liao[1]; Paul Shannon[4]; Leena Kadam,PhD[3]; Samantha Lapehn,PhD[2]; Matthew Bucher[3]; Ryan Roper, MS[4]; Cory Funk, PhD[4]; James MacDonald, MS[1]; Theo Bammler, PhD[1]; Priyanka Baloni,PhD[4]; Heather Brockway, PhD[6]; W. Alex Mason, PhD[7]; Nicole Bush, PhD[8]; Kaja Z Lewinn, ScD[8]; Catherine J Karr, PhD, MD, MS[1]; John Stamatoyannopoulos, MD[1,9]; Louis J Muglia, MD, PhD[10,11]; Helen Jones, PhD[6]; Yoel Sadovsky, MD[12]; Leslie Myatt, PhD[3]; Sheela Sathyanarayana, MD, MPH[1,2]; Nathan D. Price, PhD[4,13], on behalf of program collaborators for Environmental influences on Child Health Outcomes*

1. University of Washington, Seattle WA. 2. Seattle Children's Research Institute. Seattle WA. 3. Oregon Health and Sciences University, Portland OR. 4. Institute for Systems Biology, Seattle WA. 5. Tune Therapeutics, Seattle WA 6. University of Florida, Gainesville Florida. 7. University of Tennessee Health Sciences Center, Memphis, TN.8. University of California San Francisco, San Francisco CA 9. The Altius Institute, Seattle WA. 10. The Burroughs Wellcome Fund Research Triangle Park, NC.11. Cincinnati Children's Hospital Medical Center and Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH.12. Magee Womens Research Institute, Pittsburgh PA 13. Thorne HealthTech, New York City, NY

*See Acknowledgments for full listing of collaborators.

+ To whom correspondence should be addressed: Alison.Paquette@seattlechildrens.org

ABSTRACT

Gene regulation is essential to placental function and fetal development. We report a genome-scale transcriptional regulatory network (TRN) of the human placenta built using digital genomic footprinting and transcriptomic data. We integrated 475 transcriptomes and 12 DNase hypersensitivity datasets from placental samples to globally and quantitatively map transcription factor (TF)-target gene interactions. In an independent dataset, the TRN model predicted target gene expression with an out of sample R^2 value greater than 0.25 for 74% of target genes. We performed siRNA knockdowns of 4 TFs and achieved concordance between the predicted gene targets in our TRN and differences in expression of knockdowns with an accuracy of >0.7 for 3 of the 4 TFs. Our final model contained 113,158 interactions across 391 TFs and 7,712 target genes and is publicly available. We identified six TFs which were significantly enriched as regulators for genes previously associated with preterm birth.

INTRODUCTION

The placenta is an essential organ with multiple functions.[1] It regulates gas, nutrient and waste transport, as well as provides immunological defense. It also performs endocrine functions, producing key hormones and modulating the passage of hormonal and paracrine signals between the mother and developing fetus.[2] Thus, the placenta modulates the *in utero* environment, impacting fetal development, growth, and health.[3] This complex organ is composed of various types of trophoblast cells, including extravillous trophoblasts, cytotrophoblasts, and the syncytiotrophoblast, which are a large terminally differentiated, multi-nucleated cell type that is the primary site of nutrient uptake and transport and peptide and steroid hormone synthesis, as well as other cell types including stromal cells, fibroblasts, endothelial cells, and various immune cells.[4] With its many cell types and highly specialized functions, the placenta has a unique transcriptome,[5] which including numerous genes uniquely expressed only within the placenta. Fetal genomic variation influences placental epigenomic and transcriptomic interactions,[6] which change across gestation,[78] and in response to cues from

the maternal environment.[9] The placental transcriptome reflects the underlying physiological
functions of the organ, and understanding these changes in normal and pathological placental
tissue can provide insight into the molecular mechanisms into pathophysiology, Perturbations to
the placental transcriptome are linked to pregnancy pathologies including preterm birth[10,11] and
preeclampsia.[12] Previous studies of individual genes have revealed several temporally regulated
and cell type specific transcription factors (TFs) that play key roles in discrete developmental
stages and functions of the placenta. Throughout pregnancy, other transcription factors
including *GCM1*, *GATA2*/3 and *CREB* emerge as unique regulators of specific trophoblast
subtypes.[13] These transcription factors regulate key placental processes such as invasion and
nutrient transport that may be linked to adverse placental function and prenatal complications.
Further, understanding these transcriptional changes in normal and pathological placentas can
provide insights into the molecular mechanisms underlying gestational pathophysiology.
64
Transcriptional regulatory networks (TRNs) map TFs to the target genes they regulate.[14,15] This
type of network is context-specific; and requires capturing and quantifying the regulatory
machinery within specific cellular populations and across developmental stages.[16] TRNs
integrate information about chromatin accessibility (captured through DNaseq or ATAC seq) and
predicted transcription factor binding, using sequence motifs from a compendium of TFs and
RNA sequencing data to produce genome scale networks.[17] A TRN is made up of nodes (TFs
and their target genes) and edges that explain which transcription factors play a role in
regulating the target genes. These edges are defined by TF binding within regions of open
chromatin and established relationships in gene expression between TFs and target genes.
TRNs can be organized into units known as "regulons," which are composed of each individual
TF and the target genes they regulate. We have generated such a network previously in the
human brain,[18] which revealed (1) TFs regulated by viral abundance associated with
Alzheimer's disease,[19] (2) TFs that drive gene expression changes associated with bipolar
disorder and schizophrenia,[18] (3) regulatory functions for SNPs associated with genetic risks
associated with these disorders,[20] and (4) TF-target gene relationships underlying gene
expression changes related to Huntington's Disease.[21] This approach has demonstrated that
quantifying transcriptional networks has the potential to illuminate the etiology of a wide variety
of pathological disorders. We now extend this approach into the placenta.
83
Other network biology-based approaches have been previously deployed to gain insight into
placental function and prenatal disease, such as Weighted gene co-expression analysis
(WGCNA), which is a network strategy that identifies co-regulated gene clusters based on
correlated expression.[22] WGCNA was performed on human placental transcriptome data
collected at delivery revealed gene clusters that were involved in core placental functions and
were reproducible across trimesters (in mice).[23] An independent WGCNA constructed on a large
number of term placenta transcriptomes revealed modules associated with fetal growth
restriction.[24] Combining co-expression analyses with chromatin accessibility data is essential for
a deeper understanding of the underlying transcriptional mechanisms of regulation. A placenta
TRN constructed using DNase hypersensitivity data derived from first trimester placental data
has been used to model the transformation of embryonic stem cells to trophoblast cells;
providing key insights into transcription factors which regulate genes with regions of open
chromatin in this context.[25] An accurate model of the placental transcriptome at the end of
gestation, enhanced with DNase hypersensitivity data, will thus provide valuable insight into TF-
target gene interactions and serve as a tool for placental gene expression analyses at this
timepoint.
100
We characterized placental transcriptional regulation at term by constructing a genome scale
TRN of the human placenta, using DNAse hypersensitivity data and curated transcription factor

binding motif databases, paired with RNA sequencing data compiled from 475 placental
samples. To test the accuracy of our model, we applied a machine learning approach to
evaluate the accuracy of the model at predicting gene expression and performed experimental
validation of transcriptional regulation of a core group of TFs in isolated villous cytotrophoblasts.
Because previous transcriptomics identified fetal sex as a critical biological variable that impacts
the placental transcriptome,[26] and pathology,[27] we also characterized sex-specific differences in
TF-target gene interactions within the placenta. We hypothesized that our network could reveal
core TFs that regulated changes in expression of genes whose placental expression was
previously associated with preterm birth. We tested this hypothesis by identifying shared
regulation of genes whose placental expression was associated with preterm birth within an
independent analysis using aggregated microarray data. This approach can be broadly applied
to other types placental 'omics data (e.g. DNA methylation data), providing researchers a
different way to interpret lists of candidate genes and identify transcriptional regulators.
116
117   **RESULTS**
Reconstruction of a transcriptional regulatory network model for the human placenta.
Our placental TRN was constructed using transcriptomic data from 475 placental samples
collected in two independent birth cohorts: at Magee Womens Research Institute (MWRI) in
Pittsburgh PA and at the University of Tennessee Medical Center in Memphis TN, as a part of
the Conditions Affecting Cognitive Development and Learning in Early Childhood (CANDLE)
study. DNase hypersensitivity data was generated in 12 term human placentas from Oregon
Health & Science University in Portland OR. Complete covariate information for all placental
samples used within the model can be found in **Table 1**. The CANDLE cohort was
predominantly Black (56%); whereas the samples from MWRI were predominantly White (85%).
Overall, the demographics of RNA sequencing samples in this dataset was 46% Black, 52%
White, and 2% other Race groups. The mean gestational age of the MWH samples was 39.6
weeks, and the mean gestational age of the CANDLE cohort was 39.2 weeks. 3% of the
samples in the CANDLE cohort were born prematurely (<37 weeks), which we elected to retain
in the model to capture routine variation in the placental transcriptome related to gestational
age. We excluded samples from pregnancies with preeclampsia and multi-fetal gestations in
order to capture the placental transcriptome of infants without complicated deliveries that were
potentially medically indicated.
135
The model construction process is summarized in **Figure 1,** where we explored the impact of
multiple selection in this process. Model construction was done using genomic footprints derived
from placental DNAse hypersensitivity data (**Figure 1A**) paired with known TF binding motifs
(**Figure 1B**) to generate putative edges between TFs and target genes (**Figure 1C**). Next, we
generated a co-expression network of TFs and target genes in our training dataset to
parameterize the model (**Figure 1D**). We evaluated different parameters of the model based on
how they influenced model size and ability to predict gene expression in our test dataset
(**Figure 1E**). We constructed 6 models with different parameters for our training dataset, which
are summarized in **Supplemental Table 1**. For the "reference model" (Model 1), we elected to
use the median correlation coefficient; then included only the top 15 TFs that were correlated
with expression of each target gene in our model. We chose this module size on the basis of
analyses performed in the construction of a previous transcriptional regulatory network.[28] The
accuracy of each model was calculated as the R^2 value of the correlation coefficient (R)
between the actual and predicted gene expression in the dataset generated from LASSO
regression (out of sample R^2, or $OOSR^2$). The full range of predictions for each individual gene
is depicted in **Supplemental Figure S3**. We also assessed the accuracy of the model in our
test dataset using random forest classification, but observed minimal differences between the
two approaches. For Model 1 (the "reference" model), the out of sample R2 accuracy on the test

dataset for each gene from LASSO regression and Random forest were strongly correlated
(Cor=0.98, P <2.2x10^{-16}, Pearson correlation, see **Supplemental Figure S4**). The OOS R^2
value for LASSO regression was slightly but significantly higher than random forest in our test
dataset (Mean Random Forest OOS R^2 =0.35, Mean Lasso R2=0.37,P=3.49x10^{-7}, t test,
Supplemental Figure S4). Thus, we concluded that for these samples, LASSO regression
slightly outperformed random forest at predicting gene expression in our test dataset.
160
In parallel to our models constructed for TFs with strong correlations in genes with TF binding
sites, we also constructed three "null" models; which did not have TF binding sites and/or had
weak correlations, and these models are described in **Supplemental Table S2**. We evaluated
the accuracy of these models to predict expression in our test dataset using the same approach
(**Supplemental Figure S3**). These null models had significantly lower accuracy on predicting
gene expression in the test dataset than any of our actual models (P<0.05, Tukey Test,
Supplemental Table S2). The null models with and without TF binding sites that had
correlations between TFs and target genes less than the median training dataset performed the
poorest at predicting gene expression in a test dataset. The null models which contained TFs
that did not include binding sites at the specific region but were strongly correlated with gene
expression performed better than those with poor correlations, but were still significantly less
accurate at predicting gene expression in the test dataset than those that also included correct
binding data (p<0.05, Tukey Test). Model 9, which contained TFs with no binding sites and poor
correlations, had the worst performance for predicting gene expression based on OOS R^2 in our
testing dataset. All analyses did not include the subset of established TFs without motifs as
defined by Lambert et al.[29] We surmised that there were complex regulatory mechanisms by
which these TFs act to regulate gene expression (including acting as co-factors). Comparison
against the "null" models presented here revealed the extent to which our TRN can predict gene
expression.
180
Overall, the two parameters that altered the model the most were (1) varying the cutoff for the
correlation coefficient and (2) the number of TFs included in each model (**Supplemental Figure
S3**). Models with a more stringent cutoff contained fewer TFs, target genes and interactions
(Models 4 and 5). Model 4 used a higher correlation coefficient cutoff (correlation coefficient
>|0.5|), and overall had the highest accuracy of all the models tested (98% of samples had R^2
>0.25 in test dataset). There was no significant difference in the out of sample R^2 values
between models constructed with and without enhancers (P=1, Tukey Test, see **Supplemental
Table 1**). We constructed our model with enhancers to provide complete coverage of potential
transcriptional regulators. We elected to construct our model using the parameters established
in "Model 3," using a correlation cutoff of 0.25, as this model was among the more
comprehensive in terms of interactions, TFs, and had the highest accuracy outside of Model 4.
192
Computational Validation of Model Accuracy
After model parameterization in the training dataset, we then combined the training and testing
dataset (N=382) and used this as input data to calculate our final accuracy in the holdout
dataset (N=93). This step is critical as a guard against overfitting because it means that the final
form of the model was tested once against a holdout dataset that was completely separate from
all the variations in the model during the projects development phase,and was selected
randomly from samples in the beginning. We analyzed the correlation coefficients in the testing
set as well as the ability of the TFs to predict gene expression in the holdout set using LASSO
regression (**Figure 2**). In the holdout data, 88% of TF-target gene correlations were statistically
significant after adjusting for multiple comparisons (FDR adjusted <0.05), and 83% of TF-target
gene relationships were highly correlated with a correlation coefficient of >|0.25 |. Across all
genes, LASSO regression using the TFs identified in our model predicted gene expression with

205 a mean OOS R^2 of 0.39, and we were able to predict 73% of target genes with an OOS R^2
206 >0.25. We constructed our TRN using only target genes which we were able to predict with an
207 OOS R^2>0.25 in our holdout dataset, removing 2,802 genes from our final model.
208
209 Our final robust model contained 7,765 genes, with a subset of this model including the 3,213
210 genes (41.7%) with OOS R^2 prediction accuracy in our holdout dataset >0.5. The robust model
211 was composed of 113,356 interactions between 391 TFs and 7765 target genes, including
212 74,014 positively correlated interactions and 38,144 negatively correlated interactions. The
213 median regulon size (target genes per TF) was 156, and was heavily tailed with 75 "hub" TFs
214 which regulated >500 genes (**Figure 3**). Network centrality measures are provided in
215 **Supplemental Table S4.** The 8 most central TFs in our model based on eigenvector centrality
216 (measure of influence of node within a network for each transcription factor) were *SP1, SMAD4,*
217 *SP3, CTCF, STAT5B, NR2C2, YBX1 and CLOCK .* Our model contained 25 transcription
218 factors that were placenta specific or placenta enriched, based on the Human Protein Atlas[30],
219 including known placental transcriptional regulators such as *GCM1*, *DLX3*, *DLX4*, and *DLX5*.
220 The largest placental enriched transcriptional regulator was *TFAP2A*, which regulated 1,249
221 different target genes.
222
223 **Experimental Validation of Model Accuracy**
224 We performed an siRNA knockdown of four TFs (*GRHL2*,*CREB3L2, ESRRG, GCM1*) that were
225 selected on the basis of their characterization as "placenta enriched genes" from the Human
226 Protein Atlas [30] and which regulated >125 target genes in our model. TFs knockdown was
227 performed in 7 female and 6 male samples (with the exception of *GCM1*, which had 4 male
228 samples, and *GRHL2,* which had 5 male samples). Gene expression for predicted target genes
229 was quantified through RNA sequencing, and we compared the directionality of the knockdown
230 (log fold change of the knockdown vs. control) with the correlation between TF and target gene
231 (**Figure 4**), with the prediction that knockdown of TFs which were positive regulators in the
232 model would result in decreased expression (negative log fold change), and knockdowns of TFs
233 that were negative regulators in the model would result in increased expression (positive log fold
234 change). We achieved an accuracy of 0.96 for *GRHL2*, 0.80 for *GCM1*, 0.71 for ESRRG, and
235 0.32 for *CREB3L2*. We also analyzed the predictive accuracy and significance of our model
236 within the experimental data in a sex-stratified analysis. In the female siRNA knockdown
237 samples, we achieved a stronger validation of *GRHL2* (Accuracy=1, P=0.015) and *ESRGG*
238 (Accuracy =0.88, P=0.018), and we generally had lower accuracy in male samples
239 (**Supplemental Figure S5**). Overall, our experimental knockdowns revealed that the TRN
240 provided accurate insight into TF regulation for three of our four selected TFs.
241
242 **Identification of sex-specific differences in transcriptional regulation.**
243 Sex specific differences in TF-target gene relationships were captured by generating sex
244 stratified TF-target gene linear models from the complete placental RNA sequencing data
245 (N=475), and comparing the interaction between these 2 linear models. We observed 5,361 TF-
246 target gene relationships between 254 TFs and 1,996 target genes (See **Supplemental Table**
247 **S5**) that were significantly different between male and females after adjusting for multiple
248 comparisons (FDR adjusted <0.05). Six of these transcription factors and 81 of the target genes
249 were specifically located on the X chromosome, and the rest of the interactions were within
250 autosomal regions. The TFs with the biggest sex specific differences were *ELK3*, *SP11*, *ETV5*,
251 *ETS2*, and *TWIST1*. We compared the difference in estimates between males and females and
252 the direction of the estimate between the TF-target gene relationships and observed four distinct
253 types of sex-specific interactions, as shown in **Figure 5**. 291 positive TF-target gene
254 relationships were stronger in males than in females (Quadrant 1). 3,681 positive TF-target
255 relationships were stronger in females than in males (Quadrant 2). 1261 negative TF-target

256 gene relationships were stronger in females than in males (Quadrant 3), and 120 negative TF-
257 target gene relationships that were stronger in males than in females. (Quadrant 3) The top 3
258 biggest differences for each quadrant are depicted in **Supplemental Figure S6.** Overall, these
259 perturbed TF-target gene relationships indicate differences in regulation of gene expression
260 between female and male placentas that can be identified in our robust model..
261
Applying the Placental TRN to discover transcriptional regulators of genes associated
with preterm birth
Ultimately, the goal of the placental transcriptional regulatory network is to gain insight into
transcriptional regulation in the context of prenatal exposures or placental pathologies. As a
proof of concept, we used the placental TRN to identify key transcription factors that regulated
the group of 129 genes whose placental expression was associated with preterm birth, which
we established in an independent publication.[10] These 129 genes were regulated by a total of
223 transcription factors. We identified enriched transcription factors using hypergeometric
tests. Here, the regulons were used in lieu of a gene ontology (GO) or KEGG gene set. Seven
TFs (*AHR*, *ARID5B*, *BCL6B*, *NFE2L2*, *NR3C2*, *TBX5*, *THAP5*) were significantly enriched for
these pre-term birth associated genes (p<0.1, Fishers exact test). Together, these 7 TFs were
responsible for regulation of 35.7% of the differences in expression observed (46/129), including
some of our most significantly upregulated (*BIN2*) and downregulated genes (*ADAMTS3*).
Analysis of connectivity amongst these TFs showed two clusters (**Figure 6**). *BCL6B* and *TBX5*
formed a unique cluster regulating 4 genes together and 2 independent genes each; which were
all downregulated in preterm placenta. *BCL6B* was the most statistically significant enriched
gene in our network. *TBX5* is classified as a placental-specific gene by the Human Protein
Atlas, and it (along with BCL6B) directly regulated 2 other placenta-specific genes (KL and
FAM162B). The other five enriched TFs (*AHR*, *ARID5B*, *NFE2L2*, *NR3C2*,*THAP5*) formed a
highly complex subnetwork that regulated the remaining 38 target genes. Four of these five TFs
had identified binding sites and were positively correlated with each other (*AHR*, *ARID5B*,
NFE2L2, *THAP5*), with *THAP5, AHR, and ARID5B* emerging as the upstream regulators based
on hierarchical mapping of the network. Among these genes, *THAP5* and *AHR* had the highest
number of differentially expressed genes in total, regulating 20 and 23 genes respectively. This
subnetwork revealed that the differentially expressed genes could be attributed to a small
number of transcriptional regulators that formed distinct subnetworks.
288
Additionally, six of the 129 differentially expressed genes we previously identified from our
previous study[10] were transcription factors, and these TFs were regulated in part by the
enriched TFs. These six TFs themselves regulated downstream genes that were significantly
associated with preterm birth (**Table 2)**. *THAP5* was a regulator for two TFs (*ATF1* and
CLOCK1) that were hub genes in our network and were downregulated in preterm placentas.
These transcription factors are crucial for placental transcriptional regulation, and regulate a
number of genes in the placenta, including a subset associated with preterm birth. Our TRN
revealed that these genes themselves are regulated by a subset of transcription factors
enriched for PTB genes. It also highlights that some of the genes previously associated with
preterm birth are important to placental transcriptional regulation, based on their degree
centrality. These data demonstrate that our placental TRN can be used to identify regulatory
pathways driving differential gene expression in placentas from preterm births
301
302   **DISCUSSION**
303 In this study, we generated, for the first time, a placenta-specific TRN using established
304 genomic footprinting techniques[31] and high dimensional RNA sequencing data from 475
305 placental transcriptomes. Our network captures 113,356 interactions between 391 TFs and
306 7,765 target genes. Our model was validated against a separate transcriptomic dataset and

307 follow-up experiments for several key regulators. Our study revealed that the TFs *SP1*,
308 *ZNF148*,*SP3*, *SMAD4*, and *CLOCK* were master transcriptional regulators of placental gene
309 expression within the TRN; 25 transcription factors unique to the placenta, with TFAP2A
310 emerging as the largest regulon, and a subset of 5,361 TF-target gene interactions that
311 displayed sexual dimorphisms between male and female samples, with more significant
312 transcriptional regulation occurring within female placentas. We applied our model to identify
313 transcriptional regulators of a subset of genes previously associated with preterm birth and
314 found 7 TFs that were enriched as regulators of genes whose placental expression was
315 previously associated with preterm birth, which may indicate clinically modifiable targets.
316 Importantly, this model captures genomic regulation from placentas collected at the end of
317 pregnancy, which is generally when placental samples are readily available and typically
318 collected for large cohort studies. To facilitate additional studies in the community, we have
319 made our model publicly available as a web accessible resource.
320
321 **Comparison of the Placental TRN to other Network Models**
322 The major advancement of this study is the generation of a large and sophisticated TRN model
323 using DNAse hypersensitivity data within key placental tissues at term, and the validation of this
324 model using gene expression data captured within a large and diverse cohort. Our model also
325 complements other models of placental transcription. An independent WGCNA of the human
326 placenta constructed using only transcriptomic data revealed 5 hub modules associated with
327 fetal growth restriction (FGR).[24] One of the core genes of the module associated with both FGR
328 and fetal overgrowth was *CREB3*, which was a hub TF that regulated 738 target genes within
329 our model. Our model builds upon this by inclusion of information about chromatin accessibility
330 using DNAse hypersensitivity data. DNAse hypersensitivity data (without co-expression
331 analysis) has been used to construct a gene regulatory model of early placental transcriptional
332 regulation, including mapping of genes involved in differentiation of human embryonic stem cells
333 to trophoblast cells in mice, and identified 20 key TFs involved in placental differentiation
334 through centrality analysis.[25] Many of the genes reported as candidates for regulating placental
335 development genes are reported as hub TFs in our model, including *SMAD4*, *ETS1*, *RUNX1*,
336 *CREB1*, *SMAD3*, *PPARA*, and *JUND*. Our model captures regulation of key transcription factors
337 identified in other studies from human placental data collected at term. A WGCNA-based
338 analysis revealed that a large proportion of the placental transcriptome is organized into
339 modules of co-expressed genes that were involved in core placental functions and were
340 reproducible across trimesters.[23] Thus, our model of the placental transcriptome generated at
341 the end of gestation is concordant with previously generated network-based analyses earlier in
342 gestation, suggesting that these TFS are essential not only to placental development but also
343 placental function across gestation.
344
345 **Core Transcriptional Regulators of Placental Gene Expression**
346 Many of the highest-ranking core TFs in our model -- including *SP1*, *SP3*, *SMAD4*, and *TFAP2A*
347 – are essential regulators of placental function. *SP1* and *SP3* had the highest eigenvector and
348 degree centrality in our model. SP transcription factors are involved in placental glucocorticoid
349 signaling,[32] and placental expression of this gene has also been linked to preeclampsia.[33]
350 *SMAD4* had the second highest eigenvector centrality value, and is a signaling protein in the
351 TGF family which is activated in response to TGF-beta. This signaling pathway plays a crucial
352 role in regulation of trophoblast proliferation, differentiation and invasion, as was reviewed by
353 Adu-Gyamfi et al.[34] *TFAP2A* regulated 1255 target genes and was the largest placental specific
354 TF based on both eigenvector centrality and degree centrality. *TFAP2A* encodes the activating
355 enhancer binding protein 2-alpha (AP-2α); a transcription factor that is involved in several key
356 processes unique to the placenta.[35] *TFAP2A* expression is detectable throughout the placenta,
357 including extravillous cytotrophoblasts, cytotrophoblasts and syncytiotrophoblasts.[4] AP-2α

promotes the EGF-dependent invasion of the human trophoblast,[36] and induces human CRH
gene expression.[37] In our TRN, we were able to recapitulate some of these published findings of
TFAP2A regulation. In another study, the TFs SP1 and SP3 regulate expression of *TFAP2A* in
trophoblast cells.[38] In our model, *SP1* and SP3 were included in the top 15 TFs regulating the
expression of *TFAP2A*. TFAP2A is a positive regulator of CRH in our model; in alignment with
work showing that AP-2 induced CRH gene expression.[37] Thus, our TRN reveals the
significance of this TF for placental function as well as some core TFs that regulate many genes
within the placenta.
366
Sex Specific Regulation of Placental Gene Expression
Sex differences at the molecular level between male and female placentas may contribute in
part to differences in growth rates of male and female fetuses and differential responses to
perturbations in the in utero environment, and manifestation of disease later in life (fetal
programming).[39] In our analysis, we identified 5,361 TF-target gene relationships that were
different between male and female placental samples. Many of these TF-target gene differences
may be regulated in part by hormonal differences, as we saw a number of differences related to
nuclear hormone receptors that are responsive to sex hormones, including *ESR1*, *ESSRA*,
ESRRG, and *PGR*. We observed a general trend in our interaction terms where both negative
and positive TF-target gene relationships were stronger in female placentas compared to male
placentas. An aggregated meta-analysis of microarray data revealed 140 differentially
expressed genes in male and females samples, with the majority residing in autosomal regions,
and identified several TFs that were enriched for these genes, including *NFATC2*, *AHR*, *NKX3*,
HIF1A, and *REL*.[26] A number of these TFs were associated in our analysis with sex differences
in expression of target genes, such as the *AHR*; which was associated with 111 TF-target gene
relationships that were significantly different between males and females. Based on these
findings, we hypothesize that there may be stronger transcriptional control of gene expression in
female compared to male placentas. These differences may be driven by sex-specific
differential response to stress hormones, environmental chemicals, and nutrition, as reviewed
by Rosenfeld.[40] This is significant because there is variation in pregnancy outcomes relate to
fetal sex,[41] such as an increased rate of preterm birth in male infants.[42] To our knowledge, this
analysis is the first to specifically investigate TF-target gene differences by sex within the
context of a TRN. Further analysis needs to be performed on these relationships, including
factors that modify these relationships and the role of genomic imprinting.
391
The Placental TRN reveals Transcriptional Regulators of Prematurity
Our ultimate goal is to apply this placental TRN as a tool to help identify perturbed regulation in
advance pregnancy outcomes, or in relation to environmental exposures. To this end, we used
the TRN to identify seven different transcription factors enriched for genes associated with
preterm birth that we identified in an independent study.[10] *TBX5* and *BCL6B* regulated a
distinctive group of PTB genes within our TRN; and *BCL6B* was the most enriched TF in our
model (p=0.016). *BCBL6B* encodes the B-cell CLL/lymphoma 6 member B protein, a
transcriptional repressor which mediates VEGF signaling and angiogenesis,[43] which are both
key to placental function. *TBX5* has been previously linked to PTB,[44] and interestingly is
placentally enriched in the human protein atlas,[30] suggesting potential function in placental
development and maturation. The other 5 TFs were highly co-regulated within our model, with
THAP5, *ARID5B* and *AHR* emerging as the master regulators within this subnetwork. *THAP5*
encodes a transcriptional repressor that is not well studied due to a lack of orthologs in mice
and humans, but may have pro-apoptotic functions.[45] *ARID5* encodes AT-rich interactive
domain-containing protein 5B which is a transcriptional coactivator involved in cellular growth
and differentiation. Methylation of the *ARID5B* promoter region in the placenta has also been
associated with decreased fetal birthweight.[46]

409
Two of the TFs that were enriched for preterm birth associated genes, *AHR* and *NFE2L2*, are
important xenobiotic response genes that respond to environmental perturbations and are linked
to inflammation AHR encodes the Aryl Hydrocarbon Receptor, a nuclear hormone receptor
which is activated by exogeneous ligands. Beyond its role in cellular detoxification, this receptor
is also involved in cellular proliferation, migration and development,[47] and regulation of the
innate and adaptive immune system.[48] *AHR* is highly expressed throughout pregnancy, and
protein expression increases five-fold between first trimester and term placenta within
syncytiotrophoblasts of chorionic villi.[49] AHR agonists alter the vascular remodeling of the
placenta, resulting in reduced volume of the placental labyrinth zone.[50] AHR is hypothesized to
be involved in preterm birth through promotion of pro-inflammatory cytokines.[51] Prenatal
exposure to AHR agonist TCDD increases the incidence of preterm birth in mice[52] and induces
pro-inflammatory responses in human placental explants.[53] Placental inhibition of AHR has
been shown to inhibit cell migration as a potential mechanism.[54] Specific AHR agonists such as
cigarette smoke are associated with increased odds of developing preterm birth.[55] In our model,
AHR regulated expression of Nuclear factor erythroid 2-related factor 2 (NRF-2; encoded by
NFE2L2), which was also enriched for preterm birth genes. There is substantial cross talk
between NF-2 and AHR,[56] and our work is concordant with previous analyses demonstrating
that AHR regulates NRF-2 expression by binding to the promoter region of the gene in other
tissues.[57] We observed this same relationship within our own TRN as AHR was one of the
significant positive regulators of NRF-2 within the placenta. NRF-2 is a master regulator of
antioxidant response and its products protect cells from oxidative insults.[58] NRF-2 has been
implicated as a modulator of oxidative stress in early pregnancy.[59] NRF-2 deletion in mice leads
to increased inflammation and premature labor after treatment with LPS, changes to the volume
of the labyrinth zone of the placenta, reduction in placental volume, increased oxidative stress,
elevated inflammatory cytokines and cell death.[60,61] In human studies, alterations in NRF-2 are
attributed to changes in placental cells and preeclampsia.[60] Our TRN reveals that many of the
preterm birth associated genes are regulated by AHR and NRF-2, which could indicate
activation of these gene expression pathways in preterm placentas.
438
Applications of the Placental TRN
Placental omics data can be used to quantify perturbations that influence gene regulation and
physiological activity, providing insight into the underlying etiology related to outcomes or
exposures.[62] Many researchers studying the placenta have used omics data to generate lists of
genes which are associated with prenatal exposures as well as infant or child health outcomes,
as summarized in our recent review.[63] These genes are often contextualized using pathway
analysis approaches within KEGG or Gene Ontology (GO) gene sets. The TF enrichment
analysis we have performed using regulons is complementary to using other defined gene sets
such as KEGG or GO. TRNs are focused on the question: "*What transcription factors regulate*
these genes within the placenta?" There are other applications that provide an opportunity to
query TF binding, sites such as ChEA[64], which are based on aggregation of Chip-seq data that
is not tissue or cell type specific. Our TRN provides a placental tissue-specific version that can
be specifically queried to investigate individual TFs or target genes of interest; or to perform an
enrichment analysis similar to what we have performed using genes associated with premature
birth. We have presented the full results of the genome-scale model in the format of a
searchable web tool, PlacentalTRN, which is available at
https://github.com/AlisonPaquette/PlacentalTRNRShinyApp. This model has 3 features: (1) a
target gene search feature, (2) a transcription factor search feature, and (3) an enrichment test.
The search tools allow users to search the model for specific genes or specific transcription
factors, and evaluate model accuracy and predicted targets on a case-by-case basis. The
enrichment test tool allows users to import a list of genes (such as the list of preterm birth genes

presented here), and identify transcription factors which regulate these genes. For those
interested in accessing the TRN independently of the web interface, we provide the entire TRN
in **Supplemental Table 6**. Our intention is that other placental biology researchers will be able
to easily access the TRN and identify transcriptional regulators of their own gene lists of interest
by using this database.
465
Strengths and Limitations
Our model of placental transcriptional regulation should be interpreted with limitations in mind.
We use TF expression as a proxy for TF activity, which does not account for post transcriptional
or post translational modifications, and may be confounded by temporal and spatial
disconnects, cellular heterogeneity, and differences in proliferation,[65] which are common for all
studies that employ this approach. We anticipate confounding due to cellular heterogeneity, as
TRNs reflect epigenetic regulation dependent on regions of open chromatin and DNA
methylation which is intrinsically cell but not tissue-type specific, and our results are not built on
placental data which is a composite of different cell types. This can create false positive
relationships and dilute signal strengths between known TFs and target genes. For each target
gene, we present only the top 15 TFs that are best predicted to regulate expression. Thus, we
may not represent all the TFs with binding sites within that region. Our analyses are composed
of data solely from placentas at the end of gestation from relatively non-pathological
pregnancies without preeclampsia, and so may be inappropriate for researchers studying earlier
developmental time points, and also may not represent perturbed TF-target gene interactions
related to those pathologies. As always, it will be important to further test the model in an
entirely separate validation cohort collected from different participants as they become
available. In the future, we hope to build upon this model by including cell type specific
interactions and temporal resolution of time series changes that occur in the placental
transcriptome.
486
Our model makes a contribution as the first publicly available placental transcriptional regulatory
network, as well as one of the first models of transcriptional regulation of any organ with an
accessible and searchable database that allows researchers to perform their own analysis. We
applied rigorous methods to test the accuracy of our model, including an independent validation
dataset which is the gold standard in machine learning approach.[66,67] To our knowledge, our
model is composed of the largest placental transcriptomic dataset constructed to date. We also
provide an innovative assessment of sex differences in TF-target gene relationships in our
model, which further captures the importance of fetal sex as a biological variable in placental
omics research. Finally, we have tested our TRN using isolated villous cytotrophoblasts
collected from placentas of non-pathological women delivering at term, which are more
reflective of the term placental transcriptome than commercial placental cell lines which are
commonly used for transcriptomic analyses. This data could potentially be used in the future to
shed insight into transcriptional mechanisms related to pregnancy complications, and identify
pharmacological interventions to target specific transcription factors that alter placental function.
501
502   **MATERIALS AND METHODS**
503 **1. Sample Information**
504 To construct the TRN, we integrated transcriptomic and chromatin accessibility data from
505 placental tissue generated from 3 different sources as described below.
506
1A. Magee Women's Research Institute (MWRI) Samples:
Tissue was collected from non-pathological placentas from pregnant individuals who delivered
at the Magee Womens Hospital in Pittsburgh, PA. Inclusion criteria included: Maternal age >18
years, signed informed consent, singleton pregnancies and gestational age >37 weeks.

Exclusion criteria included pregnancy complications including, fetal growth restriction (FGR),
chromosomal abnormalities, preeclampsia, gestational diabetes, chorioamnionitis, antepartum
hemorrhage, and a previous history of preterm birth. Placental biopsies (5 mm^3) were sharply
dissected within 30-40 minutes after delivery using a region of the placenta that is midway
between the cord insertion and the placental margin, and between the chorionic and basal
plates, and free of any macroscopic lesions, as we previously detailed.[68] Tissue biopsies were
placed in RNAlater for 48 h in 4°C, then stored in -80°C until processed for RNA sequencing.
Directional RNA-seq was performed by the Genomics, Epigenomics and Sequencing Core
(GES Core) at the University of Cincinnati. Approximately 50 mg tissue was homogenized in 0.8
ml Lysis/Binding Buffer from mirVana miRNA Isolation Kit (Thermo Fisher, Grand Island, NY) by
using FastPrep-24 5G homogenizer (MP Biomedicals, Solon, OH). Total RNA extraction was
conducted according to manufacturer's protocol. RNA quality was examined on an Agilent 2100
Bioanalyzer (Santa Clara CA). rRNA was depleted using the Ribo-Zero Gold kit (Illumina, San
Diego, CA) with 1 µg of total RNA as input for the SMARTer Apollo NGS library prep system
(Takara, Mountain View, CA). The library for RNA-seq was prepared by using NEBNext Ultra II
Directional RNA Library Prep kit (New England BioLabs, Ipswich, MA) and the sequencing was
run on an Illumina HighSeq2000 with paired end reads, 101bp read length, and at 50 million
read depth.
529
1B. CANDLE Samples: Placental tissue was collected from 371 non-pathological placental
transcriptomes collected from women from Shelby County Tennessee as part of the CANDLE
(Conditions Affecting Neurocognitive Development and Learning in Early childhood) study.
Inclusion criteria included: Maternal age >18 years with signed informed consent. Exclusion
criteria included infants categorized as having severe pregnancy complications recorded in the
CANDLE medical records, including confirmed clinical chorioamnionitis, preeclampsia,
oligohydramnios, placental abruption, infraction or previa, and fetal chromosomal abnormalities.
Methods of placental sample collection and RNA sequencing processing for this cohort have
been previously described.[69] Briefly, within 15 minutes of delivery, a piece of placental villous
tissue approximately 2 x 0.5 x 0.5 cm was dissected from the placental parenchyma. The tissue
cubes were stored in 20 ml RNAlater and refrigerated at 4°C overnight (at least 8 hours but no
more than 24 hours). Following this, each cube was transferred to an individual cryovial
containing fresh RNAlater and stored at -80°C. The fetal villous tissue was manually dissected
and cleared of maternal decidua. RNA was isolated using the AllPrep DNA/RNA/miRNA
Universal Kit (Qiagen; Germantown, MD) according to the manufacturer's recommended
protocol. RNA purity was assessed by measuring $OD_{260/230}$ and $OD_{260/260}$ ratios with a NanoDrop
8000 Spectrophotometer (Thermo Fischer Scientific; Waltham MA). RNA integrity was
determined with a Bioanalyzer 2100 using RNA 6000 Nanochips (Agilent; Santa Clara, CA).
cDNA libraries were prepared from 1 µg of total RNA using the TruSeq Stranded mRNA kit
(Illumina, San Diego, CA) and the Sciclone NGSx Workstation (Perkin Elmer, Waltham, MA).
Prior to cDNA library construction, ribosomal RNA was removed by means of poly-A
enrichment. Each library was uniquely barcoded and subsequently amplified using a total of 13
cycles of PCR. Library concentrations were quantified using Qubit fluorometric quantitation (Life
Technologies, Carlsbad, CA). Average fragment size and overall quality were evaluated with the
DNA1000 assay on an Agilent 2100 Bioanalyzer. Each library was sequenced to approximately
a depth of 30 million reads on an Illumina HiSeq sequencer.
556
1C. Oregon Health Sciences Center Placental Repository (OHSU):
12 placental villi samples were derived from non-pathological, pregnancies from pregnant
individuals delivering via caesarean section at the OHSU Hospital labor and Delivery Unit.
Samples were specifically selected to be equally matched across fetal sexes, and from term
placentas (>37 weeks). Exclusion criteria included maternal BMI >25, maternal age <18 or >40,

any current pregnancy complications and current smokers. Placentas were collected and
weighed immediately following cesarean section. Five random samples of tissue (~80 grams)
were collected from each placenta and placed in PBS to be transported back to the lab. The
chorionic plate and decidua were removed from each randomly isolated placental sample,
leaving only villous tissue, which was thoroughly rinsed in PBS to remove excess blood. Primary
cytotrophoblast were isolated from villous tissue using a protocol adapted from Eis *et al.* using
trypsin/DNase digestion followed by density gradient purification. Isolated cytotrophoblast cells
($5-10 \times 10^6$ cells /ml) were then frozen in freezing media at (10% DMSO in FBS) and stored in
liquid nitrogen until usage. Placentas were collected f into a tissue repository under a protocol
approved by the Oregon Health & Sciences University Institutional Review Board with informed
consent from the patients. All tissues and clinical data were de-identified before being made
available to the investigative team.
574
2. Quantifying Placental Gene Expression in the CANDLE AND MWRI Placentas
A flowchart describing sample preprocessing is described in Supplemental **Figure S1B**. For the
RNA sequencing samples from MWRI and the CANDLE cohort, transcript abundances were
estimated using the quantification program Kallisto.[71] Initially quality control was performed on
each RNA sequencing sample using FastQC.[72] Length scaled counts were condensed to
Ensembl Gene IDs using the Bioconductor TXImport package.[73] Genes were then filtered to
remove low expressing genes using the "FilterByExprs" function of EdgeR, which by default
removes genes with <10 counts (0.5 CPM) in 70% of samples.[74] For the dataset from MWRI,
this included 17,574 genes, and from the CANDLE samples, this included 16,440 genes.
Datasets were then normalized using the trimmed mean of M,[75] and then precision weights
were generated for log-cpm values using the voom transformation.[76] The datasets were then
merged by all genes common across both datasets, generating a final transcriptomic dataset of
14,837 genes in 475 samples. Confounding related to both RNA sequencing batch and intrinsic
differences between the cohorts was eliminated using the comBat algorithm from the
Bioconductor SVA package,[77] which uses an empirical Bayes method to estimate location and
scale parameters for each batch, which are then removed to provide a consistent mean and
variance for each gene, across all batches. This approach has successfully been used to
integrate microarray datasets,[10] as well as RNA sequencing datasets.[78] RNA sequencing batch
was set as the "batch" variable to be adjusted for (The CANDLE cohort was sequenced in 4
batches, and the MWRI dataset was sequenced as one batch), and maternal race and
gestational age were included as covariate variables in the model matrix, to ensure that
differences related to these variables would be retained in the final dataset. Batch effects were
evaluated using principal components analysis (See **Supplemental Figure S2; Supplemental**
Table 1). For TRN Construction, the data was split into a holdout dataset (20% of samples-
N=93), and 20% of the remaining samples was used for the testing dataset (N=75), with 80%
used for the construction of the initial TRN (N=307). After model splitting but prior to model
construction, we tested to ensure that there were no statistical differences between the training
and holdout datasets in relation to the distribution of covariates including batch/cohort, labor
status maternal race, and fetal sex, (categorical, tested using chi squared tests) as well as
gestational age and birthweight (continuous, T-tests). In the final model, all samples were used
(N=475 samples).
606
3. DNase Hypersensitivity Data Generation
DNase hypersensitivity data was generated from the samples collected at OHSU. Altius Institute
used an established protocol for tissue processing developed as part of the ENCODE project.[79]
Samples were processed using the DHS-pseudonano assay & nuclei were sterile filtered (0.25-
6.4 million nuclei/sample), with minimal cellular debris, and treated with 3 doses of DNase (40,
60 and 80 units) followed by a stop buffer including RNAse A, followed by treatment with

proteinase K. The samples had a spot (signal portion of tags) score >0.5 except for 1 sample
(which had a score of 0.34), which is a measure of enrichment, reflecting the fraction of reads
that fall within peak calling regions. A score >0.4 is indicative of high quality DNase sequencing
data meeting ENCODE quality guidelines. Peak regions of DNase-seq data were identified
using F-seq,[80] a density estimator for high-throughput sequence tags, using a threshold
parameter of 2.5, which was selected based on previously reported performance metrics.[81] The
peak regions and DNase-seq data were both used to detect genomic "footprints"–i.e regions of
accessible–DNA using HINT,[82] a method based on hidden Markov models.
621
4. Transcription Factor Motifs & Single Gene Model Construction.
For each gene in our network, we identified putative TFs through integration of the DNAse
hypersensitivity data and TF binding motif information. Transcription factors were defined by the
HumanTFDB which had motifs from these the TF databases (667 TFs).[29] Transcription factor
motifs were identified using three established databases: JASPAR,[83] HOCOMOCO,[84] and
Swissregulon.[85] We first matched these binding sites with the footprints in the placenta
(described above) using FIMO (Find Individual Motif occurrences), applying the default 10-4
match threshold (Version 5.0.5).[86] The motifs were aggregated into position weight matrices
within the R package "MotifDB".[87] We assigned motifs to their TFs as well as those in the same
DNA-binding domain family using the TFClass database, as established in our previous work.[17]
In our analysis, we did not consider the subset of transcription factors classified by Lambert et al
with no known motifs. We included both promoter and enhancer regions in our model. The
promoter region was defined as +/- 5KB from the TSS which has been shown to maximize
target gene prediction in transcriptional regulatory networks constructed with footprinting data.[88]
The Enhancer regions were identified using genehancer (Version-4.11).[89] Genehancer is a
database that integrates reported enhancers from multiple sources, including ENCODE and
Ensembl, and uses a combinatorial likelihood method to define genes based on tissue co-
expression correlations, enhancer-targeted transcription factors, expression quantitative trait
loci, and chromatin conformation capture. We included all enhancer regions defined by
Genehancer as "elite", indicating that they had more than 2 evidence sources, and those
defined as "placental-specific" based on Genehancer analysis of tissue specific genes.
643
5. TRN Construction & Parameterization
5A. Constructing and evaluating Model in Training Data:
We integrated the data described above to generate a model consisting of nodes and edges. A
"putative edge" was identified between a TF and a target gene if a motif for that transcription
factor was present in a region of open chromatin (i.e footprint) in the gene's promoter or
enhancer (see Figure 1). We generated a co-expression matrix for each "putative edge". We
ranked each target gene based on the absolute value of the pearson correlation coefficient. We
pruned our model through an initial "baseline model" (Model 1), including only TFs with >
median absolute value of the overall Pearson correlation coefficient for each target gene, then
including only the top 15 genes based on absolute Pearson correlation coefficient. We elected
this initial module size of 15 based on similar analyses performed in the construction of a
transcriptional regulatory network constructed in the context of T-helper Cell Type 17 (Th17)
differentiation.[28] To evaluate the accuracy of our initial TRN and select model parameters for our
full model, we used LASSO regression and random forest to evaluate how the expression of the
TFs in our model predicted gene expression of each target gene in our testing dataset. The R^2
value was calculated as the squared estimate of the correlation between the predicted vs. actual
expression values. We selected the best model to use in the holdout dataset based on these
thresholds. Machine learning was performed using the "glmnet" (version 2.0-16) and "random
forest' (version 4.6-14) R packages.
663

5B. Null Model:
We created three "null" TRNs to evaluate the validity of our model to distinguish between signal
and noise, which we constructed in our training and evaluated in testing dataset. The first null
model was constructed from 15 randomly selected genes with motifs but with correlations that
were less than the median correlation coefficient. In parallel to our model constructed above, we
generated a network consisting of "null putative edges" defined as interactions between a TF
and a target gene if a motif for that transcription factor was not present in a region of open
chromatin (i.e, footprint) in the genes promoter or enhancer. We generated a co-expression
matrix for each "null putative edge" within our training dataset. Our second null model was
constructed of 15 randomly selected genes with no motifs but with correlation coefficient's > the
median value in our true dataset. Our third null model was constructed of 15 randomly selected
genes with correlation coefficients less than the median value in our true dataset and with no
motifs. In some instances, 15 genes were not available that met these criteria, and in this case,
we included all the genes available.
678
5C. Final Model Construction: After initial model parameterization in our training and testing
dataset, we moved on to combine these 2 datasets, and then test the model in our final holdout
dataset using the best parameters as identified from our original models. We did not include
target genes with an out of sample R^2 accuracy <0.25. In the full model, network connectivity
scores were calculated using the R package "igraph" (Version 1.2.4.1). To identify sex-specific
differences in TF-target gene relationships, we ran linear regression analyses for each target
gene-TF interaction in our TRN with fetal sex as an interaction term. Models were considered
statistically significantly different by sex if the FDR adjusted Q of the interaction term was <0.05.
We ran stratified models in male and female samples to calculate the coefficient estimate and
identify differences in these estimates.
689
6. Experimental Validation
We performed knock downs of selected TFs in primary villous trophoblast cells from placental
samples collected through the OHSU placental biorepository. Placenta specific TFs were
identified using the Human Protein atlas [30], then 4 TFs were selected based on module size
(>125 target genes) and median gene expression (> Median RPM). Separate transfections were
performed using the ON-Target SMARTpool siRNAs (Dharmacon) to induce silencing of four
genes: *CREB2L2*: (Catalog # L-019357-02-0005), *ESRRG* (Catalog # L-003405-00-0005),
GCM1 (Catalog # L-011491-00-0005) and *GRHL2* (Catalog # L-014515-02-0005) with a non-
targeting negative control (Catalog # D-001810-10-05). siRNAs were diluted in nuclease free
water and stored at a 50 µM stock concentration. These transfections were each performed
from villous trophoblasts collected from elective c-sections at OHSU as described above.
701
For transfection, cytotrophoblast cells were rapidly thawed in a 37^0C water bath and
immediately diluted in Iscove's modified Dulbecco's medium (25mM glucose, 4mM glutamine
and 1mM pyruvate (Gibco #12440-046) supplemented with 10% FBS (Corning #35-010-CV)
and 1% penicillin/streptomycin (Gibco #15140-122) termed as complete media. Cells were
centrifuged at 1000g for 10 min and suspended in fresh complete media. The cells were plated
at $2x10^6$ cells per well concentration in 12-well culture plates and cultured at 37^0C, 5% CO_2.
After 24 hours, cells were transiently transfected using the above siRNAs and Lipofectamine
(Invitrogen #11668-019) as per manufacturer's protocol and incubated for another 24hrs. At the
end of 24hrs (total of 48hrs from the start of culture) the transfection media was replaced with
complete media and cells were cultured for another 24hrs. After 72h of total culture time (24h
following previous complete media change) the cells were lysed in 700uL of Qiazol and RNA
was isolated using the Zymo Direct-zol Mini-prep kit and reagents (Zymo-Catalog # RD2051).
RNA integrity was determined with a Bioanalyzer and only RNA samples with a RNA Integrity

Number (RIN) > 8.6 were used for RNA-Seq analysis. cDNA libraries were prepared from 1 μg
of total RNA using the Illumina Stranded Total RNA library kit. Each library was sequenced to
approximately a depth of 50 million reads on an Illumina NovaSeq 6000 S4 (2 x100 read length)
sequencer by the OHSU sequencing core.
719
We used a similar pipeline to process the RNA sequencing data as was used for the human
placental data, as described above. Transcript abundances were estimated using Kallisto, and
length scaled counts were condensed to Ensembl Gene IDs using the Bioconductor TXImport
package.[73] Genes were then filtered to remove low expressing genes using the "FilterByExprs"
function of EdgeR.[74] Differentially expressed genes between siRNA knockdowns vs. negative
scrambled controls were identified using generalized linear models implemented within
EdgeR,[74] including only target genes that were predicted to be expressed by each TF.
Dispersion parameters were estimated using the Cox-Reid method, and then differentially
expressed genes were identified using the using quasi-likelihood F-tests, which is more robust
and produces more reliable error rates when the number of replicates is small.[90] We analyzed
only genes that were statistically significant using an unadjusted $p<0.05$. Model accuracy was
calculated at the number of positively regulated genes with a negative log fold change (i.e "true
positives") added to the number of negatively regulated target genes with a positive log fold
change (i.e "true negatives") divided by the total number of target genes for each sample.
Statistical significance was assessed using a one sided Fishers exact test on the number of
genes that were "true positives" and "true negatives" versus other genes.
736
7. Enrichment of TF Modules associated with Preterm Birth
Differentially expressed genes significantly associated with preterm birth were identified through
ancillary analyses of aggregated microarray data.[10] In this prior analysis, genes were
considered statistically significantly associated with preterm birth using an FDR adjusted Q
value <0.05. We performed an enrichment analysis[91] in which we considered the transcription
factors and all the genes they regulate (i.e regulons) as gene groups (analogous to predefined
pathways such as KEGG or GO), and compared if the list of differentially expressed genes were
enriched within specific regulons using one sided Fisher's exact test. We considered the 7712
target genes captured within our TRN as the background genes for analysis. We only included
TFs in our enrichment tests which regulated three or more differentially expressed genes. We
considered a TF to be a key regulator if its target genes were over-represented based on a P
value <0.05.
749
750   All code is publicly available in the following github repository
751   (https://github.com/AlisonPaquette/PlacentalTRN_ManuscriptCode/). All analyses were
752   performed in R (Version 3.6.1), and data was visualized using Cytoscape (Version 3.7.2). The
753   TRN was developed using the R Shiny Web Application.
754
755   **ACKNOWLEGMENTS**
We would like to acknowledge all the cohorts and biorepositories that provided placental tissue
samples, including participants and research staff and investigators who made this possible**.**
The samples from Magee Women's Research Institute were collected by the Perinatal Biology
Lab (PI, YS) from specimens by the Obstetrical Specimens Procurement Unit at Magee
Women's Hospital, based on published protocols, detailed in methods. The authors wish to
thank our ECHO colleagues, the medical, nursing and program staff, as well as the children and
families participating in the ECHO cohorts. We also acknowledge the contributions of the
following ECHO program collaborators: 1. Coordinating Center: Duke Clinical Research
Institute, Durham, North Carolina: Smith PB, Newby KL, Benjamin DK. 2. Data Analysis Center:

Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland: Jacobson
LP; Research Triangle Institute, Durham, North Carolina: Parker CB.

Disclaimer: Current Address for Dr. Brockway is the Center for Scientific Review, National
Institutes of Health, Bethesda Maryland, United States. This article was prepared while Dr.
Heather Brockway was employed by the University of Florida. The opinions expressed in this
article are the author's own and do not reflect the view of the National Institutes of Health, the
Department of Health and Human Services, or the United States government.

Funding:
This project was supported by the NICHD Human Placenta Project (R01HD091527), an NICHD
Pathway to Independence award (K99/R00 HD096112), and an ECHO Opportunity and
Infrastructure grant (OIF 0035). The Conditions Affecting Neurocognitive Development and
Learning in Early Childhood (CANDLE) study was funded by the Urban Child Institute and NIH
(R01 HL109977). ECHO PATHWAYS is funded by NIH (1UG3OD023271-01, 4UH3OD023271-
03). The University of Washington EDGE center is supported by the NIH (P30ES007033).

Author contributions:
        Conceptualization: AGP, LM, LM, SS, NP
        Methodology: YMW, PS, JP, CF, JM, TB, PB
        Software: PS, CF, YMW, RR
        Investigation: KA, YMW, JP, HL, MB, LK, SH, HB
        Resources: SS, CK, WAM, NB, KL, YS, JS
        Visualization: AGP, RR
        Supervision: NP, SS, LM, YS, HJ, LM
        Writing—original draft: AGP, NP
        Writing—review & editing: AGP, KA,YMW, JP, HL, PS, MB, LK, SL, RR,
    CF, JM, TB, PB, HB, WAM, NB, KJ, CK, JS, LM, HJ, YS, LM, SS, NP

Competing interests:
YS is on the external advisory board for Illumina. NDP serves as a scientific advisor to
Sera Prognostics, a pregnancy diagnostics company, and has stock options.

WORKS CITED

1. Burton, G. J. & Jauniaux, E. What is the placenta? *Am. J. Obstet. Gynecol.* **213**, S6.e1, S6-

    8 (2015).

2. Mesiano, S. CHAPTER 11 - The Endocrinology of Human Pregnancy and Fetoplacental

    Neuroendocrine Development A2 - Strauss, Jerome F. in *Yen & Jaffe's Reproductive*

    Endocrinology (Sixth Edition) (ed. Barbieri, R. L.) 249–281 (W.B. Saunders, 2009).

    doi:10.1016/B978-1-4160-4907-4.00011-5.

3. Myatt, L. Placental adaptive responses and fetal programming. *J. Physiol.* **572**, 25–30

    (2006).

4. Wang, Y. & Zhao, S. *Vascular Biology of the Placenta*. (Morgan & Claypool Life Sciences,

2010). doi:10.4199/C00016ED1V01Y201008ISP009.

5. Saben, J. *et al.* A Comprehensive Analysis of the Human Placenta Transcriptome. *Placenta*

35, 125–131 (2014).

6. Delahaye, F. *et al.* Genetic variants influence on the placenta regulatory landscape. *PLOS*

Genet. **14**, e1007785 (2018).

7. Lim, Y. C. *et al.* A complex association between DNA methylation and gene expression in

human placenta at first and third trimesters. *PloS One* **12**, e0181155–e0181155 (2017).

8. Sitras, V., Fenton, C., Paulssen, R., Vårtun, Å. & Acharya, G. Differences in Gene

Expression between First and Third Trimester Human Placenta: A Microarray Study. *PLOS*

ONE **7**, e33294 (2012).

9. Cox, B., Leavey, K., Nosi, U., Wong, F. & Kingdom, J. Placental transcriptome in

development and pathology: expression, function, and methods of analysis. *Am. J. Obstet.*

Gynecol. **213**, S138-151 (2015).

10. Paquette, A. G., Brockway, H. M., Price, N. D. & Muglia, L. J. Comparative transcriptomic

analysis of human placentae at term and preterm delivery. *Biol. Reprod.* **98**, 89–101 (2018).

11. Eidem, H. R., Ackerman, W. E., McGary, K. L., Abbot, P. & Rokas, A. Gestational tissue

transcriptomics in term and preterm human pregnancies: a systematic review and meta-

analysis. *BMC Med. Genomics* **8**, 27 (2015).

12. Leavey, K., Bainbridge, S. A. & Cox, B. J. Large Scale Aggregate Microarray Analysis

Reveals Three Distinct Molecular Subclasses of Human Preeclampsia. *PLoS ONE* **10**,

(2015).

13. Knöfler, M. *et al.* Human placenta and trophoblast development: key molecular mechanisms

and model systems. *Cell. Mol. Life Sci. CMLS* **76**, 3479–3496 (2019).

14. Brent, M. R. Past Roadblocks and New Opportunities in Transcription Factor Network

Mapping. *Trends Genet. TIG* **32**, 736–750 (2016).

15. He, B. & Tan, K. Understanding transcriptional regulatory networks using computational

models. *Curr. Opin. Genet. Dev.* **37**, 101–108 (2016).

16. Baur, B., Shin, J., Zhang, S. & Roy, S. Data integration for inferring context-specific gene

regulatory networks. *Curr. Opin. Syst. Biol.* **23**, 38–46 (2020).

17. Funk, C. C. *et al.* Atlas of Transcription Factor Binding Sites from ENCODE DNase

Hypersensitivity Data across 27 Tissue Types. *Cell Rep.* **32**, 108029 (2020).

18. Pearl, J. R. *et al.* Genome-Scale Transcriptional Regulatory Network Models of Psychiatric

and Neurodegenerative Disorders. *Cell Syst.* (2019) doi:10.1016/j.cels.2019.01.002.

19. Readhead, B. *et al.* Multiscale Analysis of Independent Alzheimer's Cohorts Finds

Disruption of Molecular, Genetic, and Clinical Networks by Human Herpesvirus. *Neuron* **99**,

64-82.e7 (2018).

20. Pearl, J. R. *et al.* Genome-Scale Transcriptional Regulatory Network Models of Psychiatric

and Neurodegenerative Disorders. *Cell Syst.* (2019) doi:10.1016/j.cels.2019.01.002.

21. Ament, S. A. *et al.* Transcriptional regulatory networks underlying gene expression changes

in Huntington's disease. *Mol. Syst. Biol.* **14**, e7435–e7435 (2018).

22. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

23. Buckberry, S. *et al.* Placental transcriptome co-expression analysis reveals conserved

regulatory programs across gestation. *BMC Genomics* **18**, 10 (2017).

24. Deyssenroth, M. A. *et al.* Whole-transcriptome analysis delineates the human placenta gene

network and its associations with fetal growth. *BMC Genomics* **18**, 520 (2017).

25. Liu, Y., Ding, D., Liu, H. & Sun, X. The accessible chromatin landscape during conversion of

human embryonic stem cells to trophoblast by bone morphogenetic protein 4. *Biol. Reprod.*

96, 1267–1278 (2017).

26. Buckberry, S., Bianco-Miotto, T., Bent, S. J., Dekker, G. A. & Roberts, C. T. Integrative

transcriptome meta-analysis reveals widespread sex-biased gene expression at the human

fetal-maternal interface. *Mol. Hum. Reprod.* **20**, 810–819 (2014).

27. Lorente-Pozo, S. *et al.* Influence of Sex on Gestational Complications, Fetal-to-Neonatal

Transition, and Postnatal Adaptation. *Front. Pediatr.* **6**, 63 (2018).

28. Miraldi, E. R. *et al.* Leveraging chromatin accessibility for transcriptional regulatory network

inference in T Helper 17 Cells. *Genome Res.* **29**, 449–463 (2019).

29. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

30. Uhlen, M. *et al.* Towards a knowledge-based Human Protein Atlas. *Nat. Biotechnol.* **28**,

1248–1250 (2010).

31. Funk, C. C. *et al.* Atlas of Transcription Factor Binding Sites from ENCODE DNase

Hypersensitivity Data Across 27 Tissue Types. *bioRxiv* (2018) doi:10.1101/252023.

32. Li, J. N. *et al.* The Sp1 Transcription Factor Is Crucial for the Expression of 11β-

Hydroxysteroid Dehydrogenase Type 2 in Human Placental Trophoblasts. *J. Clin.*

Endocrinol. Metab. **96**, E899–E907 (2011).

33. Sõber, S. *et al.* Extensive shift in placental transcriptome profile in preeclampsia and

placental origin of adverse pregnancy outcomes. *Sci. Rep.* **5**, 13336–13336 (2015).

34. Adu-Gyamfi, E. A., Ding, Y.-B. & Wang, Y.-X. Regulation of placentation by the transforming

growth factor beta superfamily†. *Biol. Reprod.* **102**, 18–26 (2020).

35. Cheng, Y.-H. *et al.* Critical role for transcription factor AP-2alpha in human trophoblast

differentiation. *Physiol. Genomics* **18**, 99–107 (2004).

36. Biadasiewicz, K. *et al.* Transcription factor AP-2α promotes EGF-dependent invasion of

human trophoblast. *Endocrinology* **152**, 1458–1469 (2011).

37. Cheng, Y.-H. & Handwerger, S. AP-2alpha modulates human corticotropin-releasing

hormone gene expression in the placenta by direct protein-protein interaction. *Mol. Cell.*

Endocrinol. **191**, 127–136 (2002).

38. Li, M. & Kellems, R. E. Sp1 and Sp3 Are Important Regulators of AP-2γ Gene

Transcription1. *Biol. Reprod.* **69**, 1220–1230 (2003).

39. Gabory, A., Roseboom, T. J., Moore, T., Moore, L. G. & Junien, C. Placental contribution to

the origins of sexual dimorphism in health and diseases: sex chromosomes and epigenetics.

Biol. Sex Differ. **4**, 5–5 (2013).

40. Rosenfeld, C. S. Sex-Specific Placental Responses in Fetal Development. *Endocrinology*

156, 3422–3434 (2015).

41. Eriksson, J. G., Kajantie, E., Osmond, C., Thornburg, K. & Barker, D. J. P. Boys live

dangerously in the womb. *Am. J. Hum. Biol. Off. J. Hum. Biol. Counc.* **22**, 330–335 (2010).

42. Zeitlin, J. *et al.* Fetal sex and preterm birth: are males at greater risk? *Hum. Reprod. Oxf.*

Engl. **17**, 2762–2768 (2002).

43. Wang, J. *et al.* B cell CLL/lymphoma 6 member B inhibits hepatocellular carcinoma

metastases in vitro and in mice. *Cancer Lett.* **355**, 192–200 (2014).

44. Brockway, H. M. *et al.* Unique transcriptomic landscapes identified in idiopathic

spontaneous and infection related preterm births compared to normal term births. *PloS One*

14, e0225062–e0225062 (2019).

45. Balakrishnan, M. P. *et al.* THAP5 is a DNA-binding transcriptional repressor that is regulated

in melanoma cells during DNA damage-induced cell death. *Biochem. Biophys. Res.*

Commun. **404**, 195–200 (2011).

46. Agha, G. *et al.* Birth weight-for-gestational age is associated with DNA methylation at birth

and in childhood. *Clin. Epigenetics* **8**, 118–118 (2016).

47. Larigot, L., Juricek, L., Dairou, J. & Coumoul, X. AhR signaling pathways and regulatory

functions. *Biochim. Open* **7**, 1–9 (2018).

48. Trikha, P. & Lee, D. A. The role of AhR in transcriptional regulation of immune cell

development and function. *Biochim. Biophys. Acta BBA - Rev. Cancer* **1873**, 188335

(2020).

49. Wakx, A. *et al.* Expression, Localization, and Activity of the Aryl Hydrocarbon Receptor in

the Human Placenta. *Int. J. Mol. Sci.* **19**, 3762 (2018).

50. Wu, Y. *et al.* ITE and TCDD Differentially Regulate the Vascular Remodeling of Rat

Placenta via the Activation of AhR. *PLOS ONE* **9**, e86549 (2014).

51. Li, Y. *et al.* A possible role of aryl hydrocarbon receptor in spontaneous preterm birth. *Med.*

Hypotheses **84**, 494–497 (2015).

52. Bruner-Tran, K. L. & Osteen, K. G. Developmental exposure to TCDD reduces fertility and

negatively affects pregnancy outcomes across multiple generations. *Reprod. Toxicol.*

Elmsford N **31**, 344–350 (2011).

53. Peltier, M. R. *et al.* 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) enhances placental

inflammation. *J. Reprod. Immunol.* **98**, 10–20 (2013).

54. Wu, Y. *et al.* Potential involvement of placental AhR in unexplained recurrent spontaneous

abortion. *Reprod. Toxicol. Elmsford N* **59**, 45–52 (2016).

55. Soneji, S. & Beltrán-Sánchez, H. Association of Maternal Cigarette Smoking and Smoking

Cessation With Preterm Birth. *JAMA Netw. Open* **2**, e192514–e192514 (2019).

56. Shin, S. *et al.* NRF2 Modulates Aryl Hydrocarbon Receptor Signaling: Influence on

Adipogenesis. *Mol. Cell. Biol.* **27**, 7188 (2007).

57. Miao, W., Hu, L., Scrivens, P. J. & Batist, G. Transcriptional regulation of NF-E2 p45-related

factor (NRF2) expression by the aryl hydrocarbon receptor-xenobiotic response element

signaling pathway: direct cross-talk between phase I and II drug-metabolizing enzymes. *J.*

Biol. Chem. **280**, 20340–20348 (2005).

58. He, F., Antonucci, L. & Karin, M. NRF2 as a regulator of cell metabolism and inflammation in

cancer. *Carcinogenesis* **41**, 405–416 (2020).

59. Hussain, T. *et al.* Modulatory Mechanism of Polyphenols and Nrf2 Signaling Pathway in LPS

Challenged Pregnancy Disorders. *Oxid. Med. Cell. Longev.* **2017**, 8254289 (2017).

60. Kweider, N. *et al.* The effects of Nrf2 deletion on placental morphology and exchange

capacity in the mouse. *J. Matern.-Fetal Neonatal Med. Off. J. Eur. Assoc. Perinat. Med.*

Fed. Asia Ocean. Perinat. Soc. Int. Soc. Perinat. Obstet. **30**, 2068–2073 (2017).

61. Sussan, T. E. *et al.* Nrf2 regulates gene-environment interactions in an animal model of

intrauterine inflammation: Implications for preterm birth and prematurity. *Sci. Rep.* **7**, 40194

(2017).

62. Everson, T. M. & Marsit, C. J. Integrating -Omics Approaches into Human Population-Based

Studies of Prenatal and Early-Life Exposures. *Curr. Environ. Health Rep.* (2018)

doi:10.1007/s40572-018-0204-1.

63. Lapehn, S. & Paquette, A. G. The Placental Epigenome as a Molecular Link Between

Prenatal Exposures and Fetal Health Outcomes Through the DOHaD Hypothesis. *Curr.*

Environ. Health Rep. (2022) doi:10.1007/s40572-022-00354-8.

64. Lachmann, A. *et al.* ChEA: transcription factor regulation inferred from integrating genome-

wide ChIP-X experiments. *Bioinforma. Oxf. Engl.* **26**, 2438–2444 (2010).

65. Buccitelli, C. & Selbach, M. mRNAs, proteins and the emerging principles of gene

expression control. *Nat. Rev. Genet.* **21**, 630–644 (2020).

66. Sung, J., Wang, Y., Chandrasekaran, S., Witten, D. M. & Price, N. D. Molecular signatures

from omics data: From chaos to consensus. *Biotechnol. J.* **7**, 946–957 (2012).

67. Chicco, D. Ten quick tips for machine learning in computational biology. *BioData Min.* **10**, 35

(2017).

68. Wyatt, S. M. *et al.* The correlation between sampling site and gene expression in the term

human placenta. *Placenta* **26**, 372–379 (2005).

69. Paquette Alison G. *et al.* A Comprehensive Assessment of Associations between Prenatal

Phthalate Exposure and the Placental Transcriptomic Landscape. *Environ. Health Perspect.*

129, 097003.

70. Eis, A. L., Brockman, D. E., Pollock, J. S. & Myatt, L. Immunohistochemical localization of

endothelial nitric oxide synthase in human villous and extravillous trophoblast populations

and expression during syncytiotrophoblast formation in vitro. *Placenta* **16**, 113–126 (1995).

71. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

quantification. *Nat Biotech* **34**, 525–527 (2016).

72. Brown, J., Pirrung, M. & McCue, L. A. FQC Dashboard: integrates FastQC results into a

web-based, interactive, and extensible FASTQ quality control tool. *Bioinformatics* **33**, 3137–

3139 (2017).

73. Soneson, C., Love, M. & Robinson, M. Differential analyses for RNA-seq: transcript-level

estimates improve gene-level inferences [version 1; referees: 2 approved]. *F1000Research*

4, (2015).

74. Chen, Y., Lun, A. & Smyth, G. From reads to genes to pathways: differential expression

analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline

[version 2; peer review: 5 approved]. *F1000Research* **5**, (2016).

75. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biol.* **11**, R25 (2010).

76. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights unlock linear model

analysis tools for RNA-seq read counts. *Genome Biol.* **15**, 1–17 (2014).

77. Leek, J. T. & Storey, J. D. Capturing Heterogeneity in Gene Expression Studies by

Surrogate Variable Analysis. *PLOS Genet.* **3**, e161 (2007).

78. Lachmann, A. *et al.* Massive mining of publicly available RNA-seq data from human and

mouse. *Nat. Commun.* **9**, 1366 (2018).

79. Consortium, T. E. P. An integrated encyclopedia of DNA elements in the human genome.

Nature **489**, 57–74 (2012).

80. Boyle, A. P., Guinney, J., Crawford, G. E. & Furey, T. S. F-Seq: a feature density estimator

for high-throughput sequence tags. *Bioinformatics* **24**, 2537–2538 (2008).

81. Koohy, H., Down, T. A., Spivakov, M. & Hubbard, T. A Comparison of Peak Callers Used for

DNase-Seq Data. *PLOS ONE* **9**, e96303 (2014).

82. Gusmao, E. G., Dieterich, C., Zenke, M. & Costa, I. G. Detection of active transcription

factor binding sites with the combination of DNase hypersensitivity and histone

modifications. *Bioinformatics* **30**, 3143–3151 (2014).

83. Khan, A. *et al.* JASPAR 2018: update of the open-access database of transcription factor

binding profiles and its web framework. *Nucleic Acids Res.* **46**, D260–D266 (2018).

84. Kulakovskiy, I. V. *et al.* HOCOMOCO: towards a complete collection of transcription factor

binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids*

Res. **46**, D252–D259 (2018).

85. Pachkov, M., Erb, I., Molina, N. & van Nimwegen, E. SwissRegulon: a database of genome-

wide annotations of regulatory sites. *Nucleic Acids Res.* **35**, D127–D131 (2007).

86. Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif.

Bioinforma. Oxf. Engl. **27**, 1017–1018 (2011).

87. Paul Shannon. *MotifDb: An Annotated Collection of Protein-DNA Binding Sequence Motifs.*

(2017).

88. Plaisier, C. L. *et al.* Causal Mechanistic Regulatory Network for Glioblastoma Deciphered

Using Systems Genetics Network Analysis. *Cell Syst.* **3**, 172–186 (2016).

89. Fishilevich, S. *et al.* GeneHancer: genome-wide integration of enhancers and target genes

in GeneCards. *Database J. Biol. Databases Curation* **2017**, bax028 (2017).

90. Lun, A. T. L., Chen, Y. & Smyth, G. K. It's DE-licious: A Recipe for Differential Expression

Analyses of RNA-seq Experiments Using Quasi-Likelihood Methods in edgeR. *Methods*

Mol. Biol. Clifton NJ **1418**, 391–416 (2016).

91. Falcon, S. & Gentleman, R. Hypergeometric Testing Used for Gene Set Enrichment

Analysis. in *Bioconductor Case Studies* (eds. Hahne, F., Huber, W., Gentleman, R. &

Falcon, S.) 207–220 (Springer New York, 2008). doi:10.1007/978-0-387-77240-0_14.

1011
1012

TABLES

Table 1: Covariate Data from Cohorts used for RNA sequencing and DNase hypersensitivity Data

	MWH (N=104)-RNA Sequencing				CANDLE (N=371) -RNA Sequencing				OHSU (N=12)-DNAse Seq			
	N (%)	Range	Median	SD	N (%)	Range	Median	SD	N (%)	Range	Median	SD
B. Categorical Variables												
Delivery Method												
C-Section	29 (28%)				140 (38%)				12 (100%)			
Vaginal	75 (72%)				231 (62%)							
Fetal Sex												
Female	52 (50%)				195 (53%)				6 (50%)			
Male	52 (50%)				176 (47%)				6 (50%)			
Maternal Race												
Black	11 (10%)				209 (56%)							
Other	5 (5%)				5 (1%)							
White	88 (85%)				157 (42%)				12 (100%)			
B. Continuous Variables												
Gestational age (weeks)		37-41.5	39.6	0.89		34.2-41.3	39.2	1.13		38.3-39.6	39.1	0.33

Table 2: TFs whose placental expression was previously associated with Preterm Birth

TF	TRN Results				Microarray Results[2]	
	Number of Target Genes	**Number of PTB Genes**	**Genes Previously associated with preterm Birth from microarray[2]**	**P[1](Fishers Exact Test)**	**Log Fold Change**	**FDR Adusted P**
ATF1	734	9	*DNAH2, FAM122A, TRIM45, EPC1, ZBTB44, TADA1, TC2N, SGMS2, RNF11*	0.88	-0.160	0.011
CLOCK	1345	11	*RNASEH2C, DAAM1, FIGN, ZBTB44, PPP3CB, RIC1, FBXW11, RASAL2, ARID2, SPOPL, PHIP*	1.00	-0.143	0.042
HLF	175	5	*ERAP1, SLC25A19, ACOXL, TMEM136, SLC35A5*	0.17	-0.270	0.011
IRF2	5	0		1.00	-0.119	0.017
MLX	8	0		1.00	0.139	0.004
ZBTB18	148	1	*ADAMTS3*	0.92	-0.144	0.035
	1. From Fishers Exact Test. 3Log fold change: Preterm vs. Term Samples; FDR adjusted Q, From Paquette, Brockway et al, 2018					

FIGURES

Figure 1: Overview of data and strategy used to construct TRN. First, regions of open chromatin i.e. footprints are identified from Placental DNAse hypersensitivity data (A). Concurrently, TF binding motifs were curated from publicly available databases (B). This data was overlayed (C) to identify putative regulators (TFs) of each target gene. This putative model is pruned using correlations between TFs and target genes established in the training data, and 6 models with different parameters were used. These models were tested in our test data to evaluate the best model (D). The final model was tested in our holdout data (E) to generate final accuracy for each gene and for the overall model

A. Distribution of TF-Target Gene correlations   **B.** Prediction accuracy of TFs using LASSO Regression

Figure 2: (A.) Distribution of pearson correlation coefficients in training + testing dataset (N=382 samples, pink), and holdout dataset(N=93, blue). Red lines indicate 0.25, which was the cutoff in the training dataset. (B.) Distribution of out of sample accuracy (OOS R^2) for each of the 10,984 genes calculated in our holdout dataset. Light blue box indicates genes with OOS R^2 <0.25, which were genes not included in our final model. Dark blue genes were high confidence OOS R^2.

Figure 3: (A.) Summary of network connectivity of 393 TFs based on their target genes. Orange line indicates hub TFs with >500 genes. Top 15 TFs based on degree centrality are shown in green.

Figure 4: Experimental validation for assessment of final model accuracy. Results of Experimental validation for (A) _GRHL2_, (B) _CREB3L2_, (C) _ESRRG_ and (D) _GCM1_. X axis represents the log fold change of knockout vs control, and the Y axis represents correlation between TF and target gene. Model accuracy was calculated as the number of positively regulated genes with a negative log fold change (i.e "true positives", shown in red) added to the number of negatively regulated target genes with a positive log fold change (i.e "true negatives", shown in blue) divided by the total number of target genes for each sample. Genes that were not concordant are shaded in grey. The size and shading of each target gene represents the relative rank (1-15) of the TF as a regulator in our TRN.

Figure 5: Plot depicting the 5361 sex specific interactions in the placental TRN, divided into four distinct interaction categories. The x-axis represents TF-target gene estimate values from linear regression analysis in the whole dataset. The y-axis represents the difference in estimates between male and female samples.

Figure 6: TRN Subnetwork depicting enriched TFs (P<0.1; Fishers Exact Test) (Black) and their corresponding genes that were previously associated with PTB. Color of gene indicates directionality of log fold change in association with preterm birth. Placental specific TFs, defined by the Human Protein Atlas[30] are outlined in green. Color of edge indicates correlation between TF and target gene, and number represents correlation coefficient. Transcriptional regulation between enriched TFs is highlighted by red dotted edges.