Adjusting for genetic confounders in transcriptome-wide association studies leads to reliable detection of causal genes

September 27, 2022

Siming Zhao[1*#], Wesley Crouse[2*], Sheng Qian[2], Kaixuan Luo[2], Matthew Stephens[2,3#] and Xin He[2#]

1. Department of Biomedical Data Science, Dartmouth Cancer Center, Dartmouth College, Hanover, NH 03755, USA

2. Department of Human Genetics, University of Chicago, Chicago, IL 60637, USA

3. Department of Statistics, University of Chicago, Chicago, IL 60637, USA

* Co-first authors
# Co-corresponding authors
Xin He (xinhe@uchicago.edu)
Matthew Stephens (mstephens@uchicago.edu)
Siming Zhao (siming.zhao@dartmouth.edu)

Abstract

Expression QTLs (eQTLs), provide valuable information on the functional effects of variants. Many methods have been developed to leverage eQTLs to nominate candidate genes of complex traits. These methods include colocalization analysis, transcriptome-wide association studies (TWAS) that correlate genetic components of expression with traits, and Mendelian Randomization (MR)-based methods. A fundamental problem of all these methods is that, when assessing the role of one gene in a trait using its eQTLs, nearby variants and nearby genetic components of expression of other genes can be correlated with the eQTLs of the test gene, while affecting the trait directly. These "genetic confounders" may lead to false discoveries. To address this challenge, we introduced a novel statistical framework that generalizes TWAS to adjust all genetic confounders. In our simulations, we found that existing methods based on TWAS, colocalization or MR all suffered from high false positive rates, often greater than 50%. Our method, causal-TWAS (cTWAS), in contrast, showed calibrated false positive rates while maintaining power. Application of cTWAS on several common traits highlighted the weakness of existing methods and discovered novel candidate genes. In conclusion, cTWAS is a robust statistical framework to integrate eQTL and GWAS data for gene discovery, and is extendable to other molecular QTLs.

Introduction

Genome-wide association studies (GWAS) have successfully identified a large number of loci associated with a range of complex human traits [1] [2]. It remains difficult, however, to translate these associations into knowledge of causal variants, their target genes and molecular mechanisms [3]. A common strategy to address this challenge is the use of expression Quantitative Trait Loci (eQTL) data to link variants with gene expression. Transcriptome-wide association studies (TWAS) provides a formal framework for the joint analysis of eQTL and GWAS data [4, 5]. In TWAS, researchers build predictive models of gene expression from genetic variants, usually in *cis*, and then test for associations between predicted ("imputed") expression and traits of interest. As such, TWAS identifies candidate genes for a trait, as well as the likely relevant cell/tissue contexts. TWAS requires only summary statistics, making it easy to integrate eQTL and GWAS datasets from different cohorts. Because of these benefits, TWAS has become widely used to convert GWAS associations into candidate genes [6]. TWAS is also applicable to any molecular traits, such as RNA splicing, or chromatin features, further broadening its utility.

A central question in TWAS analysis is whether the identified genes necessarily have causal effects on the phenotype. A simple analysis suggests this is not always the case (Fig. 1A). In one scenario, a non-causal gene X, has an eQTL G, that is in linkage disequilibrium (LD) with the eQTL of a nearby causal gene X'. This creates a non-causal association of the genetic component of X with the trait. In another scenario, G is in LD with a nearby causal variant, G', which acts on the trait directly, again creating a non-causal association of the genetic component of X with the trait. In all these cases, false positive findings result from association of G with the trait, through pathways other than the gene X. This is known as "horizontal pleiotropy" in the literature, and is a key challenge that reduces the utility of TWAS [6].

Alternative methods have been proposed to jointly analyze eQTL and GWAS data, but they all face similar challenges. Colocalization methods test whether gene expression and a trait are affected by the same causal variant [7, 8]. Colocalization may alleviate the problems with TWAS, but does not fully resolve them. Indeed, in both scenarios described in Fig. 1A, if LD between the eQTL variant, G, and the nearby causal variant, G', is strong enough, then G and G' are effectively indistinguishable as potential causal variants of both gene expression and phenotype, leading to a false positive finding by colocalization [9]. False positives may also occur if the eQTL variant G has direct (pleiotropic) effects on the phenotype not through X [9]. Mendelian Randomization (MR) is another strategy to nominate causal genes, treating eQTLs of a gene as instrumental variables (IV) to infer its causal effect [10]. The scenarios described above, including pleiotropic effects of instruments and their LD with causal variants, however, violate the key assumption of MR, leading to potential false positive findings.

Multiple lines of evidence suggest that the scenarios creating possible false positive findings are likely common. First, in TWAS, it is common to observe associations of multiple genes with a trait at a single locus, with most associations likely resulting from non-causal genes. Several examples were discussed in a perspective article on TWAS [6]. Colocalization analysis has a similar problem [8]. For example, in a Schizophrenia study, 282 genes in 78 risk loci, showed evidence of colocalization [11]. Secondly, at a biochemical level, co-regulation of genes by the same regulatory elements is common. For example, one study estimated that each enhancer regulates on average 2.8 genes [12]. Thirdly, eQTLs are pervasive in the genome. In GTEx samples, half of all common variants are eQTLs in at least one tissue [13]. The implication is that chance association (LD) between eQTLs of non-causal genes and causal variants may be common [14]. Finally, while eQTLs

are often associated with phenotypes, they mediate only 10-20% of heritability of complex traits [15]. This suggests that associations of eQTLs and a trait are often not due to causal effects of gene expression on the trait, but due to LD. All this evidence points to a critical need for better control of false discoveries in TWAS and other eQTL-based analysis.

Here we propose a new statistical framework to address the limitations of existing eQTL-based methods. Our approach can be viewed as a generalization of TWAS, which we term "causal-TWAS" (cTWAS). The fundamental problem of TWAS is that the imputed expression of any gene (the "focal gene") is often correlated with expression of nearby genes and with other nearby variants (Fig. 1A). If these nearby genes and variants affect the trait, they induce a correlation between the trait and imputed expression of the focal gene. That is, they act as "confounders", when testing association of the focal gene and the trait. We refer to these confounders as "genetic confounders" to distinguish them from the environmental confounders that are a common focus in the literature of causal inference and MR.

This reasoning suggests a conceptually simple solution: when assessing association between the focal gene and trait through a regression model, we should include all genetic confounders in the model. In practice, implementing this strategy is complicated by high correlations that often exist among all these variables, which creates an identifiability challenge. Our key intuition is that causal signals in a genomic region, whether via gene expression or variants, are likely sparse. This motivates a Bayesian variable selection model, which has been widely used in statistical fine-mapping [16, 17, 18, 19]. Our approach, cTWAS, generalizes standard fine-mapping methods by including imputed expression levels and genetic variants in the same regression model. In simulations and applications to real data, cTWAS greatly reduces the number of false discoveries from TWAS, colocalization and MR-based methods, laying a foundation for reliable discovery of causal genes from GWAS.

Results

Overview of the cTWAS model

We start with a formal description of standard TWAS, using the language of causal graphs [20]. We assume genetic variants, denoted as G, affect expression of a gene, X, which affects a trait Y (Fig. 1B). Both X and Y could be affected by unobserved environmental variable(s) U, such as diet. We introduce $\tilde{X}$, to denote the *cis*-genetic component of X. Importantly, the genetic variants G act on Y only through $\tilde{X}$. Under this model, the regression coefficient of $\tilde{X}$ (with Y as the response) as computed by TWAS, would give the causal effect of X on the trait. The confounder U is not a concern here because U contributes only to the non-genetic component of X, and thus does not create an association between $\tilde{X}$ and Y. Using the language of causal graphs, the potentially non-causal path from $\tilde{X}$ to Y through U, has a collider, X, which blocks the association. Following similar analysis, we can see that TWAS results are also robust to "reverse causality" where Y affects X (Fig. S1). In such a case, X is a collider in the paths from $\tilde{X}$ to Y, making the two uncorrelated.

Unfortunately, the key assumption underlying TWAS, that G is not associated with Y through other paths, is often violated. In the example discussed (Fig. 1A), $\tilde{X}$ may become correlated with a nearby variant G', or the genetic component of a nearby gene X'. These confounders would create non-causal associations, technically known as backdoor paths, between $\tilde{X}$ and Y, leading to possible false discoveries by TWAS (Fig. S1).

To control for all potential confounders, cTWAS generalizes the standard TWAS. The key idea

is to jointly model the dependence of phenotype on *all* imputed genes, and *all* variants, with their effect sizes denoted as β_j for genes and θ_m for variants, respectively (Fig. 1C). Joint estimation of all these parameters would then lead to causal effect estimates. In practice, to simplify computation, we assume that the genome can be partitioned into disjoint blocks, with imputed expression and variants independent across blocks, and perform the analysis block-by-block.

Inference in the cTWAS model presents a new challenge: the high correlations among the variables. To address this challenge we assume that in any genomic region causal effects are sparse: that is, there are likely a small number of causal effects, whether genes or variants. The analysis then becomes similar to standard fine-mapping, where one aims to identify a small number of likely causal variants among many correlated variants. An additional intuition helps explain that the model can potentially learn gene effects despite collinearity. While most variants are non-functional, gene expression traits should be more likely to have causal effects *a priori*. Also, a causal gene may have multiple eQTLs, each of which would be associated with the trait. Thus, a single gene effect would be a more parsimonious explanation of data, compared with several independent variant effects.

We thus fit our cTWAS model using the statistical machinery developed for fine-mapping. We assume sparse prior distributions of the gene and variant effects (Fig. 1C), with an Empirical Bayes strategy to estimate these prior parameters from data. Learning different prior parameters for genes vs variants allows the model to favor genes over variants when identifying possible causal effects. With the estimated prior parameters, we infer likely causal genes and variants in each block, using SuSiE, a state-of-art method for fine-mapping [16] [17]. The final results of cTWAS are expressed as Posterior Inclusion Probabilities (PIPs) for genes and variants, representing the probabilities that genes or variants have non-zero effects (Fig. 1D). While cTWAS is formulated in terms of individual level data, we have also derived the summary statistics-based model. This involves an extra step of computing correlations between imputed genes, and between imputed genes and variants (Fig. 1D).

The cTWAS model provided a general framework for joint analysis of eQTL and GWAS data. It generalizes and unifies a number of existing methods (see Discussion). We show that under a simple scenario where genes have only single eQTL variants, cTWAS reduces to existing colocalization methods (Supplemental Text). cTWAS can also be viewed as a MR method, where cis-genetic expression is used as Instrumental Variable, but it accommodates horizontal pleiotropy through the inclusion of effects from variants and other genes. Lastly, while our primary goal is gene discovery, the learned prior parameters for gene and variant effects also allow us to estimate the proportion of heritability attributable to gene expression vs. variants, informing the genetic architecture of the traits. This application of cTWAS is related to several other methods [15] [21] [22].

cTWAS controls false discoveries in simulation studies

We created simulation data to assess the performance of cTWAS and illustrate its benefits over existing methods. We used genotype data of common SNPs (minor allele frequency > 0.05) from ~ 45k samples with white British ancestry from the UK biobank [23], and imputed gene expression using the prediction models from GTEx by FUSION [5] (see Methods). We varied prior probabilities for genes and SNPs being causal, and prior effect size variances, with a total of 10 settings. We focused on two representative settings in the main results here, where the trait heritability is set at 0.5, and the percent of heritability explained by gene expression set at 10% (high gene PVE setting, where PVE stands for "percent of variance explained"), or 4% (low gene PVE). With these prior parameters, we sample causal genes, causal variants and their effect sizes, and then simulate traits

according to our model (Fig. 1C). When running cTWAS, we computed LD matrices using 2000 UK biobank samples and used summary statistics of SNP-trait associations, and the expression models as input.

We first assessed the accuracy of parameter estimation. cTWAS estimated parameters were generally close to true values (gene results in Fig. 2A, and variant results in Fig. S2). In practice, what matters most is the enrichment of causal genes vs variant effects, defined as the ratio of prior probability of non-zero gene effects to that of non-zero variant effects. The enrichment value determines the extent to which the model favors gene effects vs. variant effects. Although cTWAS slightly under-estimated some prior parameters under some settings, the estimated enrichment remains accurate (Fig. 2A). Finally, we found that cTWAS accurately estimated the proportion of trait variance explained by the gene effects (Fig. 2A).

We next assessed whether PIPs of genes computed by cTWAS are calibrated. Good calibration means that at a PIP threshold, say 0.9, we would expect at least 90% of genes above the threshold to be causal genes. Indeed, the observed proportions of causal genes from simulations were close to the expected proportions (Fig. 2B and Fig. S2). Calibration is especially good at the high PIP range (90% or higher), which is what matters most in practice.

We found cTWAS successfully removed a large number of non-causal genes with highly significant associations in standard TWAS (Fig. 3A). We systematically compared the performance of cTWAS with other methods, including the standard TWAS implemented by FUSION [5], colocalization method (coloc) [24], a MR-based method (SMR) [10], and FOCUS [25], a multi-gene analysis method. We also included the results from SMR with an additional filter HEIDI [10]. The results of all methods, except cTWAS, were severely inflated, with the majority of genes found being false positives (Fig. 3B). Importantly, cTWAS controls false positive rates with only a modest reduction of the number of discovered causal genes (Fig. 3B).

We illustrated, with two examples, how cTWAS was able to remove false positives. In the first example, the region has a single causal effect in Gene 1. However, two non-causal genes (Genes 2 and 3) also showed strong associations with the trait because of LD between their eQTLs and Gene 1's eQTLs (Fig. 3C, top). cTWAS correctly identified Gene 1 as the true signal in this region, and assigned low PIPs to the two other genes (Fig. 3C, bottom). Interestingly, coloc assigned high probability of colocalization to the non-causal Gene 3 (coloc PP4 = 0.995). In the second example, the causal signal in the region is a SNP variant, but this SNP is in LD with the eQTL of Gene 1, creating a significant association of Gene 1 with the trait (Fig. 3D, top). By including both SNPs and genes in the model, cTWAS was able to correctly identify the SNP effect as the causal signal and assigned low PIP for Gene 1 (Fig. 3D, bottom). Coloc again gave high proprobality of colocalization to Gene 1 (PP4 = 0.8).

Finally, we investigated whether cTWAS is robust to model mis-specification. We simulate traits with effect sizes for causal genes and SNPs following mixtures of normal distributions. The mixture distributions better capture the "long tails" of effect size distributions, i.e. some genes or SNPs have especially large effect sizes. Despite the mismatch of the prior used in cTWAS, the gene effect enrichment was accurately estimated and PIPs were well calibrated across different settings (Fig. S3).

cTWAS is accurate in identifying causal genes of LDL

We applied cTWAS to GWAS summary statistics of LDL cholesterol from the UK Biobank [**rapid**], and imputed expression using eQTL weights for GTEx liver tissue from PredictDB [4]

[26] [27]. After harmonizing eQTL weights with the UK Biobank in-sample LD reference panel (see Methods), we computed TWAS z-scores for 9,881 protein-coding genes. We also obtained association z-scores of ≈ 8.7 million biallelic variants with MAF>0.01. Using these z-scores and LD panel as input, cTWAS estimated that genes were over 62 times more likely than variants to be causal for LDL *a priori* (Fig. S4A). Genetic variants and imputed expression together explained 5.6% of variation of LDL (total heritability), 22.7% of which was attributable to expression. These estimates are in line with the 8.3% estimate for total heritability using LD score regression [28] and 33.5% of mediated heritability through expression using MESC [15]. The somewhat lower estimates of heritability from cTWAS likely resulted from the assumption of sparse causal effects.

Using the estimated prior parameters, cTWAS identified 35 potentially-causal liver genes (PIP > 0.8; Table S1). In contrast, standard TWAS identified 215 genes at a Bonferonni-corrected significance threshold of 0.05. To assess these results, we used 69 previously-identified LDL-related genes as "silver standard" [29] [30] (see Methods). Following earlier work [31], we assessed the performance of TWAS and cTWAS by their ability to distinguish silver standard genes from other genes within 1Mb, referred to as "bystanders", most of which are probably false positives. We limited our analysis to 46 imputable genes out of 69 silver standard genes and 539 imputed bystander genes (Table S2). Among these genes, eight have PIP > 0.8 in cTWAS, and six of them are in the silver standard, leading to a precision of 75% (Fig. 4A). Standard TWAS, on the other hand, has a precision of 31% (19 out of 61).

To illustrate how cTWAS avoids false positives, we examined two cases in detail. We first highlight the gene *HPR* in a genome-wide significant locus of LDL (Fig. 4B). This locus contains six genes significantly associated with LDL by TWAS, including *HPR* and five other genes (*ATXN1L*, *ZNF821*, *PHLPP2*, *PKD1L3*). In contrast, cTWAS identified a single candidate gene, *HPR* (PIP = 1.000) while giving no evidence (PIP < 0.01) to all other genes. This can be explained by the fact that once controlling for *HPR*, cTWAS found no additional signal in other genes. Literature evidence suggests that *HPR*, haptoglobin-related protein that binds hemoglobin and apolipoprotein-L [32], is the likely causal gene at this locus. This example thus illustrates that cTWAS avoids false positives due to confounding with nearby gene expression.

In the second case, we examined a strong GWAS locus of LDL containing the gene *POLK* (DNA polymerase kappa) (Fig. 4C). Three genes in this region show significant TWAS (Bonferroni) z-scores: *POLK, ANKDD1B,* and *POC5.* A recent method based on TWAS and Mendelian Randomization, MR-JTI [30], has highlighted *POLK* as the potential causal gene at this locus, and proposed a connection between DNA repair and regulation of LDL. The strength of associations of the three genes, however, is much lower than some nearby variants in the region (Fig. 4C). This pattern is reflected in the results of cTWAS. cTWAS selected several highly associated variants as causal signals, while giving little evidence to all three genes. This suggests that the associations of the genes with LDL are due to their correlations with true causal variants, which may act through other pathways.

To better understand these results, we inspected variant-level fine-mapping results of this locus using SuSiE [33]. SuSiE identified two causal signals in the region, with the variants in the credible set all inside or close to the gene *HMGCR*, whose expression was not imputable in our data (Fig. 4D). All the credible set variants are far from the three TWAS genes, including POLK (>200 Kb). Additionally, promoter-capture Hi-C (PC-HiC) data and the Activity-by-contact (ABC) score in liver provided no evidence linking these variants to *POLK*. Instead, the top SNP in the locus, rs12916 (PIP = 0.99) is within the 3' UTR of *HMGCR*, and 1310 bp away from one end of a chromatin

loop interacting with the HMGCR promoter, in both PCHiC and ABC (Fig. 4D). Consistent with these results, HMGCR is the rate-limiting enzyme for cholesterol synthesis and the target of statin, a key drug for reducing LDL levels [34]. All the evidence thus points to *HMGCR*, instead of *POLK* or other associated genes, as the causal gene in this region. This example demonstrates that by controlling nearby genetic variants, cTWAS is able to avoid false positive genes.

We systematically evaluated the sources of possible false positive findings from standard TWAS. We call a gene a likely false positive if it is significant under TWAS (Bonferroni threshold), but PIP < 0.5 under cTWAS. These cases are classified as "confounding by genes" or "confounding by variants", depending on whether the low PIPs of these genes were primarily driven by nearby genes or variants (see Methods). Among 170 likely false positive TWAS genes, we analyzed the 83 of these genes that can be assigned to a credible set. The majority of these genes (74.7%, Fig. 4E) were driven by confounding variants. These results show that the greatest risk of TWAS is not co-regulation of genes, i.e. shared eQTLs among nearby genes, but rather the correlation of genes with nearby variants whose effects are not manifested as eQTLs.

To seek new insights into the genetics of LDL we evaluated the functions of 35 genes with cTWAS PIP > 0.8 (Fig. 5A). We first confirmed that a large proportion of these genes represent novel findings. Only six of these genes were in the curated silver standard genes, and 20 of 35 genes were not the nearest genes of GWAS lead variants in LDL-associated loci (Fig. 5A). The 35 genes were enriched for 28 Gene Ontology Biological Process terms (FDR < 0.05), many of which are cholesterol-related (Fig. 5B, 13 non-redundant terms shown; Table S3). To assess the novelty of these GO terms, we performed GO enrichment analysis of silver standard genes (Table S4) and GWAS gene set analysis using MAGMA (Table S5). Several GO terms enriched in cTWAS genes were not found in GO terms from the silver standard or MAGMA analysis, including "peptidyl-serine phosphorylation" and "activin receptor signaling pathway". These terms suggest the importance of signal transduction in LDL regulation. Activin signaling, in particular, regulates metabolic processes including lipolysis and energy homeostasis [35, 36]. The genes associated with the two terms include well-known LDL genes, such as *CSNK1G3, TNKS,* and *GAS6,* as well as some novel and promising genes such as *ACVR1C*, an activin receptor, and *PRKD2* (Fig. 5C; Fig. S4B; see Discussion). In the cases of both *ACVR1C* and *PRKD2*, no nearby variant reaches genome-wide significance, thus the genes would be missed by standard GWAS analysis.

While cTWAS reduces false positives and identifies promising LDL candidate genes, its power seems relatively low. Out of 69 silver standard genes, 6 (8.7%) have PIP > 0.8 under cTWAS. To understand why, we categorized the outcome of cTWAS for all 69 silver standard genes, binning genes into categories such as "insignificant TWAS z-score", or "not imputed" (Fig. 5D). Many silver standard genes had no significant GWAS association signals nearby (26.1%, 18 of 69), no imputable liver expression (33.3%, 23 of 69) or insignificant TWAS z-scores (13.0%, 9 of 69). These results suggest that to improve the power of cTWAS, and eQTL-based methods in general, it is necessary to improve the power of GWAS, the power of eQTL studies, and expanding with more trait-related tissues/cell types (see Discussion).

cTWAS discovers candidate genes of several common traits

We applied cTWAS to several commonly studied traits, including Inflammatory Bowel Disease (IBD), Systolic Blood pressure (SBP) and Schizophrenia (SCZ). We used the expression prediction models of protein-coding genes from PredictDB [26, 37, 27] across 49 tissues in GTEx [38]. These models borrowed information across tissues to improve prediction accuracy in any specific tissue

[37]. The numbers of imputed protein-coding genes across tissues range from 6,591 to 11,985 (Fig. S5). We ran cTWAS analysis for a trait in each tissue separately. We summarized the results below, with a particular emphasis on IBD as a representative trait.

We first assessed the parameters learned by cTWAS. The prior probability of an imputable gene being causal ranges from 0.17% to 2.16% across tissue-trait pairs (Fig. S6, Table S6). For example, for IBD, the top tissue is whole blood, with the percent of causal genes 1.54%. The estimated proportions of heritability of traits explained by the genetic components of expression are generally small, e.g. for IBD, they range from 4% to 15% (Fig. 6A, Table S6). These estimates are generally in line with estimated values from MESC (Fig. 6A, Fig. S7, Table S7). Altogether, these results highlight the ability of cTWAS to infer important genetic parameters.

We next assessed the number of high confidence genes, at PIP > 0.8, found by cTWAS. Among the top 2 tissues per trait, cTWAS identified 13 - 26 genes (Fig. 6B). In general, the number of cTWAS genes were much smaller than those from standard TWAS (Fig. S9). For instance, for IBD, while TWAS reported 68-125 genes across 49 tissues, cTWAS identified 0-17 genes (Fig. S9). These results confirm that only a small proportion of genes associated with traits from TWAS are likely causal genes.

Given that relatively few genes were discovered in each tissue separately, we grouped related tissues into "tissue groups" and took the union of genes across all tissues within a group (Fig. 6C, Fig. S10). The top tissue groups include trait-relevant tissues, such as "Digestive" tissue for IBD (Fig. 6C), "Cardiovascular" for SBP and "Central Nervous Systems" for SCZ (Fig. S10). The number of discovered genes in the top tissue group per trait ranged from 37 (IBD) to 48 (SBP), highlighting the increased power of discovery when using data from multiple related tissues. We also assessed the novelty of the found genes. In the case of IBD, a gene is considered to be "novel" if it is not in the curated genes for IBD [39], or if it is not the nearest protein-coding gene of a lead genome-wide significant GWAS variant. Under these criteria, the majority of cTWAS genes for IBD were novel across all tissue groups (Fig. 6C).

A natural question is whether the discovered genes are limited to specific tissues. We found that most cTWAS genes were identified in a small number of tissues (Fig. S11). For instance, for the 56 IBD genes found in the "Blood/Immune", or "Digestive" tissue groups, 57.1% of them were found, at a relaxed threshold of PIP > 0.5, in 5 or fewer tissues(Fig. 6D). One caveat in interpreting these findings is that the power of discovery is low, so cTWAS may underestimate the number of tissues for discovered genes.

Finally, we examined specific genes cTWAS found (Table S8 for all traits). We focused our analysis here to 56 genes found for IBD, at PIP > 0.8, by the two biologically relevant tissue groups: digestive and blood/immune (Table S9). At a more stringent PIP > 0.9, 30 genes were found (Table 1). The set of 56 genes included some well-known IBD genes, such as *TNFSF15*, *CARD9*, *RNF186*, *ITGAL*, and *ATG16L1*. Gene set enrichment analysis revealed many IBD-related GO terms in the 56 genes (Table S10). Using Weighted Set Cover [40], a method to extract non-redundant GO terms, we identified four GO terms, such as "cytokine production", and "defense response". 29 out of 56 genes were associated with at least one of these four terms (Fig. 6E).

We highlight some of the novel genes found by cTWAS that are biologically plausible for IBD. We started with genes located within IBD associated loci. These included a number of genes with immune functions, including *IFNGR2*, *FOSL2*, *STAT3*, *FCGR2A*, *IRF8*, and *ZFP36L2* (see Discussions). cTWAS also identified novel genes outside IBD loci. As demonstrated in the TWAS literature, the ability to find such genes is one of the main attractions of TWAS. Some of these

genes have relevant functions to IBD, such as *UBE2W* (Fig. 6F), *TYMP*, *LSP1*, and *CCR5* (Fig. S12, see Discussion). For example, *UBE2W* is a ubiquitin-conjugating enzyme. Ubiquitination is a post-translational modification that controls multiple steps in autophagy, a key process implicated in IBD [41]. Indeed, *UBE2W* knockdown mice showed mucosal injuries, and its overexpression ameliorated the severity of experimental colitis, a model of IBD [42].

Discussion

Expression QTL data are commonly used to annotate the effects of genetic variants and to nominate candidate genes for complex traits. Existing methods for such analysis, however, are susceptible to finding non-causal genes. This happens because these methods fail to account for the pleiotropic effects of eQTLs or their LD with nearby causal variants. To address this challenge, we generalize the popular TWAS model by explicitly and jointly modeling the effects of all gene expression traits and genetic variants in a region. Through extensive simulations, we show that our method achieves calibrated false discovery rates. In the analysis of several common GWAS traits, our method discovered a number of candidate genes for these traits, highlighting its potential as a gene discovery tool.

Our method is related to existing methods for integrating eQTL and GWAS data but has several key advantages. When the gene of interest, X, has a single causal eQTL, denoted as G_j, and the gene is the only causal gene in a locus, our model reduces to the colocalization model [7, 8]. Indeed, in such a case, eQTL and GWAS signals colocalize to the same variant G_j. One can also show that the prior parameter in colocaliation that controls to what extent colocalization is favored, a priori, over non-colocalization, corresponds to the prior parameters under our model that favor gene effects vs. variant effects (Supplemental Methods). Despite this connection, our model has key advantages over colocalization. Colocalization typically focuses on individual variants, yet our model uses imputed gene expression, which combines the effects of multiple variants. While colocalization can be generalized to allow for multiple causal variants [43], it does not explicitly account for the combined effects of variants. Additionally, while standard colocalization analysis is sensitive to the user-selected prior parameter [44], the corresponding parameter under our model is estimated from data. Indeed, our simulations show that our model is much better at controlling false discoveries than colocalization (Fig. 3B). Our method can also be viewed as a generalization of FOCUS, which uses a fine-mapping framework, similar to ours, but includes mostly gene effects, with a very simple model of variant effects, in the model. As others have pointed out [15], and our results showed (Fig. 4E, Fig. 6A), gene expression traits mediate a relatively small percent of trait heritability, so confounding by nearby variants is a much more common source of false discoveries.

Another class of methods share the same goal of inferring causal genes from joint analysis of eQTL and GWAS. These methods are often based on Mendelian Randomization (MR), and include SMR [10], TWMR [45], PMR-Egger [46], and MR-JTI [30]. The key assumption of MR, however, is often violated in our setting because of the "horizontal pleiotropic effects" of genetic instruments. The correlation between genetic instruments due to LD adds additional complexity when applying MR. Researchers have attempted to address these challenges. PMR-Egger models potential pleiotropic effects of variants, following the Egger extension of MR. This model is, however, overly simplified, as it assumes all genetic instruments of a gene have identical pleiotropic effects. Other methods, such as TWMR and MR-JTI, used a heterogeneity filter that removes genetic instruments with likely pleiotropic effects. While this idea has mitigated the problem of horizontal pleiotropy in other

applications, its utility is questionable in the eQTL setting. Most genes have a small number of independent cis-eQTLs/instruments, so detecting heterogeneity would be difficult in most cases.

One main finding from our simulations is that the majority of genes whose expression correlated with phenotypes in TWAS are not true causal genes. This often happens due to correlation of imputed expression with nearby causal variants ("confounding by variants", Fig. 4E). These results reflect the fact that *cis*-genetic components of expression explain relatively low proportions of trait heritability [15] (Fig. 6A). Indeed, a recent study has made an evolutionary argument for this observation [47]. Namely, genetic variants with large effects on the risk genes of diseases are under strong selection and as a result, their frequencies would be low and their eQTL effects hard to detect. Despite these observations, we view eQTLs, and molecular QTLs in general, combined with our discovery framework, as an important tool for disease gene discovery. As we demonstrated, even with bulk eQTL data from a single tissue, it is possible to identify a substantial number of candidate genes for LDL (Fig. 5A). Often, these genes were found using prediction models that combine the effects of multiple, potentially weak, eQTLs, suggesting that our method is less dependent on variants with large eQTL effects. Additionally, combining data across multiple tissues can further increase power, as shown in the case of IBD (Fig. 6C). Given the strong community interest in profiling eQTLs under various conditions, and across multiple cell types, leveraging these data will likely substantially increase power to discover disease genes. Lastly, our method can be applied to other types of molecular QTL data, e.g. chromatin accessibility QTLs. These additional data may explain a larger fraction of heritability missed by eQTLs [48].

The application of cTWAS to GWAS of several complex traits suggested interesting and novel candidate genes. In the case of LDL, we identified two genes, ACVR1C and INHBB, in the signaling pathway of activin, a signaling molecule in the TGF-beta superfamily. ACVR1C is a receptor of activin A and INHBB is a subunit of activin B. Both activin A and B were found to regulate lipolysis in hepatocytes and adipocytes and alter lipid compositions [49, 35]. In an exome-wide association study, a loss-of-function variant of ACVR1C was associated with metabolic phenotypes such as triglycerides and high-density lipoprotein level [50]. We identified another signal transduction gene, a protein kinase, PRKD2, as a candidate gene affecting LDL. PRKD2 deficiency in mice triggers hyperinsulinemia, metabolic disorders and dysregulation of LDL [51].

Our study of IBD revealed a number of interesting candidate IBD risk genes (Table 1). IFNGR2 is a receptor of interferon-gamma, a proinflammatory cytokine, whose activation may result in intestinal lesions [52]. IRF8 is an important regulator of multiple immune processes ranging from antigen presentation to response to cytokines [53]. IRF8 mutations cause severe immunodeficiency, and this gene has been explored as a drug target for IBD. CCR5 is a chemokine receptor, best known for its role in HIV infection. CCR5 blockade in mice inhibited leukocyte trafficking and reduced mucosal inflammation in murine colitis [54]. TYMP encodes thymidine phosphorylase, and TYMP mutations cause Mitochondrial Neurogastrointestinal Encephalomyopathy (MNGIE), a rare recessive disease [55]. The disease causes gastrointestinal symptoms and is often misdiagnosed as celiac or Crohn's disease [55]. LSP1 is a regulator of T-cell migration, and deletion of LSP1 gene has been implicated in rheumatoid arthritis, an autoimmune disease [56].

While all our analyses were performed using eQTL data, with one tissue at a time, cTWAS is a flexible framework that can be extended to more applications. It is relatively straightforward to include more tissues or cell types in the cTWAS model. This can be done by including multiple groups of explanatory variables (imputed expressions), with different priors for different groups. Having a multi-tissue model may increase the power to detect causal genes and help identify the

"causal tissue" for these genes. For additional development of the method, it may be helpful to account for imputation errors in the model [57]. Lastly, it would be interesting to generalize the model to allow joint analysis of multiple types of molecular QTL data. For example, incorporating both eQTL and splicing QTL data may allow us to better identify causal genes and mechanisms by combining information on eQTLs and sQTLs of the same genes.

In conclusion, by modeling genetic variants and imputed gene expression jointly, cTWAS accounts for pleiotropic effects and LD, creating a robust framework for detecting causal genes. With the large amount of molecular QTL datasets available and being generated, cTWAS promises to translate genetic associations of diseases into knowledge of risk genes, disease mechanisms and potential therapeutic targets.

Code availability

Our software is available at `https://xinhe-lab.github.io/ctwas/`.

Methods

Model of individual level data. Let y be the quantitative phenotype, assumed to be standardized, of an individual. We assume that y depends on imputed gene expressions and variant genotypes of the individual. We denote X_j the expression of the gene j, $\tilde{X}_j$ as its cis-genetic component, i.e. the imputed expression, and G_m the genotype of the variant m. We have the following regression model:

$$y = \sum_j \beta_j \tilde{X}_j + \sum_m \theta_m G_m + \epsilon. \tag{1}$$

where β_j and θ_m are the effect sizes of imputed expression of gene j and the variant m, respectively. ϵ is an normally distributed error terms, i.e. $\epsilon \sim N(0, \sigma^2)$, and is assumed to be independent across individuals. In practice, we standardize both $\tilde{X}_j$ and G_m to make the variance equal to 1 for all the genes and variants.

To obtain the imputed expressions, we use existing expression prediction models. Specifically, the imputed expression of a gene j is defined as $\sum_l w_{l,j} G_l$, where G_l is the genotype of variant l, and $w_{l,j}$ is the weight of the l-th variant in gene j's expression prediction model. We assume that these weights are given at the standardized scale, i.e. the weights were derived using standardized variant genotypes. This is the case for the FUSION expression models (`http://gusevlab.org/project s/fusion/`). When the provided weights are not on the the standardized scale, for example, from PredictDB (`https://predictdb.org/`) , we will calculate the standardized weights by multiplying the provided weights with variant genotype variances.

We specify different prior distributions of gene effects β_j's, and variant effects θ_m's. To describe these priors, we first note that our model is a special case of a more general regression model, where explanatory variables from multiple groups with different distributions of effect sizes. We write the general model with K groups of explanatory variables as:

$$y = \sum_{k=1}^{K} \sum_{j \in M_k} \beta_j X_j + \epsilon, \tag{2}$$

where X_j is jth explanatory variable and $j \in M_k$ denotes that it belongs to group k. In our case, the model has two groups of variables, imputed gene expressions and genetic variants. For simplicity of notations, we will use this general model in our following discussions. We assign a spike-and-slab prior distribution for the effect of variable j, with group-specific prior parameters. Specifically, when $j \in M_k$, we denote γ_j an indicator of whether X_j has non-zero effect:

$$\beta_j | \gamma_j = 1 \sim N(0, \sigma_k^2) \tag{3}$$
$$\beta_j | \gamma_j = 0 \sim \delta_0 \tag{4}$$
$$\gamma_j \sim \text{Bernoulli}(\pi_k). \tag{5}$$

Here δ_0 is the Dirac's delta function, $\pi_k = P(\gamma_j = 1 | j \in M_k)$ is the prior probability of the jth variable from group k being casual to the trait (non-zero effect), and σ_k^2 is the prior variance of the effect size of causal variables in the group k.

Inference of the individual level model. The inference has two main steps. In the first step, we will estimate the prior parameters $\theta = \{\pi_k, \sigma_k^2, k \in \{1, 2\}\}$ for the two groups, gene effects and variants effects. In the second step, we use the estimated θ, and compute the posterior inclusion probability (PIP) of each variable, defined as the posterior probability of $\gamma_j = 1$ given all the data and parameters.

The parameter estimation is done by Maximum Likelihood. Let $\boldsymbol{y}_{n \times 1}$ be the data of the response variable, where n is the sample size. Let $\boldsymbol{X}_{n \times p} = [\boldsymbol{X}_1 \ \boldsymbol{X}_2 \ ... \ \boldsymbol{X}_p]$ be the data of all the p explanatory variables. The likelihood of our model is given by

$$L(\theta; \boldsymbol{X}, \boldsymbol{y}, \sigma) = P(\boldsymbol{y} | \boldsymbol{X}, \theta, \sigma) = \sum_{\boldsymbol{\Gamma}} P(\boldsymbol{y} | \boldsymbol{X}, \boldsymbol{\Gamma}, \theta, \sigma) P(\boldsymbol{\Gamma} | \theta), \tag{6}$$

where $\boldsymbol{\Gamma} = [\gamma_1, \gamma_2, ..., \gamma_p]$ represents the "configuration" of the causal (non-zero effect) status of all variables. We note that σ is the standard deviation of the phenotypic variance, and is assumed to be given (see below). To maximize the likelihood, we use the Expectation-Maximization(EM) algorithm. In the E-step, we obtain the expectation of log likelihood over $\boldsymbol{\Gamma}$, $\mathbb{E}_{\boldsymbol{\Gamma}} \log P(\boldsymbol{X}, \boldsymbol{y}, \boldsymbol{\Gamma} | \theta^{(t)}, \sigma)$, where $\theta^{(t)}$ is the parameter values in the t-th iteration. In the M-step, we update $\theta^{(t)}$ using the following rules to maximize the expectation from the E-step:

$$\pi_k^{(t+1)} = \frac{1}{|M_k|} \sum_{j \in M_k} \alpha_j^{(t)} \tag{7}$$

$$\sigma_k^{2,(t+1)} = \frac{\sum_{j \in M_k} \alpha_j^{(t)} \cdot \tau_j^{2,(t)}}{\sum_{j \in M_k} \alpha_j^{(t)}}, \tag{8}$$

where $|M_k|$ is the size of M_k, $\alpha_j^{(t)} = P(\gamma_j = 1 | \boldsymbol{X}, \boldsymbol{y}, \theta^{(t)}, \sigma)$ is the PIP of variable j given data and current parameter values $\theta^{(t)}$, $\tau_j^{2,(t)} = \mathbb{E}(\beta_j^2 | \gamma_j = 1, \boldsymbol{X}, \boldsymbol{y}, \theta^{(t)}, \sigma)$ is the second moment of the posterior effect size of variable j, given that it is a causal variable. The update rules have simple interpretations. The new parameter $\pi_k^{(t+1)}$ is simply the average PIP of all variables in the group k and the new $\sigma_k^{2,(t+1)}$ is the weighted average of the second moment of the posterior effect sizes. See supplementary methods for the derivation of the EM algorithm.

Computing α_j and τ_j^2 at the t-th iteration (we removed superscript t from now on for simplicity) using all variables in the genome is computationally challenging. To reduce the computational

burden, we divide the genome into Linkage Disequilibrium (LD) blocks using LDetect [58] with variants approximately independent between blocks. We assign a gene into an LD block if all SNPs in its expression prediction model fall into that block. If the variants of the prediction model of any gene span multiple LD blocks, we merge all such blocks into a new block. We will then compute α_j and τ_j^2 of the variables in each block independently, while still using all variables in the genome to update the parameters using Equation 8.

Even within a single block, there will still be hundreds to thousands of variables. This poses a challenge to compute α_j and τ_j^2, as it requires marginalization of $\boldsymbol{\Gamma}$. To address this challenge, we first notice that our problem is now reduced to standard fine-mapping or Bayesian variable selection problem, with potentially different prior distributions of the effects of different variables. So we borrow from standard fine-mapping literature to compute α_j and τ_j^2's [16, 17]. We further assume that there is at most one causal variable per block (single effect assumption). So the only possible configurations are: $\boldsymbol{\Gamma}_j$, the configuration where $\gamma_j = 1$ and $\gamma_i = 0$ if $i \neq j$, and $\boldsymbol{\Gamma}_0$, the configuration where none of the variables in this block is causal. We also assume that a single block explains a small amount of the variance of the trait, so the residual variance of the trait, σ^2 would be approximately equal to 1, the total phenotypic variance (y is assumed to be standardized). With these assumptions, we have:

$$\alpha_j = P(\gamma_j = 1|\boldsymbol{X}, \boldsymbol{y}, \theta) = P(\boldsymbol{\Gamma}_j|\boldsymbol{X}, \boldsymbol{y}, \theta) = \frac{P(\boldsymbol{y}|\boldsymbol{X}, \boldsymbol{\Gamma}_j, \theta) \cdot P(\boldsymbol{\Gamma}_j|\theta)}{P(\boldsymbol{y}|\boldsymbol{X}, \boldsymbol{\Gamma}_0, \theta) \cdot P(\boldsymbol{\Gamma}_0|\theta) + \sum_j P(\boldsymbol{y}|\boldsymbol{X}, \boldsymbol{\Gamma}_j, \theta) \cdot P(\boldsymbol{\Gamma}_j|\theta)}. \tag{9}$$

$$\tau_j^2 = \mathbb{E}(\beta_j^2|\gamma_j = 1, \theta) = \text{var}(\beta_j|\boldsymbol{X}, \boldsymbol{y}, \theta, \gamma_j = 1) + (\mathbb{E}(\beta_j|\boldsymbol{X}, \boldsymbol{y}, \theta, \gamma_j = 1))^2. \tag{10}$$

Both α_j and τ_j^2 have closed forms under the single effect assumption. Specifically, the PIP of variable j, α_j is a simple function of prior inclusion probability $p_j = P(\gamma_j = 1|\theta)$, and the Bayes factors (BFs) of the variables:

$$\alpha_j = \frac{p_j/(1 - p_j)B_j}{1 + \sum_j p_j/(1 - p_j)B_j}, \qquad \text{where } B_j = \frac{P(\boldsymbol{y}|\boldsymbol{X}, \boldsymbol{\Gamma}_j, \theta)}{P(\boldsymbol{y}|\boldsymbol{X}, \boldsymbol{\Gamma}_0, \theta)}. \tag{11}$$

The BFs can be computed using the Wakefield formula [59]. See Supplementary methods for the details.

We provided some justifications of the single effect assumption. In statistical fine-mapping, several methods have been developed to incorporate functional information of variants to set the prior inclusion probabilities. In such models, it has been found that for the purpose of estimating parameters of the prior distributions, the approximation under single effect assumption is often sufficient [8]. Furthermore, in our implementation, we prune large blocks that are likely to contain multiple causal variables in the parameter estimation step (see Supplementary methods). Finally, our simulations show that the parameters estimated under single-effect assumption are generally accurate (Figure 2 of the main text).

After we estimate the prior parameters, we apply SuSiE [16], a fine-mapping method, on all variables, including both genes and variants, in each block. Note that all blocks, including the large blocks pruned in the parameter estimation step, will be analyzed. In applying SuSiE, we set the prior probability and prior effect variance of each variable, using the estimated parameters of the group (genes or variants) that this variable belongs to. We allow multiple causal variables by setting $L = 5$ in SuSiE and assign null weight as $1 - \sum_j p_j$. SuSiE will then return PIPs of all genes and variants in each LD block.

Model of summary statistics. It is often easier to work with summary statistics from GWAS data. The summary data would include the effect size estimates of variants, $\hat{\beta}_j$, and their standard errors s_j, as well as the LD between all pairs of variants, denoted as the matrix $\boldsymbol{R}$. The effect sizes can be standardized, denoted as $\hat{z}_j = \hat{\beta}_j/s_j$. Given that the summary data has only variant information, our model would first need to expand the summary data to include gene information. Specifically, we compute the marginal association of each imputed gene with the GWAS trait, and the correlation of any gene with all other genes and all the variants. These calculations will be described below. Once computed, we will have marginal associations of all variables, including genes and variants, $\hat{\boldsymbol{z}}$, and their correlation matrix $\boldsymbol{R}$. These data would be the input of our analysis.

Following the literature [60, 17], and particularly, the summary statistics version of SuSiE (SuSiE-RSS) [17], we have the following model of $\hat{\boldsymbol{z}}$,

$$\hat{\boldsymbol{z}}|\boldsymbol{z}, \boldsymbol{R} \sim N_p(\boldsymbol{R}\boldsymbol{z}, \boldsymbol{R}), \tag{12}$$

where $\boldsymbol{z} = (z_1, z_2, ..., z_p)$ denote the "standardized" true effect sizes. We assign the same prior to $\boldsymbol{z}$ as described in Equations 3-5: when the variable j belongs to the group k,

$$z_j|\gamma_j = 1 \sim N(0, \sigma_k^2) \tag{13}$$
$$z_j|\gamma_j = 0 \sim \delta_0 \tag{14}$$
$$\gamma_j \sim \text{Bernoulli}(\pi_k). \tag{15}$$

Again, we denote $\theta = \{\pi_k, \sigma_k^2\}$ the prior parameters and $\boldsymbol{\Gamma}$ the causal configuration. The likelihood of this model is given by

$$L_Z(\theta; \hat{\boldsymbol{z}}, \boldsymbol{R}) = P(\hat{\boldsymbol{z}}|\boldsymbol{R}, \theta) = \sum_{\boldsymbol{\Gamma}} P(\hat{\boldsymbol{z}}|\boldsymbol{R}, \boldsymbol{\Gamma}, \theta)P(\boldsymbol{\Gamma}|\theta) \tag{16}$$

Inference of the summary statistics model. We estimate the prior parameters θ by MLE. This can be done with the same algorithm used for the individual level model. Specifically, following SuSiE-RSS [17], the likelihood function under the individual level data can be rewritten in terms of the sufficient statistics $\hat{\beta}_j$ and s_j^2. Then, if we make the following substitutions, the likelihood of the individual level model would be identical to that of the summary statistics model. Specifically, we change $\boldsymbol{\beta} = (\beta_1, \beta_2, ..., \beta_p)$ to $\boldsymbol{z}$, $\boldsymbol{X}^T\boldsymbol{X}$ to $\boldsymbol{R}$, $\boldsymbol{X}^T\boldsymbol{y}$ to $\hat{\boldsymbol{z}}$, $\boldsymbol{y}^T\boldsymbol{y}$ to 1 and n to 1. Also the prior model of z in the summary statistics model is the same as the prior model of β in the individual level model. Therefore, we we can use the same EM algorithm and the update rules to estimate θ. The update rules follow Equations 7-8, where the PIP of variable j is now defined as $\alpha_j^{(t)} = P(\gamma_j = 1|\hat{\boldsymbol{z}}, \boldsymbol{R}, \theta^{(t)})$ and the second moment of the posterior effect $\tau_j^{2,(t)} = \mathbb{E}(z_j^2|\gamma_j = 1, \hat{\boldsymbol{z}}, \boldsymbol{R}, \theta^{(t)})$.

Similarly as described in the inference procedures for individual level data, we divided genome into LD blocks, and used the blocks that are likely to have at most one effect, to infer the prior parameters θ. We then used these parameters to set the prior probabilities and prior variances of all variables within each block, and used SuSiE-RSS, with $\hat{\boldsymbol{z}}, \boldsymbol{R}$ as input and set $L = 5$, to obtain PIPs of all variables.

Computation of marginal associations $\hat{\boldsymbol{z}}$ and correlation matrix $\boldsymbol{R}$. When variable j is a gene, we can compute its Z-score, $\hat{z}_j$ following the S-PrediXcan formula [27]. Specifically, suppose the gene j has m SNPs with nonzero weights in its expression prediction model. Let

$\boldsymbol{w}_j = (w_{j1}, w_{j2}, .., w_{jm})$ be their standardized weights, $\hat{\boldsymbol{z}}_j^s = (\hat{z}_{j1}, \hat{z}_{j2}..., \hat{z}_{jm})^T$ be the Z scores of the associations of the variants to the phenotype, and $\boldsymbol{R}_j^s$ be the $m \times m$ correlation matrix, i.e. LD matrix, of these variants. Then $\hat{z}_j$ is basically the weighted sum of the Z-scores of all the m variants:

$$\hat{z}_j = \frac{\boldsymbol{w}_j \hat{\boldsymbol{z}}_j^s}{\sqrt{\boldsymbol{w}_j \boldsymbol{R}_j^s \boldsymbol{w}_j^T}}. \tag{17}$$

We also need $\boldsymbol{R}$, the correlation matrix, of all variables. We denote its element at row i and column j as $R_{ij} = \mathrm{Cor}(\boldsymbol{X}_i, \boldsymbol{X}_j)$, where $\boldsymbol{X}_i$ and $\boldsymbol{X}_j$ are the n-dim. genotypes (n is sample size) of the variables i and j. When both variable i and j are variants, R_{ij} is given by the variant LD matrix which contains the correlation between all variant pairs. When one of the variable is a gene, for example variable j is a gene, R_{ij} is given by:

$$R_{ij} = \frac{\mathrm{Cov}\left(\boldsymbol{X}_i, \boldsymbol{X}_j\right)}{\sqrt{\mathrm{Var}(\boldsymbol{X}_i)\mathrm{Var}(\boldsymbol{X}_j)}} = \frac{\mathrm{Cov}\left(\boldsymbol{X}_i, \boldsymbol{w}_j \boldsymbol{X}_j^s\right)}{\sqrt{\mathrm{Var}(\boldsymbol{X}_i)\mathrm{Var}(\boldsymbol{w}_j \boldsymbol{X}_j^s)}} = \frac{\boldsymbol{R}_{i,j}^s \boldsymbol{w}_j^T}{\sqrt{\boldsymbol{w}_j \boldsymbol{R}_j^s \boldsymbol{w}_j^T}}, \tag{18}$$

where $\boldsymbol{X}_j^s$ is a $n \times m$ matrix of genotypes for the m variants in gene j's prediction model, $\boldsymbol{R}_{i,j}^s$ is the m-dimensional vector of correlation between variant i with each of the m variants in the prediction model of gene j. When both variables i and j are genes,

$$R_{ij} = \frac{\mathrm{Cov}\left(\boldsymbol{X}_i, \boldsymbol{X}_j\right)}{\sqrt{\mathrm{Var}(\boldsymbol{X}_i)\mathrm{Var}(\boldsymbol{X}_j)}} = \frac{\mathrm{Cov}\left(\boldsymbol{w}_i \boldsymbol{X}_i^s, \boldsymbol{w}_j \boldsymbol{X}_j^s\right)}{\sqrt{\mathrm{Var}(\boldsymbol{w}_i \boldsymbol{X}_i^s)\mathrm{Var}(\boldsymbol{w}_j \boldsymbol{X}_j^s)}} = \frac{\boldsymbol{w}_i \boldsymbol{R}_{i,j}^s \boldsymbol{w}_j^T}{\sqrt{\boldsymbol{w}_i \boldsymbol{R}_i^s \boldsymbol{w}_i^T} \sqrt{\boldsymbol{w}_j \boldsymbol{R}_j^s \boldsymbol{w}_j^T}}, \tag{19}$$

where $\boldsymbol{X}_i^s$ is the genotype matrix and $\boldsymbol{w}_i$ is the vector of standardized weights for variants in gene i's prediction model. $\boldsymbol{R}_i^s$ is the correlation matrix of $\boldsymbol{X}_i^s$. $\boldsymbol{R}_{i,j}^s$ is now the correlation matrix between the variants in the prediction models of gene i and gene j.

Proportion of variance explained by SNPs and genes. We assume that all the explanatory variables and the response variable in the regression model are standardized, with variance equal to 1. Then the proportion of variance explained (PVE) by a single variable j, is simply $\beta_j^2 \cdot \mathrm{Var}(X_j)/\mathrm{Var}(y) = \beta_j^2$. Assuming that we use the Z-scores in the summary statistical model, the effect size is related to Z-score by: $\beta_j = z_j/\sqrt{n}$, where n is the sample size. So on average, the PVE of a variable in group k (variant or gene) is: $\mathbb{E}(z_j^2) = \sigma_k^2/n$, where σ_k is the prior variance of effect size in the group k, at the Z-score scale. The expected number of variables with non-zero effects in the group k is: $\pi_k \cdot |M_k|$, where π_k is the prior inclusion probability and $|M_k|$ the group size. Putting this together, the percent of total variance explained (PVE) by the group k is given by

$$\mathrm{PVE}_k = \sigma_k^2 \cdot \pi_k \cdot |M_k| \cdot n^{-1}. \tag{20}$$

This equation is used to compute PVE from estimated parameters using both simulated and real data.

Speeding up computation. Parameter estimation is the most time-consuming step. This is especially the case when variants are densely genotyped. We developed a "thinning" procedure to speed up this step. Specifically, we randomly select a fraction of SNPs in the single effect blocks, and only use these variants in the EM update. When most selected blocks have < 1000 variants, the EM algorithm can be done in one or two hours. We have found in our simulations that this thinning step does not affect the accuracy of results. In the final step of analyzing individual blocks, the estimated

prior probability for variants from the thinning procedure will be scaled back for the original variants data. This is done by dividing the prior probability by the fraction of selected variants. The other parameters are not affected by this procedure. To reduce computational burden of the final step, we can analyze an individual block with the original variants only when results using the thinned variants showed strong signals for gene effect, e.g the maximum PIP of genes in the block > 0.8. Our software allows the user to specify the desired thinning parameter (the fraction of variants selected) in the parameter estimation step and the PIP threshold used to determine which blocks to analyze with original variants in the final step.

Simulations. In our simulations, we used the following data:

- Genotype data. We used genotype data from UK biobank by randomly selecting 80,000 samples. We then filtered samples to only keep "White British", removed samples with missing information, mismatches between self-reported and genetic sex, or "outliers" as defined by UK Biobank. We also removed any individuals that have close relatives in the cohort. This ended up with a cohort of $n = 45,087$ samples. We used SNPs from chr 1 to chr 22 and selected those with minor allele frequency > 0.05 and at least 95% genotyping rate. After filtering, 6,228,664 SNPs remained and were used in our analysis.

- Gene expression prediction models. We used GTEx v7 Adipose tissue dataset. This dataset contains 8,021 genes with expression models. We used the lasso weights from the FUSION website (`http://gusevlab.org/projects/fusion/`).

We first impute gene expression for all samples using the prediction models. SNPs genotypes are harmonized between the expression prediction model and UK Biobank genotypes so that the reference and alternate alleles match. SNPs in the FUSION prediction models but not in UK Biobank, about 13% of all, were not used in imputing gene expression. We then sample the causal genes and SNPs under given prior inclusion probabilities π_k's, and then sample their effect sizes accordingly using the prior variance parameter σ_k^2. We then simulate y under the model defined in Equation 2. The prior parameters π_k, σ_k^2 were chosen to reflect the genetic architecture in real data. In particular, it was estimated that gene expression mediates about 10-20% trait heritability [15]. And the studies using rare variants for complex traits suggested that about 5% of protein coding genes are likely causal [61]. Given these considerations, we set the prior probability for SNPs to 10^{-4} or 2.5×10^{-4}, and PVE of SNPs to 0.3 or 0.5. For the genes, we set the prior probability 0.015 or 0.05 and PVE of genes from 0.02 to 0.1.

To run cTWAS, we performed association of individual SNPs with the trait y, to obtain summary statistics of SNPs $\hat{z}_{\mathrm{SNP}}$. We randomly selected 2000 samples from the cohort to calculate SNP genotype correlation matrix, or LD matrix $\hat{R}_{\mathrm{SNP}}$. We then ran cTWAS summary statistics version under each simulation setting with $\hat{z}_{\mathrm{SNP}}$, $\hat{R}_{\mathrm{SNP}}$ and expression prediction models as input. The software will harmonize SNP genotypes for $\hat{z}_{\mathrm{SNP}}$, $\hat{R}_{\mathrm{SNP}}$ and expression prediction models, so that the reference and alternate allele match. To further reduce computational burden in estimating parameters, we only used 1 in every 10 SNPs (SNP thinning) in the EM algorithm. When calculating PIP, we first run SuSiE-RSS with $L = 5$ in each LD Block with thinned SNPs. For each block with maximum gene PIP > 0.8, we re-run SuSiE-RSS with $L = 5$ with the original SNPs to get final gene PIPs.

We ran FUSION (`https://github.com/gusevlab/fusion_twas`) on the same simulated GWAS summary statistics data and the same gene prediction models. Bonferroni correction was applied on the result with corrected $p < 0.05$. We ran coloc (`https://cran.r-project.org/web/packa`

ges/coloc/index.html) which is provided as part of the FUSION package and used PIP4 > 0.8 as cutoff. We ran FOCUS (https://github.com/bogdanlab/focus) under the default setting and use PIP > 0.8 as the cut off. For SMR and SMR + Heidi (https://yanglab.westlake.edu.cn/software/smr/), we used the eQTL marginal association statistics provided by GTEx v7 as input. For SMR, We used BH adjusted p values < 0.05 to select significant genes. For SMR + Heidi, we further filtered the SMR results by requiring $p\_HEIDI > 0.05$, we then applied the BH procedure on the SMR p values on the filtered gene list and require adjusted $p < 0.05$ to select significant genes.

GWAS summary statistics. We obtained GWAS summary statistics of four common traits: low density lipoprotein cholesterol (LDL), inflammatory bowel disease (IBD), schizophrenia (SCZ), and systolic blood pressure (SBP). The LDL and SBP summary statistics were from the UK Biobank, computed by the Rapid GWAS project [**rapid**] using LD Score Regression [62]. These summary statistics were downloaded from the IEU OpenGWAS project [63] using GWAS IDs "ukb-d-30780_irnt" (LDL) and "ukb-a-360" (SBP). Both LDL and SBP summary statistics were based on the White British subpopulation of the UK Biobank, with sample sizes of N=343,621 and N=317,754 respectively. The IBD summary statistics were from the International IBD Genetics Consortium [64], computed by meta-analysis using METAL [65]. These summary statistics were obtained from IEU OpenGWAS using GWAS ID "ebi-a-GCST004131". IBD includes cases of both Crohn's disease and Ulcerative colitis. The IBD summary statistics were based on non-overlapping samples of European ancestry with a combined sample size of N=59,957. The SCZ summary statistics were from the Psychiatric Genetics Consortium and the CardiffCOGS study [66], computed by meta-analysis using METAL [65]. These summary statistics were obtained from the authors via the link provided in the manuscript. The SCZ summary statistics were based on non-overlapping samples of primarily European ancestry with a combined sample size of N=105,318.

LD reference data. We computed an LD reference panel of common biallelic variants using the White British subpopulation of the UK Biobank. This panel is an in-sample reference for GWAS summary statistics from the Rapid GWAS project [**rapid**]. First, we used plate and well information from the genotyping to unambiguously identify over 99% (357,654 of 361,194) of the samples used in the Rapid GWAS project in our data. To ease computation, we randomly selected 10% of these samples to serve as the LD reference panel [67]. We also limited our panel to common autosomal variants with MAF > 0.01 in the Rapid GWAS analyses. Then, we computed correlations between all pairs of variants within each of 1,700 approximately independent regions. These regions are assumed to have low LD between them, and are based on previously identified regions [58] that could be lifted over from hg37 to hg38 positions. The final LD reference panel consists of 1700 correlation matrices and contains 9,309,375 variants. This LD reference panel was used when analyzing all traits, including those that were not measured in the White British subpopulation of UK Biobank.

Prediction model of gene expression. We used expression prediction models for 49 GTEx tissues from PredictDB [4] [26] [27]. We used the mashr-based prediction models, which borrowed information across tissues to improve prediction accuracy in any specific tissue [37]. These prediction models were built using fine-mapped variants and are sparse, with a maximum of 5 eQTL per gene. The models were constructed using in-sample LD from GTEx, and the covariance between pairs of variants within each gene is reported alongside the models. We included only protein-coding genes for our analyses. We analyzed LDL using eQTL data from liver tissue. For the analyses of other traits (IBD, SCZ, SBP), we used eQTL data from all 49 tissues, performing a separate analysis

using each tissue.

Harmonization of GWAS data and expression prediction models. In order to ensure consistency between the GWAS z-scores, expression prediction models, and the LD reference panel, we performed two procedures to harmonize the datasets. The objective of harmonization was to ensure that the reference and alternate alleles of each variant are defined consistently across all three datasets [17]. In our case, we must harmonize both the GWAS z-scores and the eQTL prediction models to our LD reference, and we use a different harmonize procedure for each. These procedures are based in part on previous work [26].

To describe the two procedures, it is necessary to define several cases of inconsistencies that can occur in either dataset. The first case is a variant with its reference and alternate alleles "flipped" with respect to the LD reference. The association z-scores or eQTL weights in the prediction model of the flipped variants should have their signs reversed to be consistent with the LD reference. The second case is a variant that has had its strand "switched" with respect to the LD reference (e.g. variant is G/A in the LD reference but C/T in the other dataset). In this case, the reference and alternate alleles are the same, just named using different strands. The z-scores or weights of switched variants should not be changed, as they are already consistent with the LD reference. The third case is a variant that is "ambiguous" as to whether it is flipped or switched. This occurs when the two alleles of a variant are also complementary base pairs (A$\leftrightarrow$T substitutions or G$\leftrightarrow$C). For example, consider a variant that is A/T in the LD reference and T/A. It is unclear if this variant is flipped or switched with respect to the LD reference (both result in T/A), and it is ambiguous as to whether the signs of the z-scores or weights should be reversed. We say that variants are "unambiguous" when they do not involve substitutions of complementary base pairs.

To harmonize the z-scores from GWAS summary statistics, we first identified all inconsistencies in reference and alternate alleles between the z-scores and the LD reference. Next, we resolved all unambiguous cases of flipped and switched alleles, reversing the sign of z-scores that were flipped and taking no action for switched alleles. Then, we imputed the z-scores for ambiguous variants using all unambiguous variants in each of the LD regions [68]. If the sign of the imputed z-score did not match the sign of the observed z-score, we used the sign of the imputed z-score, reversing the sign of the observed z-score. Note that we did not perform the procedure to resolve ambiguous variants when analyzing LDL or SBP, as both the summary statistics and LD reference panel are derived from UK Biobank data, and it is unlikely that strand is inconsistent between different releases of the UK Biobank genotype data.

To harmonize the eQTL prediction models, we first identified all inconsistencies in reference and alternate alleles between the prediction models and the LD reference. Next, we resolved all unambiguous cases of flipped and switched alleles, reversing the sign of weights that were flipped and taking no action for switched alleles. These steps to resolve unambiguous variants are the same as in the z-score harmonization procedure. To resolve ambiguous variants, we leveraged correlations between ambiguous and unambiguous variants in both our LD reference panel and the LD panel used to construct the PredictDB models. PredictDB reports the covariance between pairs of variants within each gene prediction model. For gene prediction models that include both ambiguous variants and unambiguous variants, we computed the sum of correlations between each ambiguous variant and the unambiguous variants in the prediction model, using both our LD reference panel and the LD used for the prediction models. If the sign of the total correlation in the LD reference of the prediction models did not match the sign of the total correlation in our LD reference panel, we reversed the sign of the prediction model weights for the ambiguous variant. If the total correlation

in the LD reference was equal to zero, then we set the weight of the ambiguous variant to zero, as these ambiguous variants did not have any unambiguous variants in the same LD region. For gene prediction models that include only a single ambiguous variant and no unambiguous variants, we left the sign of the prediction model weight unchanged; the resulting gene z-score may have an incorrect sign, but the magnitude of the z-score will be correct. We excluded gene prediction models with multiple ambiguous variants and no unambiguous variants, as their gene z-scores could be incorrect in both sign and magnitude.

Performing cTWAS analysis in real data. We used the following cTWAS settings when analyzing real data. For parameter estimation, we used the default procedure for selecting starting values of the E-M algorithm. We then performed 30 iterations of the E-M algorithm assuming L=1 effect (at most a single causal effect) in each region, using variants that were thinned by 10% to reduce computation. For computing PIPs of genes and variants, we used thinned variants and assumed L=5 (at most 5 causal effects) in each region. For regions with maximum gene PIP > 0.8, we recomputed PIPs using all variants, allowing a maximum of 20,000 SNPs in a region to reduce computation. Unless specified otherwise, we used the threshold PIP > 0.8 for declaring significant genes.

Total heritability and percentage of heritability mediated by genes using MESC. To assess parameter estimates from cTWAS, we computed total heritability and the percentage of heritability mediated by genes using MESC [15] for comparison. We used the same summary statistics as in the cTWAS analysis, and we used GTEx v8 expression scores for individual tissues available via the MESC repository. These expression scores are derived from the same GTEx dataset as the PredictDB models. Note that expression scores for "Kidney Cortex" tissue were not available from MESC, although this tissue is included in PredictDB and our other analyses.

Silver standard and previously reported candidate genes of complex traits. To evaluate our findings, we compared the genes detected using cTWAS to curated lists of known or candidate genes for each trait. For LDL, we combined curated gene lists from two previous studies [29] [30], for a total of 69 LDL-related genes. These are genes that cause Mendelian disease or are drug targets, are in the KEGG "cholesterol metabolism" pathway, or are otherwise documented in the literature. For IBD, we used a list of 26 genes curated from literature, reported in the paper describing the activity-by-contact (ABC) model [69]. We refer to the gene lists for LDL and IBD as "silver standard" gene lists throughout the text based on the strength of their evidence. In supplemental materials, we also refer to "previously reported candidate genes" for SCZ and SBP. These are gene lists that are primarily supported by GWAS data rather than literature or experimental evidence. For SCZ, we used a list of 130 genes prioritized as SCZ-related based on proximity to GWAS, fine-mapping results and eQTL evidence [70]. For SBP, we used a list of 53 genes, supported by TWAS, gene expression, drug targets, pathway analysis and rare variant studies, reported in a large trans-ethnic GWAS (Table 3 of the paper) [71].

Evaluating methods in distinguishing silver standard and bystander genes for LDL. Following earlier work [31], we assessed the performance of TWAS and cTWAS on real data by comparing their ability to distinguish LDL silver standard genes from other nearby genes. We defined a set of "bystander" genes that were within 1Mb of a silver standard gene. These bystanderer genes would be considered the negative set. We limited our analysis to the 46 of 69 silver standard genes with imputed expression after harmonization, and the 539 imputed bystander genes that are nearby these genes. Next, we determined if these silver standard and bystander genes were significant by TWAS (Bonferroni) or cTWAS (PIP > 0.8). Then, we computed the precision of each method,

defined as: (# detected silver standard genes) / (# detected silver standard genes + # detected bystander genes).

Fine-mapping analysis of POLK / HMGCR locus. To examine the POLK / HMGCR locus, we used fine-mapping results of LDL from UK Biobank from PolyFun-SuSiE [33]. Specifically, PolyFun specifies prior probabilities for fine-mapping by estimating functional enrichments for a broad set of coding, conserved, regulatory and LD-related annotations. Then SuSiE was applied using the estimated priors with L=10 effects per locus. The summary statistics data for UKBB LDL were downloaded from: `http://data.broadinstitute.org/alkesgroup/polyfun_results /biochemistry_LDLdirect.txt.gz`.

To annotate the functions of variants their putative target genes, we used additional datasets. H3K4me1 ChIP-seq data of from human liver samples were downloaded from the ENCODE portal (ENCODE ID: "ENCFF323TFF"). Promoter Capture Hi-C (PC-HiC) data from human liver were obtained from a previous study [72].

Classifying TWAS false positive genes for LDL by source of confounding. To better understand how TWAS generated false positive findings for LDL, we classified whether TWAS false positives as primarily due to confounding by variants or confounding by genes. We defined TWAS false positives as genes that were significant by TWAS (Bonferroni) but PIP < 0.5 by cTWAS. To categorize these false positive genes, we first assigned them to credible sets. These credible sets were reported by cTWAS, using the default SuSiE setting, which means that only credible sets with sufficient "purity" are reported (i.e. all variables in a credible set are highly correlated, r > 0.5). If a false positive gene was not included in any credible set, but was highly correlated (r > 0.5) with at least one variant or gene in a credible set, that false positive gene was also assigned to the credible set. After assigning false positive genes to credible sets, for each assigned gene, we summed the PIPs of all other genes and variants in its credible set to obtain total PIPs for confounding genes and variants. If the total gene PIP was higher than that of the variants, we classified the gene as confounded by genes, otherwise, confounded by variants.

Assessing the novelty of cTWAS detected genes. If a gene was not included in the list of silver standard or previously reported genes of a trait, and was not the nearest gene of the lead variant in a genome-wide significant locus, we considered the gene a "novel" finding. We defined "nearest" genes as those with start or end positions that were the nearest out of all protein-coding genes to the lead variant of a genome-wide significant locus. To define independent lead variants for the GWAS, we performed the following procedure. First, we selected the most significant variant as a lead variant, and we removed all other variants within 500kb. Then, we iterated these selection and removal steps until there were no genome-wide significant variants remaining. We then identified the protein-coding genes that were nearest to each lead variant in the resulting list. In some analysis, we also assessed if the cTWAS detected gene were inside a genome-wide significant locus. We referred to these genes as "nearby" genes, if they were within 500kb of a genome-wide significant variant.

Gene Ontology (GO) analysis of candidate genes. We performed enrichment analysis to characterize the genes detected by cTWAS for LDL and IBD. We used Enrichr [73, 74] to identify Gene Ontology terms in the Biological Process, Molecular Function, or Cellular Component domains that are enriched for genes detected by cTWAS. Enrichr uses the Benjamini-Hochberg procedure to ensure that FDR = 0.05 in each domain. We report the full enrichment analysis results using Enrichr across all domains for LDL and IBD (Table S3; Table S11). We compared the results from cTWAS genes with similar enrichment analysis results that used silver standard genes for LDL and IBD (Table S4; Table S12). We also compared these results with enrichment results from MAGMA

[75] [76], a method that performs enrichment analysis based on GWAS data, using default settings (Table S5; Table S13). When reporting the IBD results (Table S9), we annotated detected genes with the enriched GO terms (using cTWAS genes, silver standard genes, or MAGMA) that each gene is a member of. For all three GO annotations, a maximum of 5 significant terms per gene were shown, ordered by odds ratios (cTWAS, silver standard) or p-values (MAGMA).

To visualize the GO biological process terms enriched for LDL cTWAS genes (Fig. 5B), we removed redundant terms to improve clarity. To do this, we first ranked all GO terms by their significance (p-values). A term was considered redundant if all the cTWAS genes in that term had already been included in a more significant GO term.

To visualize the GO biological process terms enriched for IBD cTWAS genes (Fig. 6E), we used the "Weighted Set Cover" tool in WebGestalt [40] to remove redundant GO terms. To further simplify visualization, we omitted GO terms whose detected genes were all included in other terms identified by Weighted Set Cover. We also report the full enrichment analysis results using WebGestalt for IBD (Table S10).

Understanding cTWAS false negatives for LDL using silver standard genes. To understand how silver standard genes for LDL were missed by cTWAS, we categorized the outcome of cTWAS for all 69 silver standard genes. We binned these genes into the following categories: detected by cTWAS ("Detected"); significant by TWAS but not detected by cTWAS ("Significant Z-score but Undetected"); insignificant by TWAS and not detected by cTWAS, with a genome-wide significant GWAS variant for LDL within 500kb ("Insignificant Z-score"); no expression prediction model ("Not Imputed"); insignificant by TWAS and not detected by cTWAS, with no genome-wide significant GWAS variant for LDL within 500kb ("No GWAS Signal"). For genes detected by cTWAS we used a PIP > 0.8 threshold, and for genes significant by TWAS we used a Bonferonni threshold.

Summarizing cTWAS results using tissue groups. To aid the interpretation of cTWAS findings, we grouped related tissues into "tissue groups" and summarized the findings within these groups. We used previously defined tissue groups that assigned 37 of the 49 tissues to one of 7 tissue groups [15]. We then took the union of genes detected at PIP > 0.8 in any tissue within each tissue group, and we used these combined lists of detected genes for downstream analyses.

References

[1] Visscher P. "10 Years of GWAS Discovery: Biology, Function, and Translation". In: *Am. J. Hum. Genet* 101 (2017), pp. 5–22.

[2] Uffelmann E. "Genome-wide association studies". In: *Nature Reviews Methods Primers* 1 (2021), pp. 1–21.

[3] Tam V. "Benefits and limitations of genome-wide association studies". In: *Nat. Rev. Genet* 20 (2019), pp. 467–484.

[4] Gamazon E. "A gene-based association method for mapping traits using reference transcriptome data". In: *Nat. Genet* 47 (2015), pp. 1091–1098.

[5] Gusev A. "Integrative approaches for large-scale transcriptome-wide association studies". In: *Nat. Genet* 48 (2016), pp. 245–252.

[6] Wainberg M. "Opportunities and challenges for transcriptome-wide association studies". In: *Nat. Genet* 51 (2019), pp. 592–599.

[7] Hormozdiari F. "Colocalization of GWAS and eQTL Signals Detects Target Genes". In: *Am. J. Hum. Genet* 99 (2016), pp. 1245–1260.

[8] Xiaoquan Wen et al. "Efficient integrative multi-SNP association analysis via deterministic approximation of posteriors". In: *The American Journal of Human Genetics* 98.6 (2016), pp. 1114–1129.

[9] Hukku A et al. "Analyzing and reconciling colocalization and transcriptome-wide association studies from the perspective of inferential reproducibility". In: *Am. J. Hum. Genet* 109 (2022), pp. 825–837.

[10] Zhu Z. "Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets". In: *Nat. Genet* 48 (2016), pp. 481–487.

[11] Liu S. "Illuminating links between cis-regulators and trans-acting variants in the human prefrontal cortex". In: (2021). DOI: 10.1101/2021.09.07.459322.

[12] Fishilevich S. "GeneHancer: genome-wide integration of enhancers and target genes in GeneCards". In: (2017), p. 28.

[13] Urbut S et al. "Flexible statistical methods for estimating and testing effects in genomic studies with multiple conditions". In: *Nat. Genet* 51 (2019), pp. 187–195.

[14] Liu B et al. "Abundant associations with gene expression complicate GWAS follow-up". In: *Nat. Genet* 51 (2019), pp. 768–769.

[15] Yao D et al. "Quantifying genetic effects on disease mediated by assayed gene expression levels". In: *Nat. Genet* 52 (2020), pp. 626–633.

[16] Wang G et al. "A simple new approach to variable selection in regression, with application to genetic fine mapping". In: *J. R. Stat. Soc. Series B Stat. Methodol* 82 (2020), pp. 1273–1300.

[17] Zou Y et al. "Fine-mapping from summary data with the "Sum of Single Effects" model". In: *PLoS Genet* 18 (2022), e1010299.

[18] Guan Y and Stephens M. "Bayesian variable selection regression for genome-wide association studies and other large-scale problems". In: 5 (2011), pp. 1780–1815.

[19]   Carbonetto P and Stephens M. "Scalable Variational Inference for Bayesian Variable Selection in Regression, and Its Accuracy in Genetic Association Studies". In: 7 (2012), pp. 73–108.

[20]   Wright S. "Correlation and causation". In: (1921).

[21]   Siewert-Rocks K et al. "Leveraging gene coregulation to identify gene sets enriched for disease heritability". In: (2021). DOI: 10.1101/2021.07.22.453442.

[22]   Zhou X, Im H, and Lee S. "CORE GREML for estimating covariance between random effects in linear mixed models for complex trait analyses". In: *Nat. Commun* 11 (2020), p. 4208.

[23]   Sudlow C. "UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age". In: *PLoS Med* 12 (2015), e1001779.

[24]   Giambartolomei C. "Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using Summary Statistics". In: *PLoS Genet* 10 (2014), e1004383.

[25]   Mancuso N. "Probabilistic fine-mapping of transcriptome-wide association studies". In: *Nat. Genet* 51 (2019), pp. 675–682.

[26]   Barbeira A. "Exploiting the GTEx resources to decipher the mechanisms at GWAS loci". In: *Genome Biol* 22 (2021), p. 49.

[27]   Barbeira A. "Exploring the phenotypic consequences of tissue specific gene expression variation inferred from GWAS summary statistics". In: *Nat. Commun* 9 (2018), p. 1825.

[28]   Browser Heritability. In: *https://nealelab.github.io/UKBB_ldsc/index.html* (accessed in 2021).

[29]   Forgetta V. "An effector index to predict target genes at GWAS loci". In: *Hum. Genet* 141 (2022), pp. 1431–1447.

[30]   Zhou D. "A unified framework for joint-tissue transcriptome-wide association and Mendelian randomization analysis". In: *Nat. Genet* 52 (2020), pp. 1239–1246.

[31]   Ndungu A et al. "A Multi-tissue Transcriptome Analysis of Human Metabolites Guides Interpretability of Associations Based on Multi-SNP Models for Gene Expression". In: *Am. J. Hum. Genet* 106 (2020), pp. 188–201.

[32]   Guthrie P. "Complexity of a complex trait locus: HP, HPR, haemoglobin and cholesterol". In: *Gene* 499 (2012), pp. 8–13.

[33]   Weissbrod O. "Functionally informed fine-mapping and polygenic localization of complex trait heritability". In: *Nat. Genet* 52 (2020), pp. 1355–1363.

[34]   Stancu C and Sima A. "Statins: mechanism of action and effects". In: *J. Cell. Mol. Med* 5 (2001), pp. 378–387.

[35]   Yogosawa S et al. "Activin receptor-like kinase 7 suppresses lipolysis to accumulate fat in obesity through downregulation of peroxisome proliferator-activated receptor γ and C/EBPα". In: *Diabetes* 62 (2013), pp. 115–123.

[36]   Osamu Hashimoto et al. "Activin E Controls Energy Homeostasis in Both Brown and White Adipose Tissues as a Hepatokine". en. In: *Cell Rep.* 25.5 (Oct. 2018), pp. 1193–1203.

[37]   Barbeira A. "Integrating predicted transcriptome from multiple tissues improves association detection". In: *PLoS Genet* 15 (2019), e1007889.

[38]   GTEx Consortium. "The GTEx Consortium atlas of genetic regulatory effects across human tissues". en. In: *Science* 369.6509 (Sept. 2020), pp. 1318–1330.

[39] Nasser J. "Genome-wide enhancer maps link risk variants to disease genes". In: *Nature* 593 (2021), pp. 238–243.

[40] Liao Y et al. "WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs". In: *Nucleic Acids Res* 47 (2019), W199–W205.

[41] Daniel B Graham and Ramnik J Xavier. "Pathway paradigms revealed from the genetics of inflammatory bowel disease". en. In: *Nature* 578.7796 (Feb. 2020), pp. 527–539.

[42] Shaoxin Wang et al. "UBE2W Improves the Experimental Colitis by Inhibiting the NF-κB Signaling Pathway". en. In: *Dig. Dis. Sci.* (Mar. 2022).

[43] Wallace C. "A more accurate method for colocalisation analysis allowing for multiple causal variants". In: *PLoS Genet* 17 (2021), e1009440.

[44] Hukku A. "Probabilistic colocalization of genetic variants from complex and molecular traits: promise and limitations". In: *Am. J. Hum. Genet* 108 (2021), pp. 25–35.

[45] Porcu E. "Mendelian randomization integrating GWAS and eQTL data reveals genetic determinants of complex and clinical traits". In: *Nat. Commun* 10 (2019), p. 3300.

[46] Yuan Z. "Testing and controlling for horizontal pleiotropy with probabilistic Mendelian randomization in transcriptome-wide association studies". In: *Nat. Commun* 11 (2020), pp. 1–14.

[47] Mostafavi H et al. "Limited overlap of eQTLs and GWAS hits due to systematic differences in discovery". In: (2022). DOI: `10.1101/2022.05.07.491045`.

[48] Baca S. "Genetic determinants of chromatin reveal prostate cancer risk mediated by context-dependent gene regulation". In: *Nat. Genet* (2022). DOI: `10.1038/s41588-022-01168-y`.

[49] Yndestad A. "A complex role of activin A in non-alcoholic fatty liver disease". In: *Am. J. Gastroenterol* 104 (2009), pp. 2196–2205.

[50] Koprulu M. "Identification of Rare Loss-of-Function Genetic Variation Regulating Body Fat Distribution". In: *J. Clin. Endocrinol. Metab* 107 (2022), pp. 1065–1077.

[51] Xiao Y. "Deficiency of PRKD2 triggers hyperinsulinemia and metabolic disorders". In: *Nat. Commun* 9 (2018), p. 2015.

[52] Langer V. "IFN-γ drives inflammatory bowel disease pathogenesis through VEcadherin-directed vascular barrier disruption". In: *J. Clin. Invest* 129 (2019), pp. 4691–4707.

[53] Ouyang X. "Transcription factor IRF8 directs a silencing programme for TH17 cell differentiation". In: *Nat. Commun* 2 (2011), p. 314.

[54] Mencarelli A. "Highly specific blockade of CCR5 inhibits leukocyte trafficking and reduces mucosal inflammation in murine colitis". In: *Sci. Rep* 6 (2016), p. 30802.

[55] Kučerová L. "Mitochondrial neurogastrointestinal encephalomyopathy imitating Crohn's disease: a rare cause of malnutrition". In: *J. Gastrointestin. Liver Dis* 27 (2018), pp. 321–325.

[56] Hwang S.-H. "Leukocyte-specific protein 1 regulates T-cell migration in rheumatoid arthritis". In: *Proc. Natl. Acad. Sci. U. S. A* 112 (2015), E6535–6543.

[57] Shi X, Yang C, and Liu J. "Using Collaborative Mixed Models to Account for Imputation Uncertainty in Transcriptome-Wide Association Studies". In: *Methods Mol. Biol* 2212 (2021), pp. 93–103.

[58] Berisa T and Pickrell J. "Approximately independent linkage disequilibrium blocks in human populations". In: *Bioinformatics* 32 (2016), pp. 283–285.

[59] Jon Wakefield. "Bayes factors for genome-wide association studies: comparison with P-values". In: *Genetic Epidemiology: The Official Publication of the International Genetic Epidemiology Society* 33.1 (2009), pp. 79–86.

[60] Xiang Zhu and Matthew Stephens. "Bayesian large-scale multiple regression with summary statistics from genome-wide association studies". In: *The annals of applied statistics* 11.3 (2017), p. 1561.

[61] F Kyle Satterstrom et al. "Large-scale exome sequencing study implicates both developmental and functional changes in the neurobiology of autism". In: *Cell* 180.3 (2020), pp. 568–584.

[62] Bulik-Sullivan B. "LD Score regression distinguishes confounding from polygenicity in genome-wide association studies". In: *Nat. Genet* 47 (2015), pp. 291–295.

[63] Ben Elsworth et al. "The MRC IEU OpenGWAS data infrastructure". en. Aug. 2020.

[64] De Lange K. "Genome-wide association study implicates immune activation of multiple integrin genes in inflammatory bowel disease". In: *Nat. Genet* 49 (2017), pp. 256–261.

[65] Willer C, Li Y, and Abecasis G. "METAL: fast and efficient meta-analysis of genomewide association scans". In: *Bioinformatics* 26 (2010), pp. 2190–2191.

[66] Pardiñas A. "Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection". In: *Nat. Genet* 50 (2018), pp. 381–389.

[67] Benner C. "Prospects of Fine-Mapping Trait-Associated Genomic Regions by Using Summary Statistics from Genome-wide Association Studies". In: *Am. J. Hum. Genet* 101 (2017), pp. 539–551.

[68] Pasaniuc B. "Fast and accurate imputation of summary statistics enhances evidence of functional enrichment". In: *Bioinformatics* 30 (2014), pp. 2906–2914.

[69] Lee A and Jung I. "Functional annotation of lung cancer-associated genetic variants by cell type-specific epigenome and long-range chromatin interactome". In: *Genomics Inform* 19 (2021), e3.

[70] Trubetskoy V. "Mapping genomic loci implicates genes and synaptic biology in schizophrenia". In: *Nature* 604 (2022), pp. 502–508.

[71] Giri A. "Trans-ethnic association study of blood pressure determinants in over 750,000 individuals". In: *Nat. Genet* 51 (2019), pp. 51–62.

[72] Jung I. "A compendium of promoter-centered long-range chromatin interactions in the human genome". In: *Nat. Genet* 51 (2019), pp. 1442–1449.

[73] Chen E. "Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool". In: *BMC Bioinformatics* 14 (2013), p. 128.

[74] Kuleshov M. "Enrichr: a comprehensive gene set enrichment analysis web server 2016 update". In: *Nucleic Acids Res* 44 (2016), W90–97.

[75] De Leeuw C et al. "MAGMA: generalized gene-set analysis of GWAS data". In: *PLoS Comput. Biol* 11 (2015), e1004219.

[76] Watanabe K et al. "Functional mapping and annotation of genetic associations with FUMA". In: *Nat. Commun* 8 (2017), p. 1826.

Figure 1. Overview of the cTWAS method.
A. Two scenarios that violate the assumptions of TWAS and lead to false positive findings. *X* is a non-causal gene of the trait *Y*. *X* is associated with the trait, shown as dashed arrows, because of the LD between its eQTL and nearby causal variants. Double-headed arrows represent LD between variants. **B.** The causal diagram implicitly assumed by TWAS. $\tilde{X}$ represents the cis-genetic component of gene expression *X*. **C.** The model of cTWAS. G_m the genotype of the mth variant; $\tilde{X_j}$ and X_j imputed expression and actual expression of the jth gene; θ_m, effect size of the mth variant on the trait *Y*; β_j, effect size of the jth gene; ϵ, error term; δ_0, point mass at 0. π_G and π_V, prior probability of being casual for genes and variants, respectively; σ^2_G and σ^2_V, variance of the effect sizes of causal gene and variants, respectively. **D.** The workflow of cTWAS. The main steps of cTWAS are shown in the red boxes. cTWAS reports PIP for all genes and variants within LD blocks. *p* values for the genes and variants from marginal association tests are shown at the top as a comparison. Red dashed lines from the output panel indicate genome-wide significance level or the PIP threshold for genes.

Figure 2. Results of parameter estimation and PIP calibration in simulations.
A. Accuracy of the estimated parameters related to gene effects. Each plot shows one parameter: π_G, prior probability for a gene being causal; enrichment (π_G/π_V), where π_V is prior probability for a SNP being causal; effect size for gene and PVE of gene. Results from two simulation settings are shown, the high and low gene PVE settings. Each dot represents the result from one out of five simulations. Horizontal bars show the true parameter values. **B.** Gene PIP calibration. Gene PIPs from all simulations are grouped into bins. The plot shows the proportion of true causal genes (Y-axis) against the average PIPs (X-axis) under each bin. A well calibrated method should produce points along the diagonal lines (red). +/- standard error is shown for each point in the vertical bars.

Figure 3. Comparison of cTWAS with other methods in simulations.

A. Comparison of standard TWAS and cTWAS. Gene PIPs from cTWAS are plotted against the marginal *p* values (-log10 scale) from TWAS. The results were from one simulation run (parameters: gene PVE 0.052, gene prior 0.025, SNP PVE 0.50, SNP prior 0.00025). Each dot represents a gene and is colored based on whether it is a causal gene. **B**. Number of genes identified by various methods. We used the suggested significance cutoff for each method: PIP > 0.8 for cTWAS, Bonferroni corrected *p* < 0.05 for FUSION, PP4 > 0.8 for coloc, PIP > 0.8 for FOCUS, FDR < 0.05 for SMR and HEIDI *p* < 0.05. We use different colors for causal genes identified by each method, and the top gray bars indicate noncausal genes. **C** and **D**. Examples of how cTWAS avoided false positive genes. Each square represents a gene and its position was determined by its transcription start site. Colors indicate whether the gene or SNP is causal (orange), noncausal but in LD with a causal effect (R^2 between SNP genotype or imputed expression > 0.4, green), or noncausal and not in LD with causal effect (purple). The eQTLs of the genes are plotted in middle tracks. Transcriptome-wide significance cut off for TWAS was indicated by the red dotted line. In the top panel, the values of PP4 (probability of colocalization) from coloc analysis were shown for each gene of interest.

Figure 4. cTWAS avoids false discoveries of candidate genes of LDL cholesterol.
A. Precision of cTWAS and TWAS in distinguishing LDL silver standard genes from nearby bystander genes. **B.** cTWAS results at the *HPR* locus. The top track depicts the -log10 p-value of variants from GWAS and genes from TWAS. Each square represents a gene with position determined by its transcription start site. Each circle represents a variant. Colors indicate LD between the focal gene (orange) and nearby genes and variants: high LD (purple; R2 > 0.4) and low LD (green, R2 <= 0.4). The red dotted line indicates the transcriptome-wide significance threshold for TWAS (Bonf.). The second track represents the positions of the eQTL for the focal gene. The third track represents the cTWAS PIPs for variants and genes at this locus. The fourth track depicts the gene bodies of all genes in the region. **C.** cTWAS results at the *POLK* locus. Figure legend is the same as in Figure 4B. **D.** Fine-mapping for the locus around *HMGCR* and *POLK* genes. The top two tracks represent the -log10 p-value of variants (with color representing LD with the lead SNP) and their PIPs from fine-mapping with PolyFun-SuSiE (with color representing credible sets). Only SNPs with reported PIPs were shown in the plot. The third track represents liver H3K4me1 peak calls from ENCODE. The bottom track shows interactions identified from liver promoter-capture HiC (PC-HiC) data. The links in red highlight regions looped to the *HMGCR* promoter. **E.** Sources of confounding for TWAS false positive findings. A TWAS gene was considered false positives if its cTWAS PIP ≤ 0.5. Only genes that can be assigned to a credible set were included in the analysis.

Figure 5. Candidate genes and pathways for LDL discovered by cTWAS.
A. PIPs of 9,881 liver genes from cTWAS analysis of LDL cholesterol. The labeled genes have PIP > 0.8, colored based on existing evidence for LDL function: silver standard genes ("Known" genes; green); nearest genes to a genome-wide significant locus for IBD ("Nearest" genes) but not in silver standard (orange); or otherwise novel ("Novel" genes; purple). **B.** GO Biological Process terms enriched among 35 detected LDL liver genes. 28 GO terms were significantly enriched at FDR < 0.05. Redundant terms were omitted for clarity (see Methods).**C.** cTWAS results at the *ACVR1C* locus. Figure legend is the same as in Figure 4B and 4C. **D.** Summary of cTWAS outcomes for all 69 silver standard genes. Silver standard genes were binned into the following categories: detected by cTWAS at PIP > 0.8 ("Detected (PIP > 0.8)"); significant by TWAS, but not detected by cTWAS ("Significant Z-score but Undetected"); insignificant by TWAS and not detected by cTWAS, with a genome-wide significant GWAS variant within 500kb ("Insignificant z-score"); gene does not have no expression prediction model ("Not imputed"); not detected by cTWAS and TWAS, and no genome-wide significant GWAS variant within 500kb ("No GWAS Signal").

Figure 6. cTWAS analysis of IBD and other traits using all GTEx tissues.
A. Estimated percentages of heritability mediated by eQTLs for IBD using MESC (X-axis) and cTWAS (Y-axis). Each dot represents the result from one GTEx tissue. The black dotted line denotes equivalence between the methods, and the blue dashed line denotes the slope relating cTWAS and MESC mediated heritability estimates. **B.** The number of cTWAS genes detected at PIP > 0.8 in the top two tissues for IBD, SBP, and SCZ. Top tissues per trait are determined by the number of detected genes. **C.** The number of cTWAS genes detected for IBD at PIP > 0.8 across major tissue groups. A gene was detected in a tissue group if it was detected in any of the tissues in that group. A gene was considered novel if it was not a silver standard gene ("Known") or if it was not the nearest gene to a genome-wide significant locus for IBD ("Nearest"). **D.** The number of tissues with cTWAS PIP > 0.5 for 56 IBD genes detected at PIP > 0.8 in the "Blood/Immune" or "Digestive" tissue groups. **E.** Non-redundant GO terms enriched among 56 detected IBD genes in the "Blood/Immune" or "Digestive" tissue groups. These terms were found using the Weight Set Cover method from WebGestalt. 29 out of 56 genes were associated with at least one of the GO terms. **F.** cTWAS results for IBD at the *UBE2W* locus using the "Colon Transverse" eQTL data. Figure legend is the same as the previous locus plots.

Gene	Position	Max PIP Tissue	Max PIP	Tissue Z-score	Other Tissues Detected	Evidence
MRPL20	chr1:1401909	Whole Blood	0.938	5.45		Novel
ERRFI1	chr1:8004404	Whole Blood	0.938	6.55		Nearest
RNF186	chr1:19814029	Colon Transverse	0.988	-7.27		Known
OAZ3	chr1:151762899	Colon Sigmoid	0.995	5.14		Novel
ADAM15	chr1:155050566	Stomach	0.917	-5.56	Cells EBV-transformed lymphocytes; Spleen; Whole Blood; Esophagus Muscularis; Small Intestine Terminal Ileum	Novel
CD244	chr1:160830160	Esophagus Mucosa	0.971	-5.59		Novel
FCGR2A	chr1:161505430	Esophagus Mucosa	0.961	9.18	Colon Sigmoid; Small Intestine Terminal Ileum	Nearest
FOSL2	chr2:28392448	Esophagus Gastroesophageal Junction	0.962	-7.39		Nearest
ZFP36L2	chr2:43222402	Spleen	0.943	-6.65		Novel
IL1R2	chr2:101991960	Colon Transverse	0.967	6.25		Novel
CCL20	chr2:227805739	Colon Sigmoid	0.968	5.19		Novel
CCR5	chr3:46370946	Whole Blood	0.925	-4.54		Novel
CD200R1	chr3:112921205	Whole Blood	0.946	-4.20		Novel
RASA2	chr3:141487027	Stomach	0.914	4.48	Esophagus Gastroesophageal Junction; Esophagus Muscularis	Novel
RGS14	chr5:177357924	Esophagus Muscularis	0.948	-6.30	Colon Sigmoid; Esophagus Gastroesophageal Junction; Small Intestine Terminal Ileum	Nearest
UBE2W	chr8:73780097	Colon Transverse	0.934	-4.18		Novel
TNFSF15	chr9:114784652	Esophagus Muscularis	1.000	-11.16	Spleen; Whole Blood	Nearest; Known
CARD9	chr9:136363956	Spleen	0.997	12.60	Whole Blood; Esophagus Muscularis	Novel
LSP1	chr11:1850904	Esophagus Muscularis	0.946	5.20	Spleen; Colon Sigmoid	Novel
EFEMP2	chr11:65866441	Stomach	0.939	4.90	Colon Sigmoid; Esophagus Gastroesophageal Junction	Novel
SOCS1	chr16:11254417	Whole Blood	0.961	-5.63		Novel
BRD7	chr16:50313487	Whole Blood	0.977	-6.86		Novel

IRF8	chr16:85899116	Colon Transverse	0.951	6.54		Nearest
CASC3	chr17:40140318	Esophagus Mucosa	0.981	-5.97	Colon Sigmoid	Novel
STAT3	chr17:42313324	Whole Blood	0.916	8.20		Nearest
SBNO2	chr19:1107637	Esophagus Mucosa	0.914	4.44		Nearest
PRKD2	chr19:46674275	Colon Transverse	0.926	-4.48		Novel
OSER1	chr20:44195939	Esophagus Mucosa	0.911	-4.75	Spleen	Novel
IFNGR2	chr21:33403413	Stomach	0.981	5.80	Colon Transverse	Nearest
TYMP	chr22:50525752	Esophagus Mucosa	0.942	-4.34		Novel

Table 1. IBD genes detected by cTWAS in the Blood/Immune and Digestive tissue groups. Genes detected by cTWAS at PIP > 0.9 in at least one of the tissues in the two tissue groups. The Max PIP Tissue, Max PIP, and Z-score columns denote the tissue with the highest PIP for each gene and its corresponding PIP and Z-score from TWAS in that tissue. The Other Tissues column lists any additional tissues with cTWAS PIP > 0.8. The Evidence column denotes whether each gene is in the silver standard gene list ("Known"); the nearest gene to genome-wide significant GWAS peak for IBD ("Nearest"); or otherwise novel ("Novel").