A spiking neural network model of cortical intraregional metastability

Siva Venkadesh[1, +], Asmir Shaikh[2], Heman Shakeri[3,4], Ernest Barreto[5, *], John D. Van Horn[1,3, *]

[1]Department of Psychology, University of Virginia, Charlottesville, VA – 22904

[2]Department of Computer Science, University of Virginia, Charlottesville, VA – 22904

[3]School of Data Science, University of Virginia, Charlottesville, VA – 22904

[4]Biomedical Engineering, University of Virginia, Charlottesville, VA – 22904

[5]Department of Physics and Astronomy, George Mason University, Fairfax, VA – 22030

[+]Correspondence: si9xr@virginia.edu

304 Gilmer Hall,

Department of Psychology,

University of Virginia,

Charlottesville, VA – 22904

*Co-senior authors

Abstract:

Transient synchronization of bursting activity in neural networks, which occurs in patterns of metastable phase relationships between neurons, is a notable feature of network dynamics observed *in vivo.* However, the mechanisms that contribute to this dynamical complexity in neural circuits are not well understood. Local circuits in cortical regions consist of populations of neurons with diverse intrinsic oscillatory features. In this study, we numerically show that the phenomenon of transient synchronization, also referred to as metastability, emerges in an inhibitory neural population when the neurons' intrinsic fast-spiking dynamics are appropriately modulated by slower inputs from an excitatory neural population. Using a compact model of a mesoscopic-scale network consisting of excitatory pyramidal and inhibitory fast-spiking neurons, our work demonstrates a relationship between the frequency of neural oscillations and the features of emergent metastability. In addition, a novel metric is formulated to characterize collective transitions in metastable networks. Finally, we discuss a blueprint to model the whole-brain resting-state dynamics using our scalable representation of intraregional network metastability.

Keywords: transient attractors, itinerancy, bursting dynamics, emergent complexity, network connectivity

1. Introduction

Computational modeling of neuronal dynamics which incorporates mesoscopic-level connectivity features can potentially offer powerful frameworks for investigating neural network mechanisms [1,2]. However, a scalable network representation of key neurodynamical features at the mesoscopic level is currently lacking. This is primarily due to limited theoretical accounts for the emergent dynamical features in a network in terms of its constituent mechanisms. A simplified model describing the complexities of between- and within-group neuronal interactions is necessary to advance our multiscale understanding of neural systems in typical and atypical conditions [3].

An experimentally observable complexity of *in vivo* networks is the metastable attractor dynamics at the level of individual neurons. More specifically, a neuron can exhibit bursting oscillations that transiently synchronize with the oscillations of other neurons, and these synchronizations can occur at different phases [4–9]. In other words, biological neural networks can realize a range of attractor states that are locally observable through different phase-locking modes between individual neurons. Such transient attractor dynamics are also referred to as metastability or itinerancy [10], and this phenomenon is hypothesized to be a necessary physical property underlying the coordinated dynamics between spatially distant neural populations [11]. Moreover, a moment of conscious experience is hypothesized to exist irresolvably (i.e., indivisible in time) for the duration between successive attractor transitions in the brain [12].

The mathematical conditions for the emergence of transient dynamics have been previously discussed [13]. More recently, conduction delays between brain areas were associated with metastable oscillations on a macroscopic spatial scale [14]. However, the biophysical features of neural networks underlying metastability at the level of neurons have not been fully elucidated.

While a previous work reported metastable attractor dynamics among intrinsically bursting

neurons [10], it was unclear if metastability could also be realized in neural populations in which

bursting dynamics are induced externally. Here, we explore the emergence and characteristics of

metastability in a fast-spiking interneuron population that exhibits bursting dynamics due to

driving inputs from a population of slower pyramidal neurons. We also examine the quantitative

characteristics of collective transitions between attractor states realized in different metastable

networks. Finally, we discuss potential applications of our work in modeling the whole-brain

resting-state dynamics to understand network physiology in neurodegenerative disorders.

2. Methods

64   Networks consisting of 100 pyramidal neurons and 50 fast-spiking interneurons were constructed

65   using Izhikevich neurons [15] (Equations $1 - 3$). The parameters $a, b, c,$ and d (Equations $2 - 3$)

$$\frac{dv}{dt} = 0.04v^2 + 5v + 140 - u + I^{ext} + I^{syn} \tag{1}$$

$$\frac{du}{dt} = a(bv - u) \tag{2}$$

$$if\ v \geq 30mV, \begin{cases} v = c \\ u = c + d \end{cases} \tag{3}$$

$$I^{syn} = \sum_{i=1}^{m_j} \delta_i \cdot w \tag{4}$$

were selected to match the intrinsic dynamics of each type of neuron. Specifically, these were set

to 0.02, 0.2, -65, and 8 respectively for the pyramidal neurons, and 0.1, 0.2, -65 and 2 for the fast-

spiking interneurons [15].

Network connections were configured as
illustrated in **Fig. 1A**. Inter- and intra-population
connections were specified using pairwise connection
probabilities between individual neurons. We studied
the network behavior for a complete range of
connection probabilities. Post-synaptic effects were
specified using an instantaneous pulse-coupling
scheme (Equation – 4), where each presynaptic spike
either increased (excitatory) or decreased (inhibitory)
the postsynaptic current by the weight parameter w
(+0.3 for excitatory connections and −0.3 for inhibitory
connections). In Equation – 4, δ_i is a function which
equals one when neuron i spikes and is zero otherwise.
The number of neurons and the weight parameters were
chosen to allow the pyramidal population to induce
sustained bursting activity in the fast-spiking
interneuron population for a range of external input
currents in the former (I^{ext} in Equation – 1). I^{ext} was
always set to zero for the fast-spiking interneurons. All
networks were simulated using Brian2 [16] for a
duration of 10s with a timestep of 0.1ms. Following the
network simulation, a low-pass filter was applied to the
voltage signal of each neuron (v in Equation – 1) to

Figure 1. A. *Left:* Network configuration consisting of pyramidal neurons (blue triangles) and fast-spiking interneurons (dark orange ovals). Connections indicated by the arrowhead and circles denote excitatory and inhibitory pulse-coupling, respectively. *Right:* Spiking frequencies of an isolated excitatory (pyramidal) neuron and an inhibitory (fast-spiking) neuron for various input currents. **B.** Raster plot showing spiking activity in pyramidal (blue) and fast-spiking neurons (orange) for one second. **C.** (top) Bursting activity (gray) in one representative fast-spiking neuron. The filtered signal, superimposed in black, captures the burst oscillations. (bottom) Burst oscillations of two representative neurons (black and red) showing transitions between in- and anti-phase transiently locked modes.

92 extract the bursting oscillations without the individual spikes (**Fig. 1C top**). Then, instantaneous

93 phases of the bursting oscillations were assigned to each neuron via Hilbert transform. **Fig. 1C**

94 **bottom** shows two representative examples.

2.1. Assessing the stability of phase relationships

To assess the nature of population-wide phase relationships among the interneurons (our

focus of interest), we calculated the phase differences between neuron pairs over time. This

resulted in an ensemble of phase difference measurements between all possible distinct pairs of

neurons ((50*49)/2=1225 pairs) at each time point over the entire duration (10s). Since time is

discrete with a resolution of 0.1ms, this ensemble consisted of N=122,500,000 phase angle

difference measurements in total. Henceforth, these angles will be referred to as ϕ_j, $j = 1,2,...,N$.

We found that the ensemble of angles ϕ_j was organized into clusters. To characterize this, we

calculated the Kuramoto-Daido order parameters Z_n [10,17,18]:

112
$$Z_n = \frac{1}{N} \sum_{j=1}^{N} e^{in\phi_j} \tag{5}$$

for n, a positive integer. These order parameters characterize clustering patterns in the ensemble

as follows. If each ensemble member is plotted as a point on the unit circle in the complex plane

at its angle, ϕ_j, then Z_1 is the centroid of the ensemble. Of interest is the magnitude $|Z_1|$, which

ranges from zero to one. If all points occur at the same angle, then $|Z_1| = 1$. In contrast, if the

points are scattered uniformly about the unit circle, then $|Z_1| = 0$. Additionally, if the ensemble of

points consists of uniformly spaced clusters, then $|Z_1| = 0$. Such arrangements can be identified

using the order parameters Z_n with $n > 1$. For example, if the phases form three ideal clusters (i.e.,

equally populated clusters 120 degrees apart), then $|Z_1| = |Z_2| = 0$, but $|Z_3| = 1$. In fact, for ideal

states consisting of c uniformly spaced and equally populated clusters, $|Z_{jc}| = 1$ for any integer

$j \geq 1$ and is zero otherwise. To correct for this redundancy, we defined a real-valued quantity G_n

$$G_n = |Z_n| \times \prod_{j=1}^{n-1}(1 - |Z_j|) \tag{6}$$

115   as in Equation 6, such that G_n is close to 1 only when the ensemble approximates an ideal n-cluster

116   state.

2.2. Characterizing exogenous transitions

118   To assess the nature of transitions between clustered states in different metastable

119   networks, the following steps were carried out: The networks were simulated twice with identical

120   initializations. The first simulation was used to establish the baseline instantaneous clustering of

121   neurons at any given time (i.e., clustering of neurons based on their instantaneous phases). In the

$$RI = \frac{g + h}{nC_2} \tag{7}$$

$$ARI = \frac{RI - E(RI)}{\max(RI) - E(RI)} \tag{8}$$

$$Collective\ response = 1 - \frac{1}{M}\sum_{i=1}^{M} ARI_i \tag{9}$$

second simulation, a fraction of randomly selected interneurons was simultaneously perturbed by

single spikes at a certain timepoint. This caused some of the neurons to change their cluster

assignments relative to the baseline. Phase cluster assignments of neurons were obtained using

Gaussian mixture models [19]. To quantify the effect of the perturbation, we measured the

similarity of the two clustering arrangements of neurons between the baseline and perturbed

conditions over time using Equations $7 - 9$. Here, RI is the "rand index," which measures the

similarity of two sets of instantaneous phase clusters, and g and h are the numbers of pairs of

neurons with the same and different cluster assignments across two groupings (i.e., the number of

pairwise agreements between two clustering arrangements). nC_2 denotes the number of possible

combinations of neuron pairs. Thus, the RI for two identical groupings of a set of neurons is one.

The adjusted rand index (ARI) defined in Equation – 8 [20–22] was used to account for chance,

and it ensures that the similarity of two random groupings of a set of neurons is zero. This is

achieved by estimating an expected RI ($E(RI)$ in Equation – 8) using random permutations of

cluster assignments.

We defined the population's collective response as in Equation – 9 to characterize the

extent to which the network (as a whole) responds to exogeneous perturbations of a fraction of its

elements. The ARIs were measured in 50 simulation trials ($M = 50$ in Equation – 9), each with a

random subset of perturbed interneurons and a random time (>0.5s) at which the perturbation was

delivered. The observed average change in ARI following the perturbation denoted the effect of

the perturbation. For instance, if there is no change to the network state following the perturbation,

then the collective response is zero. On the other hand, if a perturbation results in new phase

clusters that are equivalent to completely random cluster assignments for all the neurons compared

to the baseline condition, then the collective response is one. Finally, the magnitude of collective

response was examined in metastable networks for various numbers of perturbed neurons. It is

worth mentioning here that while each presynaptic spike influences the activity of a postsynaptic

neuron (see Equation 4), the collective response aims to characterize activity that is

spatiotemporally coarse-grained (i.e., transitions that occur among a group of neurons on a

relatively larger timescale than that of the individual spikes).

3. Results

The simulated network model incorporates a significant difference in the spiking time scales of the pyramidal and fast-spiking interneurons. The slower excitatory drive from the pyramidal neurons modulates the fast-spiking dynamics of the interneurons, resulting in bursting oscillations in the latter (**Fig. 1B&C**). The inhibitory interactions between these bursting neurons then leads to itinerant metastable behavior. That is, ongoing endogenous transitions occur between different phase-locked (attractor) states. An example is illustrated in **Fig. 1C**, which shows the bursting oscillations of a representative pair of neurons versus time. At first, the neurons are locked in phase, but then they transition to an out-of-phase state. After a short time, they return to the in-

Figure 2. The network-wide stability and the number of phase clusters in the fast-spiking interneurons as functions of pyramidal neuron frequency. Polar histograms show phase differences ϕ_j for all pairs of neurons in representative networks. Pyramidal neuron frequency was computed by taking the inverse of the average population interspike intervals.

phase state, and then switch back to the out-of-phase state. Similar persistent behavior was

described in a previous report [10] (also see Supplementary **Fig. S1**).

3.1. Frequency dependence of phase cluster stability

The frequency of pyramidal neuron spiking is determined by the value of I^{ext}. To examine

the effect of this driving frequency on the characteristics of metastability in the fast-spiking

interneuron network, we systematically varied I^{ext} and measured the distribution of phase

differences (ϕ_j) between pairs of interneurons as described in Methods **Section 2.1**. **Fig. 2** shows

a plot of G_n (n=1, 2, …, 7), which quantifies the degree to which the interneuron phase differences

group into n clusters, versus the pyramidal neuron spiking frequency (f). Above the plot are polar

histograms of the phase differences (ϕ_j) for selected f's. At low driving frequency, all the

inhibitory neurons lock in-phase, and thus their phase differences are close to zero.

Correspondingly, G_1 is high and G_2 through G_7 are low. Interestingly, as the driving frequency is

increased beyond ~25Hz, clusters appear in the phase difference distribution. We find that a well-

defined state of two clusters emerges near f=48Hz, and a three-cluster state emerges near f=77Hz.

As the driving frequency is further increased, the pattern reverses and another well-defined state

of two clusters appears near f=117Hz. Beyond f=140Hz, all G_n are small and the phase difference

distributions become more uniform. It is interesting to note that the emergence of these clustered

states occurs for driving frequencies within the gamma range. Pyramidal neurons are indeed

known to fire at gamma frequencies, although such gamma cycles have been observed to be

modulated by slower theta cycles [23,24]. These results show that the metastable attractors emerge

in an interneuron population when their spiking dynamics are appropriately driven by slower

excitatory frequencies.

Figure 3. The stability of phase clusters against network connectivity and pyramidal neuron frequency. **A.** The most stable two-cluster (left) and three-cluster (right) states realized in networks for various excitatory-inhibitory (E-I) and inhibitory-inhibitory (I-I) connection probabilities . **B.** Distributions of maximum phase cluster stability are plotted against their respective frequency bins of pyramidal neurons for two-cluster (left) and three-cluster (right) states. Width indicates the density of points for a given maximum stability.

182   The phase-clustering phenomena described above were consistently observed for a range

183   of connection probabilities (**Fig. 3**). We analyzed networks instantiated on a 50x50 grid of inter-

184   and intra-group connection probabilities in the range [0, 1]. The G_2 and G_3 versus frequency curves

185   were computed as described before for each of the 2500 networks. **Fig. 3A** shows the maximum

186   of G_2 (left) and G_3 (right) across frequencies for each network. These measures were sensitive to

187 the inter-group connection probabilities and relatively less sensitive to the intra-group connection

188 probabilities. Plotting these measures against the frequency of excitatory neurons (**Fig. 3B**)

189 revealed that the stable phase clusters emerged approximately within the range [25*Hz*, 200*Hz*].

190 Our results suggest that the emergent attractors' stability in interneurons depend on their

191 connectivity with excitatory neurons and the frequency of excitatory oscillations. There are robust

192 gradients towards maximally stable interneuron phase cluster arrangements along the dimensions

193 of pyramidal frequency and E-I connection probability (**Figs. 2&3**).

194 3.2. Collective response characteristics of metastable networks

Figure 4. Exogenous transitions in the fast-spiking interneuron population. **A.** Left and right panels show the temporal evolution of a network in unperturbed and perturbed conditions respectively. Neurons (black dots) in the left panels are sorted by their instantaneous phases at each t, and their order is preserved in the right panels to show the divergence of states caused by perturbation. t_p denotes the time of perturbation, and red circles in the right middle panel denote perturbed neurons. **B.** Ten representative ARI curves (color) computed by comparing cluster assignments of neurons between unperturbed and perturbed conditions. The black curve represents the average. Here, t=0 corresponds to the perturbation times t_p.

195   The multi-cluster states described above exhibit spontaneous and endogenous itinerant

196 dynamics (**Fig. 1C** and Supplementary **Fig. S1**). Motivated by the fact that an organism adapts its

197 internal state to the environment via external stimuli, we sought to measure how a metastable state

198 would be affected by an external perturbation. As described in Methods **Section 2.2**, the network

was simulated twice with identical initializations. The first simulation established the baseline

dynamics as shown in **Fig. 4A** (left, all rows). The panels on the left show the instantaneous phases

of individual neurons in the interneuron population at three instants of time. The neurons are

ordered along the horizontal axis by increasing phase. To mask the endogenous itinerant behavior

(see Supplementary **Fig. S1**), the neurons were reordered for each left panel in **Fig. 4A**. In the

second simulation, an external perturbation was applied. Five randomly selected interneurons each

received a single spike at time t_p. The results are shown in the right panels of **Fig. 4A**. The time

of perturbation occurred between the top and middle panels. Note that the neuron orderings from

the left columns are preserved in the right columns. Thus, any differences in the cluster assignment

of neurons are due exclusively to the perturbation. Additionally, the transitions are collective in

the sense that they occur in many non-perturbed neurons immediately following the perturbation

(**Fig. 4A** - right). This highlights the synergy of interacting elements in metastable networks.

Collective transitions may be characterized by treating network states as multiple instantaneous

phase clusters of neurons at any given time and subsequently comparing the cluster assignments

of neurons between unperturbed and perturbed conditions using the ARI (**Fig. 4B**). For example,

when $t < t_p$, the two simulation conditions (left and right panels of **Fig. 4A**) show identical cluster

arrangements of neurons resulting in ARI values of 1. However, immediately following

perturbation ($t > t_p$), the network states begin to diverge between the two simulation conditions

resulting in ARI values less than 1 (**Fig. 4B**).

Fig. 5 shows the magnitude of collective responses
(Equation – 9) in three representative metastable networks
that showed high magnitudes of stability in **Fig. 2**. The two-
cluster states that emerged near f =48Hz in Fig. 2 (see Section
3.1) showed the strongest response, reaching magnitudes of
0.4 within 1s when sufficient number of neurons were
perturbed (**Fig. 5A**). On the other hand, the three-cluster
states, which emerged near f =77Hz, elicited slower responses
to external perturbations (**Fig. 5B**), and the two-cluster states
that emerged near f =117Hz elicited the slowest response
(**Fig. 5C**). Note that f directly affects the inter-burst intervals
in interneurons and consequently shapes their multi-periodic
trajectories in the phase space. Taken together, these results
suggest that the metastability realized in interneuron
populations may arise from mechanisms [13,25,26] that
evolve depending on (a) their connectivity with excitatory
neurons and (b) the frequencies of excitatory oscillations
within a cortical region.

4. Discussion

Figure 5. Magnitude of collective response (equation – 9) characterized by the exogenous transitions in three representative networks for various numbers of perturbed neurons. The pyramidal driving frequencies used are 48Hz (A), 77Hz (B), and 117Hz (C); these correspond to two, three, and two cluster states in the interneuron population, respectively (see Fig. 2).

The modern examination of neural connectivity has enjoyed considerable attention through several
early key theoretical studies [27–30]. Along these lines, understanding the emergent dynamical
complexities in a network in terms of its constituent mechanisms remains an important goal. Here,
we studied the emergent phenomenon of metastability in spiking neural network models that

represent cortical local circuits. Metastability is characterized by the coexistence of integrated

(synchronized states) and segregated (transitions between synchronized states) behaviors in a

system of interacting elements [11]. We specifically studied how the neuron-level spiking

frequencies are related to the emergent metastability, and how these relationships are governed by

the underlying excitatory and inhibitory neuronal connectivity. Moreover, the spiking neural

network models examined in this paper provide compact mesoscopic scale approximations of

regional collective dynamics that capture a crucial biological complexity at the level of individual

neurons. A limitation of the current work is that it only considered two types of neuron populations,

whereas intraregional neural circuits consist of populations with diverse frequency profiles [31].

Nevertheless, current work demonstrates that the emergent metastability in an interneuron

population depends on their connectivity with excitatory neurons and the frequency of excitatory

drive. Their relationships are revealed in the G_n curves that show robust gradients along the

dimensions excitatory-inhibitory connection probability (**Fig. 3A**) and the frequency of the

excitatory drive (**Figs. 2 & 3B**). Thus, a numerical optimization using G_n as an objective function

can, in principle, estimate optimal intraregional connectivity configurations among groups of

neurons with heterogenous frequency profiles.

Additionally, the models and the analyses methods presented in this paper are easily

scalable to study interregional interactions in the brain. Empirically estimated anatomical

connectivity between brain regions provides useful information for modeling interregional

metastability. Note that while the current work suggests a link between two temporal scales (i.e.,

timescales of neuronal spiking and attractor transitions) of network dynamics, modeling

interregional interactions using their anatomical connectivity will further provide a venue to study

metastable brain dynamics at a macroscopic spatial scale. In the next section, we present a design

for whole-brain modeling and discuss biological observations of atypical brain physiology that can

be mechanistically interpreted in such models.

4.1. A blueprint for whole-brain modeling at the mesoscopic level:

Sustained background activity in the cortex arises from the mutual excitation of excitatory neurons

[32] such as the pyramidal neurons that project their axons to longer distances than the

interneurons. Such long-distance projections are captured in diffusion-weighted magnetic

resonance imaging (dMRI) as white matter tracts. Estimation of anatomical connectivity from

dMRI relies on the tractography-derived streamline counts between regions [33]. However, the

direction of axonal projections is not identifiable from the dMRI tractography. In other words, if

the tractography estimates that there are n streamlines between regions A and B, the fraction of n

that corresponds to the axonal projections from region A to B (and vice versa) is unidentifiable

based on the tractography alone. The net excitatory input to a region is a key determinant of its

firing frequency. Since the regional metastable dynamics may depend upon the frequency of

excitatory oscillations (See **Figs. 2&5**), we consider that resolving the directionality of fiber tracts

(i.e., incoming vs. outgoing for a region) will be important to accurately model the whole-brain

metastable dynamics.

Previous whole-brain modeling studies considered symmetric couplings between pairs of

brain regions and estimated a global coupling constant to supplement the connectivity information

derived from white matter tractography [34,35]. The unknown degrees of asymmetries in structural

connections could be estimated for the whole brain, in principle, by numerical optimization. The

network-wide stability of attractors (Equation – 6) realized across the frequency spectrum in

whole-brain models could serve as a heuristic metric to search for the directionality, or more

generally, the asymmetries in the bidirectionality, of macroscopic anatomical connections. Note

that the number of unknown parameters is directly related to the spatial resolution of structural

connectivity specification. It is uncertain if a model description at a high spatial resolution is

solvable in practice due to the computational demands. However, models can be described at

various spatial resolutions of a group-averaged whole-brain connectome, and the convergence

patterns of optimization solutions for each spatial resolution can be examined. These convergence

patterns should be compared against those of the null models with randomly shuffled anatomical

connections. The robustness of the solution convergence relative to that of the respective null

model can inform the preferred spatial resolution for the whole-brain models. This approach

considers that the brain-wide features of the human connectome evolved to be highly specific, and

they provide strong constraints to model network-wide transient attractor dynamics. The identified

directionalities can also be validated against findings from the morphological tracing studies that

report the directions of axonal projections between cortical regions using anterograde or retrograde

tracing techniques [36].

The whole-brain models of anatomically constrained mesoscopic scale metastable

dynamics are useful mathematical tools that will enable a multiscale understanding of cortical

dynamics and illuminate neural network mechanisms of pathophysiology in neurological

disorders. As an example, we briefly discuss how a whole-brain model of metastable dynamics

could be useful in elucidating the pathophysiology of Parkinson's disease (PD). The structural

degeneration of cortical and subcortical connectivity is a key factor underlying the motor and

cognitive deficits in PD [37–39]. However, the mechanisms by which the structural degeneration

alters the brain-wide dynamics in PD is not well understood. A whole-brain model of metastable

dynamics constrained by the PD-affected anatomical connectivity can potentially delineate such

mechanisms. For instance, amplified synchronization in beta frequency (8Hz – 30Hz), which has

been reported in the cortical and basal ganglia circuits of PD subjects, is hypothesized to contribute

to PD symptoms [37,40,41]. In particular, longer episodes of beta oscillations in the subthalamic

nucleus (STN) [42] and its higher synchronization with the cortical supplementary motor area

(SMA) were associated with the freezing of gait in PD [43]. It may be hypothesized that the

freezing of gait in PD is contributed by a reduction in synergistic transitions between different

attractor states realized in these areas. The increased beta synchronization between STN and SMA

observed in PD, and the hypothesized reduction in attractor transitions can be validated in whole-

brain models by examining their stability (Equation – 6) and collective response characteristics

(Equation – 9) respectively. Furthermore, by selectively targeting frequency bands for

manipulation, one can investigate causal relationships between specific frequencies of oscillations

and the dynamics of transitions between attractor states realized across multiple timescales. Such

whole-brain models could also enable identification of network mechanisms to restore optimal

brain dynamics in PD and other neurological disorders.

5. Conclusion

The numerical simulations in this study show that the inhibitory interneuron oscillations that are

modulated by slower excitatory neurons, under certain conditions, realize metastable attractors.

Furthermore, our analyses revealed frequency dependences of the emergent metastability and

quantitatively associated the features of spiking neural network dynamics at two temporal scales.

This novel cross-timescale association using a compact mathematical description captures a

biological complexity of neural circuits at the level of individual neurons and allows, in principle,

a scalable integration with empirical observations of neural network features to develop a coherent

understanding of nervous system functions.

Acknowledgments:

The authors thank Alev Erisir, Teague Henry, Giselle Petzinger, Michael Jakowec, and Dawn Schiehser for their insights and helpful discussions. The authors appreciate the complementary perspectives of Jeffrey Kopsick, Giorgio Ascoli, and Alexander Komendantov and are thankful for their many valuable discussions.

Author Contributions:

SV, EB and JDVH conceptualized the study. SV and AS contributed to the analyses. EB and HS provided critical inputs at various stages of this study. SV and EB contributed to manuscript writing. EB and JDVH contributed to supervision and JDVH contributed to funding acquisition.

Declaration of Interests:

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability:

The data that support the findings were generated from numerical simulations by the authors, and all simulation and analysis scripts are publicly available at *https://github.com/sivaven/Transient-Synchronization.*

Funding:

This work was supported in parts by the Department of Defense grant W81XWH 18 1 0665 PD170037 and an internal funding from the University of Virginia.

References:

[1] H. Markram, E. Muller, S. Ramaswamy, M.W. Reimann, M. Abdellah, C.A. Sanchez, A. Ailamaki, L. Alonso-Nanclares, N. Antille, S. Arsever, G.A.A. Kahou, T.K. Berger, A. Bilgili, N. Buncic, A. Chalimourda, G. Chindemi, J.-D. Courcol, F. Delalondre, V. Delattre, S. Druckmann, R. Dumusc, J. Dynes, S. Eilemann, E. Gal, M.E. Gevaert, J.-P. Ghobril, A. Gidon, J.W. Graham, A. Gupta, V. Haenel, E. Hay, T. Heinis, J.B. Hernando, M. Hines, L. Kanari, D. Keller, J. Kenyon, G. Khazen, Y. Kim, J.G. King, Z. Kisvarday, P. Kumbhar, S. Lasserre, J.-V. Le Bé, B.R.C. Magalhães, A. Merchán-Pérez, J. Meystre, B.R. Morrice, J. Muller, A. Muñoz-Céspedes, S. Muralidhar, K. Muthurasa, D. Nachbaur, T.H. Newton, M. Nolte, A. Ovcharenko, J. Palacios, L. Pastor, R. Perin, R. Ranjan, I. Riachi, J.-R. Rodríguez, J.L. Riquelme, C. Rössert, K. Sfyrakis, Y. Shi, J.C. Shillcock, G. Silberberg, R. Silva, F. Tauheed, M. Telefont, M. Toledo-Rodriguez, T. Tränkler, W. Van Geit, J.V. Díaz, R. Walker, Y. Wang, S.M. Zaninetta, J. DeFelipe, S.L. Hill, I. Segev, F. Schürmann, Reconstruction and Simulation of Neocortical Microcircuitry, Cell. 163 (2015) 456–492.

[2] J.D. Kopsick, C. Tecuatl, K. Moradi, S.M. Attili, H.J. Kashyap, J. Xing, K. Chen, J.L. Krichmar, G.A. Ascoli, Robust Resting-State Dynamics in a Large-Scale Spiking Neural Network Model of Area CA3 in the Mouse Hippocampus, Cognit. Comput. (2022). https://doi.org/10.1007/s12559-021-09954-2.

[3] S. Venkadesh, J.D. Van Horn, Integrative Models of Brain Structure and Dynamics: Concepts, Challenges, and Methods, Front. Neurosci. 15 (2021) 752332.

[4] M.W. Jones, M.A. Wilson, Phase precession of medial prefrontal cortical activity relative to the hippocampal theta rhythm, Hippocampus. 15 (2005) 867–873.

[5] J. Lisman, The theta/gamma discrete phase code occuring during the hippocampal phase precession may be a more general brain coding scheme, Hippocampus. 15 (2005) 913–922.

[6] W.E. Skaggs, B.L. McNaughton, Replay of neuronal firing sequences in rat hippocampus during sleep following spatial experience, Science. 271 (1996) 1870–1873.

[7] G.R. Poe, D.A. Nitz, B.L. McNaughton, C.A. Barnes, Experience-dependent phase-reversal of hippocampal neuron firing during REM sleep, Brain Res. 855 (2000) 176–180.

[8] V. Ego-Stengel, M.A. Wilson, Spatial selectivity and theta phase precession in CA1 interneurons, Hippocampus. 17 (2007) 161–174.

[9] M. Vinck, B. Lima, T. Womelsdorf, R. Oostenveld, W. Singer, S. Neuenschwander, P. Fries, Gamma-phase shifting in awake monkey visual cortex, J. Neurosci. 30 (2010) 1250–1257.

[10] S. Venkadesh, E. Barreto, G.A. Ascoli, Itinerant complexity in networks of intrinsically bursting neurons, Chaos. 30 (2020) 061106.

[11] E. Tognoli, J.A.S. Kelso, The metastable brain, Neuron. 81 (2014) 35–48.

[12] W.J. Freeman, Consciousness, intentionality and causality, Journal of Consciousness Studies. (1999). https://www.ingentaconnect.com/content/imp/jcs/1999/00000006/F0020011/995.

[13] I. Tsuda, Hypotheses on the functional roles of chaotic transitory dynamics, Chaos. 19 (2009) 015113.

[14] J. Cabral, F. Castaldo, J. Vohryzek, V. Litvak, C. Bick, R. Lambiotte, K. Friston, M.L. Kringelbach, G. Deco, Metastable oscillatory modes emerge from synchronization in the brain spacetime connectome, Communications Physics. 5 (2022) 1–13.

[15] E.M. Izhikevich, Simple model of spiking neurons, IEEE Trans. Neural Netw. 14 (2003) 1569–1572.

[16] M. Stimberg, R. Brette, D.F. Goodman, Brian 2, an intuitive and efficient neural simulator, Elife. 8 (2019). https://doi.org/10.7554/eLife.47314.

[17] H. Daido, Onset of cooperative entrainment in limit-cycle oscillators with uniform all-to-all interactions: bifurcation of the order function, Physica D: Nonlinear Phenomena. 91 (1996) 24–66. https://doi.org/10.1016/0167-2789(95)00260-x.

[18] D. Golomb, D. Hansel, The number of synaptic inputs and the synchrony of large, sparse neuronal networks, Neural Comput. 12 (2000) 1095–1139.

[19] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, É. Duchesnay, Scikit-learn: Machine Learning in Python, J. Mach. Learn. Res. 12 (2011) 2825–2830.

[20] W.M. Rand, Objective Criteria for the Evaluation of Clustering Methods, J. Am. Stat. Assoc. 66 (1971) 846–850.

[21] L. Hubert, P. Arabie, Comparing partitions, J. Classification. 2 (1985) 193–218.

[22] D. Steinley, Properties of the Hubert-Arable Adjusted Rand Index, Psychol. Methods. 9 (2004) 386–396.

[23] J. O'Keefe, M.L. Recce, Phase relationship between hippocampal place units and the EEG theta rhythm,
Hippocampus. 3 (1993) 317–330.
[24] G. Dragoi, G. Buzsáki, Temporal encoding of place sequences by hippocampal cell assemblies, Neuron. 50
(2006) 145–157.
[25] J.C. Sommerer, E. Ott, A physical system with qualitatively uncertain dynamics, Nature. 365 (1993) 138–140.
[26] E. Ott, J.C. Sommerer, J.C. Alexander, I. Kan I., J.A. Yorke, Scaling behavior of chaotic systems with riddled
basins, Phys. Rev. Lett. 71 (1993) 4134–4137.
[27] K.J. Friston, Transients, metastability, and neuronal dynamics, Neuroimage. 5 (1997) 164–171.
[28] O. Sporns, G. Tononi, G.M. Edelman, Theoretical neuroanatomy and the connectivity of the cerebral cortex,
Behav. Brain Res. 135 (2002) 69–74.
[29] G. Tononi, O. Sporns, Measuring information integration, BMC Neurosci. 4 (2003) 31.
[30] O. Sporns, G. Tononi, R. Kötter, The human connectome: A structural description of the human brain, PLoS
Comput. Biol. 1 (2005) e42.
[31] A.O. Komendantov, S. Venkadesh, C.L. Rees, D.W. Wheeler, D.J. Hamilton, G.A. Ascoli, Quantitative firing
pattern phenotyping of hippocampal neuron types, Sci. Rep. 9 (2019) 17915.
[32] W.J. Freeman, A pseudo-equilibrium thermodynamic model of information processing in nonlinear brain
dynamics, Neural Netw. 21 (2008) 257–265.
[33] S. Mori, P.C.M. van Zijl, Fiber tracking: principles and strategies - a technical review, NMR Biomed. 15
(2002) 468–480.
[34] G. Deco, V.K. Jirsa, Ongoing cortical activity at rest: criticality, multistability, and ghost attractors, J. Neurosci.
32 (2012) 3366–3375.
[35] G. Deco, A. Ponce-Alvarez, D. Mantini, G.L. Romani, P. Hagmann, M. Corbetta, Resting-state functional
connectivity emerges from structurally and dynamically shaped slow linear fluctuations, J. Neurosci. 33 (2013)
11239–11252.
[36] N.N. Foster, J. Barry, L. Korobkova, L. Garcia, L. Gao, M. Becerra, Y. Sherafat, B. Peng, X. Li, J.-H. Choi, L.
Gou, B. Zingg, S. Azam, D. Lo, N. Khanjani, B. Zhang, J. Stanis, I. Bowman, K. Cotter, C. Cao, S. Yamashita,
A. Tugangui, A. Li, T. Jiang, X. Jia, Z. Feng, S. Aquino, H.-S. Mun, M. Zhu, A. Santarelli, N.L. Benavidez, M.
Song, G. Dan, M. Fayzullina, S. Ustrell, T. Boesen, D.L. Johnson, H. Xu, M.S. Bienkowski, X.W. Yang, H.
Gong, M.S. Levine, I. Wickersham, Q. Luo, J.D. Hahn, B.K. Lim, L.I. Zhang, C. Cepeda, H. Hintiryan, H.-W.
Dong, The mouse cortico-basal ganglia-thalamic network, Nature. 598 (2021) 188–194.
[37] M.M. McGregor, A.B. Nelson, Circuit Mechanisms of Parkinson's Disease, Neuron. 101 (2019) 1042–1056.
[38] C. Atkinson-Clement, S. Pinto, A. Eusebio, O. Coulon, Diffusion tensor imaging in Parkinson's disease:
Review and meta-analysis, Neuroimage Clin. 16 (2017) 98–110.
[39] C.J. Cochrane, K.P. Ebmeier, Diffusion tensor imaging in parkinsonian syndromes: a systematic review and
meta-analysis, Neurology. 80 (2013) 857–864.
[40] J.-S. Brittain, A. Sharott, P. Brown, The highs and lows of beta activity in cortico-basal ganglia loops, Eur. J.
Neurosci. 39 (2014) 1951–1959.
[41] C. Hammond, H. Bergman, P. Brown, Pathological synchronization in Parkinson's disease: networks, models
and treatments, Trends Neurosci. 30 (2007) 357–364.
[42] M. Deffains, L. Iskhakova, S. Katabi, Z. Israel, H. Bergman, Longer β oscillatory episodes reliably identify
pathological subthalamic activity in Parkinsonism, Mov. Disord. 33 (2018) 1609–1618.
[43] J.B. Toledo, J. López-Azcárate, D. Garcia-Garcia, J. Guridi, M. Valencia, J. Artieda, J. Obeso, M. Alegre, M.
Rodriguez-Oroz, High beta activity in the subthalamic nucleus and freezing of gait in Parkinson's disease,
Neurobiol. Dis. 64 (2014) 60–65.
449