Uncovering the contributions of charge regulation to the stability of single alpha helices

Martin J. Fossat[§], Ammon E. Posey[§], Rohit V. Pappu

Department of Biomedical Engineering and the Center for Biomolecular Condensates (CBC), James McKelvey School of Engineering, Washington University in St. Louis, St. Louis, MO 63130

ABSTRACT

The single alpha helix (SAH) is a recurring motif in biology. The consensus sequence has a di-block architecture that includes repeats of four consecutive glutamate residues followed by four consecutive lysine residues. These E_4K_4 repeats are thought to contribute to helical conformations through i to $i+4$ salt bridges. Interestingly, the overall helicity of sequences with consensus E_4K_4 repeats tends to be remarkably insensitive to a wide range of pH values. This pH insensitivity cannot be explained by regular networks of salt bridges alone. Here, we use the recently introduced q-canonical ensemble, which allows us to decouple measurements of charge state and conformation, to understand how the insensitivity of SAH helicity to pH comes about. We couple the outputs from charge and conformational measurements with atomistic simulations to derive residue-specific quantifications of preferences for being in an alpha helix and for the ionizable residues to be charged vs. uncharged. We find a clear preference for accommodating uncharged Glu residues within internal positions of SAH-forming sequences. Further, while the N-terminal capping positions prefer to be negatively charged, the C-terminal Lys residues prefer being uncharged. Importantly, away from pH 7.5, there is a combinatorial increase in the number of charge states that are accessible to SAH-forming sequences, especially as the numbers of E_4K_4 repeats increase. The accessible charge states are compatible with forming conformations of high helical content pointing to conformational buffering whereby charge state heterogeneity buffers against large-scale conformational changes. This makes the overall helicity insensitive to large changes in pH.

INTRODUCTION

The alpha-helix, first discovered by Pauling, Corey, and Branson [1] is a recurring secondary structural motif in proteins [2, 3]. Considerable effort, spanning multiple decades, has gone into uncovering the physico-chemical principles of helix-coil transitions [4-7], the sequence determinants of helical stability [8-11], rules for helix capping [12, 13], the impact of solution conditions on helical stability [14-17], cooperative bundling of helices [18], and the ability of many systems to form helical structures through coupled folding and binding reactions [14, 19, 20]. Ionizable residues feature prominently in many helix-forming sequences [19, 21-23]. Among the roles attributed to ionizable residues is their ability to improve solubility [11], their contributions to helical stability through salt bridges [21, 23], their preferential accumulation at specific capping positions [13], and their ability to contribute to helical stability through pH-dependent uptake or release of protons [24-28], which we refer to as *charge regulation* [29].

Alpha helices within globular proteins are typically 10-15 residues long [30]. However, many exceptions exist and the most notable are sequences that form the Single Alpha Helix (SAH) motif. The lengths of such sequences can range from 25 to 200 residues [31, 32] (**Figure S1**). Many, although not all SAH formers feature a consensus di-block architecture comprising E_4K_4 repeats [31, 33-35]. Each repeat features four glutamic acid residues followed by four lysine residues. **Figure 1** summarizes the diverse set of functions associated with distinct variants of SAHs, all featuring

E_4K_4 repeats or variants thereof. In many systems, SAHs are thought to contribute as semiflexible rod-like elements that tether distinct functional domains. Mechanotransduction, mechanical sensing, and force generation are among the numerous functions associated with SAH forming sequences [36-38]. Further, since their discovery, SAH sequences of different lengths have been used as molecular rulers for measurements of intramolecular distances in cellular systems [39].

The stability of SAH sequences and their ability to form alpha-helical rod-like conformations have been attributed to i to $i+4$ salt bridges between negatively charged Glu residues at position i and Lys residues at position $i+4$ [37, 40]. Two observations suggest that there are likely to be complex contributions to determinants of SAH stability. First, in measurements reported to date, the helicities of SAHs show remarkable insensitivity to changes in pH [33, 40]. Given that Glu and Lys are not particularly strong acids or bases [41], it follows that charge regulation effects are likely to contribute to the stability of SAHs. Here, charge regulation refers to the alterations of charge states of ionizable residues through spontaneous binding or release of protons. This hypothesis is bolstered by results from the early work of Marqusee and Baldwin [21] who showed that the positioning of ion pairs has a modest effect on alpha helicity in alanine-based systems. They also showed that the alpha helicity in model alanine-based peptides that feature complementary ion pairs between Arg / Lys and Glu residues can be higher at extremes of pH when compared to pH 7.5. Under these conditions, the peptides have a net charge as opposed to being electroneutral. Additionally, Doig and coworkers have shown that the positioning of phosphoserine in alanine-based systems has a profound effect on helical stability [42, 43]. These observations were linked to the position-dependence of the charge state of the phosphoserine and lysine residues within the sequence. Together, these studies suggest that charge regulation is likely to be an important contributor to the stability of SAH forming sequences. Further support for the putative role of charge regulation comes from a second observation pertaining to the imperfectness of E_4K_4 repeats. Although E_4K_4 is the consensus motif, one often sees interruptions of this pattern that go beyond the substitution of Lys with Arg. Indeed, Glu and Lys are often substituted with uncharged residues such as Ala, Gln, or Leu (**Figure 1**) [31]. This can give rise to sequences with a net charge, as naively evaluated by assuming the charge states to be fixed by the pK_a values of model compounds. Additionally, while there are occasional substitutions of Glu with Asp, one never observes blocks of Asp residues [44]. The likelihood of observing four consecutive Arg residues is also lower than the likelihood of observing four consecutive Lys residues.

Why might charge regulation have a role to play in helix formation / stabilization? First, Pace and Scholtz measured the intrinsic helical propensities of naturally occurring amino acids and uncharged versions of specific ionizable residues. Averaged over three different sequence contexts, the consensus helical propensities that were derived from their measurements have the following hierarchy: Glu^0 > Ala > Leu $\approx$ Arg^+ > Lys^+ > Gln > Glu^- > Asp^0 > Asn > Asp^-. Here, Glu^0 and Asp^0 are uncharged versions of Glu and Asp, respectively. Likewise, Arg^+, Lys^+, Glu^-, and Asp^- are charged versions of Arg, Lys, Glu, and Asp, respectively. Clearly, the intrinsic helical propensities are among the highest for Glu^0. Second, the free energies of hydration, which are relevant for desolvation penalties associated with helix formation, are different for basic vs. acidic groups, and considerably more favorable for Asp^0 and Glu^0 compared to Asp^- and Glu^- [45, 46]. This points to the possibility of lowering the desolvation penalty through site-specific or context-dependent proton uptake or release. Third, a strong case has been made for the importance of reducing the linear charge density along to the helical axis [22] – a feature that is achievable through charge regulation. And fourth, local electrostatic repulsions within and between like-charged blocks are likely to disrupt or destabilize optimal i to $i+4$ salt bridges.

Figure 1: Examples of naturally occurring SAH sequences from three distinct protein families. (A) The protein name and organism. protein function, SAH amino acid sequence and structural cartoons are given for five representative SAH-containing proteins. SAH regions are colored green in the structural cartoons (column 4). For clarity, in column 3, the acidic residues are shown in red, the basic residues in blue, and the constitutively uncharged residues are shown in black. (B) An example consensus domain architecture is depicted for the "unconventional myosin" family proteins that commonly feature SAHs (indicated in green). The domains that follow the SAH are variable, as indicated. The domain architecture schematic was adapted from the work of Fili and Toseland[47] and Li et al.[48].

Taken together, the observations summarized above suggest that modulations of charge states of ionizable residues are likely to impact SAH stability through increased intrinsic helical propensities, lowered charge density along the helical axis, and decreased desolvation penalties. Further, selective, position-specific protonation of acids vs. deprotonation of bases could enhance

the specificity of helix-stabilizing i to $i+4$ salt bridges by avoiding the possibility of alternative salt bridges. To test these hypotheses, we performed systematic investigations of the pH-dependent behaviors of three systems with E_4K_4 repeats.

Our investigations were aided by the q-canonical ensemble [29], which we recently introduced. This allows us to parse measurements based on potentiometric titrations to characterize the ensemble of charge states that are prevalent in solution for a specific pH [29]. The q-canonical ensemble was applied to characterize the ensemble of charge states that peptides with E_4K_4 repeats can access as a function of pH [29]. Here, we adapt this formalism by combining the measurements of charge states with measurements of conformations to go alongside atomistic simulations based on the ABSINTH implicit solvation model and forcefield paradigm [49]. The combined approach helps us uncover the role of charge regulation as a determinant of the insensitivity of SAH stability to changes in pH. Interestingly, away from pH window of 7.0 – 7.4 we observed significant increases in charge state heterogeneity that do not compromise helical stability. Our results suggest that the effects of proton binding and unbinding are likely to add to the contributions of backbone hydrogen bonding, backbone solvation / desolvation, and inter-block salt bridges as determinants of SAH stability.

Our studies focus on three systems of the form Ac-G-$(E_4K_4)_n$-GW-NH_2, where n = 1, 2, and 3, respectively. Here, Ac refers to N-acetyl, and the tryptophan residue was used to enable precise measurements of peptide concentration. As a shorthand, we refer to the three systems as $(E_4K_4)_1$, $(E_4K_4)_2$, and $(E_4K_4)_3$, respectively. First, we present results summarizing the repeat number and pH dependence of helicity as measured by ultraviolet circular dichroism (UV-CD). Then, we summarize the main tenets of the q-canonical ensemble, specifically the parsing of the system into microstates and mesostates. This is followed by a summary of the workflow that combines potentiometric titrations, reported previously, with the UV-CD measurements to extract intrinsic helicities of different charge mesostates. Using these mesostate-specific helicities as restraints, we performed atomistic simulations based on the ABSINTH implicit solvation model to extract a full description of the pH-dependence of conformations and charge states that are consistent with the experimental data. These results were used to extract insights regarding the pH-dependent preferences for charge states of individual ionizable residues. For helical conformations, we uncovered clear, repeat number dependent downshifts in the apparent pK_a values of internal Glu residues and significant, position-specific apparent pK_a values for Lys residues. Further, for $(E_4K_4)_2$ and $(E_4K_4)_3$, we observed increased charge state heterogeneity away from pH 7.5, , that prevails despite negligible changes to the overall helicity. These results suggest that at a given pH, the overall ensemble, especially for longer sequences, is best described as being predominantly helical while accommodating significant charge state heterogeneity.

RESULTS

$(E_4K_4)_n$ systems show n- dependent and pH-independent preferences for overall helicity: We performed UV-CD measurements for each of the three $(E_4K_4)_n$ systems, where n = 1, 2, and 3 (**Figure S2**). The measurements were performed in a background solution of 10 mM KCl and 5 mM HCl at a temperature of 20˚C and titrated across pH values ranging from 2 – 12 using KOH. The overall helicity, measured using UV-CD, and plotted as the fractional helicity, is summarized in **Figure 2**. We observed the following trends in the data: For a given pH, the fractional helicity increases with the number of E_4K_4 repeats. For each construct, there is a pH window where the fractional helicity changes minimally. The size of this window increases with the number of repeats. In the pH window where the helicity varies minimally, the fractional helicity

values are ≈ 0.17 for $(E_4K_4)_1$, ≈ 0.5 for $(E_4K_4)_2$, and ≈ 0.7 for $(E_4K_4)_3$. These helicities are concordant with values reported in the literature [34]. The key result is the insensitivity of overall helicity to significant changes in pH.

Figure 2: Fractional helicity derived from the UV-CD data. The full spectra are shown in **Figure S2**. Data points and error bars represent, respectively, the mean fractional helicity from deconvolution and the standard deviation of deconvolution using three different basis sets, as described in the Methods section

Potentiometric titrations enable the measurements of charge states of $(E_4K_4)_n$ peptides: In a recent report, we published results from potentiometric titrations of the three $(E_4K_4)_n$ peptides studied in this work [29]. The salient results from these titrations are summarized in **Figure S3**. Potentiometry yields estimates of the net charge as a function of pH. We used the formalism of the q-canonical ensemble, introduced previously [29], to extract information regarding the distinct charge states that contribute to the titrations as a function of pH. The overall approach is summarized below using the $(E_4K_4)_1$ peptide as an illustrative system.

There are eight ionizable residues for the $(E_4K_4)_1$ peptide. Each Glu residue can be deprotonated or protonated. These are denoted as E and e, respectively. Likewise, each Lys residue can be protonated (K) or deprotonated (k). In all, for $(E_4K_4)_1$ there is a theoretical maximum of 2^8 or 256 charge microstates, where each microstate is denoted as a specific sequence *viz.*, eEEEKKKK, EeEEKKKK, etc. Of these, several microstates are forbidden since Lys and Glu residues are unlikely to be simultaneously deprotonated and protonated, respectively. Accordingly, for $(E_4K_4)_1$, the number of relevant charge microstates reduces to 32 as was shown previously [29]. Next, the charge microstates are grouped into mesostates. Each mesostate encompasses all microstates that have the same net charge. Mesostates are given an X_Y designation, where X refers to the net charge associated with each of the microstates within a mesostate, and Y refers to the number of thermodynamically accessible microstates within a mesostate (**Figure S4**).

The pH-dependence of the net charge is used to extract the probability that a specific mesostate will be populated at a given pH. Previous work showed that for $(E_4K_4)_1$, the mesostate namely 0_1 dominates in the pH range between 7 and 9 [29]. However, the data also indicate a downshift vs. an upshift in the pK_a values for the acids and the bases, respectively. For the longer peptides $(E_4K_4)_2$ and $(E_4K_4)_3$, there are at least two mesostates of net charge ± 1 that contribute to the measured charge profiles at every pH, even if the contributions are small (**Figure S5**). The pH dependent populations of mesostates can be used to quantify the numbers of mesostates that contribute to 95% vs. 99% of the overall population as a function of pH, as shown in **Figure S6**. The overall picture that emerges is of an ensemble of distinct charge microstates. The number of

thermodynamically relevant microstates / mesostates is set by the pH. The 0_1 mesostate becomes less important outside the pH window of 7.0 – 7.5. Accordingly, the question is if there are mesostates with a net charge that also have high intrinsic helicities. The existence of such mesostates would be one way to explain the robust pH-independence of helicity of (E_4K_4) repeats.

Results from potentiometric titrations and pH-dependent UV-CD measurements can be combined to quantify the intrinsic helicities of mesostates: The pH-dependent mesostate probabilities p_q(pH), extracted from potentiometric titrations, can be used in conjunction with the pH-dependent estimates of overall helicity, f_α(pH) plotted in **Figure 2**, to extract intrinsic helicities, $f_{\alpha,q}$, for each of the mesostates characterized by a net charge q. For this, we use the formalism summarized in Equation (1) in conjunction with a Monte Carlo fitting procedure (see details in the methods section).

$$f_\alpha(pH) = \sum_q^{N_{meso}} p_q(pH)\, f_{\alpha,q}; (1)$$

Here, f_α(pH) and p_q(pH) are inferred using data from UV-CD measurements and potentiometric titrations, respectively. These are then used in a Monte Carlo fitting procedure to extract $f_{\alpha,q}$. The workflow is summarized in panels (A) and (B) of **Figure 3**.

The values of $f_{\alpha,q}$, extracted from the fitting procedure, are summarized in **Figure 4A-4C**. Further, as shown using the solid curves in **Figure 4D**, the values obtained for $f_{\alpha,q}$ can be used to obtain fits to the measured fractional helicities as a function of pH. The key takeaway is that there are several mesostates, especially for the $(E_4K_4)_2$ and $(E_4K_4)_3$ systems whose intrinsic helicities are equivalent to or greater than that of the 0_1 mesostate. All these mesostates have a net charge. For example, the intrinsic helicities of mesostates with a net charge of ±9, ±8, ±7, ±3, ±1, and -11 are equivalent to or higher than the intrinsic helicity of the 0_1 mesostate for the $(E_4K_4)_3$ system.

The potentiometric titrations help identify the mesostates that are relevant for a coarse-grained description of the overall q-canonical ensemble. By combining the information from potentiometric titrations with pH-dependent UV-CD measurements, we quantify the intrinsic helicities of mesostates. To obtain information regarding the microstates, specifically the types of charge microstates and the conformations of microstates that contribute to the overall ensemble, we turn to simulations as summarized in **Figure 5**.

Figure 3: Workflow for deploying the full *q*-canonical ensemble using potentiometric titrations, UV-CD spectroscopy, and atomistic simulations. (A) Summary of the fitting methods used to extract mesostate probabilities as a function of pH from potentiometric titrations. The approach is that of Fossat et al. [29]. (B) Summary of the procedure to determine the intrinsic fractional helicities of mesostates. Here, we use the mesostate probabilities derived from the workflow shown in panel (A). This is combined with pH-dependent CD measurements.

Figure 4: Bar plots summarizing the intrinsic helicities, shown as fractions that lie between 0 and 1. (A) Data are shown for the relevant mesostates for each of the $(E_4K_4)_n$ peptides. In each sub-panel of the three sub-panels that make up (A), the dashed line quantifies the construct specific intrinsic helicity of the 0_1 mesostate. The intrinsic helicities were extracted using a combination of mesostate weights from potentiometric measurements and from the CD data. (B) Mean fractional helicity determined from deconvolution of the CD data. The fitting envelope corresponds to the standard deviation of the best fits from each of the 100 fits as a function of pH. These data were jointly analyzed as summarized in the main text and panel (B) of Figure 3. The analysis is based on the use of Equation 1.

Atomistic simulations identify the microstates that contribute to the experimentally observed helicity profiles as a function of pH: The simulations use atomic-level descriptions of polypeptides and solution ions based on an all-atom forcefield. The aqueous solvent is modeled implicitly using the ABSINTH model and forcefield paradigm [49]. For each system, a series of simulations were performed for each of the mesostates using the *q*-canonical Monte Carlo (*q*MC) sampling methodology [50]. Further, we used the combination of inferred intrinsic helicities of mesostates and their populations as a function of pH as inputs for reweighting the simulations for each of the mesostates. The reweighting procedure is summarized in **Figure 5**.

Simulations Reweighting

Figure 5: Summary of the procedure used to mix information from three distinct sets of simulations and reweight the ensembles, so they agree with both potentiometric titrations and CD data. Here, f_i^D, f_i^U, and f_i^H are the contributions to fractional helicities of mesostates from the disordered ensemble D, the unbiased or unmodified ensemble U, and the helical ensemble H. Likewise, W_i^D, W_i^U and W_i^H are the weights of the mixing function, which are refined through iteration. The converged values of these weights reflect the contributions from the three ensembles that yield congruence with the experimental data for fractional helicities of mesostates.

For reweighting, we mix conformations from three distinct ensembles *viz.*, conformations obtained from a highly disordered ensemble obtained by setting the simulation temperature to 350 K, a second set corresponding to unbiased simulations performed at a simulation temperature of 240 K, and the third set corresponding to the most helical ensemble. The helical ensemble is obtained by imposing harmonic constraints on the backbone, thus maintaining an overall helical

conformation. The temperature of 240 K for the unbiased simulations was determined using simulations initiated within the 0_1 mesostate and identifying the temperature at which the helicity of this mesostate equals the helicity extracted from the combination of CD and potentiometry. It is worth noting that simulation temperatures and the real temperatures in experiments do not have a one-to-one congruence for classical forcefields and implicit solvation models.

A suitable mixture of the three sets of simulated ensembles is obtained by reweighting on a per mesostate basis as summarized in **Figure 5**. For each mesostate, we assess the overall helicity in each of the three sets. We then compare the mesostate helicities of the unmodified set to the mesostate helicity obtained from experiments. If the helicity is too high for that mesostate compared to the experimentally derived value, we introduce the first set until the overall weighted sum of the two sets is within 0.001 of experiments. Conversely, if the helicity is too low for a mesostate compared to the experimentally derived value, we introduce conformations from the disordered ensemble to converge on the overall reweighted ensemble. The proportion of each ensemble used per mesostate is shown in **Figure S6**.

Using the mixing function shown in **Figure 5**, we derived weights associated with each of the three ensembles for each mesostate. We then set the mesostate free energies to those obtained from potentiometric data, while conserving the properly mixed microstate weights within mesostates. This means that the average free energy of each mesostate is concordant with potentiometrically derived values, and the free energy differences among microstates within mesostates are those derived from simulations, after proper mixing. This allows us to extract microstate weights that are also consistent with experiments thereby allowing us to generate structural insights on a per residue basis. Our use of the three sets of simulated ensembles, including the unbiased simulations allows us to obtain a per residue profile that reflects the physics of the forcefield paradigm that is used in our simulations.

The salient results from the reweighting are summarized in **Figure 6**. Here, each row shows results for a specific pH, whereas each column shows results for a specific construct. The quantities being plotted are the contributions of distinct mesostates to the position-specific helicity values. The latter were derived from the properly reweighted, pH-dependent ensembles of microstates. Along each column, we note that the heterogeneity of charge states is highest at pH 9.5 and 5.5 and lowest at pH 7.5. Further, going across each row, the contributions from charge state heterogeneity to the overall q-canonical ensemble become more pronounced with increasing numbers of E_4K_4 repeats. The implication is that as the number of repeats increases, there are more charge states that are compatible with similar degrees of helicity. This is especially true at pH values that go above or below pH 7.5.

Figure 6: Mesostate-specific contributions to helicity per residue at three different pH values. Each column contains data for a specific construct, and each row contains data for a specific pH. The stacked bars quantify the contributions from specific mesostates to the overall helicity per residue. The insets show the relevant color coding of the mesostates within a panel. Here, Z along the abscissa refers to the N- and C-terminal caps. At pH 5.5, the relevant mesostates for $(E_4K_4)_3$ have a net charge of +1, +2, and +3. The mesostates with net charge of +1, +2, and +3 have 12, 66, and 220 accessible microstates, respectively and the likelihoods of accessing these microstates are determined by the intra-mesostate microstate probabilities. In terms of their contributions to the overall helicity, the dominant mesostates follow the hierarchy, $+1_{12} > +3_{220} \approx 0_1 > +2_{66}$. At pH 7.5, there are minor contributions from the mesostates with charge of ± 1 for $(E_4K_4)_3$, and the hierarchy in terms of contributions to the overall to the overall helicity is $0_1 >> +1_{12} > -1_{12}$. Finally, for $(E_4K_4)_3$ at pH 9.5, the contributions to the overall helicity from mesostates follow the hierarchy, $-1_{12} > -2_{66} \approx 0_1 > -3_{220}$. These results illustrate the contributions of charge state heterogeneity to the maintenance of overall helicity.

How rod-like are the conformational ensembles at a given pH? Next, we quantified the distributions of the lengths of helical segments within the reweighted ensembles. The results are shown in **Figure 7** for $(E_4K_4)_3$ and in **Figure S7** for $(E_4K_4)_2$. The overall helicity is low for $(E_4K_4)_1$. Despite a high fractional helicity of ≈ 0.7 that persists across the pH-dependent ensembles, the perfectly helical rod-like conformation is not the dominant species for $(E_4K_4)_3$. Instead, the overall ensemble is a mixture of partial and fully helical conformations. The four main conformational classes are as follows: In class 1, the N- and C-terminal repeats are helical, whereas the internal repeats are disordered. In class 2, roughly 50-60% of the sequence encompassing the N-terminal

portion adopts helical conformations whereas the remaining C-terminal portion is disordered. The converse is true for class 3. Finally, in class 4, conformations correspond to the fully helical, rod-like state (**Figure 6**). The distributions of helical segments of specific lengths share commonalities across the different pH values. There are three distinct peaks corresponding to class 1, class 3, and class 4 conformations, respectively. Class 2 conformations correspond to a trough between two peaks associated with class 1 and class 3 conformations. The trough at a helical segment length of 20 residues points to a clear preference against disrupting the transition between class 3 and class 4 conformations.

Figure 7: Distributions of the lengths of helical segments extracted from the reweighted ensembles of (E$_4$K$_4$)$_3$ at pH 5.5, 7.5, and 9.5. The top row shows representative conformations as annotations of specific features of the graph. Conformations corresponding to classes 1 – 4 are depicted going from left to right.

Site specific ionization preferences are context dependent: In **Figure 8** we plot the probability of finding each residue in a charged state. For simplicity of presentation, the data are shown for (E$_4$K$_2$)$_2$. For reference, ionization curves are also shown if we were to assume fixed charge states based on the pK$_a$ values of model compounds. We define the site-specific apparent pK$_a$ as the pH at which the likelihood of being charged is 0.5. The top rows of columns A and B of **Figure 8** show insets that zoom into the ionization curves with the dashed lines highlighting the likelihood of 0.5. Overall, the apparent pK$_a$ values inferred from the ionization curves suggest that the Glu residues have systematically downshifted pK$_a$ values when compared to the model compound value of 4.34 [51]. The maximal extent of the downshift is approximately 0.8 of a pH unit. Conversely, the Lys residues show greater heterogeneity in terms of their apparent pK$_a$ values. There can be a down or upshift of the apparent pK$_a$ vis-à-vis the model compound value of 10.34 [51], and the dispersion of apparent pK$_a$ values is greater than one pH unit.

Figure 8: Site specific ionization curves for each of the positions in (E₄K₄)₂. The curves are shown by dividing the pH range into two parts. This helps visualize the titration of Glu (panel A) and the titration of Lys residues (panel B). The top row shows zoomed in versions that focus on specific pH ranges to help with identifying the pH values at which the likelihood of being charged is 0.5 for the different residues (see black dashed lines). The red and blue dashed curves correspond to the ideal case of ionizable residues titrating at pKₐ values corresponding to those of model compounds (10.34 for Lys and 4.34 for Glu [51]), which in the legend is referred to as "normal".

To quantify the preferential, site-specific protonation of ionizable residues, we introduce a parameter $\Omega_{q,r}$ for each site r along the sequence. Within a given mesostate of net charge q, the $\Omega_{q,r}$ parameter is defined as:

$$\Omega_{q,r} = \sum_{j}^{g_q} \left(p_j - \frac{1}{g_q} \right) s_{r,j}; \ (2)$$

Here, $\Omega_{q,r}$ is the preferential protonation of residue r in a mesostate with net charge q; p_j is the probability associated with microstate j; g_q is the number of microstates within the mesostate of net charge q, and $s_{r,j}$ is a switch function that quantifies the state of residue r in microstate j. The switch function has a value 1 if the residue is charged, and 0 if it is uncharged. The difference, shown in parentheses of the summand quantifies the difference between the microstate probability p_j, derived from the properly reweighted ensemble and $(g_q)^{-1}$, which is the value we would expect from the postulate of equal *a priori* probabilities, the null model which sets all microstates within a mesostate to be equiprobable. If the summand is positive, it means that the microstate in question has a higher probability of being charged when compared to the null model. Conversely, if the summand is negative, then the microstate j has a lower probability of being charged when compared to the null model.

Using the information for all $\Omega_{q,r}$ values computed across all mesostates, we obtain the pH-dependent preferential protonation values per residue. This profile is denoted as $\Omega_r(\text{pH})$. It is computed as a weighted sum over mesostates populations at a given pH and is written as:

$$\Omega_r(pH) = \sum_q \Omega_{q,r} p_q(pH); \ (3)$$

The results for $\Omega_r(pH)$ for each position r are plotted in **Figure 9** for each of the three systems at pH 7.5. Data for each system are shown along the columns. Each row shows the $\Omega_r(pH)$ profiles computed for the reweighted ensemble (top row), the helical ensemble (middle row), and the disordered ensemble (bottom row). Positive values for $\Omega_r(pH)$ imply that there is a preference to be charged, more so than would be expected based on the assumption of a fixed pK_a value for all residues of the same type. Conversely, negative values for $\Omega_r(pH)$ imply that there is a preference to be uncharged, more so than would be expected based on the assumption of a fixed pK_a value for all residues of the same type. Note that these interpretations for the positive vs. negative deviations come from the sign of the summand in Equation (2). It depends on the microstate probability as inferred from the simulations. These probabilities are referenced to the probability we would compute by assuming that all microstates within a mesostate are equiprobable.

Figure 9: Ω profiles at pH 7.5 for each of the three peptides, organized along the columns, and types of ensembles, organized along the rows. The definition of Ω is in the main text. Positive values for Ω indicate a higher propensity for that residue to be in its charged form when compared to the scenario where all residues of the same type have the same pK_a value. Negative values of Ω indicate a higher propensity for that residue to be in its uncharged form when compared to the scenario where all residues of the sane type have the same pK_a value. Along the abscissa, the Z at the N-terminal end is the label for N-acetyl and Z at the C-terminal end is for NH2. We use Z as a single letter code for both caps.

For the longest sequence, $(E_4K_4)_3$ in the reweighted or fully helical ensembles, there is a higher preference for the internal Glu residues to be uncharged, when compared to the model that assumes fixed charged states. The converse is true for the internal Lys residues. In the reweighted and fully helical ensembles, there is an increased preference for charged Glu residues at the N-terminal capping position and an increased preference for uncharged Lys residues at the C-terminal capping position. Interestingly, the Ω profiles are essentially inverted in the disordered ensemble (**Figure 9**).

Overall, the observations summarized in **Figure 9** can be traced to a series of factors that include the following: (i) The intrinsic helical propensity of Glu^0 is higher than that of Glu^- [28]. In fact, the intrinsic helical propensity of Glu^0 can even be higher than that of alanine, which is typically used as the standard for high intrinsic helical propensity [28]. This higher intrinsic helical propensity helps rationalize the increased preference for Glu^0 at internal positions. Additionally, Ermolenko et al., showed that the helical propensities of uncharged residues are independent of their positions within the middle of alpha helices [52]. This provides further support for our observation of preferential substitutions of Glu^- with Glu^0 in internal positions of sequences with (E_4K_4) repeats. (ii) Our finding of a clear positional preference for the charged Glu residues at the N-terminal end is concordant with documented preferences for Glu^- at the N-terminal caps [12, 13]. Our observation of a positional preference for uncharged Lys^0 residues at the C-terminal caps is concordant with documented statistical preferences for polar amino acids and even hydrophobic groups at C-terminal capping positions [12, 13, 53]. (iii) In the disordered ensembles, we observe a clear inversion, with one exception, of the positional preferences for being charged vs. uncharged. This reflects the dominance of local sequence context effects and contributions of conformational heterogeneity in the disordered ensembles. These effects engender a preference for maintaining Lys residues in their charged states at internal positions. Further, the increased preference of C-terminal Lys residues to be uncharged is also present in the disordered ensemble.

There is a finite probability of observing uncharged residues instead of Glu or Lys residues in naturally occurring SAH sequences: Our findings suggest that maintenance of helical conformations across a wide range of pH values comes with a significant increase in charge state heterogeneity as the pH deviates from 7.5. This is likely to be become more pronounced in longer SAH sequences given the combinatorial increase in the number of charge states. One way to reduce this increased charge state heterogeneity, if needed, would be to substitute internal Glu or Lys residues with non-ionizable residues. We quantified this probability across numerous sequences predicted to form single alpha helices (**Figure 10**). Overall, the likelihood of finding a non-ionizable residue at an internal position is only slightly lower than the likelihood of finding a Glu or Lys. This implies that there is an evolutionary drive to reduce some of the charge state heterogeneity that comes with the combinatorial increase of the numbers of charge states as the lengths of SAH forming sequences increase. Interestingly, we also observe a finite, albeit small likelihood that a Glu or Lys residue at an internal position will be occupied by a residue of the opposite charge. It is likely that these charge inverting interruptions disrupt the propagation of the SAH motif.

Figure 10: Stacked bar plots showing the substitution patterns for Glu and Lys residues within internal repeats across 240,754 predicted SAH sequences extracted from the CSAH database [54]. Conservation implies that the charge at a specific position within the idealized repeat is preserved when compared to the consensus repeat. Substitution to uncharged residues quantifies the fraction of residues in that position in the idealized repeat that have been substituted with uncharged residues in naturally occurring sequences. Finally, charge inversion quantifies the observed frequency of substituting residues in each position of the idealized repeat whereby the substitution inverts the charge. The database contains all predictions on the Trembl part of the uniprot database [55].

DISCUSSION

Motivated by the pH insensitivity of measured helicity in SAH forming sequences [34] (**Figure 2**), we have used our formulation of the q-canonical ensemble [29] to investigate the effects of charge state heterogeneity and charge regulation. Our studies are made possible by decoupling charge and conformational measurements, and using the information derived from the two sets of independent measurements as restraints for reweighting atomistic simulations.

The picture that emerges is as follows: There are several microstates and hence mesostates with a net charge that have intrinsic helicities that are equivalent to or higher than those of the single microstate within the 0_1 mesostate that has a net charge of zero. The identification of mesostates with high intrinsic helicities comes from joint analysis of the potentiometric titrations and UV-CD measurements as a function of pH. The relevant microstates within each mesostate are obtained from atomistic simulations that use the ABSINTH implicit solvation model and qMC sampling [50].

Away from a pH of 7.5, the number of microstates and hence mesostates that contribute to the q-canonical ensemble increases. These ensembles display high charge state heterogeneity and low conformational heterogeneity, with helical conformations dominating the ensemble. Even at pH 7.5, there is a finite, albeit small likelihood that the $(E_4K_4)_3$ system can sample microstates with a net charge of ± 1. As shown in **Figure 4**, the intrinsic helicities of these mesostates with a net charge are equivalent to those of the 0_1 mesostate. This leads to a different manifestation of the conformational buffering concept that was recently introduced for intrinsically disordered proteins [56]. For the SAH forming systems, several charge microstates / mesostates are compatible with forming stable helical conformations. This charge state heterogeneity or diversity buffers against large-scale conformational changes – a phenomenon we refer to as *conformational buffering*.

We also find that preferential protonation profiles at a given pH (**Figure 9**) are markedly different for the helical vs. disordered ensembles. This highlights how position-specific proton

binding and release combined with the charge state heterogeneity can tilt conformational biases toward specific types of conformations. In effect, one can think of helix formation in SAH sequences as a simultaneous minimization of enthalpy due to helix formation [17, 21] and maximization of sequence entropy through charge state heterogeneity. Our analysis of frequencies of replacement of Glu or Lys residues with uncharged residues (**Figure 10**) suggests that some of the charge state heterogeneity may be reduced for specific sequences in the SAH superfamily, and this reduction might be necessary for the different contexts in which specific SAH-forming elements are used.

From a physico-chemical perspective, we propose that there are three main factors that contribute to the linkage between proton binding and conformational equilibria appear to be as follows: (i) The intrinsic conformational biases of the neutral vs. charged forms of ionizable residues [28, 57]. (ii) The intrinsic [45] and likely sequence / conformational context dependence of the free energies of solvation of neutral vs. charged forms of ionizable residues. And (iii), the interplay between local and non-local electrostatic attractions vs. repulsions [58]. In isolated helical conformations such as the SAH, the functional groups of ionizable residues can be pushed out from the backbone, thus maximizing the extent of solvation [22]. This is important because the free energies of solvation of charged forms of ionizable residues are at least an order of magnitude more favorable than the uncharged forms [45, 46]. However, within helical conformations, there is likely to be an overlap of solvation shells of the Lys^+ and Glu^- residues [23, 59]. These overlaps are likely to be thermodynamically unfavorable because the water molecules within the overlapping solvation shells need to be oriented differently around amines vs. carboxyl groups [45]. Selective, position-specific proton binding or release can alleviate orientational restrictions by minimizing the overlap of solvation shells. (iii) The high local charge density engenders the possibility of local repulsions. These will need to be minimized while enabling favorable i to $i+4$ salt bridges to form. Overall, our findings suggest that some level of charge regulation, the extent of which depends on pH, will contribute to the complex determinants of SAH stability. This observation is consistent with the realization that the determinants of helix stability can be quite complex [60]. There is always an intricate interplay between the enthalpy and entropy of the peptide plus solvent system, and how this is achieved can be highly context specific.

There is considerable interest in the functions of SAHs given that they may be viewed as a rod-like semi-flexible entities [37]. There is also interest in using these systems as rigid molecular rulers for various intracellular conformational measurements [39]. These ideas and usages might need to account for the effects of charge state heterogeneity and accommodate the observation that for finite-sized systems, such as the $(E_4K_4)_3$ peptide studied here, the helical rod is unlikely to be the dominant conformation (**Figure 7**). This has been recently critiqued as an artefact of a specific brand of implicit solvation models [61]. In that study, the authors set up an expectation that $(E_4K_4)_n$ systems should be nearly 100% helical and perfectly rod-like. This expectation goes against what is known and well established in the helix-coil transition literature. The helical rod requires that the sequence length be infinitely long. In practical terms, the length must be closer to or beyond the higher end of the observed length distribution of SAH forming sequences [35] (**Figure S1**). Importantly, simulations based on explicit representations of solvent molecules do not show reversible transitions between helical and non-helical conformations [33]. Instead, they have been used for understanding the role of hydration in mediating i to $i+4$ salt bridges. The key takeaway is that simulations based on classical forcefields, irrespective of whether they are based on implicit vs. explicit solvation models, are likely to be confounded by site- and context-specific protonation effects. These may be accountable in next generation polarizable models such as the HIPPO model

that accounts for polarizability i.e., electron cloud deformation, and the effects of charge transfer [62]. Whether such models can be made efficient and deployed to account for charge regulation effects is unclear at this juncture. For now, the practical approach of combining information from experiments that decouple measurements of charge and conformation with simulations that explicitly account for charge state heterogeneity, best achieved using a well parameterized implicit solvation model such as ABSINTH, is, at least from our perspective, the optimal route to pursue for uncovering the effects of charge regulation. The ABSINTH framework includes the explicit representation of solution ions, and this is useful for dissecting the interplay between charge regulation and charge renormalization (see below).

It is well-known that SAH forming sequences can be misclassified as being intrinsically disordered or as drivers of coiled coils [31, 38]. This raises the formal possibility that some fraction of sequences that are predicted to be intrinsically disordered are likely to form well-ordered conformations. In such systems, especially those enriched in charged residues, there might indeed be low conformational heterogeneity that serves as a buffer against high charge state heterogeneity. This could also be true of regions predicted to be disordered that are also substrates of multisite post-translational modifications such as Ser / Thr phosphorylation, Lys acetylation, Arg methylation or citrullination [63]. What we need are combined interrogations of charge states using potentiometry, conformational states using multipronged spectroscopies, and simulations that account for heterogeneous distributions of charge states and the conversions between these states [50]. We have presented one such method here and summarized the insights that emerge from this combined methodology.

Finally, in addition to charge regulation effects, there are likely to be *charge renormalization* effects due to preferential territorial organization of solution ions around ionizable groups [64, 65]. These effects are likely to be minor at least for the 0_1 mesostate, although a detailed investigation of the interplay of charge regulation and charge renormalization requires deeper scrutiny given that the cellular milieu is a complex mixture of protons, ions, and other solutes [66].

METHODS

Materials: Potassium chloride (KCl), hydrochloric acid (HCl), potassium hydroxide (KOH), and potassium hydrogen phthalate (KHP) were purchased from Sigma (St. Louis, MO, USA). We assessed the purity of KOH using titrations of a standard with a known pK_a, which confirm that contaminants, if present are miniscule and do not affect the potentiometric titrations [29]. Peptides were purchased from GenScript (Piscataway, NJ, USA) at >95% purity as HCl salts with acetylated N-termini and amidated C-termini to avoid confounding effects from charged termini. Salt content was determined by mass spectrometry. Peptides were stored in lyophilized form at -20°C in sealed containers in the presence of desiccant until used for experiments. Three peptides were used in this study: ac-G-$(E_4K_4)_n$-GW-NH_2, where n = 1, 2, or 3. Tryptophan was used to enable precise measurements of concentration.

Potentiometry. Potentiometric titrations were carried out in CO_2-depleted solutions as previously described in an earlier report [29]. The data used here are identical to those in the previously published work [29].

UV-CD measurements: Samples were prepared for UV-CD measurements by carrying out a potentiometric titration as previously described [29], except that at regular increments throughout the titration, 40 µL aliquots were removed from the sample being titrated and set aside for UV-CD measurements. Subsequently, these aliquots were loaded sequentially into a 0.1 mm

pathlength demountable quartz cuvette, and UV-CD spectra were collected at 20°C using a JASCO J-810 spectropolarimeter. All UV-CD spectra were the average of six scans from 260-190 nm, with a 1 nm step, a 2 second response time, and 50 nm/min scanning speed. The UV-CD spectra measured for background solution at low and high pH were identical within error, so the low pH background was subtracted from all spectra. The resulting ellipticity was used to calculate the mean residue ellipticity ([θ], units of deg cm²/dmol residue) using Equation (3):

$$[\theta] = \frac{\theta}{(N-1)L_{mm}C_M} \; ; (3)$$

where θ = ellipticity (mdeg, machine units), N is the number of residues, $N–1$ is the number of backbone amides, L_{mm} is the path length of the cuvette (mm), and C_M is the protein concentration in molar units.

      Deconvolution of UV-CD data. Data were converted to units of mean residue ellipticity as described above and then analyzed by deconvolution to obtain estimates of secondary structure content. Deconvolution analysis was carried out using the CDSSTR algorithm as implemented through the Dichroweb server. Basis sets 4, 7, and SP175 were used for the deconvolution, as these are the basis sets that have been optimized for the wavelength rage of 190-240 nm. The CDSSTR algorithm consistently provided the best fits for our data and outperformed other deconvolution algorithms including CONTIN and SELCON3. The outputs of the deconvolution analysis are reported as fractional secondary structure content in six categories: alpha helix regular, alpha helix distorted, beta sheet regular, beta sheet distorted, turns, and unordered. For the sake of simplicity, we sum the two helical categories to obtain a single estimate of fractional helicity. The secondary content estimates reported in this work represent the mean and standard deviation of the values resulting from analysis using each of the three basis sets (4, 7, SP175).

Extracting intrinsic helicities of mesostates

      We recently published a protocol to extract the probabilities of individual mesostates as a function of pH [29]. As noted in Equation (1), the overall helicity profile as a function of pH is the weighted sum of the convolution of the probability of accessing a mesostate of net charge q at a given pH and the intrinsic helicity associated with the mesostate. The latter, as noted in Equation (1), is independent of pH and intrinsic to the mesostate. The intrinsic helicities of mesostates are inferred through a Monte Carlo based numerical fitting procedure designed to reproduce the measured helicity profile, which is denoted by the left-hand side of Equation (1). The mesostate populations as a function of pH are known from potentiometric titrations. Accordingly, we define a cost function as the difference in between the simulated curve and the experimental profile:

$$E_{fit} = \sum_i^{Npoints} |f_{pH_i}^{expt} - f_{pH_i}^{model}| \; ; (4)$$

      Here, pH_i is the pH at which the data point of interest was collected, $f_{pH_i}^{expt}$ is the experimental data point, and $f_{pH_i}^{model}$ is the value for the fractional helicity that we estimate at pH_i from the current set of intrinsic helicities of mesostates. The latter are chosen by sampling at random, in an interval between 0 and 1 and accepting or rejecting the proposed move in parameter space using the Metropolis-Hastings criterion defined in Equation (5) as:

$$p_{acc} = e^{-\frac{\Delta E_{fit}}{T_{fit}}} \; ; (5)$$

Here, p_{acc} is the acceptance probability of a specific move to increase or decrease the fraction helicity, ΔE_{fit} is the difference in fitting energy between the last accepted move, and the current proposed move. This step is repeated for 10^4 steps and each Monte Carlo sweep involving 10^4 steps was performed separately and restarted 10^4 times.

ABSINTH Simulations

All the simulation were performed using the ABSINTH implicit solvent model [49, 67]. We used the q-canonical Monte Carlo sampling method [50], with a few modifications. Instead of using the populations of each microstate within a single mesostate simulation to obtain the standard free energy differences, we performed individual simulations for each of the microstates. The temperature for these simulations as set to be 240 K for the unmodified ensemble, 350 K for the disordered ensemble, and 278 K for the helical ensemble. These simulation temperatures are the same for all repeat lengths. To generate the final ensemble, we set all temperatures to the experimental temperature of 298 K and mix conformations drawn from the three different ensembles. For each microstate, the ABSINTH-based Monte Carlo simulations use at 10^8 steps. Information from the first set of $5{\times}10^6$ steps was not used in the analysis, as this was the equilibration phase in the simulations. The conformation and frame specific energies were saved every 10^4 steps. We then derive the free energy differences between microstates by performing cross energy calculations. These calculations entail reassessing the energy for each frame obtained for a given microstate while substituting the potential of a different microstate. The cross energies were then used as input for the Bennett Acceptance Ratio (BAR) procedure, as implemented in the python package pymbar [68].

Reweighting of the simulations

We use the helicity per mesostates and mesostate populations as a function of pH to reweight the overall ensemble by appropriate mixing of the three simulated ensembles. The three ensembles are then mixed, and mixing is the reweighting procedure that is designed to reproduce the intrinsic helicities of individual mesostates. For each mesostate, we start by assessing the overall helicity in each of the three sets. We then compare the mesostate helicities of the unmodified set to the helicity per mesostate that we obtained from experiments. If the helicity is too high for that mesostate compared to the experimentally derived value, we introduce the first set until the overall weighted sum of the two sets are within 0.001 of experiments. Conversely, if the helicity is too low for that mesostate compared to the experimentally derived value, we introduce the third set to find the overall ensemble. The proportion of mixing of each ensemble is shown in **Figure S6**. Using the mixing function, we combined all properties of each microstate within each mesostate to match experiments. We then set the mesostate free energies to the free energies obtained from potentiometric data, while conserving the properly mixed microstate weights within mesostates. This means that the average free energy of each mesostate is the potentiometrically derived values, but the free energy differences within mesostates are those derived from simulations, after proper mixing. This allows us to extract microstate weights that are consistent with experiments, while also generating structural insights on a per residue basis. Because of the large number of simulations needed to get the full pH dependence in (E$_4$K$_4$)$_3$, we

focused on the mesostates with net charge ranging from -3 to 3. This is reasonable because potentiometry shows that these mesostates contribute to more than 99% of overall population across pH values between 5.3 to 9.7 [29]. For the constructs of $(E_4K_4)_1$ and $(E_4K_4)_2$ repeats, the computational cost is significantly lower, and we were able to obtain simulations for every microstate. The simulations were analyzed using the MDtraj python package [69]. This includes determination of the secondary structure contents, quantification of the numbers of salt bridges, etc. In our analysis, a salt bridge was approximated using a simple average of the number of counts of opposite charge ionized residue pairs within 4 Å of each other.

ASSOCIATED CONTENT

The supplemental material includes Figures S1 – S8.

AUTHOR INFORMATION

Corresponding Author

*pappu@wustl.edu

ORCID

Martin J. Fossat: 0000-0003-3078-2886

Ammon E. Posey: 0000-0002-1445-6522

Rohit V. Pappu: 0000-0003-2568-1378

Author Contributions

All authors designed the study. M.J.F., developed the *q*-canonical ensemble framework. M.J.F., designed the simulation setup, performed the simulations, and analyzed the simulation results. A.E.P., designed and deployed the potentiometric titrations and performed all the measurements. M.J.F., and A.E.P., analyzed the experimental data. All authors contributed to the writing and editing of the manuscript. R.V.P., secured funding.

§M.J.F. and A.E.P contributed equally to this work.

ACKNOWLEDGMENTS

This work was supported by grants from the Air Force Office of Scientific Research (FA9550-20-1-0241), the US National Institutes of Health (5R01NS056114), and the US National Science Foundation (MCB-1614766). We thank Mina Farag, Matthew King, Min Kyung Shinn, and Xiangze Zeng for help with the manuscript.

REFERENCES

1. Pauling, L., Corey, R.B. & Branson, H.R. The structure of proteins: Two hydrogen-bonded helical configurations of the polypeptide chain. *Proceedings of the National Academy of Sciences* **37**, 205-211 (1951).

2. Eisenberg, D. The discovery of the alpha-helix and beta-sheet, the principal structural features of proteins. *Proceedings of the National Academy of Sciences* **100**, 11207-11210 (2003).

3.	Kabsch, W. & Sander, C. Dictionary of Protein Secondary Structure - Pattern-Recognition of Hydrogen-bonded and Geometrical Features. *Biopolymers* **22**, 2577-2637 (1983).

4.	Zimm, B.H. & Bragg, J.K. Theory of the Phase Transition between Helix and Random Coil in Polypeptide Chains. *The Journal of Chemical Physics* **31**, 526-535 (1959).

5.	Lifson, S. & Roig, A. On the Theory of Helix—Coil Transition in Polypeptides. *The Journal of Chemical Physics* **34**, 1963-1974 (1961).

6.	Poland, D.C. & Scheraga, H.A. Comparison of Theories of the Helix—Coil Transition in Polypeptides. *The Journal of Chemical Physics* **43**, 2071-2074 (1965).

7.	Vitalis, A. & Caflisch, A. 50 Years of Lifson–Roig Models: Application to Molecular Simulation Data. *Journal of chemical theory and computation* **8**, 363-373 (2012).

8.	Aurora, R., Creamer, T.P., Srinivasan, R. & Rose, G.D. Local Interactions in Protein Folding: Lessons from the α-Helix *. *Journal of Biological Chemistry* **272**, 1413-1416 (1997).

9.	Marqusee, S., Robbins, V.H. & Baldwin, R.L. Unusually stable helix formation in short alanine-based peptides. *Proceedings of the National Academy of Sciences* **86**, 5286-5290 (1989).

10.	Padmanabhan, S., Marqusee, S., Ridgeway, T., Laue, T.M. & Baldwin, R.L. Relative helix-forming tendencies of nonpolar amino acids. *Nature* **344**, 268-270 (1990).

11.	Chakrabartty, A. & Baldwin, R.L. in Advances in Protein Chemistry, Vol. 46. (eds. C.B. Anfinsen, F.M. Richards, J.T. Edsall & D.S. Eisenberg) 141-176 (Academic Press, 1995).

12.	Doig, A.J. & Baldwin, R.L. N- and C-capping preferences for all 20 amino acids in α-helical peptides. *Protein science : a publication of the Protein Society* **4**, 1325-1336 (1995).

13.	Aurora, R. & Rosee, G.D. Helix capping. *Protein science : a publication of the Protein Society* **7**, 21-38 (1998).

14.	Robustelli, P., Piana, S. & Shaw, D.E. Mechanism of Coupled Folding-upon-Binding of an Intrinsically Disordered Protein. *Journal of the American Chemical Society* **142**, 11092-11101 (2020).

15.	Shiraki, K., Nishikawa, K. & Goto, Y. Trifluoroethanol-induced Stabilization of the α-Helical Structure of β-Lactoglobulin: Implication for Non-hierarchical Protein Folding. *Journal of molecular biology* **245**, 180-194 (1995).

16.	Myers, J.K., Nick Pace, C. & Martin Scholtz, J. Trifluoroethanol effects on helix propensity and electrostatic interactions in the helical peptide from ribonuclease T1. *Protein science : a publication of the Protein Society* **7**, 383-388 (1998).

17.   Lopez, M.M., Chin, D.-H., Baldwin, R.L. & Makhatadze, G.I. The enthalpy of the alanine peptide helix measured by isothermal titration calorimetry using metal-binding to induce helix formation. *Proceedings of the National Academy of Sciences* **99**, 1298-1302 (2002).

18.   Ghosh, K. & Dill, K.A. Theory for Protein Folding Cooperativity: Helix Bundles. *Journal of the American Chemical Society* **131**, 2306-2312 (2009).

19.   Rogers, J.M., Wong, C.T. & Clarke, J. Coupled Folding and Binding of the Disordered Protein PUMA Does Not Require Particular Residual Structure. *Journal of the American Chemical Society* **136**, 5197-5200 (2014).

20.   Oldfield, C.J. et al. Coupled Folding and Binding with α-Helix-Forming Molecular Recognition Elements. *Biochemistry* **44**, 12454-12470 (2005).

21.   Marqusee, S. & Baldwin, R.L. Helix stabilization by Glu-...Lys+ salt bridges in short peptides of de novo design. *Proceedings of the National Academy of Sciences* **84**, 8898-8902 (1987).

22.   Lu, H. et al. Ionic polypeptides with unusual helical stability. *Nature communications* **2**, 206 (2011).

23.   Sundaralingam, M., Sekharudu, Y.C., Yathindra, N. & Ravichandran, V. Ion pairs in alpha helices. *Proteins: Structure, Function, and Bioinformatics* **2**, 64-71 (1987).

24.   Stanley, C.B. & Strey, H.H. Osmotically Induced Helix-Coil Transition in Poly(Glutamic Acid). *Biophysical Journal* **94**, 4427-4434 (2008).

25.   Olander, D.S. & Holtzer, A. Stability of the polyglutamic acid .alpha. helix. *Journal of the American Chemical Society* **90**, 4549-4560 (1968).

26.   Nagasawa, M. & Holtzer, A. The Helix-Coil Transition in Solutions of Polyglutamic Acid. *Journal of the American Chemical Society* **86**, 538-543 (1964).

27.   Gooding, E.A. et al. pH-dependent helix folding dynamics of poly-glutamic acid. *Chemical Physics* **422**, 115-123 (2013).

28.   Nick Pace, C. & Martin Scholtz, J. A Helix Propensity Scale Based on Experimental Studies of Peptides and Proteins. *Biophysical Journal* **75**, 422-427 (1998).

29.   Fossat, M.J., Posey, A.E. & Pappu, R.V. Quantifying charge state heterogeneity for proteins with multiple ionizable residues. *Biophysical Journal* **120**, 5438-5453 (2021).

30.   Boyle, A.L. in Peptide Applications in Biomedicine, Biotechnology and Bioengineering. (ed. S. Koutsopoulos) 51-86 (Woodhead Publishing, 2018).

31.   Suveges, D., Gaspari, Z., Toth, G. & Nyitray, L. Charged single alpha-helix: a versatile protein structural motif. *Proteins: Structure, Function, Bioinformatics* **74**, 905-916 (2009).

32.   Iwata, S. et al. Complete Structure of the 11-Subunit Bovine Mitochondrial Cytochrome bc Complex. *Science (New York, N.Y.)* **281**, 64-71 (1998).

33.   Wolny, M. et al. Characterization of long and stable de novo single alpha-helix domains provides novel insight into their stability. *Scientific Reports* **7**, 44341 (2017).

34.   Baker, E.G. et al. Local and macroscopic electrostatic interactions in single α-helices. *Nature Chemical Biology* **11**, 221-228 (2015).

35.   Barnes, C.A. et al. Remarkable Rigidity of the Single α-Helical Domain of Myosin-VI As Revealed by NMR Spectroscopy. *Journal of the American Chemical Society* **141**, 9004-9017 (2019).

36.   Iwai, S., Hanamoto, D. & Chaen, S. A Point Mutation in the SH1 Helix Alters Elasticity and Thermal Stability of Myosin II *. *Journal of Biological Chemistry* **281**, 30736-30744 (2006).

37.   Spudich, J.A. & Sivaramakrishnan, S. Myosin VI: an innovative motor that challenged the swinging lever arm hypothesis. *Nature Reviews Molecular Cell Biology* **11**, 128-137 (2010).

38.   Peckham, M. & Knight, P.J. When a predicted coiled coil is really a single α-helix, in myosins and other proteins. *Soft Matter* **5**, 2493-2503 (2009).

39.   Malik, R.U., Dysthe, M., Ritt, M., Sunahara, R.K. & Sivaramakrishnan, S. ER/K linked GPCR-G protein fusions systematically modulate second messenger response in cells. *Scientific Reports* **7**, 7749 (2017).

40.   Wang, E. & Wang, C.-L.A. (i,i+ 4) Ion Pairs Stabilize Helical Peptides Derived from Smooth Muscle Caldesmon. *Archives of Biochemistry and Biophysics* **329**, 156-162 (1996).

41.   Fitch, C.A. et al. Experimental pK(a) values of buried residues: analysis with continuum methods and role of water penetration. *Biophysical Journal* **82**, 3289-3304 (2002).

42.   Andrew, C.D., Warwicker, J., Jones, G.R. & Doig, A.J. Effect of Phosphorylation on α-Helix Stability as a Function of Position. *Biochemistry* **41**, 1897-1905 (2002).

43.   Errington, N. & Doig, A.J. A Phosphoserine−Lysine Salt Bridge within an α-Helical Peptide, the Strongest α-Helix Side-Chain Interaction Measured to Date. *Biochemistry* **44**, 7553-7558 (2005).

44.   Simm, D., Hatje, K. & Kollmar, M. Distribution and evolution of stable single α-helices (SAH domains) in myosin motor proteins. *PloS one* **12**, e0174639 (2017).

45.   Fossat, M.J., Zeng, X. & Pappu, R.V. Uncovering Differences in Hydration Free Energies and Structures for Model Compound Mimics of Charged Side Chains of Amino Acids. *The Journal of Physical Chemistry B* **125**, 4148-4161 (2021).

46. Zeng, X. et al. Design of intrinsically disordered proteins that undergo phase transitions with lower critical solution temperatures. *APL Materials* **9**, 021119 (2021).

47. Fili, N. & Toseland, C.P. Unconventional Myosins: How Regulation Meets Function. *International Journal of Molecular Sciences* **21**, 67 (2020).

48. Li, J., Lu, Q. & Zhang, M. Structural Basis of Cargo Recognition by Unconventional Myosins in Cellular Trafficking. *Traffic* **17**, 822-838 (2016).

49. Vitalis, A. & Pappu, R.V. ABSINTH: A New Continuum Solvation Model for Simulations of Polypeptides in Aqueous Solutions. *Journal of computational chemistry* **30**, 673-699 (2009).

50. Fossat, M.J. & Pappu, R.V. q-Canonical Monte Carlo Sampling for Modeling the Linkage between Charge Regulation and Conformational Equilibria of Peptides. *The Journal of Physical Chemistry B* **123**, 6952-6967 (2019).

51. Platzer, G., Okon, M. & McIntosh, L.P. pH-dependent random coil 1H, 13C, and 15N chemical shifts of the ionizable amino acids: a guide for protein pKa measurements. *Journal of Biomolecular NMR* **60**, 109-129 (2014).

52. Ermolenko, D.N., Richardson, J.M. & Makhatadze, G.I. Noncharged amino acid residues at the solvent-exposed positions in the middle and at the C terminus of the α-helix have the same helical propensity. *Protein science : a publication of the Protein Society* **12**, 1169-1176 (2003).

53. Ermolenko, D.N., Thomas, S.T., Aurora, R., Gronenborn, A.M. & Makhatadze, G.I. Hydrophobic Interactions at the Ccap Position of the C-capping Motif of α-Helices. *Journal of molecular biology* **322**, 123-135 (2002).

54. Kovács, Á. et al. Detection of single alpha-helices in large protein sequence sets using hardware acceleration. *Journal of Structural Biology* **204**, 109-116 (2018).

55. Wang, Y. et al. A crowdsourcing open platform for literature curation in UniProt. *PLOS Biology* **19**, e3001464 (2021).

56. González-Foutel, N.S. et al. Conformational buffering underlies functional selection in intrinsically disordered protein regions. *Nature Structural & Molecular Biology* **29**, 781-790 (2022).

57. Pace, C.N., Grimsley, G.R. & Scholtz, J.M. Protein Ionizable Groups: pK Values and Their Contribution to Protein Stability and Solubility. *Journal of Biological Chemistry* **284**, 13285-13289 (2009).

58. Das, R.K. & Pappu, R.V. Conformations of intrinsically disordered proteins are influenced by linear sequence distributions of oppositely charged residues. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 13392-13397 (2013).

59. Sundaralingam, M. & Sekharudu, Y.C. Water-Inserted alpha-Helical Segments Implicate Reverse Turns as Folding Intermediates. *Science (New York, N.Y.)* **244**, 1333-1337 (1989).

60. Richardson, J.M., Lopez, M.M. & Makhatadze, G.I. Enthalpy of helix–coil transition: Missing link in rationalizing the thermodynamics of helix-forming propensities of the amino acid residues. *Proceedings of the National Academy of Sciences* **102**, 1413-1418 (2005).

61. Lang, E.J.M., Baker, E.G., Woolfson, D.N. & Mulholland, A.J. Generalized Born Implicit Solvent Models Do Not Reproduce Secondary Structures of De Novo Designed Glu/Lys Peptides. *Journal of chemical theory and computation* **18**, 4070-4076 (2022).

62. Rackers, J.A., Silva, R.R., Wang, Z. & Ponder, J.W. Polarizable Water Potential Derived from a Model Electron Density. *Journal of chemical theory and computation* **17**, 7056-7084 (2021).

63. Harauz, G. & Musse, A.A. A Tale of Two Citrullines—Structural and Functional Aspects of Myelin Basic Protein Deimination in Health and Disease. *Neurochemical Research* **32**, 137-158 (2007).

64. Krishnan, M. A simple model for electrical charge in globular macromolecules and linear polyelectrolytes in solution. *The Journal of Chemical Physics* **146**, 205101 (2017).

65. Netz, R.R. & Orland, H. Variational charge renormalization in charged systems. *The European Physical Journal E* **11**, 301-311 (2003).

66. Milo, R., Jorgensen, P., Moran, U., Weber, G. & Springer, M. BioNumbers—the database of key numbers in molecular and cell biology. *Nucleic Acids Research* **38**, D750-D753 (2009).

67. Mittal, A., Das, R., Vitalis, A. & Pappu, R.V. in Computational Approaches to Protein Dynamics: From Quantum to Coarse-Grained Methods. (ed. M. Fuxreiter) 181-203 (CRC Press (Taylor and Francis Group), Boca Raton, RL; 2015).

68. Shirts, M.R. & Chodera, J.D. Statistically optimal analysis of samples from multiple equilibrium states. *Journal of Chemical Physics* **129**, 124105 (2008).

69. McGibbon, Robert T. et al. MDTraj: A Modern Open Library for the Analysis of Molecular Dynamics Trajectories. *Biophysical Journal* **109**, 1528-1532 (2015).