1 **Title: Wildflower phenological escape differs by continent and spring temperature**
2

3 <u>Author list</u>: Benjamin R. Lee*[1,2,3,4], Tara K. Miller[4], Christoph Rosche[5,6], Yong Yang[7], J. Mason
4 Heberling[1,2], Sara E. Kuebbing[2,8], and Richard B. Primack[4]
5
<u>Affiliations</u>
[1]Section of Botany, Carnegie Museum of Natural History, Pittsburgh, PA
[2]Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA
[3]Holden Forests and Gardens, Kirtland, OH
[4]Biology Department, Boston University, Boston, MA
[5]Institute of Biology/Geobotany and Botanical Garden, Martin-Luther-University Halle-
Wittenberg, Halle, Germany
[6]German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Germany
[7]Co-Innovation Center for Sustainable Forestry in Southern China, College of Biology and the
Environment, Nanjing Forestry University, 159 Longpan Rd., Nanjing 210037, China
[8]Yale Applied Science Synthesis Program, The Forest School at the School of the Environment,
Yale University, 195 Prospect Street, New Haven, CT, USA
18
*Corresponding author: <u>LeeB@CarnegieMNH.org</u>
ORCID: 0000-0002-5256-0515

Keywords: Leaf-out, flowering, climate change, phenological escape, trees, wildflowers,
herbarium collections, Europe, East Asia, Eastern North America

Abstract: Temperate understory plant species are at risk from climate change and

anthropogenic threats that include increased deer herbivory, habitat loss, pollinator

declines and mismatch, and nutrient pollution. Recent work suggests that spring ephemeral

wildflowers may be at additional risk due to phenological mismatch with deciduous canopy

trees. The study of this dynamic, commonly referred to as "phenological escape", and its

sensitivity to spring temperature is limited to eastern North America. Here, we use

herbarium specimens to show that phenological sensitivity to spring temperature is

remarkably conserved for understory wildflowers across North America, Europe, and

Asia, but that canopy trees in North America are significantly more sensitive to spring

temperature compared to in Asia and Europe. We predict that advancing tree phenology

will lead to decreasing spring light windows in North America while spring light windows

will be maintained or even increase in Asia and Europe in response to projected climate

warming.

Introduction

Temperate deciduous forests are a dominant biome in the northern hemisphere, covering

extensive regions in eastern Asia, continental Europe, and eastern North America[1]. They are

characterized by winter-deciduous plant species, and the phenology of leaf out and flowering on

all three continents is sensitive to variation in average spring temperature[2–5]. Furthermore, recent

research has found disparities in phenological sensitivity among plants from different

continents[2,6], suggesting that climate change may affect temperate forests differently depending

on location.

Previous studies indicate differences in spring phenological sensitivity between woody

and herbaceous plants in forests, with direct comparisons available for plants in Asia[3] and North

America[5,7]. Evidence in Europe similarly suggests differences in sensitivities between woody

and herbaceous plants[8,9]. Herbaceous plants may be less sensitive to spring temperature than

trees because the former often overwinter underground and therefore may not respond to the

same phenological cues as the latter, which have extensive aboveground structures[10,11], but this

mechanism has only been studied in North American forests. This potential phenological

mismatch is important because spring-active wildflowers in temperate deciduous forests, as well

as some woody understory species[12–15], often rely on leafing out before the canopy closes in

order to assimilate 40-100% of their annual carbon budget[16]. The success of this strategy to

maximize the spring light window, referred to as "phenological escape"[12,17], is directly

associated with patterns of growth[10,12,17–19], survival[12,18–20], flowering[17,20], and reproductive

output[10,17,21,22]. Recent studies suggest that the duration of spring light windows in eastern North

America is likely to be significantly altered with warming climates[7,13], with herbaceous species

generally expected to experience shorter spring light windows in the coming decades.

Experimental evidence for such mismatches, however, has generally been limited to

comparisons of woody plants located in common garden experiments, which are limited in

geographical and temporal extent and have not evaluated mismatch with understory species. For

example, Zohner and Renner[6,23]have contributed valuable insights into interspecific variation in

plant phenology using woody plants growing in European botanical gardens. It is unknown,

however, if the variation in woody plant phenology found in common garden experiments are

also observed at large scales, for forest trees growing in their natal environment, and across long

time periods[24]. It is similarly unknown whether understory herbaceous plant phenological

sensitivity varies across large spatial and temporal scales.

Here we assess intercontinental, long-term data of the phenological sensitivity of canopy

and understory forest plants and the potential for phenological mismatch across temperate forests

in North America, Europe, and Asia. We evaluate the spring phenology of 5,522 herbarium

specimens collected between 1901 and 2020, representing 22 common tree and 18 common

wildflower species found in temperate deciduous forests on three continents (six wildflower

species per continent and six, six, and ten tree species in Europe, North America, and Asia,

respectively; see full species list in Table S1 and distribution of observations in Fig. S1).

Herbarium collections provide data from across large temporal (centuries) and spatial

(intercontinental) scales difficult to match with other methods. Following previously validated

methods[25,26], we model Leaf Out Date (LOD) of overstory canopy trees and First Flowering Date

(FFD) of spring-blooming understory wildflowers (perennial herbaceous species that leaf out at

approximately the same time as they flower in the early spring) in response to average spring

(March-April) temperature of the collection year and location. This period is chosen because

most (> 99%) of the observed phenology occurred during or following this period (Fig. 1a;

means and standard deviations are listed in Table S2) and because initial variable analysis

identified this temperature variable as the best for predicting tree and wildflower spring

phenology. Additionally, owing to the absence of daily weather records for the vast majority of

years and sampling locations, especially Asia, we were unable to calculate and analyze finer-

resolution daily temperature data for most geographic locations, which would be required for

intercontinental comparisons (see Supplementary Information). We use estimated model

parameter values to compare phenological sensitivities of trees and wildflowers among

continents and to project how spring light window duration will change by the end of the current

century based on projected climate trends. Our results suggest that there are differences in the

sensitivity of trees (but not wildflowers), with North American trees considerably more sensitive

to spring temperature than Asian or European trees. This difference has implications for

phenological escape and the future conservation of the species-rich herbaceous layer in

temperate deciduous forests as the climate continues to change.

Results and Discussion

We used a hierarchical Bayesian modeling approach to evaluate the relationship between

spring phenology of tree and wildflower species and various climate drivers (see Methods).

Following model selection, our final model structure included fixed effects of average spring

(March-April) temperature and elevation, as well as species-level random effects. We show

continental distributions of spring temperature values in Fig. 1b (means and standard deviations

are listed in Table S2). We report estimates for spring temperature sensitivities from the final

model structure in the main text. Parameter estimates for elevation sensitivities, as well as model

performance of other potential drivers and combinations of drivers, are reported in Tables S3 and

S4. An extended discussion of model assumptions and limitations is included in the

Supplementary Information.

Sensitivity differences by strata

Tree leaf out phenology (LOD) was substantially more sensitive to average spring

temperature in North America (mean = -3.62 days $°C^{-1}$; 95% credible interval (CI) = [-3.76, -

3.49]) than in Europe (mean = -2.79; CI = [-3.27, -2.30]) and Asia (mean = -2.62; CI = [-2.97, -

2.26]; Fig. 2). These values are consistent with previously reported phenological sensitivities in

North America[7] (-5.5 to -3.3 days $°C^{-1}$) and Europe[8] (-4.1 to -3.0 days $°C^{-1}$), as the credible

intervals from our results overlap with the reported credible intervals of prior studies. However,

the Asian LOD sensitivity was less sensitive than previously reported[27] (-3.50 to -3.03 days °C^{-1}

^{1}), potentially owing to differences in species selection[28] or model structure. Previously reported

sensitivities were determined in separate studies using either observational data[7,8] or long-term

observation-based weather station data[27]. The general consistency between our findings suggests

that phenology data from herbarium collections are good indicators of patterns in natural

systems[29–31], a point supported by a recent study of phenological sensitivity derived from

herbaria and from observed citizen science data[32]. These herbarium-based results provide

evidence that phenological sensitivity differs across the temperate forest biome (but see ref[33] for

evidence of differences in response to warming and chilling accumulation). To our knowledge,

our study is the first to contrast overstory and understory phenology across multiple continents,

and therefore to find differences in phenological sensitivity between trees and forest wildflowers

across continents. We recommend future studies explore these differences using alternative

approaches and methodologies that focus on the physiological basis for and mechanisms that

underlie these patterns.

In contrast to trees, wildflower sensitivity to spring temperature was similar across all

three continents and exhibited no strong differences (i.e., overlap in 95% Bayesian credible

intervals) among continents (means and 95% credible intervals in brackets: North America = -

3.14, [-3.28, -3.00]; Europe = -3.02, [-3.48, -2.56]; Asia = -3.12, [-3.36, -2.86]; Fig. 2). These

values are also generally consistent with those reported elsewhere in the literature (i.e., 95%

credible intervals overlap with those reported in other studies; -2.2, [-3.7, -0.76] days °C^{-1} in

North America[7] and -3.6, [-4.04, -3.18] days °C^{-1} in Europe[9]), although we are unaware of any

studies that have estimated phenological sensitivity for Asian forest wildflowers in days °C^{-1}. Ge

et al.[3] reports herbaceous plant sensitivity of -5.71 days per decade in Asia ($\pm$ 7.90 standard

deviation; based primarily on long-term observational data), which appears to be roughly

consistent with our model results, but the difference in units makes this more speculative than the

other comparisons. Discrepancies in mean responses between this study and others may be due

in part to different types of data (herbarium specimens versus field observations) and to choice in

focal taxa, as temperature sensitivity has been shown to vary widely across taxa[28].

Particularly noticeable in our results was that r^2 coefficients of predicted versus observed

phenology were much higher in North America (0.70 and 0.76 for wildflower and tree models,

respectively) compared to Asian (0.40 and 0.44, respectively) and European models (0.41 and

0.25, respectively). This difference in model performance could be due to the higher interannual

variability of spring temperatures in North America[33], leading to greater selective pressure for

strong sensitivity to spring temperatures in North American plants. This difference could explain

why North American species exhibit higher correlation of phenology with average spring

temperatures (Table S4). Alternatively, European and Asian species may have stronger

phenological responses to alternative spring forcing windows, winter chilling temperatures, or

photoperiod, relative to the March-April temperature period used in this study (see Methods).

We think the latter explanation is unlikely given the strong correlations of phenology with spring

temperature across all continents (see Supplementary Information – Justification for March-April

Temperature Window).

Herbarium-based phenological models may be improved by accounting for spatial

autocorrelation within the dataset. For example, Willems et al.[9] found that including spatial

autocorrelation significantly improved predictability of European herbaceous flowering

phenology, even when accounting for multiple drivers of spring phenology. We followed a

similar approach as their study and found similar improvements in model performance with the

addition of spatial autocorrelation (Tables S3-4) that had substantial positive effects on r^2 values

of Asian and European models. However, spatial distributions of specimens differed

substantially among continents (see Fig. S2-S4), and these differences could lead to artifacts that

make results unreliable to interpret (see Supplementary Information). Therefore, we focus here

on results for models without spatial autocorrelation while acknowledging that spatial

aggregation of herbarium specimens in Europe and Asia may be partially responsible for the

relatively lower r^2 values. We encourage other researchers to explore this question further both

with our data set and other data sets.

Climate change and spring light windows

The relative difference between wildflower and tree sensitivity varied substantially

among continents, with wildflowers being approximately equally as sensitive to spring

temperature as trees in Asia and Europe but substantially less sensitive (i.e., 95% BCI do not

overlap) than trees in North America (Fig. 2). Importantly, these differences were driven by

changes in tree phenological sensitivities among continents and resulted in different expectations

for spring light window duration (i.e., the difference in time between estimated wildflower

flowering date and canopy tree leaf out date) on different continents under current climate

conditions (Fig. 3), based on modeled leaf out and flowering under a climate scenario derived

from average climate conditions from 2009-2018 (Fig. S5).

Interestingly, the time between leaf out and flowering in North America is greater in the

north than in the south (Fig. 3c), indicating a greater spring light window duration at higher

latitudes. In addition, although there was regional variation in spring light window duration on

each continent, there was broad overlap in duration among continents estimated under current

environmental conditions (North America light window duration averages 11.7 ± 4.1 s.d. days;

European duration averages 14.7 ± 3.0 s.d. days; and Asian duration averages 8.0 ± 4.6 s.d.

days). This suggests that, under current climate conditions, wildflowers across all continents

experience similar length of spring light windows, but the impact of warming on shrinking or

expanding window size will differ among continents with uncertain impacts on wildflower

populations.

Still, differences among continents resulted in different projections for how spring light

window duration will respond to climate change over the coming century (Fig. 4), which will

likely have implications for understory plant demography[7,12,13,16]. We used climate change

projections for 2081-2100 (assuming an extreme climate change scenario; Institut Pierre-Simon

Laplace CM6A-LR climate model – shared socioeconomic pathway 585; see Methods for more

details) to forecast wildflower flowering date and canopy tree LOD for the end of the current

century (Fig. S6) and then calculated the difference between forecasted wildflower flowering and

tree leaf out phenology for the period of 2081-2100 (future spring light window duration; Fig.

S7). To estimate the change in spring light window duration between now and the end of the

century (Fig. 4), we subtracted the forecasted future light window duration from the modeled

light window duration under current climate conditions.

Dramatic differences in the projected change in spring light window duration emerged,

ranging from increasing spring light window durations for wildflowers in Asia, to minimal

change in spring light window length in Europe, to decreasing spring light window length in

North America. Importantly, these differences in phenological escape trajectories are primarily

attributable to differences in phenological sensitivities between overstory and understory species

and not to differences in projected spring temperature changes among continents. Projected

changes in March-April temperature broadly overlapped across continents (mean ± s.d. of

projected change in spring temperature is 5.41 ± 1.87 °C, 6.36 ± 2.36 °C, and 5.04 ± 2.92 °C in

Asia, Europe, and North America, respectively; Fig. S8). Intracontinental differences, however,

such as the more extreme projected changes in phenological escape in the northern regions of

each continent, are due to projected spring temperature change across latitudes, with higher

latitude regions expected to experience greater spring warming relative to lower latitudes[34,35].

The North American results are consistent with findings from a previous study which

similarly predicted reduction in access to spring light at a single site in the U.S. Northeast[7].

Although no previous studies have specifically addressed phenological escape in Asia, our

results are consistent with previous research that found higher sensitivity to spring temperature

for Asian herbaceous plants compared to Asian woody trees and shrubs[3]. Similarly, we are not

aware of European studies that have directly compared phenological sensitivity between

wildflowers and trees nor that have quantified phenological escape, but our estimates of

sensitivity are consistent with previously reported values[8,9]. Together, this supports our

conclusion that the duration of spring light windows will be differently affected by climate

change on different continents.

Importantly, it is impossible to fully measure the ecological and demographic

implications of these results using herbarium specimens. Linking changing access to light to

changes in photosynthetic carbon assimilation, growth, and survival necessitates the use of

experimental studies of living specimens such as those previously conducted in eastern North

America[7,12,13,16]. Previous studies that have done so have linked access to spring light before

canopy closure with the growth[10,12,17–19], survival[12,18–20], flowering[17,20], and reproductive

output[10,17,21,22] of understory plant species. This large body of research suggests that reductions in

access to spring light can have negative ramifications for the demographic success of understory

species.

For example, projected reductions of several days in phenological escape may not

intuitively seem like much, but the spring ephemeral wildflowers in this study have growing

seasons that last only 3-5 weeks. A decrease in one day of phenological escape can therefore be

as much as a 4% loss of total growing season length. In a study of phenological escape in sugar

maple seedlings, Kwit et al.[14] found that increasing access to spring light by three days could

increase photosynthetic carbon assimilation by more than 50%, an amount that other studies

suggest would substantially increase annual growth and probability of survival[12,13]. For spring

ephemeral herbaceous plants, Heberling et al.[16] showed that a similar three-day loss in access to

spring light resulted in up to a 30% loss in spring carbon assimilation. It is important to note,

however, that the exact impacts of change in phenological escape on plant performance in Asia

and Europe remains to be directly measured.

Thus, although our results suggest strong differences in change in performance when

assuming minimal difference in photosynthetic rates among species on different continents, it is

possible that systematic intercontinental differences could lead to different outcomes. For

example, if spring wildflowers were to be shown to have much lower spring photosynthetic rates,

differences in projected access to light could have much less impact on overall carbon

assimilation and demographic performance. Still, others have found relatively strong

relationships between spring light availability and understory plant performance on other

continents[21], suggesting that photosynthetic rates are similar enough to be comparable in this

study. We therefore conclude that the relatively small changes in spring light availability

projected here are likely to have substantial ecological consequences for the performance and

demography of understory plant species.

The future for spring ephemerals

The herbaceous layer accounts for more than 80% of plant species in temperate forests

worldwide[36] and provides a critical role in the functioning of these ecosystems[37]. Understory

wildflower species face many threats that include deer herbivory[38–40], habitat loss[41], pollinator

declines[42] and mismatch[43,44], presence of nonnative, invasive plants[45,46], and nutrient

pollution[47,48]. Furthermore, there is substantial evidence that many of these species are limited in

their dispersal ability[41,49] and populations are likely unable to shift their distributions as rapidly

as regional climates are predicted to warm. Our results show that mismatch between understory

and overstory phenology, specifically in North America, is another concern for this already

threatened group[36,50]. Reductions in spring light windows will likely lead to reduced carbon gains

each year for spring forest wildflowers[7,16], which may ultimately lead to population decline as

plant fitness declines. Therefore, we highlight the need for future conservation efforts to consider

the impact of loss of spring light in restoration planning, with simultaneous studies of overstory

and understory responses together[37]. Furthermore, this information may assist conservation

practitioners in their decisions of where to focus management or conservation efforts based on

projected changes to future forest ecosystems.

The patterns examined in this study point to fascinating distinctions in plant phenological

responses among continents, but they cannot provide a mechanistic explanation for why these

differences exist. Wildflowers and trees advance their phenology with warming spring

temperatures, for example, but why are North American trees more responsive than co-occurring

wildflower species to warming? Some speculate it is because perennial wildflowers overwinter

underground whereas tree buds overwinter aboveground; trees may therefore be more affected

by air temperature relative to wildflowers[10,11]. Although global soil and near-surface temperature

data are recently becoming available[51], there are as yet no historical estimates of soil

temperatures for use in herbarium studies. Thus, future empirical studies will be needed to

directly test this theory. Similarly, what are the mechanisms that allow North American trees to

be more sensitive to spring temperature than trees in Europe and Asia? Studies using dormant

twigs, potted seedlings, and trees growing in botanical gardens are possible methods for

investigating this. Common garden experiments with manipulated treatments may provide a

more mechanistic understanding of our observations, including to rule out potentially

confounding variables such as intercontinental differences in soil properties, precipitation, and

photoperiod.

Other recent research highlights the growing need to assess the potential for nonlinear

phenological responses to spring warming. As reviewed by Wolkovich et al.[52], future

phenological responses to spring warming may be nonlinear owing to a number of reasons

(including threshold responses to extreme environmental conditions and interactions among

drivers besides spring forcing). Accounting for factors that might drive nonlinearities in plant

phenological response to warming will necessitate controlled experiments[52,53], which was not

possible with the observational herbarium dataset used in our study. We did not find evidence for

nonlinear responses to spring temperature, but we acknowledge that experimental studies that

can experimentally manipulate and test for the impacts of spring forcing, winter chilling, and

photoperiodic effects on plant phenology could further illuminate how nonlinearity affects

phenological escape. Also, nonlinear responses may become more evident in the future if

extreme weather conditions occur more frequently.

Our study represents an early attempt to address phenological questions using herbarium

specimens at the intercontinental level. Digitization of herbarium specimens and other historical

data and citizen science programs like iNaturalist and iSpot are making data increasingly

available and enabling new research[54]. Additional research that utilizes international herbaria and

collaborations to synthesize these data at a global scale will aid in addressing pressing ecological

and climate change questions.

In sum, our results indicate that spring ephemeral wildflowers are similarly sensitive to

spring warming across temperate forests in Asia, Europe, and North America, but that deciduous

canopy trees in North America are more sensitive to spring warming compared to trees on the

other two continents. Furthermore, differences in tree leaf-out sensitivity mean that North

American wildflowers are projected to experience decreasing access to spring light under climate

change conditions whereas those in Asia and Europe are unlikely to experience substantial gains

or losses in spring light availability. These intercontinental differences suggest that climate

change may adversely affect spring ephemerals in North America more than on the other two

continents, and thus that North American wildflowers may be especially threatened by warming

springs.

<u>Methods</u>

Phenological data

Herbarium specimens from North America (see Miller et al.[5]), East Asia, and Europe

were evaluated for common species of deciduous overstory tree species and understory

deciduous wildflowers, prioritizing congeneric species living on more than one continent when

possible, and with genus and family in mind when not possible. Specimens were scored for

either Leaf Out Date[25] (LOD, tree species) or First Flowering Date[26,31] (FFD, wildflower

species), consistent with previous studies comparing phenology across forest strata[7,55]. This

approach assumes that herbaceous species flowering time is tightly correlated with timing of leaf

out for these species[7]. This approach is currently widely accepted to be the best we have to assess

shifts in phenology using herbarium specimens[31], but see also Buonaiuto et al.[56]. In particular,

we selected wildflower species that flowered and leafed out at approximately the same time in

early spring.

Specifically, we searched digitized repositories of herbarium collections on each of the

three target continents. Digitized Asian herbarium specimens were provided by the Chinese

Virtual Herbarium (https://www.cvh.ac.cn/) and Chinese Field Herbarium

(http://english.ib.cas.cn/). European specimens were located and collated using the Global

Biodiversity Information Facility (gbif; https://www.gbif.org/). In North America, digitized

specimens were collated from seven online repositories, further detailed in Miller et al.[5]. We

conducted searches for each species independently, prioritizing spring ephemeral wildflowers

and deciduous tree species with the highest number of available observations across most of the

relevant geographic range. We then filtered our initial searches by phenophase (scored by hand),

making sure we only included observations that were flowering (for the wildflowers) or newly

leafed out (trees). Next, we identified specimens that were missing information but for which

digitized images of the specimen sheet were available. For these individuals, we entered missing

phenological and georeferencing data when available (see below). Individuals with missing

information and that either were not fully digitized or that did not include the necessary

information on the digitized herbarium sheet were excluded from further study.

Specimens were georeferenced by researchers fluent in the languages in which the

specimen data were collected in. Georeferencing was completed using the most precise

geographic scale recorded at time of collection (i.e., prioritized by exact coordinates,

town/locality, county). Specimens that did not identify location to at least county (in North

America) or locality-level (in Asia and Europe) were excluded from this analysis, which is

common practice in herbarium studies given the lack of precise geolocation data for some older

specimens. Only specimens collected within each species' native range were used. Specimens

collected prior to 1901 were excluded because climate estimates were not available prior to this

year[57]. In total, after accounting for specimens that were excluded from our original search, we

collected data for a total of 40 species (22 tree species and 18 herbaceous species; Table S1)

consisting of 5,522 individual specimens. The complete dataset of herbarium specimens used in

this analysis is freely available[58].

Climate data

Climate data were extracted from the Climate Research Unit gridded Temperature Series

(CRU TS) data set[57] v4.05 using the georeferenced location for each specimen. Spring

temperature was calculated as the average of the March and April average monthly temperatures

for the year and location associated with each specimen. This metric is consistent with other

studies which found this period to be important in cueing temperate plant phenology[5,7,13,55] and

preliminary analysis indicated that this was the best spring temperature metric for modeling plant

phenological responses in our dataset (see *Justification for March-April Temperature Window*).

Daily weather records were not available for many years and many locations.

Justification for March-April Temperature Window

Prior to conducting our analysis, we explored the correlation strengths between different

spring temperature windows and observed phenology day of year (DOY). We made the decision

to use a fixed average March-April temperature window for several reasons. First, we were

limited in how we evaluated spring forcing effects by the type of data/climate estimates we were

able to access. Due to the nature of the historical herbarium specimens we used in this study, we

were limited to using historical average temperature estimates aggregated at the monthly level.

We were therefore unable to use more nuanced modeling approaches such as moving windows

or accumulation metrics like growing degree days (e.g., ref[59]).

Next, we evaluated correlation coefficients between different spring temperature

windows (one-, two-, or three-month averages) from between February and May to determine

which window led to the highest correlative strength with observed phenology. We determined

that the March-April window had consistently high correlative strength for wildflower and tree

phenology across all continents (Fig. S9), indicating that it was a robust predictor of spring

phenology. This makes intuitive sense, as the vast majority (83%) of observations in our study

occurred during or after DOY 96 (~March 15th) and nearly all observations (99%) occurring on

or after DOY 60 (~March 1st; Fig. 1a).

Lastly, using a fixed March-April temperature window allows us to directly compare our

results to other phenological escape studies using the same window[5,7,13].

Models and analysis

We modeled the day of observed phenological event (OPE) for individual i of species j

using a normal likelihood distribution:

392                                 $$OPE_{i,j} \sim N(\mu_{i,j}, \sigma^2) \tag{1}$$

394   The mean, μ, was modeled with an intercept term ($\beta 0$), slope terms representing phenological

395   sensitivity to average spring temperature ($\beta 1$) and elevation ($\beta 2$), and species random effects (α_j):

397                     $$\mu_{i,j} = \beta 0 + \beta 1 \times \text{SpringT}_i + \beta 2 \times \text{Elevation}_i + \alpha_j \tag{2}$$

We used slightly informative priors to estimate parameters: $\beta 0$, $\beta 1$, $\beta 2$, $\alpha_j \sim N(0, 1E\text{-}3)$;

$1/\sigma^2 \sim Uniform(0, 100)$. Other potential drivers of leaf-out and flowering phenology were

explored in preliminary analysis (i.e., winter temperatures, annual precipitation, and spring

precipitation), but these drivers did not generally improve model performance (Table S3) and

were thus excluded from the model structure.

Models were run separately for each stratum (i.e., tree vs. herbaceous) x continent

combination using the *R2jags* package[60] (v0.7-1) in R v4.1.0. Parameter values (means,

variances, and covariances) were estimated from posterior distributions and are considered

significantly different if the 95% Bayesian credible intervals (BCIs) of their posterior

distributions do not overlap. Model code and data used to fit each model are publicly available[58].

Description of spatial autocorrelation models

Spring leaf-out and flowering phenology has been extensively linked to many drivers

including spring and winter temperatures, interannual variability in temperature, photoperiod,

timing of snowmelt, precipitation, and elevation[2,5,7,9,12,61,62]. Importantly, many of these drivers

covary along geographical gradients and particularly latitudinal gradients. Recent work

published by Willems et al.[9] found that spatial autocorrelation among observations improved

predictability of understory plant phenology in Europe, even when a host of other drivers were

already accounted for. Their results thus suggest that there may be effects that are not commonly

accounted for in phenology models associated with spatial proximity of the observations and that

including spatial autocorrelation in such models could be important.

In this study, we first evaluated the fit of our phenological models using different

combinations of potential drivers that included average spring temperature, average winter

temperature, annual precipitation, spring precipitation, and elevation. We then assessed how

incorporating spatial autocorrelation into the model structure affected model fit and posterior

distributions of model parameters. To do so, we incorporated Integrated Nested Laplace

Approximations (INLA) that include a spatial autocorrelation term into the model structure via

the *inlabru* package[63] in *R*.

INLA modeling uses Stochastic Partial Differential Equations (SPDEs) as opposed to

MCMC approximation methods[64] but has been demonstrated to maintain high fidelity to MCMC

posterior estimates[65,66] and has the added advantage of allowing for the much faster estimation of

spatial autocorrelative effects[64,66]. To account for the potential effects of including spatial

autocorrelation on our results, we ran the original model structure using *inlabru* instead of *rjags*

without and then with the spatial autocorrelation term added to it. Model fit and performance for

inlabru models with and without spatial autocorrelation are located in Table S3. Comparison of

parameter posterior estimates are located in Table S4. Matérn correlation functions were

constructed in *inlabru* with a maximum edge length of 0.5 decimal degrees and a cutoff value of

0.25 decimal degrees. Resulting meshes can be found in Fig. S10.

A summary of results from this version of the analysis that includes spatial

autocorrelation effects, as well as a detailed justification for having excluded them in the final

analysis presented here, is included in the Supplementary Information.

Climate Change Modeling

To forecast changes in the duration of spring light windows, we compared FFD and LOD

from two different climate simulations. The first simulation represented current environmental

conditions and was estimated by taking the average of spring (March-April) temperatures from a

recent ten-year period (2009-2018). Climate and elevation data were downscaled from the CRU

TS 4.03[57] dataset using WorldClim 2.1[67] for bias correction at a resolution of 2.5 minutes.

Average monthly temperature for each month and each year was calculated as the mean of

monthly minimum and maximum temperatures.

The second simulation represents projected environmental conditions under climate

change at the end of this century (2081-2100). Climate data used in this simulation were

downscaled (using WorldClim 2.1[67]) from the Institut Pierre-Simon Laplace (IPSL) CM6A-LR

climate model[34,68], which is part of the ongoing Climate Model Intercomparison Project[35]

(CMIP6). We specifically used forecasts based on the Shared Socioeconomic Pathway (SSP)

585, which is analogous to the Representative Concentration Pathway (RCP) 8.5 from CMIP5

and earlier. This SSP is the most extreme, "business as usual", pathway used by the

Intergovernmental Panel on Climate Change (IPCC). Therefore, projections made using this

pathway represent the extreme threshold of climate change effects by the end of the century.

Model output is presented for regions where land-use is classified as > 1% temperate

deciduous forest type according to a consensus of four global land-use models[1] (uncropped

versions of figures can be found in the Supplementary Information as Figs. S11-12). Herbarium

specimens were included in the above analyses regardless of the land-use classification of the

respective georeferenced locations. We did not find consistent effects of land-use on

phenological sensitivity, so we did not include this variable as a fixed effect in the final model.

Still, we only present output for land areas currently covered by deciduous forests as those are so

far the only systems where phenological escape has been shown to be important[5,7,10,12,13,16].

Approximations of FFD and LOD (for wildflowers and trees, respectively), were

calculated using posterior mean estimates of the slopes, intercepts, and species-level random

effects of our models (Tables S4-S7). Spring light window length was then calculated as the

difference (in days) between FLD and LOD in each simulation X continent combination. Data

processing and handling was completed primarily using the *ncdf4*[69], *stars*[70], and *sp*[71] packages in

R v4.1.0.

<u>Inclusion and ethics statement:</u> The work presented here is the culmination of international

collaboration and facilitation. Herbarium specimens were collated across three continents by

researchers in those locations who spoke the language(s) in which information was recorded. All

researchers who led these efforts are coauthors on this manuscript and their employees and lab

members who assisted in this effort are listed in the Acknowledgements section. Local and

regional studies have been cited throughout this document, further acknowledging the

contributions of global scholars to our work and to science as a whole.

Data availability: The individual specimen phenology and environmental conditions data

generated in this study have been deposited in the Zenodo database at. The processed spatial

autocorrelation data are available in the same location. Past, current, and future climate estimates

were acquired from the WorldClim 2.1 [https://www.worldclim.org/data/index.html] and CRU

TS4.03 [https://crudata.uea.ac.uk/cru/data/hrg/] datasets.

Code availability: Examples of code used in this analysis is freely available at.

References:
1.   Tuanmu, M. N. & Jetz, W. A global 1-km consensus land-cover product for biodiversity
and ecosystem modelling. *Global Ecology and Biogeography* **23**, 1031–1045 (2014).
2.   Piao, S. *et al.* Plant phenology and global climate change: Current progresses and
challenges. *Global Change Biology* **25**, 1922–1940 (2019).
3.   Ge, Q., Wang, H., Rutishauser, T. & Dai, J. Phenological response to climate change in
China: A meta-analysis. *Global Change Biology* **21**, 265–274 (2015).
4.   Menzel, A. *et al.* European phenological response to climate change matches the
warming pattern. *Global Change Biology* **12**, 1969–1976 (2006).
5.   Miller, T. K., Heberling, J. M., Kuebbing, S. E. & Primack, R. B. Warming temperatures
are linked to phenological mismatch among native and non-native trees, shrubs, and wildflowers
across the Eastern Deciduous Forest of North America. *Journal of Ecology* (2022)
doi:10.1111/1365-2745.14021.
6.   Zohner, C. M. & Renner, S. S. Common garden comparison of the leaf-out phenology of
woody species from different native climates, combined with herbarium records, forecasts long-
term change. *Ecology Letters* **17**, 1016–1025 (2014).
7.   Heberling, J. M., McDonough MacKenzie, C., Fridley, J. D., Kalisz, S. & Primack, R. B.
Phenological mismatch with trees reduces wildflower carbon budgets. *Ecology Letters* **22**, 616–
623 (2019).
8.   Wang, H. *et al.* Impacts of global warming on phenology of spring leaf unfolding remain
stable in the long run. *International Journal of Biometeorology* **61**, 287–292 (2017).
9.   Willems, F. M., Scheepens, J. F. & Bossdorf, O. Forest wildflowers bloom earlier as
Europe warms: lessons from herbaria and spatial modelling. *New Phytologist* **235**, 52–65 (2022).

10.   Routhier, M. C. & Lapointe, L. Impact of tree leaf phenology on growth rates and
reproduction in the spring flowering species Trillium erectum (Liliaceae). *American Journal of*
Botany **89**, 500–505 (2002).
11.   Neufeld, H. S. & Young, D. R. Ecophysiology of the Herbaceous Layer in Temperate
Deciduous Forests. in *The Herbaceous Layer in Forests of Eastern North America* (ed. Gilliam,
F. S.) 35–95 (Oxford University Press, 2014).
doi:10.1093/acprof:osobl/9780199837656.003.0003.
12.   Lee, B. R. & Ibáñez, I. Spring phenological escape is critical for the survival of temperate
tree seedlings. *Functional Ecology* 1365-2435.13821 (2021) doi:10.1111/1365-2435.13821.
13.   Lee, B. R. & Ibáñez, I. Improved phenological escape can help temperate tree seedlings
maintain demographic performance under climate change conditions. *Global Change Biology*
gcb.15678 (2021) doi:10.1111/gcb.15678.
14.   Kwit, M. C., Rigg, L. S. & Goldblum, D. Sugar maple seedling carbon assimilation at the
northern limit of its range: the importance of seasonal light. *Canadian Journal of Forest*
Research **40**, 385–393 (2010).
15.   Augspurger, C. K. & Bartlett, E. A. Differences in leaf phenology between juvenile and
adult trees in a temperate deciduous forest. *Tree physiology* **23**, 517–525 (2003).
16.   Heberling, J. M., Cassidy, S. T., Fridley, J. D. & Kalisz, S. Carbon gain phenologies of
spring-flowering perennials in a deciduous forest indicate a novel niche for a widespread
invader. *New Phytologist* **221**, 778–788 (2019).
17.   Jacques, M. H. *et al.* Responses of two understory herbs, Maianthemum canadense and
Eurybia macrophylla, to experimental forest warming: Early emergence is the key to enhanced
reproductive output. *American Journal of Botany* **102**, 1610–1624 (2015).
18.   Augspurger, C. K. Early spring leaf out enhances growth and survival of saplings in a
temperate deciduous forest. *Oecologia* **156**, 281–286 (2008).
19.   Seiwa, K. Advantages of early germination for growth and survival of seedlings of Acer
mono under different overstorey phenologies in deciduous broad-leaved forests. *Journal of*
Ecology **86**, 219–228 (1998).
20.   Augspurger, C. K. & Salk, C. F. Constraints of cold and shade on the phenology of spring
ephemeral herb species. *Journal of Ecology* **105**, 246–254 (2017).
21.   Ida, T. Y. & Kudo, G. Timing of canopy closure influences carbon translocation and seed
production of an understorey herb, Trillium apetalon (Trilliaceae). *Annals of Botany* **101**, 435–
446 (2008).
22.   Lapointe, L. Fruit development in trillium: Dependence on stem carbohydrate reserves.
Plant Physiology **117**, 183–188 (1998).
23.   Zohner, C. M. & Renner, S. S. Innately shorter vegetation periods in North American
species explain native-non-native phenological asymmetries. *Nature Ecology and Evolution* **1**,
1655–1660 (2017).
24.   Cleland, E. E. *et al.* Phenological tracking enables positive species responses to climate
change. *Ecology* **93**, 1765–1771 (2012).
25.   Everill, P. H., Primack, R. B., Ellwood, E. R. & Melaas, E. K. Determining past leaf-out
times of New England's deciduous forests from herbarium specimens. *American Journal of*
Botany **101**, 1293–1300 (2014).
26.   Primack, D., Imbres, C., Primack, R. B., Miller-Rushing, A. J. & Del Tredici, P.
Herbarium specimens demonstrate earlier flowering times in response to warming in Boston.
American Journal of Botany **91**, 1260–1264 (2004).

27.   Ibáñez, I. *et al.* Forecasting phenology under global warming. *Philosophical Transactions*
of the Royal Society B: Biological Sciences **365**, 3247–3260 (2010).
28.   Vitasse, Y. *et al.* Leaf phenology sensitivity to temperature in European trees: Do within-
species populations exhibit similar responses? *Agricultural and Forest Meteorology* **149**, 735–
744 (2009).
29.   Miller, T. K., Gallinat, A. S., Smith, L. C. & Primack, R. B. Comparing fruiting
phenology across two historical datasets: Thoreau's observations and herbarium specimens.
Annals of Botany **128**, 159–170 (2021).
30.   Heberling, J. M. Herbaria as Big Data Sources of Plant Traits. *International Journal of*
Plant Sciences **183**, 87–118 (2022).
31.   Davis, C. C., Willis, C. G., Connolly, B., Kelly, C. & Ellison, A. M. Herbarium records
are reliable sources of phenological change driven by climate and provide novel insights into
species' phenological cueing mechanisms. *American Journal of Botany* **102**, 1599–1609 (2015).
32.   Ramirez-Parada, T. H., Park, I. W. & Mazer, S. J. Herbarium specimens provide reliable
estimates of phenological responses to climate at unparalleled taxonomic and spatiotemporal
scales. *Ecography* **2022**, (2022).
33.   Zohner, C. M., Benito, B. M., Fridley, J. D., Svenning, J. C. & Renner, S. S. Spring
predictability explains different leaf-out strategies in the woody floras of North America, Europe
and East Asia. *Ecology Letters* **20**, 452–460 (2017).
34.   Boucher, O. *et al.* Presentation and Evaluation of the IPSL-CM6A-LR Climate Model.
Journal of Advances in Modeling Earth Systems **12**, 1–52 (2020).
35.   Eyring, V. *et al.* Overview of the Coupled Model Intercomparison Project Phase 6
(CMIP6) experimental design and organization. *Geoscientific Model Development* **9**, 1937–1958
(2016).
36.   Gilliam, F. S. The ecological significance of the herbaceous layer in temperate forest
ecosystems. *BioScience* **57**, 845–858 (2007).
37.   Landuyt, D. *et al.* The functional role of temperate forest understorey vegetation in a
changing world. *Global Change Biology* **25**, 3625–3641 (2019).
38.   Frerker, K., Sabo, A. & Waller, D. Long-term regional shifts in plant community
composition are largely explained by local deer impact experiments. *PLoS ONE* **9**, 1–17 (2014).
39.   Knight, T. M., Dunn, J. L., Smith, L. A., Davis, J. & Kalisz, S. Deer facilitate invasive
plant success in a Pennsylvania forest understory. *Natural Areas Journal* **29**, 110–116 (2009).
40.   Knight, T. M. The effects of herbivory and pollen limitation on a declining population of
Trillium grandiflorum. *Ecological Applications* **14**, 915–928 (2004).
41.   Dullinger, S. *et al.* Modelling the effect of habitat fragmentation on climate-driven
migration of European forest understorey plants. *Diversity and Distributions* **21**, 1375–1387
(2015).
42.   Dicks, L. V. *et al.* A global-scale expert assessment of drivers and risks associated with
pollinator decline. *Nature Ecology and Evolution* **5**, 1453–1461 (2021).
43.   Kudo, G. & Ida, T. Y. Early onset of spring increases the phenological mismatch between
plants and pollinators. *Ecology* **94**, 2311–2320 (2013).
44.   Hegland, S. J., Nielsen, A., Lázaro, A., Bjerknes, A. L. & Totland, Ø. How does climate
warming affect plant-pollinator interactions? *Ecology Letters* **12**, 184–195 (2009).
45.   McKinney, A. M. & Goodell, K. Shading by invasive shrub reduces seed production and
pollinator services in a native herb. *Biological Invasions* **12**, 2751–2763 (2010).

46.   Gorchov, D. L. *et al.* Differential and interacting impacts of invasive plants and white-
tailed deer in eastern U.S. forests. *Biological Invasions* **23**, 2711–2727 (2021).

47.   Bobbink, R. *et al.* Global assessment of nitrogen deposition effects on terrestrial plant
diversity: A synthesis. *Ecological Applications* **20**, 30–59 (2010).

48.   Small, C. J. & McCarthy, B. C. Relationship of understory diversity to soil nitrogen,
topographic variation, and stand age in an eastern oak forest, USA. *Forest Ecology and*
Management **217**, 229–243 (2005).

49.   Miller, C. N., Papeş, M., Schilling, E. E. & Kwit, C. Reproductive traits explain
occupancy of predicted distributions in a genus of eastern North American understory herbs.
Diversity and Distributions ddi.13297 (2021) doi:10.1111/ddi.13297.

50.   Whigham, D. F. Ecology of woodland herbs in temperate deciduous forests. *Annual*
Review of Ecology, Evolution, and Systematics **35**, 583–621 (2004).

51.   Lembrechts, JJ, Aalto, J, Ashcroft, MB, et al. SoilTemp: A global database of near-
surface temperature. *Glob Change Biology* **26**, 6616– 6629 (2020).

52.   Wolkovich, E. M., Chamberlain, C. J., Buonaiuto, D. M., Ettinger, A. K. & Morales-
Castilla, I. Integrating experiments to predict interactive cue effects on spring phenology with
warming. *New Phytologist* **235**, 1719–1728 (2022).

53.   Chuine, I. & Régnière, J. Process-Based Models of Phenology for Plants and Animals.
Annu. Rev. Ecol. Evol. Syst. **48**, 159–182 (2017).

54.   Heberling, J. M., Miller, J. T., Noesgaard, D., Weingart, S. B. & Schigel, D. Data
integration enables global biodiversity synthesis. *Proceedings of the National Academy of*
Sciences of the United States of America **118**, 1–7 (2021).

55.   Miller-Rushing, A. J. & Primack, R. B. Global warming and flowering times in
Thoreau's Concord: A community perspective. *Ecology* **89**, 332–341 (2008).

56.   Buonaiuto, D. M. & Wolkovich, E. M. Differences between flower and leaf phenological
responses to environmental variation drive shifts in spring phenological sequences of temperate
woody plants. *Journal of Ecology* **109**, 2922–2933 (2021).

57.   Harris, I., Osborn, T. J., Jones, P. & Lister, D. Version 4 of the CRU TS monthly high-
resolution gridded multivariate climate dataset. *Scientific Data* **7**, 1–18 (2020).

58.   Lee, Benjamin R. Data and model code for article 'Wildflower phenological escape
differs by continent and spring temperature'. (2022) doi:10.5281/ZENODO.7080193.

59.   Roberts, A. M. I., Tansey, C., Smithers, R. J. & Phillimore, A. B. Predicting a change in
the order of spring phenology in temperate forests. *Global Change Biology* **21**, 2603–2611
(2015).

60.   Su, Y.-S. & Yajima, M. Package 'R2jags'. (2015).

61.   Ettinger, A. K. *et al.* Winter temperatures predominate in spring phenological responses
to warming. *Nature Climate Change* **10**, 1137–1142 (2020).

62.   Winkler, D. E., Butz, R. J., Germino, M. J., Reinhardt, K. & Kueppers, L. M. Snowmelt
timing regulates community composition, phenology, and physiological performance of alpine
plants. *Frontiers in Plant Science* **9**, 1–13 (2018).

63.   Bachl, F. E., Lindgren, F., Borchers, D. L. & Illian, J. B. inlabru: an R package for
Bayesian spatial modelling from ecological survey data. *Methods in Ecology and Evolution* **10**,
760–766 (2019).

64.   Krainski, E. T. *et al. Advanced spatial modeling with stochastic partial differential*
equations using R and INLA. (Chapman & Hall/CRC Press, 2019).

65.   De Smedt, T., Simons, K., Van Nieuwenhuyse, A. & Molenberghs, G. Comparing
MCMC and INLA for disease mapping with Bayesian hierarchical models. *Archives of Public*
Health **73**, O2 (2015).
66.   Rue, H., Martino, S. & Chopin, N. Approximate Bayesian inference for latent Gaussian
models by using integrated nested Laplace approximations. *Journal of the Royal Statistical*
Society. Series B: Statistical Methodology **71**, 319–392 (2009).
67.   Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces
for global land areas. *International Journal of Climatology* **37**, 4302–4315 (2017).
68.   Boucher, O. *et al.* IPSL IPSL-CM6A-LR-INCA model output prepared for CMIP6
AerChemMIP. Downloaded 20210914. (2020) doi:10.22033/ESGF/CMIP6.13581.
69.   Pierce, D. ncdf4: Interface to Unidata netCDF (Version 4 or Earlier) Format Data Files.
(2019).
70.   Pebesma, E. stars: Spatiotemporal Arrays, Raster and Vector Data Cubes. (2021).
71.   Bivand, R. S., Pebesma, E. & Gomez-Rubio, V. *Applied spatial data analysis with R*.
(Springer, 2013).

663
664
<u>Acknowledgements</u>

We thank A. Helmling, L. de Castro Machado de Souza, P.Y. Lai, C. Terry, and M. Rothendler

(North American specimens); I. Vogler, M. Schramm, V. Bastian, K. Adler, and B. Müller

(European specimens); and Z. Yang, C. Tan, L. Wang, and X. Huo (East Asian specimens) for

their efforts in databasing and georeferencing the herbarium specimens analyzed for this project.

We also thank C. Römermann, E. Welk, and B. Liu for their suggestions on the European and

Asian species list. We are grateful to the Chinese Virtual Herbarium and Chinese Field

Herbarium for providing the East Asian specimen data and digitized photos and to the

Metasequoia funding of Nanjing Forestry University [YY] which helped fund the Asian data

extraction. We thank F Lindgren and F Willems for providing guidance on INLA methodology

for the models with spatial autocorrelation. BRL, TKM, JMH, SEK, and RBP were funded by

National Science Foundation Grants Nos. DEB 1936971, 1936877, and 1936960. We

acknowledge the World Climate Research Program, which, through its Working Group on

Coupled Modelling, coordinated and promoted CMIP6. We thank the climate modeling groups

for producing and making available their model output, the Earth System Grid Federation

(ESGF) for archiving the data and providing access, and the multiple funding agencies who

support CMIP6 and ESGF. We thank Abe Miller-Rushing and Amanda S. Gallinat for providing

valuable feedback on this manuscript.

Author contributions statement: JMH, SEK, and RBP initially conceived of this project. TKM,

CR, and YY collected, scored, and georeferenced the herbarium collections used in this analysis.

BRL conducted the statistical analysis and wrote the article. All authors contributed equally to

reviewing and editing the manuscript.

Competing interests statement: The authors declare no competing interests.

Additional information: Supplementary Information is available for this paper. Correspondence

and requests for materials should be addressed to Benjamin R. Lee (LeeB@CarnegieMNH.org).

Reprints and permissions information is available at www.nature.com/reprints.

 Figures Legends/Captions:

 Figure 1: Distributions of observed understory wildflower First Flowering Date (FFD; blue) and

 canopy tree Leaf out Date (LOD; gold) in temperate forests in Asia, Europe, and North America

 by (a) observed date of phenology and (b) average spring (March-April) temperature (°C). Grey

 shading behind the violin plots in panel a indicates the March-April period used to model and

 forecast phenology.

<u>Figure 2</u>: Posterior estimated means and 95% credible intervals for spring temperature

sensitivity. Shapes represent parameter estimates for wildflower First Flower Date (FFD, blue

circles; n = 1418, 618, and 1060 for Asia, Europe, and North America, respectively) and canopy

tree Leaf Out Date (LOD, yellow triangles; n = 899, 532, and 995, for Asia, Europe, and North

America, respectively). Estimates are considered different from 0 if credible intervals do not

overlap the dashed 0 line and are considered different from each other if credible intervals do not

overlap.

<u>Figure 3</u>: Estimated mean difference between wildflower First Flower Date (FFD) and canopy

tree Leaf Out Date (LOD) (in days) under current climate conditions (averaged from 2009-2018,

see methods) in a) Asia, b) Europe, and c) North America. Negative values indicate tree LOD is

estimated to occur before wildflower FFD. Estimations were cropped by estimated area of

broadleaf and mixed-broadleaf forest (see methods). Dark gray regions indicate areas where the

consensus land classification is < 1% deciduous or mixed deciduous forest cover. An uncropped

version of this figure is available in Fig. S11.

<u>Figure 4:</u> Projected changes in spring light window duration (ΔWindow) between current

climate conditions and conditions projected for the end of the current century. Positive values

indicate regions where light window duration is expected to increase with spring warming

(particularly in northern Asia) and negative values indicate where it is expected to decrease

(particularly in northern North America). Dark gray regions indicate areas where the consensus

land classification is < 1% deciduous or mixed deciduous forest cover. An uncropped version of

this figure is available in Fig. S12.

729