Small Molecule Assembly Modulators with Pan-Cancer Therapeutic Efficacy

Authors: Anuradha F. Lingappa[#1], Olayemi Akintunde[#1], Connie Ewald[1], Markus Froehlich[1], Niloufar

Ziari[1], Shao Feng Yu[1], Maya Michon[1], Suguna Mallesh[1], Jim Lin[1], Anatoliy Kitaygorodskyy[1], Dennis Solas[1],

Jonathan C. Reed[2], Jaisri R. Lingappa[2] Andreas Mueller-Schiffmann[3], Carsten Korth[3], Dharma Prasad[1],

Aysegul Nalca[4], Emily Ashton[5], Brad Fabbri[5], Emma Petrouski[1], Debendranath Dey[1], David Andrews[6], and

Vishwanath R. Lingappa[1,7]

Affiliations:

[1]Prosetta Biosciences, San Francisco, CA USA

[2] Dept. of Global Health, University of Washington, Seattle, WA, USA.

[3]Institute of Neuropathology, Heinrich Heine University, Dusseldorf, Germany

[4]United States Army Medical Research Institute for Infectious Diseases, Fredrick MD, USA

[5]TechAccel, Overland Park, KS, USA

[6]Sunnybrook Research Institute, Toronto, ON, Canada

[7]University of California, San Francisco, CA, USA

[#]Co-first authors who contributed equally to this work

*To whom correspondence should be addressed vlingappa@prosetta.com

Abstract:

Two structurally-unrelated small molecule chemotypes, represented by compounds PAV-617 and PAV-951, with antiviral activity in cell culture against monkeypox virus (MPXV) and human immunodeficiency virus (HIV) respectively, were studied for anti-cancer efficacy. Each exhibited apparent pan-cancer cytotoxicity, reasonable pharmacokinetics, and non-toxicity in mice at active concentrations. Anti-tumor properties of each compound were validated in mouse xenografts against A549 human lung cancer. The targets of these compounds are unconventional: each binds to a different transient, energy-dependent multi-protein complex containing the protein KAP-1(TRIM28), an allosteric modulator known to broadly regulate mechanisms underlying viral and nonviral disease states including cancer. Treatment with these compounds alters the target multi-protein complexes in a manner consistent with allosteric modulation as their mechanism of action. These compounds appear to remove a block, crucial for cancer survival and progression, on the homeostatic linkage of uncontrolled cellular proliferation to apoptosis. These compounds may provide starting points for development of next-generation non-toxic, cancer therapeutics.

Introduction

The similarity of the interactions of viral infection and cancer with the healthy host has often

been noted (Javier and Butel, 2008). Both represent pathological processes of extremely diverse origin,

that overcome the complex feedback controls of homeostasis, to the detriment of the host (Morales-

Sánchez and Fuentes-Pananá, 2014). Both exploit natural selection as a powerful weapon to overcome

host defenses. Viruses have done so over deep evolutionary time through co-evolution with their hosts

and through the emergence of resistance mutation in response to the selective pressure of treatments

(Ison, 2011; Kaján et al., 2020). Cancers regularly use the latter mechanism, resulting in clonal mutants

that drive cancer progression (Usman et al., 2021).

At least seven different viruses— Epstein-Barr virus (EBV), hepatitis B virus (HBV), hepatitis C

virus (HCV), human T-lymphotropic virus 1 (HTLV-1), human papillomavirus (HPV), Kaposi sarcoma-

associated herpesvirus (KSHV or HHV-8), and Merkel cell polyomavirus (MCPyV)— are known to be

directly oncogenic through their alteration of the cellular environment and/or impairment of the host's

innate immune system defenses (Cirone, 2018; Krump and You, 2018). It has therefore been proposed

that the study of viruses could play a key role in the discovery of new cancer treatments by identifying

cellular targets that drive tumorigenesis (O'Shea, 2005).

The incompleteness in our current understanding of the dynamics of host homeostasis and its

myriad of feedback controls has been a disadvantage for efforts to design novel therapeutic

countermeasures against both viruses and cancer. However, a recent unconventional approach to

antiviral drug discovery has identified host allosteric sites essential for homeostasis that are repurposed

by viral infection (Müller-Schiffmann et al., 2022). The antiviral compounds identified by this approach

appear to restore key features essential to host homeostasis (Müller-Schiffmann et al., 2022). We

wished to determine whether those compounds might have therapeutic applicability against cancer,

given the analogy in how both viruses and cancers drive departure from homeostasis. The results to be

presented here suggest this is the case and shed light on molecular pathways relevant for both viral and

neoplastic disease, providing a strategy for development of novel cancer therapeutics.

Utilizing a cell-free protein synthesis and assembly (CFPSA) system, a library of approximately

150,000 drug-like small molecules was screened for compounds which blocked the assembly of viral

capsids without inhibiting host protein synthesis (Broce et al., 2016; U. F. Lingappa et al., 2013; Müller-

Schiffmann et al., 2022; Reed et al., 2021). The hit compounds identified have been termed "*assembly*

modulators" (Müller-Schiffmann et al., 2022, 2021a). A collection of 300 structurally-diverse assembly

modulators were assembled from hits demonstrating activity against capsid assembly in one or more

viral family (V. Lingappa et al., 2013). Antiviral assembly modulators from this collection have been

validated against live virus in cell culture for multiple viral families including *Retroviridae*, *Rhabdoviridae*,

Poxviridae, *Adenoviridae*, *Herpesviridae*, *Paramyxoviridae*, *Coronaviridae*, *Orthomyxoviradae*, and

Picornaviridae (U. F. Lingappa et al., 2013; Müller-Schiffmann et al., 2022, 2021b; Priyamvada et al.,

2021; Reed et al., 2021). In two cases, *Coronaviridae* (porcine epidemic diarrhea virus) and

Paramyxoviridae (respiratory syncytial virus), cellular antiviral activity has been confirmed in animal

disease models (Müller-Schiffmann et al., 2022).

These antiviral assembly modulators appear to target host-viral protein-protein interactions via

allosteric sites that control repurposing of host assembly machinery for viral capsid formation and which

also allow disengagement of host innate immune defenses such as autophagy (Müller-Schiffmann et al.,

2022). In one study, a class of assembly modulators was shown to change the composition of a

transient, energy-dependent multi-protein complex whose components include p62/SQSTM1, a key

regulator of autophagy (Müller-Schiffmann et al., 2022). This multi-protein complex is co-opted upon

viral infection to lose p62/SQSTM1 and to gain the viral nucleoprotein (Müller-Schiffmann et al., 2022).

Upon treatment with an antiviral assembly modulator, the multi-protein complex is restored to its

normal composition, with loss of the viral nucleoprotein component and re-engagement of

p62/SQSTM1 (Müller-Schiffmann et al., 2022).

The discovery of dynamic multi-protein complexes whose composition changes with drug

treatment offers a new means of parsing out post-translational protein heterogeneity and its relevance

for diseased states. The amount of a given protein that has been observed in the multi-protein

complexes targeted by antiviral assembly modulators comprises only a very small fraction of the total

amount of the component proteins present in a cell (Müller-Schiffmann et al., 2022). The role played by

the small subset of the particular proteins that are part of the particular complex may be related to

"moonlighting" functions observed for a growing set of cellular and viral proteins (Copley, 2012; Jeffery,

2019, 2018).

We hypothesized that if an overlap between viral and oncogenic pathways exists, some antiviral

assembly modulators might be capable of disrupting multi-protein complexes associated with the

hallmarks of cancer (Hanahan and Weinberg, 2011, 2000). To test the hypothesis, we established a

cancer-relevant counterscreen and applied it to previously-identified assembly modulating compounds.

In this paper, we describe two assembly modulators which were originally characterized for their

antiviral properties but are now shown to have potential as cancer therapeutics based on *in vivo*

screening and *in vitro* validation studies.

Results

<u>Uncontrolled cellular proliferation: a hallmark of cancer inhibited by assembly modulators PAV-617 and</u>

<u>PAV-951</u>

No hallmark of cancer is more fundamental than uncontrolled proliferation (Hanahan and

Weinberg, 2000). Uncontrolled proliferation normally triggers cell death mechanisms, including

apoptosis (Pucci et al., 2000). Therefore, to survive, a cancer must achieve a means of evading cell death

long enough to complete cell division and reset the cell death timer. This, in turn, allows further

proliferation, during which time additional mutations can occur and selection pressure will drive higher

and higher grade malignancy and, ultimately, metastasis (Pfeffer and Singh, 2018). Compounds capable

of arresting proliferation, either directly or indirectly, could make potent anti-cancer agents because the

delay in cancer progression would provide an opportunity for a patient's innate immune system and

other homeostatic mechanisms to re-establish themselves.

Abnormal signaling pathways triggered by aberrant protein-protein interactions is one way that

neoplastic cells are able to achieve uncontrolled proliferation (Pfeffer and Singh, 2018). In order to

characterize whether assembly modulators could arrest the proliferation of neoplastic cells by

redirecting key protein-protein interactions, we first sought to identify a cell line in which endogenous

apoptosis was substantially lacking. While a successful anti-cancer compound would likely exhibit both

anti-proliferative and cytotoxic efficacy, we wanted to conduct our screen under conditions where a

readout measuring the arrest of proliferation would not be obscured by downstream activation of the

normal cascade of events comprising cell death pathways.

We assessed caspase-3/7 activity in multiple tumor cell lines with an Apo-ONE assay (see **Fig.**

1A). The expected correlation between endogenous triggers of apoptotic cell death and cancer

progression was demonstrated in the LNCaP prostate cancer progression cell model, where LNCaP-C33

early (hormone sensitive) cancer cells displayed substantially more markers of apoptosis than LNCaP-

C81 late (hormone resistant) cancer cells (Igawa et al., 2002). In the Apo-ONE assay, CHO K1 cells show

very little endogenous apoptosis (see **Fig. 1A**). Hennes-20, a CHO K1 derivative into which the human

APP751 V717F mutation has been stably transfected, show even less apoptosis than their parental line

(see **Fig. 1A**).

We then counterscreened our collection of anti-viral assembly modulator in Hennes-20 cells

plated at low (500 cells/well) versus high (15,000 cells/well) densities and treated with DMSO (vehicle)

or dose-titration of compounds. The rationale for this screen is that an intrinsically toxic compound

should kill cells regardless of density, including in Hennes-20 cells. A compound that selectively triggers

the arrest of proliferation will appear cytotoxic due to inhibition of cell growth when plated at a low

density. However, it will appear non-toxic to cells plated at a high density where the cells are

approaching confluence and the readout detected by a cell viability assay is already close to the

maximum.

Two structurally-unrelated small molecules (Tanimoto similarity score of 41%), PAV-617 and

PAV-951, displayed the desired phenotype (see **Figs. 1B** and **1C**). Our findings in Hennes-20 cells at low

versus high density comparison suggested that the effect of these compounds in cells lacking

endogenous apoptosis was due to inhibition of proliferation.

To confirm that the drug-induced inhibition observed in low density Hennes 20 cells resulted

from temporary arrest of proliferation and not cell death, compound was removed after a period of

treatment (24 hour for PAV-617 and 6 hours for PAV-951), and cell growth was measured over the

subsequent two weeks for recovery potential. The cells treated with PAV-617 and PAV-951 initially

showed reduced cell density relative to the control, but once compound was removed, growth was

restored over time (See **Figs. 1D** and **1E**). By day 11, cell density in compound-treated cells caught up to

the DMSO-treated cells (See **Figs. 1D** and **1E**).

Since Hennes 20 cells do not appear to have endogenous apoptosis, we sought to use a different

cancer cell line to assess whether that inhibition of proliferation would be accompanied by cell death.

The recovery experiment was conducted in LNCaP C-33 cells which appear to have substantial

endogenous apoptosis (see **Fig. 1A**), and treatment with PAV-617 and PAV-951 did inhibit cancer cell

growth relative to the DMSO-treated control but the cells did not recover or grow significantly once

compound was removed (See **Figs. 1F** and **1G**).

Figure 1. Assay development and activity of hit compounds PAV-617 and PAV-951. Fig. 1A shows

assessment of the endogenous apoptosis response in multiple cell lines. Plates were seeded with LNCaP

C-33, LNCaP C-81, CHO K-1, and Hennes 20 cells. After three days of growth Apo-ONE reagent was added

and caspase-3/7 activity was determined by fluorescent readout. Averages and standard deviation of

observed activity in triplicate-repeated samples were calculated and graphed in Microsoft Excel. The

Hennes 20 cell line was chosen for the counterscreen based on its low levels of caspase activity. **Figs. 1B**

and 1C show activity of PAV-617 and PAV-951 in the assay, where parallel plates of Hennes 20 cells were

seeded at a high density of 15,000 cells per well and a low density of 500 cells per well and treated with

DMSO, PAV-617, or PAV-951 in triplicate-repeated dose titrations. Fluorescent reading (RFU)

corresponding to cell viability was calculated using an alamarBlue[TM] assay and the averages and standard

deviations for triplicate-repeated samples were calculated and graphed on Microsoft Excel as a

percentage of the DMSO-treated cells. PAV-617 exhibited dose-dependent inhibition of cell growth in the

low density plate, indicating an EC50 between 0.1uM and 0.5uM. PAV-617 exhibited some toxicity to cells

at the higher doses tested, indicating a CC50 slightly greater than 0.5uM. PAV-951 exhibited dose-

dependent inhibition of cell growth in the low-density plate, indicating an EC50 around 0.5uM. PAV-951

exhibited no significant inhibition of cell growth at the tested doses in the high-density plate, indicating a

CC50 greater than 0.5uM. **Figs. 1D- 1G** show recovery of cancer cell growth following removal of

compound. Hennes 20 or LNCaP C-33 cells were seeded at a low density then incubated with DMSO, PAV-

617, or PAV-951. After a period of treatment, the medium containing compound was removed and

replaced with fresh media. Plates were assessed for cell viability by alamarBlue[TM] on days 5, 8, 11, and 14

and the averages and standard deviations of triplicate-repeat samples were calculated and graphed over

time on Microsoft Excel. PAV-617 and PAV-951 treated Hennes 20 and LNCaP C-33 cells all showed

reduced viability compared to matched DMSO-treated cells on day 5. However, cell growth in the Hennes

20 cells which had been treated with compound recovered with time, while the LNCaP C-33 cells which

had been treated with compound did not recover.

<u>Investigating the activities of PAV-617 and PAV-951: from modulators of viral capsid assembly to pan-</u>

<u>cancer therapeutics</u>

The anti-proliferative compounds PAV-617 and PAV-951 had originally emerged from our CFPSA

screen as inhibitors of viral capsid formation. The CFPSA model had been validated by demonstrating

that hit antiviral assembly modulators display activity against infectious viruses in cell culture (U. F.

Lingappa et al., 2013; Müller-Schiffmann et al., 2022; Priyamvada et al., 2021; Reed et al., 2021). PAV-

617 is active against pox viruses in cell culture (see **Fig. 2A** for MPXV activity and Priyamvada et al., 2021

for chickenpox activity). The effective concentration for half maximal activity (EC50) of PAV-617 against

MPXV is approximately 300 nM (See **Fig. 2A**). PAV-951 is active against HIV in cell culture with an EC50

between 1 uM and 300 nM (see **Fig. 2B**).

Figure 2. Anti-viral and pharmacokinetic properties of PAV-617 and PAV-951. Fig. 2A shows activity of

PAV-617 against MPXV. BSC-40 cells were infected with 100 plaque forming units of MPXV Zaire 79

andtreated with PAV-617 for three days. Averages and standard deviation for plaques observed with

triplicate-repeated dose-titration of PAV-617 are shown as a percentage of the plaques observed in

untreated cells with an EC50 of approximately 300 nM. **Fig. 2B** shows activity of PAV-951 against HIV. MT-

2 cells were infected with NL4-3 Rluc HIV and treated with PAV-951 for four days. Averages and standard

deviation of viral titer observed with triplicate-repeated dose-titration of PAV-951 are shown as a

percentage of the titer observed in DMSO-treated cells, with an EC50 between 0.37 uM and 1.1 uM.

When PAV-617 and PAV-951 were identified as having anti-cancer activity in addition to anti-

viral properties, we suspected that the compounds might correct cancer-induced defects in protein

assembly that were related to the aberrant assemblies induced by viruses. To get a better

understanding of how the defects present themselves across diverse cancers, we assessed the anti-

cancer activity of these two compound chemotypes on a panel of 15 cancer cell lines from the Eurofins

OncoPanel™ ("OncoPanelTM Cell-Based Profiling Service Details," n.d.). The cell lines were derived from

a variety of tissues and were representative of male and female patients of different ages from pediatric

to senior. Both PAV-617 and PAV-951 showed pan-cancer activity with dose-dependent tumor growth

inhibition in all 15 cell lines (see **Figs. 3A** and **3B**).

Figure 3. Pan-cancer activity of PAV-617 and PAV-951. Figs. 3A and **3B** show results from a panel of

human tumor cell lines— A172 (male human glioma), BFTC-905 (female human urinary bladder

transitional cell carcinoma), COR-L105 (male human lung adenocarcinoma), DB (male human b-cell

lymphoma), FaDu (male human pharynx squamous cell carcinoma), H9 (male human t-cell lymphoma), Hs

294T (male human melanoma), MCF7 (female human breast cancer), MDA MB 436 (female human breast

cancer), MeWo (male human melanoma), MHH-PREB-1 (male human b-cell lymphoma), SJSA1-OSA (male

human osteosarcoma), SU-DHL-10 (male human b-cell lymphoma), SW1353 (female human

chondrosarcoma), and U-2 OS (female human osteosarcoma). Cells were grown for 24 hours then treated

with either vehicle, PAV-617, or PAV-951. Cell viability after 3 days of treatment was measured as

bioluminescence intensity and averages of triplicate-repeat dose-titrations with PAV-617 and PAV-951

were graphed on Microsoft Excel as percent of bioluminescence observed in DMSO-treated cells. Both

compounds showed inhibitory effects in all 15 cell lines.

<u>Animal validation of PAV-617 and PAV-951 anti-cancer efficacy</u>

With demonstrated anti-viral activity, demonstrated anti-cancer activity, and data supporting a

proliferation-based mechanism of action, we assessed mouse toxicology and pharmacokinetic (PK)

properties in order to determine suitability for efficacy studies in an animal model.

The maximum tolerated dose (MTD) estimates how much compound can be administered to an

animal without adverse effects (Gad, 2014). When administered orally to mice, both PAV-617 and PAV-

951 displayed MTDs greater than 20 mg/kg, which was the highest dose tested, as no clinical symptoms

or significant differences were observed between vehicle and treatment groups. When administered by

intraperitoneal (IP) injection, the MTD of PAV-617 was determined to be greater than 10 mg/kg, which

was the highest dose tested. When administered by IP injection, the MTD of PAV-951 was determined to

be 2.5 mg/kg because at the next-highest dose (5 mg/kg), one animal displayed dullness, a clinical sign

of toxicity, for 24 hours. The animal subsequently recovered and behaved normally for the remaining

48-hour observation period and no abnormality was observed at necropsy for any animals in the PAV-

951 5 mg/kg or 10 mg/kg treatment groups.

The PK properties are based on the absorption, distribution, metabolism, and excretion of a

compound in a living organism and are necessary to determine dosing parameters because any

compound designed for clinical use needs to achieve an efficacious concentration in a target organ

(Hughes et al., 2011; Reichel and Lienau, 2015). As an early measure of PK, we determined the

concentration of compound in the plasma of rats or mice over time following one intravenous (IV), one

IP dose, or one oral dose. Both compounds were detectable in the animals through all administration

routes, though the maximum concentration achieved (Cmax) and rate of elimination were variable

across conditions (see **Supplemental Fig. 1**).

We determined that, while both chemical series would need optimization on the PK and

toxicologic properties before being named as clinical drug-candidates, PAV-617 and PAV-951 would be

adequate for a preliminary animal efficacy study in order to validate whether or not the anti-

proliferative properties of the compounds observed in cell culture translates to anti-proliferative

properties in animals.

In the animal efficacy study, human A459 non-small cell lung cancer cells were grafted

subcutaneously onto mice. After 30 days of tumor establishment, the animals received daily treatment

with PAV-617 or PAV-951 and tumor volume was measured over time. The doses and routes of

administration for PAV-617 (10 mg/kg IP injection) and PAV-951 (1.5 mg/kg IV injection) were

determined based on their MTD and PK properties (see **Supplemental Fig. 1**). The PAV-617 study was

conducted for 28 days and the PAV-951 study was conducted for 14 days. As negative and positive

controls, both studies included a group treated with vehicle only and a group treated with Gemcitabine

Hydrochloride, an FDA-approved drug for non-small cell lung cancer ("Reference ID" 4433223-

Accessdata.fda.gov," n.d.). The Gemcitabine was administered at a dose of 100 mg/kg twice weekly.

Both PAV-617 and PAV-951 reduced tumor growth significantly compared to the vehicle-only groups

and performed comparably to Gemcitabine despite being administered at substantially lower doses (See

Figs. 4A and **4B**).

Figure 4) Anti-proliferative activities of PAV-617 and PAV-951 in animal xenograft model. Female nude

mice were injected with 0.1 mL a suspension containing 1×10^6 A549 cells. Tumor volume was initially

measured after 30 days. Subsequently randomized groups containing 6 animals each began receiving

treatment with vehicle, PAV-617, PAV-951, or Gemcitabine ("Reference ID" 4433223-

Accessdata.fda.gov," n.d.). The tumor volumes of animals within each treatment group were measured

over time. **Fig. 4A** shows tumor volume across time in the PAV-617 study where one treatment group

received daily doses of vehicle, one treatment group received daily doses of 10 mg/kg PAV-617, and one

treatment group received twice weekly doses of 100 mg/kg Gemcitabine for 28 days. After 28 days of

treatment, the Gemcitabine treated animals had a tumor volume that was, on average, 46% of tumor

volume found in the vehicle-treated animals and PAV-617-treated animals had a tumor volume that was,

on average, 47% of the tumor volume found in the vehicle-treated animals. **Fig. 4B** shows tumor volume

across time in the PAV-951 study where one treatment group received daily doses of vehicle, one

treatment group received daily doses of 1.5 mg/kg PAV-951, and one treatment group received twice

weekly doses of Gemcitabine for 14 days. After 14 days of treatment, the Gemcitabine treated animals

had a tumor volume that was, on average, 42% of tumor volume found in the vehicle-treated animals and

PAV-951-treated animals had a tumor volume that was, on average, 50% of the tumor volume found in

the vehicle-treated animals. Statistical significance is indicated with ** where the p value is < 0.01 and *

where the p value is < 0.05.

<u>Characterizing the targets of PAV-617 and PAV-951</u>

As PAV-617 and PAV-951 were identified by a phenotypic screen, their actual targets were unknown

during the early stages of compound advancement. To identify their targets, each molecule was coupled

to affigel resins from a position on the molecule unrelated to proliferation arrest activity based on

structure-activity-relationship (SAR) exploration. In that way, they could serve as target-binding ligands

for drug resin affinity chromatography (DRAC) (Tanaka, 2009). LNCaP C-33 cells were chosen for DRAC

starting material because we knew PAV-617 and PAV-951 displayed efficacy against them and we

wanted a model for our target engagement studies that would account for mechanisms of endogenous

apoptosis and innate immune system responses. Extracts were prepared from LNCaP C-33 cells that

were treated for 22 hours with either DMSO, PAV-617, or PAV-951. The extracts were applied to the

PAV-617, PAV-951, or control drug resins, washed with 100 bed volumes of buffer, and eluted with

either 100uM of PAV-617 or 100uM of PAV-951.

Samples of the DRAC eluate were sent for analysis by tandem mass spectrometry (MS-MS). The

DRAC eluate from the PAV-617 and PAV-951 resins contained large sets of proteins missing from the

control resin eluate. This included cancer-implicated proteins from the literature (see Figs. **5A**, **5B**, and

5C). 92 proteins from the DMSO-treated cell extracts were identified in the PAV-617 resin eluate that

were not present in the control resin eluate. 116 proteins from the DMSO-treated cell extracts were

identified in the PAV-951 resin eluate which were not present in the control resin eluate. Of the proteins

identified by MS-MS as unique or greatly enriched in the drug resin eluates, 38 proteins in the DMSO-

treated cell extracts were unique to PAV-617 (see **Fig. 5A**), 62 proteins in the DMSO-treated cell extracts

were unique to PAV-951 (see **Fig. 5B**), and 54 proteins from the DMSO-treated cell extracts were found

in both the PAV-617 and PAV-951 resin eluates (see **Fig. 5C**). When the proteins detected in the eluates

were searched in a database for cancer-implicated proteins, 23 of the proteins from the PAV-617 resin

eluate and 29 proteins from the PAV-951 resin eluate were known to be part of cancer-relevant

interactomes (see **Figs. 5A**, **5B**, and **5C**).

The MS-MS indicated that when LNCaP C-33 cells were treated with compound, composition of

the eluate was subsequently affected. For both the PAV-617 and the PAV-951 resin eluates, the spectral

counts of some particular proteins detected by MS-MS increased or decreased in treatment conditions

(see **Figs. 5A**, **5B**, and **5C**). However, for other proteins, the number of spectral counts detected in the

eluates remained unchanged upon treatment (see **Figs. 5A**, **5B**, and **5C**).

Figure 5. Identification of proteins in PAV-617 and PAV-951 resin eluates by MS-MS. DRAC experiments were performed where 30 ul of DMSO or compound-treated LNCaP cell extract, adjusted to a protein concentration of approximately 10 mg/ml in column buffer, was incubated on a column containing 30ul of affigel resin coupled to either PAV-617, PAV-951, or a 4% agarose matrix (control) for one hour at 4 degrees Celsius. The input material flow-through was collected and the resin was washed with 3 mL

column buffer then eluted overnight at 4 degrees Celsius in 100 ul of either 100uM PAV-617 or 100uM

PAV-951 in column buffer. **Figs. 5A-5C** show spectral counts of proteins detected by MSMS in single-point

DRAC eluates. **Fig. 5A** shows the set of proteins only detected in PAV-617 resin eluate. **Fig. 5B** shows the

set of proteins only detected in the PAV-951 resin eluate. **Fig 5C.** shows the set of proteins detected in

both the PAV-617 and the PAV-951 resin eluates. Conditional formatting has been applied where relative

abundance (by spectral count) of a given protein in a sample is visualized on a yellow-to-black scale**.**

Proteins implicated with cancer in the Bushman labs oncogene database

(http://www.bushmanlab.org/links/genelists) have been indicated in red.

When we entered the proteins identified from the DRAC eluate into a database of protein-

protein interactions, we discovered that many of the proteins eluted by either or both PAV-617 or PAV-

951 were known to be involved in networks that involved frequent interactions and associations with

each other (see **Supplemental Fig. 2**).

DRAC experiments were conducted side-by-side in triplicate with and without addition of

metabolic energy by running the experiment at 4 degrees Celsius versus 22 degrees Celsius and

supplementing with an "energy cocktail" of ribonucleotide triphosphates (1mM rATP, 1mM rGTP, 1mM

rCTP, 1mM rUTP), and 5 ug/mL creatine kinase. Changes in the amounts of proteins that bound and

eluted with PAV-617 or PAV-951 were observed by western blot in the presence of these metabolic

energy substrates (see **Figs. 6A** and **6B**). For the PAV-617 resin eluate, B-tubulin (TUBB), KRAB-

associated protein 1 (KAP1 also known as TRIM28), Methylenetetrahydrofolate dehydrogenase 1

(MTHFD1), and protein disulfide isomerase (PDI)— proteins which had previously been identified in the

eluate by MS-MS— all showed up by western blot in larger amounts under energy conditions (see **Fig.**

6A). For the PAV-951 resin eluate, MTHFD1, PDI, and heterogenous ribonucleoprotein K (hnRNPK) were

observed to be enhanced under energy conditions (see **Fig. 6B**). To the contrary, the amount of KAP1

identified in the PAV-951 resin eluate decreased in the presence of metabolic energy substrates (see **Fig.**

6B).

To confirm that the enrichment required the combination of metabolic energy substrates and

compound rather than the presence of energy substrates alone, resins were eluted side-by-side in

triplicate with either PAV-617, PAV-951, or 1% DMSO all containing the energy cocktail. By western blot,

eluates from both PAV-617 and PAV-951 resins were found to contain significant amounts of KAP1,

MTHFD1, and PDI relative to the control resin or DMSO elution (see **Figs. 6C and 6D**). The eluate of PAV-

617 resin contained TUBB which the PAV-951 resin eluate did not have in any greater amount than the

control (see **Figs. 6C and 6D**). The eluate of PAV-951 resin contained hnRNPK, not present in the PAV-

617 resin eluate in greater amount than the control (see **Figs. 6C and 6D**).

DRAC experiments were conducted under conditions where the starting extract was prepared

from LNCaP C-33 cells that had been treated with DMSO, 100 uM PAV-617, or 100 uM PAV-951 for 24

hours in order to see if the changes to the eluate observed by MS-MS would repeat with energy

supplementation (see **Fig. 5**). Western blots of triplicate-repeated samples showed the PAV-617 resin

eluate contained increased amounts of KAP1 and decreased amounts of TUBB when cells were treated

with PAV-617, meanwhile PAV-951 resin eluate showed decreased amounts of both KAP1 and hnRNPK

when cells were treated with PAV-951 (see **Figs. 6E** and **6F**).

Figure 6. Exploration of changes to PAV-617 and PAV-951 resin eluates under different conditions. DRAC experiments using LnCAP C-33 starting extract as described in **Fig. 5** were conducted under hot conditions which included supplementing the starting material and eluate with an energy cocktail to a final concentration of 1mM rATP, 1mM rGTP, 1mM rCTP, 1mM rUTP, and 0.05 ug/mL creatine kinase and running the experiment at room temperature verus cold conditions at 4 degrees Celsius and without the energy cocktail. **Figs. 6A** and **Fig. 6B** show quantitation of average integrated density of protein band detected by western blot in triplicate-repeat samples eluted with PAV-617 from the PAV-617 or eluted with PAV-951 from the PAV-951 resin when DRAC was conducted side-by-side in hot and cold conditions.

Figs. 6C and **6D** show quantitation of average integrated density of protein band detected by western blot

for TUBB, hnRNPK, KAP1, MTHFD1, and PDI in triplicate-repeat eluates under hot conditions from the

PAV-617 resin or PAV-951 resin and a single point elution with each compound from the control resin.

Resins were eluted with either compound or 1% DMSO and the amount of protein detected in the 1%

DMSO elution is subtracted from the compound elution in the figure. **Fig. 6E** shows quantitation of

average integrated density of protein band detected by western blot for KAP1 and B-tubulin in DRAC

eluates generated under hot conditions of DMSO versus PAV-617 treated cells eluted with PAV-617 in

triplicate from the PAV-617 resin and in single-point from the control resin. **Fig. 6F** shows quantitation of

average integrated density of protein band detected by western blot for KAP1 and hnRNPK in DRAC

eluates generated under hot conditions for DMSO versus PAV-951 treated cells eluted with PAV-951 in

triplicate from the PAV-951 resin or in single-point from the control resin**.**

One explanation for why the DRAC eluates contain large numbers of proteins is that drug targets

for PAV-617 and PAV-951 are themselves multi-protein complexes. To test this hypothesis and

determine which proteins directly bind the compounds and which are indirectly associated with the

compounds via protein-protein interactions involving the direct drug-binding protein(s), we modified

the drug into photocrosslinker analogs by attachment of diazirine and biotin functional groups at the

same position to the resin had previously been attached. The photocrosslinker analogs were designed so

that after an incubation with cell extract that would allow the compound to bind its target, exposure to

ultraviolet light would form a covalent bond between the diazirine moiety of the compound and the

nearest protein neighbor (MacKinnon and Taunton, 2009). The sample could then be solubilized and

precipitated with streptavidin beads (which bind biotin) to identify the drug-binding protein(s). The

streptavidin precipitation (SAP) could be done using a native sample, which would pick up the direct

drug-binding proteins(s) and with it any co-associated proteins. Or, the SAP could be done using a

denatured sample which would, by virtue of the covalent bond to biotin, identify only the drug-binding

protein(s), with all other associated proteins of the target multi-protein complex lost upon denaturation.

A549 cell extract was incubated with either 1% DMSO or the photocrosslinker analogs of PAV-

617 or PAV-951, then exposed to ultraviolet light. The samples were then divided in two equal parts,

where one part was left native and the other denatured, then both were adjusted to non-denaturing

conditions and incubated with streptavidin beads. Blots of the SAP samples for KAP1 indicate KAP1 is

only a component of the PAV-617 target under native conditions and is completely lost upon

denaturation (see **Figs. 7A** and **7B**). However, KAP1 is present to a significant extent in both native and

denatured conditions for the SAP with the PAV-951 crosslinker (see **Figs. 7A** and **7C**). The SAP samples

were sent for MS-MS analysis, however high background in the samples with no crosslinker added

rendered the data uninformative for definitive binding partner identification (data not shown).

Conventional methods of drug discovery typically involve the use of recombinant proteins to

measure affinity between a drug and its target (Hughes et al., 2011). However, we were concerned that

if protein-protein interactions between the direct drug-binding protein and other proteins comprise an

important dimension of PAV-617 and PAV-951's targets, isolated recombinant proteins would not be an

appropriate surrogate for the protein-protein interactions occurring *in vivo*. To measure target

engagement for PAV-617 and PAV-951, we returned to DRAC and determined whether passing a cell

extract over the PAV-617 or PAV-951 resins would deplete the extracts of bindable target.

We applied the flow-through of PAV-617, PAV-951, and control resins to a second copy of these

resins and demonstrated that the drug resins deplete the extract of essentially all KAP1 capable of

binding to the resins (see **Figs. 7D** and **7E**). Western blot of the resin flow-through showed that a

comparable amount of KAP1 was flowing through the PAV-617, PAV-951, and control resins without

binding (see **Fig. 7D**). However, when the flow-through that had been depleted on the PAV-951 resin

was applied onto a new PAV-951 resin, very little additional KAP1 bound to the second resin. By

contrast, application of the control resin flow-through (which was not specifically depleted of anything)

to a second PAV-951 resin, showed significant KAP1 binding to the second resin (see **Fig. 7E**). Application

of the PAV-617 resin flow-through to the PAV-951 resin showed significantly more KAP1 binding to the

second resin than from the PAV-951 flow-through, but significantly less KAP1 binding than from the

control flow-through (see **Fig. 7E**).

One notable observation was that, even though the five-fold depletion by PAV-951 resin of its

target compared to the control resin was statistically significant and reproducible, it accounted for a tiny

amount of the total KAP1 which had been detected by western blot in the starting extract. Only 0.7% of

the total amount of KAP1 detected in the starting material bound to the PAV-951 resin after passing

over a control column (see **Fig. 7E**). As a reference point to account for non-specific loss of material over

the course of the experiment (due to denaturation over time or nonspecific sticking removed during the

washing phase)— of the original PAV-951 resin that the extract was depleted with, serial elution with

PAV-951 and 1% SDS showed approximately 4% of the total KAP1 detected in the original extract had

bound to the resin (data not shown). We conclude that the compounds selectively target this small

fraction of KAP1 (less than 5% of the total KAP1 in the extract) because once the subfraction of KAP1

capable of binding to the resins is removed, the remaining extract will not bind anymore, even though it

still contains ample KAP1.

Several of the proteins found by MS-MS in the PAV-617 and PAV-951 resin eluates are known to

interact directly with KAP-1 (see **Figs. 7F** and **7G**). These KAP1 implicated proteins include some whose

relative amount in the eluate increased and/or decreased with drug treatment and are part of cancer

associated interactomes identified from the literature (see **Figs. 7F** and **7G**). The KAP1 implicated

proteins from the PAV-951 resin/eluate also contained several proteins that are associated with HIV in

the literature (see **Fig. 7G**). These associations may shed light on the cellular role played by the

subfractions of KAP1 that are targeted by PAV-617 and PAV-951.

Figure 7. KAP-1/TRIM28 as a component of the PAV-617 and PAV-951 target complexes. Fig. 7A shows

quantitation of average integrated density of KAP1 protein band detected by western blot while **Figs. 7B**

and **7C** show the western blots themselves, in triplicate-run native and denatured streptavidin

precipitations of photo crosslinked samples. Crosslinking experiments were performed where 65 uL of

A549 cell extract was adjusted to a protein concentration of approximately 1 mg/ml in column buffer and

supplemented with the energy cocktail, with either 1% DMSO or 1uM modified photo crosslinker analogs

461 of PAV-617 and PAV-951 for one hour at room temperature then 20 minutes on ice. The extracts were

462 exposed to UV light then divided into two aliquots. One aliquot was left native and one was denatured by

463 adding DTT, SDS, and boiling. 800 uL of column buffer with 0.1% triton was added to both aliquots and

464 then they were incubated with 2.5ul magnetic streptavidin beads for one hour at room temperature

465 before being denatured in loading buffer containing SDS and heated to 100°C for 3 minutes. **Figs. 7D** and

466 **7E** show the show quantitation of integrated density of KAP1 protein band detected by western blot in

467 LNCaP extract depleted on the PAV-617, PAV-951, and control resins. 230 uL of LNCaP extract was

468 incubated with 230uL of PAV-617, PAV-951, or control resins in single point for one hour in hot conditions.

469 Depleted flow-throughs were divided and put onto a subsequent PAV-951 column in triplicate or onto a

470 subsequent control column in single-point under hot conditions. Columns were eluted three times- a first

471 overnight elution with PAV-951, a second overnight elution with PAV-951, and a third elution with 1%

472 SDS. Eluates were diluted 3:1 in loading buffer and analyzed by western blot. Every western blot for the

473 eluate included a sample of the original, un-depleted starting LNCaP extract diluted 1:100 in loading

474 buffer. **Fig. 7D** shows quantitation from when the flow-throughs from each columns were blotted for

475 KAP1 to determine how much KAP1 had been depleted from each resin. **Fig. 7E** shows the amount of

476 protein detected in the eluate normalized as percent of their corresponding total sample, where the

477 amount detected by eluate samples were divided by the amount detected in the starting material sample,

478 then multiplied by 0.013 to match the concentrations. The percent detected by western blot from the two

479 overnight elutions and the SDS elutions were added together to determine the total percentage of cellular

480 KAP1 was binding to and eluting from the resins). **Figs. 7F** and **7G** show diagrams of proteins identified by

481 MS-MS in Figs. 5A-5C as comprising the PAV-617 resin/eluate and PAV-951 resin/eluate found in the

482 NURSA database of protein-protein interactions as interacting with KAP1

483 (https://dknet.org/about/NURSA_Archive) (Malovannaya et al., 2011). Red indicates KAP1 implicated

484 proteins detected in the eluate which decreased with treatment. Green indicates KAP1 implicated

485 proteins detected in the eluate which increased with treatment. Yellow shows KAP1 implicated proteins

486 detected in the eluate which were unchanged with drug treatment. Circles indicate proteins from the

487 PAV-617 and PAV-951 resin eluates implicated in cancer from the Bushman lab oncogene database

(http://www.bushmanlab.org/links/genelists). Squares indicate proteins from the PAV-951 resin

eluates implicated in HIV from the virus mentha database (https://virusmentha.uniroma2.it/).

Discussion

Our data indicate that PAV-617 and PAV-951, two assembly modulator anti-viral compounds

selected for their ability to arrest proliferation in a distinctive way by a novel screen, are cytotoxic to a

wide range of neoplastic cell lines representing both rare and common cancers. These compounds,

while early in their drug optimization, performed comparably to the commercial anti-cancer drug

Gemcitabine, at 10 and 60 fold lower doses, to inhibit the growth of an A549 tumor in immunodeficient

mice. These data suggest that these compounds are directed to two different targets common to a wide

range of cancers, and may provide a starting point for the development of novel cancer therapeutics.

DRAC and photocrosslinking experiments indicate that PAV-617 and PAV-951 interact with

proteins that are part of multi-protein complexes. These complexes are dynamic, as demonstrated by

changes in the eluate when DRAC is carried out in the presence versus absence of metabolic energy

substrates or from untreated versus compound-treated cell starting extract. Together, these findings

suggest a model for disease pathogenesis in which previously unappreciated transient multi-protein

complexes plays an important role in the dynamics linking cellular proliferation to apoptosis. We

hypothesize that cancer progression is facilitated by aberrant versions of these multi-protein complexes

and the effect of assembly modulating drugs is to restore the original version of the multi-protein

complex, possibly through an allosteric mechanism-of-action (Motlagh and Hisler, 2012), see **Fig. 8**.

Figure 8. **Cartoon diagram of proposed mechanism of action of assembly modulating compounds**. **Fig.**

8A shows the proposed model where a normal multi-protein complex that plays a role mediating the

linkage between uncontrolled proliferation and apoptosis is modified into an aberrant multi-protein

complex at a precancerous stage allowing cancer progression rather than homeostatic elimination. **Fig. 8B**

shows the proposed mechanism where treatment with an assembly modulating compound restores the

original multi-protein complex and its homeostatic functions including elimination of the cancer. Allosteric

modulation is indicated as a means by which these changes may be induced.

KAP1/TRIM28, one identified protein component of the PAV-617 and PAV-951 target complexes,

is worthy of specific mention. KAP-1 was identified by both MS-MS and western blot as being part of the

PAV-617 and PAV-951 targets (see **Figs. 5**, **6**, and **7**). Crosslinking experiments showed that for PAV-617

KAP1 is present in a complex targeted by the compound under native conditions but is lost upon

denaturation, indicating it is not a direct drug-binding protein, but rather more likely a distal component

of the target multi-protein complex (see **Fig. 7**). For PAV-951, KAP1 is also part of the complex under

native conditions and the data indicates a portion of KAP1 may be directly bound to the compound as

well. This suggests more than one copy of KAP1 per multi-protein complex (see **Fig. 7**).

KAP1 stood out as being of particular interest because it is a known allosteric modulator,

implicated in both infectious and noninfectious disease (Randolph et al., 2022, p. 1). KAP1 is involved in

a variety of protein-protein interactions and an array of functions including transcriptional activation of

HIV, T-cell development, DNA damage repair, as a transcriptional co-repressor for many genes, and as a

ligase for post translational modifications such as ubiquitination and SUMOlyation (Iyengar and

Farnham, 2011; Randolph et al., 2022). Studies have shown increased levels of KAP-1 in many types of

cancer and high levels of KAP-1 correlate with aggressive clinical phenotype and progression to

metastasis (Addison et al., 2015; Cui et al., 2014; Yu et al., 2014). However, other described functions of

KAP-1 are tumor suppressive and promote autophagy (Neo et al., 2015; Randolph et al., 2022; Yang et

al., 2013). KAP-1 directly binds with cancer-implicated proteins including MDM2, TRIM24, Fructose-1,6-

biphosphatase (FBP1), MAGE-A3, MAGE-C2, Heat shock protein 70, and TWIST1 (Fong et al., 2018; Jin et

al., 2017; Wang et al., 2005; Wei et al., 2016; Yang et al., 2013). The diversity of functions that KAP-1

displays appears to be, at least in part, through its assembly into different multi-protein complexes that

carry out different objectives.

The literature describes that KAP1 is utilized by HIV to the host's detriment but also functions as

a key component of the host's immune response in repressing HIV (Allouch et al., 2011; Randolph et al.,

2022). KAP1 is implicated as a key part of a pathway hijacked by the poxvirus p28 virulence factor

(Huang et al., 2004). We show that PAV-951 is active against HIV and PAV-617 is active against MPXV in

addition to their anti-cancer phenotypes. We hypothesize that there are shared alterations in protein

assembly in neoplastic cells and virus-infected cells. Advancing the SAR of PAV-617 and PAV-951 could

be utilized to determine whether the antiviral and anti-cancer targets are identical or merely similar. In

the latter case, the activities would separate with further SAR. Regardless, these findings suggest that

the initial CFPSA capsid assembly screen was successful in identifying novel targets relevant to both

viruses and cancer.

We suggest that PAV-617 and PAV-951 redirect protein-protein interactions, whereby particular

proteins are recruited to, and other particular proteins are expelled from, particular multi-protein

complexes in the presence of compound. KAP1 is known to both promote and suppress tumorigenesis

(Randolph et al., 2022). Therefore, a compound which selectively targets some forms of KAP1 and not

others would be important regardless of the nature of the interaction with KAP1. In some cases, as

observed for PAV-951, a portion of KAP1 is a direct drug-binding protein, although other copies of KAP1

appear not to be. In the case of PAV-617 resin KAP1 is associated with the drug only indirectly by virtue

of being a protein present in the target multi-protein complex, but at a distance from the drug-binding

site, thus present by SAP under native but not denatured conditions.

We interpret the DRAC flow-through data to mean that, by virtue of conformation or other

differences among co-associated proteins in a transient multi-protein complex, subfractions of cellular

KAP1 are selectively targeted by PAV-617 and PAV-951. Results from the cross-depletion where the

flow-through of the PAV-617 resin was applied to a new PAV-951 resin further indicate that the PAV-

617-binding subfraction of KAP1 must be distinguishable from the PAV-951-binding subfraction of KAP1

because depletion of one does not fully deplete the other (see **Fig. 7E**).

These findings about PAV-617 and PAV-951 mirror those made for PAV-104, a structurally

unrelated assembly modulator with pan-respiratory anti-viral activity (Müller-Schiffmann et al., 2022). It

appears as though assembly modulating compounds, despite structural diversity, share characteristics

including the targeting of multi-protein complexes, selectivity for a small fraction of the total amount of

a given protein found in a cell, and allosteric mechanisms of action (U. F. Lingappa et al., 2013; Müller-

Schiffmann et al., 2022, 2021b; Motlagh and Hisler, 2012). Furthermore, they appear to have

remarkable activity across broader categories of pathogens (e.g. pan-viral family and pan-cancer) than is

generally observed for existing drugs. The antiviral assembly modulators appear to have a barrier to the

development of resistance (Müller-Schiffmann et al., 2022). Further studies are needed to determine

whether this property holds true for the anti-cancer subset of assembly modulators.

Effective cancer drugs have been developed based on a number of mechanisms including

alkylating agents, antimetabolites, antimitotics, and monoclonal antibodies (Falzone et al., 2018).

However, as far as we know, no one has attempted to treat cancer through modulation of protein

assembly, giving our work with PAV-617 and PAV-951 the potential to be both risky and rewarding.

While the animal toxicity of PAV-617 and PAV-951 is higher than would be ideal for the clinic, their anti-

tumor activity is already on par with existing cancer drugs and a handful of FDA approved cancer drugs

have comparable toxicity gauged by mouse MTD— Cisplatin has a MTD of 6 mg/kg, Doxorubicin has a

MTD of 10 mg/kg, and Vinorelbine has a MTD of 10 mg/kg— and many other cancer drugs are

administered to patients despite adverse effects because of the urgency of their condition (Aston et al.,

2017; Singh and Singh, 2018). Since PAV-617 and PAV-951 are early compounds, further optimization

will likely yield chemical analogs with substantially reduced toxicity and further increased activity. We

hypothesize that driving SAR toward compounds that are selective for modulating the cancer-induced

protein assemblies will improve the therapeutic indexes.

Molecular genetic tools such as CRISPR, siRNA knock down, and even use of recombinant

protein for protein-protein interaction studies, are unable to parse out the post-translational

heterogeneity introduced into proteins as part of normal and aberrant biochemical pathways. The

methods applied here are able to do so, as evidenced by the small fraction of the total of specific

proteins such as KAP1 found in the target multi-protein complex. It is perhaps not surprising that new

tools, and the new targets they allow to be detected, make possible a path to the development of drugs

with new properties. Further work will clarify whether, as we hypothesize, the transience of our targets

reflects their involvement as a molecular basis for homeostasis. This conclusion, supported by the

consequences of assembly modulator treatment both here for cancer and previously for viruses, frames

future experiments to better understand this novel approach to more physiological disease

therapeutics.

G	animal	route	dose (mg/kg)	C_{max} (uM)	half life (h)	mean residence time (h)
	rat	IV	1	2.78	7.84	7.03
	rat	IP	5	0.01	7.04	8.31
	mouse	oral	10	2.02	5.56	7.66

H	animal	route	dose (mg/kg)	C_{max} (uM)	half life (h)	mean residence time (h)
	rat	IV	0.5	0.25	1.68	0.63
	rat	IP	2.5	0.71	nd	2.92
	mouse	oral	10	2.16	13.6	19.3

Supplemental Figure 1. Summary of PAV-617 and PAV-951 PK properties. Supplemental Figs. A-F show

plasma concentration over time for PAV-617 and PAV-951 when given to animals via three different

routes of administration. Randomized treatment groups of four male Sprague Dawley rats or three CD1

mice were administered vehicle, PAV-617, or PAV-951 either IV, IP, or orally and blood samples were

collected before injection as well as at different time points after dosing. The concentration of compound

in the plasma at different time points was measured by LC MS/MS. **Supplemental Fig. G** summarizes PK

properties observed for PAV-617 based on animal, administration route, and dose. **Supplemental Fig. H**

summarizes the PK properties observed for PAV-951 based on animal, administration route, and dose.

The maximum concentration (Cmax) was determined to be the highest measured concentration within a

dataset. The half life was calculated as time required for the concentration of compound in the animal's

plasma to decrease by half. The mean residence time was calculated as the average time the compound

remained in the animal.

Supplemental Figure 2. Known protein-protein interactions among PAV-617 and PAV-951 eluate

components. **Supplemental Fig. 2A** and **2B** show string-diagram analyses of the protein-protein

interaction network of proteins identified in **Figs. 5A-5C** by tandem MSMS as comprising the PAV-617

(**Supplemental Fig. 2A)** and PAV-951 (**Supplemental Fig. 2B**) targets. Proteins were entered into the string

database (string-db.org). The confidence mode view is shown where the confidence level was set to 0.4

(medium) and MCL clustering was applied with the inflation parameter of 3. The dotted lines represent

edges between different clusters. The thickness of the lines represent the degree of confidence for each

protein-protein interaction.

Materials and Methods

Lead Contact and Materials Availability

Further information and requests for resources and reagents should be directed to and will be fulfilled

by the Lead Contact Vishwanath Lingappa (vlingappa@prosetta.com).

Use of unique compounds PAV-617 and PAV-951 and their stable derivatives may be available upon

request by the Lead Contact if sought for experimental purposes under a valid completed Materials

Transfer Agreement.

The number of replicates carried out for each experiment is described in the figure/table legends.

Experimental Models and Subject Details

Animal models

Maximum tolerated dose (MTD) studies were conducted using female Balb/C mice, aged 8-10

weeks or female CD1 mice, aged 5-6 weeks. Treatment groups were made up of 3 animals each, unless

otherwise noted, and dosing regimens for disclosed data is provided. Animals were sacrificed at the end

of the study period using an overdose of CO_2. MTD studies were conducted at Vipragen Biosciences

Private Limited in accordance with the Committee for the Purpose of Control and Supervision of

Experiments on Animals (CPCSEA) guidelines and Animal Research: Reporting of In Vivo Experiments

(ARRIVE) guidelines.

Pharmacokinetic (PK) studies were conducted in male Sprague Dawley Rats aged 8-10 weeks or

male CD1 mice aged 5-6 weeks. Treatment groups were made up of 4 animals each and dosing regimens

for disclosed data is provided. Animals were sacrificed at the end of the study period using an overdose

of CO2. PK studies were conducted at Vipragen Biosciences Private Limited in accordance with the

Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) guidelines

and Animal Research: Reporting of In Vivo Experiments (ARRIVE) guidelines.

Tumor xenograft studies were conducted using female Athymic Nude mice, strain CrTac: Ncr-

Foxn1^{nu}, aged 6-8 weeks. Tumor transplantation occurred through subcutaneous injection of a 0.1mL

cell suspension containing 1 to 5×10^6 A549 lung cancer cells obtained from ATCC in Matrigel in PBS into

the left flank region of the mice. Treatment groups were made up of 6 animals each and dosing

regimens for disclosed data is provided. Animals were sacrificed at the end of the study period using an

overdose of isoflurane anesthesia. Both xenograft studies were conducted at Anthem Biosciences

Private Limited in Bangalore, India and were approved by the Institutional Animal Ethics committee

(IAEC) of Anthem Biosciences in accordance with the Committee for the Purpose of Control and

Supervision of Experiments on Animals (CPCSEA) guidelines and Animal Research: Reporting of *In Vivo*

Experiments (ARRIVE) guidelines.

Cell lines

The *in vivo* xenograft study utilized the A549 (male human lung cancer). Cells were grown under

sterile conditions. These studies were conducted at Anthem Biosciences Private Limited in Bangalore,

India.

The Human tumor cell proliferation assay used A172 (male human glioma), BFTC-905 (female

human urinary bladder transitional cell carcinoma), COR-L105 (male human lung adenocarcinoma), DB

(male human b-cell lymphoma), FaDu (male human pharynx squamous cell carcinoma), H9 (male human

t-cell lymphoma), Hs 294T (male human melanoma), MCF7 (female human breast cancer), MDA MB 436

(female human breast cancer), MeWo (male human melanoma), MHH-PREB-1 (male human b-cell

lymphoma), SJSA1-OSA (male human osteosarcoma), SU-DHL-10 (male human b-cell lymphoma),

SW1353 (female human chondrosarcoma), and U-2 OS (female human osteosarcoma) cell lines. This

study was conducted by Eurofins Scientific as part of their OncoPanel™.

The monkey pox virus infectious virus assay used BSC-40 cells (nonhuman primate kidney) and

MPXV Zaire 79 strain. These studies were conducted by the United States Army Medical Research

Institute of Infectious Diseases on Fort Detrick, Maryland.

Human immunodeficiency virus infectious virus assay used MT-2 cells (female human t-cell

leukemia) and the NL4-3 Rluc reporter virus. These studies were conducted at the University of

Washington in Seattle, Washington.

The *in vitro* screens for apoptosis, high-density/ low-density activity, and cell growth recovery

utilized the LNCaP C-33 (male human prostate cancer), LNCaP C-81 (male human prostate cancer), CHO-

K1 (Chinese Hamster ovary), and Hennes 20 (Chinese Hamster ovary) cell lines. Cells were grown under

sterile conditions. These studies were conducted at Prosetta Biosciences in San Francisco, California.

The *in vitro* drug resin affinity chromatography and photocrosslinking experiments utilized A549

(male human lung cancer) or LNCaP C-33 (male human prostate cancer) cell line. Cells were grown under

sterile conditions. Sterile conditions were not maintained once cells were harvested for *in vitro*

experiments. These studies were conducted at Prosetta Biosciences in San Francisco, California.

***In vitro* experiments**

Drug resin affinity chromatography and photocrosslinking experiments, and SDS-PAGE/Western

Blot analysis of the results, were conducted by Prosetta Biosciences in San Francisco, California under

conditions described in figure legends. Results from disclosed *in vitro* experiments were repeated in

triplicate unless otherwise stated. Mass spectrometry analysis of samples were conducted by MS

Bioworks in Ann Arbor, Michigan.

Method and Analysis Details

Monkey pox infectious virus assay

BSC-40 cells of 95% confluence in 24-well plates were infected with100 pfu of MPXV Zaire 79

diluted in Eagle's Minimum Essential Medium with 2% fetal bovine serum and incubated in 37 degrees

Celsius in 5% CO_2 for 1 hour. The viral inocula were removed and replaced with the test compounds in

six half log dilutions (0.1 ml per well) and the cells were overlaid with 1% methylcellulose in growth

media (1 ml per well). The media and virus control cells received growth medium containing 1%

methylcellulose. After three days of infection, when plaques appeared, cells were stained with crystal

violet for an hour and then washed with water and dried overnight. The plaques were counted the next

day and virus-only wells were compared with the compound-added wells to determine percentage

protection. Infected cells were stained with crystal violet and viral plaques were counted. Averages and

standard deviation for plaques observed under different treatment conditions were calculated in

Microsoft Excel and graphed as the percent inhibition in PAV-617 treated cells compared to untreated

cells.

Human immunodeficiency virus infectious virus assay

MT-2 cells were preseeded in 96-well plates in 100 ul of complete RPMI. Multiple

concentrations of PAV-951 were serially diluted in DMSO then into an infection media prepared by

diluting NL4-3 Rluc virus stock to 400 IU/100 ul with complete RPMI, which was transferred onto the

MT-2 cells with a final MOI of 0.02 and final DMSO concentration of 1% in infected places. One well

received DMSO only, instead of PAV-951, and one well received medium only for normalization and

background collection. Cells were incubated at 37 degrees Celsius for 96 hours. 100ul of medium was

removed and discarded and 10 ul of 15 uM EnduRen luciferase substrate was added to each well,

followed by incubation for 1.5 hours at 37 degrees Celsius. Plates were read on a luminescence plate

reader. Bioluminescence intensity was read on a Synergy H1 BioTek plate reader. Averages and standard

deviation for viral titer observed under different treatment conditions were calculated in Microsoft Excel

and graphed as the percent inhibition in PAV-951 treated cells compared to untreated cells.

Apoptosis Screen

A 96 well plate was seeded with Hennes 20 cells at 500 cells per well, CHO-K1 cells at 500 cells

per well, LNCaP C-33 cells at 2000 cells per well, and LNCaP C-81 cells at 2000 cells per well. Cells were

grown in 100uL minimum essential media for three days then three wells of each cell line received

treatment with 1% DMSO. 12 hours after drug treatment, a mixture of 25 ul media and 25 ul Apo-ONE

reagent (Promega) was added then the plate was covered and placed on a shaker at room temperature

for six hours. The plate was read on a microplate reader for fluorescence at 499/521. Values were

averaged and standard deviations were calculated for each triplicate condition and graphed on

Microsoft Excel.

High density/ low density assay

Two 96 well plates were seeded with Hennes 20 cells in parallel where one was plated at a

density of 500 cells/well and the other was plated at a density of 15,000 cells/well. 90 ul of minimum

essential media was added to each well and plates were placed in a 37 degrees Celsius incubator for 24

hours. The next day, 10ul of media containing dilutions of compound in DMSO were added to each plate

in triplicate with final concentrations of 0.025 uM PAV-617, 0.05 uM PAV-617, 0.1 uM PAV-617, 0.5 uM

PAV-617, 0.02 uM PAV-951, 0.3 uM PAV-951, 0.4 uM PAV-951, or 0.5 uM PAV-951. Six wells on each

plate received 10ul of media containing only DMSO. Each well was gently mixed 5 times with a 100ul

pipette. Plates were incubated at 37 degrees Celsius for 72 hours then 10 uL of alamarBlue was added to

each well. Wells were mixed 5 times then incubated at 37 degrees Celsius for 72 hours. Plates were then

read at 530/590. Values were averaged and standard deviations were calculated for each triplicate

condition and graphed on Microsoft Excel.

Cell growth recovery assay

A 96 well plate was seeded with either Hennes 20 or LNCaP C-33 cells at 500 cells/well in 90 uL

of minimum essential media and incubated at 37 degrees Celsius for 24 hours. 0.5% DMSO was diluted

in media and added to 6 control wells for each plate. PAV-617 was diluted in media and added to three

wells at a concentration of 0.3 uM. PAV-951 was diluted in media and added to concentration of 0.4 uM.

After 24 hours of PAV-617 treatment or 6 hours of PAV-951 treatment, the medium containing

compound was removed and replaced with fresh media. After 72 hours (day 5), plates were assayed

with alamarBlue and fluorescence was read at 530/590. The medium containing alamarBlue was

removed and replaced with fresh media. After another 72 hours (day 8) plates were assayed with

alamarBlue and fluorescence again, then medium containing alamarBlue was removed and replaced

with fresh media. After a final 72 hour incubation (day 11) plates were assayed with alamarBlue one

more time. Average fluorescence for each day and treatment condition were plotted on Microsoft Excel

with standard deviation calculated to provide error bars.

Human tumor cell proliferation assay

A panel of human tumor cell lines (A172, BFTC-905, COR-L105, DB, FaDu, H9, Hs 294T, MCF7,

MDA MB 436, MeWo, MHH-PREB-1, SJSA1-OSA, SW1353, and U2OS) were grown in RPMI 1640, 10%

FBS, 2 mM L-alanyl-L-glutamine, 1 mM Na pyruvate. Cells were seeded into 384-well plates and

incubated in a humidified atmosphere with 5% CO2 at 37C. After 24 hours of incubation DMSO, PAV-

617, or PAV-951 was added at concentrations of 5 uM, 1 uM, and 0.2 uM and plates were incubated for

3 days. Then cells were lysed with CellTiter-Glo (Promega) which generates a bioluminescence signal

relative to ATP levels and is used as a measurement of viable cells. Bioluminescence was read by a

PerkinElmer Envision microplate reader. Bioluminescence intensity was measured by a PerkinElmer

Envision microplate reader and transformed to a percent of control (POC) using the formula:

POC=(Ix/I0)*100, where Ix is the whole well signal intensity at a given treatment, and I0 is the average

intensity of the untreated vehicle wells. Values were averaged for each triplicate condition and graphed

on Microsoft Excel.

Mouse maximum tolerated dose studies

For the intraperitoneal MTD study, female Balb/c mice aged 8-10 weeks were randomly divided

into treatment groups with three animals per group. Animals in each treatment group were weighed

and received one IP injection of 0.1-0.15mL containing either vehicle (10% DMSO, 45% propylene glycol,

45% sterile water), 1mg/kg PAV-617, 2 mg/kg PAV-617, 5 mg/kg PAV-617, 10 mg/kg PAV-617, 1mg/kg

PAV-951, 2.5 mg/kg PAV-951, 5mg/kg PAV-951 or 10 mg/kg PAV-951. Animals were observed from day 0

until day 3 for clinical signs of toxicity. Animals were euthanized after 72 hours and were examined

externally and internally by a pathologist for abnormalities in organ weight and tissue damage. Blood

samples were sent for a complete blood count bioanalysis. MTD was determined to be the dose at

which no signs of toxicity were observed by any parameters.

For the oral MTD study, female CD1 mice aged 5-6 weeks were given either an oral dose of vehicle (10%

DMSO, 45% propylene glycol, 45% sterile water) or either 10 mg/kg or 20 mg/kg of PAV-617 or PAV-951.

The vehicle and 20 mg/kg groups had three animals each, while the 10 mg/kg groups only had one

animal. Animals were observed for clinical signs and after a week they were euthanized and examined

externally and internally by a pathologist for changes related to toxicity.

Pharmacokinetics studies

For the IP and IV PK studies, male Sprague Dawley rats aged 8-10 weeks were randomly divided

into treatment groups with four animals per group. Animals in each treatment group were weighed and

received one 2.4 mL intravenous dose of either vehicle (10% DMSO, 45% propylene glycol, 45% sterile

water), 1mg/kg PAV-617, or 0.5 mg/kg PAV-951, or one intraperitoneal dose of either vehicle (100%

labrasol), 5mg/kg PAV-617 or 2.5 mg/kg PAV-951 . Blood was collected from a pre-cannulated line

before dosing, and subsequently 15 minutes, 30 minutes, 1 hour, 2 hours, 4 hours, 8 hours, 12 hours, 24

hours, and 30 hours post-dosing. Concentration of drug in the plasma over time was measured using a

Waters Acquity TQD LCMS/MS. Maximum concentration (Cmax) was determined to be the maximum

concentration detected in a dataset. Half life, area under the curve, and mean residence time were

calculated with Phoenix WinNolin software.

For the oral PK studies, male CD1 mice aged 5-6 weeks were randomly divided into treatment

groups with three animals per group. Animals in each treatment group were weighed and received, via

oral gavage needle, either one oral dose of vehicle (10% DMSO, 45% propylene glycol, 45% sterile

water), 10 mg/kg PAV-617, or 10 mg/kg PAV-951. Blood was collected from a pre-cannulated line before

dosing, and subsequently 2 minutes, 5 minutes, 15 minutes, 30 minutes, 1 hour, 2 hours, 4 hours, 6

hours, 8 hours, and 24 hours post-dosing. Concentration of drug in the plasma over time was measured

using a Waters Acquity TQD LCMS/MS. Maximum concentration (Cmax) was determined to be the

maximum concentration detected in a dataset. Half life, area under the curve, and mean residence time

were calculated with Phoenix WinNolin software.

Animal xenograft study

A549 cells growing in RPMI-1640 medium were suspended with Matrigel in PBS. 0.1 ml of cell

suspension containing 1 x 106 cells were injected subcutaneously into the left flank region of female, 6-8

weeks old nude mice (CrTac: Ncr-Foxn1nu). After 30 days of tumor establishment, mice were divided

randomly into treatment groups. In the PAV-617 study, 6 animals were treated with vehicle only (10%

DMSO, 10% propylene glycol, 80% sterile water) by IP once daily, 6 animals were treated with 100

mg/kg Gemcitabine Hydrochloride by IP twice weekly, and 6 animals were treated with 10 mg/kg PAV-

617 by IP once daily for 28 days. In the PAV-951 study, 6 animals were treated with vehicle only (10%

DMSO, 10% propylene glycol, 80% sterile water) by IV once daily, 6 animals were treated with 100

mg/kg Gemcitabine by IV twice weekly, and 6 animals were treated with 1.5 mg/kg PAV-951 by IV once

daily for 14 days. In both studies, mice were weighed and their tumors were measured using a digital

Vernier caliper. Tumor volume was calculated using the formula: $(L \times W^2)/2$ where L is the largest

diameter and W is the smallest diameter of the tumor. Statistical analysis was performed using Graph

Pad Prism (Ver. 5.03). Statistical analysis of tumor growth inhibition between the Control and Treated

groups was performed by using One-way ANOVA followed by Dunnett's test.

 Drug Resin affinity chromatography

LNCaP C-33 cells were grown in minimum essential media (UCSF) with 10% FBS and 1% Penstrep

for 24 hours then treated with 500nM PAV-617, 500nM PAV-951, or DMSO for 22 hours. Cells were

scraped into cold phosphate buffered saline (PBS) (10mM sodium phosphate, 150 mM sodium chloride

pH 7.4), then spun at 1,000 rpm for 10 minutes until pelleted. The PBS was decanted and the pellet

resuspended in a low salt buffer (10mM HEPES pH 7.6, 10mM NaCl, 1mM MgAc with 0.35% Tritonx100)

then centrifuged at 10,000 rpm for 10 minutes at 4 degrees Celcius. The post-mitochondrial supernatant

was removed and adjusted to a concentration of approximately 10 mg/ml and equilibrated in a

physiologic column buffer (50 mM Hepes ph 7.6, 100 mM KAc, 6 mM MgAc, 1 mM EDTA, 4mM TGA). In

some conditions, the extract was supplemented with an energy cocktail (to a final concentration of

1mM rATP, 1mM rGTP, 1mM rCTP, 1mM rUTP, and 5 ug/mL creatine kinase). 30 ul or 230 ul of extract

was then incubated for one hour at either 4 degrees Celcius or 22 degrees Celsius on 30 ul or 230 ul of

affigel resin coupled to either PAV-617, PAV-951, or a 4% agarose matrix (control). The input material

was collected and the resin was then washed with 3 ml column bufffer. The resins were eluted overnight

at either 4 degrees Celcius or at 22 degrees Celsius in 100ul or 330ul column buffer containing either

100uM PAV-617 or 100uM PAV-951 or DMSO, with or without the energy cocktail. Eluates were run on

western blot or sent for mass spectrometry for analysis.

Chemical photocrosslinking

A549 extract was prepared as above then adjusted to a protein concentration of approximately

1 mg/ml in column buffer containing 0.01% triton and supplemented with the energy cocktail (to a final

concentration of 1mM rATP, 1mM rGTP, 1mM rCTP, 1mM rUTP, and 5 ug/mL creatine kinase).

Photocrosslinker analogs of PAV-617 and PAV-951 chemically modified to contain biotin and a diazirine

group or 1% DMSO were added to 67 ul of extract at 100uM, incubated for one hour at 22 degrees

Celsius followed by 20 minutes on ice, then exposed to ultraviolet light for 5 minutes at 22 degrees

Celsius. After crosslinking, samples were divided in two 30 ul aliquots and one set was denatured by

adding 5 ul of 10% SDS, 0.625 ul DTT, and boiling for 5 minutes. Both native and denatured aliquots

were then diluted in 800 ul column buffer containing 0.1% triton. 2.5 ul of magnetic streptavidin beads

(Pierce) were added to all samples and mixed for one hour at room temperature to capture all

biotinylated proteins and co-associated proteins. Samples were placed on a magnetic rack to hold the

beads in placed and washed three times with 800 ul of column buffer containing 0.1% triton. After

washing, beads were resuspended in 80 ul of gel loading buffer containing SDS and analyzed by western

blot or blot for affinity purified streptavidin. Samples were analyzed by western blot.

Western blotting

SDS/PAGE gels were transferred in Towbin buffer (25mM Tris, 192mM glycine, 20% w/v

methanol) to polyvinylidene fluoride membrane, blocked in 1% bovine serum albumin (BSA) in PBS,

incubated overnight at 4 degrees Celsius in a 1:1,000 dilution of 100ug/mL affinity-purified primary IGG

to KAP-1, MTHFD1, hnRNPk, TUBB, or PDI in 1% BSA in PBS containing 0.1% Tween-20 (PBST).

Membranes were then washed twice in PBST and incubated for two hours at room temperature in a

1:5000 dilution of secondary anti-rabbit or anti-mouse antibody coupled to alkaline phosphatase in

PBST. Membranes were washed two more times in PBST then incubated in a developer solution

prepared from 100 uL of 7.5 mg/mL 5-bromo-4-chloro-3-indolyl phosphate dissolved in 60% dimethyl

formamide (DMF) in water and 100ul of 15 mg/ml nitro blue tetrazolium dissolved in 70% DMF in water,

adjusted to 50mL with 0.1 Tris (pH 9.5) and 0.1 mM magnesium chloride. Membranes were scanned and

the integrated density of protein band was measured on ImageJ. Averages and the standard deviation

between repeated experiments were calculated and plotted on Microsoft Excel.

Tandem mass spectrometry

Samples were processed by SDS PAGE using a 10% Bis-ttris NuPAGE gel with the MES buffer

system. The mobility region was excised and washed with 25 mM ammonium bicarbonate followed by

15mM acetonitrile. Samples were reduced with 10 mM dithoithreitol and 60 degrees Celsius followed by

alkylation with 5o mM iodacetamide at room temperature. Samples were then digested with trypsin

(Promega) overnight (18 hours) at 37 degrees Celcius then quenched with formic acid and desalted using

an Empore SD plate. Half of each digested sample was analyzed by LC-MS/MS with a Waters

NanoAcquity HPLC system interfaced to a ThermoFisher Q Exactive. Peptides were loaded on a trapping

column and eluted over a 75 uM analytical column at 350 nL/min packed with Luna C18 resin

(Phenomenex). The mass spectrometer was operated in a data dependent mode, with the Oribtrap

operating at 60,000 FWHM and 15,000 FWHM for MS and MS/MS respectively. The fifteen most

abundant ions were selected for MS/MS.

Data was searched using a local copy of Mascot (Matrix Science) with the following parameters:

Enzyme: Trypzin/P; Database: SwissProt Human (concated forward and reverse plus common

contaminants); Fixed modification: Carbamidomethyl (C) Variable modifications: Oxidation (M), Acetyl

(N-term), Pyro-Glu (N-term Q), Deamidation (N/Q) Mass valuse: Monoisotopic; Peptide Mass Tolerance:

10 ppm; Gragment Mass Tolerance: 0.02 Da; Max Missed Cleavages: 2. The data was analyzed by label

free quantitation (LFQ) and spectral count methods. LFQ intensity values of each condition were

measured in triplicate and compared against each other to generate $\log_2$ fold change values for each

combination of conditions. Spectral counts were filtered for a 1% protein/peptide false discovery rate

requiring 2 unique peptides per protein and the data set was furher adjusted by substraction of spectral

counts for specific proteins observed in the control resin. Identified proteins were spearched in the

NURSA database of protein-protein interactions (https://dknet.org/about/NURSA_Archive) to determine if

they interact with KAP-1, the Bushman labs oncogene database

(http://www.bushmanlab.org/links/genelists) to determine if they were known to be implicated in

cancer, and the VirusMentha database (https://virusmentha.uniroma2.it/) to determine if they interact

with HIV. String diagrams of protein-protein interaction networks were generated using the STRING

database (string-db.org).

Acknowledgements:

We thank Usha F. Lingappa for help with the figures, Dmitry Temnikov for IT support, Halley McCormick

for help with data analysis, and Jairam R. Lingappa for valuable suggestions during manuscript

preparation.

Competing interests:

Vishwanath R. Lingappa is CEO of Prosetta Biosciences.

References

Addison, J.B., Koontz, C., Fugett, J.H., Creighton, C.J., Chen, D., Farrugia, M.K., Padon, R.R., Voronkova,

M.A., McLaughlin, S.L., Livengood, R.H., Lin, C.-C., Ruppert, J.M., Pugacheva, E.N., Ivanov, A.V.,

2015. KAP1 Promotes Proliferation and Metastatic Progression of Breast Cancer Cells. Cancer

Res. 75, 344–355. https://doi.org/10.1158/0008-5472.CAN-14-1561

Allouch, A., Di Primio, C., Alpi, E., Lusic, M., Arosio, D., Giacca, M., Cereseto, A., 2011. The TRIM Family

Protein KAP1 Inhibits HIV-1 Integration. Cell Host Microbe 9, 484–495.

https://doi.org/10.1016/j.chom.2011.05.004

Aston, W.J., Hope, D.E., Nowak, A.K., Robinson, B.W., Lake, R.A., Lesterhuis, W.J., 2017. A systematic

investigation of the maximum tolerated dose of cytotoxic chemotherapy with and without

supportive care in mice. BMC Cancer 17, 684. https://doi.org/10.1186/s12885-017-3677-7

Broce, S., Hensley, L., Sato, T., Lehrer-Graiwer, J., Essrich, C., Edwards, K.J., Pajda, J., Davis, C.J.,

Bhadresh, R., Hurt, C.R., Freeman, B., Lingappa, V.R., Kelleher, C.A., Karpuj, M.V., 2016.

Biochemical and biophysical characterization of cell-free synthesized Rift Valley fever virus

nucleoprotein capsids enables in vitro screening to identify novel antivirals. Biol. Direct 11, 25.

https://doi.org/10.1186/s13062-016-0126-5

Cirone, M., 2018. EBV and KSHV Infection Dysregulates Autophagy to Optimize Viral Replication, Prevent

Immune Recognition and Promote Tumorigenesis. Viruses 10, 599.

https://doi.org/10.3390/v10110599

Copley, S.D., 2012. Moonlighting is mainstream: Paradigm adjustment required. BioEssays 34, 578–588.

https://doi.org/10.1002/bies.201100191

Cui, Y., Yang, S., Fu, X., Feng, J., Xu, S., Ying, G., 2014. High Levels of KAP1 Expression Are Associated with

Aggressive Clinical Features in Ovarian Cancer. Int. J. Mol. Sci. 16, 363–377.

https://doi.org/10.3390/ijms16010363

Falzone, L., Salomone, S., Libra, M., 2018. Evolution of Cancer Pharmacological Treatments at the Turn

of the Third Millennium. Front. Pharmacol. 9, 1300. https://doi.org/10.3389/fphar.2018.01300

Fong, K., Zhao, J.C., Song, B., Zheng, B., Yu, J., 2018. TRIM28 protects TRIM24 from SPOP-mediated

degradation and promotes prostate cancer progression. Nat. Commun. 9, 5007.

https://doi.org/10.1038/s41467-018-07475-5

Gad, S.C., 2014. Maximum Tolerated Dose, in: Encyclopedia of Toxicology. Elsevier, p. 164.

https://doi.org/10.1016/B978-0-12-386454-3.00874-5

Hanahan, D., Weinberg, R.A., 2011. Hallmarks of Cancer: The Next Generation. Cell 144, 646–674.

https://doi.org/10.1016/j.cell.2011.02.013

Hanahan, D., Weinberg, R.A., 2000. The Hallmarks of Cancer. Cell 100, 57–70.

https://doi.org/10.1016/S0092-8674(00)81683-9

Huang, J., Huang, Q., Zhou, X., Shen, M.M., Yen, A., Yu, S.X., Dong, G., Qu, K., Huang, P., Anderson, E.M.,

Daniel-Issakani, S., Buller, R.M.L., Payan, D.G., Lu, H.H., 2004. The Poxvirus p28 Virulence Factor

Is an E3 Ubiquitin Ligase. J. Biol. Chem. 279, 54110–54116.

https://doi.org/10.1074/jbc.M410583200

Hughes, J., Rees, S., Kalindjian, S., Philpott, K., 2011. Principles of early drug discovery: Principles of early

drug discovery. Br. J. Pharmacol. 162, 1239–1249. https://doi.org/10.1111/j.1476-

5381.2010.01127.x

Igawa, T., Lin, F.-F., Lee, M.-S., Karan, D., Batra, S.K., Lin, M.-F., 2002. Establishment and characterization

of androgen-independent human prostate cancer LNCaP cell model. The Prostate 50, 222–235.

https://doi.org/10.1002/pros.10054

Ison, M.G., 2011. Antivirals and resistance: influenza virus. Curr. Opin. Virol. 1, 563–573.

https://doi.org/10.1016/j.coviro.2011.09.002

Iyengar, S., Farnham, P.J., 2011. KAP1 Protein: An Enigmatic Master Regulator of the Genome. J. Biol.

Chem. 286, 26267–26276. https://doi.org/10.1074/jbc.R111.252569

Javier, R.T., Butel, J.S., 2008. The History of Tumor Virology. Cancer Res. 68, 7693–7706.

https://doi.org/10.1158/0008-5472.CAN-08-3301

Jeffery, C.J., 2019. Multitalented actors inside and outside the cell: recent discoveries add to the number

of moonlighting proteins. Biochem. Soc. Trans. 47, 1941–1948.

https://doi.org/10.1042/BST20190798

Jeffery, C.J., 2018. Protein moonlighting: what is it, and why is it important? Philos. Trans. R. Soc. B Biol.

Sci. 373, 20160523. https://doi.org/10.1098/rstb.2016.0523

Jin, X., Pan, Y., Wang, L., Zhang, L., Ravichandran, R., Potts, P.R., Jiang, J., Wu, H., Huang, H., 2017.

MAGE-TRIM28 complex promotes the Warburg effect and hepatocellular carcinoma progression

by targeting FBP1 for degradation. Oncogenesis 6, e312–e312.

https://doi.org/10.1038/oncsis.2017.21

Kaján, G.L., Doszpoly, A., Tarján, Z.L., Vidovszky, M.Z., Papp, T., 2020. Virus–Host Coevolution with a

Focus on Animal and Human DNA Viruses. J. Mol. Evol. 88, 41–56.

https://doi.org/10.1007/s00239-019-09913-4

Krump, N.A., You, J., 2018. Molecular mechanisms of viral oncogenesis in humans. Nat. Rev. Microbiol.

16, 684–698. https://doi.org/10.1038/s41579-018-0064-6

Lingappa, U.F., Wu, X., Macieik, A., Yu, S.F., Atuegbu, A., Corpuz, M., Francis, J., Nichols, C., Calayag, A.,

Shi, H., Ellison, J.A., Harrell, E.K.T., Asundi, V., Lingappa, J.R., Prasad, M.D., Lipkin, W.I., Dey, D.,

Hurt, C.R., Lingappa, V.R., Hansen, W.J., Rupprecht, C.E., 2013. Host–rabies virus protein–protein

interactions as druggable antiviral targets. Proc. Natl. Acad. Sci. 110.

https://doi.org/10.1073/pnas.1210198110

Lingappa, V., Hurt, C., Garvey, E., 2013. Capsid Assembly as a Point of Intervention for Novel Anti-viral

Therapeutics. Curr. Pharm. Biotechnol. 14, 513–523.

https://doi.org/10.2174/13892010113149990201

MacKinnon, A.L., Taunton, J., 2009. Target Identification by Diazirine Photo-Cross-Linking and Click

Chemistry. Curr. Protoc. Chem. Biol. 1, 55–73.

https://doi.org/10.1002/9780470559277.ch090167

Malovannaya, A., Lanz, R.B., Jung, S.Y., Bulynko, Y., Le, N.T., Chan, D.W., Ding, C., Shi, Y., Yucer, N.,

Krenciute, G., Kim, B.-J., Li, C., Chen, R., Li, W., Wang, Y., O'Malley, B.W., Qin, J., 2011. Analysis of

the Human Endogenous Coregulator Complexome. Cell 145, 787–799.

https://doi.org/10.1016/j.cell.2011.05.006

Morales-Sánchez, A., Fuentes-Pananá, E., 2014. Human Viruses and Cancer. Viruses 6, 4047–4079.

https://doi.org/10.3390/v6104047

Motlagh, H.N., Hilser, V.J., 2012. Agonism/antagonism switching in allosteric ensembles. Proc.
Natl. Acad. Sci. U. S. A. 109, 4134–4139. https://doi.org/10.1073/pnas.1120519109

1011

Müller-Schiffmann, A., Michon, M., Lingappa, A.F., Yu, S.F., Du, L., Deiter, F., Broce, S., Mallesh, S.,

Crabtree, J., Lingappa, U.F., Macieik, A., Müller, L., Ostermann, P.N., Andrée, M., Adams, O.,

Schaal, H., Hogan, R.J., Tripp, R.A., Appaiah, U., Anand, S.K., Campi, T.W., Ford, M.J., Reed, J.C.,

Lin, J., Akintunde, O., Copeland, K., Nichols, C., Petrouski, E., Moreira, A.R., Jiang, I., DeYarman,

N., Brown, I., Lau, S., Segal, I., Goldsmith, D., Hong, S., Asundi, V., Briggs, E.M., Phyo, N.S.,

Froehlich, M., Onisko, B., Matlack, K., Dey, D., Lingappa, J.R., Prasad, M.D., Kitaygorodskyy, A.,

Solas, D., Boushey, H., Greenland, J., Pillai, S., Lo, M.K., Montgomery, J.M., Spiropoulou, C.F.,

Korth, C., Selvarajah, S., Paulvannan, K., Lingappa, V.R., 2022. A Pan-respiratory Antiviral

Chemotype Targeting a Transient Host Multiprotein Complex (preprint). Biochemistry.

https://doi.org/10.1101/2021.01.17.426875

Müller-Schiffmann, A., Torres, F., Kitaygorodskyy, A., Ramani, A., Alatza, A., Tschirner, S., Prikulis, I., Yu,

S., Dey, D., Bader, V., Rozemuller, A., Wray, S., Gopalakrishnan, J., Riek, R., Lingappa, V.R., Korth,

C., 2021a. Macrophage migration inhibitory factor is a valid drug target at the intersection of

herpes simplex virus 1 replication and Alzheimer's disease-relevant cellular pathology (preprint).

Microbiology. https://doi.org/10.1101/2021.09.11.459903

Müller-Schiffmann, A., Trossbach, S.V., Lingappa, V.R., Korth, C., 2021b. Viruses as 'Truffle Hounds':

Molecular Tools for Untangling Brain Cellular Pathology. Trends Neurosci. 44, 352–365.

https://doi.org/10.1016/j.tins.2020.11.004

Neo, S.H., Itahana, Y., Alagu, J., Kitagawa, M., Guo, A.K., Lee, S.H., Tang, K., Itahana, K., 2015. TRIM28 Is

an E3 Ligase for ARF-Mediated NPM1/B23 SUMOylation That Represses Centrosome

Amplification. Mol. Cell. Biol. 35, 2851–2863. https://doi.org/10.1128/MCB.01064-14

OncoPanelTM Cell-Based Profiling Service Details, n.d. URL

https://www.eurofinsdiscoveryservices.com/services/phenotypic-assays/oncopanel/oncopanel-

details (accessed 8.9.22).

O'Shea, C.C., 2005. Viruses: tools for tumor target discovery, and agents for oncolytic therapies – an

introduction. Oncogene 24, 7636–7639. https://doi.org/10.1038/sj.onc.1209035

Pfeffer, C., Singh, A., 2018. Apoptosis: A Target for Anticancer Therapy. Int. J. Mol. Sci. 19, 448.

https://doi.org/10.3390/ijms19020448

Priyamvada, L., Burgado, J., Baker-Wagner, M., Kitaygorodskyy, A., Olson, V., Lingappa, V.R.,

Satheshkumar, P.S., 2021. New methylene blue derivatives suggest novel anti-orthopoxviral

strategies. Antiviral Res. 191, 105086. https://doi.org/10.1016/j.antiviral.2021.105086

Pucci, B., Kasten, M., Giordano, A., 2000. Cell Cycle and Apoptosis. Neoplasia 2, 291–299.

https://doi.org/10.1038/sj.neo.7900101

Randolph, K., Hyder, U., D'Orso, I., 2022. KAP1/TRIM28: Transcriptional Activator and/or Repressor of

Viral and Cellular Programs? Front. Cell. Infect. Microbiol. 12, 834636.

https://doi.org/10.3389/fcimb.2022.834636

Reed, J.C., Solas, D., Kitaygorodskyy, A., Freeman, B., Ressler, D.T.B., Phuong, D.J., Swain, J.V., Matlack,

K., Hurt, C.R., Lingappa, V.R., Lingappa, J.R., 2021. Identification of an Antiretroviral Small

Molecule That Appears To Be a Host-Targeting Inhibitor of HIV-1 Assembly. J. Virol. 95, e00883-

20. https://doi.org/10.1128/JVI.00883-20

Reference ID" 4433223- Accessdata.fda.gov, n.d. URL

https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/020509s082lbl.pdf (accessed

8.9.22).

Reichel, A., Lienau, P., 2015. Pharmacokinetics in Drug Discovery: An Exposure-Centred Approach to

Optimising and Predicting Drug Efficacy and Safety, in: Nielsch, U., Fuhrmann, U., Jaroch, S.

(Eds.), New Approaches to Drug Discovery, Handbook of Experimental Pharmacology. Springer

International Publishing, Cham, pp. 235–260. https://doi.org/10.1007/164_2015_26

Singh, S., Singh, P., 2018. Pattern and impact of drugs targeted toward toxicity amelioration in patients

receiving cancer chemotherapy. Perspect. Clin. Res. 9, 23.

https://doi.org/10.4103/picr.PICR_156_16

Tanaka, A., 2009. Identification of the Specific Binding Proteins of Bioactive Small Compound Using

Affinity Resins, in: Koga, H. (Ed.), Reverse Chemical Genetics, Methods in Molecular Biology.

Humana Press, Totowa, NJ, pp. 181–195. https://doi.org/10.1007/978-1-60761-232-2_14

Usman, R.M., Razzaq, F., Akbar, A., Farooqui, A.A., Iftikhar, A., Latif, A., Hassan, H., Zhao, J., Carew, J.S.,

Nawrocki, S.T., Anwer, F., 2021. Role and mechanism of autophagy-regulating factors in

tumorigenesis and drug resistance. Asia Pac. J. Clin. Oncol. 17, 193–208.

https://doi.org/10.1111/ajco.13449

Wang, C., Ivanov, A., Chen, L., Fredericks, W.J., Seto, E., Rauscher, F.J., Chen, J., 2005. MDM2 interaction

with nuclear corepressor KAP1 contributes to p53 inactivation. EMBO J. 24, 3279–3290.

https://doi.org/10.1038/sj.emboj.7600791

Wei, C., Cheng, J., Zhou, B., Zhu, L., Khan, Md.A., He, T., Zhou, S., He, J., Lu, X., Chen, H., Zhang, D., Zhao,

Y., Fu, J., 2016. Tripartite motif containing 28 (TRIM28) promotes breast cancer metastasis by

stabilizing TWIST1 protein. Sci. Rep. 6, 29822. https://doi.org/10.1038/srep29822

Yang, Y., Fiskus, W., Yong, B., Atadja, P., Takahashi, Y., Pandita, T.K., Wang, H.-G., Bhalla, K.N., 2013.

Acetylated hsp70 and KAP1-mediated Vps34 SUMOylation is required for autophagosome

creation in autophagy. Proc. Natl. Acad. Sci. 110, 6841–6846.

https://doi.org/10.1073/pnas.1217692110

Yu, C., Zhan, L., Jiang, J., Pan, Y., Zhang, H., Li, X., Pen, F., Wang, M., Qin, R., Sun, C., 2014. KAP-1 is

overexpressed and correlates with increased metastatic ability and tumorigenicity in pancreatic

cancer. Med. Oncol. 31, 25. https://doi.org/10.1007/s12032-014-0025-5

1082

1083