Maternal cortisol and emotional complaints alter breast milk's microbiome composition

Nadia Deflorin[1], Ulrike Ehlert[1] Rita T. Amiel Castro[1] *

[1] Department of Clinical Psychology and Psychotherapy, Institute of Psychology, University of Zurich, Zurich, Switzerland

* Corresponding author: Dr. Rita Amiel Castro, University of Zurich, Department of Clinical Psychology and Psychotherapy, Binzmühlestrasse 14, 8050 Zurich, Switzerland. *E-mail address:* r.castro@psychologie.uzh.ch.

Data Share Statement: Data described in the manuscript, code book, and analytic code will be made available from the corresponding author upon reasonable request. Raw sequencing data for all 100 samples described in this project have been deposited in The Open Science Framework (OSF) under osf.io/r687v.

Conflicts of interest: The authors have no conflicts of interest to declare.

Funding: This work was supported by the Maiores Stiftung. The funding source had no role in the study design, execution, analyses, data interpretation, or decision to submit results.

Abbreviations:

ASV	amplicon sequence variant
AUCg	area under the curve with respect to the ground
BH	Benjamini-Hochberg
BM	Breast milk
dbRDA	distance-based redundancy analysis
EPDS	Edinburgh Postnatal Depression Scale
FDR	False discovery rate
GAS	Geburts-Angst-Skala
GHQ-12	General Health Questionnaire
HPA axis	hypothalamic-pituitary-adrenal axis
NuPDQ	Prenatal Distress Questionnaire
PSS-10	Perceived Stress Scale
STAI-S	State-Trate Anxiety Inventory, state scale

1 Abstract

Breast milk (BM) is considered the "gold standard" of nutrition due to its many benefits for

the infant. On the other hand, the challenges of pregnancy can lead to increased stress in some

women, which may affect BM quality. Although studies have demonstrated a negative link

between maternal psychopathology and child development, it remains unclear how maternal

psychobiological changes can be intergenerationally transmitted. **Aims**: We investigated the

associations between maternal biological stress, depressive symptoms, self-reported stress and

anxiety symptoms and the BM microbiome. Further, we analysed these parameters in relation

to BM glucocorticoid concentrations and explored the influence of BM glucocorticoids on

BM bacterial composition. **Methods**: N=100 women completed standardised questionnaires

(e.g. EPDS, STAI, GAS) at 34-36 weeks gestation and in the early postpartum; and sampled

saliva at 34-36 and 38 weeks gestation. BM samples were collected in the early postpartum.

Microbiota was analysed using 16S rRNA amplicon sequencing. **Results**: Prenatal GAS and

pregnancy-related symptoms (sum) were negatively correlated with *Alphaproteobacteria* (τ =

- 0.2137, FDR = 0.0199*; τ = - 0.1805 FDR = 0.0798*), whereas in the postpartum period,

the STAI-S scores were negatively correlated with different taxa. Postpartum-related

symptoms (sum) were also linked to decreased *Propionibacteriales*. Salivary cortisol AUCg

at 34-36 weeks was negatively correlated with *Actinomycetia* (τ = -0.1633, FDR = 0.0630*)

and *Clostridia* (τ = -0.1824, FDR = 0.0630*). BM cortisol was negatively correlated with

Staphylococcus (τ = - 0.2047, FDR = 0.0885*) and positively correlated with BM alpha

diversity. No associations emerged between psychobiological parameters and BM

glucocorticoids. **Conclusions**: Increased perinatal emotional complaints and prenatal cortisol

AUCg were associated with decreased commensal bacteria, whereas enhanced BM cortisol

was linked to increased alpha diversity and reduced *Staphylococcus*. These findings suggest a

negative relation between high maternal psychobiological complaints and commensal milk

microbiota.

Keywords: prenatal cortisol, prenatal emotional complaints, breast milk glucocorticoids,

breast milk microbiota

1. Introduction

Breast milk (BM) is considered the "gold standard" of nutrition due to its many benefits for the infant, including fewer respiratory infections and a lower incidence of obesity, type 1 and 2 diabetes mellitus, and gastroenteritis [1, 2]. BM composition is dynamic, highly complex and varies within one feeding, through the course of the day, across lactation periods, and between populations [3, 4]. BM has long been considered sterile, yet recent studies have shown it to be a rich source of commensal microbes that can positively influence the child's health [5]. For instance, in a recent review [6], the BM microbiome was found to be quite diverse, containing more than 800 bacterial species. *Serratia, Ralstonia, Propionibacterium, Pseudomonas, Staphylococcus, Streptococcus, Corynebacterium, Sphingomonas and Bradyrhizobium* represent the nine genera that form the core BM microbiome [5, 7] as they constitute approximately half of its microbial community [8, 9]. Evidence suggests that BM is not solely a passive reflection of maternal bioactive substances, but is rather an actively regulated secretion that may program the infant's cognitive and immunological development [10, 11]. BM microbial community reaches the child's gut via breastfeeding, influencing its colonisation [12, 13], and can affect several aspects of newborn health [12, 14–17]. Interestingly, it has recently been demonstrated that BM bacterial, hormonal and immunological composition may be negatively altered by maternal psychopathology [11, 18].

The changes and challenges of pregnancy can lead to increased stress in some women [19–22], which may in turn negatively affect BM quality [18, 23, 24]. Other determinants that have been found to interact with BM composition include maternal age, parity, body mass index (BMI), diet, use of antibiotics, and mode of delivery [5]. Although there is a considerable body of research demonstrating a negative link between perinatal maternal

psychopathology and child development and health [11, 18, 25, 26], it remains unclear how

maternal psychobiological complaints can be intergenerationally transmitted [27]. One strong

candidate in this regard is BM. BM composition can be shaped by multiple factors such as

maternal body index and diet [11]. The development of the maternal BM microbiome begins

in the third trimester of pregnancy [28, 29], with bacteria from the maternal gut probably

entering the mammary gland through bacterial translocation [30–32]. This potential

microbiome transfer route is known as the entero-mammary pathway [12, 29, 32] and can be

influenced, among others, by maternal emotional complaints [16, 33, 34] via the microbiome-

gut-brain axis [35]. The latter describes a bidirectional interaction occurring between the

gastrointestinal tract, the gut microbiome, and the central nervous system [35].

Communication pathways of this axis include vagal nerve activation, neurotransmitters,

metabolites, as well as the immune system [16]. Consistent with these pathways, experimental

studies also suggest a bidirectional communication between the gut microbiome and the

hypothalamic-pituitary-adrenal (HPA) axis [36].

The HPA axis is the main system controlling stress responses [37], in which the stress

hormone cortisol is produced by the adrenal glands in response to physiological and

psychological stress [38, 39]. Cortisol released in the systemic circulation appears to be the

primary source of BM glucocorticoids, as BM cortisol was shown to be strongly correlated

with plasma cortisol concentrations [25, 40]. Maternal stress is a substantial factor influencing

higher cortisol concentrations in BM [24, 41–43]. Although BM glucocorticoids were first

detected in the 1970s [44], few studies to date have looked at their effects on the offspring

[25].

Given that BM is an underexplored pathway underlying the relation between perinatal

emotional complaints and the offspring, the aim of the present study is threefold. First, we

investigated the associations between maternal perinatal biological stress, depressive

symptoms, self-reported stress, and anxiety symptoms and the BM microbiome. Second, we

analysed these parameters in relation to BM glucocorticoid concentrations. Finally, we

explored the influence of BM glucocorticoids concentrations on BM bacterial composition.

To the best of our knowledge, this is the first study to comprehensively examine the interplay

between the BM microbiome and glucocorticoid concentrations in colostrum (the initial BM)

in relation to maternal perinatal mental health. This is relevant to improve our understanding

of the interactions between maternal emotional complaints, endocrinological parameters and

bacterial composition [5, 10, 11]. We hypothesise that women with higher levels of perinatal

emotional complaints and biological stress will show lower bacterial diversity and less

commensal bacteria as well as elevated BM glucocorticoid concentrations. Additionally, we

hypothesise that higher levels of BM cortisol can negatively alter the BM microbiome.

2. Materials and Methods

From January 2021 to January 2022, we recruited women in the German-speaking part

of Switzerland through online platforms, mailing lists, social media, the study's homepage as

well as flyers handed to patients receiving prenatal medical care at gynaecology practices in

and around Zurich and Aargau. To meet the inclusion criteria, women had to be between 18

and 45 years old, currently in the 3rd gestational trimester, fluent in German, of European

origin and physically healthy according to self-report. Women were excluded if they had a

multifoetal gestation, a pregnancy through assisted reproductive technology, medical

complications and/or surgical interventions that might have affected ovarian function prior to

pregnancy (e.g., polycystic ovary syndrome) or a planned caesarean section. Additional

exclusion criteria were current intake of hormones, psychotropic substances, drugs (e.g.

cocaine, marijuana) and antibiotics. A total of $N = 119$ prenatal women and their children

were included in this prospective cross-sectional study.

The procedures followed were in accordance with the ethical standards of the Zurich

Cantonal Ethics Committee on human experimentation and approval was obtained prior to

study commencement. Ethical approval was granted by the Zurich Cantonal Ethics

Committee (KEK-ZH-Nr. 2020-01928) and the study was conducted at the Psychological

Institute of the University of Zurich in accordance with the Declaration of Helsinki.

The inclusion and exclusion criteria were screened using an online questionnaire.

Women who met the inclusion criteria according to this questionnaire then received the study

information and consent forms via postal mail. Next, a telephone interview was conducted

with interested participants, in which the inclusion criteria were re-checked and the Structured

Clinical Interview for DSM-5 (SCID-5-CV) was administered. Eligible participants were

invited to the first laboratory appointment between the 34th and 36th week of gestation (T1),

conducted either as a home visit or via videotelephony, depending on the respective COVID-

19 restrictions at the time. Before T1, participants were asked to sign the consent forms. T1

followed a standardised protocol. First, the procedure was explained and maternal

anthropometric measures were recorded. In the case of videotelephony appointments, this

information was taken either from the participants' maternity records or their most recent

gynaecological examination. Next, instructions regarding saliva sampling were provided, both

verbally and in writing, as participants were asked to independently collect saliva samples

during the study. Finally, any potential questions were clarified and participants were asked to

complete online questionnaires on socio-demographic, psychological and behavioural

information.

On the day after the first appointment (T1), participants were asked to collect saliva

and briefly reply to some questions on their perceived stress levels and sleep quality. Saliva

sampling took place on two consecutive days immediately after awakening, 30 min and 45

min after awakening, and before going to bed. The second saliva collection was scheduled for

the 38[th] week of pregnancy (T2), again accompanied by the same brief questionnaire as in T1.

Participants were asked to freeze the saliva samples in their home freezer. Within the first 120

hours after birth, preferably on the fourth day postpartum, the final study appointment (T3)

was conducted between 8:00 and 8:30 in the morning as a home or clinic visit. During this

visit, any remaining questions were clarified, the procedure was explained, and BM collection

was conducted. For the planned analyses, 5-15 millilitres of BM were collected using an

electric BM pump. At T3, we also collected the saliva samples that had been stored in the

participants' homes. Subsequently, participants were asked to complete questionnaires related

to birth information and some of the psychological parameters from T1.

2.1 Instruments

2.1.1   Maternal emotional complaints

Validated psychometric questionnaires were used to assess perinatal depressive

symptoms, anxiety symptoms, and self-reported stress.

We measured stress using the German version of the *Perceived Stress Scale* (PSS-10)

at T1 and T3 [45]. This is a self-report scale measuring the degree to which the respondent

perceives situations to be unpredictable, uncontrollable, and stressful referring to the past four

weeks [46]. The 10 items are rated on a five-point Likert scale (0 = "*never*" to 4 = "*very*

often"), with higher scores indicating a higher level of perceived stress [45]. Analyses were

conducted with the total score as a continuous variable. Cronbach's alpha in the present study

ranged between 0.823 (T3) and 0.828 (T1).

We used the German version of the *General Health Questionnaire* (GHQ-12) [47],

which is a reliable instrument to screen for psychological distress [48, 49]. The 12 items are

rated on a four-point Likert scale (0 = "*no, not at all*" to 4 = "*much harder than usual*") [50].

Higher scores indicate higher levels of mental distress [48]. In the present analyses, we used

the GHQ-12 total score as a continuous variable. Cronbach's alpha for this study ranged

between 0.780 (T3) and 0.808 (T1).

Pregnancy-specific stress at T1 was measured using the German version of the revised

Prenatal Distress Questionnaire (NuPDQ) [51]. Respondents are asked to indicate the extent

to which they feel "bothered, upset, or worried" about issues that become more relevant as

pregnancy progresses (e.g., concerns about caring for a newborn) [52], with items rated on a

three-point Likert scale (from 0 = "*not at all*" to 2 = "*very much*"). In the present study, the

total score was used as a continuous variable for statistical analyses. Cronbach's alpha was

.073.

The German version of the *Edinburgh Postnatal Depression Scale* (EPDS) was

applied to assess maternal depressive symptoms at T1 and T3 [53]. The questionnaire was

developed to assess depressive symptoms in pre- and postnatal women. It consists of 10 items

rated on a four-point Likert scale (0 = "*yes, most of the time*" to 3 = "*no, never*") relating to

how one has felt in the last seven days. In our analyses, we used the total score as a

continuous variable. Cronbach's alpha for this study ranged between 0.798 (T1) and 0.880

(T3).

Pregnancy/birth anxiety was assessed at T1 using the *Birth Anxiety Scale* (Geburts-

Angst-Skala, GAS) [54]. The scale comprises 25 items assessing the extent to which

respondents are afraid of certain aspects related to pregnancy and birth, rated on a four-point

Likert scale (0 = "no fear" to 3 = "strong fear") [55]. For the present analyses, the total score

was used as a continuous variable. Cronbach's alpha was 0.871.

Trait anxiety and state anxiety were measured using the German version of the State-

Trait Anxiety Inventory (STAI) [56, 57]. The State scale (STAI-S) addresses the level of

anxiety in a particular situation [56] and consists of 20 items rated on a four-point Likert scale

(1 = "*not at all*" to 4 = "*very much*"), with higher scores indicating higher state anxiety [58].

Cronbach's alpha for this scale ranged between 0.891 (T1) and 0.924 (T3). The Trait scale

(STAI-T) relates to the overall disposition to experience worries, fears, and anxieties in

different situations [59]. It likewise consists of 20 items rated on four-point Likert scale (1 =

"*almost never*" to 4 = "*almost always*"), and asks respondents how they feel in general.

Higher scores indicate higher trait anxiety. Cronbach's alpha for the current study ranged

between 0.890 (T3) and 0.897 (T1). Trait and State anxiety total scores were scored separately

and treated as continuous variables in our analyses.

Maternal emotional complaints can be considered as a multidimensional concept, with

women reporting various different symptoms (e.g., anxiety, stress, and/or depression) [18, 60,

61]. Therefore, we created summary variables from the questionnaires described above. To

capture pregnancy-specific emotional complaints, the questionnaires NuPDQ, GAS, and

EPDS were merged, as they all specifically target the prenatal period. To do so, we used the

total scores from each questionnaire at T1 and summed them together to create a pregnancy-

related sum score (continuous variable). Similarly, a multidimensional summary variable was

created to capture postpartum emotional complaints. Total scores derived from the GHQ-12,

PSS-10, EPDS, and STAI-S were combined into a postpartum sum score (continuous

variable), as they account for depressive symptoms, anxiety symptoms and self-reported

distress.

2.1.2   Maternal prenatal salivary cortisol

During the course of the study, participants were required to collect saliva samples for

the assessment of cortisol concentrations. Under standardised conditions, participants were

required to collect 8 saliva samples per time point (T1 and T2), corresponding to a total of 16

saliva samples per participant. The three morning and one evening samples were collected

using the passive drool method in SaliCap sampling tubes with 2 mL capacity (SaliCaps, IBL

international GmBH, Hamburg, Germany). For this purpose, the women were instructed to

collect saliva immediately after awakening, 30 min after awakening, 45 min after awakening

and before going to bed. After sampling, they were instructed to store the samples in their

freezers until the home or clinic visit (T3). The day before sampling, participants were asked

to avoid exercise, alcohol, coffee, black tea, and chewing gum. Additionally, they were asked

not to eat, smoke, or brush their teeth during the morning saliva collection sessions. After

completion of study participation, samples were transported by a member of the study team

using a cool box and stored at -20 degrees in the biochemical laboratory of the University of

Zurich. For subsequent cortisol analyses, samples were then thawed, centrifuged, and

analysed biochemically using an enzyme-linked immunosorbent assay (ELISA; IBL

international GmBH, Hamburg, Germany). Individual values beyond the range of the kit were

excluded from the analyses ($N = 2$).

2.1.3   Breast milk collection

Within the first 120 hours after birth, preferably on the fourth day postpartum, the final

study appointment (T3) was performed between 8:00 and 8:30 in the morning as a home or

clinic visit, as BM cortisol and cortisone concentrations follow the diurnal rhythm of the HPA

axis [62]. The method of BM collection was similar to that used by McGuire et al. (2017)

[63]. Accordingly, participants were asked to wash their hands and dry them with a disposable

towel provided by the study team. Both the participant and the study team member had to

disinfect their hands and put on disposable gloves. As the researcher had to unwrap the

materials and give them to the woman, this procedure ensured that the materials were as

sterile as possible. Once the appropriate BM pump set was selected, the breast was then

stimulated. Following this, the breast was cleaned twice with a newly opened pre-packaged

castile soap towelette (Pdi Inc. Pyd41900 Castile Soap Towelettes). We used an electric BM

pump (Medela Symphony) and disposable pump sets (Medela Inc.) to collect milk samples

into a single-use, sterile, 35 ml milk collection container (Medela Inc.). To ensure the best

possible milk flow, the breast chosen for pumping was the one that was not used during the

last feeding or that the participant felt to be fuller. If not enough milk was collected from one

breast, participants were asked to express milk from both breasts; in these cases, the

previously described cleaning procedure was repeated with the other breast.

For the BM microbiome sample, we collected 5-10 ml of BM in a milk collection

container. In a second container, a further 2-5 ml were expressed for the analyses of BM

glucocorticoids. Immediately after collection, for the endocrinological analyses, BM was

transferred from the 35 ml milk collection container into 2 ml microtubes (Sarstedt AG) by a

member of the study team using a sterile pipette. All samples were immediately placed in a

cool box to be transported by members of the research team to the biochemical laboratory at

the University of Zurich, where they were stored at -80 degrees until shipment. $N = 97$ BM

samples were collected, which amounted to 1.64 litres of BM. Samples for microbiome

analysis were shipped to the Clinical Microbiomics Laboratory in Copenhagen (Denmark).

Samples for cortisol and cortisone measurements were shipped to the Laboratory of

Endocrinology, University Medical Centers (UMC), Amsterdam (Netherlands).

2.2 BM cortisol and cortisone concentrations

To evaluate cortisol and cortisone concentrations in BM, isotope dilution liquid

chromatography-tandem mass spectrometry (LC-MS/MS) was used, as previously described

in Van der Voorn et al. (2015) [64]. Briefly, milk samples were washed with hexane after

internal standards (13C3-labelled cortisol and cortisone) were added to the samples. Samples

were then extracted with isolute plates (Biotage, Uppsala, Sweden) and analysed by LC-

MS/MS (Acquity with Quattro Premier XE, Milford MA, USA, Waters Corporation). The

intra-assay coefficient of variation (CV) was 4-5% for cortisol and 5 % for cortisone at

different values. The inter-assay coefficient of variation for both, cortisol and cortisone was <

9%, with the lower limit of quantitation of 0.5 nmol/L.

2.3 BM 16S ribosomal RNA (rRNA) sequencing and quality control

At the Clinical Microbiomics laboratory, DNA was extracted from ~5 mL aliquots of

the $N = 97$ BM samples. Prior to DNA extraction, cells from the aliquots were pelleted by

centrifugation (2 x 10 minutes at 6000 rpm). In a next step, DNA extraction from the pellets

was performed using the NucleoSpin 96 Soil (Macherey-Nagel) kit. Bead beating was

performed horizontally on a Vortex-Genie 2 at 2700 rpm for 5 minutes. Negative and positive

(mock) controls were incorporated into the analysis. Products from nested PCR were pooled

based on band intensity, before the resulting library was cleaned with magnetic beads. Pooled

library DNA concentration was measured fluorometrically. Sequencing was performed on an

Illumina MiSeq desktop sequencer using the MiSeq Reagent Kit V3 (Illumina) for 2x 300 bp

paired-end sequencing. We performed PCR using the forward primer S-D-Bact-0341-b-S-17

and the reverse primer S-D-Bact-0785-a-A-21 [65] with Illumina adapters attached. In fact,

these are universal bacterial 16S rDNA primers targeting the V3-V4 region. The following

PCR program was applied: 98 °C for 30 sec, 29x (98° C for 10 s, 55 °C for 20 s, 72 °C for 20

s), 72°C for 5 min. Amplification was verified by running the products on an agarose gel.

Indices were added in a subsequent PCR, utilising an Illumina Nextera kit with the

subsequent PCR program: 98 °C for 30 sec, 8x (98° C for 10 s, 55 °C for 20 s, 72 °C for 20 s),

72 °C for 5 min. Attachment of the indices was validated by testing the products on an

agarose gel. To account for variations in detection ability due to differences in sequencing

depth, all samples were rarefied to a fixed number of reads. Rarefied count and abundance

matrices were generated by random sampling of reads without replacement with a target

number of 4063 read counts. Any samples with a lower number of reads were excluded.

2.4 Statistical analysis

To define the minimum sample size required to test the study hypothesis, an a priori

power analysis was performed with G*Power version 3.1.9.7 [66]. All statistical analyses

were conducted with the IBM Statistical Package for the Social Sciences (SPSS version 28 for

Windows) and the software program R, using the adonis2 function from the vegan R package.

Normality of the sample was tested with the Shapiro-Wilk test, and non-parametric tests were

used for non-normally distributed data. For associative analysis with salivary cortisol

concentrations, we used the mean value of the area under the curve with respect to the ground

(AUCg) of the two consecutive days of saliva sampling. The AUCg was itself calculated

using the trapezoid formula, which is routinely calculated to include multiple time points [67].

Moreover, we used a mean value of total cortisol decline of the two consecutive days,

calculated as follows: Δ in the morning - Δ during the day, in which Δ in the morning

corresponds to the highest morning value - baseline morning value and Δ during the day

corresponds to the highest morning value - the evening value. Both procedures, the AUCg and

the total decline, were conducted separately for T1 and T2. Spearman's rank correlation was

used to examine correlations between and within psychological variables (PSS-10, GHQ-12,

NuPDQ, EPDS, GAS, STAI-S, pregnancy-related sum, postpartum sum), salivary cortisol and

BM glucocorticoid concentrations at all time points. Additionally, a Kruskal-Wallis test was

run to test for significant differences in BM glucocorticoid concentrations depending on the

extraction breast (e.g. right, left or both). Next, the cortisone-to-(cortisol + cortisone) ratio of

glucocorticoids in breast milk was calculated [62, 68, 69] in order to describe the activity of

11β-hydroxysteroid dehydrogenase type 2 (11β-HSD2) enzyme, which converts active

cortisol into inactive cortisone [68, 69]. The following formula was used for this purpose: BM

cortisone/BM cortisone + BM cortisol. Associations between the cortisone-to-(cortisol +

cortisone) ratio and the biological or psychological stress markers were evaluated using

Spearman's rank correlation. Both BM glucocorticoids and salivary cortisol results were

reported in nmol/l.

For the analysis of the BM microbiome, BM composition was first summarised in an

amplicon sequence variant (ASV) abundance table using a customised pipeline based on

dada2 [70]. The taxonomic classification of the detected ASVs was performed in two steps.

First, ASV sequences were compared with full-length 16S sequences in an internal reference

database (CM_16S_27Fto1492R_v1.0.0) using a naive Bayesian classifier. Then, we

optimised the taxonomic assignments based on the exact sequence identity percentages

between the discovered ASVs and the reference amplicons in an internal V3V4 amplicon

database (CM_16S_341Fto785R_v1.0.0.rds). Subsequently, we explored the diversity of the

BM microbiome communities within and between samples. For this purpose, bacterial alpha

diversity of the samples was calculated using the number of detected entities (richness) and

the Shannon index. Bray-Curtis dissimilarity and weighted UniFrac distances were used to

calculate bacterial beta diversity. Both alpha and beta diversity estimates were calculated from

rarefied abundance matrices. Additionally, when conducting statistical analyses, the

Benjamini-Hochberg (BH) method was applied to control for the false discovery rate (FDR)

at a level of 10% to control for multiple testing [71].

Associations for beta diversity – including Bray-Curtis dissimilarity and weighted

UniFrac distances as dependent variables and psychological variables as independent

variables – were performed using a distance-based redundancy analysis (dbRDA) test

conducted with 1000 permutations. We ran associative analyses with alpha diversity and

taxon abundance as dependent variables and psychological continuous variables as

independent variables using Kendall rank correlations. Unrarefied data were used for the

associations involving abundance of microbial taxons, for which 97 BM samples were

included. For the analyses of alpha and beta diversity parameters, we used 96 BM samples

analysed with rarefied data. This difference is due to the rarefaction method (see 2.5, BM

sequencing and quality control). Finally, to investigate the potential influence of several

covariates on the BM microbiome, PERMANOVA (for categorical variables) and dbRDA

(for continuous variables) were performed, following the approach suggested by Falony et al.,

2016 [72]. Only covariates that demonstrated a significant relationship with the BM

microbiome were further analysed for their predictive effects.

2.4.1   Covariates

Maternal age, parity, maternal BMI, the newborn's weight and length, and number of

hours between last feeding and BM collection were analysed as continuous variables.

Categorical variables were education (coded as 1 = university degree or higher and 0 = other)

and mode of delivery, considering membrane rupture during labour (coded as 0 = no and 1 =

yes). All covariates were identified based on previous literature [5, 12, 63, 73–76].

361 3. Results

362 3.1 Sample characteristics

Descriptive characteristics of the sample are shown in Table 1. Most of the

participants were married (66%) and highly educated, with 67% having a university degree.

71% of the women experienced a natural vaginal birth whereas 16% underwent an emergency

caesarean section. Of those who had a caesarean section, 43.75% ($N = 7$) had membrane

rupture prior to the intervention. The median time of BM sample collection was 97 hours after

birth (IQR = 30).

$N = 19$ participants voluntarily withdrew or were excluded from the study (e.g.

pregnancy complications), resulting in a final sample of 100 pregnant women (see Figure 1)

with an age range from 22 to 40 years ($MD = 32.50$, $IQR = 4.00$). Our power analysis [66]

suggested a required sample size of $N = 84$ to reach 80% power to detect a medium effect (f =

0.15), at a significance level of α = .05. Thus, the achieved sample size of $N = 100$ is

sufficient to test the study hypothesis. Moreover, we were unable to collect BM samples from

$N = 3$ participants, who had little or no milk (N=2) or were prevented from attending the final

appointment due to problems with the newborn baby (N=1).

379   [Add Figure 1 here]
380   [Add Table 1 here]
381

383 3.2 Analysis of the BM microbiome and BM glucocorticoids

For a taxonomic overview, Figure 1 and Figure 2 show the relative abundance of

bacterial taxa aggregated at the genus and family levels in the BM. We found that the BM

microbiome is dominated by the genera *Staphylococcus* (41.17%), *Streptococcus* (35.26%)

and *Gemella* (5.74%) (Figure 1). At the family level (Figure 2), we found that most of the

assigned BM taxa were from *Staphylococcaceae* (41.17%), *Streptococcaceae* (35.38%) and

Gemellaceae (5.74%). Both *Staphylococcus* as well as *Streptococcus* belong to the previously

described core BM microbiome [5–7]. Overall, the 97 BM samples had an average of 107.5

thousand (K) read pairs per sample and a minimum of 0.3 K read pairs. An average of 52.4 K

read pairs per sample could be mapped to the ASV catalogue, representing an average of 48.1

% of the high-quality reads.

395   [Add Figure 2 here]

397   [Add Figure 3 here]

At postpartum, a total of 423.5 millilitres of BM for glucocorticoid analyses were

obtained from all 97 participants. The median BM cortisol concentration was 11.16 nmol/l

(IQR = 9.12) whereas the median BM cortisone concentration was 52.96 nmol/l (IQR =

17.53). Furthermore, a significant positive correlation between these two concentrations was

found (rho = .283**, *p* = .005). Interestingly, there were no significant differences in BM

cortisol (*H* = 5.376, *p* = .068) and cortisone (*H* = 1.297, *p* = .532) concentrations based on

which breast was selected for sampling. The median of the BM cortisone-to-(cortisol +

cortisone) ratio was 0.803 nmol/l with an IQR of 0.100.

3.3 Association of maternal emotional complaints with the BM microbiome

In our initial analyses, no significant associations were found between any of the

covariates and BM microbial composition (*p* > 0.05). Therefore, no covariates were included

in the associative analyses. We explored BM bacterial composition in relation to perinatal

maternal emotional complaints from class to species level. After applying the Benjamini-

Hochberg (BH) method to control for the FDR [71] at a level of 10%, we found that prenatal

GAS and pregnancy-related sum were significantly negatively correlated with the class of

Alphaproteobacteria (τ = - 0.2137, FDR = 0.0199*; τ = - 0.1805 FDR = 0.0798*). Moreover,

in the postpartum period (T3), the class *Alphaproteobacteria*, the order *Propionibacteriales*,

as well as the genera *Cutibacterium* and *Pseudomonas_*N were negatively correlated only

with the STAI-S values and not with the STAI-T (see Table 2). However, no significant

associations were identified between prenatal and postnatal maternal emotional complaints

and BM alpha (Shannon diversity index range: τ = - 0.0671, FDR = 0.359 to τ = 0.0006, FDR

= 0.994) or beta diversity (Bray-Curtis dissimilarity range: Sum of Squares = 0.3812, Pr(>F)

= 0.295 to Sum of Squares = 0.1757, Pr(>F) = 0.999; weighted UniFrac distances range: Sum

of Squares = 0.1808, Pr(>F) = 0.305 to Sum of Squares = 0.0925, Pr(>F) = 0.903).

425   [Add Table 2 here]

3.4 Associations between salivary cortisol concentrations and the BM microbiome

We examined the associations of salivary cortisol concentrations at T1 and T2,

measured by AUCg and total decline, with each of the taxa found (from class to species

level). Again, the Benjamini-Hochberg (BH) method was applied to control for the FDR [71].

Among others, AUCg at T1 was negatively correlated with *Actinomycetia* (τ = -0.1633, FDR

= 0.0630*) and *Clostridia* (τ = -0.1824, FDR = 0.0630*). No associations were found between

total decline and any bacterial taxa (p > 0.05). Moreover, no significant associations emerged

between BM alpha and beta diversity in relation to salivary cortisol concentrations (p > 0.05).

All significant results of the BM microbiome in relation to salivary cortisol concentrations

can be seen in Table 3.

438   [Add Table 3 here]

3.5 Associations of maternal emotional complaints and biological stress with BM

glucocorticoids

We found significant correlations between and within T1 and T3 for the PSS-10,

GHQ-12, EPDS, and STAI-S ($p < .001$**). Spearman's rho analyses between the PSS-10,

GHQ-12, NuPDQ, EPDS, GAS, STAI-S, pregnancy-related sum at T1 and BM cortisol and

cortisone concentrations yielded null findings ($p > .05$). Similarly, no significant associations

were found between BM glucocorticoids and the GHQ-12, PSS-10, EPDS, STAI-S, and

postpartum sum at T3 ($p > .05$).

Spearman's rho correlations between T1 and T2 for each salivary cortisol

measurement (maternal salivary cortisol AUCg: rho = .706**, $p < .001$; total decline: rho =

.507**, $p < .001$) were significant. However, both salivary cortisol AUCg and total decline at

T1 and T2 were not significantly associated with any psychological parameters at T1 and T3

($p > .05$). Similarly, no significant results were found between the maternal salivary cortisol

AUCg and total decline measurements and BM glucocorticoids ($p > .05$).

The BM cortisone-to-(cortisol + cortisone) ratio was negatively correlated with the

amount of BM cortisol (rho = - .914**, $p < .001$) and not associated with BM cortisone ($p >$

.05). This suggests a decreased conversion of cortisol to cortisone in initial breast milk.

Moreover, we did not find any significant associations between salivary cortisol (AUCg and

total decline at T1 or T2), maternal emotional complaints (at T1 and T3), and the BM

cortisone-to-(cortisol + cortisone) ratio. All p-values were > .05.

3.6 Relationship between the BM microbiome and BM glucocorticoids

Kendall's rank correlations indicated that BM cortisol concentration was significantly

associated with BM alpha diversity ($\tau = 0.1671$, FDR = 0.0352*). After adjusting for the FDR

[71], BM cortisol was significantly correlated with the family *Staphylococcaceae* (τ = -

0.2047, FDR = 0.0577*) and the genus *Staphylococcus* (τ = - 0.2047, FDR = 0.0885*; see

Table 4).

Conversely, no significant associations were found between BM cortisone concentrations

and alpha diversity (Shannon diversity index; τ = - 0.0535, FDR = 0.4417, richness; τ = -

0.0595, FDR = 0.4417) or individual BM microbiome taxa (FDR > 0.1). Moreover, both BM

cortisol and cortisone showed no significant correlation with the beta diversity of the BM

microbiome at T3 (*p* > 0.05).

[Add Table 4 here]

4. Discussion

In this cross-sectional study involving perinatal women, we investigated whether there

are any associations between maternal perinatal biological stress, perinatal emotional

complaints, and the BM microbiome. Moreover, we analysed psychobiological parameters in

relation to BM glucocorticoids and explored the influence of BM glucocorticoid levels on BM

bacterial composition. Our main findings suggest that: the BM microbiome is dominated by

the genera *Staphylococcus*, *Streptococcus* and *Gemella;* increased perinatal maternal

emotional complaints and salivary AUCg are associated with decreased specific bacterial

taxa; and enhanced BM cortisol is linked to alpha diversity and *Staphylococcus.* No

associations were found between psychobiological parameters and BM glucocorticoids.

Our results support the view that *Staphylococcus* and *Streptococcus* are predominant

bacterial species in BM [77, 78] and reinforce the finding that BM can be an important source

of protective bacteria to the infant gut [79]. *Staphylococcus* and *Streptococcus* represented,

respectively, 41.1% and 35.2% of the total bacterial isolates in our sample and have been

previously identified in stool samples of breast-fed infants [80, 81]. Human milk contains

commensal bacteria, mainly non-pathogenic, which compete for nutrients and prevent growth

of pathogens by producing acid, hydrogen peroxide or bacteriocins [82]. Conversely, whereas

previous studies have identified *Lactobacillus* and *Bifidobacteria* - often considered as part of

a healthy microbiota - as common members of BM microbiota [83], we observed very few

sequences from these microbes. This difference may be attributable to geographical or

environmental differences among the studies, use of colostrum instead of transitional or

mature milk, differences in sample processing, analytical techniques, or diet [9].

Studies in both humans and animal models indicate that maternal psychosocial distress

is related to negative alterations in gut microbiota [84–87]. This is partially in line with our

findings, since breastfeeding may adversely modify infant gut microbiota and BM is

recognised as one of the most important elements modulating a child's gut health [88]. We

demonstrated that increased birth-specific anxiety and pregnancy-related psychological

symptoms (sum) were associated with a decrease in BM *Alphaproteobacteria*, whereas

heightened postpartum anxiety symptoms and postpartum psychological symptoms (sum)

were linked to decreased *Alphaproteobacteria*, *Propionibacteriales*, *Cutibacterium* and

Pseudomonas. All of these bacterial communities, although not among the most abundant

genera in our BM samples, have been previously reported as members of the human milk or

skin microbiota inducing protective responses on the host's immune system [89, 90]. It was

shown that the presence of *Pseudomonas* in BM, alone or in combination with other bacteria,

does not define an unhealthy microbial state [91]. This finding suggests that higher maternal

emotional complaints contribute to the decrease of commensal, commonly found, gram-

positive bacteria in BM. Human milk is in itself a source of commensal bacteria. Commensal

strains generally represent ecologically important inhabitants of the human mucosal surfaces

[92]. Therefore, this negative alteration in BM microbiota might, at least in part, be driven by

increased maternal emotional complaints in pregnancy and postpartum. Importantly, however,

not all studies observed this association [90].

Our results revealed that enhanced salivary cortisol, measured by AUCg, was

associated with lower *Actinomycetia*, *Clostridia*, *Stenotrophomonas, and Kocuria* whereas

increased BM cortisol was related to decreased *Staphylococcus*. All these bacteria exert a

great impact on regulation of immunophysiological functions, including metabolism and

pathogen defense [90, 91]. Here too, we see negative effects of maternal prenatal and

postnatal emotional complaints in terms of altering BM microbial composition, and

particularly in reducing commensal bacteria. This finding is novel, as prenatal salivary

cortisol and BM cortisol have not been previously addressed as potential factors that may

influence milk bacterial composition. Notably, we observed a link between higher BM

cortisol and increased alpha diversity and richness. Microbiota diversity plays an important

role in the maintenance of human immune balance while promoting intestinal immune system

maturation [93]. Our finding conflicts with the only previous study to have addressed the

effects of postpartum psychosocial stress on BM microbiota, which reported high milk

diversity in women with lower psychosocial stress levels [18]. The few other studies assessing

the effects of self-reported stress described similar findings in relation to infant gut microbiota

diversity [94, 95]. We are the first to publish data specifically targeting BM glucocorticoid

effects on BM bacterial composition, thus precluding any direct comparison with previous

works. The actual impact of the observed BM alterations remains uncertain, since different

taxonomic profiles may retain microbiota functionality due to metabolic plasticity [96].

Future studies examining BM cortisol alterations in BM bacteriome in clinical and non-

clinical populations may increase our understanding of the mechanisms linking maternal

mental health and BM microbiota.

Finally, we did not find any significant associations between maternal perinatal

emotional complaints, salivary cortisol and BM glucocorticoids. Although BM cortisol was

previously shown to be strongly correlated with plasma cortisol concentrations and maternal

emotional complaints [25, 40], this finding was not replicated in our sample. It is important to

acknowledge that some studies also failed to reveal an association between psychological and

biological stress during the perinatal phase. For instance, in a single assessment at 28 weeks

gestation, Wadhwa and colleagues (1996) [97] found no association between maternal plasma

cortisol and stressful life events, perceived stress, or self-reported pregnancy anxiety.

Similarly, Shea et al. (2007) [98] examined the cortisol awakening response in pregnant

women at 20 weeks gestation and reported no significant associations with self-reported

depression or anxiety. Finally, other researchers [99, 100] reported that pregnant women with

high stress levels presented with lower cortisol morning levels and a flatter diurnal decline in

late pregnancy, with associations between psychological stress and cortisol decline also being

observed previously [101]. In contrast, a recent study found that women with higher

psychosocial stress had increased levels of BM cortisol [18]. The underlying mechanisms of

these opposing effects are unclear, and more studies on glucocorticoid and mineralocorticoid

receptor activity are thus needed.

This study is not without limitations. The use of a single sampling time point

prevented us from studying intra-individual variability and from exploring changes in BM

bacterial profiles over time. Moreover, the high sensitivity of the next-generation sequencing

used in our analyses can detect minimal amounts of contaminant DNA, especially in samples

carrying a low bacterial load such as BM, making it difficult to characterise BM microbes

without the influence of other DNA sources (e.g. skin). Strengths of our study include our

large sample size, the consistent methods for sample collection, processing and sequencing,

and the robust statistical analysis.

In sum, we provide initial insights into the dynamics of how maternal

psychobiological aspects in pregnancy and postpartum may affect BM composition. We

report that increased perinatal emotional complaints and prenatal salivary cortisol AUCg are

associated with decreased specific commensal bacteria, whereas enhanced BM cortisol is

linked to higher BM alpha diversity and reduced *Staphylococcus*. No associations were found

between psychobiological parameters and BM glucocorticoids. Future studies exploring the

consequence of these alterations on the infant gut and development are warranted.

Acknowledgements

We wish to express our gratitude to all women who participated in this study. We are grateful to our study team members Annika Schneider, Eileen Stephan, Nathalie Appenzeller, Rhea Sarah Gesù, Tamara Grossrieder, Tamara Lovrinovic, Seraina Weber, Stéphanie Loeffel, Simon Hühni and Firouzeh Farahmand for their research assistantship. Special thanks to Rausch AG, Medela Schweiz AG, MAM Baby AG and Bimbosan that freely provided gift sets for the study participants. Finally, we would like to thank the Swiss Federal Food Safety and Veterinary Office as well as the Research Group for Food Perception from the ZHAW School of Life Sciences for their support.

Authors' Contributions

RAC and UE conceived the presented idea. ND and RAC conducted research. ND analysed the data. RAC and UE verified the analytical methods and contributed to the interpretation of the results. RAC supervised the manuscript. ND took the lead in writing the manuscript. All authors provided critical feedback and contributed to the final version of the manuscript.

References

[1] E. J. Mayer-Davis, S. L. Rifas-Shiman, L. Zhou, F. B. Hu, G. A. Colditz, and M. W. Gillman, "Breast-feeding and risk for childhood obesity: does maternal diabetes or obesity status matter?," *Diabetes care*, vol. 29, no. 10, pp. 2231–2237, 2006, doi: 10.2337/dc06-0974.

[2] C. G. Victora *et al.,* "Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect," *The Lancet*, vol. 387, no. 10017, pp. 475–490, 2016, doi: 10.1016/S0140-6736(15)01024-7.

[3] T. M. Samuel, Q. Zhou, F. Giuffrida, D. Munblit, V. Verhasselt, and S. K. Thakkar, "Nutritional and Non-nutritional Composition of Human Milk Is Modulated by Maternal, Infant, and Methodological Factors," *Frontiers in nutrition*, vol. 7, p. 576133, 2020, doi: 10.3389/fnut.2020.576133.

[4] O. Ballard and A. L. Morrow, "Human milk composition: nutrients and bioactive factors," *Pediatric clinics of North America*, vol. 60, no. 1, pp. 49–74, 2013, doi: 10.1016/j.pcl.2012.10.002.

[5] V. Notarbartolo, M. Giuffrè, C. Montante, G. Corsello, and M. Carta, "Composition of Human Breast Milk Microbiota and Its Role in Children's Health," *Pediatric gastroenterology, hepatology & nutrition*, vol. 25, no. 3, pp. 194–210, 2022, doi: 10.5223/pghn.2022.25.3.194.

[6] A. Togo, J.-C. Dufour, J.-C. Lagier, G. Dubourg, D. Raoult, and M. Million, "Repertoire of human breast and milk microbiota: a systematic review," *Future microbiology*, vol. 14, pp. 623–641, 2019, doi: 10.2217/fmb-2018-0317.

[7] C. A. Moubareck, "Human Milk Microbiota and Oligosaccharides: A Glimpse into Benefits, Diversity, and Correlations," *Nutrients*, vol. 13, no. 4, 2021, doi: 10.3390/nu13041123.

[8] A. Diez-Sampedro, M. Flowers, M. Olenick, T. Maltseva, and G. Valdes, "Women's Choice Regarding Breastfeeding and Its Effect on Well-Being," *Nursing for women's health*, vol. 23, no. 5, pp. 383–389, 2019, doi: 10.1016/j.nwh.2019.08.002.

[9] K. M. Hunt *et al.,* "Characterization of the diversity and temporal stability of bacterial communities in human milk," *PloS one*, vol. 6, no. 6, e21313, 2011, doi: 10.1371/journal.pone.0021313.

[10] M. Kortesniemi *et al.,* "Human milk metabolome is associated with symptoms of maternal psychological distress and milk cortisol," *Food chemistry*, vol. 356, p. 129628, 2021, doi: 10.1016/j.foodchem.2021.129628.

[11] M. G. Di Benedetto, C. Bottanelli, A. Cattaneo, C. M. Pariante, and A. Borsini, "Nutritional and immunological factors in breast milk: A role in the intergenerational transmission from maternal psychopathology to child development," *Brain, behavior, and immunity*, vol. 85, pp. 57–68, 2020, doi: 10.1016/j.bbi.2019.05.032.

[12] H. Demmelmair, E. Jiménez, M. C. Collado, S. Salminen, and M. K. McGuire, "Maternal and Perinatal Factors Associated with the Human Milk Microbiome," *Current developments in nutrition*, vol. 4, no. 4, nzaa027, 2020, doi: 10.1093/cdn/nzaa027.

[13] R. Enaud *et al.,* "The Gut-Lung Axis in Health and Respiratory Diseases: A Place for Inter-Organ and Inter-Kingdom Crosstalks," *Frontiers in cellular and infection microbiology*, vol. 10, p. 9, 2020, doi: 10.3389/fcimb.2020.00009.

[14] S. K. Dogra, J. Doré, and S. Damak, "Gut Microbiota Resilience: Definition, Link to Health and Strategies for Intervention," *Frontiers in microbiology*, vol. 11, p. 572921, 2020, doi: 10.3389/fmicb.2020.572921.

[15] K. Hufnagl, I. Pali-Schöll, F. Roth-Walter, and E. Jensen-Jarolim, "Dysbiosis of the gut and lung microbiome has a role in asthma," *Seminars in immunopathology*, vol. 42, no. 1, pp. 75–93, 2020, doi: 10.1007/s00281-019-00775-y.

[16] K. G. Margolis, J. F. Cryan, and E. A. Mayer, "The Microbiota-Gut-Brain Axis: From Motility to Mood," *Gastroenterology*, vol. 160, no. 5, pp. 1486–1501, 2021, doi: 10.1053/j.gastro.2020.10.066.

[17] G. Oikonomou *et al.,* "Milk Microbiota: What Are We Exactly Talking About?," *Frontiers in microbiology*, vol. 11, p. 60, 2020, doi: 10.3389/fmicb.2020.00060.

[18] P. D. Browne *et al.,* "Human Milk Microbiome and Maternal Postnatal Psychosocial Distress," *Frontiers in microbiology*, vol. 10, p. 2333, 2019, doi: 10.3389/fmicb.2019.02333.

[19] M. W. Groer, M. W. Davis, and J. Hemphill, "Postpartum stress: current concepts and the possible protective role of breastfeeding," *Journal of obstetric, gynecologic, and neonatal nursing : JOGNN*, vol. 31, no. 4, pp. 411–417, 2002, doi: 10.1111/j.1552-6909.2002.tb00063.x.

[20] R. J. Ruiz and J. T. Fullerton, "The measurement of stress in pregnancy," *Nursing & health sciences*, vol. 1, no. 1, pp. 19–25, 1999, doi: 10.1046/j.1442-2018.1999.00004.x.

[21] B. R. H. van den Bergh *et al.,* "Prenatal developmental origins of behavior and mental health: The influence of maternal stress in pregnancy," *Neuroscience and biobehavioral reviews*, vol. 117, pp. 26–64, 2020, doi: 10.1016/j.neubiorev.2017.07.003.

[22] S. M. Woods, J. L. Melville, Y. Guo, M.-Y. Fan, and A. Gavin, "Psychosocial stress during pregnancy," *American journal of obstetrics and gynecology*, vol. 202, no. 1, 61.e1-7, 2010, doi: 10.1016/j.ajog.2009.07.041.

[23] A. Kawano, Y. Emori, and S. Miyagawa, "Association between stress-related substances in saliva and immune substances in breast milk in puerperae," *Biological research for nursing*, vol. 10, no. 4, pp. 350–355, 2009, doi: 10.1177/1099800409331892.

[24] S. Pundir *et al.,* "Maternal influences on the glucocorticoid concentrations of human milk: The STEPS study," *Clinical nutrition (Edinburgh, Scotland)*, vol. 38, no. 4, pp. 1913–1920, 2019, doi: 10.1016/j.clnu.2018.06.980.

[25] J. J. Hollanders, A. C. Heijboer, B. van der Voorn, J. Rotteveel, and M. J. J. Finken, "Nutritional programming by glucocorticoids in breast milk: Targets, mechanisms and possible implications," *Best practice & research. Clinical endocrinology & metabolism*, vol. 31, no. 4, pp. 397–408, 2017, doi: 10.1016/j.beem.2017.10.001.

[26] A. Stein *et al.,* "Effects of perinatal mental disorders on the fetus and child," *The Lancet*, vol. 384, no. 9956, pp. 1800–1819, 2014, doi: 10.1016/S0140-6736(14)61277-0.

[27] K. M. Sawyer, P. A. Zunszain, P. Dazzan, and C. M. Pariante, "Intergenerational transmission of depression: clinical observations and molecular mechanisms," *Molecular psychiatry*, vol. 24, no. 8, pp. 1157–1177, 2019, doi: 10.1038/s41380-018-0265-4.

[28] L. Drago, M. Toscano, R. de Grandi, E. Grossi, E. M. Padovani, and D. G. Peroni, "Microbiota network and mathematic microbe mutualism in colostrum and mature milk collected in two different geographic areas: Italy versus Burundi," *The ISME journal*, vol. 11, no. 4, pp. 875–884, 2017, doi: 10.1038/ismej.2016.183.

[29] L. Fernández *et al.,* "The human milk microbiota: origin and potential roles in health and disease," *Pharmacological research*, vol. 69, no. 1, pp. 1–10, 2013, doi: 10.1016/j.phrs.2012.09.001.

[30] A. J. Macpherson and T. Uhr, "Induction of protective IgA by intestinal dendritic cells carrying commensal bacteria," *Science (New York, N.Y.)*, vol. 303, no. 5664, pp. 1662–1665, 2004, doi: 10.1126/science.1091334.

[31] P. F. Perez *et al.,* "Bacterial imprinting of the neonatal immune system: lessons from maternal cells?," *Pediatrics*, vol. 119, no. 3, e724-32, 2007, doi: 10.1542/peds.2006-1649.

[32] L. F. Stinson *et al.,* "The human milk microbiome: who, what, when, where, why, and how?," *Nutrition reviews*, vol. 79, no. 5, pp. 529–543, 2021, doi: 10.1093/nutrit/nuaa029.

[33] E. A. Mayer and E. Y. Hsiao, "The Gut and Its Microbiome as Related to Central Nervous System Functioning and Psychological Well-being: Introduction to the Special Issue of Psychosomatic Medicine," *Psychosomatic medicine*, vol. 79, no. 8, pp. 844–846, 2017, doi: 10.1097/PSY.0000000000000525.

[34] G. de Palma, S. M. Collins, P. Bercik, and E. F. Verdu, "The microbiota-gut-brain axis in gastrointestinal disorders: stressed bugs, stressed brain or both?," *The Journal of physiology*, vol. 592, no. 14, pp. 2989–2997, 2014, doi: 10.1113/jphysiol.2014.273995.

[35] V. Osadchiy, C. R. Martin, and E. A. Mayer, "The Gut-Brain Axis and the Microbiome: Mechanisms and Clinical Implications," *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association*, vol. 17, no. 2, pp. 322–332, 2019, doi: 10.1016/j.cgh.2018.10.002.

[36] B. Misiak *et al.,* "The HPA axis dysregulation in severe mental illness: Can we shift the blame to gut microbiota?," *Progress in neuro-psychopharmacology & biological psychiatry*, vol. 102, p. 109951, 2020, doi: 10.1016/j.pnpbp.2020.109951.

[37] A. Harris and J. Seckl, "Glucocorticoids, prenatal stress and the programming of disease," *Hormones and behavior*, vol. 59, no. 3, pp. 279–289, 2011, doi: 10.1016/j.yhbeh.2010.06.007.

[38] G. A. Kaltsas and G. P. Chrousos, Eds., *The neuroendocrinology of stress,* 3rd ed. Cambridge, UK: Cambridge University Press, 2007.

[39] R. M. Sapolsky, L. M. Romero, and A. U. Munck, "How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions," *Endocr Rev*, vol. 21, no. 1, pp. 55–89, 2000, doi: 10.1210/edrv.21.1.0389.

[40] F. R. Patacchioli *et al.,* "Maternal plasma and milk free cortisol during the first 3 days of breast-feeding following spontaneous delivery or elective cesarean section," *Gynecologic and obstetric investigation*, vol. 34, no. 3, pp. 159–163, 1992, doi: 10.1159/000292751.

[41] M. Aparicio *et al.,* "Human milk cortisol and immune factors over the first three postnatal months: Relations to maternal psychosocial distress," *PloS one*, vol. 15, no. 5, e0233554, 2020, doi: 10.1371/journal.pone.0233554.

[42] S. Hart, L. Boylan, B. Border, S. R. Carroll, D. McGunegle, and R. M. Lampe, "Breast milk levels of cortisol and Secretory Immunoglobulin A (SIgA) differ with maternal mood and infant neuro-behavioral functioning," *Infant Behavior and Development*, vol. 27, no. 1, pp. 101–106, 2004, doi: 10.1016/j.infbeh.2003.06.002.

[43] M. Romijn *et al.,* "The Association between Maternal Stress and Glucocorticoid Rhythmicity in Human Milk," *Nutrients*, vol. 13, no. 5, 2021, doi: 10.3390/nu13051608.

[44] H. A. Tucker and J. W. Schwalm, "Glucocorticoids in mammary tissue and milk," *Journal of animal science*, vol. 45, no. 3, pp. 627–634, 1977, doi: 10.2527/jas1977.453627x.

[45] E. M. Klein *et al.,* "The German version of the Perceived Stress Scale - psychometric characteristics in a representative German community sample," *BMC psychiatry*, vol. 16, p. 159, 2016, doi: 10.1186/s12888-016-0875-9.

[46] S. Cohen, T. Kamarck, and R. Mermelstein, "A Global Measure of Perceived Stress," *Journal of Health and Social Behavior*, vol. 24, no. 4, p. 385, 1983, doi: 10.2307/2136404.

[47] N. Schmitz, J. Kruse, and W. Tress, "Psychometric properties of the General Health Questionnaire (GHQ-12) in a German primary care sample," *Acta psychiatrica Scandinavica*, vol. 100, no. 6, pp. 462–468, 1999, doi: 10.1111/j.1600-0447.1999.tb10898.x.

[48] M. Romppel, E. Braehler, M. Roth, and H. Glaesmer, "What is the General Health Questionnaire-12 assessing? Dimensionality and psychometric properties of the General Health Questionnaire-12 in a large scale German population sample," *Comprehensive psychiatry*, vol. 54, no. 4, pp. 406–413, 2013, doi: 10.1016/j.comppsych.2012.10.010.

[49] D. P. Goldberg and B. Blackwell, "Psychiatric illness in general practice. A detailed study using a new method of case identification," *British medical journal*, vol. 1, no. 5707, pp. 439–443, 1970, doi: 10.1136/bmj.2.5707.439.

[50] M. Hankins, "The factor structure of the twelve item General Health Questionnaire (GHQ-12): the result of negative phrasing?," *Clinical practice and epidemiology in mental health : CP & EMH*, vol. 4, p. 10, 2008, doi: 10.1186/1745-0179-4-10.

[51] S. M. Ibrahim and M. Lobel, "Conceptualization, measurement, and effects of pregnancy-specific stress: review of research using the original and revised Prenatal Distress Questionnaire," *Journal of behavioral medicine*, vol. 43, no. 1, pp. 16–33, 2020, doi: 10.1007/s10865-019-00068-7.

[52] M. Lobel, D. L. Cannella, J. E. Graham, C. DeVincent, J. Schneider, and B. A. Meyer, "Pregnancy-specific stress, prenatal health behaviors, and birth outcomes," *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*, vol. 27, no. 5, pp. 604–615, 2008, doi: 10.1037/a0013242.

[53] A. M. Bergant, T. Nguyen, K. Heim, H. Ulmer, and O. Dapunt, "Deutschsprachige Fassung und Validierung der "Edinburgh postnatal depression scale"," (in ger), *Deutsche*

medizinische Wochenschrift (1946), vol. 123, no. 3, pp. 35–40, 1998, doi: 10.1055/s-2007-1023895.

[54] H. Lukesch, *Geburts-Angst-Skala G-A-S*. Göttingen: Hogrefe, 1983.

[55] A. Bratsikas, "Assessment of psychobiological stress reactivity and its relation to postpartum mood states," University of Zurich, 2006.

[56] L. Laux, *Das State-Trait-Angstinventar (STAI) : theoretische Grundlagen und Handanweisung*: Beltz, 1981. [Online]. Available: https://fis.uni-bamberg.de/handle/uniba/26756

[57] C. Spielberger and S. Sydeman, *State-trait anxiety inventory and state-trait anger expression inventory. The use of psychological testing for treatment planning and outcome assessment*, 1994. [Online]. Available: https://scholar.google.com/citations?user=l5bq0t4aaaaj&hl=en&oi=sra

[58] SPIELBERGER C. D., "Manual for the State-trait Anxietry, Inventory," *Consulting Psychologist*, 1970. [Online]. Available: https://cir.nii.ac.jp/crid/1572261549249149184

[59] Y. Gidron, "Trait Anxiety," in *Encyclopedia of Behavioral Medicine*, M. Gellman and J. R. Turner, Eds., New York, NY: Springer New York, 2016, pp. 1–2.

[60] J. Heron, T. G. O'Connor, J. Evans, J. Golding, and V. Glover, "The course of anxiety and depression through pregnancy and the postpartum in a community sample," *Journal of affective disorders*, vol. 80, no. 1, pp. 65–73, 2004, doi: 10.1016/j.jad.2003.08.004.

[61] I. Nast, M. Bolten, G. Meinlschmidt, and D. H. Hellhammer, "How to measure prenatal stress? A systematic review of psychometric instruments to assess psychosocial stress during pregnancy," *Paediatric and perinatal epidemiology*, vol. 27, no. 4, pp. 313–322, 2013, doi: 10.1111/ppe.12051.

[62] B. van der Voorn, M. de Waard, J. B. van Goudoever, J. Rotteveel, A. C. Heijboer, and M. J. Finken, "Breast-Milk Cortisol and Cortisone Concentrations Follow the Diurnal Rhythm of Maternal Hypothalamus-Pituitary-Adrenal Axis Activity," *The Journal of nutrition*, vol. 146, no. 11, pp. 2174–2179, 2016, doi: 10.3945/jn.116.236349.

[63] M. K. McGuire *et al.,* "What's normal? Oligosaccharide concentrations and profiles in milk produced by healthy women vary geographically," *The American journal of clinical nutrition*, vol. 105, no. 5, pp. 1086–1100, 2017, doi: 10.3945/ajcn.116.139980.

[64] B. van der Voorn *et al.,* "Determination of cortisol and cortisone in human mother's milk," *Clinica chimica acta; international journal of clinical chemistry*, vol. 444, pp. 154–155, 2015, doi: 10.1016/j.cca.2015.02.015.

[65] A. Klindworth *et al.,* "Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies," *Nucleic acids research*, vol. 41, no. 1, e1, 2013, doi: 10.1093/nar/gks808.

[66] F. Faul, E. Erdfelder, A. Buchner, and A.-G. Lang, "Statistical power analyses using G*Power 3.1: tests for correlation and regression analyses," *Behavior research methods*, vol. 41, no. 4, pp. 1149–1160, 2009, doi: 10.3758/BRM.41.4.1149.

[67] J. C. Pruessner, C. Kirschbaum, G. Meinlschmid, and D. H. Hellhammer, "Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change," *Psychoneuroendocrinology*, vol. 28, no. 7, pp. 916–931, 2003, doi: 10.1016/S0306-4530(02)00108-7.

[68] P. La Marca-Ghaemmaghami, R. La Marca, S. M. Dainese, M. Haller, R. Zimmermann, and U. Ehlert, "The association between perceived emotional support, maternal mood, salivary cortisol, salivary cortisone, and the ratio between the two compounds in response to acute stress in second trimester pregnant women," *Journal of psychosomatic research*, vol. 75, no. 4, pp. 314–320, 2013, doi: 10.1016/j.jpsychores.2013.08.010.

[69] R. Benediktsson, A. A. Calder, C. R. Edwards, and J. R. Seckl, "Placental 11 beta-hydroxysteroid dehydrogenase: a key regulator of fetal glucocorticoid exposure," *Clinical endocrinology*, vol. 46, no. 2, pp. 161–166, 1997, doi: 10.1046/j.1365-2265.1997.1230939.x.

[70] B. J. Callahan, P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes, "DADA2: High-resolution sample inference from Illumina amplicon data," *Nature methods*, vol. 13, no. 7, pp. 581–583, 2016, doi: 10.1038/nmeth.3869.

[71] Y. Benjamini and Y. Hochberg, "Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing," *Journal of the Royal Statistical Society: Series B (Methodological)*, vol. 57, no. 1, pp. 289–300, 1995, doi: 10.1111/j.2517-6161.1995.tb02031.x.

[72] G. Falony *et al.,* "Population-level analysis of gut microbiome variation," *Science (New York, N.Y.)*, vol. 352, no. 6285, pp. 560–564, 2016, doi: 10.1126/science.aad3503.

[73] E. Bayar, P. R. Bennett, D. Chan, L. Sykes, and D. A. MacIntyre, "The pregnancy microbiome and preterm birth," *Seminars in immunopathology*, vol. 42, no. 4, pp. 487–499, 2020, doi: 10.1007/s00281-020-00817-w.

[74] S. L. Gillespie, A. M. Mitchell, J. M. Kowalsky, and L. M. Christian, "Maternal parity and perinatal cortisol adaptation: The role of pregnancy-specific distress and implications

for postpartum mood," *Psychoneuroendocrinology*, vol. 97, pp. 86–93, 2018, doi: 10.1016/j.psyneuen.2018.07.008.

[75] C. Gomez-Gallego, I. Garcia-Mantrana, S. Salminen, and M. C. Collado, "The human milk microbiome and factors influencing its composition and activity," *Seminars in fetal & neonatal medicine*, vol. 21, no. 6, pp. 400–405, 2016, doi: 10.1016/j.siny.2016.05.003.

[76] N. T. Mueller *et al.,* "Birth mode-dependent association between pre-pregnancy maternal weight status and the neonatal intestinal microbiome," *Scientific reports*, vol. 6, p. 23133, 2016, doi: 10.1038/srep23133.

[77] L. CARROLL, "BACTERIOLOGICAL CRITERIA FOR FEEDING RAW BREAST-MILK TO BABIES ON NEONATAL UNITS," *The Lancet*, vol. 314, no. 8145, pp. 732–733, 1979, doi: 10.1016/s0140-6736(79)90654-8.

[78] A. I. Eidelman and G. Szilagyi, "Patterns of bacterial colonization of human milk," *Obstetrics and gynecology*, vol. 53, no. 5, pp. 550–552, 1979.

[79] M. P. Heikkilä and P. E. J. Saris, "Inhibition of Staphylococcus aureus by the commensal bacteria of human milk," *Journal of applied microbiology*, vol. 95, no. 3, pp. 471–478, 2003, doi: 10.1046/j.1365-2672.2003.02002.x.

[80] P. V. Kirjavainen, E. Apostolou, T. Arvola, S. J. Salminen, G. R. Gibson, and E. Isolauri, "Characterizing the composition of intestinal microflora as a prospective treatment target in infant allergic disease," *FEMS immunology and medical microbiology*, vol. 32, no. 1, pp. 1–7, 2001, doi: 10.1111/j.1574-695X.2001.tb00526.x.

[81] C. F. Favier, E. E. Vaughan, W. M. de Vos, and A. D. L. Akkermans, "Molecular monitoring of succession of bacterial communities in human neonates," *Applied and environmental microbiology*, vol. 68, no. 1, pp. 219–226, 2002, doi: 10.1128/AEM.68.1.219-226.2002.

[82] P. Huovinen, "Bacteriotherapy: the time has come : Bacterial interference is an increasingly attractive approach to prevention and therapy," *BMJ : British Medical Journal*, vol. 323, no. 7309, pp. 353–354, 2001.

[83] M. C. Collado, S. Delgado, A. Maldonado, and J. M. Rodríguez, "Assessment of the bacterial diversity of breast milk of healthy women by quantitative real-time PCR," *Letters in applied microbiology*, vol. 48, no. 5, pp. 523–528, 2009, doi: 10.1111/j.1472-765X.2009.02567.x.

[84] J. F. Cryan and T. G. Dinan, "Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour," *Nature reviews. Neuroscience*, vol. 13, no. 10, pp. 701–712, 2012, doi: 10.1038/nrn3346.

[85] J. D. Galley, N. M. Parry, B. M. M. Ahmer, J. G. Fox, and M. T. Bailey, "The commensal microbiota exacerbate infectious colitis in stressor-exposed mice," *Brain, behavior, and immunity*, vol. 60, pp. 44–50, 2017, doi: 10.1016/j.bbi.2016.09.010.

[86] J. D. Galley *et al.,* "Exposure to a social stressor disrupts the community structure of the colonic mucosa-associated microbiota," *BMC microbiology*, vol. 14, p. 189, 2014, doi: 10.1186/1471-2180-14-189.

[87] C. Hechler *et al.,* "Association between Psychosocial Stress and Fecal Microbiota in Pregnant Women," *Scientific reports*, vol. 9, no. 1, p. 4463, 2019, doi: 10.1038/s41598-019-40434-8.

[88] R. Cabrera-Rubio, M. C. Collado, K. Laitinen, S. Salminen, E. Isolauri, and A. Mira, "The human milk microbiome changes over lactation and is shaped by maternal weight and mode of delivery," *The American journal of clinical nutrition*, vol. 96, no. 3, pp. 544–551, 2012, doi: 10.3945/ajcn.112.037382.

[89] L. Ruiz *et al.,* "Comparison of Two Approaches for the Metataxonomic Analysis of the Human Milk Microbiome," *Frontiers in cellular and infection microbiology*, vol. 11, p. 622550, 2021, doi: 10.3389/fcimb.2021.622550.

[90] A. Ojo-Okunola *et al.,* "Influence of Socio-Economic and Psychosocial Profiles on the Human Breast Milk Bacteriome of South African Women," *Nutrients*, vol. 11, no. 6, 2019, doi: 10.3390/nu11061390.

[91] R. W. Hyman, M. Fukushima, L. Diamond, J. Kumm, L. C. Giudice, and R. W. Davis, "Microbes on the human vaginal epithelium," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, no. 22, pp. 7952–7957, 2005, doi: 10.1073/pnas.0503236102.

[92] L. Grozdanov *et al.,* "Analysis of the genome structure of the nonpathogenic probiotic Escherichia coli strain Nissle 1917," *Journal of bacteriology*, vol. 186, no. 16, pp. 5432–5441, 2004, doi: 10.1128/JB.186.16.5432-5441.2004.

[93] M. Selma-Royo, J. Calvo Lerma, E. Cortés-Macías, and M. C. Collado, "Human milk microbiome: From actual knowledge to future perspective," *Seminars in perinatology*, vol. 45, no. 6, p. 151450, 2021, doi: 10.1016/j.semperi.2021.151450.

[94] S. R. Knowles, E. A. Nelson, and E. A. Palombo, "Investigating the role of perceived stress on bacterial flora activity and salivary cortisol secretion: a possible mechanism underlying susceptibility to illness," *Biological psychology*, vol. 77, no. 2, pp. 132–137, 2008, doi: 10.1016/j.biopsycho.2007.09.010.

[95] A. Kato-Kataoka *et al.,* "Fermented Milk Containing Lactobacillus casei Strain Shirota Preserves the Diversity of the Gut Microbiota and Relieves Abdominal Dysfunction in Healthy Medical Students Exposed to Academic Stress," *Applied and environmental microbiology*, vol. 82, no. 12, pp. 3649–3658, 2016, doi: 10.1128/AEM.04134-15.

[96] A. Moya and M. Ferrer, "Functional Redundancy-Induced Stability of Gut Microbiota Subjected to Disturbance," *Trends in microbiology*, vol. 24, no. 5, pp. 402–413, 2016, doi: 10.1016/j.tim.2016.02.002.

[97] P. D. Wadhwa, C. Dunkel-Schetter, A. Chicz-DeMet, M. Porto, and C. A. Sandman, "Prenatal psychosocial factors and the neuroendocrine axis in human pregnancy," *Psychosomatic medicine*, vol. 58, no. 5, pp. 432–446, 1996, doi: 10.1097/00006842-199609000-00006.

[98] A. K. Shea, D. L. Streiner, A. Fleming, M. V. Kamath, K. Broad, and M. Steiner, "The effect of depression, anxiety and early life trauma on the cortisol awakening response during pregnancy: preliminary results," *Psychoneuroendocrinology*, vol. 32, 8-10, pp. 1013–1020, 2007, doi: 10.1016/j.psyneuen.2007.07.006.

[99] S. F. Suglia, J. Staudenmayer, S. Cohen, M. B. Enlow, J. W. Rich-Edwards, and R. J. Wright, "Cumulative Stress and Cortisol Disruption among Black and Hispanic Pregnant Women in an Urban Cohort," *Psychological trauma : theory, research, practice and policy*, vol. 2, no. 4, pp. 326–334, 2010, doi: 10.1037/a0018953.

[100] T. G. O'Connor, W. Tang, M. A. Gilchrist, J. A. Moynihan, E. K. Pressman, and E. R. Blackmore, "Diurnal cortisol patterns and psychiatric symptoms in pregnancy: short-term longitudinal study," *Biological psychology*, vol. 96, pp. 35–41, 2014, doi: 10.1016/j.biopsycho.2013.11.002.

[101] C. Obel, M. Hedegaard, T. B. Henriksen, N. J. Secher, J. Olsen, and S. Levine, "Stress and salivary cortisol during pregnancy," *Psychoneuroendocrinology*, vol. 30, no. 7, pp. 647–656, 2005, doi: 10.1016/j.psyneuen.2004.11.006.

Table legends

Table 1. Demographic and obstetric outcomes of the total study sample ($N = 100$)

Table 2. BM microbiome taxa in relation to maternal prenatal emotional complaints ($N = 97$)

Table 3. BM microbiome taxa in relation to maternal salivary cortisol concentrations ($N = 96$)

Table 4. BM microbiome features in relation to BM cortisol concentrations ($N = 97$ BM samples for associations with microbial taxon abundances, $N = 96$ for alpha and beta diversity measures)

Figure Legends

Figure 1. Flow diagram for study participants

Figure 2. Taxonomic overview of the breast milk at genus level per sample for unrarefied data

Figure 3. Taxonomic overview of the breast milk at family level per sample for unrarefied data

Table 1. Demographic and obstetric outcomes of the total study sample ($N = 100$)

	N (%)	M ± SD	MD	IQR
Maternal characteristics at T1				
Maternal age	100 (100%)		32.50	4.00
Maternal BMI	100 (100%)		25.95	3.70
Maternal education				
Other	33 (33%)			
University/higher education	67 (67%)			
Marital status				
Single	30 (30%)			
Married	66 (66%)			
Divorced/separated	4 (4%)			
Maternal psychological parameters				
EPDS	100 (100%)		3.00	5.00
PSS-10	100 (100%)		20.00	7.00
STAI-S	100 (100%)		31.00	11.75
GHQ-12	100 (100%)		8.00	5.00
NuPDQ	100 (100%)		8.00	5.00
GAS	100 (100%)	16.78 ± 4.15		
Maternal salivary cortisol				
Mean AUCg (nmol/l)	100 (100%)		418.87	191.21
Mean total decline (nmol/l)	100 (100%)		- 4.94	3.56
Maternal characteristics at T2				
Maternal salivary cortisol				
Mean AUCg (nmol/l)	96 (96%)	436.09 ± 119.04		
Mean total decline (nmol/l)	96 (96%)		- 3.90	2.83
Maternal characteristics at T3				
Maternal psychological parameters				
EPDS	100 (100%)		3.00	6.75
PSS-10	100 (100%)	21.60 ± 5.50		
STAI-S	100 (100%)		31.50	13.75
GHQ-12	100 (100%)		9.00	5.75
Hours between last feeding and BM collection	97 (97%)		1.25	2.00
Breast used for breast milk samples	97 (97%)			
Right	41 (41%)			
Left	41 (41%)			
Both	15 (15%)			
Duration of breast milk sampling (seconds)	96 (96%)		420.00	300.00
Breast milk cortisol nmol/l	97 (97%)		11.16	9.13

Breast milk cortisone nmol/l	97 (97%)	52.96	17.52
Cortisone-to-(cortisol+cortisone) ratio	97 (97%)	0.83	0.10
Newborn characteristics at T3			
Gestational age	100 (100%)	40.14	1.57
Gender			
Male	46 (46%)		
Female	54 (54%)		
Weight (grams)	100 (100%)	3412.50	563
Length (centimetres)	100 (100%)	50.00	2
Membrane rupture during labour			
Yes	91 (91%)		
No	9 (9%)		

Note: N = number of participants; M = Mean; SD = standard deviation; MD = Median; IQR = Interquartile range; EPDS = Edinburgh Postnatal Depression Scale; GHQ-12 = General Health Questionnaire; PSS-10 = Perceived Stress Scale; STAI-S = State-Trait Anxiety Inventory - State scale; GAS = Birth Anxiety Scale (Geburts-Angst-Skala); NuPDQ = revised Prenatal Distress Questionnaire.

Table 2. BM microbiome taxa in relation to maternal prenatal emotional complaints (*N* = 97)

Category	Item	Variable (T1)	Kendall's tau	*p*-value	FDR
Class	Alphaproteobacteria	GAS	-0.2137	0.0028	0.0199*
Class	Alphaproteobacteria	Pregnancy-related sum	-0.1805	0.0114	0.0798*
Category	**Item**	**Variable (T3)**	**Kendall's tau**	***p*-value**	**FDR**
Class	Alphaproteobacteria	STAI-S	-0.1909	0.0079	0.0551*
Order	Propionibacteriales	STAI-S	-0.2413	0.0009	0.0173*
Order	Propionibacteriales	Postpartum sum	-0.2154	0.0028	0.0541*
Family	Propionibacteriaceae	STAI-S	-0.2413	0.0009	0.0273*
Family	Propionibacteriaceae	Postpartum sum	-0.2154	0.0028	0.0855*
Genus	Cutibacterium	STAI-S	-0.2413	0.0009	0.0419*
Genus	Pseudomonas_N	STAI-S	-0.2322	0.0019	0.0444*
Species	Cutibacterium acnes	STAI-S	-0.2488	0.0006	0.0497*
Species	Pseudomonas_N sp003640395	STAI-S	-0.2322	0.0019	0.0742*

Note: *p*-value = statistical significance, * *p* < .05; *FDR* = FDR-adjusted p-value; GAS = Birth Anxiety Scale (Geburts-Angst-Skala); STAI-S = State-Trait Anxiety Inventory - State scale; Postpartum sum = GHQ-12, PSS-10, EPDS, STAI-S at T3; Pregnancy-related sum = NuPDQ, GAS and EPDS at T1

Table 3. BM microbiome taxa in relation to maternal salivary cortisol concentrations (*N* = 96)

Category	Item	Variable (T1)	Kendall's tau	*p*-value	FDR
Class	Actinomycetia	AUCg	-0.1633	0.0180	0.0630*
Class	Clostridia	AUCg	-0.1824	0.0157	0.0630*
Genus	Stenotrophomonas	AUCg	-0.2433	0.0010	0.0481*
Species	Stenotrophomonas bentonitica_A	AUCg	-0.2557	0.0007	0.0526*
Category	**Item**	**Variable (T2)**	**Kendall's tau**	***p*-value**	**FDR**
Genus	Kocuria	AUCg	-0.2720	0.0012	0.0570*
Species	Kocuria rhizophila	AUCg	-0.2720	0.0012	0.0941*

Note: *p*-value = statistical significance, * *p* < .05; *FDR* = FDR-adjusted p-value; AUCg = area under the curve with respect to the ground.

Table 4. BM microbiome features in relation to BM cortisol concentrations ($N = 97$ BM samples for associations with microbial taxon abundances, $N = 96$ for alpha and beta diversity measures)

Category	Item	Variable (T3)	Kendall's tau	p-value	FDR
Alpha Diversity	Shannon diversity index	Cortisol	0.0967	n.s	n.s
Alpha Diversity	Richness	Cortisol	0.1671	0.0176*	0.0352*

Category	Item	Variable (T3)	Sum of Squares	Pr(>F)	FDR
Beta Diversity	Bray-Curtis dissimilarity	Cortisol	0.4374	n.s	n.s
Beta Diversity	Weighted UniFrac	Cortisol	0.2497	n.s	n.s

Category	Item	Variable (T3)	Kendall's tau	p-value	FDR
Family	Gemellaceae	Cortisol	0.2040	0.0038*	0.0577*
Family	Staphylococcaceae	Cortisol	- 0.2047	0.0030*	0.0577*
Genus	Gemella	Cortisol	0.2040	0.0038*	0.0885*
Genus	Staphylococcus	Cortisol	- 0.2047	0.0030*	0.0885*

Note: *p*-value = statistical significance, * $p < .05$; *Pr(>F)* = *p*-value associated with the F statistic; *FDR* = FDR-adjusted p-value; Cortisol nmol/l = BM cortisol concentration in nmol per litre.

Figure 1. Flow diagram for study participants.

Figure 2. Taxonomic overview of the breast milk at genus level per sample for unrarefied data. The numbers in parentheses refer to the number of ASVs aggregated for a given taxon. Bar plots display the relative abundance of the top 10 taxa with the highest average abundance across all samples. Light grey (Other) indicates the total relative abundance of ASVs that are not in the top 10 most abundant taxa; BM= breast milk.

Figure 3. Taxonomic overview of the breast milk at family level per sample for unrarefied data. The numbers in parentheses refer to the number of ASVs aggregated for a given taxon. Bar plots display the relative abundance of the top 10 taxa with the highest average abundance across all samples. Light grey (Other) indicates the total relative abundance of ASVs that are not in the top 10 most abundant taxa; BM= breast milk.