Chromosome-scale reference genome and RAD-based genetic map of yellow starthistle

(*Centaurea solstitialis*) reveal putative structural variation and QTLs associated with invader

traits

Running Title: Reference genome, genetic map, and invasion QTL in *Centaurea solstitialis*

Authors and affiliations:

Bryan Reatini[1,3]*, F. Alice Cang[1], Qiuyu Jiang[1], Michael T. W. McKibben[1,4], Michael S. Barker[1,5],

Loren H. Rieseberg[2,6], Katrina M. Dlugosch[1,7]

[1]Department of Ecology and Evolutionary Biology, University of Arizona

[2]Department of Botany, University of British Columbia

[3]ORCID:0000-0001-8643-4296

[4]ORCID:0000-0002-1342-0085

[5]ORCID:0000-0001-7173-1319

[6]ORCID:0000-0002-2712-2417

[7]ORCID:0000-0002-7302-6637

* *To whom correspondence should be addressed:* reati1bs@gmail.com

Abstract: Yellow starthistle (*Centaurea solstitialis*) is a highly invasive species that has been a

model system for the potential contribution of evolution to invader traits. Here, we report the

construction of a chromosome-scale reference genome for *C. solstitialis* using a combination of

PacBio HiFi and Dovetail Omni-C technologies, and functional gene annotation with RNAseq.

We validate the reference genome using a restriction site-associated DNA (RAD)-based genetic

map from an F2 mapping population. We find that syntenic comparisons to other taxa in the

Asteraceae reveal a chromosomal fusion in the lineage of *C. solstitialis*, and widespread fission

in globe artichoke (*Cynara cardunculus*). Using a QTL analysis from the mapping population

(derived from a cross between native and invading parents) we identify 13 QTL underpinning

size traits that are associated with adaptation in the invaded range, including a putative large-

scale chromosomal inversion that has a pleiotropic and overdominant effect on key invader

traits.

Keywords: reference genome, QTL, genetic map, invasive species, Centaurea solstitialis

Introduction

Yellow starthistle (*Centaurea solstitialis* L., Asteraceae) is a highly invasive plant on at least four continents and occupies a broad distribution throughout its native range in Eurasia (Maddox et al., 1985). Due to its ability to occupy a wide range of environments, and its profoundly negative ecological and economic impact in the invaded range, *C. solstitialis* has become a model system for understanding how invasive species might exploit open niches and dominate the communities in which they invade (e.g. Barker et al., 2017; Dlugosch et al., 2015a; Dukes et al., 2011; Graebner et al., 2012; Hierro et al., 2006; Zavaleta and Hulvey, 2004). This species is also well-suited to genomic study of the genetic basis of its success, as an annual, diploid (1N = 8), species of modest genome size (860 Mbp) (Bancheva & Greilhuber, 2006; Heiser Jr. & Whitaker, 1948; Irimia et al., 2017). Here, we present a chromosome-scale reference genome and associated analyses to facilitate genomic investigations of this model invader.

Centaurea solstitialis was introduced to South America in the 1600s and then to North America in the 1800s, and has since spread aggressively throughout the Western US, Chile, and Argentina (Barker et al., 2017; Gerlach, 1997). For invasive populations in California in particular, severe invasions are associated with the evolution of increased biomass and reproduction (Barker et al., 2017; Widmer et al., 2007). Within this invasion, plant size and biomass show evidence of adaptive clines along local environmental gradients (Dlugosch, Cang, et al., 2015)**.** Larger plant size is associated with increased reproductive fitness (Dlugosch, Cang, et al., 2015) and potentially an increased competitive ability (Montesinos et al., 2019;

Montesinos & Callaway, 2017). The former is predicted to lead to higher population growth

rates for invader genotypes relative to native genotypes (Dlugosch, Cang, et al., 2015).

Together, these observations suggest that the evolution of increased plant size in

Californian *C. solstitialis* populations may have directly contributed to the invasiveness of this

species, which makes it an attractive system to investigate how local adaptation can contribute

to biological invasions. Understanding the genetic basis of putative invasion traits is a core

objective of invasion biology (Bock et al., 2015; Dlugosch, Anderson, et al., 2015). However,

genomic resources are currently limited for many invasive species, which has impeded

investigations of the genetic basis of local adaptation for important invasion traits (Bock et al.,

2015; Dlugosch, Anderson, et al., 2015).

Here, we present a chromosome-scale reference genome for *C. solstitialis*, with

annotations of repetitive elements and functional genes. We validate the reference genome

assembly by comparing it with a genetic map constructed from an F2 mapping population, and

by carrying out gene synteny comparisons between starthistle and other representatives of

Asteraceae. We then demonstrate the utility of these genomic resources by investigating the

regions of the genome underpinning increased plant size in Californian populations via QTL

analysis, identifying candidate genes within those QTL regions.

Materials and Methods

Genomic reference sample collection and sequencing

Plant tissue used for the reference genome was grown from seed collected in August

2018 from the native range near Canales, Spain (site code 'CAN'; Lat: 41.00033, Long: -4.89718).

A reference voucher from the same locality is archived at the University of Arizona herbarium

(ARIZ #425375). The plant was reared to the early bolting stage in a greenhouse at the

University of Arizona (Tucson, Arizona, USA) under ambient light conditions from January to

July 2020, and placed in the dark for 24 hours before harvesting leaves directly into liquid

nitrogen for DNA extraction.

Leaf tissue was sent to Dovetail Genomics (Scotts Valley, California, USA) for reference

genome sequencing and assembly using a combination of PacBio HiFi and Dovetail Omni-C

approaches. Specifically, PacBio HiFi reads were generated by first constructing a ~20kb library

using SMRTbell Express Template Prep Kit 2.0 (PacBio, Menlo Park, CA, USA) using the

manufacturer's recommended protocol. The library was then bound to polymerase using the

Sequel II Binding Kit 2.0 (PacBio) and sequenced on PacBio Sequel II 8M SMRT cells, generating

45 Gb of data. Dovetail Omni-C libraries were prepped by fixing chromatin with formaldehyde,

then digesting chromatin with DNAse I. Proximity ligation was then performed, crosslinks were

reversed, and DNA was purified. Sequencing libraries were generated using NEBNext Ultra

enzymes and Illumina-compatible adapters, followed by PCR enrichment. Libraries were

sequenced on an Illumina HiSeqX platform, at ~30X coverage.

Genome assembly

The initial HiFi assembly was constructed *de novo* from PacBio reads using Wtdbg2

(Ruan & Li, 2020). Potential contamination in the initial assembly was identified and removed

based on BLAST v2.9 results against the NT database using Blobtools v1.1.1 (Laetsch & Blaxter,

2017). Haplotypic duplications were then filtered from the assembly using purge_dups v1.1.2

(Guan et al., 2020). The filtered initial assembly was used as input along with the Dovetail

Omni-C reads for scaffolding using the HiRise pipeline (Putnam et al., 2016). Briefly, Omni-C

reads were aligned to the draft assembly, a likelihood model for genomic distance between

read pairs was produced, and that model was then used to make joins and break misjoins in the

input draft assembly. The completeness of the final HiRise assembly was evaluated using the

255 Benchmarking Universal Single-Copy Orthologs (BUSCOs) from the Eukaryota dataset

(eukaryote_odb10) using BUSCO v4.0.5 (Manni et al., 2021).

Genome annotation

Repeat family identification and classification were performed on the final genome

assembly using RepeatModeler v2.0.1 (http://www.repeatmasker.org/RepeatModeler), relying

on RECON v1.08 (Bao & Eddy, 2002) and RepeatScout v1.0.6 (Price et al., 2005) for *de novo*

identification of repeats. Custom repeat libraries produced by RepeatModeler were then used

to identify and mask repeats in the final assembly using RepeatMasker v4.1.0

(https://www.repeatmasker.org/RepeatMasker/). High-confidence transposable elements were

then identified and classified by running RepeatMasker v4.1.0 with the plant TE database

nrTEplantsApril2020 (Contreras-Moreira et al., 2021) as the input library. In brief, this library

combines multiple plant TE databases and prunes the resulting library to remove redundant

sequences and minimize overlap with protein-coding domains in nucleotide-binding, leucine-

rich repeat (NLR) genes (Contreras-Moreira et al., 2021). Repetitive elements were then soft-

masked in the reference.

For gene identification, transcripts from six *C. solstitialis* individuals were sequenced

using RNAseq. Tissues included leaves from the same individual used for the reference genome,

whole shoots of two additional small seedlings from the same population ('CAN'), and three

individuals (again including two seedlings and a mature bolting individual) from seed collected

in September 2016 near Gilroy, California, USA in the invaded range (site code 'GIL'; Lat:

37.03373, Long: -121.53674; reference voucher ARIZ #425113). All seeds were grown in early

2020 in greenhouses at the University of Arizona under ambient light conditions. Tissue was

flash-frozen in liquid nitrogen and sent to Genewiz (Azenta Life Sciences, South Plainfield, NJew

Jersey, USA) for RNA extraction and sequencing. RNA libraries were prepared via rRNA-

depletion, and paired-end 2x150 reads were generated on the Illumina HiSeq platform.

Gene annotation was performed on the final repeat masked genome using a

combination of AUGUSTUS v2.5.5 (Stanke et al., 2006) and SNAP v2006-07-28 (Korf, 2004).

Specifically, coding sequences from *Cynara cardunculus* (Acquadro et al., 2020), *Lactuca sativa*

(Reyes-Chin-Wo et al., 2017), and *Helianthus annuus* (Badouin et al., 2017) were used to train

two independent *ab initio* models for *C. solstitialis* using AUGUSTUS and SNAP. RNAseq data

were then aligned to the final genome assembly using the STAR aligner software v2.7 (Dobin et

al., 2013) and AUGUSTUS was used to generate intron hints using the *bam2hints* tool. Gene

predictions were generated using MAKER (Holt & Yandell, 2011), SNAP, and AUGUSTUS, using

Swiss-Prot peptide sequences from the UniProt database to guide prediction and generate

peptide evidence in the MAKER pipeline. The final set of genes was filtered to contain only

genes predicted by both AUGUSTUS and SNAP. Putative gene function was assessed by

performing a BLAST search of the peptide sequences against the UniProt database, and tRNA

predictions were generated using tRNAscan-SE v2.05 (Chan et al., 2021). The quality of the

annotation was evaluated by plotting annotation edit distance from MAKER2 and gene synteny

was investigated between *C. solstitialis*, *Cynara cardunculus,* and *Lactuca sativa* using MCScanX

via TBTools v1.098696 (Chen et al., 2020) and visualized using SynVisio (Bandi & Gutwin, 2020).

The genome assembly and annotation of globe artichoke were downloaded from

http://www.artichokegenome.unito.it and the lettuce assembly (Lsat_Salinas_v7) and

annotation were downloaded from NCBI (BioProject: PRJNA173551).

Putative centromeric regions were identified using a sliding window analysis of gene and

repeat density using custom Python scripts (*YST_genome.ipynb*; zenodo citation). Specifically,

gene density from the gene annotation, unique repetitive element density from the repeat

annotation, and Gypsy/Copia long terminal repeat retrotransposon (LTR-RT) density from the

curated TE database were all calculated within 5Mbp sized windows with a step size of 100,000

bp for each chromosome. The midpoint of each element was used to define its location in the

genome. Counts for each of these four categories were normalized by dividing the count for the

window by the maximum count for the category within the chromosome, yielding a

measurement of the relative density of each category across the chromosome. Given that each

of the categories differs in their predicted association with centromere proximity (i.e. gene

density is predicted to be low near centromeres but LTR-RT density is often predicted to be

high), the direction of the effect was rescaled such that higher values always corresponded with

putative association with the centromere for all four categories. The average of the four

categories was then calculated for each window in order to assign a score ranging from 0-1 to

the window, with a score of 1 indicating a perfect score across all four categories (minimum

gene density, minimum unique repeat density, and maximum LTR-RT density for both Copia

and Gypsy elements). This score was then plotted across each chromosome to estimate the

location of centromeric regions.

Mapping population collection and sequencing

170   An F2 mapping population was created using an initial cross between a single native

171   range maternal parent and a single invaded range paternal parent. The native parent was

172   grown from seeds collected near Kirklareli, Turkey (site code 'TK23'; Lat: 41.751233, Long:

173   27.247883) in September 2008. The invader parent was grown from seeds collected near

174   Mariposa, California, USA (site code 'TRI'; Lat: 37.46178, Long: -119.79218) in 2008. Vouchers

175   from each of these populations were deposited in the ARIZ herbarium (#425116-425117).

176   Parental plants were grown in a greenhouse at the University of British Columbia

(Vancouver, British Columbia, Canada) and hand pollinated to produce F1 seed. Crosses were

performed by covering flowering heads (capitula) with white organza bags before the heads

opened to prevent pollination, then clipping heads presenting pollen and using these to brush

against heads presenting a large fraction of receptive stigmas, before re-covering receiving

heads until seed maturation. Of the resulting F1 progeny, four were selected at random and

reared to a large size in 11.4 L pots of commercial potting soil with 1 mL / L of 13:13:13

Osmocote fertilizer (Scotts Miracle-Gro, Marysville, Ohio, USA). The four F1 plants were

reciprocally crossed to one another in two pairs, and samples of their leaves stored at -80 C for

later genotyping.

Resulting F2 seeds were germinated on moist potting soil in an environmental room set

to 12 hr days and 16-18 C days / 14 C nights, and misted by hand every other day. A total of

2901 F2 seedlings were transplanted to 410 ml Deepots (Steuwe & Sons, Tangent, Oregon, USA)

in a 50:50 mix of silica sand and potting soil, with 2 mL of 13:13:13 Osmocote fertilizer. Deepots

were watered daily from below on a flood table in a greenhouse at the University of British

Columbia, under ambient lighting. Of the 2901 F2 plants, 300 were selected for genotyping

based on measurements of maximum leaf length at the first time point (see trait measurements

below), including 150 plants with the longest maximum leaf length and 150 plants with the

shortest maximum leaf length from the F2 generation.

Leaves from the 300 selected F2s, all four F1s, and six siblings of each parent (from field-

collected seeds) were used for genotyping. DNA was extracted using a modified CTAB/PVP

extraction protocol (Webb & Knapp, 1990) and sent to Floragenex (Beaverton, Oregon, USA) for

single digest Restriction Site Associated DNA (RAD) sequencing (Miller et al., 2007). RAD library

preparation was performed with the CpNpG 5-methylcytosine sensitive enzyme, *PstI*, and 1x80

bp reads were sequenced on the Illumina Genome Analyzer II platform, yielding an average of

29.8X coverage per sample.

Reference-guided genetic map

Raw single-end reads from the F2 mapping population were trimmed with the

process_radtags function in STACKS v2.60 (Rochette et al., 2019) using the *c*, *q*, and *r* flags and

specifying the *Pst*I restriction enzyme used in RAD library preparation. Trimmed reads were

aligned to the largest eight scaffolds of the final reference assembly (reference chromosomes,

see Results) using the *mem* function of the Burrows-Wheeler Aligner (BWA) (Li & Durbin, 2009)

and the resulting alignments were coordinate-sorted using samtools v1.10 (Li et al., 2009). RAD

loci were built and genotyped using the reference-aligned pipeline of STACKS via the *ref_map.pl*

wrapper. The *populations* function of STACKS was then used to filter the resulting genotype

data, requiring that a locus be present in each generation (P1, F1, and F2) and with a minimum

of 50% individuals represented at genotyped sites in order to be kept. Filtered data were

exported in variant call format, and parental genotypes for genetic mapping were called for

each RAD marker using an allele frequency-based approach. A custom python script

(*genmap_preprocessvcf.py*; zenodo citation) used the genotypes of six siblings of each parent to

polarize the alleles present in F1s as derived from either the native or invading population,

using the following criteria: for each allele present in the F1s at each RAD locus, the allele

frequency among native (TK) sibs and invader (TRI) sibs was compared and if freq(TRI)>freq(TK)

the allele was identified as predominantly invaded range (denoted 'A'), whereas if

freq(TK)>freq(TRI) the allele was identified as predominantly native range (denoted 'B'). For a

marker to be retained, all alleles in F1s needed to be successfully polarized, and there could be

no more than two missing F1 genotypes or one missing parental population genotype. Filtered

alleles were collapsed into parental classes (A and B), and F2 genotypes were exported to R/qtl

format using these genotype calls.

Importantly, although the reference genome was used to call genotypes at each RAD

marker, it was not used to group markers into linkage groups for genetic map construction.

Instead, initial linkage groups for the genetic map were built in R/qtl v1.50 (Broman et al., 2003)

using the recombination fractions observed among F2 genotypes. Specifically, a maximum

recombination fraction of 0.35 and minimum LOD score of 15 were used to group markers into

initial linkage groups. Markers showing evidence of significant segregation distortion were

removed using a significance cutoff of p=1E-5. The eight largest linkage groups were used for

the genetic map, assuming these linkage groups corresponded with the eight chromosomes of

C. solstitialis. The genetic distances between markers in the genetic map were then estimated

using R/qtl2 v0.28 (Broman et al., 2019). Specifically, genetic distances were first estimated

using the *est_map* function of R/qtl2 under a range of genotype error rates [0.001, 0.01, 0.025,

0.05, 0.075, 0.1]. The most likely error rate (0.075) was estimated using the $\log_{10}$ likelihood of

each estimated map and this rate was used to estimate distances in the final genetic map.

239   To validate the linear sequence of the chromosomes in the reference genome, we

plotted the collinearity between the genetic map and the physical reference using custom

Python scripts (*YST_genome.ipynb*; zenodo citation). Abberant markers in the genetic map -

those that fell on different chromosomes than the rest of the linkage group - were quantified to

determine the consistency between the reference genome and genetic map, and these

abberant markers where ultimately removed from downstream analyses. Collinearity patterns

were then compared with centromere genome scans in order to evaluate whether patterns of

recombination were consistent with the estimated centromere positions, again using custom

Python scripts (*YST_genome.ipynb*; zenodo citation).

Trait measurements and QTL analyses

250   The length of the longest leaf and the total number of leaves were recorded for all 2901

plants at 3.5 weeks and 5 weeks of age. Potential confounding factors during trait

measurement were accounted for by fitting linear models for each trait measurement using the

aov function in R, with fixed effects of the day measurements were collected, the individual

collecting the data, the spatial location in the greenhouse (block), the location of the plant

within the block, and the identity of the specific F1 mother of each F2. Residuals of the model

were then extracted and added to the grand mean to obtain corrected values of maximum leaf

length and total number of leaves at both time points for use in QTL analysis.

Additional representatives of both parental populations and F1 crosses between these

populations were included in the experiment for phenotypic comparison to the F2s. Parental

population plants were grown from field-collected seed including 28 'TRI' plants from the

invasion and 29 'TK23' plants from the native range. F1 seeds came from additional controlled

crosses, and included 17 plants with 'TRI' genotypes as the maternal parents and 24 plants with

'TK23' genotypes as the maternal parents. Germination and rearing were concurrent with the

F2 populations, under the same conditions.

Genome scans to identify QTL were performed using both Haley-Knott regression (Haley

& Knott, 1992) and the leave one chromosome out (LOCO; Yang et al., 2014) approach in R/qtl2

using the final genetic map and corrected phenotypic data as input. Both models were used to

qualitatively gauge support for QTL by comparing overlap between them. Given that the LOCO

method accounts for kinship when crosses are generated with parents from divergent source

populations – as was the case for the present mapping population – the LOCO model was used

as the primary model for candidate gene identification. For each method, two thresholds were

used to define QTL peaks using a permutation-based approach in R/qtl2: a standard significance

threshold of 0.05 and a suggestive threshold of 0.1. Separate peaks on the same chromosome

were identified using a minimum LOD decline of 1 between peaks. Candidate genes within QTL

peaks were identified by extracting all genes within a decline of 1 LOD on either side of the

peak.

Results

Genome assembly and annotation

The final chromosome-scale reference genome contained 8 primary scaffolds

corresponding with the haploid chromosome number of *C. solstitialis* (n=8) totaling ~725 Mbp.

The 8 primary scaffolds composed 94.8% of the total sequence content of the reference (L90=8,

Figure 1A), and 84.4% of the average genome size of *C. solstitialis* (860 Mbp) estimated by flow

cytometry (Irimia et al., 2017). An additional 1072 unplaced scaffolds totaled ~39 Mbp. Of the

255 BUSCOs in the eukaryote_odb10 dataset, 237 (92.9%) complete BUSCOs were found in the

8 primary scaffolds: 188 (73.7%) were complete and single-copy and 49 (19.2%) were complete

and duplicated (Table 1). None of the BUSCOs that were missing or fragmented in the 8 primary

scaffolds were located in the trailing scaffolds, but 4 of the BUSCOs that were found in single-

copy in the 8 primary scaffolds were duplicated in the trailing scaffolds. Given these results, we

hereafter refer to the 8 primary scaffolds as chromosomes, numbered by size.

Roughly 63.3% of the reference was repetitive DNA, with 29.04% identified as class I

transposable elements (retrotransposons) and 2.07% identified as class II transposable

elements (DNA transposons) (Table 1). A total of 481,954 retroelements - composing 12% of

the genome - were successfully classified using the high-confidence nrTEplants database

including 122,163 Ty1/Copia and 172,604 Gypsy LTR-RTs.

The complete reference annotation included 34,323 predicted gene models

encompassing a total of 59.8 Mbp of the reference sequence (Table 1). Of the 32,431 genes

that fell on the 8 putative chromosomes, 24,011 (74%) were able to be functionally annotated

using the UniProt database. A total of 616 tRNA prediction models were identified on the 8

putative chromosomes using tRNAscan-SE, which is similar to the 639 identified in *Arabidopsis*

thaliana (Chan et al., 2021). Using the cumulative frequency of annotation edit distance (from 0

being perfect support to 1 being no support) as a metric of annotation quality, the majority of

genes in the complete annotation are well supported by overlapping aligned RNA-seq and

protein homology data (Supplemental Figure 1).

Gene synteny analyses between the *C. solstitialis* genome and that of globe artichoke

(*Cynara cardunculus*) – the most closely related thistle (subfamily *Carduoideae*) whose genome

has been assembled to the chromosome-scale – using MCScanX revealed large blocks of

syntenic genes across the genome (Figure 2). Conserved synteny was also evident in

comparisons between the *C. solstitialis* genome and that of *Lactuca sativa* (subfamily

Cichorioideae), albeit with more evidence for interchromosomal rearrangements (Figure 2).

Three-way comparisons between *C. solstitialis*, *Cynara cardunculus,* and *Lactuca sativa* revealed

a complex history of chromosome evolution within the thistle subfamily *Carduoideae* (Figure X).

In particular, by using *Lactuca sativa* as an outgroup, multiple chromosome fissions are evident

within the *Cynara cardunculus* lineage (Figure 2C), whereas chromosome 3 of *C. solstitialis*

shows evidence of being derived from chromosome fusion (Figure 2D), and chromosome 7 is

highly conserved across all three taxa (Figure 2E).

For the sliding window analysis to detect putative centromeric regions, the raw counts

for each of the individual categories varied in their ability to predict centromere location for

each of the 8 chromosomes (Supplemental Figure 2). However, the average score of the four

categories revealed a clear signal indicating the location of putative pericentromeric regions for

each of the 8 chromosomes (Figure 3). Sliding window analysis of gene density alone revealed

roughly 2X higher gene density at the ends of chromosome arms relative to putative

centromeric regions for the 8 presumed chromosomes of the *C. solstitialis* genome

(Supplemental Figure 2).

Reference-guided genetic map

A total of 1,765 RAD markers across the 300 F2 individuals were successfully polarized

and were used as input for initial linkage group formation in the genetic map. After filtering for

segregation distortion, 1,064 markers remained in the 8 linkage groups of the genetic map.

Each of the 8 linkage groups corresponded with one of the 8 chromosomes in the *C. solstitialis*

reference genome (Figure 4B), with only 5 aberrant markers from a single linkage group falling

on a different chromosome than the rest of the markers from their linkage group. These

aberrant markers were pruned in the final map. Strong linkage and low estimated

recombination were evident between markers within each chromosome, whereas linkage was

weak and recombination was estimated to be frequent between markers on different

chromosomes (Figure 4A).

Collinearity analyses between the genetic map and physical map revealed megabase-

scale regions of low recombination in the center of each chromosome, with chromosomes 1, 5,

7, and 8 showing low recombination at one of the two extremes of the chromosome as well

(Figure 3A). To evaluate whether these regions of low recombination correspond with

pericentromeric regions, we compared our collinearity analyses with our centromere genome

scans (Figure 3A,B). For all chromosomes except 1 and 7, regions of low recombination in the

genetic map corresponded closely with estimated pericentromeric regions (Figure 3A,B).

Moreover, these same regions corresponded with lower RAD marker density (Figure 3A;

Supplemental Figure 2). Chromosomes 1 and 7 displayed low recombination across the majority

of the chromosome - in putative pericentromeric regions and on one of the distal arms of each

chromosome (Figure 3A,B).

Trait differences and QTL analyses

Invader parental genotypes had greater total number of leaves and greater maximum

leaf length than native parental genotypes at both time points (Supplemental Figure 3), which is

consistent with previous studies of invasive and native *C. solstitialis* populations (Dlugosch,

Cang, et al., 2015). F1 individuals displayed heterosis for maximum leaf length at both time

points, and for total number of leaves at the second time point (Supplemental Figure 3),

whereas F2 individuals showed the greatest range in trait values (Supplemental Figure 3). Given

that the 300 genotyped F2s were selected from the ends of the distribution of longest leaf

length at time point 1, they showed an expected bimodal distribution for this trait

(Supplemental Figure 4). However, total number of leaves followed a roughly normal

distribution for both time points within the F2 mapping population used for downstream QTL

analyses (Supplemental Figure 4).

Haley-Knott regression and LOCO genome scans revealed a total of 13 distinct QTL

peaks with varying degrees of support (summarized in Table 2). A total of seven peaks were

associated with maximum leaf length alone, four were associated with total number of leaves

alone, and two peaks were associated with both traits (Table 2). For all four peaks associated

with total number of leaves, the dominance pattern appeared to be additive, and the invader

alleles at each QTL were associated with a greater number of leaves (Table 2). Of the seven

peaks associated with maximum leaf length, mixed patterns of dominance (additive, complete,

and overdominance) were observed and the parental allele driving larger leaves varied

between loci (Table 2). One of the two pleiotropic QTL displayed complete dominance favoring

the native allele, and the other pleiotropic QTL displayed overdominance. The latter QTL

corresponded with the large region of low recombination outside of the putative centromeric

region on chromosome 1 (Figure 3). Similarly, one of the suggestive QTL for total number of

leaves corresponded with the other large region of low recombination outside of the

pericentromeric region on chromosome 7 (Figure 3). The number of candidate genes under QTL

peaks depended on the size of the QTL peak, varying from 16 genes within 0.2 Mbp to 3532

genes within the 88.9 Mbp region of low recombination on chromosome 1 (Table 2).

Discussion

We constructed a chromosome-scale genome assembly, gene and repeat annotations,

and genetic map for yellow starthistle (*Centaurea solstitialis*), and we validated these genomic

resources using a combination of assembly statistics, collinearity analyses, and gene synteny

analyses. This is the first non-cultivated species to be assembled to chromosome-scale in the

thistle subfamily (*Carduoideae*) of *Asteraceae*, with the only other representative genome of

Carduoideae being cultivated globe artichoke.

Following genome assembly using PacBio HiFi reads and scaffolding using Dovetail

Omni-C data, the 8 pseudohaploid chromosomes of the assembly made up 94.8% of the total

sequence content and contained 92.9% complete BUSCO genes (Table 1). The lack of any

additional BUSCO genes within the 1062 trailing scaffolds - in addition to these scaffolds only

comprising 5.2% of the total sequence content of the genome (Figure 1A) - suggest that these

trailing scaffolds are likely composed of contamination or mis-assemblies due to heterozygosity

in the outbred wild genotype used for genome assembly. The Omni-C contact map of the 8

pseudohaploid chromosomes of the assembly (Figure 1B) also demonstrates two expected

patterns of chromosome-scale assemblies: 1) link density is higher within chromosomes than

between, and 2) link density decays with physical distance within a chromosome (Lajoie et al.,

2015). Each of the 8 linkage groups in the genetic map also corresponded with one of the 8

chromosomes of the assembly (Figure 4), and the recombination frequencies between markers

in the genetic map were consistent with the linear sequence of each chromosome (Figure 3A).

Together, these results suggest that the 8 pseudohaploid chromosomes of the assembly

represent the complete set of chromosomes of *C. solstitialis*.

Estimated recombination was generally highest at the ends of each chromosome and

decreased towards putative pericentromeric regions located roughly in the center of each

chromosome (Figure 3A,B) - although chromosomes 5 and 8 had highly asymmetric centromere

positions. The general locations of these putative pericentromeric regions were estimated using

genome scans of gene density, unique repeat density, and LTR-RT density (Figure 3B).

Pericentromeric regions also corresponded with low RAD marker density in the genetic map

(Figure 3A).

Interestingly, chromosomes 1 and 7 displayed recombination patterns that did not

appear to be explained by the location of centromeres alone. For both of these chromosomes,

no recombination was observed across one of the chromosome arms, despite the central

location of the centromeres (Figure 3A,B). Similar collinearity patterns have been explained by

chromosomal inversions in a range of plant and animal taxa (Huang et al., 2020; Kirubakaran et

al., 2016; Lee et al., 2017; Tong et al., 2016), suggesting that large-scale chromosomal

rearrangements may exist between native and invasive genotypes of *C. solstitialis* on these

chromosomes.

The *C. solstitialis* assembly is the second chromosome-scale reference genome for a

member of the *Carduoideae*, which provides new opportunities to examine genome structural

evolution in the Asteraceae. Gene synteny analyses between *C. solstitialis*, *Cynara cardunculus,*

and *Lactuca sativa* revealed large regions of conserved synteny between all three genomes

(Figure 2). The ancestral haploid chromosome number of *Asteraceae* is estimated to be n=9

(Mota et al., 2016), which is retained in *Lactuca sativa* (subfamily *Chicoriae*; n=9) but not *C.*

solstitialis (1N = 8) or *Cynara cardunculus* (1N = 17). By using *Lactuca sativa* as an outgroup, we

attempted to assess the most parsimonious history of chromosomal evolution within the *C.*

solstitialis and *Cynara cardunculus* ancestral lineages. *Cynara cardunculus* has over double the

chromosome number as *C. solstitialis*, yet Scaglione et al. (2016) found that no whole-genome

duplication appears to have occurred after speciation between *Chicoriae* and *Carduoideae*,

using the distribution of Ks values between *Cynara cardunculus* and *Lactuca*. Consistent with

Scaglione et al. (2016), our gene synteny analyses revealed evidence for multiple chromosome

fissions in *Cynara cardunculus* rather than whole-genome duplication (Figure 2A; exemplified in

Figure 2C). Karyotype analyses of *Cynara cardunculus* have revealed that the chromosomes

display an unusual size distribution with discrete size categories of large, medium, and small

chromosomes (Falistocco, 2016) - perhaps due to multiple fission events of larger ancestral

chromosomes into smaller chromosomes as our results suggest. Our gene synteny analyses also

suggest that the reduction in the chromosome number of *C. solstitialis* (n=8) relative to the

base number for *Asteraceae* (n=9) may be due to the fusion of two ancestral chromosomes

within the lineage leading up to *C. solstitialis* (Figure 2D) - although there appear to be many

other interchromosomal rearrangements between these genomes (Figure 2B).

Invasive *C. solstitialis* populations show evidence for adaptive evolution of increased size

and reproduction, genetic changes which are thought to have aided in its invasion success

(Barker et al., 2017; Dlugosch, Cang, et al., 2015). Using the reference genome and genetic

mapping population described here, we identified 13 QTL associated with two phenotypic

components of increased plant size in this species. We identified four QTL peaks associated

with the total number of leaves, and all of these loci displayed greater leaf number for invader

genotypes relative to the native genotypes (Table 2) - a pattern consistent with differences

observed between parental phenotypes (Supplementary Figure 3). We identified seven QTL

associated with maximum leaf length, for which the dominance pattern and favored allele

depended on the locus, suggesting that both native and invasive populations have segregating

variation for larger leaf length at different loci.

Two pleiotropic QTL were identified by our QTL analyses, one displaying overdominance

and the other favoring the native genotype. Interestingly, the overdominant pleiotropic QTL on

chromosome 1 corresponded with the putative inversion identified by collinearity analyses

(Figure 3). Given that heterosis is observed in the F1 generation for both traits (Supplemental

Figure 3), it is plausible that overdominance (and pleiotropy) at this QTL could be driven by an

inversion (Faria et al., 2019). Specifically, a large-scale inversion would prevent recombination

across a large portion of the chromosome, as appears to be the case in this region of

chromosome 1 (Figure 3A), which would "fix" heterozygous loci affecting both traits across its

length in the F2 generation. The inverted region - and all phenotypically relevant genes within it

- would act as a single heterotic and pleiotropic locus. Balancing selection on overdominant

inversions is a classic form of local adaptation in general (Faria et al., 2019), making this QTL a

particularly intriguing candidate for local adaptation in this system. Alternatively,

overdominance at this QTL could be explained by heterozygous loci across the inversion

masking deleterious alleles (Connallon & Olito, 2022; Jay et al., 2022), which are likely to

accumulate during biological invasions due to strong genetic drift (i.e. expansion load: Gilbert et

al., 2017; Peischl et al., 2015). In this case, the maintenance of the inversion polymorphism

within invasive populations would not need to be driven by local adaptation *per se*, but it could

still be an important variant contributing to invasion success if invader genotypes that are

heterozygous for the inversion have higher fitness than homozygotes.

The putative inversion on chromosome 7 (identified by collinearity analyses) was also

identified as a suggestive QTL using Haley-Knott regression, though with additive effects

(favoring more leaves in the invader). It is perhaps not surprising that QTL are associated with

both of the large inversions that we inferred from our genetic map and reference genome,

given that any trait loci in these chromosomal regions will be linked to the entire inversion.

Nevertheless, loci that have large phenotypic effects - such as inversions containing many

tightly linked genes affecting a trait - are predicted to be favored by selection during invasions

due to the relative strength of genetic drift (Bock et al., 2015; Dlugosch, Anderson, et al., 2015),

particularly when gene flow is involved (Reatini & Vision, 2020). Inversions are increasingly

being identified by genomic studies as playing important roles in adaptation (Huang and

Rieseberg, 2020) and have recently been shown to contribute to the success of a plant invasion

(Battlay et al., 2022). Our results suggest that these structural variants may play a role in

phenotypic evolution in invasive populations of *C. solstitialis* and merit further investigation.

Investigating the genetic basis of invasion traits remains a central objective of invasion

biology, and here we advance that goal in the *Centaurea solstitialis* system. Together, the

reference genome assembly, annotation, genetic map, and QTL candidate regions presented

here provide essential genomic resources for studies of *C. solstitialis* invasion. Moreover, these

genomic resources will be useful for investigating the history of genome evolution across the

agriculturally and evolutionarily important Asteraceae family.

References

Acquadro, A., Portis, E., Valentino, D., Barchi, L., & Lanteri, S. (2020). "Mind the Gap": Hi-C Technology

Boosts Contiguity of the Globe Artichoke Genome in Low-Recombination Regions. *G3:*

Genes|Genomes|Genetics, *10*(10), 3557–3564. https://doi.org/10.1534/g3.120.401446

Badouin, H., Gouzy, J., Grassa, C. J., Murat, F., Staton, S. E., Cottret, L., Lelandais-Brière, C., Owens, G. L.,

Carrère, S., Mayjonade, B., Legrand, L., Gill, N., Kane, N. C., Bowers, J. E., Hubner, S., Bellec, A.,

Bérard, A., Bergès, H., Blanchet, N., … Langlade, N. B. (2017). The sunflower genome provides

insights into oil metabolism, flowering and Asterid evolution. *Nature*, *546*(7656), 148–152.

https://doi.org/10.1038/nature22380

Bancheva, S., & Greilhuber, J. (2006). Genome size in Bulgarian Centaurea s.l. (Asteraceae). *Plant*

Systematics and Evolution, *257*(1), 95–117. https://doi.org/10.1007/s00606-005-0384-7

Bandi, V., & Gutwin, C. (2020). Interactive Exploration of Genomic Conservation. *Graphics Interface*.

https://doi.org/10.20380/GI2020.09

Bao, Z., & Eddy, S. R. (2002). Automated De Novo Identification of Repeat Sequence Families in

Sequenced Genomes. *Genome Research*, *12*(8), 1269–1276. https://doi.org/10.1101/gr.88502

Barker, B. S., Andonian, K., Swope, S. M., Luster, D. G., & Dlugosch, K. M. (2017). Population genomic

analyses reveal a history of range expansion and trait evolution across the native and invaded

range of yellow starthistle (Centaurea solstitialis). *Molecular Ecology*, *26*(4), 1131–1147.

https://doi.org/10.1111/mec.13998

Battlay, P., Wilson, J., Bieker, V. C., Lee, C., Prapas, D., Petersen, B., Craig, S., Boheemen, L. van, Scalone,

R., Silva, N. P. de, Sharma, A., Konstantinović, B., Nurkowski, K. A., Rieseberg, L., Connallon, T.,

Martin, M. D., & Hodgins, K. A. (2022). *Large haploblocks underlie rapid adaptation in an*

invasive weed (p. 2022.03.02.482376). bioRxiv. https://doi.org/10.1101/2022.03.02.482376

Bock, D. G., Caseys, C., Cousens, R. D., Hahn, M. A., Heredia, S. M., Hübner, S., Turner, K. G., Whitney, K.

D., & Rieseberg, L. H. (2015). What we still don't know about invasion genetics. *Molecular*

Ecology, *24*(9), 2277–2297. https://doi.org/10.1111/mec.13032

Broman, K. W., Gatti, D. M., Simecek, P., Furlotte, N. A., Prins, P., Sen, Ś., Yandell, B. S., & Churchill, G. A.

(2019). R/qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and

Multiparent Populations. *Genetics*, *211*(2), 495–502.

https://doi.org/10.1534/genetics.118.301595

Broman, K. W., Wu, H., Sen, Ś., & Churchill, G. A. (2003). R/qtl: QTL mapping in experimental crosses.

Bioinformatics, *19*(7), 889–890. https://doi.org/10.1093/bioinformatics/btg112

Chan, P. P., Lin, B. Y., Mak, A. J., & Lowe, T. M. (2021). tRNAscan-SE 2.0: Improved detection and

functional classification of transfer RNA genes. *Nucleic Acids Research*, *49*(16), 9077–9096.

https://doi.org/10.1093/nar/gkab688

Chen, C., Chen, H., Zhang, Y., Thomas, H. R., Frank, M. H., He, Y., & Xia, R. (2020). TBtools: An Integrative

Toolkit Developed for Interactive Analyses of Big Biological Data. *Molecular Plant*, *13*(8), 1194–

1202. https://doi.org/10.1016/j.molp.2020.06.009

Connallon, T., & Olito, C. (2022). Natural selection and the distribution of chromosomal inversion

lengths. *Molecular Ecology*, *31*(13), 3627–3641. https://doi.org/10.1111/mec.16091

Contreras-Moreira, B., Filippi, C. V., Naamati, G., García Girón, C., Allen, J. E., & Flicek, P. (2021). K-mer

counting and curated libraries drive efficient annotation of repeats in plant genomes. *The Plant*

Genome, *14*(3), e20143. https://doi.org/10.1002/tpg2.20143

Dlugosch, K. M., Anderson, S. R., Braasch, J., Cang, F. A., & Gillette, H. D. (2015). The devil is in the

details: Genetic variation in introduced populations and its contributions to invasion. *Molecular*

Ecology, *24*(9), 2095–2111. https://doi.org/10.1111/mec.13183

Dlugosch, K. M., Cang, F. A., Barker, B. S., Andonian, K., Swope, S. M., & Rieseberg, L. H. (2015).

Evolution of invasiveness through increased resource use in a vacant niche. *Nature Plants*, *1*.

https://doi.org/10.1038/nplants.2015.66

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras,

T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

https://doi.org/10.1093/bioinformatics/bts635

Dukes, J. S., Chiariello, N. R., Loarie, S. R., & Field, C. B. (2011). Strong response of an invasive plant

540   species (Centaurea solstitialis L.) to global environmental changes. *Ecological Applications*, *21*(6),

541   1887–1894. https://doi.org/10.1890/11-0111.1

542 Falistocco, E. (2016). Cytogenetic characterization of cultivated globe artichoke (Cynara cardunculus var.

543   Scolymus) and cardoon (C. cardunculus var. Altilis). *Caryologia*, *69*(1), 1–4.

544   https://doi.org/10.1080/00087114.2015.1109935

545 Faria, R., Johannesson, K., Butlin, R. K., & Westram, A. M. (2019). Evolving Inversions. *Trends in Ecology*

546   *& Evolution*, *34*(3), 239–248. https://doi.org/10.1016/j.tree.2018.12.005

547 Gerlach, J. D. (1997). How the West Was Lost: Reconstructing the Invasion Dynamics of Yellow

548   Starthistle and Other Plant Invaders of Western Rangelands and Natural Areas. *Calif. Exot. Pest*

549   *Plant Counc. Symp. Proc.*, 6.

550 Gilbert, K. J., Sharp, N. P., Angert, A. L., Conte, G. L., Draghi, J. A., Guillaume, F., Hargreaves, A. L.,

551   Matthey-Doret, R., & Whitlock, M. C. (2017). Local Adaptation Interacts with Expansion Load

552   during Range Expansion: Maladaptation Reduces Expansion Load. *The American Naturalist*,

553   *189*(4), 368–380. https://doi.org/10.1086/690673

554 Graebner, R. C., Callaway, R. M., & Montesinos, D. (2012). Invasive species grows faster, competes

555   better, and shows greater evolution toward increased seed size and growth than exotic non-

556   invasive congeners. *Plant Ecology*, *213*(4), 545–553. https://doi.org/10.1007/s11258-012-0020-x

557 Guan, D., McCarthy, S. A., Wood, J., Howe, K., Wang, Y., & Durbin, R. (2020). Identifying and removing

558   haplotypic duplication in primary genome assemblies. *Bioinformatics*, *36*(9), 2896–2898.

559   https://doi.org/10.1093/bioinformatics/btaa025

560 Haley, C. S., & Knott, S. A. (1992). A simple regression method for mapping quantitative trait loci in line

561   crosses using flanking markers. *Heredity*, *69*(4), 315–324. https://doi.org/10.1038/hdy.1992.131

562 Heiser Jr., C. B., & Whitaker, T. W. (1948). Chromosome Number, Polyploidy, and Growth Habit in

563   California Weeds. *American Journal of Botany*, *35*(3), 179–186. https://doi.org/10.1002/j.1537-

2197.1948.tb05204.x

Hierro, J. L., Villarreal, D., Eren, Ö., Graham, J. M., & Callaway, R. M. (2006). Disturbance Facilitates

Invasion: The Effects Are Stronger Abroad than at Home. *The American Naturalist*, *168*(2), 144–

156. https://doi.org/10.1086/505767

Holt, C., & Yandell, M. (2011). MAKER2: An annotation pipeline and genome-database management tool

for second-generation genome projects. *BMC Bioinformatics*, *12*(1), 491.

https://doi.org/10.1186/1471-2105-12-491

Huang, K., Andrew, R. L., Owens, G. L., Ostevik, K. L., & Rieseberg, L. H. (2020). Multiple chromosomal

inversions contribute to adaptive divergence of a dune sunflower ecotype. *Molecular Ecology*,

29(14), 2535–2549. https://doi.org/10.1111/mec.15428

Huang, K., & Rieseberg, L. H. (2020). Frequency, Origins, and Evolutionary Role of Chromosomal

Inversions in Plants. *Frontiers in Plant Science*, *11*.

https://www.frontiersin.org/articles/10.3389/fpls.2020.00296

Irimia, R.-E., Montesinos, D., Eren, Ö., Lortie, C. J., French, K., Cavieres, L. A., Sotes, G. J., Hierro, J. L.,

Jorge, A., & Loureiro, J. (2017). Extensive analysis of native and non-native Centaurea solstitialis

L. populations across the world shows no traces of polyploidization. *PeerJ*, *5*, e3531.

https://doi.org/10.7717/peerj.3531

Jay, P., Tezenas, E., Véber, A., & Giraud, T. (2022). Sheltering of deleterious mutations explains the

stepwise extension of recombination suppression on sex chromosomes and other supergenes.

PLOS Biology, *20*(7), e3001698. https://doi.org/10.1371/journal.pbio.3001698

Kirubakaran, T. G., Grove, H., Kent, M. P., Sandve, S. R., Baranski, M., Nome, T., De Rosa, M. C., Righino,

B., Johansen, T., Otterå, H., Sonesson, A., Lien, S., & Andersen, Ø. (2016). Two adjacent

inversions maintain genomic differentiation between migratory and stationary ecotypes of

Atlantic cod. *Molecular Ecology*, *25*(10), 2130–2143. https://doi.org/10.1111/mec.13592

Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics*, *5*(1), 59.

https://doi.org/10.1186/1471-2105-5-59

Laetsch, D. R., & Blaxter, M. L. (2017). *BlobTools: Interrogation of genome assemblies* (6:1287).

F1000Research. https://doi.org/10.12688/f1000research.12232.1

Lajoie, B. R., Dekker, J., & Kaplan, N. (2015). The Hitchhiker's Guide to Hi-C Analysis: Practical guidelines.

Methods (San Diego, Calif.), *72*, 65–75. https://doi.org/10.1016/j.ymeth.2014.10.031

Lee, C.-R., Wang, B., Mojica, J. P., Mandáková, T., Prasad, K. V. S. K., Goicoechea, J. L., Perera, N.,

Hellsten, U., Hundley, H. N., Johnson, J., Grimwood, J., Barry, K., Fairclough, S., Jenkins, J. W., Yu,

Y., Kudrna, D., Zhang, J., Talag, J., Golser, W., … Mitchell-Olds, T. (2017). Young inversion with

multiple linked QTLs under selection in a hybrid zone. *Nature Ecology & Evolution*, *1*(5), 1–13.

https://doi.org/10.1038/s41559-017-0119

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics, *25*(14), 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., & Durbin, R.

(2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079.

https://doi.org/10.1093/bioinformatics/btp352

Maddox, D. M., Mayfield, A., & Poritz, N. H. (1985). Distribution of Yellow Starthistle (Centaurea

solstitialis) and Russian Knapweed (Centaurea repens). *Weed Science*, *33*(3), 315–327.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO Update: Novel and

Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of

Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution*, *38*(10), 4647–

4654. https://doi.org/10.1093/molbev/msab199

Miller, M. R., Dunham, J. P., Amores, A., Cresko, W. A., & Johnson, E. A. (2007). Rapid and cost-effective

polymorphism identification and genotyping using restriction site associated DNA (RAD)

markers. *Genome Research*, *17*(2), 240–248. https://doi.org/10.1101/gr.5681207

Montesinos, D., & Callaway, R. M. (2017). Inter-regional hybrids of native and invasive Centaurea

solstitialis display intermediate competitive ability. *Ecography*, *40*(7), 801–802.

https://doi.org/10.1111/ecog.02653

Montesinos, D., Graebner, R. C., & Callaway, R. M. (2019). Evidence for evolution of increased

competitive ability for invasive Centaurea solstitialis, but not for naturalized C. calcitrapa.

Biological Invasions, *21*(1), 99–110. https://doi.org/10.1007/s10530-018-1807-z

Mota, L., Torices, R., & Loureiro, J. (2016). The Evolution of Haploid Chromosome Numbers in the

Sunflower Family. *Genome Biology and Evolution*, *8*(11), 3516–3528.

https://doi.org/10.1093/gbe/evw251

Peischl, S., Kirkpatrick, M., & Excoffier, L. (2015). Expansion Load and the Evolutionary Dynamics of a

Species Range. *The American Naturalist*, *185*(4), E81–E93. https://doi.org/10.1086/680220

Price, A. L., Jones, N. C., & Pevzner, P. A. (2005). De novo identification of repeat families in large

genomes. *Bioinformatics (Oxford, England)*, *21 Suppl 1*, i351-358.

https://doi.org/10.1093/bioinformatics/bti1018

Putnam, N. H., O'Connell, B. L., Stites, J. C., Rice, B. J., Blanchette, M., Calef, R., Troll, C. J., Fields, A.,

Hartley, P. D., Sugnet, C. W., Haussler, D., Rokhsar, D. S., & Green, R. E. (2016). Chromosome-

scale shotgun assembly using an in vitro method for long-range linkage. *Genome Research*,

26(3), 342–350. https://doi.org/10.1101/gr.193474.115

Reatini, B., & Vision, T. J. (2020). Genetic architecture influences when and how hybridization

contributes to colonization. *Evolution*, *74*(8), 1590–1602. https://doi.org/10.1111/evo.13972

Reyes-Chin-Wo, S., Wang, Z., Yang, X., Kozik, A., Arikit, S., Song, C., Xia, L., Froenicke, L., Lavelle, D. O.,

Truco, M.-J., Xia, R., Zhu, S., Xu, C., Xu, H., Xu, X., Cox, K., Korf, I., Meyers, B. C., & Michelmore, R.

W. (2017). Genome assembly with in vitro proximity ligation data and whole-genome triplication

in lettuce. *Nature Communications*, *8*(1), 14953. https://doi.org/10.1038/ncomms14953

Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2: Analytical methods for paired-end

sequencing improve RADseq-based population genomics. *Molecular Ecology*, *28*(21), 4737–

4754. https://doi.org/10.1111/mec.15253

Ruan, J., & Li, H. (2020). Fast and accurate long-read assembly with wtdbg2. *Nature Methods*, *17*(2),

155–158. https://doi.org/10.1038/s41592-019-0669-3

Scaglione, D., Reyes-Chin-Wo, S., Acquadro, A., Froenicke, L., Portis, E., Beitel, C., Tirone, M., Mauro, R.,

Lo Monaco, A., Mauromicale, G., Faccioli, P., Cattivelli, L., Rieseberg, L., Michelmore, R., &

Lanteri, S. (2016). The genome sequence of the outbreeding globe artichoke constructed de

novo incorporating a phase-aware low-pass sequencing strategy of F1 progeny. *Scientific*

Reports, *6*(1), 19427. https://doi.org/10.1038/srep19427

Stanke, M., Keller, O., Gunduz, I., Hayes, A., Waack, S., & Morgenstern, B. (2006). AUGUSTUS: Ab initio

prediction of alternative transcripts. *Nucleic Acids Research*, *34*(Web Server issue), W435–W439.

https://doi.org/10.1093/nar/gkl200

Tong, C., Li, H., Wang, Y., Li, X., Ou, J., Wang, D., Xu, H., Ma, C., Lang, X., Liu, G., Zhang, B., & Shi, J.

(2016). Construction of High-Density Linkage Maps of Populus deltoides × P. simonii Using

Restriction-Site Associated DNA Sequencing. *PLOS ONE*, *11*(3), e0150692.

https://doi.org/10.1371/journal.pone.0150692

Webb, D. M., & Knapp, S. J. (1990). DNA extraction from a previously recalcitrant plant genus. *Plant*

Molecular Biology Reporter, *8*(3), 180–185. https://doi.org/10.1007/BF02669514

Widmer, T. L., Guermache, F., Dolgovskaia, M. Y., & Reznik, S. Ya. (2007). Enhanced Growth and Seed

Properties in Introduced vs. Native Populations of Yellow Starthistle (Centaurea solstitialis).

Weed Science, *55*(5), 465–473.

Yang, J., Zaitlen, N. A., Goddard, M. E., Visscher, P. M., & Price, A. L. (2014). Advantages and pitfalls in

the application of mixed-model association methods. *Nature Genetics*, *46*(2), 100–106.

https://doi.org/10.1038/ng.2876

Zavaleta, E. S., & Hulvey, K. B. (2004). Realistic Species Losses Disproportionately Reduce Grassland

Resistance to Biological Invaders. *Science*, *306*(5699), 1175–1177.

https://doi.org/10.1126/science.1102643

Acknowledgements

666   We thank B.M. Anderson, S. Lin and K. Nurkowski for help with data collection and D.

667   Kaplan for assistance with greenhouse logistics. This work was supported by funding from a

668   Natural Sciences and Engineering Research Council of Canada grant #353026 to L.H.R., United

669   States National Science Foundation grants #1750280 to KMD and #2109625 to BR.

Author Contributions

672   BR, KD, MB, and LR conceived the work. BR, KD, AC, QJ, and MM conducted the research

673   and analyzed the resulting data. BR and KD drafted the manuscript, and all authors provided

674   revisions and feedback.

Tables and Figures

Tables

678   Table 1. Statistics of the *Centaurea solstitialis* genome assembly and annotation.

Metric	Initial PacBio assembly	Final Omni-C Assembly
Contiguity		
Total length	764,896,586	765,086,668
Total number of scaffolds	2,969	1,080
N50	1,356,249	100,696,929
L50	127	4
L90	873	8

Complete BUSCOs	231 (90.98%)	237 (92.9%)
Complete and single copy BUSCOs	185 (72.55%)	184 (72.1%)
Annotation		
Total repetitive DNA	--	63.31%
Transposable elements	--	31.1%
Simple repeats	--	1.16%
Total number of genes	--	34,323
Total coding region (bp)	--	59,845,476
Average gene length (bp)		1,743

680 Table 2. Summary of QTL peaks associated with maximum leaf length (len) and total number of leaves

681 (num) at time points 1 and 2 (e.g. len1 and len2, respectively, or just len if the QTL is associated with the

682 trait at both time points). The identification method, location in the reference, size of the peak, favored

683 allele (i.e. TRI for invader, TK for native, and H for overdominance), dominance pattern at the locus, and

684 number of candidate genes within 1 LOD drop of the peak are all given for each peak.

PeakID	method	trait	chr	start	end	size (Mbp)	favored_ allele	dominance_p attern	#candid ates
1	LOCO	len_n um	Chr_1	74748	889798 19	88.90 507	H	overdominan t	3532
2	HK	len	Chr_1	106564 625	106770 853	0.206 228	H	overdominan t	16
3	LOCO	num	Chr_2	435326 1	125267 28	8.173 467	TRI	additive	495
4	LOCO	num2	Chr_2	809990 70	990916 74	18.09 26	TRI	additive	880
5	LOCO	num1	Chr_2	953343 65	102637 104	7.302 739	TRI	additive	370
6	LOCO	len	Chr_4	179749 5	747097 6	5.673 481	TK	additive	407
7	LOCO	len	Chr_4	206908 37	755431 91	54.85 235	TK	additive	2198
8	LOCO	len1	Chr_5	602372 73	683941 49	8.156 876	TK	complete	391
9	LOCO	len	Chr_6	759346 95	814965 78	5.561 883	TK	complete	364
10	LOCO	len	Chr_7	614837 0	984538 5	3.697 015	TRI	complete	282
11	HK	num2	Chr_7	192456 30	744199 50	55.17 432	TRI	additive	2102

| 12 | LOCO | len | Chr_8 | 1027923 | 2687378 | 1.659455 | TK | complete | 147 |
| 13 | LOCO | len_num | Chr_8 | 10425376 | 16906147 | 6.480771 | TK | complete | 353 |

Figures

Figure 1. Scaffolding to the chromosome-level of the *C. solstitialis* genome using Omni-C data via the dovetail HiRise pipeline. A) Cumulative length of scaffolds in the final HiRise assembly versus the initial PacBio HiFi assembly. B) Link density of read pairs from the Dovetail Omni-C data mapped to the final *C. solstitialis* assembly. X and Y-axes represent the mapping positions of the first and second read in each read pair, respectively. Positions are binned, and link density within bins are represented as a heat map. Grey bars denote boundaries between scaffolds.

Figure 2. Gene synteny between yellow starthistle (*C. solstitialis*; n=8), globe artichoke (*Cynara cardunculus*; n=17), and lettuce (*Lactuca sativa*; n=9) genomes reveal chromosome evolution within the thistle subfamily (*Carduoideae*). A) Genome-wide synteny between yellow starthistle and artichoke displaying synteny blocks with match score of at least 2866. B) Genome-wide synteny between yellow starthistle and lettuce displaying synteny blocks with match score of at least 981. C) Putative fission of an ancestral chromosome into three separate chromosomes in globle artichoke revealed by three-way comparisons of gene synteny. D) Putative chromosome fusion between two ancestral chromosomes into one chromosome in yellow starthistle revealed by three-way comparisons of gene synteny. E) Example of a highly conserved chromosome between all three taxa revealed by three-way comparisons of gene synteny.

Figure 3. Summary of genome structure and QTL analyses. Panel A) Collinearity between the physical map on the x-axis and genetic map on the y-axis reveals suppressed recombination in putative pericentromeric regions (vertical grey bars) for all chromosomes and putative structural variation on the arms of chromosomes 1 and 7. Panel B) centromere genome scan score heatmap of the average gene density, unique repetitive element density, Gypsy LTR-RT density, and Copia LTR-RT density across each chromosome. Panel C) QTL LOD scores for maximum leaf length at time point 1 (black lines) and time point 2 (green lines) using the HK method (dashed lines) and LOCO method (solid lines). Panel D) QTL LOD scores for total number of leaves at time point 1 (black lines) and time point 2 (green lines) using the HK method (dashed lines) and LOCO method (solid lines). Horizontal lines represent the suggestive QTL threshold of 0.1 for each time point and method.

Figure 4. Visualization of the reference-aligned genetic map. A) Heatmap of recombination fractions (RF) in the top left of the diagonal and logarithm of odds (LOD) scores in the bottom right of the diagonal for ordered markers in linkage groups 1-8 reveal strong linkage and low recombination for markers within linkage groups and weak linkage with high recombination for markers on different linkage groups. Each linkage group is labeled with the corresponding chromosome number from the reference genome. B) Physical distribution of markers from the genetic map across chromosomes 1-8 of the reference genome, with each linkage group displayed as a different color. Putative centromere locations from genome scans are indicated with black dots.

714   **Supplemental Material**

Supplemental Figure 1. Annotation Edit Distance (AED) for the *C. solstitialis* genome annotation.

Supplemental Figure 2. Normalized density of genes, unique repetitive elements, Gypsy LTR-RT elements, and Copia LTR-RT elements across the genome.

Supplemental Figure 3. Phenotypic values for invader parental genotype (TRI), native parental genotype (TK), F1, and F2 genotypes of the mapping population.

Supplemental Figure 4. Distribution of corrected values for maximum leaf length and total number of leaves at both time points for the 300 genotyped F2s used as input for QTL analysis.