USP22 overexpression fails to augment tumor formation in MMTV-ERBB2 mice but loss of

function impacts MMTV promoter activity

Xianghong Kuang[1,2], Andrew Salinger[1, 2], Fernando Benavides[1, 2], William J. Muller[3, 4, 5], Sharon

Y. R. Dent[1, 2, 6] * and Evangelia Koutelou[1, 2] *

[1]Department of Epigenetics and Molecular Carcinogenesis, The University of Texas MD

Anderson Cancer Center, Houston, TX 77054, USA

[2] Center for Cancer Epigenetics, The University of Texas MD Anderson Cancer Center, Houston,

TX 77030, USA

[3] Rosalind and Morris Goodman Cancer Institute, McGill University, Montreal, Canada.

[4] Department of Biochemistry, McGill University, Montreal, Canada.

[5] Faculty of Medicine, McGill University, Montreal, Canada.

[6] The University of Texas MD Anderson Cancer Center/UTHealth Houston Graduate School of

Biomedical Sciences

*****Corresponding authors:** Sharon Y. R. Dent and Evangelia Koutelou, Department of Epigenetics

and Molecular Carcinogenesis, The University of Texas MD Anderson Cancer Center, PO Box

301429, Unit 1951 Houston, Texas 77230, USA. Phone number: 713-563-7742. Email:

sroth@mdanderson.org (S.Y.R.D.); ekoutelou@mdanderson.org (E.K.)

Running title: Usp22 impacts on MMTV-NIC tumorigenesis

Key Words: Usp22, MMTV, Mammary gland tumorigenesis, ERBB2, Neu

Funding resource: The National Institutes of Health grant R01HD094400 to S.Y.R. Dent.

Abstract

The USP22 deubiquitinase, a component of the SAGA histone modifying complex, is overexpressed in multiple human cancers, but how USP22 impacts tumorigenesis is not clear. We reported previously that *Usp22* loss in mice impacts execution of several signaling pathways driven by growth factor receptors such as ERBB2. To determine whether changes in USP22 expression affects ERBB2-driven tumorigenesis, we introduced conditional overexpression or deletion alleles of *Usp22* into mice bearing the MMTV-NIC transgene, which drives both rat ERBB2/NEU expression and Cre recombinase activity from the MMTV promoter resulting in mammary tumor formation. We found that USP22 overexpression in mammary glands did not further enhance primary tumorigenesis in MMTV-NIC female mice, but increased lung metastases were observed. However, deletion of *Usp22* significantly decreased tumor burden and increased survival of MMTV-NIC mice. These effects were associated with markedly decreased levels of both *Erbb2* mRNA and protein, indicating *Usp22* loss impacts MMTV promoter activity. *Usp22* loss had no impact on ERBB2 expression in other settings, including MCF10A cells bearing a CMV-driven *ERBB2* transgene or in HER2+ human SKBR3 and HCC1953 cells. Decreased activity of the MMTV promoter in MMTV-NIC mice correlated with decreased expression of known regulatory factors, including the glucocorticoid receptor (GR), the progesterone receptor (PR), and the chromatin remodeling factor BRG1. Together our findings indicate that increased expression of USP22 does not augment the activity of an activated ERBB2/NEU transgene, but impacts of Usp22 loss on tumorigenesis cannot be assessed in this model due to unexpected effects on MMTV-driven Erbb2/Neu expression.

Significance

Overexpression of USP22 alone is not sufficient to promote MMTV-ERBB2-driven tumor formation but may enhance metastasis. *Usp22* loss impacts expression of GR, PR and BRG1, decreasing MMTV promoter activity.

Introduction

USP22 overexpression occurs in several types of highly aggressive human cancers and is associated with a "death from cancer" gene signature for therapy resistance and stem cell like phenotypes (1-3). How USP22 impacts cancer development and progression, however, is still not well defined.

USP22 provides the catalytic activity of the deubiquitinase (DUB) module of the SAGA histone modifying complex. SAGA complex is multifunctional, housing both DUB and histone lysine acetyltransferase (KAT) activities, along with factors that interact directly with either sequence specific or general transcription factors to activate genes. Both the KAT and DUB modules of SAGA have been implicated in human cancers (1). The catalytic subunit of the KAT module, GCN5/KAT2A, is an important co-activator of MYC target genes during normal mouse development and in human cancer cells (4-6). Deletion or inhibition of *Gcn5* impacts growth and survival of MYC-driven cancers, including lung cancers and lymphoma, in both mouse models and human cells (5,7). USP22 functions have also been linked to MYC and other oncogenes, including BMI1, SIRT1, cyclin B1 and KDM1A (8-11). These findings support a role for USP22 and SAGA in promoting cell proliferation. USP22 impact on proliferation is further indicated by connections between USP22, the androgen receptor and c-MYC in prostate cancer cells and models (12). Other studies, however, indicate that USP22 also functions to suppress tumor formation in particular contexts, such as colorectal cancer (13). In all cases, the functions of USP22 in cancer may reflect its activity towards histone substrates and gene regulation or to removal of ubiquitin from non-histone substrates by USP22 that may impact other processes, such as telomere stability or DNA repair (14,15).

In order to better define USP22 functions in cancer, we created mouse models to determine its functions during normal development. Deletion of *Usp22* leads to embryonic lethality due to defects in placental vasculature (16). Gene expression changes indicate that these defects are linked to aberrant execution of multiple receptor tyrosine kinase signaling cascades as well as TGFβ signaling that drive propagation of endothelial cells, specification of perivascular cells, and cell migration (16). These same pathways and processes are often associated with tumorigenesis, so we also created a Lox-Stop-Lox mouse model that overexpresses USP22 in all tissues to determine whether overexpression is sufficient for induction of these pathways. We observed over branching of the mammary glands in USP22 overexpressing mice, along with

aberrant up regulation of many of the same signaling pathways that were down regulated in *Usp22*

null embryos, including estrogen receptor, ERK/MAPK, and TGFβ signaling (16,17). However,

USP22 overexpression was not sufficient to induce tumors in the mammary gland or in any other

tissues (17), suggesting this DUB likely works with specific oncogenes to promote tumorigenesis.

ERBB2 was one of the signaling pathways impacted by *Usp22* loss (16). Overexpression of

ERBB2 (also known as HER2/NEU) is associated with human breast cancers, and mouse models

of HER2+ tumors, including MMTV-NIC transgenic mice (expressing an activated oncogenic

human *ERBB2*), have provided many insights into how ERBB2 induces tumors and promotes

metastases (18-20). One previous study by others indicated that USP22 regulates the unfolded

protein response in HER2+ tumors (21). Here we combine the MMTV-NIC transgene with either

USP22 overexpression or *Usp22* loss to determine whether and how this DUB impacts ERBB2

driven oncogenesis in mice.

Materials and Methods

Mouse models

To generate bigenic mice that specifically overexpress Usp22 in mouse mammary gland

epithelium (*Usp22*$^{OE/OE}$; MMTV-NIC**)**, we crossed our lab generated *Rosa26KI-Usp22-LSL* mice

(congenic FVB/N background) (17) with MMTV-NIC (MMTV-Neu (NDL2-5)-IRES-Cre, JAX Strain

032576) mice on pure FVB/N background (generous gift from Dr. William J. Muller) (22). We

identified mice bearing desired genotypes by PCR, using allele-specific primers, 5'

AAAGTCCCTATTGGCGTTACTA 3' (forward) and 5' AAAGTCGCTCTGAGTTGTTATC 3'

(reverse) for the knock-in allele with a product size of 388 bp. PCR primers for the wild-type

Rosa26 locus had the sequences 5' GGAGCGGGAGAAATGGATAT 3' (forward) and 5'

AAAGTCGCTCTGAGTTGTTATC 3' (reverse), producing a PCR product size of 602 bp. The

PCR program ran 35 cycles of 95°C, 5 min, for denaturing; 55°C, 40 sec, for annealing; and 72°C,

1 min, for extension.

To generate mice bearing a *Usp22* floxed allele, we obtained sperm bearing the *Usp22*$^{tm1a(KOMP)Wtsi}$

allele (C57BL/6N) from the Wellcome Trust KOMP consortium (www.komp.org) and performed *in*

vitro fertilization. The *Usp22*$^{tm1a(KOMP)Wtsi}$ allele contains two LoxP sites surrounding exon 2 of

Usp22 and a neomycin selection cassette surrounded by two short flippase recognition target

(FRT) sites. We crossed mice bearing the Usp22[tm1a(KOMP)Wtsi] allele with FLPeR mice (obtained

from the Genetically Engineered Mouse Facility, MD Anderson Cancer Center) expressing a

flippase expression construct (23), in order to remove the neomycin cassette. The resulting

Usp22[FL/+] or Usp22[FL/FL] mice were backcrossed several times onto FVB/NHsd (Envigo,

Indianapolis, IN) using marker-assisted (speed congenics) crosses (Laboratory Animal Genetic

Services, MD Anderson Cancer Center), and then crossed with MMTV-NIC mice (also in the

FVB/N background) to achieve mammary gland epithelial-specific Usp22 deletion. Genotypes

were determined by PCR using primers CSD-Usp22-F (forward): 5'

ACTTTGGGAAAGCTCGTATGTGTGC 3' and CSD-Usp22-ttR (reverse): 5'

GATGACATCATCACATTCCCACGCC 3' yielding a PCR product of 486bp for the wildtype allele

and 623bp for the floxed allele.

Detection of mammary gland tumors

Mouse mammary gland tumor onset in female mice was monitored by palpation twice weekly

starting from 10-weeks of age. Tumor size was measured by caliber and mice were sacrificed

when the largest tumor diameter reached 2 cm. Tumor volume was estimated with the formula:

volume = (width)^2 X length/2 (cm^3). All animal experiments were conducted under the guidelines

of the Institutional Animal Care and Use Committee of UT MD Anderson Cancer Center and

approved animal protocols.

Isolation of primary mouse mammary gland epithelial cells (MECs)

The left and right 2nd, 3rd, and 4th mammary glands were dissected from 100-day-old female

Usp22[FL/FL]-NIC and wildtype-NIC mice, with lymph nodes removed from the 4th glands. Organoids

were prepared from the dissected tissue as described (17,24) with modifications. Briefly, glands

were minced into paste and incubated in 5 mL DME/F-12 medium (HyClone, #SH30023.1) with

2 mg/mL collagenase A (Roche #10103578001), 100 units/mL hyaluronidase (Sigma-Aldrich,

H3506) for 1 h at 37°C with 180 rpm rotation. The resulting cell mixture was collected by

centrifugation and then washed with DME/F-12 medium and centrifuged twice at 450 g for 10 min

to remove the fatty layer. The cells were next incubated in 3 mL DME/F-12 medium with 2 units/mL

DNase I (Sigma-Aldrich, D4263) at room temperature for 3 min and centrifuged at 450 g for 10

min. The cell pellets were resuspended in 4 ml Red Blood Cell Lysis buffer (Ammonium Chloride

solution, Stemcell Technology, #07800) on ice for 2 min and washed with 6 mL DME/F-12 medium

at 450 g for 10 min. Differential centrifugation (pulse centrifugation to 450 g for 10 seconds) was

used to separate stroma from organoids. The supernatant from the second spin containing
fibroblasts, macrophages, and endothelial cells were discarded. The pellets from 2nd round pulse
centrifugations were organoids consisting mainly of mammary gland epithelial cells. The
organoids were then digested with 3 mL 0.05% Trypsin/EDTA (HyClone, #SH30236.1) with 2
unit/mL of DNase I for 3 min. An equal volume of DME/F-12 containing 10% fetal bovine serum
(FBS, Gibco, #10437-028) was added and the cells were dissociated by pipetting 20 times with a
1000 µL (size) tip. Isolated epithelial cells were then collected by 3 min centrifugation at 600g.

Cell lines and cell culture

MCF10A cells bearing either a vector control or an *ERBB2* overexpression construct (OE) were
generously provided by Dr. Dihua Yu (25). Cells were cultured in DME/F12 media supplemented
with 5% horse serum (Thermo Fisher Scientific, 16050122), 20 ng/mL hEGF (Sigma, E9644), 100
ng/mL cholera toxin (Sigma, C8052), 0.5 µg/mL hydrocortisone (Sigma, H0888), 10 µg/mL insulin
(Sigma, I1882), 50 units/mL penicillin, and 50 µg/mL streptomycin (Hyclone, #SV30010). DMEM
media (Hyclone #SH3002201) with 10% FBS was used to culture SKBR3 (the American Type
Culture Collection (ATCC), #HTB-30) and HCC1954 cells (ATCC, #CRL-2338).

shRNA knockdown *USP22* in SKBR3, HCC1954 and MCF10A-*ERBB2* OE cells

HEK293T cells at 70% confluency were co-transfected with lentiviral vectors expressing human
USP22 shRNA. Vectors used included *pGIPZ-GFP-Puro* (see Supplementary Materials) and
psPAX.2 and *pMD2.G* (Addgene). Cell transfections were done in Opti-MEM (Life Technologies,
#31985-070) using Lipofectamine 2000 (Life Technologies, Cat. # 11668019) following the
manufacturers' transfection reagent manual. Eight hours after the transfection, the medium was
changed to DMEM complete medium (DMEM medium with 10% FBS), and 48 hours post
transfection, the DMEM medium containing viral particles was collected and filtered through 0.45
µm filters. The virus containing medium with 8 µg/mL polybrene (Sigma-Aldrich, Cat #H9268) was
used in 1:1 dilution with the culture medium to infect SKBR3 and HCC1954 cells. 48 hours after
infection, the viral medium was changed to DMEM complete medium containing puromycin (EMD
Millipore, #540222) at 2 µg/mL for selection of infected cells. Since MCF10A-*ERBB2* cells are
already puromycin resistant, transfected cells were sorted on a BD FACSAria Fusion flow
cytometer (BD Biosciences) using a 100 µm tip at 70 psi to select for GFP+ cells (comparing

transfected cells to un-transfected cells) and were stained with Propidium Iodide to monitor

viability. Selected GFP+ cells were collected in a 15mL conical tube with 2mL FBS and then

expanded by cell culture in MCF10A culture medium.

CRISPR-Cas9 *USP22* knock out in MCF10A-ERBB2 cells

USP22 was deleted in MCF10A-*ERBB2* OE cells using Synthego Gene Knock out Kit v2,

Transfection Optimization Kit and Lipofectamine CRISPRMAX Transfection Reagent (Invitrogen,

#CMAX00003) according to Synthego's instructions (www.synthego.com/resources). Briefly, 70%

confluent MCF10A-*ERBB2* OE cells were trypsinized, washed and resuspended in OPTI-MEM

media. Ribonucleoprotein (RNP) complexes that consist of purified Cas9 nuclease duplexed with

chemically modified synthetic multi-guide sgRNA (sgRNA Usp22 KO, from Synthego's Gene

Knockout Kit v2) at a ratio of 1:1.3, or a non-sgRNA control in Opti-MEM media were then mixed

with lipofectamine CRISPRMAX transfection reagent at a volume ratio of 50:3. The RNP-

transfection solution mixture was then mixed with MCF10A-*ERBB2* OE cell suspensions and

seeded on cell culture plates. *USP22* knock out cells and non-sgRNA control cells were collected

and knock out efficiencies were validated by immunoblot analyses.

Immunoblotting

Isolated MECs, pelleted cells or tumor samples were lysed in RIPA buffer (150 mM NaCl, 0.1%

SDS, 50mM Tris (pH 7.4), 0.5% deoxycholic acid, 1% Nonidet P-40) with protease inhibitor

cocktail (Sigma, #P8340) and phosphatase inhibitors PhosSTOP (Roche, #04906837001). The

extracts were mixed with Laemmli buffer and boiled at 95°C for 5 min. The lysates (20 - 30 μg of

total protein) were subjected to electrophoresis on NuPAGE 4-12% Bis-Tris gels (Invitrogen,

NP0322BOX), transferred to PVDF membranes (BIO-RAD, #1620177), and probed with primary

and secondary antibodies. Antibody information is available in Supplementary Materials. ECL

reagents were from Amersham (#RPN2236).

RNA isolation and qRT-PCR

Total RNA from mouse tissues, cell lines or mammary gland epithelial cells were extracted using

RNeasy Plus Mini Kit (Qiagen, #74136). Extracts were then treated with RNase free DNase

(Qiagen, #79254). qRT-PCR products were measured using Power SYBR Green RNA-to-Ct 1-

step kit (Applied Biosystem, #4389986) and the 7500 Fast Real-Time PCR System (Applied

Biosystems). Sequences for gene-specific qRT-PCR primers used are listed in the

Supplementary Materials. Expression of *GAPDH* (human) or *Pbgd* (mouse) was used as internal

control. Data were normalized by the expression levels of internal control and the relative

expression of RNAs was calculated using the comparative Ct method.

Statistical analysis

All data were represented as mean ±standard error of mean. Calculations for qRT-PCR were

performed in Microsoft Excel. The Log-rank (Mantel–Cox) test was used as statistical analysis for

Kaplan-Meier tumor-free survival curves. All other statistical analysis was performed in Prism

9 (GraphPad Prism Software 9.0) using two-tailed unpaired t test. * p value <0.05, ** p< 0.01 and

***p<0.001.

Results

USP22 Overexpression in MMTV-NIC mice

To determine whether Usp22 overexpression might enhance ERBB2-driven formation of

mammary tumors, we introduced our previously described *Lox-Stop-Lox (LSL) Usp22* allele (17)

into mice carrying the MMTV-NIC transgene (**Fig. 1A**). The *LSL Usp22* allele inserted at the

Rosa26 locus allows tissue-specific overexpression (OE) of USP22 upon exposure to Cre

recombinase (17). The USP22 protein expressed from the *OE/LSL* allele (hereafter referred to

as *OE*) is fully functional and can fully rescue lethality of *Usp22* null embryos, but it is not sufficient

to induce tumors on its own (17). The MMTV-NIC transgene expresses both the ERBB2/NEU

oncoprotein and Cre from the MMTV promoter, assuring Cre expression, and thereby USP22

overexpression, in the same mammary epithelial cells at the same time as ERBB2/NEU

overexpression.

We monitored mammary tumor formation in MMTV-NIC mice bearing only the endogenous allele

of *Usp22* (*Usp22*[+/+]) or bearing one (*Usp22*[OE/+]) or two (*Usp22*[OE/OE]) copies of the *Usp22 OE*

transgene in addition to the endogenous allele (**Fig. 1B**). As expected (22,26), the MMTV-NIC

mice all developed mammary epithelial tumors between ~120-150 days (T_{50}, median latency

period of 126 days). The kinetics and frequency of tumor formation were not altered by Usp22

overexpression, in either the *Usp22*[OE/+]; NIC or *Usp22*[OE/OE]; NIC mice (**Fig. 1B**). No significant

changes were observed in tumor weight, tumor volume, or numbers of tumors per mouse upon

Usp22 overexpression (**Fig. 1C, D** and **E**).

Consistent with these results, no increase in ERBB2, phospho-ERBB2, or ERα protein levels

were observed in mammary epithelial cells isolated from USP22 overexpressing mice, relative to

cells isolated from MMTV-NIC mice (**Fig. 2A**). Interestingly, a trend for increased incidence of

lung metastases was observed in both $Usp22^{OE/+}$ - NIC or $Usp22^{OE/OE}$ - NIC mice (**Fig. 2B**), which

may be related to decreased expression of E-CADHERIN and increased level of SLUG in

$Usp22^{OE/OE}$ mice (**Fig. 2C**). Expression of other markers of the epithelial-mesenchymal transition,

however, were not significantly changed upon USP22 overexpression. Altogether, these results

indicate that increased expression of USP22 does not further enhance the tumorigenicity of the

activated ERBB2/NEU oncogene in MMTV-NIC mice.

Usp22 **deletion impacts tumor formation in MMTV-NIC mice**

We next sought to determine whether *Usp22* is required for tumorigenesis in MMTV-NIC mice

using the $Usp22^{flox}$ ($Usp22^{FL}$) allele, which allows tissue specific deletion of *Usp22* upon exposure

to Cre (**Fig. 3A**). Loss of one allele of *Usp22* ($Usp22^{FL/+}$) had no impact on the kinetics of tumor

formation in MMTV-NIC mice (**Fig. 3B**). However, complete deletion of *Usp22* significantly

delayed tumor formation (T_{50} 214 days) with some mice remaining tumor free for almost 300 days

(**Fig. 3B**). Loss of *Usp22* impacted both weight and volume of tumors and substantially reduced

the numbers of tumors formed per mouse (**Fig. 3C, D** and **E**). These data are consistent with

those of Prokakis et al (21), who also observed reduced tumor burden upon loss of *Usp22* using

a different MMTV-Neu mouse model.

Decreased expression of ERBB2 in MMTV-NIC mice

To determine the molecular mechanisms underlying decreased tumor formation, we isolated

mammary epithelial cells (MECs) from MMTV-NIC mice with or without the $Usp22^{FL}$ allele at an

early time point (100 days), before any tumors were observed in the MMTV-NIC mice.

Immunoblots confirmed partial and complete loss of USP22 protein in the $Usp22^{FL/+}$-NIC and

$Usp22^{FL/FL}$-NIC MECs, respectively. Remarkably, levels of GCN5 and ATXN7L3 were also

reduced in $Usp22^{FL/FL}$- NIC MECs, indicating potential loss of both the histone acetyltransferase

and the deubiquitinase modules in the SAGA complex upon *Usp22* deletion.

Loss of USP22 protein expression correlated with loss of *Usp22* mRNA in *Usp22^{FL/FL}*-NIC MECs

(**Fig. 4B**). Surprisingly, exogenous *Erbb2* RNA, protein, and phosphorylation levels were also

dramatically reduced upon loss of *Usp22* (**Fig. 4C** and **D**). Expression of endogenous mouse

ERBB3 was also decreased in *Usp22^{FL/FL}*- NIC MECs, consistent with cross regulation of ERBB2

and ERBB3 stability reported by others (22). No changes were observed in expression of heat

shock protein family members, HSP70, HSP90, or HSP90ab1 either in MECs or in end-stage

tumors isolated from *Usp22^{FL/FL}* - NIC mice (**Fig. 4D** and **E**). These findings are in contrast to

those of Prokakis et al, who found no changes in ERBB2/HER2 expression by

immunohistochemistry of ERBB2/NEU induced tumors from *Usp22^{FL/FL}* mice(21). This

discrepancy may reflect the different time points examined, as our analyses focused on events

before tumor formation rather than on the later tumor stages that did form in these mice.

The decreased expression of ERBB2/NEU observed in our *Usp22^{FL/FL}*-NIC mice is highly

consistent with the decreased tumor burden of these mice. However, these data reveal that the

impact of *Usp22* loss on tumorigenesis in this model system, and likely other MMTV driven NEU

mouse models, is due to changes in the activity of the MMTV promoter that drives expression of

the *Erbb2/Neu* transgene RNA (**Fig. 4C**).

***Usp22* depletion does not affect ERBB2/HER2 expression in human cells**

To assess whether *USP22* loss might also impact ERBB2 protein levels independently of *ERBB2*

gene transcription changes, we depleted *USP22* in human MCF-10A cells that overexpress

ERBB2. Two independent shRNAs targeting *USP22* significantly depleted both USP22 RNA and

protein, with no effects on ERBB2 RNA or protein expression (**Figure 5A, B** and **C**). We further

confirmed these results using CRISPR/Cas9 targeting of *USP22*, and again saw no impact on

ERBB2 expression upon deletion of *USP22* in the MCF-10A-ERBB2 OE cells (**Fig. 5D**). Finally,

we examined ERBB2 levels with and without *USP22* depletion in two HER2-positive human

breast cancer cell lines, SKBR3 and HCC1954, and again saw no impact on ERBB2 protein levels

upon *USP22* loss (**Fig. 5E**). These results indicate that *USP22* does not impact ERBB2

expression driven by non-MMTV exogenous promoters or by the endogenous *ERBB2* promoter.

However, our results do not rule out effects on downstream factors in the ERBB2 signaling

cascade that impact tumorigenesis (27).

Usp22 loss of function impacts multiple factors required for MMTV promoter activity

MMTV promoter activity is regulated by nuclear hormone receptors including glucocorticoid receptor (GR), progesterone receptor (PR), androgen receptor, as well as by ATP-dependent chromatin remodeling factors (28). The receptors bind to hormone response elements located in the MMTV promoter and then recruit the SWI/SNF BRG1 complex and other coactivators to activate the promoter (29-33). We compared expression of GR, PR, and BRG1 in MECs isolated from 100 day old _Usp22_$^{+/+}$- NIC and _Usp22_$^{FL/FL}$- NIC mice. Loss of Usp22 resulted in decreased levels of _Nr3c_1 transcript and protein (GR), and decreased levels of _PGR_ transcript _(PR)_ (**Fig. 6A, B** and **C**). Although _Smarca4_ transcript (Brg1) was increased in the absence of _Usp22_, protein levels were decreased (**Fig. 6C and D**), suggesting USP22 regulates BRG1 at a post-transcriptional level, directly or indirectly. Diminished levels of GR, PR and BRG1 are consistent with decreased MMTV promoter activity and decreased levels of Erbb2 observed in _Usp22_$^{FL/FL}$ - NIC mic_e._

Discussion

The role of USP22 overexpression in tumorigenesis and metastasis is an important question, as it is associated with highly aggressive, therapy resistant cancers. Our mouse models clearly established that overexpression of USP22 on its own is not sufficient to induce tumors (17). Here, we report that USP22 overexpression does not accelerate or amplify tumorigenesis in MMTV-NIC mice. This lack of effect could indicate a different mode of action for USP22 in tumor formation or progression, or it may reflect the already very high activity and expression of the rat NEU oncogene in these mice. The trend towards increased metastasis to the lung in USP22 OE MMTV-NIC mice may indicate that facilitates downstream steps in cancer progression, such as the epithelial-mesenchymal transition. Additional studies using a more attenuated oncogene are needed to determine if USP22 augments both tumorigenesis or metastasis.

Our finding that loss of _Usp22_ drastically hampers tumor formation in MMTV-NIC mice is consistent with the results of Prokaris et al, who used a similar mouse model wherein ERBB2 expression and Cre were driven by the MMTV promoter, but in separate transgenes. Our data clearly indicate that loss of _Usp22_ strongly impacts _Erbb2/Neu_ RNA and protein expression in

MECs isolated from 100-day old MMTV-NIC mice, before the time of the earliest tumor formation,
reflecting decreased activity of the MMTV promoter. Loss of exogenous EBB22/NEU expression
also impacted endogenous ERBB3 expression, further reducing tumorigenesis. Reduced levels
of ERBB2/NEU protein were also observed in end stage tumors from our MMTV-NIC *Usp22* null
mice, although more variability in expression was evident between mice and between tumors. In
contrast, Prokaris et al (21) reported no change in ERBB2 levels in their mice, although they only
presented measurements of levels of the oncogene in tumors by histology and did not examine
pre-tumor cells or tissues. These authors also reported down regulation of HSPA5, an HSP70
family member involved in the unfolded protein response, in tumors from *Usp22* null mice,
consistent with reports that USP22 mediates deubiquitination of HSPA5 in human prostate cancer
cells (15). Knock down of *USP22* in HCC1954 cells led to loss of HSPA5, indicating USP22 may
stabilize HSPA5 to suppress the unfolded protein response. Whether *Usp22* loss in pretumor
cells might impact tumor formation through changes in the UPR is not yet clear. Other studies by
this group indicated USP22 impacts stability of HSP90AB1 in prostate cancer and breast cancer
cells, including HCC1954 cells (34). We saw no effects on HSP90AB1, HSP90, or HSP70 protein
levels in our *Usp22* null MMTV-NIC mice at 100 days or on HSP90AB1 levels in end stage tumors
from these mice.
383
The impact on ERBB2/NEU expression levels that we observe is limited to the MMTV-NIC
transgene. Loss of *USP22* had no effect on *ERBB2* RNA or protein expression driven by the
CMV promoter in MCF10A cells or on endogenous ERBB2 protein expression in two HER2-
positive breast cancer cell lines, SKBR3 and HCC1954. However, these findings do not preclude
an impact of *Usp22* loss on downstream components of Erbb2 signaling. Indeed, deletion of
Usp22 in mice or *in vitro*, in induced endothelial cells, led to decreased expression of several
common intermediates to receptor tyrosine kinase signaling, including Gab1 and Gab2, and
decreased phosphorylation of AKT and ERK1/2 (16). Additional studies using different, non-
MMTV transgene mouse models are needed to determine the impact of *Usp22* loss on these
factors in mammary tumors and to define the mechanisms underlying such effects.
394
Multiple preclinical studies in HER2-positive cells have been performed to understand the
molecular mechanism of therapy resistance, but the use of cancer cell lines as the primary model
system in those studies does not recapitulate the complexity and heterogeneity of tumorigenesis
in organisms. The limitations of conventional cell culture and mouse xenograft studies are well-
recognized as an obstacle to the effective translation of preclinical findings into therapy (35).

Mouse models for HER2-driven mammary tumorigenesis have been used extensively and the
MMTV-NIC mouse is among the most commonly analyzed (36). Our work draws attention to
caveats in the interpretations of Erbb2 regulation when it is under the control of exogenous
promoters like MMTV. Loss of *Usp22* impacts MMTV promoter activity, likely through down
regulation of BRG1, PR and GR.　These factors themselves impact cancer formation and
progression. PR is a well-established breast cancer biomarker in evaluating prognosis and
response to treatment. GR plays key roles in breast cancer metastasis and response to treatment
(37). TCGA analysis reveals high frequency of BRG1 overexpression, but low frequency of BRG1
mutation, in invasive breast cancer (38,39). Given the importance of each of these factors in gene
regulation and cancer, further delineation of USP22 functions in regulation of BRG1, PR, and GR
could provide new insights to its roles in cancer formation and progression. The recent report of
novel USP22 specific inhibitors (40) provides strong impetus for fully understanding the functions
of this DUB in both normal cells and in cancers.

413
414
415　**Conflicts of Interest**

416　The authors have no competing financial interests to disclose.

417
418　**Acknowledgments**
419

We would like to thank Pam Whitney at the Flow Cytometry and Cell Imaging Core Facility-
Department of Epigenetics and Molecular Carcinogenesis (EMC), MD Anderson Cancer Center
and CPRIT Core Facility Support Grant (RP170628) for the support on GFP+ cell sorting
experiments. We thank Debra Hollowell for assistance with mouse *in vitro* fertilization. We thank
Dr. Lezlee Coghlan, Amanda Martin and animal facility staff for their help in determining tumor
size and animal husbandry. We also thank Dr. Dihua Yu (MD Anderson Cancer Center) for
providing us MCF10A-vec and MCF10A-*ERBB2* OE cell lines, and Dr. László Tora for providing
the anti-ATXN7L3 antibody. We would also like to thank all members of the Dent lab for their
advice in experimental procedures. This work was largely supported by the National Institutes of
Health grant R01HD094400 to S.Y.R. Dent. This study made use of the Research Animal Support
Facility-Smithville (including Laboratory Animal Genetic Services), supported by P30 CA016672
DHHS/NCI Cancer Center Support Grant to MD Anderson Cancer Center.

432
433
434

Authors' Contributions

Conception and design: X. Kuang, S.Y.R. Dent and E. Koutelou.

Methodology: X. Kuang, F. Benavides, W.J. Muller, S.Y.R. Dent and E. Koutelou.

Data curation: X. Kuang and E. Koutelou.

Formal analysis: X. Kuang., S.Y.R. Dent. and E. Koutelou.

Investigation: X. Kuang.

Validation: X. Kuang., A. Salinger.

Project administration: X. Kuang, S.Y.R. Dent. and E. Koutelou.

Supervision: S.Y.R. Dent and E. Koutelou.

Funding acquisition: S.Y.R. Dent.

Writing—original draft: X. Kuang, S.Y.R. Dent and E. Koutelou.

Writing, review, and/or revision of the manuscript: X. Kuang, S.Y.R. Dent and E. Koutelou. All

authors have read and agreed to this submitted version of the manuscript.

    Figure legends

    Figure 1.

    Overexpression of USP22 does not augment mammary gland tumor formation in MMTV-NIC

    mice. **A**. Schematic representation of *Rosa26KI-Usp22-LSL* mice that were crossed with MMTV-

    NIC mice to generate USP22-OE-MMTV-NIC progeny. **B**. Kaplan-Meier curve showing tumor-

    free survival, comparing *Usp22$^{+/+}$*- NIC (n=41), *Usp22$^{OE/+}$*- NIC (n=37) and *Usp22$^{OE/OE}$*- NIC (n=39)

    mice. p=0.012 was determined by Log-rank (Mantel–Cox) test, comparing *Usp22$^{OE/OE}$*- NIC mice

    with *Usp22$^{+/+}$*-NIC mice. **C-E**. Tumor weight (**C**)**,** tumor volume (**D**) and **n**umber of tumors per

    animal (**E**) were measured in *Usp22$^{+/+}$*-NIC (n=7), *Usp22$^{OE/+}$*-NIC (n=5) and *Usp22$^{OE/OE}$*-NIC mice

    (n=6). Tumors were harvested 45-55 days after tumor was first palpable (tumor onset). ns: No

    statistical significance when compared *Usp22$^{OE/OE}$-NIC* mice with *Usp22$^{+/+}$-NIC* mice using two-

    tailed unpaired t test.

    Figure 2.

    Expression levels of ERBB2 and ERα proteins and incidence of lung metastases in *Usp22 $^{OE/OE}$*-

    NIC mice. **A.** Immunoblot analyses using lysates from endpoint tumors (n=3 for each group)

    showed no difference in the levels of ERBB2 and ERα protein levels upon overexpression of

    USP22 in *Usp22$^{OE/OE}$*- NIC mice. Levels of phosphorylated ERBB2 protein are also unaffected by

    USP22 overexpression. **B.** Incidence of lung metastasis observed in *Usp22$^{+/+}$*-NIC (n=19),

    Usp22$^{OE/+}$-NIC (n=22) and *Usp22$^{OE/OE}$*- NIC (n=22) mice. **C**. Immunoblot analysis using lysates

    from endpoint tumors (n=3 for each group) showed decreased protein levels of E-CADHERIN and

    increased levels of SLUG protein in *Usp22 $^{OE/OE}$* NIC/+ mice when compared with *Usp22 $^{+/+}$*- NIC

    mice. EMT markers like OCCLUDIN, VIMENTIN and TWIST were not significantly affected by

    USP22 overexpression.

    Figure 3.

    Delayed tumor formation upon conditional deletion of *Usp22* in mammary gland epithelial cells in

    MMTV-NIC mice. **A.** Schematic representation *Usp22* floxed mice crossed with MMTV-NIC mice

    to generate of *Usp22$^{FL/FL}$-NIC* progeny. **B.** Kaplan-Meier curve showing tumor-free survival of

    animals, comparing *Usp22$^{+/+}$*- NIC (n=20), *Usp22$^{FL/+}$*- NIC (n=18) and *Usp22$^{FL/FL}$*- NIC (n=21)

    mice. p<0.0001 was determined by Log-rank (Mantel–Cox) test. **C-E**. Tumor weight (**C**), **t**umor

    volume (**D**) and number of tumors per animal (**E**) measured in *Usp22$^{+/+}$*- NIC (n=11; p=0.0015),

    Usp22$^{FL/+}$- NIC (n=5; p=0.0027) and *Usp22$^{FL/FL}$*- NIC mice (n=14; p<0.0001). P values were

513 determined by two-tailed unpaired t test when compared *Usp22^{FL/FL}* -NIC mice with *Usp22^{+/+}*-NIC

514 mice.

516 **Figure 4.**

517 Decreased expression of SAGA components, ERBB2, and ERBB3 upon deletion of *Usp22* in

518 mammary gland epithelial cells (MECs) collected from 100-day old NIC mice. **A.** Immunoblot

519 analyses showed decreased levels of USP22, GCN5 and ATXN7L3 proteins in MECs collected

520 from *Usp22^{FL/FL}-NIC* mice compared to *Usp22^{+/+}* -NIC and *Usp22^{FL/+}* -NIC mice. n=3 for each

521 group. **B, C.** qRT-PCR analysis of total RNA collected from *Usp22^{FL/FL}- NIC* MECs confirmed that

522 mRNA levels of *Usp22* and *Erbb2* were drastically decreased compared to *Usp22^{+/+}- NIC*.

523 p<0.0001 for both *Usp22* and *Erbb2*, as determined by two-tailed unpaired t test. n=5 for each

524 group. Each sample was measured in triplicates **D.** Immunoblot analyses showed decreased

525 levels of ERBB2 and ERBB3 proteins in MECs collected from *Usp22^{FL/FL}- NIC* mice compared to

526 *Usp22^{+/+}- NIC* and *Usp22^{FL/+}-* NIC mice. HSP70, HSP90AB1 and HSP90 protein levels were not

527 affected when analyzed by western blotting in the same lysates. n=3 for each group. **E.**

528 Immunoblot analyses using lysates from endpoint tumors from *Usp22^{FL/FL}- NIC* mice showed that

529 levels of USP22, ERBB2, ERBB3, phospho-ERK and H2Bub decreased relative to other

530 genotypes, but levels of HSP90AB1 and GCN5 did not change. n=3 for each group.

532 **Figure 5.**

533 Depletion or deletion of *USP22* in MCF10A cells that stably overexpress ERBB2 did not affect the

534 levels of ERBB2. **A.** Efficient depletion of *USP22* by lentiviral shRNA knock down in MCF10A-

535 ERBB2 cells as verified by qRT-PCR analysis. **B.** qRT-PCR analysis indicate levels of *ERBB2*

536 mRNA did not change upon *USP22* shRNA knock down. ns: No statistical significance. **C.**

537 Immunoblot analysis of MCF10A-ERBB2 cell lysates showed that ERBB2 protein levels are not

538 changed upon *USP22* shRNA knock down. **D.** Immunoblot analysis of MCF10A-ERBB2 cell

539 lysates indicate that ERBB2 protein levels are not changed upon *USP22* CRISPR/Cas9 knockout.

540 **E.** Immunoblot analysis of HER2-positive SKBR3 and HCC1954 cell lysates showed no difference

541 in ERBB2 protein levels upon *USP22* shRNA knock down.

543 **Figure 6.**

544 Expression levels of *Nr3c1* (glucocorticoid receptor), *Pgr* (progesterone receptor) and *Smarca4*

545 *(*BRG1*)* upon loss of *Usp22*. **A, B, C.** qRT-PCR analysis of total RNA collected from *Usp22^{FL/FL}-*

546 *NIC* MECs revealed decreased levels of *Nr3c1* and *Pgr,* but increased levels of *Smarca4 (Brg1)*

547   compared to *Usp22 $^{+/+}$- NIC* mice. p=0.0026 for *Nr3c1*, p=0.0053 for *PGR* and p=0.0012 for

548   *Smarca4* were determined by two-tailed unpaired t test. n=3-4 for each group. **D.** Immunoblot

549   analyses indicate decreased levels of GR and BRG1 proteins in MECs collected from *Usp22$^{FL/FL}$*-

550   NIC mice compared to *Usp22 $^{+/+}$*- NIC and *Usp22 $^{FL/+}$*- NIC mice. n=3 for each group.

References

1. Wang L, Dent SY. Functions of SAGA in development and disease. Epigenomics
2014;6:329-39
2. Glinsky GV. Death-from-cancer signatures and stem cell contribution to metastatic
cancer. Cell Cycle **2005**;4:1171-5
3. Glinsky GV, Berezovska O, Glinskii AB. Microarray analysis identifies a death-from-
cancer signature predicting therapy failure in patients with multiple types of cancer. J
Clin Invest **2005**;115:1503-21
4. Farria AT, Mustachio LM, Akdemir ZHC, Dent SYR. GCN5 HAT inhibition reduces
human Burkitt lymphoma cell survival through reduction of MYC target gene expression
and impeding BCR signaling pathways. Oncotarget **2019**;10:5847-58
5. Farria AT, Plummer JB, Salinger AP, Shen J, Lin K, Lu Y, *et al.* Transcriptional
Activation of MYC-Induced Genes by GCN5 Promotes B-cell Lymphomagenesis.
Cancer Res **2020**;80:5543-53
6. Wang L, Koutelou E, Hirsch C, McCarthy R, Schibler A, Lin K, *et al.* GCN5 Regulates
FGF Signaling and Activates Selective MYC Target Genes during Early Embryoid Body
Differentiation. Stem Cell Reports **2018**;10:287-99
7. Mustachio LM, Roszik J, Farria AT, Guerra K, Dent SY. Repression of GCN5 expression
or activity attenuates c-MYC expression in non-small cell lung cancer. Am J Cancer Res
2019;9:1830-45
8. Liu YL, Jiang SX, Yang YM, Xu H, Liu JL, Wang XS. USP22 acts as an oncogene by
the activation of BMI-1-mediated INK4a/ARF pathway and Akt pathway. Cell Biochem
Biophys **2012**;62:229-35
9. Lin Z, Yang H, Kong Q, Li J, Lee SM, Gao B, *et al.* USP22 antagonizes p53
transcriptional activation by deubiquitinating Sirt1 to suppress cell apoptosis and is
required for mouse embryonic development. Mol Cell **2012**;46:484-94
10. Gennaro VJ, Stanek TJ, Peck AR, Sun Y, Wang F, Qie S, *et al.* Control of CCND1
ubiquitylation by the catalytic SAGA subunit USP22 is essential for cell cycle
progression through G1 in cancer cells. Proc Natl Acad Sci U S A **2018**;115:E9298-e307
11. Zhou A, Lin K, Zhang S, Chen Y, Zhang N, Xue J, *et al.* Nuclear GSK3β promotes
tumorigenesis by phosphorylating KDM1A and inducing its deubiquitylation by USP22.
Nature cell biology **2016**;18:954-66
12. Schrecengost RS, Dean JL, Goodwin JF, Schiewer MJ, Urban MW, Stanek TJ, *et al.*
USP22 regulates oncogenic signaling pathways to drive lethal cancer progression. Cancer
Res **2014**;74:272-86
13. Kosinsky RL, Zerche M, Saul D, Wang X, Wohn L, Wegwitz F, *et al.* USP22 exerts
tumor-suppressive functions in colorectal cancer by decreasing mTOR activity. Cell
Death Differ **2020**;27:1328-40
14. Atanassov BS, Evrard YA, Multani AS, Zhang Z, Tora L, Devys D, *et al.* Gcn5 and
SAGA regulate shelterin protein turnover and telomere maintenance. Mol Cell
2009;35:352-64
15. McCann JJ, Vasilevskaya IA, Poudel Neupane N, Shafi AA, McNair C, Dylgjeri E, *et al.*
USP22 Functions as an Oncogenic Driver in Prostate Cancer by Regulating Cell
Proliferation and DNA Repair. Cancer Res **2020**;80:430-43

16. Koutelou E, Wang L, Schibler AC, Chao HP, Kuang X, Lin K, *et al.* USP22 controls
multiple signaling pathways that are essential for vasculature formation in the mouse
placenta. Development **2019**;146

17. Kuang X, McAndrew MJ, Mustachio LM, Chen YC, Atanassov BS, Lin K, *et al.* Usp22
Overexpression Leads to Aberrant Signal Transduction of Cancer-Related Pathways but
Is Not Sufficient to Drive Tumor Formation in Mice. Cancers (Basel) **2021**;13

18. Muller WJ, Sinn E, Pattengale PK, Wallace R, Leder P. Single-step induction of
mammary adenocarcinoma in transgenic mice bearing the activated c-neu oncogene. Cell
1988;54:105-15

19. Moody SE, Sarkisian CJ, Hahn KT, Gunther EJ, Pickup S, Dugan KD, *et al.* Conditional
activation of Neu in the mammary epithelium of transgenic mice results in reversible
pulmonary metastasis. Cancer Cell **2002**;2:451-61

20. Ursini-Siegel J, Schade B, Cardiff RD, Muller WJ. Insights from transgenic mouse
models of ERBB2-induced breast cancer. Nat Rev Cancer **2007**;7:389-97

21. Prokakis E, Dyas A, Grün R, Fritzsche S, Bedi U, Kazerouni ZB, *et al.* USP22 promotes
HER2-driven mammary carcinoma aggressiveness by suppressing the unfolded protein
response. Oncogene **2021**;40:4004-18

22. Siegel PM, Ryan ED, Cardiff RD, Muller WJ. Elevated expression of activated forms of
Neu/ErbB-2 and ErbB-3 are involved in the induction of mammary tumors in transgenic
mice: implications for human breast cancer. Embo j **1999**;18:2149-64

23. Farley FW, Soriano P, Steffen LS, Dymecki SM. Widespread recombinase expression
using FLPeR (flipper) mice. Genesis **2000**;28:106-10

24. Bray K, Gillette M, Young J, Loughran E, Hwang M, Sears JC, *et al.* Cdc42
overexpression induces hyperbranching in the developing mammary gland by enhancing
cell migration. Breast Cancer Res **2013**;15:R91

25. Lu J, Guo H, Treekitkarnmongkol W, Li P, Zhang J, Shi B, *et al.* 14-3-3zeta Cooperates
with ErbB2 to promote ductal carcinoma in situ progression to invasive breast cancer by
inducing epithelial-mesenchymal transition. Cancer Cell **2009**;16:195-207

26. Ursini-Siegel J, Hardy WR, Zuo D, Lam SH, Sanguin-Gendreau V, Cardiff RD, *et al.*
ShcA signalling is essential for tumour progression in mouse models of human breast
cancer. Embo j **2008**;27:910-20

27. Feng Y, Spezia M, Huang S, Yuan C, Zeng Z, Zhang L, *et al.* Breast cancer development
and progression: Risk factors, cancer stem cells, signaling pathways, genomics, and
molecular pathogenesis. Genes Dis **2018**;5:77-106

28. Kinyamu HK, Archer TK. Modifying chromatin to permit steroid hormone receptor-
dependent transcription. Biochim Biophys Acta **2004**;1677:30-45

29. Chen J, Kinyamu HK, Archer TK. Changes in attitude, changes in latitude: nuclear
receptors remodeling chromatin to regulate transcription. Mol Endocrinol **2006**;20:1-13

30. Fryer CJ, Archer TK. Chromatin remodelling by the glucocorticoid receptor requires the
BRG1 complex. Nature **1998**;393:88-91

31. Huang ZQ, Li J, Sachs LM, Cole PA, Wong J. A role for cofactor-cofactor and cofactor-
histone interactions in targeting p300, SWI/SNF and Mediator for transcription. Embo j
2003;22:2146-55

32. Johnson TA, Elbi C, Parekh BS, Hager GL, John S. Chromatin remodeling complexes
interact dynamically with a glucocorticoid receptor-regulated promoter. Mol Biol Cell
2008;19:3308-22

33.   Trotter KW, Archer TK. Reconstitution of glucocorticoid receptor-dependent
674   transcription in vivo. Mol Cell Biol **2004**;24:3347-58
675   34.   Kosinsky RL, Helms M, Zerche M, Wohn L, Dyas A, Prokakis E*, et al.* USP22-
676   dependent HSP90AB1 expression promotes resistance to HSP90 inhibition in mammary
677   and colorectal cancer. Cell Death Dis **2019**;10:911
678   35.   Sharpless NE, Depinho RA. The mighty mouse: genetically engineered mouse models in
679   cancer drug development. Nat Rev Drug Discov **2006**;5:741-54
680   36.   Taneja P, Frazier DP, Kendig RD, Maglic D, Sugiyama T, Kai F*, et al.* MMTV mouse
681   models and the diagnostic values of MMTV-like sequences in human breast cancer.
682   Expert Rev Mol Diagn **2009**;9:423-40
683   37.   Dietrich N, Hoffman JA, Archer TK. BAF Complexes and the Glucocorticoid Receptor
684   in Breast Cancers. Curr Opin Endocr Metab Res **2020**;15:8-14
685   38.   Bai J, Mei P, Zhang C, Chen F, Li C, Pan Z*, et al.* BRG1 is a prognostic marker and
686   potential therapeutic target in human breast cancer. PLoS One **2013**;8:e59772
687   39.   Marquez SB, Thompson KW, Lu L, Reisman D. Beyond Mutations: Additional
688   Mechanisms and Implications of SWI/SNF Complex Inactivation. Front Oncol
689   **2014**;4:372
690   40.   Morgan M, Ikenoue T, Suga H, Wolberger C. Potent macrocycle inhibitors of the human
691   SAGA deubiquitinating module. Cell Chem Biol **2022**;29:544-54.e4
692
693
694

A

Nr3c1 (Glucocorticoid receptor)

B

PGR (Progesterone receptor)

C

Smarca 4 (Brg1)

D