Asynchronous haltere input drives specific wing and head movements in *Drosophila*

Running title: Asynchronous haltere input

Michael J Rauscher (mjr67@case.edu) and Jessica L Fox (jlf88@case.edu)

Case Western Reserve University Dept. Biology

Keywords:

Animal Behavior, Drosophila, Haltere, Insect Flight, Wing Hitch, Multisensory Integration

Asynchronous haltere input drives specific wing and head movements in *Drosophila*

Running title: Asynchronous haltere input

Abstract

Halteres are multifunctional mechanosensory organs unique to the true flies (Diptera). A

set of reduced hindwings, the halteres beat at the same frequency as the lift-generating forewings

and sense inertial forces via mechanosensory campaniform sensilla. Though haltere ablation

makes stable flight impossible, the specific role of wing-synchronous input has not been

established. Using small iron filings attached to the halteres of tethered flies and an alternating

electromagnetic field, we experimentally decoupled the wings and halteres of flying *Drosophila*

and observed the resulting changes in wingbeat amplitude and head orientation. We find that

asynchronous haltere input results in fast amplitude changes in the wing ("wing hitches"), but

does not appreciably move the head. In multi-modal experiments, we find that wing and gaze

optomotor responses are disrupted differently by asynchronous input. These effects of wing-

asynchronous haltere input suggest that specific sensory information is necessary for maintaining

wing amplitude stability and adaptive gaze control.

Introduction

Animal locomotion, from crawling [1] to swimming [2] to walking [3] to flight [4],

typically requires coordinated motion of multiple body segments or appendages. Though this

coordination can stem from activity in a central pattern generator, proprioceptive input

modulates body motions to adapt each step [5,6]. Specific contraction timing of muscles

actuating multiple appendages can significantly change gait, and thus the mechanisms ensuring

limb coordination are essential to structuring adaptive locomotion.

In flying flies, wing [7] and haltere [8] proprioceptors provide crucial feedback allowing for active control of each wingstroke to counteract flies' inherent pitch instability [9]. The halteres are small club-shaped organs which have evolved from hindwings in true flies (Diptera). Long known to be required for stable flight [10], the halteres do not generate significant lift, but instead primarily detect inertial forces using campaniform sensilla at their base [8,11,12]. Afferent neurons from haltere campaniform sensilla entrain their firing to a preferred phase of the haltere stroke cycle [13,14], which in *Drosophila* and most other fly taxa is biomechanically linked to the wing stroke cycle with an antiphase synchrony [15,16].

Stroke-synchronous feedback from halteres is functionally important for wing-steering motoneurons and muscles [8,17,18]. In tethered flight, haltere ablation results in variable wing amplitudes and diminished wing optomotor responses [19–21]. In the blowfly *Calliphora*, some haltere afferents project directly to the motoneuron controlling wing steering muscle wB1 [22]. Stimulation of the haltere nerve can entrain firing of the motoneuron, which is normally more sensitive to wing reafference, if it occurs within a specified time relative to the wing nerve's firing [23]. Optogenetic activation of haltere muscles resets the firing phase of both the wB1 and wB2 muscles [18]. For these reasons, we hypothesized that asynchronous haltere input is insufficient for stable wing muscle output, and predicted that changing the phase of the haltere's stroke would cause wing amplitude instabilities distinct from those observed with ablation.

Neck motoneurons also receive input from haltere afferents [24,25], and mediate a similar suite of gaze-stabilizing reflexes [26]. Following bilateral haltere ablation, flies cannot modulate the gain of optomotor gaze responses [27]. However, the head's movements are driven primarily by vision and not halteres [28]. In some neck motoneurons, spiking responses to visual input are gated, but not modified, by haltere input: they require haltere oscillation but are

invariant to haltere frequency [29]. We thus hypothesized that stroke-synchronous haltere
information is not necessary for head movements.

Haltere amplitude manipulation results in smooth changes in wingstroke amplitude and
gaze direction [28]. However, flies often execute sharp changes in wing stroke amplitude that
can take on a variety of time courses. Fast, unilateral "wing hitches" can occur spontaneously
[30,31]. These contrast with longer saccades [32,33], which can be either spontaneous or driven
by visual input [34–37] and involve coordinated syndirectional motion of both wings and the
head [38–41]. Both saccades and wing hitches are distinct outputs of wing and neck motor
pathways, and observing how sensory modulation influences them can reveal pathways for
processing sensory input [42]

To examine how halteres drive wing and gaze behaviors, we experimentally decoupled
wing and haltere movement with both static and dynamic perturbations. By adding a small iron
mass to one haltere, we immobilized it, abolishing it as a source of mechanosensory information.
Then, using electromagnets, we drove the haltere at specific frequencies, moving it throughout
its natural range of motion asynchronously with the wings. An asynchronously-driven haltere
increased the likelihood of unilateral wing hitches as well as noise in the wing's amplitude, but
this did not occur in movements of the head. Optomotor responses of both head and wings were
significantly diminished. Our results show that asynchronous haltere input is insufficient for
wing amplitude stabilization and adaptive optomotor responses, and results in head responses of
diminished amplitude.

Materials and Methods

Animal care, haltere manipulations, and experimental trials

Animals were reared from a colony of wild-caught *Drosophila melanogaster*. Female

flies (3-5 days post-eclosion) were cold-anaesthetized and attached to tungsten pins using UV-

curing cement. For magnetic stimulation experiments, a single iron filing was attached to the

right haltere with UV-curing cement. Only iron filings of sufficient mass to immobilize the

haltere (in excess of ~12μg, [28]) were used for experiments. For ablation experiments, one or

both halteres were completely removed using fine forceps. For filing-treated animals, successive

presentations of different haltere stimuli were preceded and followed by control epochs with the

magnets disengaged lasting a minimum of 250ms. For each animal, 1-6 epochs from the same

experimental condition were drawn for analysis.

Haltere stimulation

To control haltere kinematics, two iron-core electromagnets were positioned anterior-

dorsal and posterior-ventral to the fly (figure 1*a*; [28]). Each electromagnet was powered from a

variable power supply (TP1803D, Tekpower, Montclair, CA, USA) switched through a power

transistor (D1276A, Panasonic, Kadoma, Japan) and controlled via 5V TTL-level output from a

data acquisition board (USB-6434, National Instruments, Austin TX, USA). Pulse trains

consisted of 50% duty cycle square wave signals alternating at 25, 36, 50, 72, 100, or 167Hz.

Signals to the magnets were 180° out of phase to produce a dorsal-ventral oscillation at the

commanded frequency.

Visual stimulation

To elicit optomotor responses, we used a wide-field vertical stripe pattern moving in the

yaw aspect under the control of a 1Hz triangle wave, presented using a modular LED flight arena

[43]. Stripes were one or two pixels wide and pseudo-randomly spaced such that average

luminance of the wide-field panorama was 50% [44]. Different pattern velocities were achieved

by varying the amplitude of the triangle wave whilst keeping the period constant.

High-speed video and kinematic measurements

Haltere and wing kinematics were recorded using two synchronized high-speed cameras

(TS4 or IL5, Fastec Imaging, San Diego, CA, USA) with sagittal views of the fly from the left

and right. Recordings were made at 1000 or 2000fps for all treated animals and the majority of

untreated animals; some untreated animals were recorded at 750fps. In each video, the haltere

bulb was digitized using the DLTdv8 [45] or DeepLabCut [46] software packages. Wingbeat

elevation time series were extracted using a custom MATLAB script. A third camera recording

at 100fps (Point Grey Chameleon3, FLIR, Wilsonville, OR, USA) provided a dorsal view to

measure head and wing kinematics. Head yaw and wing downstroke envelope kinematic time

series were computed using a custom MATLAB program called Flyalyzer

(https://github.com/michaelrauscher/flyalyzer; [28]).

Wing hitch detection and analysis

Wing hitchesCellini et al., 2021 were identified from baseline-subtracted video segments

using a custom MATLAB script based around the *findpeaks* function, selecting peaks with a

minimum amplitude of 4° (after the saccade detection algorithm in [39]). To capture hitches as

described by Götz and Heide [30,31] while excluding saccades, we screened putative hitches for

a minimum topographic prominence of 4°, and a maximum duration of 40ms. All detected

hitches were used to calculate the hitch rates shown in figure 2*c-d*, figure 4*c-d*, and

supplementary figure 2*d-i*. For the event-triggered average (ETA) responses shown in figure 2*e-*

h (and supplementary figure 2*a-e*) as well as the cyclic average responses shown in

supplementary figure 3, hitches were included only from video segments with concurrent high-

speed haltere and wing tracking from the lateral aspect cameras. Each 80ms event-triggered

window used to produce the ETA was baseline-subtracted and matched to a control window

selected with a randomly drawn frame index from the same video segment. To produce the

histograms shown in figure 3*b-g* and the heatmaps shown in supplementary figure 4, data were

pooled from all available asynchronously-driven stimulus categories (up to and including

167Hz). Each frame from the 100fps camera was assigned a haltere position and wing

instantaneous phase value at frame start by finding the single nearest-match frame recorded on

the high-speed lateral aspect cameras (1000 or 2000 fps). To assign a single angular velocity

value to each frame from the 100fps camera, the root-mean-square value of the angular velocity

time series was computed for all concurrent frames from the lateral aspect camera. Then, the sign

(positive or negative) of the lateral aspect frame with the largest absolute value was applied to

the result. A graphical summary of this process is included in supplementary figure 4*a*.

Frequency-domain analysis

For estimation of visual response magnitude (figure 5), kinematic time series were DC-

leveled by fitting a least-squares regression line and then subtracting it out to remove linear

trends, then zero-padded to 4096 samples. Magnitude estimates from resultant FFTs were then

divided by the number of samples in each time series prior to zero-padding to normalize

magnitudes for trials of varying length. Only segments longer than 64 samples were used,

resulting in a coarsest possible spectral resolution of 1.5Hz.

Results

Electromagnetic control of haltere stroke kinematics

To establish magnetic control over one haltere, an iron filing was attached to the bulb

(figure 1*a*). With the electromagnets deactivated, this mass immobilized the haltere and

prevented the fly from oscillating it. Activation of the electromagnets at 25, 36, 50, 72, 100, and

167 Hz resulted in movement of the haltere at the commanded frequency (figure 1*b-f*,

supplemental video 1). This method resulted in haltere movement both within and outside of its

naturalistic range (figure 1*c*), with all possible phase relationships to the wing. In all observed

trials, the untreated haltere maintained full stroke amplitude (figure 1*g*). At the highest frequency

tested (167 Hz), 13 out of 14 flies were unable to maintain a haltere amplitude that was within

the range of those observed naturally (figure 1*g*, 22 of 27 replicate trials). We thus excluded 167

Hz trials from further analyses that compared across conditions (see Methods). All conditions

achieved peak angular velocities within the range of those observed in untreated animals (figure

1*h*). Wingbeat frequencies were statistically indistinguishable from untreated controls for all

treatments (figure 1*i*), showing that manipulations did not impair the gross functioning of the

wing motor system.

Asynchronous haltere movement increases wing hitch rate

Haltere-manipulated flies showed significantly higher rates of wing hitches than

untreated control animals, for hitches reflecting wing amplitude increase as well as for hitches of

decreasing amplitude (figure 2*a-d*, supplemental video 2). All treatments exhibited higher rates

of ipsilateral wing hitches than untreated controls. For both amplitude-increasing hitches (figure

2*c*) as well as amplitude-decreasing hitches (figure 2*d*), dynamic haltere stimulation at any

frequency resulted in ipsilateral wing hitch rates comparable to those seen with bilateral haltere

ablation, whereas haltere immobilization resulted in intermediate hitch rates comparable to those

seen with unilateral ablation.

Were ipsilateral hitches accompanied by coordinated movement of the contralateral wing

and head, as seen in saccades? To investigate this, we examined the responses of the

contralateral wing and head immediately following an ipsilateral hitch (event-triggered averages;

figure 2*e-h)*. We observed that under every condition, each ipsilateral wing hitch (figure *2e,*

supplemental figure 2*a*) was accompanied by very small changes in the amplitude of the

contralateral wing (figure 2*f,* supplemental figure 2*b*) and no meaningful change in head yaw

(figure 2*g,* supplemental figure 2*c*). The small changes in contralateral stroke amplitude very

rarely met identification criteria for a wing hitch (supplemental figure 2*d-e*). While all treatment

groups did exhibit higher rates of contralateral hitches than untreated controls, these hitches were

not coordinated with the ipsilateral hitches. Accordingly, the rate of head yaw deviations of

equivalent magnitude to a wing hitch was virtually zero under all treatment conditions

(supplemental figure 2*e*). Wing and head kinematics during wing hitches therefore did not

resemble the coordinated syndirectional movement of the head and wings seen in saccades,

emphasizing that these two behaviors are categorically different.[39]

Haltere kinematic parameters influence hitch initiation

If haltere manipulations directly drive hitches, the rate should increase with increasing

stimulus rate. However, wing hitch rate was uniform across stimulation frequencies, and

indistinguishable from that seen with bilateral haltere ablation. Despite this, ETAs of the

ipsilateral wing amplitude (figure 2*e*) and wing instantaneous frequency (figure 2*h*) oscillated in

time with the driving stimulus. As well, fast Fourier transforms (FFTs) of wing and head

kinematic time series (supplemental figure 1) showed elevated power at the magnetic stimulation

frequency (or its baseband alias, for frequencies which exceeded the Nyquist limit of the

kinematics camera). Similarly, stimulus-triggered cyclic averages aligned on each magnetic

stimulation pulse showed that hitch probability varied in time with the stimulus for both

amplitude-increasing and amplitude-decreasing wing hitches (figure 3*a*). Taken together, these

data suggest that hitches occurred at specific times relative to the driving stimulus but were not

elicited upon every stroke cycle.

Cyclic averages also illustrated that whereas the treated haltere's frequency and

amplitude were reliably commanded with the electromagnets, the oscillation assumed a diversity

of phase relationships with the stimulus across animals and between trials (supplemental figure

3*b-d*). This technical limitation implies that stimulus timing could not necessarily predict specific

haltere positions or phases: these had to be measured directly from high-speed video.

After observing phase-locked wing hitches, we examined whether specific haltere

kinematic parameters drove hitch initiation. We assessed the haltere's angular position, angular

velocity, and phase relationship to the wingstroke cycle (figure 3*a*). We found that our

manipulations resulted in values both within and outside of the range of values observed in

untreated animals, allowing us to observe responses to both naturally-occurring stimuli and

stimuli that do not occur in flight.

We then created histograms assessing the probability of hitch initiation as a function of

each kinematic parameter under each condition, performing separate analyses for amplitude-

increasing (figure 3*b-d*) and amplitude-decreasing hitches (figure 3*e-g*). Amplitude-increasing

hitch probability was not related to haltere angular position under any condition (figure 3*b*),

however amplitude-decreasing hitches were comparatively more likely for the dynamically-

stimulated conditions when the haltere was in a downstroke position that exceeded its naturalistic

stroke limits (figure 3*e,* right). In untreated animals, the haltere's angular velocity was almost

always high, whether the haltere was moving up or down (figure 3*c*, gray bars); in haltere-

immobilized animals, the angular velocity was always low. Hitch probability was also low in

these static conditions, although higher in immobilized animals. When we dynamically perturbed

the haltere in a velocity range either faster or slower than its natural activity, however, we saw a

dramatic increase in amplitude-increasing hitch probability (figure 3*c*, right). For both

amplitude-increasing and amplitude-decreasing hitches, there was no relationship between wing

phase and hitch probability under any condition (figure 3*d,g*). Multivariate probability heatmaps

of multiple kinematic parameters did not reveal any trends over and above those outlined in

univariate analysis (supplemental figure 4). Our data therefore show that flies respond to

unnatural haltere velocities by initiating ipsilateral wing hitches, with little discernable influence

from other measured kinematic parameters.

Haltere-evoked hitches are not influenced by concurrent visual stimulation

The fly thoracic nervous system integrates inertial sensory information from the halteres

with information from the fly's visual system [25,47,48]. Tethered flies adjust gaze and

wingstroke amplitude in response to visual movement [26,49] using a motor pool shared with

reflexes driven by the halteres [41,50]. Could visual input influence wing hitches evoked by

haltere stimulation?

Concurrently with each haltere treatment, we presented a wide-field pattern of high-

contrast vertical stripes with a pattern velocity of 90°/s (figure 4*a-b*, supplemental video 3). We

then subdivided each stimulus epoch into periods of clockwise and anticlockwise visual motion

(figure 4*a-b*) and calculated hitch rate during each period (as in figure 2). In both visual stimulus

directions and for both amplitude-increasing and amplitude-decreasing hitches, visual

stimulation did not affect hitch rate under most haltere stimulus conditions (figure 4*c-d*).

Notably, however, concurrent visual stimulation in either direction reduced the rate of amplitude-

increasing ipsilateral hitches when the haltere was immobilized (figure 4*c*). For amplitude-

decreasing hitches, this was observed only for stimulation in the anticlockwise direction (figure

4*d*). Similar results were observed for the contralateral wing (supplemental figure 5). Our results

suggest that descending visual input cannot "rescue" the wing from the effects of asynchronous

haltere input.

Asynchronous haltere movement disrupts wing and head optomotor responses

Halteres adjust the gain of wing and gaze optomotor equilibrium reflexes

[20,21,27,41,51]. Is asynchronous input sufficient to drive these? We estimated the optomotor

response amplitude (following [51]) by measuring the magnitude of the FFT of each kinematic

time series at 1Hz, the frequency of the visual stimulus (figure 5*a-c*). For the ipsilateral wing and

head (figure 5*d,f*), all haltere manipulations resulted in smaller visual response amplitudes than

those of untreated flies. For the contralateral wing, only the bilateral ablation was statistically

distinct (figure 5*e*).

Static haltere manipulations (bilateral ablation and immobilization) impair the ability of

flies to adjust their head responses to widefield visual stimuli, locking the fly into a fixed low-

amplitude response to all salient pattern velocities [27]. We presented a subset of flies with a

series of visual stimuli that moved at a range of pattern velocities, 30-240°/s. Untreated flies

exhibited an optomotor tuning curve that peaked at 180°/s for wing responses (figure 5*g-h,*

supplemental figure 6*a-b*) and 120°/s for head responses (figure 5*i,* supplemental figure 6*c*), in

agreement with previous studies [27,49]. Similar tuning curves were observed for flies with

immobilized halteres (figure 5*g-i*) although with significantly lower response amplitudes than

untreated flies (supplemental figure 6*d-f*). Under dynamic stimulation (magnets engaged at

36Hz), optomotor tuning was abolished for both contralateral and ipsilateral wings. Visual

pattern velocity was not a significant predictor of response amplitude for these flies (figure 5*g-*

h). In comparison, head responses showed lower gain relative to untreated controls, at levels

comparable to those seen with haltere immobilization (figure 5*i,* supplemental figure 6*c,f*).

Discussion

For locomotion, coordinated appendages are so important that it is often difficult to force

uncoordinated movement: in flying flies, passive thoracic biomechanics restrict halteres and

wings to oscillate in antiphase [16]. We used stimuli both within and outside the bounds of

natural haltere motions to demonstrate that wing steering reflexes are critically impaired by

asynchronous haltere input, with increased ipsilateral wing hitches following "unnatural" haltere

movements (figure 2,3). In doing so, we showed that asynchronous haltere input is uniquely

deleterious to ipsilateral wing control.

Though we hypothesized that any haltere input would be sufficient for full-amplitude

visually guided head movements, we instead observed that asynchronous haltere input results in

lower amplitude movements (figure 5). The head's movements, though driven by a more diverse

set of inputs and occurring at lower speeds, still show deleterious effects of asynchronous haltere

input. Asynchronous haltere input did not evoke head yaw changes comparable to those seen in

the wing (figure 2), congruent with previous findings that abrupt changes in head yaw (as part of

a saccade) are primarily evoked by proprioceptive inputs [39]. We show that dynamic haltere

perturbations drive "wing hitches", short-duration changes in amplitude that can occur

unilaterally. These hitches may be a small component of the body saccade, a coordinated

movement of the head and both wings which is used to reorient the flying fly.

Asynchronous haltere input increases wing hitches

295   Asynchronous dynamic haltere input increases the rate of ipsilateral wing hitches (figure

2), particularly when the haltere moves at angular velocities outside of its natural operating

range. The halteres are sensitive to the body's angular velocity [19]; here, we show that they are

sensitive to their own angular velocity as well (figure 3). Our experiment with one immobilized

haltere shows that hitches are not driven by specific motions of the haltere: loss of one or both

halteres also results in an elevated rate of hitches of similar amplitude. Therefore, another

possible interpretation is that hitches emerge spontaneously from activity of the wing motor

system, and that synchronous haltere input acts to suppress them.

303   [23][51]Campaniform sensilla are sensitive to compressive and shearing forces [8,52,53],

which arise from the haltere's movement and scale with the bulb's angular velocity [54]. We

might therefore predict that hitches are more or less likely to be initiated as a function of the

haltere's angular position. Our results suggest that amplitude-decreasing hitches are more likely

when the haltere is hyperextended past its natural downstroke position, we did not otherwise

observe a strong correspondence between haltere angular position and hitch initiation (figure 3).

This suggests, but does not conclusively show, that the halteres suppress spontaneous hitches

rather than driving hitches at specific positions. However, the extreme sensitivity of the sensilla

[14,55] requires us to consider the possibility that we were not able to stimulate accurately

enough to reach a conclusion. The overall effect on flight behavior is the same under both

hypotheses: the halteres have a damping effect on wing steering and thus on body rotations,

reflecting prior experimental and theoretical work [4,56,57].

Coordinated movement and haltere influence on the contralateral wing

In previous work [28], manipulating haltere stroke amplitude without compromising

haltere-wing synchrony evoked a smooth wing-steering response. Here, responses to haltere

perturbation were less stereotyped, but in neither the present nor previous work was there a clear

relationship between instantaneous wing phase and haltere-evoked changes in wing kinematic

parameters (figure 3). This was surprising given the organization of the wing-steering motor

system, a major component of which are tonic muscles which fire once every stroke cycle and

control muscle tension by varying firing phase within it [30,58–60]. While our stimulation

sampled the entire range of wing/haltere phases, a limitation of our experimental paradigm is that

we could not stimulate with a particular phase combination for consecutive stroke cycles, which

may be important for the force production properties of these muscles. Another hypothesis is that

the untreated contralateral haltere provides enough input to supersede an absent or

malfunctioning ipsilateral haltere.

Our findings that asynchronous haltere input increases the rate of contralateral wing

hitches independent of movements accompanying ipsilateral hitches (figure 2f, supplemental

figure 2b,*d-g*) are consistent with previous findings that each wing receives information from

both halteres, but that this information can be used to drive the wings independently. Flies are

capable of apparently normal flight following unilateral haltere ablation [61]. In *Calliphora,*

input from a single haltere is sufficient to entrain firing of the contralateral first basalare muscle

following ablation of the ipsilateral haltere and both wing nerves [17]. Experiments rotating the

fly's entire body showed that both amplitude and frequency components of haltere-wing reflexes

are impaired following bilateral—but not unilateral—haltere ablation [19]. Similarly,

manipulating the bulb mass of a single haltere produced comparable changes in both clockwise

and counterclockwise body saccades [42].

This bilateral integration implicates the contralateral haltere interneurons, or cHINs

[24,62], which receive input from haltere afferents and represent the only known anatomical

pathway through which haltere information crosses the body axis. Though conserved across fly

taxa, these neurons have eluded physiological characterization, leaving questions as to the

sensory representations and transformations they mediate *en route* to their motor targets. One

possibility is that the predominately unilateral hitches observed in our data might represent

"building blocks" that the thoracic nervous system can assemble into more complicated

maneuvers such as saccades that coordinate tightly across the body axis.

Haltere contribution to head and wing optomotor responses

As descending visual inputs concurrently target wing and gaze motoneurons [24,25,63]

we also examined multisensory integration of haltere and visual inputs. We show that haltere-

wing asynchrony critically impairs the wing optomotor response to a degree comparable with

bilateral haltere ablation. While this impairment may be unsurprising for wings, it is less obvious

how the head might respond to asynchronous haltere input. Bilateral haltere ablation prevents

flies from adjusting the gain of the head optomotor response to different pattern velocities [27].

Here, we show that loss of one haltere (by immobilization) or asynchronous input from one

haltere reduces the optomotor response gain, but does not abolish pattern velocity tuning (figure

5).

In some neck motoneurons of quiescent *Calliphora*, an absence of haltere input prevents

spiking responses to visual input, but spiking can be restored with haltere movement at any

frequency, even as low as 10 Hz [29]. If the halteres are necessary only to provide excitatory

drive, then asynchronous input would produce gaze optomotor tuning curves with similar gain to

a control. However, we found that asynchronous drive resulted in lower gain.

There are two main hypotheses that could explain why neck motoneurons that can be

excited by any haltere input in quiescence fail to produce high-gain visually-guided head

movements when stimulated asynchronously in tethered flight. First, wing sensory inputs to neck

motoneurons are still unknown, raising the possibility that full neck activity may require both

wing and haltere input, like the wing motoneuron mnb1 [22]. However, flies with ablated wings

execute high-gain optomotor responses in tethered "flight" (Mureli et al., 2017).

Second, the "control loop," in which descending neurons provide excitatory input to

haltere muscles, [18,63] may explain this result. Our experiment breaks the control loop by

nullifying the relationship between haltere muscle activity and haltere sensory input: visual input

to the haltere muscles no longer produces haltere input that can be used to initiate proper

reflexes. Both hypotheses could explain our results that visual input cannot "rescue" the wing

from high hitch rates (figure 4), that visual tuning is ablated in the ipsilateral wing (figure 5g),

and that visual tuning in the head remains intact but at lower gain (figure 5). Future recordings of

electrophysiological properties of neck motoneurons under different behavioral conditions and of

haltere movement responses to visual inputs could help untangle these possibilities.

Acknowledgments

We would like to thank Chenxin Bi and Brandon Chun for help with data collection and

annotation, as well as Noah DeFino, Jesse Fritz and Lauren Metz for fly care. We would also like

to thank Hillel Chiel, Bradley Dickerson, Jeremiah Didion, Alexandra Gurgis, Nicholas

Kathman, Kristianna Lea, Mark Willis, Gabriella Wolff, and Alexandra Yarger for helpful

commentary.

Competing interests

No competing interests declared.

Funding

This work was supported by United States Air Force Office of Scientific Research

[FA9550-14-0398 and FA9550-16-1-0165 to JLF]; and National Science Foundation [1754412

to JLF].

Data Availability

All processed data is available in the associated Dryad repository. Raw video data

available upon request. Kinematic time series for some untreated control animals in Figures 1-5

are included with a previously published publicly available dataset [64].

References Cited

1. Heckscher ES *et al.* 2015 Even-skipped+ interneurons are core components of a sensorimotor circuit that maintains left-right symmetric muscle contraction amplitude. *Neuron* **88**, 314–329.

2. Wiggin TD, Peck JH, Masino MA. 2014 Coordination of fictive motor activity in the larval zebrafish is generated by non-segmental mechanisms. *PLoS One* **9**, e109117.

3. Wannier T, Bastiaanse C, Colombo G, Dietz V. 2001 Arm to leg coordination in humans during walking, creeping and swimming activities. *Experimental Brain Research 2001 141:3* **141**, 375–379.

4. Dickinson MH, Muijres FT. 2016 The aerodynamics and control of free flight maneuvers in Drosophila. *Philosophical transactions of the Royal Society of London B: Biological Sciences* **371**, 20150388.

5. Hiebert GW, Pearson KG. 1999 Contribution of sensory feedback to the generation of extensor activity during walking in the decerebrate Cat. *J Neurophysiol* **81**, 758–70.

6. Noah JA, Quimby L, Frazier SF, Zill SN. 2004 Walking on a 'peg leg': Extensor muscle activities and sensory feedback after distal leg denervation in cockroaches. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **190**, 217–231.

7. Dickinson MH. 1990 Comparison of encoding properties of campaniform sensilla on the fly wing. *Journal of Experimental Biology* **261**, 245–261.

8. Pringle JWS. 1948 The gyroscopic mechanism of the halteres of Diptera. *Philosophical Transactions of the Royal Society B: Biological Sciences* **233**, 347–384.

9. Sun M, Wang J, Xiong Y, Sun M, Wang J, Xiong Y. 2007 Dynamic flight stability of hovering insects. *Acta Mech Sin* **23**, 231–246.

10. Derham W. 1714 *Physico-theology: or, a demonstration of the being and attributes of God, from His works of creation. Being the substance of XVI sermons preached in St. Mary Le Bow-church, London, at the honble Mr. Boyle's lectures, in the years 1711 and 1712.* W. and J. Innys, London, 1714.

11. Gnatzy W, Grünert U, Bender M. 1987 Campaniform sensilla of Calliphora vicina (Insecta, Diptera). *Zoomorphology* **106**, 312–319.

12. Agrawal S, Grimaldi D, Fox JL. 2017 Haltere morphology and campaniform sensilla arrangement across Diptera. *Arthropod Struct Dev* **46**, 215–229.

13. Fox JL, Daniel TL. 2008 A neural basis for gyroscopic force measurement in the halteres of Holorusia. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **194**, 887–897.

14. Yarger AM, Fox JL. 2018 Single mechanosensory neurons encode lateral displacements using precise spike timing and thresholds. *Proceedings of the Royal Society B: Biological Sciences* **285**, 20181759.

15. Hall JM, McLoughlin DP, Kathman ND, Yarger AM, Mureli S, Fox JL. 2015 Kinematic diversity
suggests expanded roles for fly halteres. *Biol Lett* **11**, 20150845.

16. Deora T, Singh AK, Sane SP. 2015 Biomechanical basis of wing and haltere coordination in flies.
Proc Natl Acad Sci U S A **112**, 1481–6.

17. Heide G. 1983 Neural mechanisms of flight control in Diptera. In *Biona-report 2* (ed G Fischer),
pp. 35–52. Stuttgart, New York.

18. Dickerson BH, de Souza AM, Huda A, Dickinson MH. 2019 Flies regulate wing motion via active
control of a dual-function gyroscope. *Current Biology* **29**, 3517–3524.

19. Dickinson MH. 1999 Haltere-mediated equilibrium reflexes of the fruit fly, Drosophila
melanogaster. *Philosophical Transactions of the Royal Society B: Biological Sciences* **354**, 903–
916.

20. Mureli S, Fox JL. 2015 Haltere mechanosensory influence on tethered flight behavior in
Drosophila. *Journal of Experimental Biology* , 2528–2537.

21. Bartussek J, Lehmann F-O. 2016 Proprioceptive feedback determines visuomotor gain in
Drosophila. *R Soc Open Sci* **3**, 150562.

22. Fayyazuddin A, Dickinson MH. 1996 Haltere afferents provide direct, electrotonic input to a
steering motor neuron in the blowfly, Calliphora. *J Neurosci* **16**, 5225–5232.

23. Fayyazuddin a, Dickinson MH. 1999 Convergent mechanosensory input structures the firing
phase of a steering motor neuron in the blowfly, Calliphora. *J Neurophysiol* **82**, 1916–1926.

24. Strausfeld NJ, Seyan HS. 1985 Convergence of visual, haltere, and prosternal inputs at neck motor
neurons of Calliphora erythrocephala. *Cell Tissue Res* **240**, 601–615.

25. Namiki S, Dickinson MH, Wong AM, Korff W, Card GM. 2018 The functional organization of
descending sensory-motor pathways in Drosophila. *Elife* **7**, e34272.

26. Hengstenberg R, Sandeman DC, Hengstenberg B. 1986 Compensatory head roll in the blowfly
Calliphora during flight. *Proceedings of the Royal Society B: Biological Sciences* **227**, 455–482.

27. Mureli S, Thanigaivelan I, Schaffer ML, Fox JL. 2017 Cross-modal influence of mechanosensory
input on gaze responses to visual motion in Drosophila. *Journal of Experimental Biology* **220**,
2218–2227.

28. Rauscher MJ, Fox JL. 2021 Haltere and visual sensory inputs sum linearly to predict wing (but not
head) motor output in tethered flying Drosophila. *Proceedings of the Royal Society B* **288**,
20202374.

29. Huston SJ, Krapp HG. 2009 Nonlinear integration of visual and haltere inputs in fly neck motor
neurons. *Journal of Neuroscience* **29**, 13097–13105.

30. Heide G, Götz KG. 1996 Optomotor control of course and altitude in Drosophila melanogaster is
correlated with distinct activities of at least three pairs of flight steering muscles. *Journal of*
Experimental Biology **199**, 1711–1726.

31. Götz KG. 1979 The optomotor equilibrium of the Drosophila navigation system. *Journal of*
Comparative Physiology A **99**, 187–210.

32. Land MF, Collett TS. 1974 Chasing behaviour of houseflies (Fannia canicularis). *Journal of*
comparative physiology 1974 89:4 **89**, 331–357.

33. Land MF. 1973 Head movement of flies during visually guided flight. *Nature* **243**, 299–300.

34. Bender J a, Dickinson MH. 2006 Visual stimulation of saccades in magnetically tethered
Drosophila. *J Exp Biol* **209**, 3170–3182.

35. Mongeau JM, Frye MA. 2017 Drosophila Spatiotemporally Integrates Visual Signals to Control
Saccades. *Current Biology* **27**, 2901-2914.e2.

36. Kim AJ, Fitzgerald JK, Maimon G. 2015 Cellular evidence for efference copy in Drosophila
visuomotor processing. *Nat Neurosci* **18**, 1247–1255.

37. Kim AJ, Fenk LM, Lyu C, Maimon G, Otsuna H, Ito K, Borst A, Reiff DF. 2017 Quantitative
predictions orchestrate visual signaling in Drosophila. *Current Biology* **168**, 280–294.

38. Schilstra C, van Hateren JH. 1998 Stabilizing gaze in flying blowflies. *Nature* **395**, 654.

39. Cellini B, Salem W, Mongeau JM. 2021 Mechanisms of punctuated vision in fly flight. *Current*
Biology **31**, 4009-4024.e3.

40. Cellini B, Mongeau JM. 2020 Active vision shapes and coordinates flight motor responses in flies.
Proc Natl Acad Sci U S A **117**, 23085–23095.

41. Cellini B, Mongeau JM. 2022 Nested mechanosensory feedback actively damps visually guided
head movements in Drosophila. *Elife* **11**.

42. Bender JA, Dickinson MH. 2006 A comparison of visual and haltere-mediated feedback in the
control of body saccades in Drosophila melanogaster. *Journal of Experimental Biology* **209**, 4597–
606.

43. Reiser MB, Dickinson MH. 2008 A modular display system for insect behavioral neuroscience. *J*
Neurosci Methods **167**, 127–139.

44. Fox J, Aptekar J, Zolotova N, Shoemaker P, Frye M. 2014 Figure-ground discrimination behavior in
Drosophila. I. Spatial organization of wing-steering responses. *J Exp Biol* **217**, 558–69.

45. Hedrick TL. 2008 Software techniques for two- and three-dimensional kinematic measurements
of biological and biomimetic systems. *Bioinspir Biomim* **3**, 34001.

46. Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018 DeepLabCut:
markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* **21**,
1281–1289.

47. Suver MP, Huda A, Iwasaki N, Safarik S, Dickinson MH. 2016 An array of descending visual
interneurons encoding self-motion in Drosophila. *The Journal of Neuroscience* **36**, 11768–11780.

48.   Namiki S, Ros IG, Morrow C, Rowell WJ, Card GM, Korff W, Dickinson MH. 2022 A population of
descending neurons that regulates the flight motor of Drosophila. *Current Biology* **32**, 1–8.

49.   Duistermars BJ, Reiser MB, Zhu Y, Frye MA. 2007 Dynamic properties of large-field and small-field
optomotor flight responses in Drosophila. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol*
193, 787–799.

50.   Schwyn DA, Heras FJH, Bolliger G, Parsons MM, Krapp HG, Tanaka RJ. 2011 Interplay between
feedback and feedforward control in fly gaze stabilization. *IFAC Proceedings Volumes (IFAC-*
PapersOnline) **18**, 9674–9679.

51.   Rimniceanu M, Currea JP, Frye MA. 2023 Proprioception gates visual object fixation in flying flies.
Current Biology **33**, 1459-1471.e3.

52.   Zill SN, Moran DT. 1981 The exoskeleton and insect proprioception. I. Responses of tibial
campaniform sensilla to external and muscle-generated forces in the American cockroach,
Periplaneta americana. *Journal of Experimental Biology* **91**, 1–24.

53.   Chapman KM, Smith RS. 1963 A linear transfer function underlying impulse frequency
modulation in a cockroach mechanoreceptor. *Nature 1963 197:4868* **197**, 699–700.

54.   Nalbach G. 1993 The halteres of the blowfly Calliphora. *Journal of Comparative Physiology A* **173**,
293–300.

55.   Haag J, Wertz A, Borst A. 2010 Central gating of fly optomotor response. *Pnas* **107**, 20104–20109.

56.   Elzinga MJ, Dickson WB, Dickinson MH. 2012 The influence of sensory delay on the yaw dynamics
of a flapping insect. *J R Soc Interface* **9**, 1685–1696.

57.   Sherman A, Dickinson MH. 2004 Summation of visual and mechanosensory feedback in
Drosophila flight control. *Journal of Experimental Biology* **207**, 133–142.

58.   Lindsay T, Sustar A, Dickinson MH. 2017 The function and organization of the motor system
controlling flight maneuvers in flies. *Current Biology* **27**, 345–358.

59.   Tu MS, Dickinson MH. 1996 The control of wing kinematics by two steering muscles of the
blowfly (Calliphora vicina). *Journal of Comparative Physiology A 1996 178:6* **178**, 813–830.

60.   Balint CN, Dickinson MH. 2001 The correlation between wing kinematics and steering muscle
activity in the blowfly Calliphora vicina. *J Exp Biol* **204**, 4213–4226.

61.   Fraenkel G. 1939 The function of the halteres of flies (Diptera). *Proceedings of the Zoological*
Society of London **A109**, 69–78.

62.   Trimarchi JR, Murphey RK. 1997 The shaking-B2 mutation disrupts electrical synapses in a flight
circuit in adult Drosophila. *Journal of Neuroscience* **17**, 4700–4710.

63.   Chan WP, Prete F, Dickinson MH. 1998 Visual input to the efferent control system of a fly's
'gyroscope'. *Science* **280**, 289–292.

64.   Rauscher MJ, Fox JL. 2020 Weighted haltere and imposed haltere stroke reduction tethered flying
540   Drosophila kinematics. *Dryad, Dataset*
541   (doi:https://datadryad.org/stash/dataset/doi:10.5061/dryad.g4f4qrfnr)

Figure 1. Electromagnetic control of haltere movement asynchronous from the wings.
(a) Schematic of iron filing treatment (top, inset) and electromagnetic stimulation system (bottom). *(b)* Representative traces showing movement of untreated haltere at wingbeat frequency (~200Hz), immobilized haltere treated with iron filing (ii), and actively-driven haltere at a range of frequencies (iii-viii) controlled by electromagnets *(c)* Haltere tracking by condition, registered into a normalized body coordinate system defining the haltere base as the origin, represented by circle and the anterior wing base as (0,1), represented by asterisk. Colors reflect the Z-normalized wingstroke elevation at each measurement point. *(d)* Mean FFT (95% confidence interval shaded) of all high-speed (1000 or 2000fps) time series with the haltere immobilized (black), driven at 25Hz (red), and 50Hz (pink). *(e)* same as *(d)* for 36Hz (blue) and 72Hz (teal) frequencies. *(f)* same as *(d)* and *(e)* for 100Hz (orange) and 167Hz (magenta) frequencies. *(g-i)* Haltere stroke amplitude *(g)*, peak angular velocity *(h)* and baseline wingbeat frequencies *(i)* observed under each haltere manipulation. Lines show median for each group. 167Hz responses from outlier animal shown with +. Letters denote statistical groupings from multiple Bonferroni-corrected Wilcoxon rank-sum test (5% alpha).

Figure 2. Haltere manipulation increases ipsilateral wing hitch rate

(a) Schematic of measured kinematic outputs, recorded at 100fps, showing downstroke amplitude (angle of wing envelope leading edge) for each wing and yaw movement of the head. Arcs denote the sign (positive or negative) of clockwise and anticlockwise rotations for each parameter *(b)* Example ipsilateral wing amplitude traces showing fast wing hitches from untreated animal *(i)*, similar in amplitude but distinct in temporal dynamics from a wing saccade *(ii)*, and increased hitch rate observed in an animal with an asynchronously-driven haltere *(iii)*. Hitches are annotated with filled circles. Saccade with gray triangle *(c)* Rate of all ipsilateral wing hitches of increasing amplitude across all haltere treatments (see Methods). Left and right wing responses for symmetrical manipulations (untreated, bilateral ablation) are pooled. Lines show median for each group. Letters denote statistical groupings from multiple Bonferroni-corrected Wilcoxon rank-sum test (5% alpha). *(d)* Same as *(c)* but for amplitude-decreasing hitches *(e)* Event-triggered average (ETA) responses for each condition, generated from time-aligned, baseline-subtracted kinematic timeseries. Gray triangles show the hitch peak, around which each constitutive video segment of the ETA is indexed, with amplitude-increasing hitch ETAs shown with solid lines and amplitude-decreasing hitch ETAs shown with dashed lines (Shade, 95% confidence interval). Gray traces show controls triggered from random indices within the same time series. *(f-h)* Same as *(e)* but for concurrent movements of the contralateral wing *(f)*, head yaw *(g)* and baseline-subtracted instantaneous wingbeat frequency *(h)*.

Figure 3. Ipsilateral wing saccades are more likely when the haltere exceeds normal kinematic parameters
(a) Summary of haltere angular position, angular velocity, and wing instantaneous phase from all high speed data, separated by condition. *(ai)* Untreated animals, showing natural antiphase synchrony of haltere and wings. Inset shows unified body coordinate scheme, with haltere upstrokes typically occupying the 45-90° range and downstrokes typically occupying the 235-270° range *(aii)* Frames from haltere immobilized (magnets OFF) condition. Green dashed line shows extent of the naturalistic parameter space observed in the untreated animal. *(aiii)* High-speed frames aggregated from all magnetic stimulation frequencies, showing broad overlap with the naturalistic range of haltere angular velocities and angular positions. *(b)* Histograms of ipsilateral wing amplitude video frames showing probability of amplitude-increasing hitch as a function of haltere angular position at frame onset for each condition. Light green shaded region outlines the natural range of haltere angular positions. *(c)* Same as *(b)* but for haltere signed-RMS angular velocity throughout the frame (see Methods). Light green shaded regions outline the naturalistic range of negative (upstroke) and positive (downstroke) angular velocities. *(d)*. Same as *(b)* but for wing instantaneous phase at frame onset. *(e-g)* Same as *(b-d)* but for amplitude-decreasing hitches.

Figure 4. Concurrent visual input does not influence wing hitch rate during most haltere manipulations
(a) Schematic of visual stimulus, showing irregular stripe pattern moving on an LED panel under the control of a 1Hz triangle-wave with a pattern velocity of 90°/s *(b)* Example ipsilateral wing amplitude responses to one cycle of the visual stimulus under several haltere treatment conditions. *(c)* Rate of increasing-amplitude ipsilateral wing hitches across visual conditions, showing no change between uniform visual stimulus (reproduced from figure 2), clockwise visual stimulus epochs, and anticlockwise epochs in most haltere treatment conditions. Mean value for each visual stimulus condition represented by dash, upward pointing triangle, and downward pointing triangle respectively. *(d)* Same as a *(c)* but for amplitude-decreasing ipsilateral wing hitch rate (** = significance at 1% alpha level, *** = significance at 0.1% alpha level, multiple Bonferroni-corrected Wilcoxon rank-sum test).

Figure 5. Haltere manipulations influence visually-guided wing and head movements.
(a) Schematic of visual stimulus, as in figure 4A. *(b)* Example ipsilateral wing amplitude responses to multiple cycles of the visual stimulus under several haltere treatment conditions. *(c)* FFTs of traces from *(b)*, showing peak at the frequency of the visual stimulus. *(d)* Ipsilateral wing response amplitude at visual stimulus frequency for all tested haltere treatments. For symmetrical manipulations (untreated, bilateral ablation), left and right wing responses are pooled. Letters denote statistical groupings from multiple Bonferroni-corrected Wilcoxon rank-sum test (5% alpha). *(e,f)* Same as *(d)* for contralateral wing and head yaw responses, respectively. Lines show median for each group. *(g)* Ipsilateral wing visual response magnitude, as estimated by measuring FFT magnitude at visual stimulus frequency, for untreated flies, haltere-immobilized flies, and 36Hz magnetic stimulation flies. Trendlines connect median data from each pattern velocity, with shaded region showing 1st and 3rd quartiles. Inset, results of Kruskal-Wallis omnibus test indicate whether pattern velocity is a significant predictor of response amplitude within each treatment group at the 5% alpha level (*) or not (ns). *(h-i)* Same as *(g)*, but for contralateral wing and head yaw amplitude, respectively.

(a) electromagnet

IR cameras

tethered fly

electromagnet

iron filing glued to haltere

(b)
(i) untreated Haltere
(normalized wingbeat)
(ii) haltere Immobilized
100°| 10 ms
(iii) magnets 25Hz
(iv) magnets 36Hz
(v) magnets 50Hz
(vi) magnets 72Hz
(vii) magnets 100Hz
(viii) magnets 167Hz

(c)

normalized wing elevation (Z-Score)

| untreated | halt imm | mag 25Hz | mag 36Hz | mag 50Hz | mag 72Hz | mag 100Hz | mag 167Hz |

(0,1)
(0,0)

n = 135377 frames
n = 312241
n = 141798
n = 96700
n = 136774
n = 139187
n = 57311
n = 98478

(d)

haltere stroke FFT magnitude (°)

halt imm n = 186
mag 25Hz n = 52
mag 50Hz n = 41

frequency (Hz)

(e)

halt imm n = 186
mag 36Hz n = 44
mag 72Hz n = 38

(f)

halt imm n = 186
mag 100Hz n = 16
mag 167Hz n = 26

(g)

haltere stroke amplitude (°)

a n=38
a n=34
b n=186
c n=52
c n=44
c n=41
c n=38
c n=16
d n=22

untreated   contra haltere   halt imm   mag 25Hz   mag 36Hz   mag 50Hz   mag 72Hz   mag 100Hz   mag 167Hz

(h)

haltere peak angular velocity (°/ms)

a b n=38
a b n=34
c n=186
a d e n=52
a d e f n=44
a d e f n=41
a d e f n=38
a d e f n=16
a d f n=22

untreated   contra haltere   halt imm   mag 25Hz   mag 36Hz   mag 50Hz   mag 72Hz   mag 100Hz   mag 167Hz

(i)

wingbeat frequency (Hz)

a b n=120
a n=200
a b n=57
a b n=46
a b n=46
a b n=45
a b n=16
a b n=27

untreated   halt imm   mag 25Hz   mag 36Hz   mag 50Hz   mag 72Hz   mag 100Hz   mag 167Hz

(a)

+ head yaw +
contra. wing   ipsi. wing
untreated haltere   treated haltere

(b)

i. hitches   ii. saccade   iii. magnets 25Hz

10° 250ms

(c) amplitude-increasing hitches

ipsi wing hitch rate (Hz)

a b b c c c c c c
n = 242 284 33 56 57 33 52 49 19

untreated, halt imm., single abl, both abl, mag 25Hz, mag 36Hz, mag 50Hz, mag 72Hz, mag 100Hz

(d) amplitude-decreasing hitches

a b b c c c c c c
n = 242 284 33 56 57 33 52 49 19

untreated, halt imm, single abl, both abl, mag 25Hz, mag 36Hz, mag 50Hz, mag 72Hz, mag 100Hz

	untreated flies	haltere imm	single abl	both abl	mag 25Hz	mag 36Hz	mag 50Hz	mag 72Hz	mag 100Hz

(e) ipsi wing amplitude (°)

(f) contra wing amplitude (°)   3°

(g) head yaw (°)

(h) wingbeat frequency (Hz)   1Hz

$n_{(+)}$ = amplitude-increase
$n_{(-)}$ = amplitude-decrease

$n_{(+)}$ = 60
$n_{(-)}$ = 91

$n_{(+)}$ = 1081
$n_{(-)}$ = 575

$n_{(+)}$ = 972
$n_{(-)}$ = 850

$n_{(+)}$ = 3081
$n_{(-)}$ = 2137

$n_{(+)}$ = 797
$n_{(-)}$ = 640

$n_{(+)}$ = 675
$n_{(-)}$ = 518

$n_{(+)}$ = 760
$n_{(-)}$ = 566

$n_{(+)}$ = 662
$n_{(-)}$ = 530

$n_{(+)}$ = 280
$n_{(-)}$ = 220

(a) 1Hz visual stimulus

(b) anticlockwise visual stimulus epoch

clockwise visual stimulus epoch

1Hz visual stimulus position (not to scale)

untreated halt Imm. mag. 50Hz

ipsi. wing amplitude (°)

time (s)

(c) amplitude-increasing hitches

(d) amplitude-decreasing hitches

ipsi. wing hitch rate (Hz)

n =	n =	n =	n =	n =	n =	n =
242	284	56	57	33	52	49
68	97	36	25	18	24	22
68	97	36	25	18	24	22

— No Visual Stimulus
△ Anticlockwise
▽ Clockwise

***

untreated halt imm. both ABL mag. 25Hz mag. 36Hz mag. 50Hz mag. 72Hz