**Asynchronous haltere input drives specific wing and head movements in *Drosophila***

**Running title: Asynchronous haltere input**

Michael J Rauscher (mjr67@case.edu) and Jessica L Fox (jlf88@case.edu)

Case Western Reserve University Dept. Biology

**Keywords:**

Animal Behavior, Drosophila, Haltere, Insect Flight, Saccade, Multisensory Integration

**Abstract**

Halteres are multifunctional mechanosensory organs unique to the true flies (Diptera). A set of reduced hindwings, the halteres beat at the same frequency as the lift-generating forewings and sense inertial forces via mechanosensory campaniform sensilla. Though it is well-established that haltere ablation makes stable flight impossible, the specific role of wing-synchronous input has not been established. Using small iron filings attached to the halteres of tethered flies and an alternating electromagnetic field, we experimentally decoupled the wings and halteres of flying *Drosophila* and observed the resulting changes in wingbeat amplitude and head orientation. We find that asynchronous haltere input results in fast saccades in the wing but not the head. We also find that wing and gaze optomotor responses are disrupted differently by asynchronous input. These effects of wing-asynchronous haltere input show that specific sensory information is necessary for maintaining wing amplitude stability and adaptive gaze control.

**Introduction**

Animal locomotion, from crawling (Heckscher et al., 2015) to swimming (Wiggin et al., 2014) to walking (Wannier et al., 2001) to flight (Dickinson and Muijres, 2016), typically requires the coordinated motion of multiple body segments or appendages. Though this coordination can stem from the activity of a central pattern generator, sensory input from proprioceptors modulates body motions to adapt each step (Hiebert and Pearson, 1999; Noah et al., 2004). The specific contraction timing of muscles actuating multiple appendages can significantly change gait and posture, and thus the mechanisms ensuring limb coordination are essential to structuring adaptive locomotion.

In flying flies, proprioceptors on the wing (Dickinson, 1990) and haltere (Pringle, 1948)

provide crucial feedback allowing for active control of each wingstroke to counteract flies'

inherent pitch instability (Sun et al., 2007). The halteres are small club-shaped organs which

have evolved from the hindwings of the true flies (Diptera). Long known to be required for

stable flight (Derham, 1714), the halteres do not generate significant lift, but instead primarily

detect inertial forces using campaniform sensilla at their base (Agrawal et al., 2017; Gnatzy et

al., 1987; Pringle, 1948). Campaniform sensilla are a key class of insect proprioceptor, sensitive

to strain forces acting on and within the cuticle (Chapman and Smith, 1963; Zill and Moran,

1981). Afferent neurons from haltere campaniform sensilla entrain their firing to a preferred

phase of the haltere stroke cycle (Fox and Daniel, 2008; Yarger and Fox, 2018), which in

*Drosophila* and most other fly taxa is biomechanically linked to the wing stroke cycle with an

antiphase synchrony (Deora et al., 2015; Hall et al., 2015).

Continuous, stroke-synchronous feedback from the halteres is functionally important for

wing-steering motoneurons and muscles (Dickerson et al., 2019; Heide, 1983; Pringle, 1948). In

tethered flight, haltere ablation results in highly variable wing amplitudes and diminished wing

optomotor responses (Bartussek and Lehmann, 2016; Dickinson, 1999; Mureli and Fox, 2015).

The wing motoneurons are a primary target of haltere afferents, mediating a suite of

compensatory equilibrium reflexes. In the blowfly *Calliphora*, some haltere afferents project

directly to the motoneuron controlling the wing steering muscle wB1 (Fayyazuddin and

Dickinson, 1996). Direct stimulation of the haltere nerve shows that it can entrain the firing of

the wB1 motoneuron, which is normally more sensitive to reafference from the wing, if

stimulation occurs within a specified time after the wing nerve's time of last firing (Fayyazuddin

and Dickinson, 1999). In flying *Drosophila*, optogenetic manipulation of the haltere muscles

resets the firing phase of both the wB1 and wB2 muscles (Dickerson et al., 2019). What is the

impact on the wings in flight if their motoneurons receive asynchronous haltere input? We

hypothesize that the haltere's phase-locked input provides stabilizing input to the wing muscles,

and that changing the phase of the haltere's stroke relative to the wing will result in wing

amplitude instabilities.

Neck motoneurons also receive input from haltere afferents (Namiki et al., 2018;

Strausfeld and Seyan, 1985), and mediate a similar suite of gaze-stabilizing reflexes

(Hengstenberg et al., 1986). Following bilateral haltere ablation, flies are incapable of

modulating the gain of optomotor gaze responses (Mureli et al., 2017). In recent work, we found

that the head's movements are driven primarily by vision and not by the halteres (Rauscher and

Fox, 2021). In larger flies, spiking responses to visual input are gated, but not modified, by

haltere input in a subset of neck motoneurons: suprathreshold action potentials in these cells

require haltere oscillation but are invariant to changes in haltere frequency (Huston and Krapp,

2009). We hypothesize that stroke-synchronous haltere information may be dispensable during

visually-guided behaviors, with haltere input acting solely as a gating source of excitatory drive

onto neck motoneurons. Under this hypothesis, we would expect both wing-synchronous and

wing-asynchronous haltere input to be capable of modulating gaze optomotor behavior at similar

or equivalent levels.

Previously, we showed that synchronous manipulation of the haltere's amplitude resulted

in smooth changes in wingstroke amplitude and gaze direction (Rauscher and Fox, 2021).

However, periods of smooth control over these kinematic outputs in fly flight are punctuated by

abrupt changes in heading and gaze direction called saccades, analogous to the rapid stabilizing

eye movements of vertebrates. These serve to minimize motion blur on the eyes and have been

extensively characterized in free flight (Land and Collett, 1974), tethered flight with free rotation

about the yaw axis (Cellini and Mongeau, 2022; Land, 1973) and rigidly tethered flight (Cellini

and Mongeau, 2020; Cellini et al., 2021; Kim et al., 2015). Saccades can occur spontaneously or

be evoked by visual input, motivating study of the sensorimotor elements mediating saccades.

Because saccades in tethered flight are larger and appear more frequently than saccades in free

flight, their functional significance can be difficult to interpret. However, they are a distinct

output of the wing and neck motor pathways, and observing how sensory modulation influences

saccades can reveal pathways for processing sensory input. For example, Bender (2006) showed

that changes to the bulb mass of the haltere (and thus the "gain" of the haltere's strain-sensitive

campaniform sensilla) alters the dynamics of saccade termination for visually-initiated saccades.

To examine how haltere input drives smooth and saccadic wing and gaze behaviors, we

experimentally decoupled the movement of the wings and haltere. By adding a small iron mass

to one haltere, we immobilized it, abolishing it as a source of stroke-synchronous information.

Then, using electromagnets, we drove the haltere at specific frequencies, moving it throughout

its natural range of motion in a manner that was periodic and predictable but asynchronous with

the wings. This movement allowed us to observe responses to wing-haltere phase relationships

that do not naturally occur. We show that asynchronously-driven halteres increase the likelihood

of saccadic steering changes in the wing motor output as well as the broadband noise levels of

the wing's stroke amplitude. Movements of the head, by contrast, are not accompanied by an

increase in broadband noise levels. However, the optomotor responses of both head and wings

are significantly diminished by asynchronous haltere input. Our results show that synchronous

haltere input is necessary for wing amplitude (but not gaze) stabilization and adaptive optomotor

responses.

**Materials and Methods**

*Animal care, haltere manipulations, and experimental trials*

Animals were reared from a colony of wild-caught *Drosophila melanogaster*. Adult female flies (aged 3-5 days post-eclosion) were cold-anaesthetized and attached to rigid tungsten pins using UV-curing dental cement. For magnetic stimulation experiments, a single iron filing was attached to the right haltere with UV-curing dental cement prior to tethering. Only iron filings of sufficient mass to immobilize the haltere with the magnets disengaged (filings in excess of 12μg, as determined in Rauscher and Fox, 2021) were used for experiments. For ablation experiments, one or both halteres were gently and completely removed using a pair of fine forceps.

Flies were given at least ten minutes to recover from haltere manipulation and tethering prior to usage in tethered flight behavioral experiments. Haltere stimulation periods lasted 3-4 seconds, which frequently evoked grooming of the haltere with the ipsilateral metathoracic leg. Time indices for grooming epochs were annotated manually and these video segments were excluded from analysis of head and wing kinematics. For filing-treated animals, successive presentations of different haltere stimulation conditions were preceded and followed by control epochs with the magnets disengaged lasting a minimum of 250ms (~50 wingbeats). For each animal, 1-6 epochs from the same experimental condition (from replicate stimulus presentations and intervening between grooming bouts) were drawn for subsequent analysis.

*Haltere stimulation*

To facilitate control over haltere stroke kinematics, two iron-core electromagnets were

positioned anterior-dorsal and posterior-ventral to a tethered flying *Drosophila* (Fig. 1A). Each

electromagnet was powered from a variable power supply (TP1803D, Tekpower, Montclair, CA,

USA) switched through a power transistor (D1276A, Panasonic, Kadoma, Japan) and controlled

via 5V TTL-level output from a data acquisition board (USB-6434, National Instruments, Austin

TX, USA). Pulse trains consisted of 50% duty cycle square wave signals alternating at 25, 36,

50, 72, 100, or 167Hz. Signals sent to the two magnets were 180° out of phase with one another

to produce a dorsal-ventral oscillation of the filing-treated haltere at the commanded frequency.

Electromagnets were hand-wound using electromagnet wire and, as previously estimated

(Rauscher and Fox, 2021), the field strength of each magnet was approximately 2000μT at the

distance of the fly's body from the axial tip of each iron core (2mm).

*Visual stimulation*

To elicit optomotor responses, we used a wide-field vertical stripe pattern moving in the

yaw aspect under the control of a 1Hz triangle wave, which was presented to the fly using a

modular LED flight arena (Reiser and Dickinson, 2008). Stripes were one or two pixels wide and

pseudo-randomly spaced such that average luminance of the wide-field panorama was 50%; i.e.

the pattern consisted of an equivalent number of bright and dark pixels, where one pixel

encompassed 3.75° of the circumference of the display (Fox et al., 2014). Different pattern

velocities were achieved by varying the amplitude of the triangle wave whilst keeping the period

constant.

*High-speed video and kinematic measurements*

Haltere and wing kinematics were recorded using two synchronized high-speed cameras

(TS4 or IL5, Fastec Imaging, San Diego, CA, USA) with sagittal views of the fly from the left

and right aspect. Recordings were made at 1000 or 2000fps for all treated animals and the

majority of untreated animals, however some untreated animals were recorded at 750fps. In each

video, the position of the haltere bulb was digitized using the DLTdv8 (Hedrick, 2008) or

DeepLabCut (Mathis et al., 2018) software packages. Static points were manually labeled for

each video at the position of the haltere base and anterior wing base. These were used to

construct a common body-normalized coordinate framework for registering tracking data across

animals, with the haltere base becoming the origin and the anterior wing base becoming

coordinate (1,0). Wingbeat elevation time series were extracted using a custom MATLAB script.

Haltere stroke amplitudes were estimated by measuring the range of each angular position time

series. For asynchronously-driven trials, video segments shorter than the minimum period of a

single cycle of the given frequency were excluded. A third camera recording at 100fps (Point

Grey Chameleon3, FLIR, Wilsonville, OR, USA) provided a dorsal view used to measure head

and wing kinematics. Head yaw with respect to the body axis and wing downstroke envelope

kinematic time series were computed using a custom MATLAB program called Flyalyzer

(https://github.com/michaelrauscher/flyalyzer), introduced previously in Rauscher and Fox,

2021. Shutter signals from each camera were recorded on the DAQ to facilitate alignment and

synchronization of kinematic time series with one another and with the magnet control pulses

and visual stimuli from the LED flight arena.

*Saccade detection and analysis*

Wing and head saccades were identified in kinematic traces from the 100fps camera

using the amplitude (4° amplitude change in a single frame) and duration (10-100ms) thresholds

described in Cellini et al., 2021. All detected saccades were used to calculate the saccade rates

shown in Fig. 2C-E and Fig. 5C-D. For the saccade-triggered average (STA) responses shown in

Fig. 2F-G (and Fig. S2B) as well as the stimulus-aligned cyclic average responses shown in Fig

S2B-E, saccades were only included from 100fps video segments with concurrent high-speed

tracking of the haltere and wingbeat on the lateral aspect cameras. Each 80ms saccade-triggered

window used to produce the STA was baseline-subtracted and matched to a control window

selected with a randomly drawn frame index from the same video segment. To produce the

histograms shown in Fig. 3 and the heatmaps shown in Fig. S3, data were pooled from all

available asynchronously-driven stimulus categories (up to and including 167Hz). Each frame

from the 100fps camera was assigned a haltere position and wing instantaneous phase value at

frame start by finding the single nearest-match frame recorded on the high-speed lateral aspect

cameras (1000 or 2000 fps). Haltere angular velocity was computed as the numerical derivative

of each haltere angular position time series. To assign a single angular velocity value to each

frame from the 100fps camera, the root-mean-square value of the angular velocity time series

was computed for all concurrent frames from the lateral aspect camera. Then, the sign (positive

or negative) of the lateral aspect frame with the largest absolute value was applied to the result.

A graphical summary of this process is included in Fig. S3A.

*Frequency-domain analyses*

For the estimation of broadband noise (Fig. S4), the fast Fourier transform (FFT) was

calculated using only the first 150 samples of applicable kinematic time series from the dorsal-

view camera to provide a consistent spectral resolution of 0.67Hz. Time series were DC-leveled

by fitting a least-squares regression line and then subtracting it out to remove linear trends. The

area under the curve (AUC) of the resultant FFT was then estimated using the trapezoid method.

For magnetic stimulation trials, we excluded a 4Hz range centered on the stimulus frequency

from the AUC determination. For 50Hz trials, which were at the Nyquist limit for the dorsal

view camera, an interval spanning 46-50Hz was instead excluded from the area under the curve

measurement. For trials at frequencies exceeding the Nyquist limit for the dorsal view camera

(72Hz, 100Hz, 167Hz), the excluded range encompassed the baseband alias of the driven haltere

frequency. A range spanning 23-27Hz was excluded from trials without magnetic stimulation

(untreated, haltere immobilized, single or double haltere ablation).

For estimation of visual response magnitude (Fig. 5), kinematic time series were DC-

leveled as described above and zero-padded to 4096 samples. Magnitude estimates from

resultant FFTs were then divided by the number of samples in each time series prior to zero-

padding to normalize magnitudes for trials of varying length. Only segments longer than 64

samples were used, resulting in a coarsest possible spectral resolution of 1.5Hz.

**Results**

*Electromagnetic control of haltere stroke kinematics*

To establish magnetic control over one of the fly's two halteres, an iron filing was

attached to the bulb (see Methods, Fig. 1A). With the electromagnets deactivated, this mass

immobilized the haltere and prevented the fly from oscillating it. Activation of the

electromagnets at 25, 36, 50, 72, 100, and 167 Hz resulted in movement of the haltere at the

commanded frequency (Fig. 1B-F, Video S1). The natural oscillation of the halteres and wings

produces an out-of-phase relationship and commanding the haltere's frequency allowed us to

sample wingbeats with all possible phase relationships. Haltere tracking data was registered into

a unified coordinate framework, normalized using the haltere-thorax and wing-thorax joints as

common control points, allowing us to survey movement of the haltere both within and in excess

of its naturalistic range of motion (Fig 1C). In all observed trials, the untreated haltere

maintained its full stroke amplitude (Fig. 1G). At the highest frequency tested, 13 out of 14 flies

were unable to maintain a haltere amplitude that was within the range of those observed naturally

(Fig 1G, 22 of 27 replicate trials). Because this commanded haltere motion was significantly

smaller than those observed at lower frequencies, we excluded 167 Hz trials from further

analyses that compared across conditions (see Methods). All conditions (save for the

immobilized haltere) achieved peak angular velocities within the range of those observed in

untreated animals (Fig 1H). Baseline wingbeat frequencies were statistically indistinguishable

from untreated controls for all treatments (Fig. 1I), showing that our manipulations did not

impair the gross functioning of the wing motor system.

*Asynchronous haltere movement evokes wing saccades*

Haltere primary afferents form synaptic connections with head- and wing-steering

motoneurons either directly (Fayyazuddin and Dickinson, 1996; Fayyazuddin and Dickinson,

1999) or via interneurons (Strausfeld and Seyan, 1985; Trimarchi and Murphey, 1997). These

thoracic microcircuits mediate a suite of compensatory equilibrium reflexes stabilizing the

orientation of the body and head during flight (Dickinson, 1999; Sherman and Dickinson, 2003;

Sherman and Dickinson, 2004). To investigate the impact of asynchronous haltere input on these

reflexes, we measured each animal's head yaw and wingstroke amplitudes (Fig. 2A, Video S2)

using machine vision software developed previously in Rauscher and Fox, 2021.

We noted that treated flies showed significantly higher rates of wing saccades than the

untreated control animals. Saccades in *Drosophila* and other flies are abrupt changes in heading,

analogous to the optokinetic reflexes of vertebrates (Cellini and Mongeau, 2020; Cellini et al.,

2021; Land, 1973; Land and Collett, 1974). In tethered flight, saccades appear as sharp changes

in wing amplitude (Fig. 2B, Video S2). For each condition, we calculated the saccade rate (after

Cellini et al., 2021) and found that all treatments exhibited higher ipsilateral wing saccade rates

than untreated controls. Active asynchronous drive of the haltere at any frequency resulted in

ipsilateral wing saccade rates comparable to those seen with bilateral haltere ablation, whereas

haltere immobilization resulted in intermediate saccade rates comparable to those seen with

unilateral ablation (Fig 2C). For contralateral wing saccades (Fig 2D), all treatment groups

exhibited higher saccade rates than untreated control animals. In contrast, the head saccade rate

was insensitive to haltere manipulation under all tested conditions save for bilateral ablation (Fig

2E), in agreement with previous findings that proprioceptive mechanosensory inputs, likely in

the neck, are the primary driver of head saccades (Cellini et al., 2021).

*Haltere kinematic parameters influence saccade initiation*

In individual trials, fast Fourier transforms (FFTs) of wing and head kinematic time series

(Fig S1) showed elevated spectral power at the magnetic stimulation frequency (or its baseband

alias, for frequencies which exceeded the Nyquist limit of the camera measuring the wingstroke

amplitude). Does the increased saccade rate explain the increase in power at the stimulation

frequency? If halteres manipulations drive saccades, the saccade rate should increase with

increasing stimulus rate. Surprisingly, the average rate of ipsilateral wing saccades was uniform

across magnetic stimulation frequencies, and that rate was indistinguishable from that seen with

bilateral haltere ablation. Despite this, saccade-triggered average responses of the ipsilateral

wing amplitude (Fig. 2F) and instantaneous wingbeat frequency (Fig. 2G) followed the

frequency of the driving stimulus. As well, stimulus-triggered cyclic averages aligned on each

magnetic stimulation pulse showed that saccade probability varied in time with the stimulus (Fig.

S2A). Taken together, these    data suggest that saccades occurred at specific times relative to the

driving stimulus, but were not reliably elicited upon every stroke cycle. Saccades accompanying

all haltere treatments were categorically of larger amplitude than those observed in untreated

animals (Fig. S2B). Cyclic averages also illustrated that while the treated haltere's frequency and

amplitude could be reliably commanded with the magnetic stimulation, the induced oscillation

assumed a diversity of phase relationships with the magnetic stimulus across animals and

between trials (Fig. S2C-E). This technical limitation implies that the timing of the stimulus

could not necessarily predict the specific positions or phases of the haltere: these had to be

measured directly from the high-speed video.

After observing phase-locked saccades, we sought to determine whether specific

kinematic parameters of the haltere drove saccade initiation. Using all available high-speed

frames, we assessed the haltere's angular position, angular velocity, and phase relationship to the

wingstroke cycle (Fig 3A) and found that our manipulations resulted in values both within and

outside of the range of values observed in untreated animals. This afforded the opportunity to

observe responses to both naturally-occurring stimuli and stimuli that do not occur in flight.

We then created histograms assessing the probability of saccade initiation as a function of

each kinematic parameter under each condition (Fig. 3B-C, Fig S3).  Saccade probability was not

related to haltere angular position under any condition (Fig. 3B). Similarly, there was no

relationship between the wing phase and the saccade probability under any condition (Fig. 3D).

In untreated animals, the haltere's angular velocity was almost always high, whether the haltere

was moving up or down (Fig. 3C, gray bars); in haltere-immobilized animals, the angular

velocity was always low. Saccade probability was also low in these conditions, although higher

than in untreated animals. When we drove the haltere in a velocity range either faster or slower

than it's natural activity, however, we saw a dramatic increase in saccade probability (Fig. 3C,

right). Multivariate probability heatmaps of multiple kinematic parameters did not reveal any

trends over and above those outlined in univariate analysis (Fig. S3). Our data therefore show

that flies respond to unnatural haltere velocities by initiating wing saccades, with little

discernable influence from other measured kinematic parameters.

*Asynchronous haltere movement increases wingstroke amplitude variability*

295           Bilateral haltere ablation increases wingstroke amplitude variability (Dickinson, 1999;

Mureli and Fox, 2015). Could the wing's amplitude stability be "rescued" with any periodic

excitation from the halteres, or must this input be synchronous with the wings? Because

variability in the time domain includes periodic input from the driven haltere (reflecting the

saccadic steering changes discussed above), we estimated the broadband noise level in the

frequency domain. This allowed us to estimate changes in the baseline amplitude variability,

independent of the saccades, by measuring the area under the curve of the spectrum, excluding a

narrow frequency band centered on the driven haltere frequency (see Methods, Fig. S5A-B).

303           Haltere immobilization increased the broadband noise in the ipsilateral wing amplitude

(Fig. S5C) to a level comparable to that seen with unilateral ablation, whereas active

asynchronous drive using the electromagnets increased it further to levels equivalent to those

seen with bilateral ablation. Similar responses of a smaller magnitude were observed for the

contralateral wing (Fig. S5D), with all unilateral manipulations being distinct from those seen

with untreated animals and bilateral ablation. Thus, not only does wing-asynchronous periodic

input fail to stabilize the wing amplitude, it is categorically worse than no input at all. No haltere

manipulation resulted in head yaw noise levels distinct from untreated animals, similar to our

results when measuring saccade rate (Fig. S5E).

*Haltere-evoked saccades are not influenced by concurrent visual stimulation*

A major task of the fly thoracic nervous system is to integrate inertial sensory

information from wingbeat-synchronous haltere afferents with descending information from the

fly's motion vision system (Namiki et al., 2018; Namiki et al., 2022; Suver et al., 2016).

Tethered flies adjust their gaze and wingstroke amplitude in response to widefield visual

movement (Duistermars et al., 2007; Hengstenberg et al., 1986) acting through a motor pool

shared with complementary reflexes driven by the halteres. This circuit architecture has been

modeled as a "nested feedback" control system, in which "inner" feedback loops mediated by

haltere-wing and haltere-head projections are hierarchically nested within an "outer loop"

mediated by vision (Cellini and Mongeau, 2022; Schwyn et al., 2011). Within this framework,

visual inputs can modulate haltere-mediated reflexes. We asked: does descending visual input

influence wing saccades evoked by haltere stimulation?

Concurrently with each class of haltere treatment, we presented flies with a wide-field

pattern of high-contrast vertical stripes moving at 1Hz in the yaw aspect with a pattern velocity

of 90°/s (Fig. 4A-B,Video S3). We then subdivided each visual stimulus epoch into periods of

clockwise and anticlockwise visual motion (Fig. 4A-B) and calculated the saccade rate during

each of these (as in Fig. 2).  In both visual stimulus directions and for both wings, visual

stimulation did not affect the saccade rate under the majority of haltere stimulus conditions (Fig.

4C-D). Notably, however, concurrent visual stimulation in either direction reduced the rate of

ipsilateral wing saccades when the haltere was immobilized. For the contralateral wing, the

saccade rate was significantly lower only for the anticlockwise direction, although the saccade

rate in the clockwise condition was itself statistically indistinguishable from both the

anticlockwise and uniform visual stimulus conditions. Our results suggest that descending visual

input cannot "rescue" the wing from the destabilizing saccades resulting from asynchronous

haltere input.

*Asynchronous haltere movement disrupts wing and head optomotor responses*

Halteres adjust the gain of wing and gaze optomotor equilibrium reflexes (Bartussek and

Lehmann, 2016; Cellini and Mongeau, 2022; Mureli and Fox, 2015; Mureli et al., 2017;

Rimniceanu et al., 2023). Is synchronous haltere input necessary for this function? To determine

this, we estimated the optomotor response amplitude (following Rimniceanu et al., 2023) by

measuring the magnitude of the FFT of each kinematic time series at 1Hz, the frequency of the

visual stimulus (Fig. 5A-C). For the ipsilateral wing and head (Fig. 5D,F), all haltere

manipulations resulted in categorically smaller visual response amplitudes than those seen in

untreated flies. For the contralateral wing, only the bilateral ablation was statistically distinct

from untreated animals (Fig. 5E).

Bilateral haltere ablation (as well as haltere immobilization) impairs the ability of flies to

adjust their head responses to widefield visual stimuli, instead locking the fly into a fixed low-

amplitude response to all salient pattern velocities (Mureli et al., 2017). To assess the influence

of our manipulations on optomotor tuning, we presented a subset of flies with a series of visual

stimuli that moved at a range of pattern velocities spanning 30-240°/s. Untreated flies exhibited

an optomotor tuning curve in response to these stimuli that peaked at 180°/s for wing responses

(Fig. 5G-H) and 120°/s for head responses (Fig. 5I), in broad agreement with previous studies

(Duistermars et al., 2007; Mureli et al., 2017). Similar tuning curves were observed for flies with

immobilized halteres (Fig. 5G-I) although with significantly lower response amplitudes than the

untreated flies (Fig. S5D). With the magnets engaged at 36Hz, optomotor tuning was abolished

for both contralateral and ipsilateral wings. Visual pattern velocity was not a significant predictor

of response amplitude for these flies (Fig 5G-H).  In comparison, head responses showed lower

gain relative to untreated controls, at levels comparable to those seen with haltere immobilization

(Fig. 5I,S5C,F).

**Discussion**

Coordinated motion of multiple appendages is so important that it is often very difficult

to force them to move in uncoordinated ways. In flies, the biomechanics of the thorax restricts

the halteres and wings to oscillate in perfect antiphase with no input from the nervous system:

elevating the wing of a dead fly with forceps depresses the haltere (Deora et al., 2015). We used

stimuli both within and outside the bounds of natural haltere motions to demonstrate the effects

of asynchronous haltere perturbations on wing and head movements in flying flies. In previous

studies, haltere ablation or immobilization was demonstrated to be deleterious to wing amplitude

stability (Dickinson, 1999; Mureli and Fox, 2015) and optomotor reflexes (Bartussek and

Lehmann, 2016; Mureli and Fox, 2015; Mureli et al., 2017), but these works left unresolved how

specific haltere information might contribute to each of these behavioral outputs. Consistent with

our hypothesis that altering the phase of the haltere and wing should destabilize the wing motor

output, our data demonstrate that both ipsilateral and contralateral wing steering reflexes are

critically impaired by asynchronous haltere input, with increased saccades following "unnatural"

haltere movements.

Though we hypothesized that haltere input of any kind would be sufficient for full-

amplitude visually guided head movements, we instead observed that synchronous haltere input

is necessary for attaining naturalistic levels of optomotor response gain. The head's movements,

though receiving a more diverse set of inputs from both descending and ascending neurons and

occurring at a lower speed, still shows deleterious effects of asynchronous haltere input.

Asynchronous haltere input did not lead to higher rates of head saccades, congruent with

previous findings showing that head saccades are primarily evoked by proprioceptive inputs to

"reset" the head when it reaches the limits of its range of travel (Cellini et al., 2021).

*Asynchronous haltere input releases wing saccades, elevates wingstroke variability*

Our data show that asynchronous haltere input increases the saccade rate of the ipsilateral

wing, and that these saccades are more likely when the haltere moves at angular velocities

outside of its natural operating range. Experiments in which the entire body was rotated showed

that the halteres are most sensitive to body angular velocity (Dickinson, 1999); our results extend

this to show a relationship between the haltere's own angular velocity and movements of the

wings. Are these saccades being driven by specific motions of the haltere? Our experiment with

immobilized haltere suggests not: loss of one or both halteres also results in an elevated rate of

saccades of similar amplitude. Therefore, another possible interpretation is that saccades emerge

spontaneously from the activity of the wing motor system, and that synchronous haltere input at

specific haltere positions and velocities acts to suppress them.

This view accords well with our other finding that wingstroke amplitude variability also

increases with the loss of input from one or both halteres in a stepwise fashion. This corroborates

physiological recordings of the motoneuron controlling the first basalare muscle in *Calliphora*

showing higher timing variability in spikes evoked by reafferent stimulation of the wing nerve as

compared with those evoked by stimulation of the haltere nerve (Fayyazuddin and Dickinson,

1999). It is further corroborated by recent work in which behavior was recorded from the same

individual flies under both a rigid body-fixed paradigm as well as a magnetic-suspension

paradigm in which the body can rotate in the yaw plane (Rimniceanu et al., 2023). In the fixed

condition, the wingstroke amplitude variability is categorically higher than when the same flies

are freed to rotate about the yaw axis.

Campaniform sensilla are sensitive to compressive and shearing forces acting on and

within the cuticle in which they are embedded (Chapman and Smith, 1963; Pringle, 1948; Zill

and Moran, 1981). Such forces arise from the haltere's movement through its natural stroke

plane and therefore scale proportionately with the bulb's angular velocity (Nalbach, 1993). As

hypothesized by Pringle (1948) and predicted in subsequent modeling (Mohren et al., 2019),

different fields of campaniform sensilla should also experience forces preferentially in-line with

their receptive fields at different points within the stroke cycle. We might therefore predict that

saccades are more or less likely to be initiated as a function of the haltere's angular position. Our

results suggest that saccades may be slightly more likely when the asynchronously-driven haltere

was in the downstroke position, but we did not observe a strong correspondence between haltere

angular position and saccade initiation.

Though our data support the hypothesis that halteres suppress spontaneous saccades,

rather than drive endogenous saccades, the extreme sensitivity of the campaniform sensilla

forces us to consider the possibility that we were not able to stimulate or measure the haltere

stimuli accurately enough to reach this conclusion. The firing properties of primary haltere

afferents (Yarger and Fox, 2018) and downstream neck motoneurons (Haag et al., 2010)change

significantly when the haltere oscillates naturally versus when its motion is driven exogenously,

even in similar planes and frequencies. Additionally, our experimental setup was not optimized

to measure all wingstroke parameters that have been shown to be relevant to steering output.

While we have measured the peak downstroke angle of the wing as our proxy for steering effort,

the total force produced by each wingstroke is a complex function of not just the downstroke

angle, but also the wing joint deviation angle and the spanwise rotation angle of the wing

(Muijres et al., 2015).

*Haltere influence on the contralateral wing*

In previous work (Rauscher and Fox, 2021), we manipulated the stroke amplitude of a

beating haltere without compromising the haltere-wing stroke synchrony and evoked a consistent

wing-steering response to the magnetic stimulation. Here, we found the response to haltere

perturbation was less stereotyped and more probabilistic, but in neither the present nor previous

work did we observe a clear relationship between the instantaneous wing phase and haltere-

evoked changes in wing kinematic parameters. This was surprising given the organization of the

wing-steering motor system downstream of haltere primary afferents, a major component of

which are tonic muscles which fire once every stroke cycle and control muscle tension by

varying their firing phase within it (Balint and Dickinson, 2001; Heide and Götz, 1996; Lindsay

et al., 2017; Tu and Dickinson, 1996). One possibility is that the haltere-evoked saccades in our

data are mediated primarily by phasic muscles, for which the wing phase is comparatively less

important for force generation. Another hypothesis is that the untreated opponent haltere (present

in all of our manipulations) provides redundant input to contralateral wing motoneurons that can

supersede an absent or malfunctioning ipsilateral haltere.

Our findings that asynchronous haltere input evokes wing saccades and increases the

stroke amplitude variability of the contralateral wing are consistent with previous findings that

each wing receives information from both halteres. Flies are capable of apparently normal flight

following unilateral haltere ablation (Fraenkel, 1939). In *Calliphora,* input from a single haltere

is sufficient to entrain the firing of the contralateral first basalare muscle following the ablation

of the ipsilateral haltere and both wing nerves (Heide, 1983).  Experiments rotating the fly's

entire body showed that both amplitude and frequency components of haltere-wing reflexes are

impaired following bilateral—but not unilateral—haltere ablation (Dickinson, 1999). Similar

results were observed using a magnetic-suspension tethering system in which the fly initiated its

own rotations: manipulating the bulb mass of a single haltere produced comparable changes in

both clockwise and counterclockwise body saccades (Bender and Dickinson, 2006).

This bilateral integration implicates the contralateral haltere interneurons, or cHINs

(Strausfeld and Seyan, 1985; Trimarchi and Murphey, 1997), which receive input from haltere

afferents and represent the only known anatomical pathway through which haltere information

crosses the body axis for both the wing and gaze motor control systems. Though anatomically

conserved across fly taxa, these neurons have eluded physiological characterization to date,

leaving many questions unresolved as to the sensory representations and transformations they

mediate *en route* to their downstream motor targets. As we were unable to reliably elicit tethered

flight following any manipulation of both halteres aside from total bilateral ablation, behavioral

explorations of this circuit's function are unlikely to be successful. Further exploration of each

haltere's influence on the contralateral wing will likely remain contingent on a more complete

description of the anatomy and physiology of haltere-wing thoracic microcircuits.

*Haltere contribution to head and wing optomotor responses*

As descending visual inputs concurrently target wing and gaze motoneurons (Chan et al.,

1998; Namiki et al., 2018; Strausfeld and Seyan, 1985), we also examined multisensory

integration of haltere and visual inputs. We show that haltere-wing asynchrony critically impairs

the wing optomotor response to a degree comparable with bilateral haltere ablation. While this

impairment may be unsurprising for wings, it is less obvious how the head might respond to

asynchronous haltere input. Bilateral haltere ablation prevents flies from adjusting the gain of the

head optomotor response to different pattern velocities (Mureli et al., 2017). Here, we show that

loss of one haltere (by immobilization) or asynchronous input from one haltere reduces the

optomotor response gain, but does not abolish pattern velocity tuning (Fig. 6).

In some of the neck motoneurons of quiescent *Calliphora*, an absence of haltere input

prevents spiking responses to visual input, but spiking can be restored with haltere movement at

any frequency, even as low as 10 Hz (Huston and Krapp, 2009). If the halteres are necessary

only to provide flight-state-dependent excitatory drive onto the neck motor system, then

asynchronous input would produce gaze optomotor tuning curves with similar or equivalent gain

to an unmanipulated animal. Our data do not show this. Instead, we found that asynchronous

drive of a single haltere resulted in head responses with lower gain. There are two main

hypotheses that could explain why neck motoneurons that can be excited by asynchronous

haltere input in a quiescent state fail to produce high-gain visually-guided head movements when

stimulated asynchronously in tethered flight.

First, the wing sensory inputs to the neck motoneurons are still unknown, raising the

possibility that full neck motoneuron activity may require both wing and haltere input, like the

wing motoneuron mnb1 (Fayyazuddin and Dickinson, 1996). This would allow the neck

motoneurons to operate differently in flying and quiescent states. However, flies with ablated

wings are still able to execute high-gain optomotor responses in tethered "flight" (thorax and

halteres oscillating; Mureli et al., 2017) ).  Though this would not preclude a role for wing

reafference in head optomotor responses, as some mechanosensors (e.g. chordotonal organs at

the base) may still function in the absence of wing aerodynamic loading, the complete wing

sensory complement is unnecessary.

Second, the "control loop," in which descending neurons provide excitatory input to

haltere muscles, (Chan et al., 1998; Dickerson et al., 2019) may explain this result. Our

experiment breaks the control loop by nullifying the relationship between haltere muscle activity

and haltere sensory input: visual input to the haltere muscles no longer produces haltere input

that can be used to initiate proper reflexes. Both hypotheses could explain our results that visual

input cannot "rescue" the wing from high noise and high saccade rates induced by haltere

asynchrony (Fig. 4), and that visual tuning in the head remains intact but at lower gain. Future

recordings of the electrophysiological properties of neck motoneurons under different behavioral

conditions and of the haltere movement responses to visual inputs could help untangle these

possibilities.

**Acknowledgments**

We would like to thank Chenxin Bi and Brandon Chun for help with data collection and

annotation, as well as Noah DeFino, Jesse Fritz and Lauren Metz for fly care. We would also like

to thank Hillel Chiel, Bradley Dickerson, Jeremiah Didion, Alexandra Gurgis, Nicholas

Kathman, Kristianna Lea, Mark Willis, Gabriella Wolff, and Alexandra Yarger for helpful

commentary.

**Competing interests**

No competing interests declared.

**Funding**

This work was supported by United States Air Force Office of Scientific Research [FA9550-14-

0398 and FA9550-16-1-0165 to JLF]; and National Science Foundation [1754412 to JLF].

**Data Availability**

All processed data is available in the associated Dryad repository. Raw video data available upon

request. Kinematic time series for some untreated control animals in Figures 1-5 are included

with a previously-published publicly available dataset (Rauscher and Fox, 2020).

**References Cited**

**Agrawal, S., Grimaldi, D. and Fox, J. L.** (2017). Haltere morphology and campaniform sensilla
arrangement across Diptera. *Arthropod Struct Dev* **46**, 215–229.

**Balint, C. N. and Dickinson, M. H.** (2001). The correlation between wing kinematics and steering muscle
activity in the blowfly Calliphora vicina. *J Exp Biol* **204**, 4213–4226.

**Bartussek, J. and Lehmann, F.-O.** (2016). Proprioceptive feedback determines visuomotor gain in
Drosophila. *R Soc Open Sci* **3**, 150562.

**Bender, J. A. and Dickinson, M. H.** (2006). A comparison of visual and haltere-mediated feedback in the control of body saccades in Drosophila melanogaster. *Journal of Experimental Biology* **209**, 4597–606.

**Cellini, B. and Mongeau, J. M.** (2020). Active vision shapes and coordinates flight motor responses in flies. *Proc Natl Acad Sci U S A* **117**, 23085–23095.

**Cellini, B. and Mongeau, J. M.** (2022). Nested mechanosensory feedback actively damps visually guided head movements in Drosophila. *Elife* **11**,.

**Cellini, B., Salem, W. and Mongeau, J. M.** (2021). Mechanisms of punctuated vision in fly flight. *Current Biology* **31**, 4009-4024.e3.

**Chan, W. P., Prete, F. and Dickinson, M. H.** (1998). Visual input to the efferent control system of a fly's "gyroscope". *Science* **280**, 289–292.

**Chapman, K. M. and Smith, R. S.** (1963). A linear transfer function underlying impulse frequency modulation in a cockroach mechanoreceptor. *Nature 1963 197:4868* **197**, 699–700.

**Deora, T., Singh, A. K. and Sane, S. P.** (2015). Biomechanical basis of wing and haltere coordination in flies. *Proc Natl Acad Sci U S A* **112**, 1481–6.

**Derham, W.** (1714). *Physico-theology: or, a demonstration of the being and attributes of God, from His works of creation. Being the substance of XVI sermons preached in St. Mary Le Bow-church, London, at the honble Mr. Boyle's lectures, in the years 1711 and 1712.* W. and J. Innys, London, 1714.

**Dickerson, B. H., de Souza, A. M., Huda, A. and Dickinson, M. H.** (2019). Flies regulate wing motion via active control of a dual-function gyroscope. *Current Biology* **29**, 3517–3524.

**Dickinson, M. H.** (1990). Comparison of encoding properties of campaniform sensilla on the fly wing. *Journal of Experimental Biology* **261**, 245–261.

**Dickinson, M. H.** (1999). Haltere-mediated equilibrium reflexes of the fruit fly, Drosophila melanogaster. *Philosophical Transactions of the Royal Society B: Biological Sciences* **354**, 903–916.

**Dickinson, M. H. and Muijres, F. T.** (2016). The aerodynamics and control of free flight maneuvers in Drosophila. *Philosophical transactions of the Royal Society of London B: Biological Sciences* **371**, 20150388.

**Duistermars, B. J., Reiser, M. B., Zhu, Y. and Frye, M. A.** (2007). Dynamic properties of large-field and small-field optomotor flight responses in Drosophila. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **193**, 787–799.

**Fayyazuddin, A. and Dickinson, M. H.** (1996). Haltere afferents provide direct, electrotonic input to a steering motor neuron in the blowfly, Calliphora. *J Neurosci* **16**, 5225–5232.

**Fayyazuddin, a and Dickinson, M. H.** (1999). Convergent mechanosensory input structures the firing phase of a steering motor neuron in the blowfly, Calliphora. *J Neurophysiol* **82**, 1916–1926.

**Fox, J. L. and Daniel, T. L.** (2008). A neural basis for gyroscopic force measurement in the halteres of Holorusia. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **194**, 887–897.

**Fraenkel, G.** (1939). The function of the halteres of flies (Diptera). *Proceedings of the Zoological Society of London* **A109**, 69–78.

**Gnatzy, W., Grünert, U. and Bender, M.** (1987). Campaniform sensilla of Calliphora vicina (Insecta, Diptera). *Zoomorphology* **106**, 312–319.

**Haag, J., Wertz, A. and Borst, A.** (2010). Central gating of fly optomotor response. *Pnas* **107**, 20104–20109.

**Hall, J. M., McLoughlin, D. P., Kathman, N. D., Yarger, A. M., Mureli, S. and Fox, J. L.** (2015). Kinematic diversity suggests expanded roles for fly halteres. *Biol Lett* **11**, 20150845.

**Heckscher, E. S., Zarin, A. A., Faumont, S., Clark, M. Q., Manning, L., Fushiki, A., Schneider-Mizell, C. M., Fetter, R. D., Truman, J. W., Zwart, M. F., et al.** (2015). Even-skipped+ interneurons are core components of a sensorimotor circuit that maintains left-right symmetric muscle contraction amplitude. *Neuron* **88**, 314–329.

**Hedrick, T. L.** (2008). Software techniques for two- and three-dimensional kinematic measurements of biological and biomimetic systems. *Bioinspir Biomim* **3**, 34001.

**Heide, G.** (1983). Neural mechanisms of flight control in Diptera. In *Biona-report 2* (ed. Fischer, G.), pp. 35–52. Stuttgart, New York.

**Heide, G. and Götz, K. G.** (1996). Optomotor control of course and altitude in Drosophila melanogaster is correlated with distinct activities of at least three pairs of flight steering muscles. *Journal of Experimental Biology* **199**, 1711–1726.

**Hengstenberg, R., Sandeman, D. C. and Hengstenberg, B.** (1986). Compensatory head roll in the blowfly Calliphora during flight. *Proceedings of the Royal Society B: Biological Sciences* **227**, 455–482.

**Hiebert, G. W. and Pearson, K. G.** (1999). Contribution of sensory feedback to the generation of extensor activity during walking in the decerebrate Cat. *J Neurophysiol* **81**, 758–70.

**Huston, S. J. and Krapp, H. G.** (2009). Nonlinear integration of visual and haltere inputs in fly neck motor neurons. *Journal of Neuroscience* **29**, 13097–13105.

**Kim, A. J., Fitzgerald, J. K. and Maimon, G.** (2015). Cellular evidence for efference copy in Drosophila visuomotor processing. *Nat Neurosci* **18**, 1247–1255.

**Land, M. F.** (1973). Head movement of flies during visually guided flight. *Nature 1973 243:5405* **243**, 299–300.

**Land, M. F. and Collett, T. S.** (1974). Chasing behaviour of houseflies (Fannia canicularis). *Journal of comparative physiology 1974 89:4* **89**, 331–357.

**Lindsay, T., Sustar, A. and Dickinson, M. H.** (2017). The function and organization of the motor system controlling flight maneuvers in flies. *Current Biology* **27**, 345–358.

**Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W. and Bethge, M.** (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* **21**, 1281–1289.

**Mohren, T. L., Daniel, T. L., Eberle, A. L., Reinhall, P. G. and Fox, J. L.** (2019). Coriolis and centrifugal
forces drive haltere deformations and influence spike timing. *J R Soc Interface* **16**, 20190035.

**Muijres, F. T., Elzinga, M. J., Iwasaki, N. A. and Dickinson, M. H.** (2015). Body saccades of Drosophila
consist of stereotyped banked turns. *Journal of Experimental Biology* **218**, 864–875.

**Mureli, S. and Fox, J. L.** (2015). Haltere mechanosensory influence on tethered flight behavior in
Drosophila. *Journal of Experimental Biology* 2528–2537.

**Mureli, S., Thanigaivelan, I., Schaffer, M. L. and Fox, J. L.** (2017). Cross-modal influence of
mechanosensory input on gaze responses to visual motion in Drosophila. *Journal of Experimental*
*Biology* **220**, 2218–2227.

**Nalbach, G.** (1993). The halteres of the blowfly Calliphora. *Journal of Comparative Physiology A* **173**,
293–300.

**Namiki, S., Dickinson, M. H., Wong, A. M., Korff, W. and Card, G. M.** (2018). The functional organization
of descending sensory-motor pathways in drosophila. *Elife* **7**, e34272.

**Namiki, S., Ros, I. G., Morrow, C., Rowell, W. J., Card, G. M., Korff, W. and Dickinson, M. H.** (2022). A
population of descending neurons that regulates the flight motor of Drosophila. *Current Biology*
**32**, 1–8.

**Noah, J. A., Quimby, L., Frazier, S. F. and Zill, S. N.** (2004). Walking on a "peg leg": Extensor muscle
activities and sensory feedback after distal leg denervation in cockroaches. *J Comp Physiol A*
*Neuroethol Sens Neural Behav Physiol* **190**, 217–231.

**Pringle, J. W. S.** (1948). The gyroscopic mechanism of the halteres of Diptera. *Philosophical Transactions*
*of the Royal Society B: Biological Sciences* **233**, 347–384.

**Rauscher, M. J. and Fox, J. L.** (2020). Weighted haltere and imposed haltere stroke reduction tethered
flying Drosophila kinematics. *Dryad, Dataset*.

**Rauscher, M. J. and Fox, J. L.** (2021). Haltere and visual sensory inputs sum linearly to predict wing (but
not head) motor output in tethered flying Drosophila. *Proceedings of the Royal Society B* **288**,
20202374.

**Reiser, M. B. and Dickinson, M. H.** (2008). A modular display system for insect behavioral neuroscience.
*J Neurosci Methods* **167**, 127–139.

**Rimniceanu, M., Currea, J. P. and Frye, M. A.** (2023). Proprioception gates visual object fixation in flying
flies. *Current Biology* **33**, 1459-1471.e3.

**Schwyn, D. A., Heras, F. J. H., Bolliger, G., Parsons, M. M., Krapp, H. G. and Tanaka, R. J.** (2011).
Interplay between feedback and feedforward control in fly gaze stabilization. *IFAC Proceedings*
*Volumes (IFAC-PapersOnline)* **18**, 9674–9679.

**Sherman, A. and Dickinson, M. H.** (2003). A comparison of visual and haltere-mediated equilibrium
reflexes in the fruit fly Drosophila melanogaster. *Journal of Experimental Biology* **206**, 295–302.

**Sherman, A. and Dickinson, M. H.** (2004). Summation of visual and mechanosensory feedback in
Drosophila flight control. *Journal of Experimental Biology* **207**, 133–142.

**Strausfeld, N. J. and Seyan, H. S.** (1985). Convergence of visual, haltere, and prosternal inputs at neck
motor neurons of Calliphora erythrocephala. *Cell Tissue Res* **240**, 601–615.

**Sun, M., Wang, J., Xiong, Y., Sun, M., Wang, J. and Xiong, Y.** (2007). Dynamic flight stability of hovering
insects. *Acta Mech Sin* **23**, 231–246.

**Suver, M. P., Huda, A., Iwasaki, N., Safarik, S. and Dickinson, M. H.** (2016). An array of descending
visual interneurons encoding self-motion in Drosophila. *The Journal of Neuroscience* **36**, 11768–
11780.

**Trimarchi, J. R. and Murphey, R. K.** (1997). The shaking-B2 mutation disrupts electrical synapses in a
flight circuit in adult Drosophila. *Journal of Neuroscience* **17**, 4700–4710.

**Tu, M. S. and Dickinson, M. H.** (1996). The control of wing kinematics by two steering muscles of the
blowfly (Calliphora vicina). *Journal of Comparative Physiology A 1996 178:6* **178**, 813–830.

**Wannier, T., Bastiaanse, C., Colombo, G. and Dietz, V.** (2001). Arm to leg coordination in humans during
walking, creeping and swimming activities. *Experimental Brain Research 2001 141:3* **141**, 375–379.

**Wiggin, T. D., Peck, J. H. and Masino, M. A.** (2014). Coordination of fictive motor activity in the larval
zebrafish is generated by non-segmental mechanisms. *PLoS One* **9**, e109117.

**Yarger, A. M. and Fox, J. L.** (2018). Single mechanosensory neurons encode lateral displacements using
precise spike timing and thresholds. *Proceedings of the Royal Society B: Biological Sciences* **285**,
20181759.

**Zill, S. N. and Moran, D. T.** (1981). The exoskeleton and insect proprioception. I. Responses of tibial
campaniform sensilla to external and muscle-generated forces in the American cockroach,
Periplaneta americana. *Journal of Experimental Biology* **91**, 1–24.

**Figure 1. Electromagnetic control of haltere movement asynchronous from the wings.**
(**A**) Schematic of iron filing treatment (top, inset) and electromagnetic stimulation system (bottom). (**B**)
Representative traces showing movement of untreated haltere at wingbeat frequency (~200Hz), immobilized haltere
treated with iron filing (ii), and actively-driven haltere at a range of frequencies controlled by electromagnets (iii-
viii) (**C**) Haltere tracking by condition, registered into a normalized body coordinate system defining the haltere base
as the origin, represented by circle and the anterior wing base as (0,1), represented by asterisk. Colors reflect the z-
normalized wingstroke elevation at each measurement point. (**D**) Mean FFT (95% confidence interval shaded) of all
high-speed (1000 or 2000fps) time series with the haltere immobilized (black), driven at 25Hz (red), and 50Hz
(pink). (**E**) same as (D) for 36Hz (blue) and 72Hz (teal) frequencies. (**F**) same as (D) and (E) for 100Hz (orange) and
167Hz (magenta) frequencies. (**G-I**) Haltere stroke amplitude (G), peak angular velocity (H) and baseline wingbeat
frequencies (I) observed under each haltere manipulation. Lines show median for each group. 167Hz responses from
outlier animal shown with +. Letters denote statistical groupings from multiple Bonferroni-corrected Wilcoxon rank-
sum test (5% alpha).

**Figure 2. Haltere manipulation increases saccade rate in kinematic outputs downstream of halteres.**
**(A).** Schematic of measured kinematic outputs, recorded at 100fps, showing downstroke amplitude (angle of wing
envelope leading edge) for each wing and yaw movement of the head. **(B)** Example ipsilateral wing amplitude traces
under haltere treatment conditions. Positive-going angles denote counterclockwise changes in the projected wing
envelope leading edge angle which for the right-hand wing (as in all depicted examples) signifies an increase in
downstroke amplitude. **(C)** Ipsilateral wing saccade rates from all haltere treatments (see Methods). Left and right
wing responses for symmetrical manipulations (untreated, bilateral ablation) are pooled. Lines show median for each
group. Letters denote statistical groupings from multiple Bonferroni-corrected Wilcoxon rank-sum test (5% alpha).
**(D,E)** Same as (C) for contralateral wing and head saccade rates, respectively. **(F)** Saccade-triggered average
ipsilateral wing amplitude responses for each condition (Shade, 95% confidence interval). Gray triangles show the
saccade peak, around which each constitutive video segment of the STA is indexed. Gray traces show controls
triggered from random indices within the same time series. **(G)** Same as (F) but for the baseline-subtracted
instantaneous wingbreat frequency.

697
**Figure 3. Ipsilateral wing saccades are more likely when the haltere exceeds normal kinematic parameters**
**(A)** Summary of haltere angular position, angular velocity, and wing instantaneous phase from all high speed data,
separated by condition. (Ai) Untreated animals, showing natural antiphase synchrony of haltere and wings. Inset
shows unified body coordinate scheme, with haltere upstrokes typically occupying the 45-90° range and
downstrokes typically occupying the 235-270° range (Aii) Frames from haltere immobilized (magnets OFF)
condition. Green dashed line shows extent of the naturalistic parameter space observed in the untreated animal.
(Aiii) High-speed frames aggregated from all magnetic stimulation frequencies, showing broad overlap with the
naturalistic range of haltere angular velocities and angular positions. **(B)** Histograms of ipsilateral wing amplitude
video frames showing probability of saccade as a function of haltere angular position at frame onset for each
condition. Light green shaded region outlines the natural range of haltere angular positions **(C).** Same as (B) but for
haltere signed-RMS angular velocity throughout the frame (see Methods). Light green shaded regions outline the
naturalistic range of negative (upstroke) and positive (downstroke) angular velocities. **(D)**. Same as (B) but for wing
instantaneous phase at frame onset.
711

**Figure 4. Concurrent visual input does not influence saccade rate during most haltere manipulations**
(**A**) Schematic of visual stimulus, showing irregular stripe pattern moving on an LED panel under the control of a
1Hz triangle-wave with a pattern velocity of 90°/s (**B**) Example ipsilateral wing amplitude responses to one cycle of
the visual stimulus under several haltere treatment conditions. (**C**) Ipsilateral wing saccade rate across visual
conditions, showing no change between uniform visual stimulus (reproduced from Fig. 2), clockwise visual stimulus
epochs anticlockwise epochs in most haltere treatment conditions. Mean value for each visual stimulus condition
represented by dash, upward pointing triangle, and downward pointing triangle respectively. (**D**) Same as a (C) but
for contralateral wing saccade rate (*** = significance at 0.1% alpha level, multiple Bonferroni-corrected Wilcoxon
rank-sum test) (D) Same as a (C) but for contralateral wing saccade rate. (* = significance at 5% alpha level,
multiple Bonferroni-corrected Wilcoxon rank sum test)

723
**Figure 5. Haltere manipulations influence visually-guided wing and head movements.**
(**A**) Schematic of visual stimulus, as in Fig. 4A (**B**) Example ipsilateral wing amplitude responses to multiple cycles
of the visual stimulus under several haltere treatment conditions. (**C**) FFTs of traces from (B), showing peak at the
frequency of the visual stimulus. (**D**) Ipsilateral wing response amplitude at visual stimulus frequency for all tested
haltere treatments. For symmetrical manipulations (untreated, bilateral ablation), left and right wing responses are
pooled. Letters denote statistical groupings from multiple Bonferroni-corrected Wilcoxon rank-sum test (5% alpha).
(**E,F**) Same as (D) for contralateral wing and head yaw responses, respectively. Lines show median for each group.
(**G**) Ipsilateral wing visual response magnitude, as estimated by measuring FFT magnitude at visual stimulus
frequency, for untreated flies, haltere-immobilized flies, and 36Hz magnetic stimulation flies. Trendlines connect
median data from each pattern velocity, with shaded region showing 1$^{st}$ and 3$^{rd}$ quartiles. Inset, results of Kruskal-
Wallis omnibus test indicate whether pattern velocity is a significant predictor of response amplitude within each
treatment group at the 5% alpha level (*) or not (ns). (**H-I**) Same as (G), but for contralateral wing and head yaw
amplitude, respectively.