Xanomeline restores endogenous nicotinic acetylcholine receptor signaling in mouse prefrontal cortex

S. K. Power[1], S. Venkatesan[1], E. K. Lambe[1,2,3]

1. Department of Physiology, University of Toronto, Toronto ON, Canada

2. Department of OBGYN, University of Toronto, Toronto ON, Canada

3. Department of Psychiatry, University of Toronto, Toronto ON, Canada

Corresponding author:

E. K. Lambe, Ph.D.
Depts of Physiology, OBGYN, and Psychiatry
Temerty Faculty of Medicine
University of Toronto
1 King's College Circle
Toronto ON
Canada M5S 1A8
evelyn.lambe@utoronto.ca

Abstract: 197 (250 max)

Word count: 3996 (4000 max)

Figures: 5

Supplemental Table: 1

Supplemental Methods: 1

Conflicts: The authors declare no conflicts of interest.

Abstract

Cholinergic synapses in prefrontal cortex are vital for attention, but this modulatory system undergoes substantial pre- and post-synaptic alterations during adulthood. To examine the integrated impact of these changes, we optophysiologically probe cholinergic synapses *ex vivo*, revealing a clear decline in neurotransmission in middle adulthood. Pharmacological dissection of synaptic components reveals a selective reduction in postsynaptic nicotinic receptor currents. Other components of cholinergic synapses appear stable, by contrast, including acetylcholine autoinhibition, metabolism, and excitation of postsynaptic muscarinic receptors. Pursuing strategies to strengthen cholinergic neurotransmission, we find that positive allosteric modulation of nicotinic receptors with NS9283 is effective in young adults but wanes with age. To boost nicotinic receptor availability, we harness the second messenger pathways of the preserved excitatory muscarinic receptors. The muscarinic agonist and cognitive-enhancer xanomeline significantly restores nicotinic signaling in older mice in a muscarinic- and PKC-dependent manner. The rescued nicotinic component regains youthful sensitivity to allosteric enhancement: treatment with xanomeline and NS9283 restores cholinergic synapses in older mice to the strength, speed, and receptor mechanism of young adults. Our results reveal a new and efficient strategy to rescue age-related nicotinic signaling deficits, demonstrating a novel pathway for xanomeline to restore cognitively-essential endogenous cholinergic neurotransmission.

Introduction

Even in healthy aging, there are declines in sustained attention [1,2], cue detection [3,4], and executive function [5,6]. These abilities all rely on cholinergic control of the prefrontal cortex [7], with basal forebrain projections releasing acetylcholine in deep cortical layers [8–11]. Not surprisingly, it is hypothesized that age-related cognitive changes arise from disruption of the cholinergic system [12–14]. Human clinical investigations reveal age-related reductions in basal forebrain volume [15,16], decreases in nicotinic binding [17,18], and possible changes in muscarinic binding [19,20] and acetylcholinesterase [21,22]. Targeting the cholinergic system has shown some promise in treating age-related cognitive changes in humans [23–26][27–29][30–32]). However, treatment strategy is limited by the substantial knowledge gap in the integrated functional impact of age-related changes to prefrontal cholinergic synapses. This uncertainty complicates the identification of specific treatments capable of restoring youthful functional parameters.

Similar to humans, mice show progressive cognitive changes that occur during adulthood [33,34], beginning as early as 4-5 months [35]. Transgenic mice have been used to investigate the complexity of cortical cholinergic synapses [36–39], revealing the functional roles of multiple pre- and post-synaptic receptors and regulatory enzymes [36–39]. However, previous work has not examined the impact of age on these complex synapses. The need to study the integrated impact of age-related changes on endogenous cholinergic signaling is underscored by research suggesting that cholinergic receptor expression is dynamic [40,41] and there may be interactions across receptor subtypes [42–44].

Here, we use optogenetic tools to investigate the impact of age on integrated functioning of prefrontal cholinergic synapses from young to middle adulthood. We demonstrate a striking and selective reduction of the postsynaptic nicotinic optophysiological response while revealing the stability of pre- and postsynaptic muscarinic responses. We discover that the age-related nicotinic functional deficit is not amenable to positive allosteric modulation. Therefore, we harness signaling pathways of the well-preserved muscarinic receptors as an unorthodox approach to nicotinic potentiation. Overall, we demonstrate the strength, speed, and receptor mechanism of cholinergic signaling can be restored in aging and discover a new and efficient strategy to rescue age-related nicotinic signaling deficits.

Materials and Methods

Methods are summarized briefly below. See **Supplemental Methods** for additional description.

Animals

The University of Toronto Faculty of Medicine Animal Care Committee approved all experiments in accordance with the guidelines of the Canadian Council on Animal Care (protocol #20011621). We performed experiments on transgenic mice expressing channelrhodopsin in cholinergic afferents (ChAT-ChR2; RRID:IMSR_JAX:014546, Jackson Laboratory [45]). This transgenic line labels cholinergic neurons and avoids labeling neurons only transiently cholinergic in development [46]. This approach also avoids use of anesthetic agents known to perturb nicotinic receptors [47,48] and surgery known to impact cognitive function [49]. A total of 55 mice were used for this study (age range: ~P50-350), with balanced numbers of males

64 and females. This age range was selected because mice show progressive, age-dependent
65 changes in behavior through adulthood [33–35].

Electrophysiology and Optogenetics

For whole-cell patch-clamp electrophysiology, Layer 6 pyramidal neurons in medial prefrontal
cortex were identified by size, morphology, and proximity to white matter. Responses from a
homogenous population of layer 6 regular spiking corticothalamic pyramidal neurons [39] were
examined in voltage-clamp at -75mV and in current-clamp.

To excite chanelrhodopsin-containing cholinergic afferents, blue light (470nm) was delivered
in 5 ms pulses with an LED (2mW; Thorlabs) through at 60x objective lens. As developed
previously in our lab, this opto-ACh stimulus is delivered in a frequency-accommodating (50Hz
to 10Hz) eight pulse train to stimulate cholinergic axons [39] to mimic the activation pattern of
cholinergic neurons [50–52].

Experimental design, analysis, and statistics

Sex-balanced mice were used across ages and in pharmacological interventions. Experiments
draw from at least 3 mice per intervention, and recordings were made from 1-3 neurons per
brain slice, with 2-3 slices per mouse. Unless otherwise stated, pharmacological agents were
pre-applied for 10 minutes and co-applied during optogenetic stimuli and recovery period.
Magnitude of cholinergic currents were measured from exponential fit by peak amplitude (pA)
and charge transferred (pC) into the cell, measured by area under the current response for 5s
from opto-ACh onset. Opto-ACh responses plotted in a histogram by age were analyzed
statistically by nonlinear robust regression (Graphpad Prism 9.3.1). Age differences in the
presence of pharmacological acetylcholine blockers were analyzed with unpaired and paired,
two-tailed *t*-tests, where applicable. Pharmacological manipulations to improve nicotinic
responses were evaluated with respect to age by two-way ANOVA and one-way ANOVA, where
applicable.

Results

Postsynaptic responses to optogenetic release of endogenous acetylcholine decrease
with age

To stimulate release of endogenous acetylcholine (opto-ACh, **Fig 1A,B**), we use a pattern of
optogenetic stimulation [39] designed to mimic burst firing with spike frequency accommodation
seen in cholinergic neurons innervating the prefrontal cortex [50–52]. With whole cell
electrophysiology, we measure opto-ACh responses in regular spiking pyramidal neurons in
layer 6 with voltage-clamp at -75 mV. Inward currents are resilient to combined application of
glutamatergic synaptic blockers CNQX (20µM) and D-APV (50µM), (peak amplitude, opto-ACh:
54.5 ±21.22pA, opto-ACh+synaptic blockers: 55.5 ±18.5pA; paired *t*-test, $t_6 = 0.2$, $P = 0.9$) and
are abolished by combined cholinergic receptor antagonists atropine (200nM) and DHβE (3µM),
(0 ±0% opto-ACh current remaining with cholinergic blockers; paired *t*-test, $t_{21} = 4.8$, $P = $
0.0001). This pharmacological assessment suggests opto-ACh currents are directly mediated
by cholinergic receptors on the recorded pyramidal neurons.

To explore age-related changes in this functional cholinergic signal, we record whole-cell
responses to opto-ACh across a broad age distribution in adulthood. We find a significant
decline in opto-ACh responses with age (**Fig 1C,D,E)**, as measured by peak amplitude of the
response (R^2 = 0.165; P = 0.0001, **Fig 1D** *inset*) and the area-integrated measure of charge
transfer (R^2 = 0.141; P = 0.0004, **Fig 1E** *inset*). Fitting a nonlinear robust regression to peak
amplitude and charge transfer reveals a fast decline in opto-ACh responses in early adulthood
(until ~P150) before responses plateau into middle adulthood and beyond (**Fig 1D, E**). With the
emergence of two distinct periods in this age range, we compare younger and older responses
using P150 as a cut-off point, revealing that the older response is ~50% lower than the younger
response in terms of peak amplitude (younger, 42.7 ±3.5pA; older, 22.6 ±2.8pA, unpaired *t*-test,
t_{183} = 4.5, P < 0.0001) and charge transfer (younger, 48.6 ±4.1pC; older, 27.8 ±2.8pC, unpaired
t-test, t_{183} = 4.2, P < 0.0001). From this grouped perspective, we find the decline in cholinergic
neurotransmission occurs in the absence of commensurate changes in postsynaptic neuron
intrinsic properties (**Supplemental Table S1**), although the older group showed a modest
increase in resting membrane potential (~2mV; unpaired *t*-test: t_{213} = 3, P = 0.004). Overall,
there is a net diminishment of the integrated response to opto-ACh across the age range
examined.

To probe components of the cholinergic synapse that may contribute to this decline in
cholinergic responses, we interrogate the functional role of the autoinhibitory muscarinic M2
receptor in the opto-ACh response. Located presynaptically, inhibitory M2 receptors modulate
cholinergic responses [53–55], restricting responses to opto-ACh at the prefrontal synapse [39].
We find that blocking muscarinic-mediated autoinhibition with muscarinic M2 receptor
antagonist AF-DX 116 (300nM) significantly increased opto-ACh responses peak amplitude
(paired *t*-test, t_{14} = 4.7, P = 0.0004) and charge transfer (paired *t*-test, t_{14} = 8, P < 0.0001) in
neurons across the age range. However, there were no significant age-related differences in the
increase in peak amplitude (unpaired *t*-test, t_{12} = 1.2, P = 0.3; **Fig 1F**) nor charge transfer (t_{12} =
0.3, P = 0.8; **Fig 1G**). These results suggest that muscarinic autoinhibition exerts a consistent
effect on synaptic cholinergic signaling across the age range examined.

To further examine synaptic components that exert control over cholinergic responses, we
consider the functional role of acetylcholinesterase in the opto-ACh response. Investigations of
acetylcholinesterase in pyramidal neurons of post-mortem human tissue show changing levels
in aging, without consensus on direction of change [22,56] or examination of the functional
impact of these changes. We show that blocking acetylcholinesterase with inhibitor galantamine
(1µM) significantly amplified and prolonged opto-ACh responses peak amplitude (paired *t*-test,
t_{16} = 3.7, P = 0.002) and charge transfer (paired *t*-test, t_{16} = 4.6, P = 0.0003) in neurons across
the age ranged examine. However, there were no significant age-related differences in increase
in peak amplitude (unpaired *t*-test, t_{15} = 0.5, P = 0.7; **Fig 1H**) nor charge transfer (unpaired *t*-
test, t_{15} = 0.8, P = 0.4; **Fig 1I**). These results suggest that acetylcholinesterase activity exerts a
consistent effect on synaptic cholinergic signaling across the age range examined.

Both presynaptic autoinhibition and acetylcholinesterase suppression at prefrontal cholinergic
synapses are intact in aging, indicating that the decline in opto-ACh responses may have a
postsynaptic origin.

Postsynaptic selectivity: age-related decline in nicotinic signal, stable muscarinic signal

To identify whether postsynaptic nicotinic and muscarinic components of opto-ACh responses
each show an age-dependent decline, we pharmacologically isolated these two
optophysiological signals in separate experiments. Our strategy was based on evidence that the
opto-ACh response is abolished by co-application of atropine and DHβE, as shown above and
in [39]. Therefore, we examined the isolated nicotinic component after muscarinic blockade with
atropine and the isolated muscarinic component after nicotinic blockade with DHβE.

The isolated nicotinic opto-ACh current differed significantly and substantially with age-group
(**Fig 2A,B**). The older group displayed ~30% of the younger nicotinic response in terms of both
peak amplitude (unpaired t-test, t_{27} = 3.3, P = 0.003; **Fig 2C**), and charge transfer (unpaired t-
test, t_{27} = 3.3, P = 0.003; **Fig 2D**). Under this condition, there was also a significant age-group
difference in rising slope (unpaired t-test, t_{27} = 3, P = 0.005; **Fig 2E**). In a subset of recordings,
the nicotinic identity of the residual optogenetic response was confirmed with application of
selective antagonist DHβE which abolished the residual current response (paired t-test, t_{21} =
4.8, P = 0.0001). The isolated nicotinic currents differ markedly by age-group.

By contrast, the isolated muscarinic opto-ACh currents do not differ by age-group (**Fig 2F,G**).
The older group displayed ~105% of the younger muscarinic response for peak amplitude
(unpaired t-test, t_{14} = 0.2, P = 0.7; **Fig 2H**) and ~95% for charge transfer (t_{14} = 0.2, P = 0.9; **Fig
2I**). The kinetics of the slower muscarinic response was better captured by 10-90 rise time,
which did not show a significant age-group difference (t_{14} = 1.6, P = 0.1; **Fig 2J**).

Since muscarinic receptors are state-dependent and exert stronger responses when the cell
is depolarized, near threshold, or firing action potentials [42,43], we also investigated the impact
of opto-ACh on postsynaptic neurons held in current clamp, with injection of current to elicit ~2-3
Hz action potential firing at baseline. Under these conditions, opto-ACh increased the frequency
of action potential firing (**Fig 2K**), and this response was insensitive to application of nicotinic
receptor blocker DHβE (peak: paired t-test, t_{10} = 0.6, P = 0.6; duration: paired t-test, t_{10} = 0.02, P
= 0.9; data not shown). Opto-ACh firing response did not differ significantly between younger
and older groups (peak: unpaired t-test, t_{10} = 0.3, P = 0.8; **Fig 2L**) duration: t_{10} = 0.3, P = 0.8;
Fig 2M). There was also no significant age-group difference in time to peak firing response (t_{10} =
1.8, P = 0.1, **Fig 2N**). These results in current-clamp are consistent with those in voltage-clamp,
confirming that the optogenetic muscarinic response is preserved across age groups.

These results support the conclusion that a change in postsynaptic nicotinic receptors, not in
muscarinic receptors, is predominantly responsible for the age-dependent decline in opto-ACh
responses.

Treatment impact of nicotinic positive allosteric modulation diminishes with age

Towards restoring a strong nicotinic signal, we apply a nicotinic positive allosteric modulator
NS9283 [57,58] known to improve cognition in young animals [59,57]. The success of this
treatment in aging animals will depend on there being sufficient postsynaptic nicotinic receptors
available for potentiation. We find that NS9283 (1µM) strongly potentiates younger opto-ACh
responses, but this potentiation is less robust in older responses (**Fig 3A, B**) with a significant
interaction of NS9283 and age on peak amplitude of opto-ACh (repeated measures two-way
ANOVA, Interaction: $F_{1,24}$ = 4.9, P = 0.04; **Fig 3C**). Sidak *post hoc* tests confirm significant
NS9283 potentiation in the younger group (t_{14} = 5.1, P < 0.0001) that is not observed in the
older group (t_{13} = 2, P = 0.1). Considering the NS9283 effect on opto-ACh charge transfer, there
are significant main effects of age and drug (Two-way ANOVA, age: $F_{1,24}$ = 9.2, P = 0.005;

nicotinic PAM: $F_{1,24}$ = 22.3, P < 0.0001; **Fig 3D**). Again, Sidak *post hoc* tests show NS9283
potentiation of opto-ACh charge transfer in the younger group (t_{14} = 4.7, P = 0.0001) and not the
older group (t_{13} = 2, P = 0.1). Improving nicotinic receptor function with NS9283 is effective in
young adults but this efficacy diminishes in older adults. The decrease in effectiveness of the
nicotinic PAM with age points toward a loss of nicotinic receptor availability. Therefore, we next
investigate a candidate mechanism to improve the available population of nicotinic receptors.

Harnessing muscarinic receptor PKC signaling to improve nicotinic neurotransmission

The signaling pathways activated by muscarinic receptors [60–62] have independently been
shown to improve nicotinic receptor insertion in the cell membrane [63,64]. To enhance nicotinic
receptor availability, we therefore pursue an unorthodox strategy of stimulating muscarinic
receptors to activate their secondary messenger pathways. We hypothesize that activating
intact M1 muscarinic receptor signaling with direct agonist xanomeline [65] will upregulate
postsynaptic nicotinic receptor availability in older mice (**Fig 4A**).

In prefrontal brain slices from older mice (postnatal 210 ±5days), application of xanomeline
(300nM, 10 min) significantly increased opto-ACh responses (peak amplitude: paired *t*-test, t_{14} =
3.2, P = 0.006; charge transfer: t_{14} = 3, P = 0.01, (**Fig 4B**) with negligible impact on baseline
holding current (-0.8 ±1pA). Following xanomeline treatment, we found opto-ACh responses
continued to increase, with most neurons reaching a peak response 10-20 minutes post-
xanomeline. Over time, xanomeline yielded a greater significant increase in opto-ACh peak
amplitude (paired *t*-test: t_{14} = 3.9, P = 0.002; **Fig 4C**) and charge transfer (paired two-tailed *t*-
test: t_{14} = 4.7, P = 0.0003; **Fig 4D**). In the absence of xanomeline, control experiments recorded
over the same time course did not show potentiation of opto-ACh response (control change in
peak with time: paired *t*-test, t_{14} = 0.6, P = 0.6; control change in charge transfer with time: t_{14} =
0.3, P = 0.8; data not shown). Consistent with our hypothesis that xanomeline boosts opto-ACh
response by increasing the nicotinic component, nicotinic receptor antagonist DHβE eliminated
the potentiated opto-ACh response (**Fig 4E**), significantly decreasing peak amplitude (paired *t*-
test, t_9 = 6.6, P = 0.0001; **Fig 4E** *inset*) and charge transfer (t_9 = 4, P = 0.003, data not shown).
These results demonstrate that xanomeline enhances the optogenetic nicotinic response in
older mice.

To investigate the mechanisms underlying xanomeline-potentiation of the nicotinic
component, we contrast the effects of xanomeline alone and after pathway manipulations in
brain slices from the same mice (**Fig 4G,H**). We found that muscarinic receptor blockade with
atropine eliminated xanomeline's potentiation of opto-ACh responses, consistent with a
requirement for activation of muscarinic receptors. Likewise, inhibition of PKC with either
chelerythrine (1µM) or Go6983 (300nM) also severely disrupted xanomeline potentiation of
opto-ACh responses, consistent with dependence on PKC. Analyzing these experiments
together, one-way ANOVA illustrates that either manipulation is sufficient to disrupt xanomeline
potentiation of peak amplitude ($F_{2,14}$ = 7.3, P = 0.007; Dunnett's multiple comparisons, xano vs
+AT: q_{14} = 3.1, P = 0.02; xano vs +PKC inhib: q_{14} = 3.6, P = 0.006; **Fig 4G,H**). Together, these
experiments illustrate that xanomeline-elicited potentiation requires downstream activation of
PKC by muscarinic receptors, increasing the nicotinic component of the opto-ACh response in
older adult mice.

Full restoration with combined treatment of xanomeline and NS9283

To examine the sensitivity of the restored nicotinic response to further amplification, we applied
xanomeline followed by NS9283 in brain slices from an additional group of older mice (postnatal
332 ±3days). To achieve a similar time-course to previous experiments with NS9283 (**Fig 3**),
slices were treated with xanomeline then neurons were patched for paired recordings before
and after application of NS9283 (**Fig 5A**). Application of nicotinic PAM after xanomeline yielded
a highly significant increase in peak amplitude (paired *t*-test, $t_{11} = 5.3$, $P = 0.0003$; **Fig 5B**),
charge transfer (paired *t*-test, $t_{11} = 4.7$, $P = 0.0007$; **Fig 5C**) and rise speed (paired *t*-test, $t_{11} =$
5.7, $P = 0.0001$; **Fig 5D**). This striking potentiation of the older opto-ACh response by NS9283
further supports the conclusion that xanomeline increases postsynaptic nicotinic receptor
availability.

Next, we normalized opto-ACh responses from this group of much older mice to the baseline
mean of the younger mice (**Fig 5E**). This normalization illustrates the replication of the decline in
opto-ACh response with age. It also demonstrates the ability of combination treatment of
xanomeline with NS9283 to restore older responses into the typical range of young adults (one-
way ANOVA, $F_{2,41} = 16.49$, $P < 0.0001$; Tukey's *post hoc*: older vs xanomeline+NS9283: $q_{28} = 8$,
$P < 0.0001$). Even in year-old mice, the deficit in opto-ACh signal can be restored to younger
levels via combined treatment of xanomeline and NS9283 (**Fig 5F,G**).

Overall, we present a novel strategy to restore nicotinic receptor neurophysiology in aging
informed by careful functional dissection of pre- and post-synaptic components of prefrontal
cholinergic synapses.

Discussion

Optogenetic activation of prefrontal cholinergic synapses reveals a significant decline in
signaling during adulthood. Across the age range examined, multiple aspects of cholinergic
neurotransmission are stable, with the critical exception of the decline in postsynaptic nicotinic
receptor signal. While direct potentiation of nicotinic receptors with positive allosteric modulator
NS9283 improves nicotinic responses in young adults, it imparts minimal benefit in older mice.
We demonstrate that harnessing preserved muscarinic signaling via xanomeline upregulates
nicotinic receptor availability in prefrontal cortex, restoring sensitivity to allosteric modulation.
Combining xanomeline with nicotinic allosteric potentiation rescues the speed, strength, and
receptor mechanism of endogenous cholinergic responses to younger levels.

How age impacts pre- and post-synaptic aspects of the prefrontal cholinergic system

The net effect of aging on cholinergic neurons and cortical signaling has been a subject of
considerable debate. Multiple, sometimes conflicting, alterations in pre-and post-synaptic
cholinergic components have been described during adulthood [40,66–73], leaving a knowledge
gap in their integrated functional consequences. Here, we identify an age-related selective
postsynaptic nicotinic receptor deficit in functional prefrontal cholinergic neurotransmission. This
finding is consistent with earlier preclinical research showing age-dependent cortical decreases
in nicotinic receptor binding [40,69,71,74] and in response to exogenous agonist [41]. A parallel
decline in cortical nicotinic receptor binding is observed in human aging [17,18,75],
underscoring the translational relevance of this work.

In contrast to the declining nicotinic signal, we found optogenetically-evoked postsynaptic muscarinic signaling remained stable across the age range studied. This stability was observed for two electrophysiological measures mediated by distinct cellular mechanisms [43,76–78], shedding new light on earlier research [66,79–82]. In contrast to prior reports at the whole tissue level [67,70,72,73], regulation of prefrontal cholinergic synapses by autoinhibition and acetylcholinesterase remain similar between age groups. These age-stable measures, together with successful rescue of nicotinic signaling, suggest presynaptic acetylcholine remains intact. In prefrontal cortex of healthy mice, there appears to be steady maintenance of trophic support mechanisms for cholinergic afferents over the age range we examined [83]. In healthy human aging, there are only gradual changes in cholinergic neurons and axons [84–86], with large or rapid declines in cholinergic afferents increasingly viewed as the domain of pathological aging [83,87,88].

Implications for interventions to improve cholinergic signaling

Progressive decline in certain aspects of cognition during adulthood has been documented clinically [1–6] and preclinically [33–35]. Due to the importance of cholinergic signaling for executive function [7,8,10], acetylcholine receptors have been targeted extensively by therapeutic manipulations for cognitive improvement [23–29]. Our results indicate that improving nicotinic receptor signaling may allow for cognitive restoration. Direct stimulation of nicotinic receptors is complicated by desensitization [89–91]. Nicotinic receptor positive allosteric modulators, such as NS9283, present less risk of desensitization [92] and are pro-cognitive in young animals [57,59]. Our results (**Fig 3**) suggest the cognitive success of allosteric modulation in older subjects would be constrained and (**Fig 4**) highlight a strategy of first boosting nicotinic receptor availability via intracellular transduction pathways.

In cell systems, PKC upregulates nicotinic receptor trafficking [63,64] and phosphorylation [93]. This important kinase is activated by excitatory muscarinic receptors [60–62] and optophysiological experiments reveal proximity of muscarinic receptors to prefrontal cholinergic synapses (**Fig 2**; [36,39,78]. We reveal that activating muscarinic signalling via direct agonist xanomeline increases functional nicotinic responses in a PKC-dependent manner (**Fig 4**). This is consistent with a previous report that stimulating PKC through a different $G\alpha_q$-coupled receptor increased electrically-evoked nicotinic responses without altering the probability of presynaptic acetylcholine release [94]. Here, we find xanomeline-upregulated nicotinic receptors restored youthful sensitivity of prefrontal cholinergic synapses to allosteric potentiation (**Fig 5**).

Caveats and future research directions

Tools to examine cholinergic neurotransmission are not without limitations. Here, we discuss caveats for choice of stimulus pattern and model selection. We employed optogenetic stimuli in a descending frequency pattern to mimic firing properties of the major subgroup of cortically-projecting cholinergic neurons [50–52]. Relatively little is known, however, about how these firing patterns change in adulthood, as previous work has focused predominantly on younger animals [50–52]. There is some evidence, however, that cholinergic firing patters do not change substantially with age from juvenile to young adult [95]. Complications in cholinergic stimulation include limitations in achieving opsin expression in cholinergic neurons. The two major strategies in the field are suspected to alter cholinergic release in different directions (increased [96]; decreased [97]) and are associated with cognitive and behavioural changes from wildtype [96,97]. In previous electrophysiological work, we have examined the different models and

found many similarities in their prefrontal cholinergic synapses in autoinhibition, metabolism,
and postsynaptic receptor signaling [39], extending work from other groups [36,78]. Here, our
finding of age-related functional decline in nicotinic responses is consistent with work in wildtype
rodents showing decreased nicotinic binding and diminished responses to exogenous agonist
[40]. Furthermore, current discovery that xanomeline upregulates postsynaptic nicotinic
signaling in a PKC dependent manner matches the consequences of electrical activation of a
different cholinergic circuit in wildtype mice [94].

There are also caveats to consider in receptor and second messenger pathway specificity.
Our results give new insight into a previously-unknown mechanism underlying xanomeline-
mediated cognitive enhancement. The atropine-sensitivity and PKC-dependence of its nicotinic
receptor upregulation points to an excitatory muscarinic mechanism, but xanomeline has also
been shown to inhibit M4 receptors [98,99] and could potentially disinhibit glutamate release.
Our data implicate the involvement of PKC, typically activated by $G\alpha_q$-signaling, but this
signaling also triggers activation of PLC [100] and may regulate GPI-anchored lynx family
proteins [101] that also modulate nicotinic receptors [102,103] directly [104–106] or indirectly
[105,107]. While the complex action of pharmacological intervention explored here warrants
further investigation, our results demonstrate, for the first time, that muscarinic signaling can be
harnessed to restore deficits in postsynaptic nicotinic receptors.

Clinical relevance

Muscarinic agonism as a treatment for cognitive impairment has long been of interest. The
direct M1 agonist xanomeline was first found to reduce cognitive impairments in Alzheimer's
disease in clinical trials 25 years ago, but presented adverse peripheral effects of muscarinic
agonism [27,108]. Recently, xanomeline together with a peripheral blocker has been revisited in
clinical trial, showing improved cognition in schizophrenia with less-severe adverse effects
[109]. Another muscarinic agonist has also recently been shown to be pro-cognitive in pre-
clinical work, tolerable in clinical trial, and improve human fMRI signals of hippocampal
activation during spatial cue encoding [110]. These investigations highlight the importance of
more fully understanding the mechanisms by which muscarinic signaling improves cognition.

Summary of findings and implications

Our work investigates, for the first time, the functional consequences of aging on prefrontal
cholinergic neurotransmission. We identify age-dependent functional deficits in postsynaptic
nicotinic receptor activation. Using xanomeline to improve nicotinic signaling takes advantage of
well-preserved muscarinic receptors and their second messenger pathways. Combining this
treatment with nicotinic allosteric potentiation rescues endogenous nicotinic signaling. This
reveals a novel mechanism for improving cognition through muscarinic receptors, taking
advantage of a pharmacological agent already well investigated in clinical trials.

Acknowledgments

We greatly appreciate the insight and feedback on earlier versions of this manuscript from Dr.
Peter Carlen, Dr. JoAnne McLaurin, and Dr. Amy Ramsey of the University of Toronto.

Author Contributions

Conceptualization, SKP, EKL; Methodology, SKP, SV, EKL; Data acquisition SKP, Analysis, SKP, SV, EKL; Writing, SKP EKL; Review & editing, SKP, SV, EKL; Funding, SKP, SV, EKL; Supervision, EKL. All authors have critically reviewed and approved of the manuscript.

Funding

This research was generously supported by the Canadian Institutes of Health Research (CIHR; MOP89825 and CIHR PJT-153101; EKL), the Canada Research Chair in Developmental Cortical Physiology (EKL), a Banting and Best Doctoral Canada Graduate Scholarship (SKP), and Ontario Graduate Scholarships (SKP, SV).

Competing interests

The authors have nothing to disclose.

References

1. Filley CM, Cullum CM. Attention and vigilance functions in normal aging. Appl Neuropsychol. 1994;1:29–32.
2. Mani TM, Bedwell JS, Miller LS. Age-related decrements in performance on a brief continuous performance test. Arch Clin Neuropsychol. 2005;20:575–586.
3. Gamboz N, Zamarian S, Cavallero C. Age-related differences in the attention network test (ANT). Exp Aging Res. 2010;36:287–305.
4. Kaufman DAS, Sozda CN, Dotson VM, Perlstein WM. An Event-Related Potential Investigation of the Effects of Age on Alerting, Orienting, and Executive Function. Front Aging Neurosci. 2016;8:99.
5. Zhou S, Fan J, Lee TMC, Wang C, Wang K. Age-related differences in attentional networks of alerting and executive control in young, middle-aged, and older Chinese adults. Brain Cogn. 2011;75:205–210.
6. Hedden T, Van Dijk KRA, Shire EH, Sperling RA, Johnson KA, Buckner RL. Failure to Modulate Attentional Control in Advanced Aging Linked to White Matter Pathology. Cereb Cortex. 2012;22:1038–1051.
7. Dalley JW, Theobald DE, Bouger P, Chudasama Y, Cardinal RN, Robbins TW. Cortical cholinergic function and deficits in visual attentional performance in rats following 192 IgG-saporin-induced lesions of the medial prefrontal cortex. Cereb Cortex N Y N 1991. 2004;14:922–932.
8. Passetti F, Dalley JW, O'Connell MT, Everitt BJ, Robbins TW. Increased acetylcholine release in the rat medial prefrontal cortex during performance of a visual attentional task. Eur J Neurosci. 2000;12:3051–3058.
9. Parikh V, Kozak R, Martinez V, Sarter M. Prefrontal acetylcholine release controls cue detection on multiple time scales. Neuron. 2007;56:141–154.
10. Gritton HJ, Howe WM, Mallory CS, Hetrick VL, Berke JD, Sarter M. Cortical cholinergic signaling controls the detection of cues. Proc Natl Acad Sci. 2016;113:E1089–E1097.
11. Howe WM, Gritton HJ, Lusk NA, Roberts EA, Hetrick VL, Berke JD, Sarter M. Acetylcholine Release in Prefrontal Cortex Promotes Gamma Oscillations and Theta–Gamma Coupling during Cue Detection. J Neurosci. 2017;37:3215–3230.
12. Bartus RT, Dean RL, Beer B, Lippa AS. The Cholinergic Hypothesis of Geriatric Memory Dysfunction. Science. 1982;217:408–417.
13. Dumas JA, Newhouse PA. The Cholinergic Hypothesis of Cognitive Aging Revisited Again: Cholinergic Functional Compensation. Pharmacol Biochem Behav. 2011;99:254–261.

406   14.   Gallagher M, Colombo PJ. Ageing: the cholinergic hypothesis of cognitive decline. Curr Opin
407   Neurobiol. 1995;5:161–168.
408   15.   Hanyu H, Asano T, Sakurai H, Tanaka Y, Takasaki M, Abe K. MR analysis of the substantia
409   innominata in normal aging, Alzheimer disease, and other types of dementia. AJNR Am J
410   Neuroradiol. 2002;23:27–32.
411   16.   Ikonomovic MD, Abrahamson EE, Isanski BA, Wuu J, Mufson EJ, DeKosky ST. Superior frontal
412   cortex cholinergic axon density in mild cognitive impairment and early Alzheimer disease. Arch
413   Neurol. 2007;64:1312–1317.
414   17.   Mitsis EM, Cosgrove KP, Staley JK, Bois F, Frohlich EB, Tamagnan GD, Estok KM, Seibyl JP, van
415   Dyck CH. Age-related decline in nicotinic receptor availability with [123I]5-IA-85380 SPECT.
416   Neurobiol Aging. 2009;30:1490–1497.
417   18.   Kendziorra K, Wolf H, Meyer PM, Barthel H, Hesse S, Becker GA, Luthardt J, Schildan A, Patt M,
418   Sorger D, Seese A, Gertz H-J, Sabri O. Decreased cerebral α4β2* nicotinic acetylcholine receptor
419   availability in patients with mild cognitive impairment and Alzheimer's disease assessed with
420   positron emission tomography. Eur J Nucl Med Mol Imaging. 2011;38:515–525.
421   19.   Dewey SL, Volkow ND, Logan J, MacGregor RR, Fowler JS, Schlyer DJ, Bendriem B. Age-related
422   decreases in muscarinic cholinergic receptor binding in the human brain measured with positron
423   emission tomography (PET). J Neurosci Res. 1990;27:569–575.
424   20.   Pilar-Cuéllar F, Paniagua MA, Díez-Alarcia R, Dos Anjos S, Montori S, Pérez CC, Fernández-
425   López A. Muscarinic receptor changes in the gerbil thalamus during aging. Brain Res.
426   2008;1243:38–46.
427   21.   Terry AV, Buccafusco JJ. The cholinergic hypothesis of age and Alzheimer's disease-related
428   cognitive deficits: recent challenges and their implications for novel drug development. J
429   Pharmacol Exp Ther. 2003;306:821–827.
430   22.   Janeczek M, Gefen T, Samimi M, Kim G, Weintraub S, Bigio E, Rogalski E, Mesulam M-M, Geula
431   C. Variations in Acetylcholinesterase Activity within Human Cortical Pyramidal Neurons Across
432   Age and Cognitive Trajectories. Cereb Cortex N Y N 1991. 2018;28:1329–1337.
433   23.   Poltavski DV, Petros T. Effects of transdermal nicotine on attention in adult non-smokers with and
434   without attentional deficits. Physiol Behav. 2006;87:614–624.
435   24.   Howe WM, Ji J, Parikh V, Williams S, Mocaër E, Trocmé-Thibierge C, Sarter M. Enhancement of
436   attentional performance by selective stimulation of alpha4beta2(*) nAChRs: underlying cholinergic
437   mechanisms. Neuropsychopharmacol. 2010;35:1391–1401.
438   25.   Hong LE, Thaker GK, McMahon RP, Summerfelt A, Rachbeisel J, Fuller RL, Wonodi I, Buchanan
439   RW, Myers C, Heishman SJ, Yang J, Nye A. Effects of moderate-dose treatment with varenicline
440   on neurobiological and cognitive biomarkers in smokers and nonsmokers with schizophrenia or
441   schizoaffective disorder. Arch Gen Psychiatry. 2011;68:1195–1206.
442   26.   Preskorn SH, Gawryl M, Dgetluck N, Palfreyman M, Bauer LO, Hilt DC. Normalizing effects of
443   EVP-6124, an α-7 nicotinic partial agonist, on event-related potentials and cognition: a proof of
444   concept, randomized trial in patients with schizophrenia. J Psychiatr Pract. 2014;20:12–24.
445   27.   Bodick NC, Offen WW, Levey AI, Cutler NR, Gauthier SG, Satlin A, Shannon HE, Tollefson GD,
446   Rasmussen K, Bymaster FP, Hurley DJ, Potter WZ, Paul SM. Effects of xanomeline, a selective
447   muscarinic receptor agonist, on cognitive function and behavioral symptoms in Alzheimer disease.
448   Arch Neurol. 1997;54:465–473.
449   28.   Shekhar A, Potter WZ, Lightfoot J, Lienemann J, Dubé S, Mallinckrodt C, Bymaster FP, McKinzie
450   DL, Felder CC. Selective muscarinic receptor agonist xanomeline as a novel treatment approach
451   for schizophrenia. Am J Psychiatry. 2008;165:1033–1039.
452   29.   Brannan SK, Sawchak S, Miller AC, Lieberman JA, Paul SM, Breier A. Muscarinic Cholinergic
453   Receptor Agonist and Peripheral Antagonist for Schizophrenia. N Engl J Med. 2021;384:717–726.
454   30.   Salloway S, Ferris S, Kluger A, Goldman R, Griesing T, Kumar D, Richardson S, Donepezil 401
455   Study Group. Efficacy of donepezil in mild cognitive impairment: a randomized placebo-controlled
456   trial. Neurology. 2004;63:651–657.
457   31.   Doody RS, Ferris SH, Salloway S, Sun Y, Goldman R, Watkins WE, Xu Y, Murthy AK. Donepezil
458   treatment of patients with MCI: a 48-week randomized, placebo-controlled trial. Neurology.
459   2009;72:1555–1561.

32.   Fang X, Zhang J, Zhao J, Wang L. Effect of Resveratrol Combined with Donepezil Hydrochloride
on Inflammatory Factor Level and Cognitive Function Level of Patients with Alzheimer's Disease. J
Healthc Eng. 2022;2022:9148650.
33.   Palliyaguru DL, Vieira Ligo Teixeira C, Duregon E, di Germanio C, Alfaras I, Mitchell SJ, Navas-
Enamorado I, Shiroma EJ, Studenski S, Bernier M, Camandola S, Price NL, Ferrucci L, de Cabo
R, SLAM Investigators. Study of Longitudinal Aging in Mice: Presentation of Experimental
Techniques. J Gerontol Ser A. 2021;76:552–560.
34.   Yanai S, Endo S. Functional Aging in Male C57BL/6J Mice Across the Life-Span: A Systematic
Behavioral Analysis of Motor, Emotional, and Memory Function to Define an Aging Phenotype.
Front Aging Neurosci. 2021;13.
35.   Shoji H, Takao K, Hattori S, Miyakawa T. Age-related changes in behavior in C57BL/6J mice from
young adulthood to middle age. Mol Brain. 2016;9:11.
36.   Hedrick T, Waters J. Acetylcholine excites neocortical pyramidal neurons via nicotinic receptors. J
Neurophysiol. 2015;113:2195–2209.
37.   Hay YA, Lambolez B, Tricoire L. Nicotinic Transmission onto Layer 6 Cortical Neurons Relies on
Synaptic Activation of Non-α7 Receptors. Cereb Cortex N Y N 1991. 2016;26:2549–2562.
38.   Sparks DW, Tian MK, Sargin D, Venkatesan S, Intson K, Lambe EK. Opposing Cholinergic and
Serotonergic Modulation of Layer 6 in Prefrontal Cortex. Front Neural Circuits. 2017;11:107.
39.   Venkatesan S, Lambe EK. Chrna5 is Essential for a Rapid and Protected Response to
Optogenetic Release of Endogenous Acetylcholine in Prefrontal Cortex. J Neurosci Off J Soc
Neurosci. 2020;40:7255–7268.
40.   Ghimire M, Cai R, Ling L, Hackett TA, Caspary DM. Nicotinic Receptor Subunit Distribution in
Auditory Cortex: Impact of Aging on Receptor Number and Function. J Neurosci. 2020;40:5724–
5739.
41.   Patel AV, Codeluppi SA, Ervin KSJ, St-Denis MB, Choleris E, Bailey CDC. Developmental Age
and Biological Sex Influence Muscarinic Receptor Function and Neuron Morphology within Layer
VI of the Medial Prefrontal Cortex. Cereb Cortex. 2021:bhab406.
42.   Gulledge AT, Park SB, Kawaguchi Y, Stuart GJ. Heterogeneity of phasic cholinergic signaling in
neocortical neurons. J Neurophysiol. 2007;97:2215–2229.
43.   Tian MK, Bailey CDC, De Biasi M, Picciotto MR, Lambe EK. Plasticity of prefrontal attention
circuitry: upregulated muscarinic excitability in response to decreased nicotinic signaling following
deletion of α5 or β2 subunits. J Neurosci. 2011;31:16458–16463.
44.   Tian MK, Bailey CDC, Lambe EK. Cholinergic excitation in mouse primary vs. associative cortex:
region-specific magnitude and receptor balance. Eur J Neurosci. 2014;40:2608–2618.
45.   Zhao S, Ting JT, Atallah HE, Qiu L, Tan J, Gloss B, Augustine GJ, Deisseroth K, Luo M, Graybiel
AM, Feng G. Cell-type Specific Optogenetic Mice for Dissecting Neural Circuitry Function. Nat
Methods. 2011;8:745–752.
46.   Nasirova N, Quina LA, Agosto-Marlin IM, Ramirez J-M, Lambe EK, Turner EE. Dual recombinase
fate mapping reveals a transient cholinergic phenotype in multiple populations of developing
glutamatergic neurons. J Comp Neurol. 2020;528:283–307.
47.   Tassonyi E, Charpantier E, Muller D, Dumont L, Bertrand D. The role of nicotinic acetylcholine
receptors in the mechanisms of anesthesia. Brain Res Bull. 2002;57:133–150.
48.   Solt K, Kelly EW, Cotten JF, Raines DE. Inhibition of human alpha4beta2 neuronal nicotinic
acetylcholine receptors by volatile aromatic anesthetics depends on drug hydrophobicity. Anesth
Analg. 2010;110:455–460.
49.   Zhong J, Li J, Miao C, Zuo Z. A Novel Individual-based Determination of Postoperative Cognitive
Dysfunction in Mice. Aging Dis. 2020;11:1133–1145.
50.   Laszlovszky T, Schlingloff D, Hegedüs P, Freund TF, Gulyás A, Kepecs A, Hangya B. Distinct
synchronization, cortical coupling and behavioral function of two basal forebrain cholinergic neuron
types. Nat Neurosci. 2020;23:992–1003.
51.   Lee MG, Hassani OK, Alonso A, Jones BE. Cholinergic Basal Forebrain Neurons Burst with Theta
during Waking and Paradoxical Sleep. J Neurosci. 2005;25:4365–4369.
52.   Unal CT, Golowasch JP, Zaborszky L. Adult mouse basal forebrain harbors two distinct cholinergic
populations defined by their electrophysiology. Front Behav Neurosci. 2012;6:21.
53.   Brown. Muscarinic acetylcholine receptors (mAChRs) in the nervous system: some functions and
mechanisms. J Mol Neurosci MN. 2010;41:340–346.

54. Iannazzo L, Majewski H. M(2)/M(4)-muscarinic receptors mediate automodulation of acetylcholine
outflow from mouse cortex. Neurosci Lett. 2000;287:129–132.
55. Zhang W, Basile AS, Gomeza J, Volpicelli LA, Levey AI, Wess J. Characterization of central
inhibitory muscarinic autoreceptors by the use of muscarinic acetylcholine receptor knock-out
mice. J Neurosci. 2002;22:1709–1717.
56. Bu J. Age-related changes in calbindin-D28k, calretinin, and parvalbumin-immunoreactive neurons
in the human cerebral cortex. Exp Neurol. 2003;182:220–231.
57. Grupe M, Grunnet M, Laursen B, Bastlund JF. Neuropharmacological modulation of the P3-like
event-related potential in a rat two-tone auditory discrimination task with modafinil and NS9283, a
positive allosteric modulator of α4β2 nAChRs. Neuropharmacology. 2014;79:444–455.
58. Olsen JA, Ahring PK, Kastrup JS, Gajhede M, Balle T. Structural and Functional Studies of the
Modulator NS9283 Reveal Agonist-like Mechanism of Action at α4β2 Nicotinic Acetylcholine
Receptors. J Biol Chem. 2014;289:24911–24921.
59. Timmermann D, Sandager-Nielsen K, Dyhring T, Smith M, Jacobsen A-M, Nielsen E, Grunnet M,
Christensen J, Peters D, Kohlhaas K, Olsen G, Ahring P. Augmentation of cognitive function by
NS9283, a stoichiometry-dependent positive allosteric modulator of α2- and α4-containing nicotinic
acetylcholine receptors. Br J Pharmacol. 2012;167:164–182.
60. Caccamo A, Oddo S, Billings LM, Green KN, Martinez-Coria H, Fisher A, LaFerla FM. M1
Receptors Play a Central Role in Modulating AD-like Pathology in Transgenic Mice. Neuron.
2006;49:671–682.
61. Huang C-C, Hsu K-S. Activation of muscarinic acetylcholine receptors induces a nitric oxide-
dependent long-term depression in rat medial prefrontal cortex. Cereb Cortex N Y N 1991.
2010;20:982–996.
62. Matsuoka H, Harada K, Mashima K, Inoue M. Muscarinic receptor stimulation induces TASK1
channel endocytosis through a PKC-Pyk2-Src pathway in PC12 cells. Cell Signal.
2020;65:109434.
63. Gopalakrishnan M, Molinari EJ, Sullivan JP. Regulation of human alpha4beta2 neuronal nicotinic
acetylcholine receptors by cholinergic channel ligands and second messenger pathways. Mol
Pharmacol. 1997;52:524–534.
64. Nashmi R, Dickinson ME, McKinney S, Jareb M, Labarca C, Fraser SE, Lester HA. Assembly of
alpha4beta2 nicotinic acetylcholine receptors assessed with functional fluorescently labeled
subunits: effects of localization, trafficking, and nicotine-induced upregulation in clonal mammalian
cells and in cultured midbrain neurons. J Neurosci. 2003;23:11554–11567.
65. Wood MD, Murkitt KL, Ho M, Watson JM, Brown F, Hunter AJ, Middlemiss DN. Functional
comparison of muscarinic partial agonists at muscarinic receptor subtypes hM1, hM2, hM3, hM4
and hM5 using microphysiometry. Br J Pharmacol. 1999;126:1620–1624.
66. Biegon A, Hanau M, Greenberger V, Segal M. Aging and brain cholinergic muscarinic receptor
subtypes: an autoradiographic study in the rat. Neurobiol Aging. 1989;10:305–310.
67. Araujo DM, Lapchak PA, Meaney MJ, Collier B, Quirion R. Effects of aging on nicotinic and
muscarinic autoreceptor function in the rat brain: relationship to presynaptic cholinergic markers
and binding sites. J Neurosci. 1990;10:3069–3078.
68. Mitsushima D, Mizuno T, Kimura F. Age-related changes in diurnal acetylcholine release in the
prefrontal cortex of male rats as measured by microdialysis. Neuroscience. 1996;72:429–434.
69. Ferrari R, Pedrazzi P, Algeri S, Agnati LF, Zoli M. Subunit and region-specific decreases in
nicotinic acetylcholine receptor mRNA in the aged rat brain. Neurobiol Aging. 1999;20:37–46.
70. Das A, Dikshit M, Nath C. Profile of acetylcholinesterase in brain areas of male and female rats of
adult and old age. Life Sci. 2001;68:1545–1555.
71. Tribollet E, Bertrand D, Marguerat A, Raggenbass M. Comparative distribution of nicotinic receptor
subtypes during development, adulthood and aging: an autoradiographic study in the rat brain.
Neuroscience. 2004;124:405–420.
72. Décossas M, Doudnikoff É, Bloch B, Bernard V. Aging and subcellular localization of m2
muscarinic autoreceptor in basalocortical neurons in vivo. Neurobiol Aging. 2005;26:1061–1072.
73. Strauss V, Rey Moreno MC, Vogt J, Dammann M, Schneider S, Gröters S, van Ravenzwaay B.
Acetylcholinesterase measurement in various brain regions and muscles of juvenile, adolescent,
and adult rats. Toxicol Mech Methods. 2017;27:666–676.

74. Rogers SW, Gahring LC, Collins AC, Marks M. Age-Related Changes in Neuronal Nicotinic
Acetylcholine Receptor Subunit α4 Expression Are Modified by Long-Term Nicotine
Administration. J Neurosci. 1998;18:4825–4832.
75. Marutle A, Warpman U, Bogdanovic N, Nordberg A. Regional distribution of subtypes of nicotinic
receptors in human brain and effect of aging studied by (+/-)-[3H]epibatidine. Brain Res.
1998;801:143–149.
76. Gulledge AT, Bucci DJ, Zhang SS, Matsui M, Yeh HH. M1 Receptors Mediate Cholinergic
Modulation of Excitability in Neocortical Pyramidal Neurons. J Neurosci. 2009;29:9888–9902.
77. Proulx É, Suri D, Heximer SP, Vaidya VA, Lambe EK. Early stress prevents the potentiation of
muscarinic excitation by calcium release in adult prefrontal cortex. Biol Psychiatry. 2014;76:315–
323.
78. Baker AL, O'Toole RJ, Gulledge AT. Preferential cholinergic excitation of corticopontine neurons. J
Physiol. 2018;596:1659–1679.
79. James TC, Kanungo MS. Alterations in atropine sites of the brain of rats as a function of age.
Biochem Biophys Res Commun. 1976;72:170–175.
80. Morin AM, Wasterlain CG. Aging and rat brain muscarinic receptors as measured by quinuclidinyl
benzilate binding. Neurochem Res. 1980;5:301–308.
81. Waller SB, London ED. Age-differences in choline acetyltransferase activities and muscarinic
receptor binding in brain regions of C57BL/6J mice. Exp Gerontol. 1983;18:419–425.
82. Pedigo NW, Minor LD, Krumrei TN. Cholinergic drug effects and brain muscarinic receptor binding
in aged rats. Neurobiol Aging. 1984;5:227–233.
83. Perez SE, He B, Muhammad N, Oh K-J, Fahnestock M, Ikonomovic MD, Mufson EJ.
Cholinotrophic basal forebrain system alterations in 3xTg-AD transgenic mice. Neurobiol Dis.
2011;41:338–352.
84. Geula C, Nagykery N, Nicholas A, Wu C-K. Cholinergic neuronal and axonal abnormalities are
present early in aging and in Alzheimer disease. J Neuropathol Exp Neurol. 2008;67:309–318.
85. Wolf D, Grothe M, Fischer FU, Heinsen H, Kilimann I, Teipel S, Fellgiebel A. Association of basal
forebrain volumes and cognition in normal aging. Neuropsychologia. 2014;53:54–63.
86. Peter J, Mayer I, Kammer T, Minkova L, Lahr J, Klöppel S, Grothe MJ, Orth M. The relationship
between cholinergic system brain structure and function in healthy adults and patients with mild
cognitive impairment. Sci Rep. 2021;11:16080.
87. Wu H, Williams J, Nathans J. Complete morphologies of basal forebrain cholinergic neurons in the
mouse. ELife. 2014;3.
88. Fernández-Cabello S, Kronbichler M, Van Dijk KRA, Goodman JA, Spreng RN, Schmitz TW.
Basal forebrain volume reliably predicts the cortical spread of Alzheimer's degeneration. Brain.
2020;143:993–1009.
89. Wang H, Sun X. Desensitized nicotinic receptors in brain. Brain Res Brain Res Rev. 2005;48:420–
437.
90. Picciotto MR, Addy NA, Mineur YS, Brunzell DH. It's not "either/or": activation and desensitization
of nicotinic acetylcholine receptors both contribute to behaviors related to nicotine addiction and
mood. Prog Neurobiol. 2008;84:329–342.
91. Henderson BJ, Lester HA. Inside-out neuropharmacology of nicotinic drugs. Neuropharmacology.
2015;96:178–193.
92. Williams DK, Wang J, Papke RL. Positive allosteric modulators as an approach to nicotinic
acetylcholine receptor- targeted therapeutics: advantages and limitations. Biochem Pharmacol.
2011;82:915–930.
93. Wecker L, Pollock VV, Pacheco MA, Pastoor T. Nicotine-induced up regulation of α4β 2 neuronal
nicotinic receptors is mediated by the PKC-dependent phosphorylation of α4 subunits.
Neuroscience. 2010;171:12–22.
94. Sun Y-G, Rupprecht V, Zhou L, Dasgupta R, Seibt F, Beierlein M. mGluR1 and mGluR5
Synergistically Control Cholinergic Synaptic Transmission in the Thalamic Reticular Nucleus. J
Neurosci. 2016;36:7886–7896.
95. López-Hernández GY, Ananth M, Jiang L, Ballinger EC, Talmage DA, Role LW.
Electrophysiological properties of basal forebrain cholinergic neurons identified by genetic and
optogenetic tagging. J Neurochem. 2017;142 Suppl 2:103–110.

96.   Kolisnyk B, Guzman MS, Raulic S, Fan J, Magalhães AC, Feng G, Gros R, Prado VF, Prado
627   MAM. ChAT-ChR2-EYFP mice have enhanced motor endurance but show deficits in attention and
628   several additional cognitive domains. J Neurosci. 2013;33:10427–10438.
629   97.   Lhopitallier C, Perrault C, Chauveau F, Saurini F, Berrard S, Granon S, Faure A. Characterization
630   of social behavior in young and middle-aged ChAT-IRES-Cre mouse. PloS One.
631   2022;17:e0272141.
632   98.   Thorn CA, Popiolek M, Stark E, Edgerton JR. Effects of M1 and M4 activation on excitatory
633   synaptic transmission in CA1. Hippocampus. 2017;27:794–810.
634   99.   Montani C, Canella C, Schwarz AJ, Li J, Gilmour G, Galbusera A, Wafford K, Gutierrez-Barragan
635   D, McCarthy A, Shaw D, Knitowski K, McKinzie D, Gozzi A, Felder C. The M1/M4 preferring
636   muscarinic agonist xanomeline modulates functional connectivity and NMDAR antagonist-induced
637   changes in the mouse brain. Neuropsychopharmacology. 2021;46:1194–1206.
638   100.   Larocca JN, Rodriguez-Gabin AG, Rashbaum WK, Weidenheim KM, Lyman WD. Muscarinic
639   receptor-dependent activation of phospholipase C in the developing human fetal central nervous
640   system. Brain Res. 1994;653:9–15.
641   101.   Miwa JM, Anderson KR, Hoffman KM. Lynx Prototoxins: Roles of Endogenous Mammalian
642   Neurotoxin-Like Proteins in Modulating Nicotinic Acetylcholine Receptor Function to Influence
643   Complex Biological Processes. Front Pharmacol. 2019;10:343.
644   102.   Venkatesan S, Chen T, Liu Y, Turner EE, Tripathy S, Lambe EK. Chrna5 is a marker of
645   acetylcholine super-responder subplate neurons with specialized expression of nicotinic modulator
646   proteins. 2022:2022.06.20.496737.
647   103.   Tekinay AB, Nong Y, Miwa JM, Lieberam I, Ibanez-Tallon I, Greengard P, Heintz N. A role for
648   LYNX2 in anxiety-related behavior. Proc Natl Acad Sci U S A. 2009;106:4477–4482.
649   104.   Wu M, Puddifoot CA, Taylor P, Joiner WJ. Mechanisms of Inhibition and Potentiation of α4β2
650   Nicotinic Acetylcholine Receptors by Members of the Ly6 Protein Family. J Biol Chem.
651   2015;290:24509–24518.
652   105.   Moriwaki Y, Kubo N, Watanabe M, Asano S, Shinoda T, Sugino T, Ichikawa D, Tsuji S, Kato F,
653   Misawa H. Endogenous neurotoxin-like protein Ly6H inhibits alpha7 nicotinic acetylcholine
654   receptor currents at the plasma membrane. Sci Rep. 2020;10:11996.
655   106.   Puddifoot CA, Wu M, Sung R-J, Joiner WJ. Ly6h Regulates Trafficking of Alpha7 Nicotinic
656   Acetylcholine Receptors and Nicotine-Induced Potentiation of Glutamatergic Signaling. J Neurosci.
657   2015;35:3420–3430.
658   107.   Nichols WA, Henderson BJ, Yu C, Parker RL, Richards CI, Lester HA, Miwa JM. Lynx1 Shifts
659   α4β2 Nicotinic Receptor Subunit Stoichiometry by Affecting Assembly in the Endoplasmic
660   Reticulum. J Biol Chem. 2014;289:31423–31432.
661   108.   Bymaster FP, Whitesitt CA, Shannon HE, DeLapp N, Ward JS, Calligaro DO, Shipley LA, Buelke-
662   Sam JL, Bodick NC, Farde L, Sheardown MJ, Olesen PH, Hansen KT, Suzdak PD, Swedberg
663   MDB, Sauerberg P, Mitch CH. Xanomeline: A selective muscarinic agonist for the treatment of
664   Alzheimer's disease. Drug Dev Res. 1997;40:158–170.
665   109.   Karuna Therapeutics. A Phase 2, Randomized, Double-blinded Study to Assess the Safety,
666   Tolerability, and Efficacy of KarXT in Hospitalized Adults With DSM-5 Schizophrenia.
667   clinicaltrials.gov; 2020.
668   110.   Brown, Bradley SJ, Marshall FH, Brown GA, Bennett KA, Brown J, Cansfield JE, Cross DM, de
669   Graaf C, Hudson BD, Dwomoh L, Dias JM, Errey JC, Hurrell E, Liptrot J, Mattedi G, Molloy C,
670   Nathan PJ, Okrasa K, Osborne G, Patel JC, Pickworth M, Robertson N, Shahabi S, Bundgaard C,
671   Phillips K, Broad LM, Goonawardena AV, Moirarty SR, Browning M, Perini F, Dawson GR, Deakin
672   JFW, Smith RT, Sexton PM, Warneck J, Vinson M, Tasker T, Tehan BG, Teobald B, Christopoulos
673   A, Langmead CJ, Jazayeri A, Cooke RM, Rucktooa P, Congreve MS, Weir M, Tobin AB. From
674   structure to clinic: Design of a muscarinic M1 receptor agonist with potential to treatment of
675   Alzheimer's disease. Cell. 2021;184:5886-5901.e22.
676

Figure Legends

Figure 1. Responses to optogenetically-released acetylcholine decrease with age
(**A**) Image depicts coronal brain section taken for whole-cell electrophysiological recordings. The area outlined in pink shows layer 6 of the prefrontal (prelimbic and infralimbic) cortex. The recording electrode is represented in black and the delivered optogenetic stimulus is represented in blue. (**B**) Schematic represents the cholinergic prefrontal synapse receiving optogenetic stimulus (0.5s pulse train of decreasing frequency). Afferents release endogenous acetylcholine (ACh) which binds to nicotinic and muscarinic receptors, exciting postsynaptic pyramidal neurons. (**C**) Example recordings of opto-ACh current from a neuron from younger and older mice, measured in voltage clamp (Vm = -75mV, stimuli in blue, triple exponential fit in red). (**D, E**) Graphs show the decrease with age of the peak amplitude and the charge transfer of the opto-Ach current (n = 185 neurons, 43 mice). (**F, G**) Graphs show the increase in opto-ACh current peak and charge transfer with M2 muscarinic antagonist AF-DX 116 are not different between younger (< P150) and older (≥ P150) responses. (**H, I**) Graphs show the change in peak amplitude and charge transfer of the opto-ACh current with galantamine are not different between younger and older responses.

Figure 2. Nicotinic response is decreased and muscarinic response is preserved in aging
(**A**) Schematic illustrates muscarinic receptor blockade to isolate the nicotinic response. (**B**) Example optogenetically-elicited nicotinic responses recorded in voltage clamp (Vm = -75mV) in the presence of muscarinic receptor blocker atropine (AT) from younger and older mice. Graphs show significant age-group decreases in opto-ACh nicotinic responses: (**C**) peak amplitude, (**D**) charge transfer and (**E**) rising slope (**P <0.01, unpaired t-test, n = 29 neurons, 18 mice). (**F**) Schematic illustrates nicotinic receptor blockade to isolate the postsynaptic muscarinic response. (**G**) Example responses recorded in voltage clamp (Vm = -75mV) in the presence of nicotinic receptor blocker DHβE show muscarinic currents in neurons from younger and older mice. Graphs show no significant age group differences in opto-ACh muscarinic currents: (**H**) peak amplitude, (**I**) charge transfer, or (**J**) 10-90 rise time. (**K**) Example responses recorded in current clamp with depolarizing current injected to bring the neuron to fire action potentials in the presence of nicotinic receptor blocker DHβE show muscarinic firing response in neurons from younger and older mice. Graphs show no significant age-group difference in (**L**) peak firing frequency, (**M**) duration of opto-ACh firing change, or (**N**) time to peak firing.

Figure 3. Older opto-ACh responses are less sensitive to nicotinic allosteric potentiation
(**A**) Schematics show acetylcholine activation of nicotinic receptors at baseline and with NS9283 positive allosteric modulation. (**B**) Paired recordings in voltage clamp (Vm = -75mV) show opto-ACh current responses before (black) and after (blue) bath-application of nicotinic positive allosteric modulator (PAM) NS9283 in neurons from younger and older mice. (**C**) Graph shows peak amplitude of paired opto-ACh currents before and after NS9283 in neurons from younger and older mice (*P < 0.05, Interaction, two-way ANOVA; ****P < 0.0001, Sidak's multiple comparisons). Inset graph shows the change in current amplitude imparted by NS9283 is significantly greater in younger than older mice (*P <0.05, unpaired t-test). (**D**) Graph shows charge transfer of paired opto-ACh currents before and after NS9283 in neurons from younger and older mice (****P < 0.0001, NS9283, **P < 0.01, Age, two-way ANOVA; ***P < 0.001, Sidak's multiple comparisons). Inset graph shows the change in charge transfer imparted by NS9283 trends towards a difference between younger and older mice (t_{24} = 1.8, $^{\#}P$ = 0.08).

724
Figure 4. Aged nicotinic responses improved by muscarinic M1 agonist xanomeline
(**A**) Schematic of cholinergic receptor activation in prefrontal layer 6 pyramidal neurons in younger, older, and older with intervention to improve nicotinic receptor availability. (**B**) Paired recordings in voltage clamp (Vm = -75mV) in older mouse show opto-ACh response before (black) and after (green) bath-application of muscarinic M1 agonist xanomeline. (**C, D**) Graphs show peak amplitude and charge transfer of older responses significantly increases with xanomeline (**$P < 0.01$, ***$P < 0.001$, paired t-test). (**E**) Paired recordings in voltage clamp (Vm = -75mV) show opto-ACh response with xanomeline (green) is greatly reduced by nicotinic receptor blocker DHβE (orange). Graph (*inset*) shows peak amplitude of responses with xanomeline are significantly reduced with application of DHβE (***$P < 0.001$, paired t-test). (**F, G**) Graphs show potentiation with xanomeline is blocked in the presence of muscarinic receptor blocker atropine (AT) as well as PKC inhibitors (chelerythrine or Go6983). There is a significant drug effect on the percent change of peak amplitude, (**$P < 0.01$, one-way ANOVA; *$P < 0.05$, ** $P < 0.01$, Dunett's *post hoc* tests) and charge transfer (*$P < 0.05$, one-way ANOVA; ** $P < 0.01$, Dunett's *post hoc* tests).

Figure 5. Older opto-ACh responses rescued to younger levels by combined treatment of M1 agonist xanomeline and nicotinic PAM NS9283
(**A**) Paired recordings in voltage clamp (Vm = -75mV) show opto-ACh response before (black), after bath-applied xanomeline (green), and after bath-applied NS9283 (teal) from an older mouse. (**B**) Peak amplitude, (**C**) charge transfer, and (**D**) rising slope of older responses with xanomeline (xano) significantly increases with NS9283 (NS) (***$P < 0.001$, paired t-test). (**E**) Graph shows peak amplitude of opto-ACh responses normalized to the mean peak amplitude of the younger opto-ACh responses, with grey denoting one standard deviation of younger opto-ACh responses. There is a significant effect of pharmacological interventions in rescuing older responses (****$P < 0.0001$, one-way ANOVA; *** $P < 0.001$, Tukey's *post hoc* tests). (**F**) Schematic shows model for nicotinic receptor changes with age and the effects of combined xanomeline and NS9283 pharmacological intervention. (**G**) Example exponential fits of opto-ACh responses from younger, older, and older neuron treated with xanomeline and NS9283. Combined pharmacological intervention rescues older opto-ACh responses to younger levels.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5