Metatranscriptomic response of the wheat holobiont to decreasing soil water content

Pranav M. Pande[1], Hamed Azarbad[2], Julien Tremblay[3], Marc St-Arnaud[4], Etienne Yergeau[1]*

[1]Institut national de la recherche scientifique, Centre Armand-Frappier Santé Biotechnologie, Laval (Québec), H7V 1B7, Canada

[2] Philipps-University Marburg, Department of Biology, Evolutionary Ecology of Plants, Marburg, Germany

[3]National Research Council of Canada, Energy Mining and Environment, Montréal (Québec), Canada.

[4]Institut de recherche en biologie végétale, Université de Montréal et Jardin Botanique de Montréal, Montréal (Québec), Canada.

*Corresponding author : Institut national de la recherche scientifique, Centre Armand-Frappier Santé Biotechnologie, 531 boulevard des Prairies, Laval (Québec), H7V 1B7, Canada, Phone : +1 450 687-5010 ext. 8881, Etienne.Yergeau@inrs.ca

Abstract

Crops associate with microorganisms that help their resistance to biotic. However, it is not clear
how the different partners of this association react during exposure to stresses. This knowledge is
needed to target the right partners when trying to adapt crops to climate change. Here, we grew
wheat in the field under rainout shelters that let through 100%, 75%, 50% and 25% of the
precipitation. At the peak of the growing season, we sampled plant roots and rhizosphere, and
extracted and sequenced their RNA. We compared the 100% and the 25% treatments using
differential abundance analysis. In the roots, most of the differentially abundant (DA) transcripts
belonged to the fungi, and most were more abundant in the 25% precipitation treatment. About
10% of the DA transcripts belonged to the plant and most were less abundant in the 25%
precipitation treatment. In the rhizosphere, most of the DA transcripts belonged to the bacteria
and were generally more abundant in the 25% precipitation treatment. Taken together, our results
show that the transcriptomic response of the wheat holobiont to decreasing precipitation levels is
more intense for the fungal and bacterial partners than for the plant.

Introduction

Drought is one of the most significant threats to crops and will become more frequent and

intense with climate change [1]. Both the plant and its microbiota respond to decreasing soil

water content, which affects the fitness of the plant. However, because of a lack of studies

integrating plant and microorganisms, the best targets for improving crop resistance to water

stress are not clear. Many *Actinobacteria* and *Proteobacteria* can improve plant tolerance to

drought- or salinity-related stresses [2-5]. Fungal endophytes can also improve plant

performance under abiotic stress [6-8]. Mycorrhizal fungi can improve water use efficiency and

reduce drought stress in wheat [9], oat [10], and corn [11]. Interestingly, endophytic and

rhizospheric microorganisms isolated from environments prone to drought tend to confer plants

with a better resistance to drought [8, 12]. Many mechanisms are involved in the enhancement of

plant drought tolerance by microbes. These include modulation of plant drought stress genes

[13], reduction of the stress hormone ethylene levels through degradation of its precursor 1-

aminocyclopropane-1-carboxylic acid (ACC) by the bacterial enzyme ACC deaminase [2, 3],

stimulation of the expression of plant genes related to osmolytes and osmoprotectants by

bacterial volatile organic compounds [14] and modulation of the plant epigenetics response to

drought [15]. Plants also directly respond to water stress through genetic, molecular and

physiological mechanisms [16].

A host and its microbiota form an holobiont, and their combined genomes is the

hologenome [17]. Although the concept has been debated [18-21], it is useful in emphasising the

role that microbial communities play in the host biology [for more details on these concepts, see

17]. The hologenome theory of evolution [22] considers the hologenome as one evolutionary

unit, which provides an interesting framework for studying the adaptation of holobionts to

stressful conditions. It implies that there are microbial-driven means by which holobionts can

adapt to new environmental conditions [23-25]. The hologenome can change through 1)

recruitment of new microbial partners from external sources, 2) amplification or reduction of the

microbial partners already in place, and 3) horizontal gene transfer (HGT) from the external

communities to the microbial partners already in place. These are coherent with the mechanisms

of ecological community change put forward in the theory of ecological communities [26, 27],

namely 1) migration, 2) selection, 3) speciation and 4) drift. At the transcriptomic level, the

microbial response can stem from two mechanisms: 1) changes in the metagenome (by the three

mechanisms listed above) and 2) changes in the gene expression of the members of the

community. Although these two mechanisms cannot be disentangle using metatranscriptomics,

the result will be the same: a change in the genes expressed within the holobiont. For the host,

the transcriptomic response is limited to shifts in gene expression. We therefore hypothesized

that most of the transcriptomic response of the wheat holobiont to decreasing soil water

availability will be microbial. To test this hypothesis, we grew wheat under rainout shelters that

let through 25, 50, 75 or 100% of the natural precipitation. Plant roots and rhizosphere were

sampled, their RNA extracted and sequenced.

Materials and methods

Experimental design and sampling

Four rainfall manipulation treatments were set-up in 2016 at the Armand-Frappier Santé

Biotechnologie Centre (Laval, Québec, Canada) using rain-out shelters that passively let through

25%, 50%, 75% and 100% of the natural precipitation. The rainfall exclusion treatments were

performed using 2m x 2m rain-out shelters, which were covered with nine, six, three, or zero 2m

x 16.7cm sheets of transparent plastic for the 25%, 50%, 75% and 100% treatments, respectively.

The rain was intercepted by the plastic sheeting and guided in a gutter and downspout and

collected in 20L buckets that were manually emptied when they were full. Two wheat genotypes

were seeded under these shelters (drought sensitive, *Triticum aestivum* cv. AC Nass and drought

tolerant, *Triticum turgidum spp. durum* cv. Strongfield), and the experiment was replicated over

6 fully randomized blocks, resulting in 48 plots (4 treatments x 2 genotypes x 6 blocks). Plots

were seeded at a density of 500 seeds per m^2 on May 18 (2016) and May 23 (2017). Seeds

harvested from each of the plots were re-seeded in the exact same plot the following year. For

the current manuscript, only the Strongfield cultivar was used, from which rhizosphere soil and

root samples were taken on July 26, 2017. For rhizosphere sampling, a plant was randomly

selected (avoiding the edge of the plots), uprooted and shaken vigorously to remove the loosely

attached soil. Soil tightly adhering to roots after shaking was considered as rhizosphere soil and

was collected in sterile 1.5 ml microcentrifuge tubes. After collecting the rhizosphere soil, roots

were washed with distilled water, separated from the plant and stored in sterile 15 ml Falcon

tubes. Collected rhizosphere soil and root samples were flash frozen in liquid nitrogen within a

span of 2 minutes after uprooting the plant to maintain the RNA integrity. Tubes were stored at -

80ºC until the samples were processed for RNA extraction. At sampling, we also collected a bulk

soil sample from the center of each plot for soil water content measurement. We measured soil

water content by weighing soils before and after drying overnight at 105ºC.

RNA extraction and sequencing

Total RNA was extracted from 2 g of rhizosphere soil using the RNeasy PowerSoil Total RNA

Kit (QIAGEN, Canada) and 0.5 g roots using RNeasy Plant Mini Kit (QIAGEN, Canada).

Extracted RNA was treated with DNAse (ThermoFisher, Canada) to remove the DNA prior to

sequencing. The absence of DNA was confirmed by the lack of PCR amplification using 16S

rRNA gene specific primers. Total RNA was sent for Illumina HiSeq4000 2 x 100 bp pair end

sequencing at the Centre d'Expertise et de Services Génome Québec (Montréal, Québec).

Libraries for rhizosphere samples were created using a microbial ribosome subtraction approach

to capture all microbial transcripts, whereas libraries for root samples were created using a poly-

dT reverse transcription approach to focus on the plant transcripts. The raw data produced in this

study was deposited in the NCBI under Bioproject accession PRJNA880647.

Bioinformatics

The metatranscriptome sequencing of the 24 root and 24 rhizosphere samples resulted in 2,639M

reads resulting in 264 giga bases which were processed together through our metatranscriptomics

bioinformatics pipeline [28]. Briefly, sequencing adapters were removed from each read and

bases at the end of reads having a quality score less than 30 were cut off (Trimmomatic v0.32)

[29] and scanned for sequencing adapters contaminants reads using DUK

(http://duk.sourceforge.net/) to generate quality controlled (QC) reads. QC-passed reads from

each sample were co-assembled using Megahit v1.1.2 [30] with iterative kmer sizes of

31,41,51,61,71,81 and 91 bases. Gene prediction was performed by calling genes on each

assembled contig using Prodigal v2.6.2 [31]. Genes were annotated following the JGI's

guidelines [32] including the assignment of KEGG orthologs (KO). QC-passed reads were

mapped (BWA mem v0.7.15) (unpublished - http://bio-bwa.sourceforge.net) against contigs to

assess quality of metagenome assembly and to obtain contig abundance profiles. Alignment files

in bam format were sorted by read coordinates using samtools v1.2 [33] and only properly

aligned read pairs were kept for downstream steps. Each bam file (containing properly aligned

paired-reads only) was analyzed for coverage of called genes and contigs using bedtools

(v2.17.0) [34] using a custom bed file representing gene coordinates on each contig. Only paired

reads both overlapping their contig or gene were considered for gene counts. Coverage profiles

of each sample were merged to generate an abundance matrix (rows = contig, columns =

samples) for which a corresponding CPM (Counts Per Million – normalized using the TMM

method) (edgeR v3.10.2) [35]. Each contig was blasted (BLASTn v2.6.0+) against NCBI's nt

database (version downloaded from NCBI's server on January 9th 2019) and the best hit's

taxonomic identifier was used to assign a taxonomic lineage to the contig. Taxonomic summaries

were performed using MicrobiomeUtils v0.9 (github.com/microbiomeutils). The

metatranscriptome co-assembly, gene abundance, read count summaries and mapping statistics

and other results generated by our bioinformatic workflow are provided in the companion online

Zenodo archive (https://doi.org/10.5281/zenodo.7121038).

Statistical analyses

All statistical analyses were performed in R version 4.1.0. [36]. Transcript differential abundance

analyses between the 100% and 25% precipitation treatments were carried out using the EBTest

function of the EBSeq library with a false discovery rate (FDR) of 0.05. Anovas were performed

using the aov function of the stats package. All graphs were generated using ggplot2. The R

project folder containing the R code used for data manipulation, statistical analyses, and tables

and figure generation is available on our lab GitHub repository (https://github.com/le-labo-

yergeau/MT_Holobiont_Wheat). The associated transcript abundance and annotation tables, the

metadata, and the soil water content files used with the R code are available on Zenodo:

https://doi.org/10.5281/zenodo.7096909.

Results

Soil water content (SWC)

There was a significant difference (p = 0.000367) between the mean SWC across the four treatments. The water content was highest in plots exposed to 100% of the natural precipitation and gradually decreased in plots receiving 75%, 50% and 25% of the natural precipitation (Fig. 1a). The SWC was of 11% at its lowest (in the 25% precipitation treatment) and of 23% at its highest (in the 100% treatment). The rest of our analyses focus on the two most extreme conditions, the 25% and 100% precipitation treatments.

Responses of the holobiont partners

We retrieved 1,069,108,624 clean sequencing reads (per sample, mean: 22,746,992, max: 43,912,775, min: 13,943,395) that were assembled in a total of 1,269,055 transcripts, among which 1,193,501 and 1,136,587 transcripts were found in the rhizosphere soil and wheat roots, respectively. Among the wheat root transcripts, 12,792 (1.1%) belonged to the Archaea, 365,435 (32.2%) to Bacteria, 200,233 (17.6%) to Fungi, 200,823 (17.7%) to plants, 132,313 (11.6%) to other Eukaryotes, 3,660 (0.3%) to viruses and 221,331 (19.5%) were not identified at the kingdom level (Fig 1b). Among the rhizosphere soil transcripts, 14,943 (1.3%) belonged to the archaea, 430,984 (36.1%) to the bacteria, 186,745 (15.6%) to the fungi, 169,788 to the plants (14.2%), 132,042 (11.1%) to other eukaryotes, 4,255 (0.4%) to viruses, and 254,744 (21.3%) were not classified at the kingdom level (Fig. 1b).

In the roots, among the 1,136,587 transcripts, 42,001 (3.70%) were differentially abundant (DA) at a false discovery rate (FDR) of 0.05. Among these DA transcripts, 2,309

belonged to the bacteria (5.50%), 23,274 to the fungi (55.41%), 4,357 to the plants (10.37%),

5,303 were not classified at the kingdom level (12.63%) and 6,758 belonged to other taxa

(16.09%) (Fig. 1c and 2a). For bacteria and fungi, most of the DA transcripts were more

abundant in the 25% treatment as compared to the 100% treatment (23,042 and 2,231 more

abundant vs. 232 and 78 less abundant for fungi and bacteria, respectively), whereas it was the

inverse for plant (1,295 genes more abundant vs. 3,061 less abundant) (Fig. 1c and 2a).

In the rhizosphere, among the 1,193,501 transcripts, 21,765 (1.82%) were differentially

abundant at an FDR of 0.05. Among these DA transcripts, 14,178 belonged to the bacteria

(65.14%), 159 to the archaea (0.73%), 402 to the fungi (1.85%), 219 to the plants (1.01%), 5,224

were not classified at the kingdom level (24.00%) and 1,583 belonged to other taxa (7.27%) (Fig.

1c and 2b). For bacteria, slightly more DA transcripts were more abundant in the 25% treatment

as compared to the 100% treatment (7,938 more abundant vs. 6,240 less abundant), whereas it

was the inverse for plant (41 more abundant vs. 178 less abundant) and fungi (149 more

abundant vs. 253 less abundant) (Fig. 1c and 2b).

High level taxonomy and functions of the DA transcripts

We compared the taxonomic affiliations at the phylum/class levels for all transcripts vs. for the

positive and negative DA transcripts in the roots and the rhizosphere (Fig. 3). Since the DA

analyses result in a single list of DA transcripts per plant compartment, we are not able to test

statistically for the differences in the representation of the taxa in the different subsets. However,

interesting trends emerged. Some taxa were relatively less abundant among DA transcripts than

among all transcripts, suggesting a lack of response to the precipitation exclusion treatments. The

Sordariomycetes, Chloroflexi, Gemmatimonadetes, among others, were in this situation across

all compartments, together with the Acidobacteria in the roots and the Dothideomycetes in the

rhizophere (Fig. 3). Other taxa were overrepresented among the positive DA transcripts and

underrepresented among the negative DA transcripts, suggesting an increase in relative

abundance or an upregulation of several genes under lower soil water content. The

Actinobacteria in both compartments, the Ascomycota in the rhizosphere, and the

Dothideomycetes in the roots were in that situation (Fig. 3). In contrast, some taxa were

overrepresented among the negative DA transcripts and underrepresented among the positive DA

transcripts, suggesting a decrease in relative abundance or a downregulation of several genes

under lower soil water content. The Proteobacteria, Bacteroidetes, and Eurotiomycetes in both

compartments, the Acidobacteria in the rhizosphere and the Agaricomycetes in the roots showed

this pattern (Fig. 3).

As for COG (clusters of orthologous genes) categories, some were overrepresented in the

positive DA transcripts and underrepresented in the negative DA transcripts (Fig. 4), suggesting

high-level categories that are generally upregulated following a reduction of soil water content.

Among these were "Carbohydrate transport and metabolism" and "Lipid metabolism" in the

roots, " "Cell envelope biogenesis, outer membrane", "Signal transduction mechanisms", and

"Transcription" in the rhizosphere (Fig. 4). The COG categories overrepresented in the negative

DA transcripts included "Translation, ribosomal structure and biogenesis" in both the

rhizosphere and the roots and "Posttranslational modifications, protein turnover, chaperones" in

the roots (Fig. 4). These would be COG categories that are generally downregulated with

decreasing soil water content. Some COG categories were relatively less abundant among

positive and negative DA transcripts than among all transcripts, suggesting a lack of response to

the precipitation exclusion treatments. This included "Amino acid transport and metabolism" in

the rhizosphere and "Cell envelope biogenesis, outer membrane", "DNA replication,

recombination and repair", "Signal transduction mechanisms" and "Transcription" in the roots

(Fig. 3

Most differentially abundant transcripts

For DA analyses of the root samples, there were many transcripts that had a P-value=0, so we

sorted them by mean abundance and are showing the top 50 transcripts in Table 1 and Figure 5A.

Twenty-seven transcripts among the top 50 transcripts belonged to the *Agaricomycetes*, mostly

Coprinopsis cinerea, and were almost all more abundant in the 25% precipitation treatment

(Table 1 and Fig. 5A). Seven transcripts could be related to the wheat tribe (*Triticum aestivum* or

Aegilops tauschii), all of which were less abundant in the 25% precipitation treatment (Table 1

and Fig. 5A). Many of the most significantly more abundant transcripts in the 25% precipitation

treatment were related to amino acid and carbohydrate transport and metabolism, with transcripts

such as "Amino acid transporters", "Glycerol uptake facilitator and related permeases", "Beta-

glucanase/Beta-glucan synthetase", "Dipeptide/tripeptide permease", "Fucose permease",

"Neutral trehalase" and "Hexokinase" (Table 1). In contrast, many of the most significantly less

abundant transcripts in the 25% precipitation treatments were linked to the COG categories

"Posttranslational modification, protein turnover, chaperones" and "Secondary metabolites

biosynthesis, transport and catabolism" (Table 1).

For the rhizosphere, as not that many DA transcripts had a P-value=0, we are presenting

the 50 lowest P-values observed in Table 2 and Figure 5B. Half the DA transcripts with the

lowest P-values belonged to the *Proteobacteria*, mainly the *Alpha-* and *Delta-* classes (Table 2

and Fig 5B). Many of the most significantly less abundant transcripts in the 25% precipitation

treatment were linked to the COG categories "Cell motility and secretion" and "Intracellular

trafficking and secretion", with functions related to pilus, flagella and type II and VI secretion

systems (Table 2). Similar to what we observed in the roots, the fungal transcripts more abundant

in the 25% precipitation treatment belonged to the *Agaricomycetes* and were related to

carbohydrate and amino acid transport and metabolism (e.g. "monoamine oxidase",

"hexokinase") (Table 2).

DA transcripts common to roots and rhizosphere

We looked for DA transcripts that showed a common DA response in roots and the rhizosphere.

Among the 37,242 and 10,565 positive DA transcripts in roots and the rhizosphere, respectively,

513 were shared (Fig. 6). Out of these 513 transcripts, 392 were affiliated to the Actinobacteria,

27 to the Basidiomycota, 12 to the Ascomycota and 11 to the Proteobacteria (Table S1). The

most represented COG category were "Translation, ribosomal structure and biogenesis" (43

transcripts), "Transcription" (33 transcripts), "Carbohydrate transport and metabolism" (29

transcripts), "Posttranslational modification, protein turnover, chaperones" (26 transcripts) and

"Amino acid transport and metabolism" (14 transcripts) (Table S1). Among the 4,758 and 11,200

negative DA transcripts for roots and rhizosphere, respectively, 47 transcripts were shared (Fig.

6). Most of these transcripts were not affiliated at the phylum level (26 transcripts), followed by

transcripts affiliated to Streptophyta (7 transcripts) and Basidiomycota (3 transcripts) (Table S2).

For COG categories, again, most of the transcripts were not affiliated with a category, and the

rest were mostly affiliated to "Cytoskeleton" (5 transcripts), "Energy production and conversion"

(2 transcripts), and "Translation, ribosomal structure and biogenesis" (2 transcripts) (Table S2).

Discussion

We wanted to know how the wheat holobiont would respond to change in soil water availability at the transcriptomic level, and which of the partners would be more responsive. We were successful in reducing soil water content in a field experiment using rainout shelters, and found that, when comparing the two most contrasting treatments, most of the differentially abundant (DA) genes were linked to the fungi in the roots and to the bacteria in the rhizosphere. In the roots, most of the DA fungal transcripts were more abundant, whereas about half the DA bacterial transcripts in the rhizosphere were more abundant and the other half less abundant. These DA transcripts belonged to specific taxa and many of them could be related to genes known to help plants and microorganisms cope with water stress. Our results agree with one of our previous studies of the willow holobiont that showed that the root fungi are the strongest responders to soil contamination [37]. Bacteria responded mainly by expressing pollutant degradation genes, whereas plants did not show large transcriptomic responses [37]. Plant gene expression in the roots was more variable across plant genotypes than between contaminated and non-contaminated soils, in contrast to the strong response of bacteria to soil contamination [38].

The microbial component of the hologenome (the metagenome) is much more dynamic and plastic than the host genome [24]. Indeed, the microbial metagenome can be modified rapidly by changing the relative abundance of the community members, by recruiting new members from the environment or through mechanisms such as horizontal gene transfer (HGT) [24]. The host genome cannot be modified in response to environmental stress within a single generation. This could explain why most of the DA transcripts were microbial, as it combines changes in microbial gene expression and in the metagenome. The response of the host is limited to changing gene expression levels. The changes detected in plant gene expression could still

affect important physiological processes, including root exudation [39]. As root exudates

influence the transcriptome of bacteria [40], the microbial transcriptomic response to decreasing

soil water content could have been mediated by the plant. The water depletion caused by our

rainout shelters did not result in extremely low soil water content (around 12% soil water content

at the lowest), which did not result in any visible stress on wheat. The wheat variety used is also

water stress resistant, and this could explain the lack of a strong transcriptomic response. It

would be interesting to contrast our results to the transcriptomic response of sensitive plant

holobionts when exposed to much more extreme stress levels.

With the method used here, it is difficult to disentangle the metatranscriptomic response

due to shifts in the composition of the microbial community and in the gene expression within

the same community. The *Actinobacteria* well exemplify this. There was an overrepresentation

of the *Actinobacteria* among positive DA transcripts in the rhizosphere, and most of the

transcripts that were positively DA in both the roots and the rhizosphere were from this phylum.

This phylum increases in relative abundance when soils get drier [41-45]. Inversely, the

Proteobacteria and *Acidobacteria* were overrepresented among the negative DA transcripts and

underrepresented among the positive DA transcripts, in line with their heightened sensitivity to

water stress [45, 46]. In these two cases, the shifts observed are likely a combination of shifts in

the relative abundance and of gene expression. Therefore, we referred to our differential

expression analysis as a transcript differential abundance analysis. Looking only at high-level

functional categories, like in Figure 4, could partly solve this problem, as general trends in gene

expression at this level is less likely to be influenced by shifts in community composition.

Nevertheless, we argue that whatever the underlying mechanisms are, variation in the

rhizosphere and root metatranscriptome complement will have functional consequences on the

holobiont adaptation to stress.

Many of the most positive DA transcripts in the roots under 25% precipitation regime,

were related to amino acid and carbohydrate transport and metabolism. Amino acids, such as

proline, glutamine, and glycine, betaine, and carbohydrates, such as trehalose and ectoine can be

used as osmolytes [47] to maintain cellular turgor and protect macromolecular structures [48].

Gram-negative bacteria produce osmolytes purely as a drought-inducible response, whereas

Gram-positive bacteria tend to produce osmolytes, at least partially, on a constitutive basis [49],

which could explain some of the differences in the transcriptomic response of different taxa

observed here. It would be interesting to know how much this higher abundance of transcripts is

beneficial to the microbes vs. the host plant. There is some evidence that microbial endophytes

and rhizobacteria can increase plant osmolyte concentration [50, 51], including proline [52], and

some studies have reported that microbes can exude these compounds in the plant environment

[53, 54], enabling them to directly contribute to the plant osmolyte concentration during water

stress. For instance, *Coprinopsis* were often reported as endophytes of plants, including

Arabidopsis [55] and were found here among the root fungi that showed the strongest response

to decreasing soil water content, with many of their more abundant transcripts related to

carbohydrate or amino acid transport and metabolism.

Other important transcripts were affected by the precipitation treatments. Among the

rhizosphere bacteria, transcripts related to pilus and flagella formation were less abundant with

decreasing soil water content, which might be indicative of a switch from a free-living to a

biofilm lifestyle. Biofilm formation is a well-known mechanism that bacteria use to cope with

environmental stresses [56]. Transcripts related to heat shock proteins were more abundant in the

rhizosphere and the roots under low water content, in line with their important roles for microbes

and plants under water stress [57-59]. In both root and rhizosphere, there was an

overrepresentation of genes related to translation among the negative DA transcripts. A similar

down-regulation of the protein biosynthesis machinery was observed in a recent soil warming

metatranscriptomic study [60]. The author suggested that the increased enzymatic activity and

overall metabolism caused by warming could call for a lower energy investment in ribosomes,

thus optimizing resource allocation [60]. In contrast, during soil drying, *Acidobacteria* and

Verrocumicrobia reduced their ribosomal content, whereas the *Actinobacteria* increased it [61].

Similarly, among a general decrease in translation-related transcripts, we observed here that for

positive DA transcripts found in both the roots and in the rhizosphere, translation-related

transcripts affiliated to the Actinobacteria was the most represented category. This differential

regulation of translation among microbial groups could explain the dominance of Actinobacteria

under reduce soil water availability.

In conclusion, holobionts are posited to respond in a coordinated fashion to stressful

events. In our case, the microbial partners were clearly the strongest responders to decreasing

water content, being responsible for most of the DA transcripts across the wheat holobiont. We

had hypothesised that this would be the case since transcriptomic shifts in the microbiome

combines changes in the metagenome and in gene expression, something that is not possible for

the host. These transcriptomic shifts were related to microbial genes and taxa, such as the

Actinobacteria and osmolyte-related genes, that are known to be beneficial to plants under water

stress. Because of their dynamic response and beneficial potential, the microbiome should be

considered as central in efforts to adapt crop holobionts to water stress.

Acknowledgments

We would like to thank all the members, past and current, of the mECO:LABS for their help in setting up and maintaining the field experiment. We are also grateful to Dr. Sara Correa-Garcia for her help in making the R code portable and for submission to GitHub. This work was funded by the Natural Sciences and Engineering Research Council of Canada (Discovery grant RGPIN-2014-05274 and Strategic grant for projects STPGP 494702 to EY). We also wish to acknowledge Compute Canada for access to the University of Waterloo's High-Performance Computing (HPC) infrastructure (Graham system) through a resources allocation granted to EY.

Conflict of interest

The authors declare no conflict of interest.

References

1. IPCC (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, vol. In Press. Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA p.

2. Mayak S, Tirosh T, Glick BR. Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiol Biochem* 2004; **42:** 565-572.

3. Mayak S, Tirosh T, Glick BR. Plant growth-promoting bacteria that confer resistance to water stress in tomatoes and peppers. *Plant Sci* 2004; **166:** 525-530.

4. Cheng Z, Park E, Glick BR. 1-Aminocyclopropane-1-carboxylate deaminase from *Pseudomonas putida* UW4 facilitates the growth of canola in the presence of salt. *Can J Microbiol* 2007; **53:** 912-918.

5. Saravanakumar D, Samiyappan R. ACC deaminase from *Pseudomonas fluorescens* mediated saline resistance in groundnut (*Arachis hypogea*) plants. *J Appl Microbiol* 2007; **102:** 1283-1292.

6. Singh LP, Gill SS, Tuteja N. Unraveling the role of fungal symbionts in plant abiotic stress tolerance. *Plant Sign Behav* 2011; **6:** 175-191.

7. Hubbard M. Fungal endophytes that confer heat and drought tolerance to wheat. PhD Thesis. University of Saskatchewan: Saskatoon, 176 p.

8. Redman RS, Kim YO, Woodward CJ, Greer C, Espino L, Doty SL *et al.* Increased fitness of rice plants to abiotic stress via habitat adapted symbiosis: a strategy for mitigating impacts of climate change. *PLoS One* 2011; **6:** e14823.

9. Al-Karaki G, McMichael B, Zak J. Field response of wheat to arbuscular mycorrhizal fungi and drought stress. *Mycorrhiza* 2004; **14:** 263-269.

10. Khan IA, Ahmad S, Ayub N. Response of oat (Avena sativa) to inoculation with vesicular arbuscular mycorrhizae (VAM) in the presence of phosphorus. *Asian J Plant Sci* 2003.

11. Subramanian K, Charest C, Dwyer L, Hamilton R. Arbuscular mycorrhizas and water relations in maize under drought stress at tasselling. *New Phytol* 1995; **129:** 643-650.

12. Timmusk S, Paalme V, Pavlicek T, Bergquist J, Vangala A, Danilas T *et al.* Bacterial distribution in the rhizosphere of wild barley under contrasting microclimates. *PLoS One* 2011; **6:** e17968.

13. Timmusk S, Wagner EGH. The plant-growth-promoting rhizobacterium *Paenibacillus polymyxa* induces changes in *Arabidopsis thaliana* gene expression: a possible connection between biotic and abiotic stress responses. *Mol Plant Microbe Interact* 1999; **12:** 951-959.

14. Murzello C. Volatile organic compounds from *Bacillus subtilis* GBO3 promote osmotic and drought tolerance in *Arabidopsis thaliana* (Col-0). MS Thesis. Texas Tech University p.

15. Hubbard M, Germida JJ, Vujanovic V. Fungal endophyte colonization coincides with altered DNA methylation in drought-stressed wheat seedlings. *Can J Plant Sci* 2014; **94:** 223-234.

16. Tuberosa R, Grillo S, Ellis R. Unravelling the genetic basis of drought tolerance in crops. *Abiotic stresses in plants*. 2003. Springer, pp 71-122.

17. Rosenberg E, Zilber-Rosenberg I (2013). *The hologenome concept: human, animal and plant microbiota*. Springer: Heidelberg, 187 p.

18. Bordenstein SR, Theis KR. Host biology in light of the microbiome: Ten principles of holobionts and hologenomes. *PLoS Biol* 2015; **13:** e1002226.

19. Doolittle WF, Booth A. It's the song, not the singer: an exploration of holobiosis and evolutionary theory. 2017; **32:** 5-24.

20. Moran NA, Sloan DB. The Hologenome Concept: Helpful or Hollow? 2015; **13:** e1002311.

21. Theis KR, Dheilly NMM, Klassen JL, Brucker RM, Baines JF, Bosch TCG *et al.* Getting the hologenome concept right: An eco-evolutionary framework for hosts and their microbiomes. *mSystems* 2016; **1:** e00028-16.

22. Zilber-Rosenberg I, Rosenberg E. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Rev* 2008; **32:** 723-735.

23. Voss JD, Leon JC, Dhurandhar NV, Robb FT. Pawnobiome: Manipulation of the Hologenome Within One Host Generation and Beyond. *Front Microbiol* 2015; **6**.

24. Zilber-Rosenberg I, Rosenberg E. Microbial-driven genetic variation in holobionts. *FEMS Microbiol Rev* 2021.

25. Rosenberg E, Zilber-Rosenberg I. Bacterial Bleaching of Corals Leads to Hologenome Concept. *Microbe Mag* 2016.

26. Vellend M (2016). *The theory of ecological communities*. Princeton University Press, 229 p.

27. Agoussar A, Yergeau E. Engineering the plant microbiota in the context of the theory of ecological communities. *Curr Opin Biotechnol* 2021; **70:** 220-225.

28. Tremblay J, Schreiber L, Greer CW. High resolution shotgun metagenomics: the more data, the better? *biorXiv* 2022**:** 2022.04.19.488797.

29. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 2014; **30:** 2114-2120.

30. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 2015; **31:** 1674-1676.

31. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinfo* 2010; **11:** 119.

32. Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Tennessen K *et al.* The standard operating procedure of the DOE-JGI Metagenome Annotation Pipeline (MAP v. 4). *Stand Genom Sci* 2016; **11:** 17.

33. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N *et al.* The sequence alignment/map format and SAMtools. *Bioinformatics* 2009; **25:** 2078-2079.

34. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 2010; **26:** 841-842.

35. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 2010; **26:** 139-140.

36. R_Core_Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing: Vienna, Austria.

37. Gonzalez E, Pitre F, Pagé A, Marleau J, Guidi Nissim W, St-Arnaud M *et al.* Trees, fungi and bacteria: Tripartite metatranscriptomics of a root microbiome responding to soil contamination. *Microbiome* 2018; **6:** 53.

38. Yergeau E, Tremblay J, Joly S, Labrecque M, Maynard C, Pitre FE *et al.* Soil contamination alters the willow rhizosphere and root metatranscriptomes and the root-rhizosphere interactome. *ISME J* 2018; **12:** 869-884.

39. Williams A, de Vries FT. Plant root exudation under drought: implications for ecosystem functioning. *New Phytol* 2020; **225:** 1899-1905.

40. Mark GL, Dow JM, Kiely PD, Higgins H, Haynes J, Baysse C *et al.* Transcriptome profiling of bacterial responses to root exudates identifies genes involved in microbe-plant interactions. *Proc Natl Acad Sci USA* 2005; **102:** 17454-17459.

41. Naylor D, DeGraaf S, Purdom E, Coleman-Derr D. Drought and host selection influence bacterial community dynamics in the grass root microbiome. *ISME J* 2017.

42. Meisner A, Samuel J, Snoek BL, Ten Hooven FC, Van Der Putten WH. Drought legacy effects on the composition of soil fungal and prokaryote communities. *Front Microbiol* 2018; **9:** 294.

43. Preece C, Verbruggen E, Liu L, Weedon JT, Peñuelas J. Effects of past and current drought on the composition and diversity of soil microbial communities. *Soil Biol Biochem* 2019; **131:** 28-39.

44. Azarbad H, Tremblay J, Bainard LD, Yergeau E. Relative and Quantitative Rhizosphere Microbiome Profiling Results in Distinct Abundance Patterns. *Front Microbiol* 2022; **12:** 798023.

45. Azarbad H, Tremblay J, Giard-Laliberté C, Bainard L, Yergeau É. Four decades of soil water stress history together with host genotype constrain the response of the wheat microbiome to soil moisture. *FEMS Microbiol Ecol* 2020; **96:** fiaa098.

46. Agoussar A, Azarbad H, Tremblay J, Yergeau É. The resistance of the wheat microbial community to water stress is more influenced by plant compartment than reduced water availability. *FEMS Microbiol Ecol* 2021; **97:** fiab149.

47. Bouskill NJ, Wood TE, Baran R, Ye Z, Bowen BP, Lim H *et al.* Belowground response to drought in a tropical forest soil. I. Changes in microbial functional potential and metabolism. *Front Microbiol* 2016; **7:** 525.

48. Welsh DT. Ecological significance of compatible solute accumulation by micro-organisms: from single cells to global climate. *FEMS Microbiol Rev* 2000; **24:** 263-290.

49. Schimel J, Balser TC, Wallenstein M. Microbial stress-response physiology and its implications for ecosystem function. *Ecology* 2007; **88:** 1386–1394.

50. Gagné-Bourque F, Bertrand A, Claessens A, Aliferis KA, Jabaji S. Alleviation of drought stress and metabolic changes in timothy (Phleum pratense L.) colonized with Bacillus subtilis B26. *Front Plant Sci* 2016; **7:** 584.

51. Khan N, Bano A, Rahman MA, Guo J, Kang Z, Babar MA. Comparative Physiological and Metabolic Analysis Reveals a Complex Mechanism Involved in Drought Tolerance in Chickpea (Cicer arietinum L.) Induced by PGPR and PGRs. *Sci Rep* 2019; **9:** 2097.

52. Vílchez JI, Niehaus K, Dowling DN, González-López J, Manzanera M. Protection of pepper plants from drought by Microbacterium sp. 3J1 by modulation of the plant's glutamine and α-ketoglutarate content: a comparative metabolomics approach. *Front Microbiol* 2018; **9:** 284.

497   53.   Madkour MA, Smith LT, Smith GM. Preferential osmolyte accumulation: a mechanism
498   of osmotic stress adaptation in diazotrophic bacteria. *Appl Environ Microbiol* 1990; **56:**
499   2876-2881.
500   54.   Paul D, Nair S. Stress adaptations in a plant growth promoting rhizobacterium (PGPR)
501   with increasing salinity in the coastal agricultural soils. *J Basic Microbiol* 2008; **48:** 378-
502   384.
503   55.   Mesny F, Miyauchi S, Thiergart T, Pickel B, Atanasova L, Karlsson M *et al.* Genetic
504   determinants of endophytism in the Arabidopsis root mycobiome. 2021; **12:** 7227.
505   56.   Rudrappa T, Biedrzycki ML, Bais HP. Causes and consequences of plant-associated
506   biofilms. *FEMS Microbiol Ecol* 2008; **64:** 153-166.
507   57.   Yu T-F, Xu Z-S, Guo J-K, Wang Y-X, Abernathy B, Fu J-D *et al.* Improved drought
508   tolerance in wheat plants overexpressing a synthetic bacterial cold shock protein gene
509   SeCspA. *Sci Rep* 2017; **7:** 44050.
510   58.   Augustine SM. Function of heat-shock proteins in drought tolerance regulation of plants.
511   *Drought Stress Tolerance in Plants, Vol 1*. 2016. Springer, pp 163-185.
512   59.   Castiglioni P, Warner D, Bensen RJ, Anstrom DC, Harrison J, Stoecker M *et al.* Bacterial
513   RNA chaperones confer abiotic stress tolerance in plants and improved grain yield in
514   maize under water-limited conditions. *Plant Physiol* 2008; **147:** 446-455.
515   60.   Söllinger A, Séneca J, Dahl MB, Motleleng LL, Prommer J, Verbruggen E *et al.* Down-
516   regulation of the bacterial protein biosynthesis machinery in response to weeks, years,
517   and decades of soil warming. *Sci Adv* 2022; **8:** eabm3230.
518   61.   Barnard RL, Osborne CA, Firestone MK. Responses of soil bacterial and fungal
519   communities to extreme desiccation and rewetting. *ISME J* 2013; **7:** 2229-2241.

Figure legends

Figure 1: (A) mean soil water content at the time of sampling for the four different precipitation manipulation treatments, (B) kingdom-level taxonomic affiliation of the transcripts retrieved for all roots and rhizosphere samples, (C) kingdom-level affiliation of the differentially abundant (DA) transcripts together with information if they were more or less abundant in the 25% treatment as compared to the 100% treatment.

Figure 2: Volcano plot of transcripts log2 fold change vs. mean relative abundance, with significantly differently abundant (DA) transcripts highlighted by colors corresponding to their kingdom-level taxonomy for (A) roots and (B) rhizosphere soil. Blue: fungi, green: plant, red: bacteria, yellow: others, purple: unclassified, pink: archaea.

Figure 3: Stack bar chart comparing the taxonomical affiliations of transcripts associated to (A) fungi and to (B) bacteria in the roots and the rhizosphere across all samples vs. among transcripts positively (DA+) or negatively (DA-) differentially abundant. The transcripts not classified at this level ("NULL") were removed.

Figure 4: Stack bar chart comparing the functional affiliations (COG category) of transcripts associated in (A) roots and (B) the rhizosphere across all samples vs. among transcripts positively (DA+) or negatively (DA-) differentially abundant. The transcripts not classified at this level ("NULL") were removed.

Figure 5: Heatmaps for the top 50 most differentially abundant transcripts for (A) roots and (B) rhizosphere samples.

Figure 6: Upset plot showing the shared and unique genes between the root and rhizosphere transcripts positively (DA+) or negatively (DA-) differentially abundant.

1   **Table 1:** Mean abundance, log2 fold change (25% vs 100% precipitation), COG function and category and taxonomic affiliation for the top 50

2   most abundant genes with a P-value=0 in the roots.

Gene id	mean	log2	P-value	COG function	COG category	Class	Species
751566	805.00	9.57	0	Neutral trehalase	Carbohydrate transport and metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
148591	759.45	8.04	0	Dipeptide/tripeptide permease	Amino acid transport and metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
130818	659.18	8.17	0	Molecular chaperone, HSP90 family	Posttranslational modification, protein turnover, chaperones	*Agaricomycetes*	*Coprinopsis cinerea*
945131	441.27	8.04	0	NAD-dependent aldehyde dehydrogenases	Energy production and conversion	*Agaricomycetes*	*Hebeloma cylindrosporum*
560	359.64	7.75	0	Uncharacterized conserved protein	Function unknown	*Agaricomycetes*	*Coprinopsis cinerea*
934948	310.09	8.19	0	Hexokinase	Carbohydrate transport and metabolism	*Agaricomycetes*	*Gymnopilus dilepis*
1189377	300.36	8.14	0	Phosphatidylserine synthase	Lipid metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
956836	243.82	7.64	0	Nucleoside permease	Nucleotide transport and metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
121608	241.82	7.83	0	Amidases related to nicotinamidase	Secondary metabolites biosynthesis, transport, and catabolism	*Agaricomycetes*	*Coprinopsis cinerea*
92085	224.27	8.94	0	6-phosphogluconate dehydrogenase	Carbohydrate transport and metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
86792	213.27	8.87	0	Phosphatidylserine/phosphatidylglycerophosphate/cardiolipin synthases and related enzymes	Lipid metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
1107703	211.18	7.98	0	Polyketide synthase modules and related proteins	Secondary metabolites biosynthesis, transport, and catabolism	*Sordariomycetes*	*Stachybotrys chartarum*
2979	194.00	9.32	0	Thiamine pyrophosphate-requiring enzymes [acetolactate synthase, pyruvate dehydrogenase (cytochrome), glyoxylate carboligase, phosphonopyruvate decarboxylase]	Amino acid transport and metabolism / Coenzyme metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
183197	190.18	7.70	0	Glycerol uptake facilitator and related permeases (Major Intrinsic Protein Family)	Carbohydrate transport and metabolism	*Agaricomycetes*	*Mycena chlorophos*
14009	178.73	8.61	0	NAD-dependent aldehyde dehydrogenases	Energy production and conversion	*Agaricomycetes*	*Coprinopsis cinerea*
636334	165.64	10.09	0	Predicted aminopeptidases	General function prediction only	*Agaricomycetes*	*Panaeolus cyanescens*
196702	159.00	9.03	0	Nucleoside-diphosphate-sugar epimerases	Cell envelope biogenesis, outer membrane / Carbohydrate transport and metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
47528	150.45	8.37	0	Ca2+-binding actin-bundling protein fimbrin/plastin (EF-Hand superfamily)	Cytoskeleton	*Agaricomycetes*	*Coprinopsis cinerea*
657208	147.45	8.34	0	Amino acid transporters	Amino acid transport and metabolism	*Agaricomycetes*	*Dichomitus squalens*
166571	140.73	8.86	0	Predicted hydrolases or acyltransferases (alpha/beta hydrolase superfamily)	General function prediction only	*Sordariomycetes*	*Sodiomyces alkalinus*
488208	136.36	8.81	0	RNA-binding proteins (RRM domain)	General function prediction only	*Agaricomycetes*	*Coprinopsis cinerea*
222991	134.27	9.79	0	Nicotinic acid mononucleotide adenylyltransferase	Coenzyme metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
165517	107.55	-Inf	0	Ca2+-binding protein (EF-Hand superfamily)	Signal transduction mechanisms / Cytoskeleton / Cell division and chromosome partitioning / General function prediction only	*Liliopsida*	*Aegilops tauschii*
210176	102.55	8.40	0	O-Glycosyl hydrolase	Cell envelope biogenesis, outer membrane	*Sordariomycetes*	*Sodiomyces alkalinus*

202276	99.73	9.36	0	Cytochrome P450	Secondary metabolites biosynthesis, transport, and catabolism	*Agaricomycetes*	*Coprinopsis cinerea*
414450	98.55	9.34	0	NAD-dependent aldehyde dehydrogenases	Energy production and conversion	*Agaricomycetes*	*Laccaria amethystina*
660987	96.09	8.30	0	Asp-tRNAAsn/Glu-tRNAGln amidotransferase A subunit and related amidases	Translation, ribosomal structure and biogenesis	*Agaricomycetes*	*Coprinopsis cinerea*
1184958	79.64	9.03	0	Beta-glucanase/Beta-glucan synthetase	Carbohydrate transport and metabolism	*Agaricomycetes*	*Coprinopsis cinerea*
572300	78.27	-Inf	0	Predicted membrane protein	Function unknown	*Agaricomycetes*	*Panaeolus cyanescens*
210781	76.73	-Inf	0	Ribosomal protein S3	Translation, ribosomal structure and biogenesis	*Magnoliopsida*	*Populus trichocarpa*
720478	74.91	-Inf	0	FKBP-type peptidyl-prolyl cis-trans isomerases 1	Posttranslational modification, protein turnover, chaperones	*Magnoliopsida*	*Populus trichocarpa*
547100	64.45	-Inf	0	Chromosome segregation ATPases	Cell division and chromosome partitioning	*Clitellata*	*Hirudo medicinalis*
466738	61.55	-Inf	0	Dehydrogenases with different specificities (related to short-chain alcohol dehydrogenases)	Secondary metabolites biosynthesis, transport, and catabolism / General function prediction only	*Liliopsida*	*Aegilops tauschii*
1041684	60.91	Inf	0	Fucose permease	Carbohydrate transport and metabolism	*Dothideomycetes*	*Ascochyta rabiei*
1022565	60.55	-Inf	0	Putative multicopper oxidases	Secondary metabolites biosynthesis, transport, and catabolism	*Magnoliopsida*	*Populus trichocarpa*
82978	60.27	-Inf	0	Ribosomal protein S19E (S16A)	Translation, ribosomal structure and biogenesis	*Magnoliopsida*	*Populus trichocarpa*
140966	58.09	Inf	0	Uncharacterized stress protein (general stress protein 26)	General function prediction only	*Sordariomycetes*	*Stachybotrys chartarum*
539196	57.45	-Inf	0	Beta-fructosidases (levanase/invertase)	Carbohydrate transport and metabolism	*Liliopsida*	*Aegilops tauschii*
665062	55.18	-Inf	0	Cytochrome P450	Secondary metabolites biosynthesis, transport, and catabolism	*Liliopsida*	*Triticum aestivum*
200364	55.00	-Inf	0	Ubiquitin-protein ligase	Posttranslational modification, protein turnover, chaperones	*Liliopsida*	*Triticum aestivum*
690880	54.64	Inf	0	Peptidyl-prolyl cis-trans isomerase (rotamase) - cyclophilin family	Posttranslational modification, protein turnover, chaperones	*Insecta*	*Onthophagus taurus*
222916	54.55	-Inf	0	Secreted trypsin-like serine protease	Posttranslational modification, protein turnover, chaperones	*Arachnida*	*AcarusAcarus siro*
1044721	52.00	-Inf	0	FOG: WD40 repeat	General function prediction only	*Liliopsida*	*Aegilops tauschii*
715083	49.45	-Inf	0	Zn-dependent oligopeptidases	Amino acid transport and metabolism	*Liliopsida*	*Aegilops tauschii*
1245890	48.91	Inf	0	Deoxyribodipyrimidine photolyase	DNA replication, recombination, and repair	*Agaricomycetes*	*Leucoagaricus sp. SymC.cos*
247073	48.91	-Inf	0	Ca2+-binding protein (EF-Hand superfamily)	Signal transduction mechanisms / Cytoskeleton / Cell division and chromosome partitioning / General function prediction only	*NA*	*NA*
113358	48.82	Inf	0	Probable taurine catabolism dioxygenase	Secondary metabolites biosynthesis, transport, and catabolism	*Agaricomycetes*	*Coprinopsis cinerea*
258573	47.82	-Inf	0	Secreted trypsin-like serine protease	Posttranslational modification, protein turnover, chaperones	*Insecta*	*Aethina tumida*
1203022	47.00	-Inf	0	Uncharacterized conserved protein	Function unknown	*Liliopsida*	*Aegilops tauschii*
1010199	46.18	-Inf	0	Ribosomal protein HS6-type (S12/L30/L7a)	Translation, ribosomal structure and biogenesis	*Magnoliopsida*	*Populus trichocarpa*

3   **NA: Not Available; the transcript had no significant match to the database.**

1 **Table 2:** Mean abundance, log2 fold change (25% vs 100% precipitation), COG function and category and taxonomic affiliation for
2 the top 50 microbial genes with the lowest P-values in the rhizosphere.

gene ID	mean	log2	P-value	COG function	COG category	Class	Species
1150899	22.42	Inf	0	Monoamine oxidase	Amino acid transport and metabolism	Agaricomycetes	Leucoagaricus sp.
1238692	18.00	-Inf	0	Heme/copper-type cytochrome/quinol oxidase, subunit 3	Energy production and conversion	Clitellata	Enchytraeus albidus
190232	9.83	-Inf	0	Ferritin-like protein	Inorganic ion transport and metabolism	Insecta	Sipha flava
25017	12.17	-Inf	0	Zn-dependent protease with chaperone function	Posttranslational modification, protein turnover, chaperones	Gammaproteobacteria	Buchnera aphidicola
252300	20.67	-2.34	0	Cold shock proteins	Transcription	Deltaproteobacteria	Minicystis rosea
25926	15.83	Inf	0	Hexokinase	Carbohydrate transport and metabolism	Agaricomycetes	Gymnopilus dilepis
346718	25.00	-Inf	0	Myosin heavy chain	Cytoskeleton	Clitellata	Helobdella robusta
368534	11.33	-Inf	0	Ribosomal protein L20A (L18A)	Translation, ribosomal structure and biogenesis	Clitellata	Helobdella robusta
53704	38.00	-Inf	0	Cytochrome b subunit of the bc complex	Energy production and conversion	Clitellata	Enchytraeus cf. crypticus
685694	25.67	-1.87	0	Phage tail sheath protein FI	General function prediction only	Deltaproteobacteria	Haliangium ochraceum
90220	31.67	-2.15	0	Actin and related proteins	Cytoskeleton	NA	Dictyostelium discoideum
125923	554.00	-7.44	6.66E-16	Putative intracellular protease/amidase	General function prediction only	Microbotryomycetes	Rhodotorula graminis
625425	18.00	-2.05	1.89E-15	Uncharacterized protein conserved in bacteria	Function unknown	Deltaproteobacteria	Sorangium cellulosum
1093457	8.67	-Inf	3.11E-15	Glutathione S-transferase	Posttranslational modification, protein turnover, chaperones	Insecta	Aphis citricidus
900820	41.58	-1.77	3.55E-15	Predicted permease	General function prediction only	NA	Verrucomicrobia bacterium
988040	16.50	-3.51	5.00E-15	Translation elongation factor EF-1alpha (GTPase)	Translation, ribosomal structure and biogenesis	NA	Stramenopile sp.
438624	22.83	-1.49	1.17E-14	Outer membrane protein (porin)	Cell envelope biogenesis, outer membrane	Betaproteobacteria	Betaproteobacteria bacterium
1155936	36.50	-1.21	1.35E-14	Serine/threonine protein kinase	General function prediction only / Signal transduction mechanisms / Transcription / DNA replication, recombination, and repair	Deltaproteobacteria	Minicystis rosea
843326	19.50	-2.46	1.83E-14	Response regulator containing CheY-like receiver, AAA-type ATPase, and DNA-binding domains	Signal transduction mechanisms	NA	Nitrosopumilales archaeon
963336	20.17	-2.61	7.27E-14	ABC-type sugar transport system, periplasmic component	Carbohydrate transport and metabolism	Actinobacteria	Streptomyces sp.
920879	10.00	Inf	1.98E-13	Predicted integral membrane protein	Function unknown	Sphingobacteriia	Mucilaginibacter gotjawali
1143951	141.75	1.51	2.50E-13	Molecular chaperone (small heat shock protein)	Posttranslational modification, protein turnover, chaperones	Alphaproteobacteria	Skermanella aerolata
666947	19.08	-2.73	3.36E-13	Tfp pilus assembly protein PilE	Cell motility and secretion / Intracellular trafficking and secretion	NA	Candidatus Latescibacteria
998558	36.75	-2.31	4.82E-13	Type II secretory pathway, pseudopilin PulG	Cell motility and secretion / Intracellular trafficking and secretion	Spartobacteria	Chthoniobacter flavus
308612	18.92	-1.54	5.51E-13	Carbon dioxide concentrating mechanism/carboxysome shell protein	Secondary metabolites biosynthesis, transport, and catabolism / Energy production and conversion	Deltaproteobacteria	Haliangium ochraceum

5291	7.42	-3.34	1.34E-12	Predicted component of the type VI protein secretion system	Intracellular trafficking, secretion, and vesicular transport	Deltaproteobacteria	Minicystis rosea
91601	17.67	-2.29	2.07E-12	Type II secretory pathway, pseudopilin PulG	Cell motility and secretion / Intracellular trafficking and secretion	NA	Acidobacteria bacterium
993945	45.58	-1.31	2.20E-12	Uncharacterized conserved protein	Function unknown	Caldilineae	Caldilineae bacterium
731684	22.67	1.82	4.13E-12	Putative intracellular protease/amidase	General function prediction only	Actinobacteria	Micromonospora auratinigra
920153	13.50	-1.91	4.87E-12	DNA-directed RNA polymerase, alpha subunit/40 kD subunit	Transcription	Deltaproteobacteria	Haliangium sp.
389139	16.67	-1.66	7.06E-12	Flagellin and related hook-associated proteins	Cell motility and secretion	Alphaproteobacteria	Asticcacaulis taihuensis
647044	11.50	-2.74	9.44E-12	Glutamine synthetase	Amino acid transport and metabolism	Deltaproteobacteria	Geobacter sp.
1215469	48.33	-1.06	9.81E-12	Heme/copper-type cytochrome/quinol oxidases, subunit 1	Energy production and conversion	Deltaproteobacteria	Sorangiineae bacterium
758119	14.33	-1.92	1.03E-11	Nitrogen regulatory protein PII	Amino acid transport and metabolism	Deltaproteobacteria	Anaeromyxobacter dehalogen
1188088	7.67	-Inf	1.25E-11	Translation elongation factor EF-1alpha (GTPase)	Translation, ribosomal structure and biogenesis	Clitellata	Enchytraeus sp. Enc
372651	28.00	-1.24	1.32E-11	S-adenosylmethionine synthetase	Coenzyme metabolism	Deltaproteobacteria	Labilithrix luteola
1084785	7.67	-Inf	1.33E-11	Ribosomal protein L23	Translation, ribosomal structure and biogenesis	Polychaeta	Sipunculus nudus
377149	14.92	-2.15	1.43E-11	Biopolymer transport proteins	Intracellular trafficking and secretion	Deltaproteobacteria	Deltaproteobacteria bacteriur
1018589	19.17	-1.44	2.74E-11	F0F1-type ATP synthase, alpha subunit	Energy production and conversion	Clostridia	Butyricicoccus pullicaecorun
303242	13.33	-2.98	2.76E-11	Competence protein ComGC	Intracellular trafficking and secretion	NA	Acidobacteria bacterium
1232742	49.17	-2.26	2.90E-11	Type II secretory pathway, pseudopilin PulG	Cell motility and secretion / Intracellular trafficking and secretion	NA	Acidobacteria bacterium
1041075	20.58	-1.77	3.42E-11	Cytochrome c peroxidase	Inorganic ion transport and metabolism	NA	Planctomycetes bacterium
720588	9.50	-2.61	4.18E-11	DNA-directed RNA polymerase, beta subunit/140 kD subunit	Transcription	Deltaproteobacteria	Haliangium ochraceum
208638	48.17	-8.17	5.39E-11	Subtilisin-like serine proteases	Posttranslational modification, protein turnover, chaperones	Collembola	Orchesella cincta
754729	12.83	-2.58	6.88E-11	Type VI protein secretion system component Hcp (secreted cytotoxin)	Intracellular trafficking, secretion, and vesicular transport	Deltaproteobacteria	Chondromyces crocatus
1082904	17.33	-2.68	7.54E-11	Uncharacterized conserved protein	Function unknown	Sordariomycetes	Sporothrix schenckii
136368	51.50	-1.91	7.75E-11	Type II secretory pathway, component PulD	Cell motility and secretion / Intracellular trafficking and secretion	NA	Armatimonadetes bacterium
609055	9.50	-2.23	7.95E-11	Response regulators consisting of a CheY-like receiver domain and a winged-helix DNA-binding domain	Signal transduction mechanisms / Transcription	Deltaproteobacteria	Sorangium cellulosum
336735	35.50	-1.28	9.77E-11	Uncharacterized protein conserved in bacteria	Function unknown	Bacilli	Anaerobacillus isosaccharini
36391	17.17	-1.53	1.05E-10	GTPases - translation elongation factors	Translation, ribosomal structure and biogenesis	Deltaproteobacteria	Sorangium cellulosum

3   **NA: Not Available; the transcript had no significant match to the database.**