Eat or Be Eaten: Implications of potential exploitative competition between wolves and

humans across predator- savvy and -naive deer populations

ELLEN M. CANDLER*, *University of Minnesota, 135 Skok Hall, 2003 Upper Buford Circle, St.*

Paul, MN 55108, USA, ORCID: 0000-0002-5810-1942

STOTRA CHAKRABARTI, *Macalester College, Olin-Rice Science Center 211, St. Paul, MN*

55105, USA, ORCID: 0000-0001-8741-4780

WILLIAM J. SEVERUD, *South Dakota State University, Department of Natural Resource*

Management, Brookings, South Dakota USA 57007, USA, ORCID: 0000-0003-0150-

5986

JOSEPH K. BUMP, *University of Minnesota, 135 Skok Hall, 2003 Upper Buford Circle, St.*

Paul, MN 55108, USA, ORCID: 0000-0002-4369-7990

23 *Corresponding Author: belle130@umn.edu

Abstract

1. Recolonization of predators to their former ranges is becoming increasingly prevalent. Such recolonization places predators amongst their prey once again; the latter having lived without predation (from such predators) for considerable time. This renewed coexistence creates opportunities to explore predation ecology at both fundamental and applied levels.

2. We used a paired experimental design to investigate white-tailed deer risk allocation in the Upper and Lower Peninsulas (UP and LP) in Michigan, USA. Wolves are functionally absent in the LP, while deer in the UP coexist with a re-established wolf population. We treated 15 sites each in UP and LP with wolf olfactory cues, and observed deer vigilance, activity, and visitation rates at the interface of habitat covariates using remote cameras. Such a paired design across wolf versus no-wolf areas allowed us to examine indirect predation effects while accounting for confounding parameters such as the presence of other predators and human activity.

3. While wolf urine had no effect across most metrics in both UP and LP, we observed differences in deer activity in areas with versus without wolves. Sites treated with wolf urine in the UP showed a reduction in crepuscular deer activity, compared to control/novel-scent treated sites. Further, we observed a strong positive effect of vegetation cover on deer vigilance in these sites. This indicates that simulated predator cues likely affect deer vigilance more acutely in denser habitats, which presumably facilitates predation success. Such responses were however absent among deer in the LP that are presumably naïve towards wolf predation.

4. Where human and non-human predators hunt shared prey, such as in Michigan, predators

may constrain human hunting success by increasing deer vigilance. Hunters may avoid

such exploitative competition by choosing hunting/bait sites located in open areas. Our

results pertaining to fundamental predation ecology have strong applied implications that

can promote human-predator coexistence.

Key words: behavioral manipulation, human-wildlife conflict, olfactory cues, optimal foraging,

predator-prey interactions, predator recolonization, predator reintroduction

Introduction

Predators can alter prey populations directly through lethal offtake and/or behaviorally mediated

indirect effects (Kuijper et al., 2013; McNay et al., 2022; Palmer et al., 2021). While some prey

succumb, others avoid predation by changing their behavior or life-histories (Kohl et al., 2018;

Sabal et al., 2021). Predator-effects on prey are a fundamental focus in ecology, although,

dynamic predator-prey interactions in a fast-changing Anthropocene warrant more experimental

examinations of the underlying mechanisms of these direct and indirect effects (Parsons et al.,

2022; Scrosati, 2021). While predator populations are declining in many parts of the world

(Ripple et al., 2014), some areas are witnessing their comebacks (Gigliotti et al., 2019; Jarausch

et al., 2021; Scharf & Fernández, 2018). Such contrasting levels of predator presence create (and

re-create) intriguing interactions. Predators recolonizing former habitats may encounter

potentially naive prey that have lost anti-predatory strategies (Berger, 1999; Berger et al., 2001).

Anti-predator responses are defense strategies that prey species employ when they

perceive predation risk via visual, auditory, tactile, and/or olfactory stimuli. When at risk, prey

respond through multiple mechanisms either proactively, such as by increasing vigilance,

avoiding risky areas, social grouping, and birth synchrony, or reactively through evasion,

camouflage, and defense-herding (Flagel et al., 2016; Ims, 1990; Kohl et al., 2018; Mech, 2007;

Rowland et al., 2020). Successful anti-predatory responses, while providing survival benefits,

can have severe fitness costs for individuals. For example, ungulates that intensify vigilance as a

response to predator cues limited durations of feeding or avoided high-risk areas even if they are

rich with resources (Chamaillé-Jammes et al., 2014; Kuijper et al., 2013; Melchiors & Leslie,

1985; Sahlén et al., 2016; Wikenros et al., 2015). Furthermore, certain herbivores are known to

incur major reproductive costs while courting at breeding grounds that also attract a host of

predators (Cherry et al., 2016; Jhala & Isvaran, 2016). However, perception and response to

predation risk have significant decision-making implications that can result from a mix of

instinctive and learnt behaviors. Thus, the risk of predation can vary greatly amongst

experienced prey (prey that live under constant predation pressure) versus naive prey (prey that

live without or under very low predation pressure). For example, in areas where wolves (*Canis*

lupus) and grizzly bears (*Ursus horribilis*) had been extirpated, moose (*Alces alces*) did not

exhibit avoidance to predator olfactory or auditory cues, while those that had been exposed to

both predators showed avoidance (Berger et al., 2001). Similarly, species on predator-free

islands have been observed to 'adaptively forget' anti-predatory strategies, resulting in prey

naivety (Blumstein et al., 2004)

The Upper and Lower Peninsulas (UP and LP) of Michigan, USA offer a novel

opportunity to explore these drivers of predator-effects. In the LP, white-tailed deer (*Odocoileus*

virginianus) occur where wolves have been absent since the mid-1800s. Individual wolves have

been detected (although rarely) in the LP but remain functionally absent as predators of deer.

Alternatively, deer in the nearby UP coexist with a re-established wolf population since the mid-

2010s (O'Neil et al., 2017). Both peninsulas have a similar guild of other deer predators - black

bears *U. americanus*, coyotes *C. latrans*. Also, white-tailed deer are a subsistence species for

indigenous communities and a highly valued hunting species in this region. In both UP and LP,

deer are typically hunted by humans at specific locations where deer are food conditioned with

bait consisting of corn, fruits, or vegetables – creating artificially provisioned resource rich sites

on the landscape. These resource-rich but localized hunter-bait sites are intended to attract and

condition deer to revisit these sites frequently. According to behavioral optimality, prey species

need to balance the conflicting demands of efficient foraging and predator avoidance (Lima &

Bednekoff, 1999; Sih, 1980). Thus, deer behavior at these hunter-bait sites is expected to be

mediated by the perceived risk of predation because wolves are attracted to bait sites as well

(Bump et al., 2013), which in turn should be influenced by prey experience or naivety towards

predators.

Through a paired experimental design, that accounted for confounding effects of other

predators and human hunters, we tested for differences in risk perception by examining the

immediate behavioral effects of wolf olfactory cues on naive versus experienced deer at hunter-

bait sites. Since we expected no response from naive deer (LP deer; no functional wolf

population) to wolf scent (urine), we predicted:

1. deer that generally experience wolf predation (UP) would increase group size, vigilance

intensity, and occur less often at bait sites with wolf cues (scent),

2. the immediate effects of predator cues on deer vigilance will be further modulated by

habitat covariates such as vegetation cover at the sites, with higher vigilance to be

expected in denser habitats. Prey are often cautious, and tend to avoid 'risky' areas that

lack good visibility and offer ambushing opportunities for predators even where food is

abundant (Edwards, 1983; Kuijper et al., 2013; Ripple & Beschta, 2004),

3. predation-experienced deer would adjust their activity patterns to avoid crepuscular times

when wolves are most active, while naive deer would not make such adjustments. This

prediction is in accordance with the predation risk allocation hypothesis that suggests

temporal variation in predation risk severity affects foraging and vigilance behavior in

prey (Lima & Bednekoff, 1999).

Materials and Methods

Study Area

We conducted this study in Michigan's UP and LP from September 21, 2018 to November 10,

2018. The UP study area was entirely within the Hiawatha National Forest and was interspersed

with unpaved, US Forest Service roads. The bait sites in the LP study area were all located in

Mackinaw State Forest of Michigan, and the area was also interspersed with private agriculture,

forestland, and paved, secondary roads (Fig. 1). The main vegetation cover, derived from the

National Land Cover Database, at the UP study sites was evergreen (5 sites), woody wetland (3

sites), grassland (3 sites), deciduous (2 sites), and mixed forest (2 sites), while the LP study sites

were deciduous (6 sites), scrub/shrub (4 sites), grassland (3 sites), and evergreen (2 sites; Fig. 1)

(Homer et al., 2015).

Study Design

We established 30 deer bait sites between the two study areas, 15 in the UP and 15 in the LP

(Fig. 1). To mimic recreational hunters, we selected sites based on deer hunting desirability such

as habitat openness and/or proximity to a deer trail (Peterson, 2015). Sites were at least 1 km

away from each other to avoid duplication and dependence (Winke, 2012).

We baited each site with 3.8 liters of corn spread in a 3 by 3-meter- square plot, as per the

Michigan deer baiting regulations (Michigan Department of Natural Resources, 2018). For the

duration of the study, all sites were baited each week in order to maintain 3.8 liters of corn on the

ground to replicate actual hunter behavior and to reduce the confounding effect of bait

availability on deer behavior. We deployed one remote camera (Reconyx Hyperfire series or

Cuddeback E) on a tree 4 meters from the center of the bait site. Each camera was programmed

to take 5 burst images with no delay between triggers to record the number of deer present, their

posture, sex (when possible), and presence/absence of fawn(s). We replaced batteries and

memory cards weekly.

BACI Experimental Setup

We used a 10 week before-after-control-impact (BACI) design for the experimental treatment of

both areas. This BACI design was conducted so that all sites were paired with themselves with

all other factors (e.g., other predators, human visitation, environmental effects) remaining

comparable. This allowed us to tease apart the effects of the treatment. Before experimental

treatments, we collected data for one week preceding the six-week period to establish a baseline

condition of deer visitation to the bait sites (Stewart-Oaten et al., 1986). For the first 3 weeks

(September 28-October 20), we baited sites and recorded deer occurrence and behavior via

remote cameras with no scent treatments applied. For the following three weeks we continued to

bait and record deer after treating the sites with three scents. In each area, UP and LP, 5 sites

were treated each with wolf urine (predator treatment), lemon juice (novel scent; not a native

plant), or distilled water (control). We assigned site treatment randomly (Atkins et al., 2016;

Wikenros et al., 2015). When we treated sites in the field, we visited all 'control' sites in an area

first followed by the 'novel scent' sites, and finally 'experimental/wolf urine' sites to avoid

cross-site spread of scents. We also used different treatment application tools (i.e., pipets, travel

containers) for each treatment type to avoid cross contamination of treatments. At each site, we

applied 10 mL of the given treatment (experimental, novel, or control) by dripping it from a pipet

onto the bait site. This was done to mimic the amount and application of scent marking of a pair

of wolves (Peters & Mech, 1975).

Vegetation Cover

We also recorded predator concealment or hiding potential (horizontal cover) at each site to

account for site-specific differences in habitat openness - an index of perceived predation threat.

We used a 2 m cover pole with 20 sections, each 10 cm long, to measure predator hiding

potential at each site (Kuijper et al., 2014; Severud et al., 2019). From the center of each bait

pile, we measured the horizontal cover in each cardinal direction. While one person held the

cover pole 10-meters away from the center of the bait pile, another took a picture of the cover

pole with a camera stationed on a 1-meter high pole at the center of the bait pile, mimicking deer

visual height (Severud et al., 2019). We conducted this procedure twice during the study, once at

the beginning and once when treatment started, to account for change in hiding potential with

loss of leaves later in the year. For each of the 20 (10 cm) sections on the 2 m cover pole, we

estimated obscurement to the nearest 25% (Severud et al., 2019). We calculated a single mean

and standard error value for each site and each measurement time (beginning and middle of

study).

Analysis

Photo Analysis

Images were tagged in batches using the DigiKam photo editing software (Niedballa et

al., 2016). Each batch was defined by any set of images taken within 5 minutes of the prior

image. In each batch, for deer images only, we recorded if fawns were present, labelled all sexes

when possible, and tallied a total count. This increased the likelihood that we accounted for all

individuals that visited the bait pile and avoided miscounting any that were out of the camera

frame. For individual pictures within batches, in addition to generic behaviours (e.g., fighting,

nursing, foraging), we recorded the number of individuals with their heads above their shoulders,

indicating vigilance/state of alertness (Fig. 2; Flagel et al., 2016; Schuttler et al., 2017). We also

labelled batches for different species that were captured at site visits including squirrels

(*Sciurus* spp.), raccoons (*Procyon lotor*), wild turkeys (*Meleagris gallopavo*), and coyotes. We

did not detect additional scent marking by other species.

Treatment Impact on Deer Vigilance, Group Size and Visitation Rate

We explored the impact of different treatments in the two areas (UP and LP) on the

number of deer at each site, proportion of group that was vigilant, and the number of visits made

to the site. To account for the paired nature of our treatment design, we first averaged the

variable values for each site individually and calculated a difference in values before and during

treatment application. We then averaged across all sites within a given treatment. We also

calculated a vigilance intensity metric using equation 1. This metric includes both group

vigilance and event duration.

$$I = \frac{\Sum \frac{v}{g}}{e}$$

199   (Eq 1)

Where I is the vigilance intensity metric for a given event, v is the number of deer vigilant in a

single image, g is the group size in a single image, and e is the total time of the event. Hence,

vigilance intensity simply standardizes individual vigilance across different group sizes and time

spent in front of the camera. Similar to the previous analysis, we calculated a difference in

vigilance intensity before and after treatment for each treatment type in each peninsula.

Diel Activity

To analyze possible temporal variability in the use of bait sites in the UP and LP before

and after treatment, we used nonparametric kernel density estimation (Prugh et al., 2019; Wang

et al., 2015). We converted times to radians and used a kernel density estimator to create a

probability density distribution for each before or after period (Ridout & Linkie, 2009). We

calculated the proportion of temporal overlap between the two treatment periods for each

treatment in an area (Wang et al., 2015). We used a Δ_4 with a smoothing parameter of 1

because our sample size for all analyses was greater than 50 (Ridout & Linkie, 2009). We

conducted this analysis using the overlap package (Meredith & Ridout, 2018; Wang et al., 2015)

in R (R Development Core Team 2013).

We applied Watson's U2 statistic with the CircStats package to test for homogeneity

between the two samples of interest (i.e., test for/detect a statistically significant shift in the diel

pattern before and after treatment) (Lashley et al., 2018; Lund & Agostinelli, 2012). If deer

significantly shifted their temporal pattern between the two treatment periods, we expected

Watson's U2 statistic would be greater than the critical value (0.19 for an α value of 0.05) and P

< 0.05. In the UP, we predicted that there would be no shift in temporal visitation by deer at the

control and lemon treated sites (high Δ_4, U2 ≤ 0.19), while a shift to more nocturnal activity at

the wolf urine treated sites would be detected (lower Δ_4, U2 > 0.19; Kohl et al. 2018) to

potentially avoid wolves that are known to be typically less active at night (Kohl et al., 2018). In

the LP, we expected that we would not see a significant shift in any treatment sites because deer

in the LP are ostensibly naive to wolf predation.

Effect on Vigilance Behaviour

We fitted generalized linear models using pooled data from both the UP and LP to check

for the effect on overall vigilance intensity by treatment effect (before/after) and type

(predator/novel/control), area (UP or LP) and vegetation cover (i.e., predator hiding potential;

S1; Fležar et al., 2019; Prugh et al., 2019) through additive and interactive effects between

parameters of interest. Subsequently, we fitted models to examine the relationship between

vigilance intensity and vegetation cover at each treatment type within both the UP and LP before

and after treatment.

Using generalized linear model, we further examined the effects of sex, vegetation cover,

and presence of young on vigilance intensity, and included additive and interactive effects with

area to detect any difference between the UP and LP. We ranked models based on AICc.

Results

We obtained 286,436 images over seven weeks. After removing images taken during the pre-

experimental baseline week and images that did not contain deer, we retained 213,264 images for

analysis with 85,675 images taken in the LP and 127,589 images taken in the UP.

Impact on Deer Vigilance, Group Size and Visitation Rate

We observed a near 30% increase in group size at sites treated with wolf urine in the UP but this

increase translates to less than half a deer - an increase that is not ecologically meaningful.

Additionally, we observed a slight increase (<3%) in the number of visits at both lemon and

control treated sites in the UP and a marginal decrease (<4%) in the proportion of the group that

was vigilant at lemon treated sites in the LP (Table 1, Fig. 3). We also observed a decrease in

vigilance intensity of deer at wolf urine treated sites in the LP (Fig. 3). We did not detect a

significant change before and after treatments for any other treatment sites in either area for the

four response variables, i.e., the proportion of group that is vigilant, number of visits/visitation

rate, group size and vigilance intensity (Fig. 3).

Table 1. Perceived effect of wolf (*Canis lupus*) predation risk on deer behavior. Mean difference

before treatment period and during treatment period in group vigilance, group size, number of

visits, and vigilance intensity (sum of proportion of group vigilance in each event divided by

total time of event) from September to November 2018 in the Upper Peninsula (UP) and Lower

Peninsula (LP) Michigan, USA. *Differences before treatment period and during treatment

periods were significant if the 95% confidence interval did not include 0.

Area	Response variable	Treatment	Mean difference	95% CI
UP	Group vigilance	wolf	-0.044	-0.101 to 0.013
		lemon	-0.032	-0.082 to 0.017
		control	-0.020	-0.071 to 0.031
	Group size	wolf	0.293*	0.093 to 0.492
		lemon	0.238	-0.159 to 0.635
		control	0.166	-0.018 to 0.351
	Number of visits	wolf	0.011	-0.020 to 0.043
		lemon	0.024*	0.011 to 0.037
		control	0.007*	0.003 to 0.011
	Vigilance intensity	wolf	0.001	-0.021 to 0.024
		lemon	-0.008	-0.024 to 0.007
		control	-0.010	-0.023 to 0.003
LP	Group vigilance	wolf	-0.007	-0.083 to 0.068
		lemon	-0.035*	-0.044 to -0.025
		control	0.016*	0.002 to 0.034
	Group size	wolf	-0.341	-1.294 to 0.611
		lemon	0.133	-0.061 to 0.327
		control	-0.110	-0.626 to 0.405
	Number of visits	wolf	0.004	-0.019 to 0.012
		lemon	0.001	0.002 to 0.004
		control	0.000	-0.065 to 0.065
	Vigilance intensity	wolf	-0.028	-0.055 to 0.001
		lemon	0.002	-0.012 to 0.008
		control	0.008	-0.010 to 0.026

Diel Activity

Deer changed their diel activity in the LP and UP, and in all treatment types, before and after

treatment application (i.e., all U2 > 0.19). We did not detect a consistent directional pattern in the

temporal shifts for most sites. However, we did observe a difference between the UP and LP diel

patterns wherein the deer in the UP tended to reduce activity between 6:00 and 18:00 both before

and after treatment application while the deer in the LP did not have a consistent inactivity

period (Fig. 4). Additionally, deer at wolf-treated sites in the UP reduced bait site visitation in

the evening hours (i.e., 1800h; Fig. 4).

Impact of Vegetation Cover

Deer vigilance intensity increased by 1.3 times for each unit increase in vegetation cover

in the UP but not in the LP (Fig. 5; S1). Though vigilance intensity did not increase with

vegetation cover in the LP, the base vigilance intensity (model intercepts) increased significantly

after all treatment applications (S2).

We detected an increase in vigilance based on sex (males showing higher vigilance than

females) and when young were present, however, these effects did not vary between the UP and

the LP (Table 2). The most parsimonious model from our analysis indicates that the variation in

vigilance intensity was caused by difference in vegetation cover, modulated by area (Table 2,

model 1).

284　**Table 2.** Performance of nine individual, additive, and interactive models predicting vigilance intensity for white-tailed deer

285　(*Odocoileus virginianus*) from September to November 2018 in the Upper Peninsula (UP) and Lower Peninsula (LP) Michigan, USA.

Model	Predictor	Estimate	Standard Error	p	Null deviance	Residual deviance	AICc	Delta_ AICc
1. vigilance intensity ~ percent vegetation cover*area	Intercept	4.68E-02	4.08E-03	<0.05	4.61	4.47	-6553.27	0
	percent vegetation cover	1.82E-05	1.08E-04	0.87				
	area.UP	-2.07E-02	5.28E-03	<0.05				
	percent vegetation cover:area.UP	6.10E-04	1.35E-04	<0.05				
2. vigilance intensity ~ percent vegetation cover	Intercept	3.36E-02	2.58E-03	<0.05	4.61	4.52	-6536.6	16.66
	percent vegetation cover	4.09E-04	6.51E-05	<0.05				
3. vigilance intensity ~ percent vegetation cover+area	Intercept	3.31E-02	2.75E-03	<0.05	4.61	4.52	-6534.93	18.34
	percent vegetation cover	4.07E-04	6.52E-05	<0.05				
	area.UP	1.22E-03	2.12E-03	0.57				
4. vigilance intensity ~ sex*area	Intercept	0.04	4.39E-03	<0.05	4.61	4.54	-6515.33	37.94
	Female	6.62E-03	4.84E-03	0.17				
	Male	0.02	5.22E-03	<0.05				
	unknown sex	9.33E-03	8.10E-03	0.25				
	areaUP	1.86E-03	5.59E-03	0.74				
	Female:area.UP	6.14E-03	6.31E-03	0.33				
	Male:area.UP	-7.14E-03	6.93E-03	0.30				
	Unknown sex:area.UP	-8.92E-03	1.11E-02	0.42				
5. vigilance intensity ~ sex	Intercept	0.04	2.72E-03	<0.05	4.61	4.56	-6513.61	39.66
	Female	9.33E-03	3.09E-03	<0.05				
	Male	0.01	3.41E-03	<0.05				
	unknown sex	4.66E-03	5.54E-03	0.40				
6. vigilance intensity ~ sex+area	Intercept	0.04	3.03E-03	<0.05	4.61	4.56	-6513.24	40.03
	Female	9.70E-03	3.11E-03	<0.05				

	Male	0.02	3.42E-03	<0.05				
	unknown sex	4.97E-03	5.54E-03	0.37				
	areaUP	2.73E-03	2.14E-03	0.20				
7. vigilance intensity ~ young present	Intercept	0.05	1.24E-03	<0.05	4.61	4.60	-6501.93	51.34
	Young present	-5.17E-03	2.44E-03	<0.05				
8. vigilance intensity ~ young present+area	Intercept	0.05	1.64E-03	<0.05	4.61	4.60	-6500.7	52.57
	Young present	-5.16E-03	2.44E-03	<0.05				
	areaUP	1.89E-03	2.14E-03	0.38				
9. vigilance intensity ~ young present*area	Intercept	0.05	1.76E-03	<0.05	4.61	4.60	-6499.83	53.44
	Young present	-7.76E-03	3.45E-03	<0.05				
	areaUP	5.43E-04	2.48E-03	0.83				
	Young presentarea.UP	5.20E-03	4.89E-03	0.29				

Discussion

Predator absence for long durations can lead to the gradual erosion of anti-predatory

responses of prey (Berger, 1999; Berger et al., 2001). Subsequent recolonization of predators to

such landscapes raises questions about the response that naive prey will have once predators

return, particularly in prey populations valued and pursued by human hunters. We investigated

the effects of simulated predator presence on experienced versus naive deer at food-rich hunter-

bait sites in Michigan. We did not find a significant effect of wolf scent treatment on deer

vigilance, number of visits, or vigilance intensity among predator-savvy/experienced deer in the

UP but did note a slight increase in group size, though the difference was not ecologically

meaningful. A similar lack of response to wolf urine has been documented in deer populations in

the Netherlands and Poland where little support was found for effects of predator scents in both

predator savvy and predator naive prey populations even when three times the volume of wolf

urine was used (van Ginkel, Smit, et al., 2019). This lack of reaction to wolf scent cues in both

areas may also be a result of decreasing response over time to the wolf urine as it degrades

(Bytheway et al., 2013; Kuijper et al., 2013; Wikenros et al., 2015). Introducing cues that

indicate immediate wolf proximity, such as howling or visual, are likely to produce different

results (Liley & Creel, 2008; van Ginkel, Smit, et al., 2019). The absence of any response to wolf

urine may also be a result of predator hunting strategies. Cursorial predators might fail to evoke

strong anti-predatory responses in deer, as opposed to pronounced effects as observed in the case

of cues from ambush predators (Preisser et al., 2007; Sunde et al., 2022).

We also observed a significant shift in diel patterns at all sites before and after treatment

but there was no consistency to the shift in activity for most treatment sites. However, deer at

sites treated with wolf scent in the UP (where they are sympatric with wolves) reduced their

visitation in the evening hours. This finding is in accordance with other research that suggests

that ungulate prey shift their activity to avoid riskier times (Gehr et al., 2018; Kohl et al., 2018).

For example, deer in Minnesota, USA demonstrated a more diurnal pattern under the effects of

wolf scent to likely avoid the riskier crepuscular times when wolves tend to hunt (Palmer et al.,

2021). It is also noteworthy that before and after treatment application, all UP sites exhibited a

decrease in deer activity at bait sites between 6:00 and 18:00 hours, while the LP sites had

inconsistent patterns. This suggests that deer in the UP are perhaps avoiding peak wolf

activity/hunting hours regardless of predator cues at sites, while LP deer do not have as large of

an inactivity period, likely because they do not coexist with wolves (S2; Kohl et al., 2018).

Our results mirror findings from similar studies that found that deer vigilance was

dependent on sex and fawn presence (Cherry et al., 2015; Gulsby et al., 2018; Higdon et al.,

2019; Lashley et al., 2014). When young were present, vigilance increased regardless of wolf

presence on the landscape. Similarly, sex played an important role where males showed more

vigilance than females (Table 2). However, we found that vegetation cover was the largest

contributor to the variance in vigilance, and this difference in vigilance was modulated by wolf

presence or absence in that area (Cherry et al., 2015; Olson et al., 2021).

Our results show a strong effect of vegetation cover on deer vigilance intensity where

wolves are established. Increase in vegetation cover at the UP sites were characterized by intense

vigilance activity in deer, regardless of the treatment. Denser vegetation likely represents more

predator hiding potential and difficult escape routes, likely causing an increase in deer vigilance

regardless of olfactory cues at the site (Fig. 5; Dellinger et al., 2019; van Ginkel, Kuijper, et al.,

2019; Wikenros et al., 2015). Research comparing white-tailed deer and mule deer (*Odocoileus*

hemionus) in Oregon's expanding wolf range reported that white-tailed deer did not typically

avoid areas with wolves but did favor areas with less dense vegetation that facilitates an easy

escape because they relied on flight and early detection of predators (Dellinger et al., 2019).

Other studies also demonstrated the importance of escape routes for white-tailed deer, indicating

that white-tailed deer flee rather than stand their ground in the presence of a predator (Lingle &

Pellis, 2002). This phenomenon has also been noted in other species such as chacma baboons

(*Papio ursinus*) in Gorongosa National Park, Mozambique where vocalization and vigilance

were heightened in denser habitats (Hammond et al., 2022). Our findings further highlight the

importance of considering prey behavior and escape strategies, along with habitat characteristics

for a mechanistic understanding of predator-effects.

While other studies, investigating questions similar to ours, have found that sex and

presence of young were the strongest predictors of deer vigilance, several key aspects in our

research differed from these studies, and might explain our subtly different results (Cherry et al.,

2015, 2017; Gulsby et al., 2018; Higdon et al., 2019; Lashley et al., 2014). In the cited studies,

vegetation was standardized or the time of year the research was conducted was different than

our study period. If we were to conduct this experiment again during spring when fawns are

more susceptible to coyote predation, the influence of vegetation may be masked with intensified

vigilance because coyotes occur in both areas. Consequently, a season-specific antipredator

response, recorded in many of the aforementioned studies, could potentially be present in our

study system (Cherry et al., 2017; Gulsby et al., 2018).

Though vegetation cover did not have a significant effect at LP sites, deer at sites treated

with wolf urine demonstrated higher baseline vigilance intensity after treatment was applied but

intensity did not vary across vegetation cover. This likely suggests that naive deer may

demonstrate an innate fear of wolves, also reported in East Limestone Island, British Columbia,

Canada (Chamaillé-Jammes et al., 2014). However, when the novel scent (lemon) and control

sites were assessed in the same context, we observed a similar pattern indicating that something

other than predator scent might be affecting baseline vigilance later in the study. Because the

second half of the study was conducted during archery hunting season, this shift in vigilance may

indicate a response to increased human presence (Benhaiem et al., 2008). A study in Poland

found that red deer (*Cervus elaphus*) demonstrated heightened vigilance during hunting season

when hunters were likely pursuing them (Proudman et al., 2020). Hunting seasons occurred

simultaneously in the UP and LP making our study an appropriate comparison between the two

locations. Similarly, other predators (e.g., black bears) occur in both areas, reducing any

confounding effects such predators might have had. This shift in baseline vigilance could also be

attributed to the onset of the deer breeding season, an observation made in other white-tailed deer

studies (Cherry et al., 2015; Schuttler et al., 2016).

Our findings offer partial support to the predator risk allocation hypothesis by

demonstrating a shift in activity by prey with predator experience to a less risky feeding time -

when predators are less active (Kohl et al., 2018; Lima & Bednekoff, 1999). This hypothesis

may also explain the lack of difference in vigilance detected in the UP-deer population in

relation to the wolf urine treatment. Because this population is under constant predation pressure

from wolves and frequently encounters wolf cues, deer in the UP may demonstrate a change of

vigilance especially at food rich locations such as hunter-bait sites, in order to take advantage of

the food source as quickly as possible. Deer in the UP are also likely, as we observed, to show

more vigilance in areas with higher predation potential, such as with dense vegetation cover

(Dellinger et al., 2019; Kunkel & Pletscher, 2001).

Finally, through our experimental study we highlight the importance of considering both

spatial and temporal factors when understanding predator effects on experienced versus naive

prey. Past research has suggested that naive prey do not maintain the innate ability to react to

predator olfactory cues but do have the ability to learn and adjust to predators within one

generation (Berger et al., 2001; Steindler et al., 2020). Though predator olfactory cues did not

have a significant effect on most variables we investigated, we observed a strong predator effect

depending on vegetation cover that ostensibly modifies risk.

The use of a BACI experimental method in a predator free and predator savvy population

allowed us to account for the complexity of the systems and make a few key conclusions: 1.

Wolf scent had little effect on number of visits, group vigilance, or group size but it did impact

the time that predator-savvy deer visited bait sites (deer decreased visitation in the evenings), 2.

The overall diel patterns of predator-savvy deer generally avoided bait visitation between 6:00

and 18:00, whereas predator naïve deer did not show a similar avoidance pattern, and 3. As

predator hiding potential (vegetation cover) increased, so did vigilance intensity among predator-

savvy deer, but not in the predator-naïve deer. Because the only predator difference between the

two populations is wolves, we can suppose this difference in vigilance intensity is due to wolf

presence.

The recreational hunting tradition of baiting is controversial inside and outside the

hunting community. Baiting increases the opportunity for a successful shot and more immediate

kill, but also creates a predictable food source that concentrates deer and deer scent, which in

turn attracts wolves (Hughes et al., 2010). As large predators continue to recolonize their former

ranges, they will inevitably overlap with human hunters (Bled et al., 2022; Grima et al., 2021).

Our current research can shed light on the impact a predator has on a predator savvy and predator

naive prey population as it relates to hunting by humans. We show that predator cues can affect

deer vigilance at bait-sites with denser vegetation, seemingly making deer cautious and difficult

to hunt, thereby potentially affecting the experience and satisfaction reported in hunter surveys.

However, we strongly urge readers, managers, and the hunting community to interpret our results

cautiously and constructively. Our results can be formative in aiding carnivore-human

coexistence wherein human hunters can have more success if they potentially choose not to

bait/hunt deer at vegetatively dense sites.

Author Contributions

Ellen M. Candler conceived the idea, developed the sampling design and methodology,

preformed data collection, conducted the formal analysis, oversaw project administration,

acquired resources, supervised field work, validated, and visualized data, and wrote, reviewed,

and edited the original draft. Stotra Chakrabarti helped develop the sampling design, and

methodology, analyzed, validated, and visualized data, wrote, reviewed and edited drafts;

William J. Severud developed the sampling design and methodology, reviewed and edited drafts,

Joseph K. Bump conceived the idea, developed the sampling design, was in charge of project

administration and supervision, reviewed and edited drafts, acquired funding and resources.

Statement on inclusion

The team of authors includes members representing both the Global North and South countries,

including scientists based in the country where the study was carried out. Furthermore, both the

lead and the last authors have significant association with the state where this study was

conducted. All authors were engaged early on with the research and study design to ensure that

the diverse sets of perspectives they represent was considered from the onset. Whenever

relevant, literature published by scientists from the region as well as from other parts of the

world were cited.

Acknowledgements

This material is based upon work supported by the National Science Foundation Graduate

Research Fellowship Program under Grant No. 00039218. Any opinions, findings, and

conclusions or recommendations expressed in this material are those of the authors and do not

necessarily reflect the views of the National Science Foundation. This research was also

supported by grants from the National Science Foundation to J. K. Bump (NSF ID#1545611,

NSF ID#1556676). We thank M. Dean, K. Rajski, and A. Schubel for field assistance.

Conflict of Interests

No conflict of interests.

Data Availability Statement

Data has been uploaded to the Dryad Digital Repository.

https://datadryad.org/stash/share/LhDJMgwhDbAX3LDhATW7XPgJHpHdfZQRIAIB5lTfwQo

Literature Cited

Atkins, R., Blumstein, D. T., Moseby, K. E., West, R., Hyatt, M., & Letnic, M. (2016). Deep
evolutionary experience explains mammalian responses to predators. *Behavioral Ecology*
and Sociobiology, *70*(10), 1755–1763. https://doi.org/10.1007/s00265-016-2181-4

Benhaiem, S., Delon, M., Lourtet, B., Cargnelutti, B., Aulagnier, S., Hewison, A. J. M.,
Morellet, N., & Verheyden, H. (2008). Hunting increases vigilance levels in roe deer and
modifies feeding site selection. *Animal Behaviour*, *76*(3), 611–618.
https://doi.org/10.1016/j.anbehav.2008.03.012

Berger, J. (1999). Anthropogenic extinction of top carnivores and interspecific animal behaviour:
Implications of the rapid decoupling of a web involving wolves, bears, moose and ravens.
Proceedings of the Royal Society of London. Series B: Biological Sciences, *266*(1435),
2261–2267. https://doi.org/10.1098/rspb.1999.0917

Berger, J., Swenson, J. E., & Persson, I.-L. (2001). Recolonizing carnivores and naïve prey:
Conservation lessons from pleistocene extinctions. *Science*, *291*(5506), 1036–1039.
https://doi.org/10.1126/science.1056466

Bled, F., Cherry, M. J., Garrison, E. P., Miller, K. V., Conner, L. M., Abernathy, H. N.,
454  Ellsworth, W. H., Margenau, L. L. S., Crawford, D. A., Engebretsen, K. N., Kelly, B. D.,
455  Shindle, D. B., & Chandler, R. B. (2022). Balancing carnivore conservation and sustainable
456  hunting of a key prey species: A case study on the Florida panther and white-tailed deer.
457  *Journal of Applied Ecology*, *n/a*(n/a). https://doi.org/10.1111/1365-2664.14201

458 Blumstein, D. T., Daniel, J. C., & Springett, B. P. (2004). A test of the multi-predator hypothesis:
459  Rapid loss of antipredator behavior after 130 years of isolation. *Ethology*, *110*(11), 919–934.
460  https://doi.org/10.1111/j.1439-0310.2004.01033.x

461 Bump, J. K., Murawski, C. M., Kartano, L. M., Beyer, D. E. Jr., & Roell, B. J. (2013). Bear-
462  baiting may exacerbate wolf-hunting dog conflict. *PLoS ONE*, *8*(4), e61708.
463  https://doi.org/10.1371/journal.pone.0061708

464 Bytheway, J. P., Carthey, A. J. R., & Banks, P. B. (2013). Risk vs. reward: How predators and
465  prey respond to aging olfactory cues. *Behavioral Ecology and Sociobiology*, *67*(5), 715–725.
466  https://doi.org/10.1007/s00265-013-1494-9

467 Chamaillé-Jammes, S., Malcuit, H., Saout, S. L., & Martin, J.-L. (2014). Innate threat-sensitive
468  foraging: Black-tailed deer remain more fearful of wolf than of the less dangerous black bear
469  even after 100 years of wolf absence. *Oecologia*, *174*(4), 1151–1158.
470  https://doi.org/10.1007/s00442-013-2843-0

471 Cherry, M. J., Conner, L. M., & Warren, R. J. (2015). Effects of predation risk and group
472  dynamics on white-tailed deer foraging behavior in a longleaf pine savanna. *Behavioral
473  Ecology*, *26*(4), 1091–1099. https://doi.org/10.1093/beheco/arv054

474 Cherry, M. J., Morgan, K. E., Rutledge, B. T., Conner, L. M., & Warren, R. J. (2016). Can
475  coyote predation risk induce reproduction suppression in white-tailed deer? *Ecosphere*,
476  *7*(10), e01481. https://doi.org/10.1002/ecs2.1481

477 Cherry, M. J., Warren, R. J., & Conner, L. M. (2017). Fire-mediated foraging tradeoffs in white-
478  tailed deer. *Ecosphere*, *8*(4), e01784. https://doi.org/10.1002/ecs2.1784

479 Dellinger, J. A., Shores, C. R., Craig, A., Heithaus, M. R., Ripple, W. J., & Wirsing, A. J.
480  (2019). Habitat use of sympatric prey suggests divergent anti-predator responses to
481  recolonizing gray wolves. *Oecologia*, *189*(2), 487–500. https://doi.org/10.1007/s00442-018-
482  4323-z

483 Edwards, J. (1983). Diet shifts in moose due to predator avoidance. *Oecologia*, *60*(2), 185–189.
484  https://doi.org/10.1007/BF00379520

485 Flagel, D. G., Belovsky, G. E., & Beyer, D. E. (2016). Natural and experimental tests of trophic
486  cascades: Gray wolves and white-tailed deer in a Great Lakes forest. *Oecologia*, *180*(4),
487  1183–1194. https://doi.org/10.1007/s00442-015-3515-z

488 Fležar, U., le Roux, E., Kerley, G. I. H., Kuijper, D. P. J., te Beest, M., Druce, D. J., Prinsloo, D.,
489  & Cromsigt, J. P. G. M. (2019). Simulated elephant-induced habitat changes can create

dynamic landscapes of fear. *Biological Conservation*, *237*, 267–279.
https://doi.org/10.1016/j.biocon.2019.07.012

Gehr, B., Hofer, E. J., Ryser, A., Vimercati, E., Vogt, K., & Keller, L. F. (2018). Evidence for
nonconsumptive effects from a large predator in an ungulate prey? *Behavioral Ecology*,
29(3), 724–735. https://doi.org/10.1093/beheco/ary031

Gigliotti, L. C., Matchett, M. R., & Jachowski, D. S. (2019). Mountain lions on the prairie:
Habitat selection by recolonizing mountain lions at the edge of their range. *Restoration*
Ecology, *27*(5), 1032–1040. https://doi.org/10.1111/rec.12952

Grima, N., Brainard, J., & Fisher, B. (2021). Are wolves welcome? Hunters' attitudes towards
wolves in Vermont, USA. *Oryx*, *55*(2), 262–267.
https://doi.org/10.1017/S0030605319000061

Gulsby, W. D., Cherry, M. J., Johnson, J. T., Conner, L. M., & Miller, K. V. (2018). Behavioral
response of white-tailed deer to coyote predation risk. *Ecosphere*, *9*(3), e02141.
https://doi.org/10.1002/ecs2.2141

Hammond, P., Lewis-Bevan, L., Biro, D., & Carvalho, S. (2022). Risk perception and
terrestriality in primates: A quasi-experiment through habituation of chacma baboons (Papio
ursinus) in Gorongosa National Park, Mozambique. *American Journal of Biological*
Anthropology, *179*(1). https://doi.org/10.1002/ajpa.24567

Higdon, S. D., Diggins, C. A., Cherry, M. J., & Ford, W. M. (2019). Activity patterns and
temporal predator avoidance of white-tailed deer (Odocoileus virginianus) during the
fawning season. *Journal of Ethology*, *37*(3), 283–290. https://doi.org/10.1007/s10164-019-
00599-1

Homer, C., Dewitz, J., Yang, L., Jin, S., Danielson, P., Coulston, J., Herold, N., Wickham, J., &
Megown, K. (2015). Completion of the 2011 national land cover database for the
conterminous United States – representing a decade of land cover change information.
PHOTOGRAMMETRIC ENGINEERING, 11.

Hughes, N. K., Price, C. J., & Banks, P. B. (2010). Predators are attracted to the olfactory signals
of prey. *PLOS ONE*, *5*(9), e13114. https://doi.org/10.1371/journal.pone.0013114

Ims, R. A. (1990). On the adaptive value of reproductive synchrony as a predator-swamping
strategy. *The American Naturalist*, *136*(4), 485–498. https://doi.org/10.1086/285109

Jarausch, A., Harms, V., Kluth, G., Reinhardt, I., & Nowak, C. (2021). How the west was won:
Genetic reconstruction of rapid wolf recolonization into Germany's anthropogenic
landscapes. *Heredity*, *127*(1), Article 1. https://doi.org/10.1038/s41437-021-00429-6

Jhala, Y. V., & Isvaran, K. (2016). Behavioural ecology of a grassland antelope, the blackbuck
antilope cervicapra: Linking habitat, ecology and behaviour. In F. S. Ahrestani & M.
Sankaran (Eds.), *The Ecology of Large Herbivores in South and Southeast Asia* (pp. 151–
176). Springer Netherlands. https://doi.org/10.1007/978-94-017-7570-0_6

Kohl, M. T., Stahler, D. R., Metz, M. C., Forester, J. D., Kauffman, M. J., Varley, N., White, P.
J., Smith, D. W., & MacNulty, D. R. (2018). Diel predator activity drives a dynamic
landscape of fear. *Ecological Monographs*, *0*(0), 1–15. https://doi.org/10.1002/ecm.1313

Kuijper, D. P. J., de Kleine C., Churski M., van Hooft P., Bubnicki J., & Jędrzejewska B. (2013).
Landscape of fear in Europe: Wolves affect spatial patterns of ungulate browsing in
Białowieża Primeval Forest, Poland. *Ecography*, *36*(12), 1263–1275.
https://doi.org/10.1111/j.1600-0587.2013.00266.x

Kuijper, D. P. J., Verwijmeren, M., Churski, M., Zbyryt, A., Schmidt, K., Jędrzejewska, B., &
Smit, C. (2014). What cues do ungulates use to assess predation risk in dense temperate
forests? *PLOS ONE*, *9*(1), e84607. https://doi.org/10.1371/journal.pone.0084607

Kunkel, K., & Pletscher, D. H. (2001). Winter Hunting Patterns of Wolves in and Near Glacier
National Park, Montana. *The Journal of Wildlife Management*, *65*(3), 520–530.
https://doi.org/10.2307/3803105

Lashley, M. A., Chitwood, M. C., Biggerstaff, M. T., Morina, D. L., Moorman, C. E., &
DePerno, C. S. (2014). White-Tailed Deer Vigilance: The Influence of Social and
Environmental Factors. *PLOS ONE*, *9*(3), e90652.
https://doi.org/10.1371/journal.pone.0090652

Lashley, M. A., Cove, M. V., Chitwood, M. C., Penido, G., Gardner, B., DePerno, C. S., &
Moorman, C. E. (2018). Estimating wildlife activity curves: Comparison of methods and
sample size. *Scientific Reports*, *8*(1), 1–11. https://doi.org/10.1038/s41598-018-22638-6

Liley, S., & Creel, S. (2008). What best explains vigilance in elk: Characteristics of prey,
predators, or the environment? *Behavioral Ecology*, *19*(2), 245–254.
https://doi.org/10.1093/beheco/arm116

Lima, S. L., & Bednekoff, P. A. (1999). Temporal variation in danger drives antipredator
behavior: The predation risk allocation hypothesis. *American Naturalist*, *153*(6), 649–659.
https://doi.org/10.1086/303202

Lingle, S., & Pellis, S. (2002). Fight or flight? Antipredator behavior and the escalation of coyote
encounters with deer. *Oecologia*, *131*(1), 154–164. https://doi.org/10.1007/s00442-001-
0858-4

Lund, U., & Agostinelli, C. (2012). *CircStats: Circular statistics, from "topics in circular
statistics" (2001)* (R-Package 0.2-4) [Computer software].

McNay, R. S., Lamb, C. T., Giguere, L., Williams, S. H., Martin, H., Sutherland, G. D., &
Hebblewhite, M. (2022). Demographic responses of nearly extirpated endangered mountain
caribou to recovery actions in Central British Columbia. *Ecological Applications*, *n/a*(n/a),
e2580. https://doi.org/10.1002/eap.2580

Mech, L. D. (2007). Possible use of foresight, understanding, and planning by wolves hunting
muskoxen. *Arctic*, *60*(2), 145–149.

Melchiors, M. A., & Leslie, C. A. (1985). Effectiveness of Predator Fecal Odors as Black-Tailed
Deer Repellents. *The Journal of Wildlife Management*, *49*(2), 358–362.
https://doi.org/10.2307/3801531

Meredith, M., & Ridout, M. (2018). *overlap: Estimates of coefficient of overlapping for animal*
activity patterns (0.3.2) [Computer software].

Michigan Department of Natural Resources. (2018). *Michigan Hunting Digest*.

Niedballa, J., Sollmann, R., Courtiol, A., & Wilting, A. (2016). camtrapR: An R package for
efficient camera trap data management. *Methods in Ecology and Evolution*, *7*(12), 1457–
1462. https://doi.org/10.1111/2041-210X.12600

Olson, L. O., Van Deelen, T. R., Storm, D. J., & Crimmins, S. M. (2021). Understanding
environmental patterns of canid predation on white-tailed deer (Odocoileus virginianus).
Canadian Journal of Zoology, *99*(10), 912–920. https://doi.org/10.1139/cjz-2021-0024

O'Neil, S. T., Bump, J. K., & Beyer, D. E. (2017). Spatially varying density dependence drives a
shifting mosaic of survival in a recovering apex predator (*Canis lupus*). *Ecology and*
Evolution, *7*(22), 9518–9530. https://doi.org/10.1002/ece3.3463

Palmer, M. S., Portales-Reyes, C., Potter, C., Mech, L. D., & Isbell, F. (2021). Behaviorally-
mediated trophic cascade attenuated by prey use of risky places at safe times. *Oecologia*,
195(1), 235–248. https://doi.org/10.1007/s00442-020-04816-4

Parsons, M. A., Newsome, T. M., & Young, J. K. (2022). The consequences of predators without
prey. *Frontiers in Ecology and the Environment*, *20*(1), 31–39.
https://doi.org/10.1002/fee.2419

Peters, R. P., & Mech, L. D. (1975). Scent-Marking in wolves: Radio-tracking of wolf packs has
provided definite evidence that olfactory sign is used for territory maintenance and may
serve for other forms of communication within the pack as well. *American Scientist*, *63*(6),
628–637.

Peterson, T. J. (2015, September 9). *5 best funnel & pinch-point setups for hunting whitetails*.
Game & Fish. https://www.gameandfishmag.com/editorial/5-best-funnel-pinch-point-setups-
for-hunting-whitetails/191246

Preisser, E. L., Orrock, J. L., & Schmitz, O. J. (2007). Predator Hunting Mode and Habitat
Domain Alter Nonconsumptive Effects in Predator–Prey Interactions. *Ecology*, *88*(11),
2744–2751. https://doi.org/10.1890/07-0260.1

Proudman, N. J., Churski, M., Bubnicki, J. W., Nilsson, J.-Å., Kuijper, D. P. J., Proudman, N. J.,
Churski, M., Bubnicki, J. W., Nilsson, J.-Å., & Kuijper, D. P. J. (2020). Red deer allocate
vigilance differently in response to spatio-temporal patterns of risk from human hunters and
wolves. *Wildlife Research*, *48*(2), 163–174. https://doi.org/10.1071/WR20059

Prugh, L. R., Sivy, K. J., Mahoney, P. J., Ganz, T. R., Ditmer, M. A., van de Kerk, M., Gilbert,
S. L., & Montgomery, R. A. (2019). Designing studies of predation risk for improved
inference in carnivore-ungulate systems. *Biological Conservation*, *232*, 194–207.
https://doi.org/10.1016/j.biocon.2019.02.011

Ridout, M. S., & Linkie, M. (2009). Estimating overlap of daily activity patterns from camera
trap data. *Journal of Agricultural, Biological, and Environmental Statistics*, *14*(3), 322–337.
https://doi.org/10.1198/jabes.2009.08038

Ripple, W. J., & Beschta, R. L. (2004). Wolves and the ecology of fear: Can predation risk
structure ecosystems? *BioScience*, *54*(8), 755–766. https://doi.org/10.1641/0006-
3568(2004)054[0755:WATEOF]2.0.CO;2

Ripple, W. J., Estes, J. A., Beschta, R. L., Wilmers, C. C., Ritchie, E. G., Hebblewhite, M.,
Berger, J., Elmhagen, B., Letnic, M., Nelson, M. P., Schmitz, O. J., Smith, D. W., Wallach,
A. D., & Wirsing, A. J. (2014). Status and ecological effects of the world's largest
carnivores. *Science*, *343*(6167), 1241484. https://doi.org/10.1126/science.1241484

Rowland, H. M., Burriss, R. P., & Skelhorn, J. (2020). The antipredator benefits of postural
camouflage in peppered moth caterpillars. *Scientific Reports*, *10*(1), Article 1.
https://doi.org/10.1038/s41598-020-78686-4

Sabal, M. C., Boyce, M. S., Charpentier, C. L., Furey, N. B., Luhring, T. M., Martin, H. W.,
Melnychuk, M. C., Srygley, R. B., Wagner, C. M., Wirsing, A. J., Ydenberg, R. C., &
Palkovacs, E. P. (2021). Predation landscapes influence migratory prey ecology and
evolution. *Trends in Ecology & Evolution*, *36*(8), 737–749.
https://doi.org/10.1016/j.tree.2021.04.010

Sahlén, E., Noell, S., DePerno, C. S., Kindberg, J., Spong, G., & Cromsigt, J. P. G. M. (2016).
Phantoms of the forest: Legacy risk effects of a regionally extinct large carnivore. *Ecology*
and Evolution, *6*(3), 791–799. https://doi.org/10.1002/ece3.1866

Scharf, A. K., & Fernández, N. (2018). Up-scaling local-habitat models for large-scale
conservation: Assessing suitable areas for the brown bear comeback in Europe. *Diversity and*
Distributions, *24*(11), 1573–1582. https://doi.org/10.1111/ddi.12796

Schuttler, S. G., Parsons, A. W., Forrester, T. D., Baker, M. C., McShea, W. J., Costello, R., &
Kays, R. (2016). Deer on the lookout: How hunting, hiking and coyotes affect white-tailed
deer vigilance. *Journal of Zoology*, *301*(4), 320–327. https://doi.org/10.1111/jzo.12416

Scrosati, R. A. (2021). Nonconsumptive predator effects on prey demography: Recent advances
using intertidal invertebrates. *Frontiers in Ecology and Evolution*, *9*.
https://www.frontiersin.org/article/10.3389/fevo.2021.626869

Severud, W. J., DelGiudice, G. D., & Obermoller, T. R. (2019). Association of moose parturition
and post-parturition habitat with calf survival. *The Journal of Wildlife Management*, *83*(1),
175–183. https://doi.org/10.1002/jwmg.21570

Sih, A. (1980). Optimal behavior: Can foragers balance two conflicting demands? *Science*, *210*(4473), 1041–1043. https://doi.org/10.1126/science.210.4473.1041

Steindler, L. A., Blumstein, D. T., West, R., Moseby, K. E., & Letnic, M. (2020). Exposure to a novel predator induces visual predator recognition by naïve prey. *Behavioral Ecology and Sociobiology*, *74*(8), 102. https://doi.org/10.1007/s00265-020-02884-3

Stewart-Oaten, A., Murdoch, W. W., & Parker, K. R. (1986). Environmental impact assessment: "Pseudoreplication" in time? *Ecology*, *67*(4), 929–940. https://doi.org/10.2307/1939815

Sunde, P., Böcker, F., Rauset, G. R., Kjellander, P., Chrenkova, M., Skovdal, T. M., van Beeck Calkoen, S., Mayer, M., & Heurich, M. (2022). Mammal responses to predator scents across multiple study areas. *Ecosphere*, *13*(8), e4215. https://doi.org/10.1002/ecs2.4215

van Ginkel, H. A. L., Kuijper, D. P. J., Schotanus, J., & Smit, C. (2019). Wolves and tree logs: Landscape-scale and fine-scale risk factors interactively influence tree regeneration. *Ecosystems*, *22*(1), 202–212. https://doi.org/10.1007/s10021-018-0263-z

van Ginkel, H. A. L., Smit, C., & Kuijper, D. P. J. (2019). Behavioral response of naïve and non-naïve deer to wolf urine. *PLOS ONE*, *14*(11), e0223248. https://doi.org/10.1371/journal.pone.0223248

Wang, Y., Allen, M. L., & Wilmers, C. C. (2015). Mesopredator spatial and temporal responses to large predators and human development in the Santa Cruz Mountains of California. *Biological Conservation*, *190*, 23–33. https://doi.org/10.1016/j.biocon.2015.05.007

Wikenros, C., Kuijper, D. P. J., Behnke, R., & Schmidt, K. (2015). Behavioural responses of ungulates to indirect cues of an ambush predator. *Behaviour*, *152*(7–8), 1019–1040. https://doi.org/10.1163/1568539X-00003266

Winke, B. (2012, March 9). *Can deer tell how far away a source of an odor Is?* Petersen's Bowhunting. https://www.bowhuntingmag.com/editorial/can-deer-tell-how-far-away-a-source-of-an-odor-is/309382

Roads

 Control

★ Novel Scent

⚡ Wolf Urine

Figure 1. Locations of 30 experimental deer bait sites with camera traps deployed between

September and November 2018 in Michigan, USA. The top inset map (solid magenta line)

indicates the 15 sites in the Upper Peninsula (UP; wolves [*Canis lupus*] present) and the lower

inset map (black dashed line) indicates the 15 sites in the Lower Peninsula (LP; wolves absent).

The different symbols indicate the different treatments (wolf urine, lemon juice and distilled

water) assigned to each site.

Figure 2. Camera-trap detections of white-tailed deer (*Odocoileus virginianus*) at simulated deer

hunter-bait sites September to November 2018 in Michigan, USA. The representative images

indicate vigilance (A) and non-vigilance (B) behavior.

Figure 3. Perceived effect of wolf (*Canis lupus*) predation risk on deer behavior. Difference

before treatment period and during treatment period in group size, group vigilance, number of

visits, and vigilance intensity (sum of proportion of group vigilance in each event divided by

682   total time of event) from September to November 2018 in Michigan, USA. Symbols indicate the

683   mean difference and error bars indicate 95% confidence interval. Blue shading represents the

684   control sites (water), yellow indicates sites treated with novel scent (lemon), and grey indicates

685   wolf-urine treated sites. Positive values indicate an increase in the metric. Asterix indicates p <

686   0.05.

 Figure 4. Temporal activity patterns of white-tailed deer (*Odocoileus virginianus*) populations

 before (solid line) and during treatment application (dashed line) in the Upper Peninsula (UP)

 and Lower Peninsula (LP) of Michigan, USA between September and November 2018. Shaded

 area indicates the temporal overlap between the 2 periods. The orange circle highlights the

 decrease in deer activity at wolf (*Canis lupus*) urine treated sites in the UP. The reported U2

 statistic is evaluated against the test statistic U2 = 0.19.

706 **Figure 5.** Effects graph showing the predicted vigilance intensity (sum of proportion of group

707 vigilant in each event divided by total time of event) of white-tailed deer (*Odocoileus*

708 *virginianus*) by percent vegetation cover for all treatment sites across the Upper Peninsula (UP)

709 of Michigan, USA (A; wolves present); and all treatment sites across the Lower Peninsula (LP)

710 of Michigan, USA (B; wolves [*Canis lupus*] absent).

Roads

✚ Control

★ Novel Scent

⚡ Wolf Urine