Eat or Be Eaten: Implications of potential exploitative competition between wolves and

humans across predator- savvy and -naive deer populations

ELLEN M. CANDLER, *University of Minnesota, 135 Skok Hall, 2003 Upper Buford Circle, St.*

Paul, MN 55108, USA, ORCID: 0000-0002-5810-1942

STOTRA CHAKRABARTI, *Macalester College, Olin-Rice Science Center 211, St. Paul, MN*

55105, USA, ORCID: 0000-0001-8741-4780

WILLIAM J. SEVERUD, *South Dakota State University, Department of Natural Resource*

Management, Brookings, South Dakota USA 57006, USA, ORCID: 0000-0003-0150-

5986

JOSEPH K. BUMP, *University of Minnesota, 135 Skok Hall, 2003 Upper Buford Circle, St.*

Paul, MN 55108, USA, ORCID: 0000-0002-4369-7990

Abstract

Predator recolonization often evokes innate as well as learnt anti-predatory responses in prey. To test white-tailed deer risk allocation amidst an expanding wolf population in Michigan, we used a paired experimental approach to investigate deer behavior in areas with and without wolves. We treated sites with wolf urine, a novel scent (lemon juice), and a control (distilled water), and used remote cameras to observe deer vigilance, diel activity, and abundance before and after treatment. We also examined the effect of vegetation cover/predator ambush potential on deer vigilance. For most deer-response metrics, wolf urine had no effect irrespective of actual wolf presence or absence. However, we observed a difference in the overall diel pattern of deer in areas with wolf presence versus without. Further, we detected significant shifts in diel activity in both areas pre- and post-treatment. While such shifts were not consistent, we could detect a reduction in crepuscular deer activity in areas where deer coexist with wolves. While deer vigilance was typically unaffected by treatments, habitat cover was found to be pivotal. We observed a significant positive effect of vegetation cover on deer vigilance at sites where wolves were present. These results likely indicate that predator cues affect deer vigilance more acutely in denser habitats, which presumably facilitate higher predator success. We did not observe a similar response in the *predator-naive* deer population. In areas where human and non-human predators hunt the same prey, predator presence may thus negatively impact human hunting success through exploitative competition by making deer more vigilant. However, hunters may avoid such competition by choosing hunting/bait sites located in open areas. Our results on fundamental predation ecology have strong applied implications for fostering human-predator coexistence.

Key words: behavioural manipulation, human-wildlife conflict, olfactory cues, optimal

foraging, predator-prey interactions, predator reintroduction

Background

Predators can alter prey populations directly through lethal offtake and/or behaviourally

mediated indirect effects [1–3]. While some prey succumb, others avoid predation by changing

their behaviour and life-histories [4,5]. Predator-effects on prey are a fundamental focus in

ecology, although, dynamic predator-prey interactions in a fast-changing Anthropocene warrant

more experimental examinations of the underlying mechanisms of these direct and indirect

effects [6,7]. While predator populations are declining in many parts of the world [8], some areas

are witnessing their comebacks [9–11]. Such contrasting levels of predator presence create (and

re-create) intriguing interactions. Predators recolonizing previously predator-free habitats may

encounter potentially naive prey that have lost anti-predatory strategies [12,13].

58   Anti-predator responses are defence strategies that prey species employ when they

perceive predation risk via visual, auditory, tactile, and/or olfactory stimuli. When at risk, prey

respond through multiple mechanisms either proactively, such as by increasing vigilance,

avoiding risky areas, social grouping, and birth synchrony, or reactively through evasion,

camouflage, and defence-herding [5,14–17]. Successful anti-predatory responses, while

providing survival benefits, can have severe fitness costs for individuals. For example, ungulates

that are known to intensify vigilance as a response to predator cues limit durations of feeding or

avoid high-risk areas even if they are rich with resources [2,18–21]. Furthermore, certain

herbivores are known to incur major reproductive costs while courting at breeding grounds that

also attract a host of predators [22,23]. However, perception and response to predation risk have

significant decision-making implications that can result from a mix of instinctive and learnt

behaviours. Thus, the risk of predation can vary greatly amongst experienced prey (prey that live

under constant predation pressure) versus naive prey (prey that live without or under low

predation pressure). For example, in areas where wolves (*Canis lupus*) and grizzly bears (*Ursus*

horribilis) had been extirpated, moose (*Alces alces*) did not exhibit avoidance to predator

olfactory or auditory cues, while those that had been exposed to both predators showed

avoidance [12]. Similarly, species on predator-free islands have been observed to 'adaptively

forget' anti-predatory strategies, resulting in prey naivety [24]

76   The Upper and Lower Peninsula (UP and LP) of Michigan, USA offers a novel

opportunity to explore these drivers of predator-effects. In the LP, white-tailed deer (*Odocoileus*

virginianus) occur where wolves have been functionally absent since the mid-1800s; while deer

in the nearby UP coexist with a re-established wolf population since the mid-2010s [25]. Also,

white-tailed deer are a subsistence species for indigenous populations and a highly valued

hunting species in this region. In Michigan, deer are typically hunted by humans at specific

locations where deer are acclimatized with bait consisting of corn, fruits, or vegetables – creating

artificially provisioned resource rich pockets on the landscape. These resource-rich but localized

hunter-bait sites are intended to attract and condition deer to revisit these sites frequently.

According to behavioural optimality, prey species need to balance the conflicting demands of

efficient foraging and predator avoidance [26,27]. Thus, deer behaviour at these hunter-bait sites

is expected to be mediated by the perceived risk of predation because wolves are attracted to bait

sites as well [28], which in turn should be influenced by prey experience or naivety towards

predators.

90   Through a paired experimental design, we tested for differences in risk perception by

examining the immediate behavioural effects of wolf olfactory cues on naive versus experienced

deer at hunter-bait sites. Whereas we expected no response from naive deer (LP deer; no wolf

population) to wolf scent (urine), we predicted deer that generally experience wolf predation

(UP) would increase group size, intensity of vigilance, and occur less often at bait sites with wolf

cues (scent). We also predicted that the immediate effects of predator cues on deer vigilance will

be further modulated by habitat covariates such as vegetation cover at the sites. Prey are often

cautious, and tend to avoid 'risky' areas that lack good visibility and offer ambushing

opportunities for predators even where food is abundant [2,29,30]. Additionally, in accordance

with the predation risk allocation hypothesis that suggests temporal variation in predation risk

severity will affect foraging and vigilance behaviour in prey [27], we predicted that experienced

deer would adjust their activity patterns to avoid crepuscular times when wolves are most active,

while naive deer would not make such adjustments.

Methods

Study Area

We conducted the study in Michigan's UP and LP from September 21, 2018 to November 10,

2018. The UP study area was entirely within the Hiawatha National Forest and was interspersed

with unpaved, forest service roads. While the bait sites in the LP study area were all located in

Mackinaw State Forest of Michigan, the study area was also interspersed with private

agriculture, forestland, and paved, secondary roads (Figure 1). The main vegetation cover,

derived from the National Land Cover Database (NLCD), at the UP study sites was evergreen (5

sites), woody wetland (3 sites), grassland (3 sites), deciduous (2 sites), and mixed forest (2 sites),

while the LP study sites were deciduous (6 sites), scrub/shrub (4 sites), grassland (3 sites), and

evergreen (2 sites; Figure 1) [31].

Study Design

We established 30 deer bait sites between the two study areas, 15 in the UP and 15 in the LP

(Figure 1). To mimic recreational hunters, we selected sites based on deer hunting desirability

such as habitat openness and/or proximity to a deer trail [32]. Sites were at least one km away

from each other to avoid duplication and dependence [33].

We baited each site with 3.8 litres of corn spread in a 3 by 3-meter- square plot, as per the

Michigan deer baiting regulations [34]. For the duration of the study, all sites were baited each

week in order to maintain 3.8 litres of corn on the ground to replicate actual hunter behaviour

and to reduce the confounding effect of bait availability on deer behaviour. We deployed one

remote camera (Reconyx Hyperfire series or Cuddeback E) on a tree 4 meters from the centre of

the bait site. Each camera was programmed to take 5 burst images with no delay between

triggers to record the number of deer present, their posture, sex (when possible), and

presence/absence of fawn(s). We replaced batteries and memory cards weekly.

We used a 10 week before-after-control-impact (BACI) design for the experimental

treatment of both areas with one week preceding the six-week period to establish a baseline

condition of deer visitation to the bait sites [35]. For the first three weeks (September 28-October

20), we baited sites, recorded deer occurrence and behaviour via remote cameras with no scent

treatments applied. For the following three weeks we continued to bait and record deer after

treating sites with three scents. In each area, UP and LP, 5 sites were treated each with wolf urine

(predator treatment), lemon juice (novel scent; not a native plant), or distilled water (control).

We assigned site treatment randomly [20,36]. When we treated sites in the field, we visited all

'control' sites in an area first followed by the 'novel scent' sites, and finally 'experimental/wolf

urine' sites to avoid cross-site spread of scents. We also used different treatment application

tools (i.e. pipets, travel containers) for each treatment type to avoid cross contamination of

treatments. At each site, we applied 10 mL of the given treatment (experimental, novel, or

control) by dripping it from a pipet onto the bait site. This was done to mimic the amount and

application of scent marking of a pair of wolves [37].

We also recorded predator concealment or hiding potential (horizontal cover) at each site

to account for site-specific differences in habitat openness - an index of perceived predation

threat. We used a 2 m cover pole with 20 sections, each 10 cm long, to measure predator hiding

potential at each site [38,39]. From the centre of each bait pile, we measured the horizontal cover

in each cardinal direction. While one person held the cover pole 10-meters away from the centre

of the bait pile, another took a picture of the cover pole with a camera stationed on a 1-meter

high pole at the centre of the bait pile, intended to mimic deer visual height [39]. We conducted

this procedure twice during the study, once at the beginning and once when treatment started, to

account for change in hiding potential with loss of leaves later in the year. For each of the 20 (10

cm) sections on the 2 m cover pole, we estimated obscurement to the nearest 25% [39]. We

calculated a single mean and standard error value for each site and each measurement time

(beginning and middle of study).

Analysis

Photo Analysis

Images were tagged in batches using the DigiKam photo editing software [40]. Each

batch was defined by any set of images taken within 5 minutes of the prior image. In each batch,

for deer only, we recorded if fawns were present, labelled all sexes when possible, and tallied a

total count. For individual pictures within batches, in addition to generic behaviours (e.g.,

fighting, nursing, foraging), we recorded the number of individuals with their heads above their

shoulders, indicating vigilance/state of alertness (Figure 2) [14,41]. We also labelled batches for

different species that were captured at site visits including squirrels (*Sciurus* spp.), raccoons

(*Procyon lotor*), turkeys (*Meleagris gallopavo*), and coyotes (*Canis latrans*). We did not detect

additional scent marking by other species.

Treatment Impact

We explored the impact of different treatments in the two areas (UP and LP) on the

number of deer at each site, proportion of group that was vigilant, and the number of visits made

to the site. To account for the paired nature of our treatment design, we first averaged the

variable values for each site individually and calculated a difference in values before and during

treatment application. We then averaged across all sites within a given treatment. We also

calculated a vigilance intensity metric using equation 1. This metric includes both group

vigilance and event duration.

172
$$I = \frac{\Sigma \frac{v}{g}}{e}$$

173   (Eq 1)

Where *I* is the vigilance intensity metric for a given event, *v* is the number of deer vigilant in a

single image, *g* is the group size in a single image, and *e* is the total time of the event. Hence,

vigilance intensity simply standardizes individual vigilance across different group sizes and time

spent in front of the camera. Similar to the previous analysis, we calculated a difference in

vigilance intensity before and after treatment for each treatment type in each peninsula.

Diel Activity

To analyse possible temporal variability in the use of bait sites in the UP and LP before

and after treatment, we used a nonparametric kernel density estimation procedure [42,43]. We

converted times to radians and used a kernel density estimator to create a probability density

distribution for each before or after period [44]. We calculated the proportion of temporal

overlap between the two treatment periods for each treatment in an area [42]. We used a $\hat{\Delta}_4$ with

a smoothing parameter of 1 because our sample size for all analyses was greater than 50 [44].

We conducted this analysis using the overlap package [42,45] in R.

We applied Watson's U2 statistic with the CircStats package to test for homogeneity

between the two samples of interest (i.e. test for/detect a statistically significant shift in the diel

pattern before and after treatment) [46,47]. If deer significantly shifted their temporal pattern

between the two treatment periods, we expected Watson's U2 statistic would be greater than the

critical value (0.19 for an α value of 0.05) and $P < 0.05$. In the UP, we predicted that there would

be no shift in temporal visitation by deer at the control and lemon treated sites (high $\hat{\Delta}_4$, U2 ≤

0.19), while a shift to more nocturnal activity at the wolf urine treated sites would be detected

(lower $\hat{\Delta}_4$, U2 > 0.19; Kohl et al. 2018) to potentially avoid wolves that are known to be typically

less active at night [5]. In the LP, we expected that we would not see a significant shift in any

treatment sites because deer in the LP are ostensibly naive to wolf predation.

Vegetation Cover Analysis

We fitted generalized linear models using pooled data from both the UP and LP to check

for the effect on overall vigilance intensity by treatment effect (before/after) and type

(predator/novel/control), area (UP or LP) and vegetation cover (i.e., predator hiding potential;

S1) [43,48] through additive and interactive effects between parameters of interest.

Subsequently, we fitted models to examine the relationship between vigilance intensity and

vegetation cover at each treatment type within both the UP and LP before and after treatment.

Results

We obtained 286,436 images over seven weeks. After removing images taken during the pre-experimental, baseline week and images that did not contain deer, we had 213,264 images for analysis with 85,675 images taken in the LP and 127,589 images taken in the UP.

We observed a near 30% increase in group size at sites treated with wolf urine in the UP but this increase translates to less than half a deer, an increase that is not ecologically meaningful. Additionally, we observed a slight increase (<0.03) in the number of visits at both lemon and control treated sites in the UP and a marginal decrease (<0.04) in the proportion of the group that was vigilant at lemon treated sites in the LP (Table 1, Figure 3). We also observed a decrease in vigilance intensity of deer at wolf urine treated sites in the LP (Figure 3). We did not detect a significant change before and after treatments for any other treatment sites in either area for the four response variables, i.e., the proportion of group that is vigilant, number of visits, and vigilance intensity (Figure 3).

Table 1. Perceived effect of wolf (*Canis lupus*) predation risk on deer behaviour. Mean difference before treatment period and during treatment period in group vigilance, group size, number of visits, and vigilance intensity (sum of proportion of group vigilance in each event divided by total time of event) from September to November 2018 in the Upper Peninsula (UP) and Lower Peninsula (LP) Michigan, USA. *Differences before treatment period and during treatment periods were significant if the 95% confidence interval did not include 0.

Area	Response variable	Treatment	Mean difference	95% CI
UP	Group vigilance	wolf	-0.044	-0.101 to 0.013
		lemon	-0.032	-0.082 to 0.017
		control	-0.020	-0.071 to 0.031
	Group size	wolf	0.293*	0.093 to 0.492
		lemon	0.238	-0.159 to 0.635
		control	0.166	-0.018 to 0.351

	Number of visits	wolf	0.011	-0.020 to 0.043
		lemon	0.024*	0.011 to 0.037
		control	0.007*	0.003 to 0.011
	Vigilance intensity	wolf	0.001	-0.021 to 0.024
		lemon	-0.008	-0.024 to 0.007
		control	-0.010	-0.023 to 0.003
LP	Group vigilance	wolf	-0.007	-0.083 to 0.068
		lemon	-0.035*	-0.044 to -0.025
		control	0.016*	0.002 to 0.034
	Group size	wolf	-0.341	-1.294 to 0.611
		lemon	0.133	-0.061 to 0.327
		control	-0.110	-0.626 to 0.405
	Number of visits	wolf	0.004	-0.019 to 0.012
		lemon	0.001	0.002 to 0.004
		control	0.000	-0.065 to 0.065
	Vigilance intensity	wolf	-0.028	-0.055 to 0.001
		lemon	0.002	-0.012 to 0.008
		control	0.008	-0.010 to 0.026

Deer changed their diel activity in the LP and UP, and in all treatment types, before and

after treatment application (i.e., all U2 > 0.19). We did not detect a consistent directional pattern

in the temporal shifts for most sites. However, we did observe a difference between the UP and

LP diel patterns where the deer in the UP tended to reduce activity between 6:00 and 18:00 both

before and after treatment application while the deer in the LP did not have a consistent

inactivity period (Figure 4). Additionally, deer at wolf-treated sites in the UP reduced bait site

visitation in the evening hours (i.e., 1800h; Figure 4).

Deer vigilance intensity increased by 1.3 times for unit increase in vegetation cover in the

UP but not in the LP (Figure 5; S1). Though vigilance intensity did not increase with vegetation

cover in the LP, the base vigilance intensity (model intercepts) increased significantly after all

treatment applications (S2).

Discussion

Predator absence for long durations can lead to the gradual erosion of anti-predatory

responses of prey [12,13]. Subsequent recolonization of predators to such landscapes raises

questions about the response that naive prey will have once predators return, particularly in prey

populations valued and pursued by human hunters. We investigated the effects of simulated

predator presence on experienced versus naive deer at food-rich hunter-bait sites. We did not

find a significant effect of wolf scent treatment on deer vigilance, number of visits, or vigilance

intensity among predator-savvy/experienced deer in the UP but did note a slight increase in

group size, though the difference was not ecologically meaningful. A lack of response to wolf

urine has been documented in deer populations in the Netherlands and Poland where little

support was found for effects of predator scents in both predator savvy and predator naive prey

populations even when three times the volume of wolf urine was used [49]. This lack of reaction

to wolf scent cues in both areas may also be a result of decreasing response over time to the wolf

urine as it degrades [2,20,50]. Introducing cues that indicate immediate wolf proximity, such as

howling or visual, are likely to produce different results [49,51]. The absence of any response to

wolf urine may also be a result of predator hunting strategies. Cursorial predators might fail to

evoke strong anti-predatory responses in deer, as opposed to pronounced effects as observed in

the case of cues from ambush predators [52,53].

We also observed a significant shift in diel patterns at all sites before and after treatment

but there was no consistency to the shift in activity for most treatment sites. However, deer at

sites treated with wolf scent in the UP (where they coexist with wolves) reduced their visitation

in the evening hours. This finding is in accordance with other research that suggests that ungulate

prey shift their activity to avoid riskier times [5,54]. For example, deer in Minnesota, USA

demonstrated a more diurnal pattern under the effects of wolf scent to likely avoid the riskier

crepuscular times when wolves tend to hunt [1]. It is also noteworthy that before and after

treatment application, all UP sites exhibited a decrease in deer activity at bait sites between 6:00

and 18:00 hours, while the LP sites had inconsistent patterns. This suggests that deer in the UP

are perhaps avoiding peak wolf activity/hunting hours regardless of cues at sites, while LP deer

do not have as large of an inactivity period, likely because they do not share habitat with wolves

(S2; [5]).

Our results show a strong correlation of vigilance intensity with vegetation cover where

wolves are established. Increase in vegetation cover at the UP sites were characterized by intense

vigilance activity in deer, regardless of treatment. Thicker vegetation likely represents more

predator hiding potential and difficult escape routes, likely causing an increase in deer vigilance

regardless of olfactory cues at the site (Figure 5) [20,55,56]. Research comparing white-tailed

deer and mule deer (*Odocoileus hemionus*) in Oregon's expanding wolf range reported that

white-tailed deer did not typically avoid areas with wolves but did favour areas with less dense

vegetation that facilitates an easy escape because they relied on flight and early detection of

predators [55]. Alternatively, mule deer avoided areas where wolves were present altogether

because they relied on a stotting gait that is advantageous in uneven areas with more obstacles.

This phenomenon has also been noted in other species such as chacma baboons (*Papio ursinus*)

in Gorongosa National Park, Mozambique where vocalization and vigilance were heightened in

denser habitats [57]. Our findings further highlight the importance of considering prey behaviour

and escape strategies, along with habitat characteristics for a mechanistic understanding of

predator-effects.

Though vegetation cover did not have a significant effect at LP sites, deer at wolf urine

treated sites did demonstrate higher baseline vigilance intensity after treatment was applied but it

did not vary across vegetation cover. This suggests that naive deer may demonstrate an innate

fear of wolves, also reported in East Limestone Island, Canada [19]. However, when the novel

scent (lemon) and control sites were assessed in the same context, we saw a similar pattern

indicating that something other than predator scent might be affecting baseline vigilance later in

the study. Since the second half of the study was conducted during archery hunting season, this

shift in vigilance may indicate a response to increased human presence [58]. Research in Poland

found that red deer (*Cervus elaphus*) demonstrated heightened vigilance during hunting season

when hunters were likely pursuing them [59].

Our findings give partial support to the predator risk allocation hypothesis by showing a

shift in activity by prey with predator experience to a less risky feeding time for prey that

experience a pause in predation pressure when predators are less active [5,27]. This hypothesis

may also explain the lack of difference in vigilance detected in the UP-deer population in

relation to the wolf urine treatment. Because this population is under constant predation pressure

from wolves and frequently encounters wolf cues, deer in the UP may demonstrate a change of

vigilance especially at food rich site such as hunter-bait sites, in order to take advantage of the

food source as quickly as possible. Deer in the UP are also likely, as we observed, to show more

vigilance intensity in areas with higher predation potential, such as high vegetation cover [55].

Finally, through our experimental study we highlight the importance of considering both

spatial and temporal factors when understanding predator effects on experienced versus naive

prey. Past research has suggested that naive prey do not maintain the innate ability to react to

predator olfactory cues but do have the ability to learn and adjust to predators within one

generation [12,60]. Though olfactory cues did not have a significant effect on most variables we

investigated, we did observe a strong predator effect depending on vegetation cover that

ostensibly modifies risk.

The recreational hunting tradition of baiting is controversial inside and outside the

hunting community. Baiting increases the opportunity for a successful shot and more immediate

kill, but also creates a predictable food source that concentrates deer and deer scent, which in

turn attracts wolves [61]. As large predators continue to recolonize their former ranges, they will

inevitably overlap with human hunters [62,63]. Our current research can shed light on the impact

a predator has on a predator savvy and predator naive prey population as it relates to hunting by

humans. We show that predator cues can affect deer vigilance at bait-sites with thicker

vegetation, seemingly making deer wary and difficult to hunt, thereby potentially affecting the

experience and satisfaction reported in hunter surveys. However, we urge readers, managers, and

the hunting community to interpret our results cautiously and constructively. Our results can be

formative in aiding carnivore-human coexistence wherein human hunters can have more success

if they potentially choose not to bait/hunt deer at vegetatively dense sites.

Acknowledgements
This material is based upon work supported by the National Science Foundation Graduate

Research Fellowship Program under Grant No. 00039218. Any opinions, findings, and

conclusions or recommendations expressed in this material are those of the authors and do not

necessarily reflect the views of the National Science Foundation. This research was also

supported by grants from the National Science Foundation to J. K. Bump (NSF ID#1545611,

NSF ID#1556676). We thank M. Dean, K. Rajski, and A. Schubel for field assistance.

Author Contributions

E.M.C.: idea formulation, sampling design, data collection, formal analysis, methodology,

project administration, resources, supervision, validation, visualization, writing—original draft,

writing—review and editing; S.C.: sampling design, methodology, validation, visualization,

writing—review and editing; W.J.S.: sampling design, methodology, writing—review and

editing, J.K.B.: idea formulation, sampling design, project administration, writing—review and

editing, funding acquisition, supervision, resources

Literature Cited

1. Palmer MS, Portales-Reyes C, Potter C, Mech LD, Isbell F. 2021 Behaviorally-mediated
trophic cascade attenuated by prey use of risky places at safe times. *Oecologia* **195**, 235–
248. (doi:10.1007/s00442-020-04816-4)

2. Kuijper DPJ, de Kleine C., Churski M., van Hooft P., Bubnicki J., Jędrzejewska B. 2013
Landscape of fear in Europe: Wolves affect spatial patterns of ungulate browsing in
Białowieża Primeval Forest, Poland. *Ecography* **36**, 1263–1275. (doi:10.1111/j.1600-
0587.2013.00266.x)

3. McNay RS, Lamb CT, Giguere L, Williams SH, Martin H, Sutherland GD, Hebblewhite M.
2022 Demographic responses of nearly extirpated endangered mountain caribou to recovery
actions in Central British Columbia. *Ecological Applications* **n/a**, e2580.
(doi:10.1002/eap.2580)

4. Sabal MC *et al.* 2021 Predation landscapes influence migratory prey ecology and evolution.
Trends in Ecology & Evolution **36**, 737–749. (doi:10.1016/j.tree.2021.04.010)

5. Kohl MT, Stahler DR, Metz MC, Forester JD, Kauffman MJ, Varley N, White PJ, Smith
DW, MacNulty DR. 2018 Diel predator activity drives a dynamic landscape of fear.
Ecological Monographs **0**, 1–15. (doi:10.1002/ecm.1313)

6. Scrosati RA. 2021 Nonconsumptive predator effects on prey demography: Recent advances
using intertidal invertebrates. *Frontiers in Ecology and Evolution* **9**.

7. Parsons MA, Newsome TM, Young JK. 2022 The consequences of predators without prey.
Frontiers in Ecology and the Environment **20**, 31–39. (doi:10.1002/fee.2419)

8. Ripple WJ *et al.* 2014 Status and ecological effects of the world's largest carnivores. *Science*
343, 1241484. (doi:10.1126/science.1241484)

9. Scharf AK, Fernández N. 2018 Up-scaling local-habitat models for large-scale conservation:
Assessing suitable areas for the brown bear comeback in Europe. *Diversity and Distributions*
24, 1573–1582. (doi:10.1111/ddi.12796)

10. Jarausch A, Harms V, Kluth G, Reinhardt I, Nowak C. 2021 How the west was won: genetic
reconstruction of rapid wolf recolonization into Germany's anthropogenic landscapes.
Heredity **127**, 92–106. (doi:10.1038/s41437-021-00429-6)

11. Gigliotti LC, Matchett MR, Jachowski DS. 2019 Mountain lions on the prairie: Habitat
selection by recolonizing mountain lions at the edge of their range. *Restoration Ecology* **27**,
1032–1040. (doi:10.1111/rec.12952)

12. Berger J, Swenson JE, Persson I-L. 2001 Recolonizing carnivores and naïve prey:
Conservation lessons from pleistocene extinctions. *Science* **291**, 1036–1039.
(doi:10.1126/science.1056466)

13. Berger J. 1999 Anthropogenic extinction of top carnivores and interspecific animal
behaviour: Implications of the rapid decoupling of a web involving wolves, bears, moose and
ravens. *Proc. R. Soc. Lond. B* **266**, 2261–2267. (doi:10.1098/rspb.1999.0917)

14. Flagel DG, Belovsky GE, Beyer DE. 2016 Natural and experimental tests of trophic
cascades: Gray wolves and white-tailed deer in a Great Lakes forest. *Oecologia* **180**, 1183–
1194. (doi:10.1007/s00442-015-3515-z)

15. Ims RA. 1990 On the adaptive value of reproductive synchrony as a predator-swamping
strategy. *The American Naturalist* **136**, 485–498. (doi:10.1086/285109)

16. Rowland HM, Burriss RP, Skelhorn J. 2020 The antipredator benefits of postural camouflage
in peppered moth caterpillars. *Sci Rep* **10**, 21654. (doi:10.1038/s41598-020-78686-4)

17. Mech LD. 2007 Possible use of foresight, understanding, and planning by wolves hunting
muskoxen. *Arctic* **60**, 145–149.

18. Melchiors MA, Leslie CA. 1985 Effectiveness of Predator Fecal Odors as Black-Tailed Deer
Repellents. *The Journal of Wildlife Management* **49**, 358–362. (doi:10.2307/3801531)

19. Chamaillé-Jammes S, Malcuit H, Saout SL, Martin J-L. 2014 Innate threat-sensitive
foraging: Black-tailed deer remain more fearful of wolf than of the less dangerous black bear
even after 100 years of wolf absence. *Oecologia* **174**, 1151–1158. (doi:10.1007/s00442-013-
2843-0)

20. Wikenros C, Kuijper DPJ, Behnke R, Schmidt K. 2015 Behavioural responses of ungulates
to indirect cues of an ambush predator. *Behaviour* **152**, 1019–1040. (doi:10.1163/1568539X-
00003266)

21. Sahlén E, Noell S, DePerno CS, Kindberg J, Spong G, Cromsigt JPGM. 2016 Phantoms of
the forest: Legacy risk effects of a regionally extinct large carnivore. *Ecol Evol* **6**, 791–799.
(doi:10.1002/ece3.1866)

22. Jhala YV, Isvaran K. 2016 Behavioural ecology of a grassland antelope, the blackbuck
antilope cervicapra: Linking habitat, ecology and behaviour. In *The Ecology of Large*
Herbivores in South and Southeast Asia (eds FS Ahrestani, M Sankaran), pp. 151–176.
Dordrecht: Springer Netherlands. (doi:10.1007/978-94-017-7570-0_6)

23. Cherry MJ, Morgan KE, Rutledge BT, Conner LM, Warren RJ. 2016 Can coyote predation
risk induce reproduction suppression in white-tailed deer? *Ecosphere* **7**, e01481.
(doi:10.1002/ecs2.1481)

24. Blumstein DT, Daniel JC, Springett BP. 2004 A test of the multi-predator hypothesis: Rapid
loss of antipredator behavior after 130 years of isolation. *Ethology* **110**, 919–934.
(doi:10.1111/j.1439-0310.2004.01033.x)

25. O'Neil ST, Bump JK, Beyer DE. 2017 Spatially varying density dependence drives a shifting
mosaic of survival in a recovering apex predator (*Canis lupus*). *Ecology and Evolution* **7**,
9518–9530. (doi:10.1002/ece3.3463)

26. Sih A. 1980 Optimal behavior: Can foragers balance two conflicting demands? *Science* **210**,
1041–1043. (doi:10.1126/science.210.4473.1041)

27. Lima SL, Bednekoff PA. 1999 Temporal variation in danger drives antipredator behavior:
the predation risk allocation hypothesis. *The American Naturalist* **153**, 649–659.
(doi:10.1086/303202)

28. Bump JK, Murawski CM, Kartano LM, Beyer DEJr, Roell BJ. 2013 Bear-baiting may
exacerbate wolf-hunting dog conflict. *PLOS ONE* **8**, e61708.
(doi:10.1371/journal.pone.0061708)

29. Edwards J. 1983 Diet shifts in moose due to predator avoidance. *Oecologia* **60**, 185–189.
(doi:10.1007/BF00379520)

30. Ripple WJ, Beschta RL. 2004 Wolves and the ecology of fear: Can predation risk structure
ecosystems? *BioScience* **54**, 755–766. (doi:10.1641/0006-
3568(2004)054[0755:WATEOF]2.0.CO;2)

31. Homer C, Dewitz J, Yang L, Jin S, Danielson P, Coulston J, Herold N, Wickham J, Megown
K. 2015 Completion of the 2011 national land cover database for the conterminous United
States – representing a decade of land cover change information. *PHOTOGRAMMETRIC*
ENGINEERING , 11.

32. Peterson TJ. 2015 5 best funnel & pinch-point setups for hunting whitetails. *Game & Fish*.
See https://www.gameandfishmag.com/editorial/5-best-funnel-pinch-point-setups-for-
hunting-whitetails/191246.

33. Winke B. 2012 Can deer tell how far away a source of an odor Is? *Petersen's Bowhunting*.
See https://www.bowhuntingmag.com/editorial/can-deer-tell-how-far-away-a-source-of-an-
odor-is/309382.

34. Michigan Department of Natural Resources. 2018 Michigan Hunting Digest.

35. Stewart-Oaten A, Murdoch WW, Parker KR. 1986 Environmental impact assessment:
'Pseudoreplication' in time? *Ecology* **67**, 929–940. (doi:10.2307/1939815)

36. Atkins R, Blumstein DT, Moseby KE, West R, Hyatt M, Letnic M. 2016 Deep evolutionary
experience explains mammalian responses to predators. *Behav Ecol Sociobiol* **70**, 1755–
1763. (doi:10.1007/s00265-016-2181-4)

37. Peters RP, Mech LD. 1975 Scent-Marking in wolves: Radio-tracking of wolf packs has
provided definite evidence that olfactory sign is used for territory maintenance and may
serve for other forms of communication within the pack as well. *American Scientist* **63**, 628–
637.

38. Kuijper DPJ, Verwijmeren M, Churski M, Zbyryt A, Schmidt K, Jędrzejewska B, Smit C.
2014 What cues do ungulates use to assess predation risk in dense temperate forests? *PLOS*
ONE **9**, e84607. (doi:10.1371/journal.pone.0084607)

39. Severud WJ, DelGiudice GD, Obermoller TR. 2019 Association of moose parturition and
post-parturition habitat with calf survival. *The Journal of Wildlife Management* **83**, 175–183.
(doi:https://doi.org/10.1002/jwmg.21570)

40. Niedballa J, Sollmann R, Courtiol A, Wilting A. 2016 camtrapR: an R package for efficient
camera trap data management. *Methods in Ecology and Evolution* **7**, 1457–1462.
(doi:https://doi.org/10.1111/2041-210X.12600)

41. Schuttler SG, Parsons AW, Forrester TD, Baker MC, McShea WJ, Costello R, Kays R. 2017
Deer on the lookout: How hunting, hiking and coyotes affect white-tailed deer vigilance.
Journal of Zoology **301**, 320–327. (doi:https://doi.org/10.1111/jzo.12416)

42. Wang Y, Allen ML, Wilmers CC. 2015 Mesopredator spatial and temporal responses to
large predators and human development in the Santa Cruz Mountains of California.
Biological Conservation **190**, 23–33. (doi:10.1016/j.biocon.2015.05.007)

43. Prugh LR, Sivy KJ, Mahoney PJ, Ganz TR, Ditmer MA, van de Kerk M, Gilbert SL,
Montgomery RA. 2019 Designing studies of predation risk for improved inference in
carnivore-ungulate systems. *Biological Conservation* **232**, 194–207.
(doi:10.1016/j.biocon.2019.02.011)

44. Ridout MS, Linkie M. 2009 Estimating overlap of daily activity patterns from camera trap
data. *JABES* **14**, 322–337. (doi:10.1198/jabes.2009.08038)

45. Meredith M, Ridout M. 2018 overlap: Estimates of coefficient of overlapping for animal
activity patterns.

46. Lund U, Agostinelli C. 2012 CircStats: Circular statistics, from 'topics in circular statistics'
(2001).

47. Lashley MA, Cove MV, Chitwood MC, Penido G, Gardner B, DePerno CS, Moorman CE.
2018 Estimating wildlife activity curves: Comparison of methods and sample size. *Scientific*
Reports **8**, 1–11. (doi:10.1038/s41598-018-22638-6)

48. Fležar U, le Roux E, Kerley GIH, Kuijper DPJ, te Beest M, Druce DJ, Prinsloo D, Cromsigt
JPGM. 2019 Simulated elephant-induced habitat changes can create dynamic landscapes of
fear. *Biological Conservation* **237**, 267–279. (doi:10.1016/j.biocon.2019.07.012)

49. van Ginkel HAL, Smit C, Kuijper DPJ. 2019 Behavioral response of naïve and non-naïve
deer to wolf urine. *PLOS ONE* **14**, e0223248. (doi:10.1371/journal.pone.0223248)

50. Bytheway JP, Carthey AJR, Banks PB. 2013 Risk vs. reward: How predators and prey
respond to aging olfactory cues. *Behav Ecol Sociobiol* **67**, 715–725. (doi:10.1007/s00265-
013-1494-9)

51. Liley S, Creel S. 2008 What best explains vigilance in elk: Characteristics of prey, predators,
or the environment? *Behavioral Ecology* **19**, 245–254. (doi:10.1093/beheco/arm116)

52. Preisser EL, Orrock JL, Schmitz OJ. 2007 Predator Hunting Mode and Habitat Domain Alter
Nonconsumptive Effects in Predator–Prey Interactions. *Ecology* **88**, 2744–2751.
(doi:10.1890/07-0260.1)

53. Sunde P, Böcker F, Rauset GR, Kjellander P, Chrenkova M, Skovdal TM, van Beeck
Calkoen S, Mayer M, Heurich M. 2022 Mammal responses to predator scents across multiple
study areas. *Ecosphere* **13**, e4215. (doi:10.1002/ecs2.4215)

54. Gehr B, Hofer EJ, Ryser A, Vimercati E, Vogt K, Keller LF. 2018 Evidence for
nonconsumptive effects from a large predator in an ungulate prey? *Behav Ecol* **29**, 724–735.
(doi:10.1093/beheco/ary031)

55. Dellinger JA, Shores CR, Craig A, Heithaus MR, Ripple WJ, Wirsing AJ. 2019 Habitat use
of sympatric prey suggests divergent anti-predator responses to recolonizing gray wolves.
Oecologia **189**, 487–500. (doi:10.1007/s00442-018-4323-z)

56. van Ginkel HAL, Kuijper DPJ, Schotanus J, Smit C. 2019 Wolves and tree logs: Landscape-
scale and fine-scale risk factors interactively influence tree regeneration. *Ecosystems* **22**,
202–212. (doi:10.1007/s10021-018-0263-z)

57. Hammond P, Lewis-Bevan L, Biro D, Carvalho S. In press. Risk perception and terrestriality
in primates: A quasi-experiment through habituation of chacma baboons (Papio ursinus) in
Gorongosa National Park, Mozambique. *American Journal of Biological Anthropology* **n/a**.
(doi:10.1002/ajpa.24567)

58. Benhaiem S, Delon M, Lourtet B, Cargnelutti B, Aulagnier S, Hewison AJM, Morellet N,
Verheyden H. 2008 Hunting increases vigilance levels in roe deer and modifies feeding site
selection. *Animal Behaviour* **76**, 611–618. (doi:10.1016/j.anbehav.2008.03.012)

59. Proudman NJ *et al.* 2020 Red deer allocate vigilance differently in response to spatio-
temporal patterns of risk from human hunters and wolves. *Wildl. Res.* **48**, 163–174.
(doi:10.1071/WR20059)

60. Steindler LA, Blumstein DT, West R, Moseby KE, Letnic M. 2020 Exposure to a novel
predator induces visual predator recognition by naïve prey. *Behav Ecol Sociobiol* **74**, 102.
(doi:10.1007/s00265-020-02884-3)

61. Hughes NK, Price CJ, Banks PB. 2010 Predators are attracted to the olfactory signals of
prey. *PLOS ONE* **5**, e13114. (doi:10.1371/journal.pone.0013114)

62. Bled F *et al.* 2022 Balancing carnivore conservation and sustainable hunting of a key prey
species: A case study on the Florida panther and white-tailed deer. *Journal of Applied*
Ecology **n/a**. (doi:10.1111/1365-2664.14201)

63. Grima N, Brainard J, Fisher B. 2021 Are wolves welcome? Hunters' attitudes towards
wolves in Vermont, USA. *Oryx* **55**, 262–267. (doi:10.1017/S0030605319000061)

Figure Captions

Figure 1. Locations of 30 experimental deer bait sites with camera traps deployed between September and November 2018 in Michigan, USA. The top inset map (solid magenta line) indicates the 15 sites in the Upper Peninsula (UP; wolves [*Canis lupus*] present) and the lower inset map (black dashed line) indicates the 15 sites in the Lower Peninsula (LP; wolves absent). The different symbols indicate the different treatments (wolf urine, lemon juice and distilled water) assigned to each site.

Figure 2. Camera-trap detections of white-tailed deer (*Odocoileus virginianus*) at simulated deer hunter-bait sites September to November 2018 in Michigan, USA. The representative images indicate vigilance (A) and non-vigilance (B) behaviour.

Figure 3. Perceived effect of wolf (*Canis lupus*) predation risk on deer behaviour. Difference before treatment period and during treatment period in group size, group vigilance, number of visits, and vigilance intensity (sum of proportion of group vigilance in each event divided by total time of event) from September to November 2018 in Michigan, USA. Symbols indicate the mean difference and error bars indicate 95% confidence interval. Blue shading represents the control sites (water), yellow indicates sites treated with novel scent (lemon), and grey indicates wolf-urine treated sites. Positive values indicate an increase in the metric. Asterix indicates $p < 0.05$.

Figure 4. Temporal activity patterns of white-tailed deer (*Odocoileus virginianus*) populations before (solid line) and during treatment application (dashed line) in the Upper Peninsula (UP)

and Lower Peninsula (LP) of Michigan, USA between September and November 2018. Shaded

area indicates the temporal overlap between the 2 periods. The orange circle highlights the

decrease in deer activity at wolf (*Canis lupus*) urine treated sites in the UP. The reported U2

statistic is evaluated against the test statistic U2 = 0.19.

Figure 5. Effects graph showing the predicted vigilance intensity (sum of proportion of group

vigilant in each event divided by total time of event) of white-tailed deer (*Odocoileus*

virginianus) by percent vegetation cover for all treatment sites across the Upper Peninsula (UP)

of Michigan, USA (A; wolves present); and all treatment sites across the Lower Peninsula (LP)

of Michigan, USA (B; wolves [*Canis lupus*] absent).

Figure 1

587 Figure 2

Figure 3

627 Figure 4

647   Figure 5