Common brain areas for processing physiologically and socially 'needed' stimuli

Bosulu, J[1]., Luo, Y[2]. Hétu, S.[1]

[1]Faculté des arts et des sciences, Université de Montréal
[2]School of Psychology and Cognitive Science, East China Normal University

ABSTRACT

We looked at the overlap between brain areas related to perception of physiologically and socially needed stimuli and how they might regulate serotonin levels. First, we conducted separate ALE meta-analyses on published results pertaining to brain activation patterns when participants perceived food while hungry or water while thirsty, and social interactions while being excluded. This allowed us to identify common consistent brain activation patterns for physiological and social needed stimuli. We also looked at significant spatial association between the common network and serotonin receptor distribution. We found that regions within the mid-posterior insula, the anterior cingulate cortex and the caudate are at the intersection of physiological (hunger and thirst) and social (exclusion) aspects of 'needing'. Furthermore, we found a significant positive spatial correlation between that common network and 5HT4 serotonin receptor. Our study suggests there is a common brain pattern during the processing of physiologically and socially needed stimuli, which modulates serotonin neurons possibly through a feedback loop from the PFC to the serotonergic neurons in the dorsal raphe nuclei.

KEYWORDS: need, deprivation, meta-analysis, fmri, hunger, thirst, social exclusion, interoception, serotonin, neuroscience

INTRODUCTION

Needs are related to states of deprivation of a biologically significant stimuli or events (Bouton, 2016). They can be related to physiologically relevant stimuli and events such as food and eating for hunger, or water and drinking for thirst, but also to social stimuli and events like belongingness for social exclusion and meaningful social contact for social isolation (Maslow,

1943; Baumeister & Leary, 1995; Maner et al., 2007). Not responding to a need might lead to some adverse consequences such as physiological or psychological suffering which go beyond mere frustration (MacGregor 1960; Baumeister & Leary, 1995). Maslow (1943) (likely influenced by Hull's (1943) drive theory) was convinced that there are motivations which are driven by basic needs instead of external rewards. He ranked these needs which later became Maslow's pyramid. At the base, there are needs that are physiological, then needs of security, belonging, esteem, (more social needs) and actualization. When the lower order needs are satisfied to a certain extent, the higher ones appear gradually. Despite the fact that evidences seem to go against some definite hierarchy of needs both on an individual and on a societal level (Goebel & Brown, 1981; Hofstede, 1984; Tay & Diener, 2011), Maslow's ideas indicate that physiological and social needs have something in common: they have some negative effect if not met, and hence motivate people to choose stimuli (or actions) that satisfy them. Indeed, in the brain, physiological needs and social needs can both alter the affective value of their relevant/significant needed stimuli (Chen et al., 2016; De Araujo, 2003). Furthermore, Tomova et al. (2020) recently showed some overlap in brain activations between hunger while observing food stimuli and social isolation while observing social interactions. However, the overlap between brain areas processing physiologically (e.g., through hunger and thirst) and socially (e.g., through exclusion and isolation) needed stimuli have rarely been tested, especially on a meta-analytic level, which would provide a quantitative comparison of the processing of physiologically vs. socially needed stimuli.

The idea of needs can be considered to include two components : the state itself and the (needed) stimulus that alleviate that state. Since processing of the needed stimuli are more likely to control behavior, either as cues or outcomes, than the deprivation states themselves (see Bindra, 1974; Toates, 1994), we will focus on brain responses associated with this processing. Hence, here we conceptualize the brain response to the perception of a stimulus, when deprived of it, as 'needing'. For hunger and thirst, brain studies that measured brain response to (viewing, tasting, or smelling) food or (either ingesting drops of water or viewing beverages) while hungry or thirsty have reported activations in the insula (for hunger/food: van der Lan et al., 2011; Goldstone, et. al., 2009; Siep et. al., 2009; for thirst/water: De Araujo et al., 2003; Egan et al., 2003; Becker et al., 2017; Farrell et al., 2011), orbitofrontal cortex (OFC) (for hunger/food: van

der Lan et al., 2011; Goldstone, et. al., 2009; Siep et. al., 2009; Führer et. al., 2008; for thirst/water: De Araujo et al., 2003; Saker et al., 2014), dorsal striatum (for hunger/food: van der Laan, et. al., 2011; Siep et. al., 2009), anterior cingugale cortext (ACC) (for hunger/food: Goldstone, et. al., 2009; Siep et. al., 2009; Führer et. al., 2008; for thirst/water: De Araujo et al., 2003; Becker et al., 2015, Becker et al., 2017; Farrell et al., 2011; Saker et al., 2013), amygdala and parahippocampal gyrus (for hunger/food: van der Laan, et. al., 2011; LaBar et. al., 2001; Führer, 2008; Goldstone, 2009; Mohanty, et. al., 2008; Chen et. al., 2020; for thirst/water: Becker, 2015), and posterior cingulate cortex (PCC) (for thirst/water: Farrell et al., 2011). Hence, although the brain networks supporting different types of physiological needs (here hunger and thirst) are not the same, they partially overlap in the ACC and amygdala, as well as OFC and the insular cortex; specifically the mid-posterior insula which has been shown to code for needed stimuli (see Bosulu et al., 2022; Livneh et al., 2020 ; Livneh et al., 2017). Furthermore, the insular cortex and ACC are viewed as common regions for conscious perception of both hunger and thirst (Mckinley et al., 2019), suggesting that they are core regions for homeostatic related perception and motivation (Craig, 2003).

Studies and meta-analyses of social exclusion/belongingness that have looked at brain responses to viewing others interact, or viewing (pleasant and/or close other) social stimuli while excluded from social interaction or after social isolation/loneliness have reported activity in the ACC (for social exclusion: Eisenberger, et al., 2003; Masten et al., 2009; Bolling et al., 2011; Vijayakumar et al., 2017; Mwilambwe-Tshilobo & Spreng, 2021), the posterior cingulate cortex (PCC) (for social exclusion: Vijayakumar et al., 2017; Mwilambwe & Spreng, 2021); the insula (for social exclusion: Masten et al., 2009; Bolling et al., 2011; Mwilambwe & Spreng, 2021; for social isolation: Tomova et al., 2020), the ventral striatum (for special exclusion: Masten et al., 2009; Vijayakumar et al., 2017; for social isolation: Inagaki et al., 2016), and the OFC (Cacioppo et al., 2013; Vijayakumar et al., 2017; for social isolation: Tomova et al., 2020). Some studies on social isolation/loneliness while viewing (pleasant and/or close other) social stimuli have also found decreased activity or no activation at all in the ventral striatum (Cacioppo et al., 2009; D'agostino et al., 2019). Overall, looking at these various results, social exclusion and isolation related responses to social cues/interactions seem to activate the insula, ACC, and OFC.

Overall, these different results suggest that brain networks supporting physiological and social needing at least partially overlap. Indeed, qualitatively the activation patterns of hunger/food, thirst/water, social exclusion/belongingness and isolation/meaningful contact often show activity within the insula (e.g. van der Lan et al., 2011; De Araujo et al., 2003; Mwilambwe-Tshilobo & Spreng, 2021; Tomova et al., 2020), ACC (ex. Goldstone, et. al., 2009; De Araujo et al., 2003; Mwilambwe-Tshilobo & Spreng, 2021) and OFC (ex. van der Lan et al., 2011; De Araujo et al., 2003; Vijayakumar et al., 2017; Tomova et al., 2020). However, in a recent study directly testing for common brain areas when participants observed food versus social interactions, after respectively 10 hours of hunger or 10 hours of social isolation, only activations of regions conaining mesolimbic and nigrostriatal dopamine neurons of the ventral tegmental area (VTA) and substantia nigra (SN) were found (Tomova et al., 2020). These conflicting observations suggest that there is still some uncertainty with regard to if and which regions may be recruited by both the processing of physiological (hunger and thirst) and social needs (exclusion/isolation) Hence, it seems timely to tackle this question using a meta-analysis which provides a quantitative analysis of the existing literature.

Tomova et al. (2020) study seems to suggest that, because VTA/SN are related to dopamine, this neurotransmitter would be a neurotransmitter commonly involved in perception of physiologically and socially needed stimuli, likely turning deprivation into reward seeking for both types of needs. Beyond dopamine, we can wonder what other neurotransmitters might be related to the processing of physiological and social needed stimuli. Serotonin is a good candidate for overall need processing, as it is central for biologically important sensory events (Sizemore, 2020). Indeed, low serotonin levels in the brain have been related to higher sensitivity to food (van Galen et al., 2021) and to higher reactivity to social exclusion (Preller et al., 2015); and serotonin level is negatively altered by social isolation (Dankoski et al., 2014). As low serotonin is often associated with aversive processing (Dayan and Huys, 2009) (which characterize deprivation); serotonin secretion is said to indicate how beneficial the current state is (Luo et al., 2016; Liu et al., 2020). In that sense, (low) serotonin level might signal states of deprivation of both biologically and socially significant stimuli/events. However such a mechanism has not been elucidated.

Hence, the present study aimed at investigating the common and specific brain activation patterns for, on one hand, the processing of physiologically needed stimuli and on the other socially needed stimuli. As an exploratory objective, to help us understand how the brain processes relevant stimuli in a deprived state, we also tested for the spatial correlation between areas recruited during both the processing for physiologically and socially needed stimuli and brain regions associated with the spatial distribution of neurotransmitter receptors, with a specific interest in serotonin receptors, given its inherent link with well-being (see Luo et al., 2016; Liu et al., 2020).

METHODOLOGY

Meta-analysis of brain coordinate

We used a meta-analytic approach to quantitatively assess brain activation patterns of both physiological and social aspects of 'needing' using large collections of data. Meta-analyses make it possible to investigate questions across different paradigms, samples and analysis approaches. Specifically, we first conducted two meta-analyses to quantitatively summarize results from functional magnetic resonance imaging (fMRI) published studies on physiological needs: hunger/food and thirst/water (Physiological-Need); when participants perceived food while hungry or water while thirsty (that we will refer to as Physiological-Need) ; and on social need: exclusion(isolation)/interaction; when participants perceived social interactions (that s/he was supposed to be part of) while being excluded (or isolated) from that social interaction (that we will refer to as Social-Need). (Of note, we looked for other kinds of physiological and social needs, but did not find a sufficient number of experiments that satisfied the selection criteria.) These meta-analyses will help us look at brain regions that are consistently recruited for processing of either physiologically or socially needed stimuli. Second, we used the single meta-analytic results for contrasts analysis as well as a conjunction analysis to identify differences and overlaps in consistent brain activation for physiological and social 'needing'.

We used the PRISMA framework and the following keywords to identify articles related to hunger: ("hunger" OR "food deprivation") AND ("fMRI"). For thirst, we used: ("thirst" OR

"water deprivation") AND ("fMRI"). For social exclusion, we used: ("social" AND ("exclusion" OR "isolation") AND "fmri"). Keywords were entered on PubMed (February 2021) for physiological needs and social needs. Additional articles were found by checking the articles references lists and review articles. For both physiological and social aspect of 'needing' the following inclusion criteria were used: healthy subjects; whole-brain analyses (with or without SVC), MNI or Talairach Coordinates (all Talairach coordinates were converted to MNI SPM152 in Ginger ALE using Lacanster transform; Lancaster et al., 2007); maps were corrected (or cluster level corrected); activation contrast only (see table 1 for selection criteria). The database returned 150 articles for hunger and 19 for thirst; 7 and 3 additional articles were found through other articles and reviews for hunger and thirst, respectively. In line with our objective, specific for physiological needs articles, we used two additional main criteria : 1) the participants were in a food or water deprivation state; 2) the participant was perceiving (visual, taste, odor, etc.) some food (while hungry) or water (while thirsty) stimulus. We found a total of 16 articles for hunger and 4 for thirst that met all inclusion criteria (see table 2 and 3).

Table 1. Selection Criteria

Criteria	Hunger and Thirst	Social exclusion
Privation contrast	Yes	Yes
Presence of the relevant rewarding cue	Yes	Yes
Passive viewing	Yes	Yes
Healthy individuals only	Yes	Yes
MNI or Talairach Coordinates	Yes	Yes
Whole brain contrast (with or without SVC)	Yes	Yes
Corrected	Yes	Yes

Activation contrast only	Yes	Yes
Excluded: MRI and resting states; cognitive conjunction analysis; and functional connectivity results	Yes	Yes

Table indicating selection criteria. The red coloured "yes" means the criterion is crucial for the definition of either 'physiological or social needing.

Table 2. Hunger selected articles

Paper	Contrast	Stimuli	Task	Healthy Participants
Jiang, T., Soussignan, R., Schaal, B., & Royet, J. P. (2014). Reward for food odors: an fMRI study of liking and wanting as a function of metabolic state and BMI. Social cognitive and affective neuroscience, 10(4), 561-568.	Liking – Wanting (hunger) Wanting – Liking (hunger) Food - NFood (hunger)	Odor	Odor presentation and rating	12
Green, E., Jacobson, A., Haase, L., & Murphy, C. (2015). Neural correlates of taste and pleasantness evaluation in the metabolic syndrome. Brain research, 1620, 57-71.	Hunger-Satiety: Control Sucrose>Caffeine (during hunger) Caffeine>Sucrose (hunger)	Taste	Swallowing aqueuos solution	15
Martens, M. J., Born, J. M., Lemmens, S. G., Karhunen, L., Heinecke, A., Goebel, R., ... & Westerterp-Plantenga, M. S.	Fasted: F>NF Fasted: stimuli - subject group Fasted: correlation	Visual	Viewing food and non food pictures	40

(2013). Increased sensitivity to food cues in the fasted state and decreased inhibitory control in the satiated state in the overweight. The American journal of clinical nutrition, 97(3), 471-479.	F>NF with BMI			
LaBar, K. S., Gitelman, D. R., Parrish, T. B., Kim, Y. H., Nobre, A. C., & Mesulam, M. (2001). Hunger selectively modulates corticolimbic activation to food stimuli in humans. Behavioral neuroscience, 115(2), 493.	Hungry-Satieted	Visual	Viewing food and tool images	17
Harding, I. H., Andrews, Z. B., Mata, F., Orlandea, S., Martinez-Zalacain, I., Soriano-Mas, C., ... & Verdejo-Garcia, A. (2018). Brain substrates of unhealthy versus healthy food choices: influence of homeostatic status and body mass index. International Journal of Obesity, 42(3), 448.	Healthy versus unhealthy food choice: fasted>satieted	Visual	participants were asked to select an option using a two-button response box.	30
Frank, S., Laharnar, N., Kullmann, S., Veit, R., Canova, C., Hegner, Y. L., ... & Preissl, H. (2010). Processing of food pictures: influence of hunger, gender and calorie content. Brain research, 1350, 159-166.	HiCal hungry vs. HiCal satiated	Visual	One-back task: press a button, either to indicate that the seen image was the same or another button to indicate that the picture was not the same	12

Holsen, L. M., Zarcone, J. R., Thompson, T. I., Brooks, W. M., Anderson, M. F., Ahluwalia, J. S., ... & Savage, C. R. (2005). Neural mechanisms underlying food motivation in children and adolescents. Neuroimage, 27(3), 669-676.	Premeal : Food > non-food	Visual	Viewing pictures of food, animals, and baseline control images	9
Jacobson, A., Green, E., Haase, L., Szajer, J., & Murphy, C. (2019). Differential Effects of BMI on Brain Response to Odor in Olfactory, Reward and Memory Regions: Evidence from fMRI. Nutrients, 11(4), 926.	Odor during the hunger condition	Odor/Taste	Odor stimuli deliverred to the tongue	40
Haase, L., Green, E., & Murphy, C. (2011). Males and females show differential brain activation to taste when hungry and sated in gustatory and reward areas. Appetite, 57(2), 421-434.	Hunger x male x Sucrose > water Hunger x female x NaCl > water Hunger x female x CAFFEINE > water Hunger x female x SUCROSE > water Hunger x female x CITRIC ACID > water	Taste	Taste stimuli presentation	21
Führer, D., Zysset, S., & Stumvoll, M. (2008). Brain activity in hunger and satiety: an exploratory visually stimulated FMRI study. Obesity, 16(5), 945-950.	Hunger > Satiety	Visual	Viewing pictrures and two-back task: he subject was asked to press a button when the same letter was shown two steps	12

			earlier	
Haase, L., Cerf-Ducastel, B., & Murphy, C. (2009). Cortical activation in response to pure taste stimuli during the physiological states of hunger and satiety. Neuroimage, 44(3), 1008-1021.	Hunger > satiety x sucrose Hunger > satiety x Caffeine Hunger > satiety x Saccharin Hunger > satiety x Citric acid	Taste	Stimulus presentation delivered to the tip of the tongue	18
Uher, R., Treasure, J., Heining, M., Brammer, M. J., & Campbell, I. C. (2006). Cerebral processing of food-related stimuli: effects of fasting and gender. Behavioural brain research, 169(1), 111-119.	Fasting > satiety x food-related stimuli (Chocolate AND Chicken) Fasting > satiety x food-related stimuli (Chocolate) Fasting > satiety x food-related stimuli (Chicken)	Visual	Viewing photographs of food	18
Holsen, L. M., Zarcone, J. R., Brooks, W. M., Butler, M. G., Thompson, T. I., Ahluwalia, J. S., ... & Savage, C. R. (2006). Neural mechanisms underlying hyperphagia in Prader‑Willi syndrome. Obesity, 14(6), 1028-1037.	HW x Pre‑meal x food > non‑food Hw x Pre‑meal x non‑food > food	Visual	Viewing pictures of food, animals and control	9
Jacobson, A., Green, E., & Murphy, C. (2010). Age-related functional changes in gustatory and reward processing regions: An fMRI study. Neuroimage, 53(2), 602-610.	Hunger x Older adults x Sucrose Hunger x Young adults x Sucrose Hunger x Older adults x Citric acid Hunger x Younger adults x Citric acid Hunger x Older	Taste	Stimuli were delivered orally	38

	adults x NaCl Hunger x Younger adults x NaCl Hunger x Older adults x Caffeine Hunger x Younger adults x Caffeine			
Cheah, Y. S., Lee, S., Ashoor, G., Nathan, Y., Reed, L. J., Zelaya, F. O., ... & Amiel, S. A. (2014). Ageing diminishes the modulation of human brain responses to visual food cues by meal ingestion. International Journal of Obesity, 38(9), 1186.	FASTED > FED x Visual food cue	Visual	Viewing food and non food	24
He, Q., Huang, X., Zhang, S., Turel, O., Ma, L., & Bechara, A. (2019). Dynamic causal modeling of insular, striatal, and prefrontal cortex activities during a food-specific Go/NoGo task. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *4*(12), 1080-1089.	Hungry > Satiated	Visual	Food pictures	45

List of articles that were selected. for hunger..

Table 3. thirst selected articles

Paper	Stimulus contrats	Healthy participants
Becker, C. A., Schmälzle, R., Flaisch, T., Renner, B., & Schupp, H. T. (2015). Thirst and the state-dependent representation of incentive stimulus value in human motive circuitry. Social cognitive and affective neuroscience, 10(12), 1722-1729.	(Beverages : Thirst > Chairs : Thirst) vs (Beverages : No-thirst > Chairs : No-thirst)	24
De Araujo, I. E., Kringelbach, M. L., Rolls, E. T., & McGlone, F. (2003). Human cortical responses to water in the mouth, and the effects of thirst. Journal of neurophysiology, 90(3), 1865-1876.	water-control water (pre and post satiety)	11
Becker, C. A., Flaisch, T., Renner, B., & Schupp, H. T. (2017). From thirst to satiety: the anterior mid-cingulate cortex and right posterior insula indicate dynamic changes in incentive value. Frontiers in human neuroscience, 11, 234.	(BOLD) activity proportional to the amount of water ingested	24
Saker, P., Farrell, M. J., Egan, G. F., McKinley, M. J., & Denton, D. A. (2018). Influence of anterior midcingulate cortex on drinking behavior during thirst and following satiation. Proceedings of the National Academy of Sciences, 115(4), 786-791.	Thirst > No-Thirst	20

List of articles that were selected. for thirst.

Regarding social exclusion/isolation, we found 129 articles, 22 were selected, and 4 more articles were found through other articles and reviews. Similarly to Physiological-Need, two additional main criteria were used for social needs : 1) the participant was in a social deprivationstate (i.e., s/he was either isolated from others or experienced social exclusion) ; 2) the participant was perceiving some social interaction s/he was excluded from, or after being isolated. It's important to note that these criteria resulted in all of our included articles for social needs using the cyber ball task (a virtual ball tossing game with other individuals from which the participant is excluded (Williams et al., 2000)), and were thus related to social exclusion (not isolation). Social exclusion (as threat to fundamental social needs in humans) will cause

emotional distress (Williams, 2007), and it has been proposed that the cyberball exclusion paradigm can induce need-like emotional distress (Bernstein and Claypool (2012). Of the few human studies on the perception of social cues in social isolation state (see Tomova et al., 2021), none met the inclusion criteria. Based on these criteria and theories, we selected a total of 26 articles (for social exclusion) for the meta-analysis (see table 4).

Table 4. social exclusion selected articles

Bach, P., Frischknecht, U., Bungert, M., Karl, D., Vollmert, C., Vollstädt-Klein, S., ... & Hermann, D. (2019). Effects of social exclusion and physical pain in chronic opioid maintenance treatment: fMRI correlates. European Neuropsychopharmacology, 29(2), 291-305.	Exclusion > Inclusion	Cyberball task	21
Bolling, D. Z., Pelphrey, K. A., & Vander Wyk, B. C. (2015). Trait-level temporal lobe hypoactivation to social exclusion in unaffected siblings of children and adolescents with autism spectrum disorders. Developmental cognitive neuroscience, 13, 75-83.	Social Exclusion > Fair Play	Cyberball task	15
Bolling, D. Z., Pitskel, N. B., Deen, B., Crowley, M. J., Mayes, L. C., & Pelphrey, K. A. (2011). Development of neural systems for processing social exclusion from childhood to adolescence. Developmental science, 14(6), 1431-1444.	comparison of social exclusion and fair play	Cyberball task	26
Bolling, D. Z., Pitskel, N. B., Deen, B., Crowley, M. J., McPartland, J. C., Kaiser, M. D., ... & Pelphrey, K. A. (2011). Enhanced neural responses to rule violation in children with autism: a comparison to social exclusion. Developmental Cognitive Neuroscience, 1(3), 280-294.	Social exclusion > fair play	Cyberball task	24
Bolling, D. Z., Pitskel, N. B., Deen, B., Crowley, M. J., McPartland, J. C., Mayes, L. C., & Pelphrey, K. A. (2011). Dissociable brain mechanisms for processing social exclusion and rule violation. NeuroImage, 54(3), 2462-2471.	Social exclusion > Fair play	Cyberball	26
Cheng, T. W., Vijayakumar, N., Flournoy, J. C., de Macks, Z. O., Peake, S. J., Flannery, J. E., ... & Pfeifer, J. H. (2020). Feeling left out or just surprised? Neural correlates of social exclusion and overinclusion in adolescence. Cognitive, Affective, & Behavioral Neuroscience, 1-16.	Exclusion>Inclusion	Cyberball task	97

Cogoni, C., Carnaghi, A., & Silani, G. (2018). Reduced empathic responses for sexually objectified women: An fMRI investigation. Cortex, 99, 258-272.	Self (exclusion > inclusion).	social pain task	36
de Water, E., Mies, G. W., Ma, I., Mennes, M., Cillessen, A. H., & Scheres, A. (2017). Neural responses to social exclusion in adolescents: Effects of peer status. Cortex, 92, 32-43.	Exclusion (popular + average players) > Inclusion Ball (popular + average players) Exclusion (popular + average players) > Inclusion No Ball (popular + average players)	Cyberball task	52
DeWall, C. N., Masten, C. L., Powell, C., Combs, D., Schurtz, D. R., & Eisenberger, N. I. (2012). Do neural responses to rejection depend on attachment style? An fMRI study. Social cognitive and affective neuroscience, 7(2), 184-192.	exclusion vs inclusion	Cyberball task	25
Eisenberger, N. I., Lieberman, M. D., & Williams, K. D. (2003). Does rejection hurt? An fMRI study of social exclusion. Science, 302(5643), 290-292.	explicit social exclusion implicit social exclusion		13
Falk, E. B., Cascio, C. N., O'Donnell, M. B., Carp, J., Tinney Jr, F. J., Bingham, C. R., ... & Simons-Morton, B. G. (2014). Neural responses to exclusion predict susceptibility to social influence. *Journal of Adolescent Health*, *54*(5), S22-S31.	Exclusion>inclusion	Cyberball task	36
Gradin, V. B., Waiter, G., Kumar, P., Stickle, C., Milders, M., Matthews, K., ... & Steele, J. D. (2012). Abnormal neural responses to social exclusion in schizophrenia. PloS one, 7(8), e42608.	Activations with increasing social exclusion	Cyberball task	20
Gilman, J. M., Curran, M. T., Calderon, V., Schuster, R. M., & Evins, A. E. (2016). Altered neural processing to social exclusion in young adult marijuana users. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *1*(2), 152-159.	Exclusion > Fair Play	Cyberball task	42
Gonzalez, M. Z., Beckes, L., Chango, J., Allen, J. P., & Coan, J. A. (2015). Adolescent neighborhood quality predicts adult dACC response to social exclusion. *Social cognitive and affective neuroscience*, *10*(7), 921-928.	Exclusion > inclusion	Cyberball task	85
Le, T. M., Zhornitsky, S., Wang, W., & Li, C. S. R. (2020). Perceived burdensomeness and neural responses to ostracism in the Cyberball task. Journal of Psychiatric Research, 130, 1-8.	Activations to social exclusion.	Cyberball task	64

Luo, S., Yu, D., & Han, S. (2016). Genetic and neural correlates of romantic relationship satisfaction. Social cognitive and affective neuroscience, 11(2), 337-348.	social exclusion vs inclusion	Cyberball task	42
Moor, B. G., Güroğlu, B., de Macks, Z. A. O., Rombouts, S. A., Van der Molen, M. W., & Crone, E. A. (2012). Social exclusion and punishment of excluders: neural correlates and developmental trajectories. Neuroimage, 59(1), 708-717.	No Ball- exclusion game > Ball- inclusion game No Ball- exclusion game > No Ball- inclusion game	Cyberball game	51
Nishiyama, Y., Okamoto, Y., Kunisato, Y., Okada, G., Yoshimura, S., Kanai, Y., ... & Yamawaki, S. (2015). fMRI Study of social anxiety during social ostracism with and without emotional support. PloS one, 10(5), e0127426.	Exclusion -Inclusion	Cyberball task	46
Novembre, G., Zanon, M., & Silani, G. (2015). Empathy for social exclusion involves the sensory-discriminative component of pain: a within-subject fMRI study. Social cognitive and affective neuroscience, 10(2), 153-164.	Main effect of social pain: Self (Exclusion > Inclusion)	Cyberball task	23
Puetz, V. B., Kohn, N., Dahmen, B., Zvyagintsev, M., Schüppen, A., Schultz, R. T., ... & Konrad, K. (2014). Neural response to social rejection in children with early separation experiences. Journal of the American Academy of Child & Adolescent Psychiatry, 53(12), 1328-1337.	Controls SoEx>TechEx	Cyberball task	26
Radke, S., Seidel, E. M., Boubela, R. N., Thaler, H., Metzler, H., Kryspin-Exner, I., ... & Derntl, B. (2018). Immediate and delayed neuroendocrine responses to social exclusion in males and females. Psychoneuroendocrinology, 93, 56-64.	Social exclusion > Inclusion	Social exclusion task and cover story (ball tossing game)	80
van der Meulen, M., Steinbeis, N., Achterberg, M., van IJzendoorn, M. H., & Crone, E. A. (2018). Heritability of neural reactions to social exclusion and prosocial compensation in middle childhood. Developmental cognitive neuroscience, 34, 42-52.	Exclusion > Inclusion	Prosocial Cyberball Game	283
Wagels, L., Bergs, R., Clemens, B., Bauchmüller, M., Gur, R. C., Schneider, F., ... & Kohn, N. (2017). Contextual exclusion processing: an fMRI study of rejection in a performance-related context. Brain imaging and behavior, 11(3), 874-886.	Exclusion > Inclusion	mental visualisation abilities while playing a ball-tossing game	40

Will, G. J., van Lier, P. A., Crone, E. A., & Güroğlu, B. (2016). Chronic childhood peer rejection is associated with heightened neural responses to social exclusion during adolescence. Journal of abnormal child psychology, 44(1), 43-55.	Social exclusion (Exclusion: not receiving the ball > Inclusion: receiving the ball) Social exclusion (Exclusion: not receiving the ball > Inclusion: not receiving the ball)	Cyberball task	44
Chester, D. S., Lynam, D. R., Milich, R., & DeWall, C. N. (2018). Neural mechanisms of the rejection–aggression link. *Social cognitive and affective neuroscience*, *13*(5), 501-512.	Rejection > Acceptance	Cyberball task	60
Wudarczyk, O. A., Kohn, N., Bergs, R., Gur, R. E., Turetsky, B., Schneider, F., & Habel, U. (2015). Chemosensory anxiety cues moderate the experience of social exclusion–an fMRI investigation with Cyberball. Frontiers in psychology, 6, 1475.	Exclusion > Inclusion for: (a) sports chemosensory condition;	Cyberball task	24

List of articles that were selected. for social exclusion.

Meta-analyses were conducted with the activation likelihood estimation (ALE) approach using the Brainmap's GingerALE application. The revised ALE meta-analysis by Eickhoff and colleagues (2009) treats activation foci not as single point, but as spatial probability distributions centered at the given coordinates (Eickhoff et. al, 2012). It models spatial uncertainty by using an estimation of the inter-subject and inter-laboratory variability (typically observed in neuroimaging experiments). An ALE map is obtained by computation of union of activation probabilities for each voxel of all included experiments; and a permutation procedure is used to test for true convergence vs. random clustering (Eickhoff, et. al, 2012). The inference is done through the use of random-effects analysis that calculates the above-chance clustering between experiments. Furthermore, the algorithm gives more weight to gray matter compared to white matter by limiting the meta-analysis to an anatomically constrained space specified by a gray matter mask. For each single meta-analysis, we used the MNI152 coordinate system and the less conservative (larger) mask size. For hunger and thirst, there were 20 articles, 44 experiments, 856 subjects and 612 foci. (Hunger and thirst were merged together as physiological 'needing'). For social exclusion, there were 26 articles, 33 experiments, 1511 subjects and 342 foci. In our study, for main individual meta-analyses, all maps were thresholded using a cluster-level family-wise error (cFWE) correction ($P < 0.05$) with a cluster-forming threshold of $P <$

0.001(uncorrected at the voxel level) (Eklund et al., 2016; Woo et al., 2014), and 1000 permutations. The contrasts analyses ([Physiological-Need] > [Social-Need] and [Social-Need] > [Physiological-Need]) compared the two different datasets (i.e. the ALE results from the Physiological-Need and Social-Need meta-analyses) for statistically significant differences. The conjunction analysis ([Physiological-Need] AND [Social-Need]) was performed by intersecting the thresholded maps for physiological and social needs and allowed us to identify potential brain areas that were consistently activated during both physiological and social needs. Maps from meta-analyses were overlaid on a MNI template using Mango (http://ric.uthscsa.edu/mango/).

Spatial correlation with neurotransmitters

Using the conjunction results ([Physiological-Need] AND [Social-Need]) , we also looked at possible spatial (topographical) relationships between this intersection map and neurotransmitters distribution whole-brain maps available in the JuSpace (as of december 2021). The latter is a MATLAB based toolbox introduced by Dukart and colleagues (2021) which allows for cross-modal correlation of spatial patterns of MRI or fMRI based measures with PET derived biological distribution of specific tissue properties, i.e. receptor density estimates covering dopaminergic, serotonergic, noradrenergic receptors and/or transporters (Dukart et al., 2021). Significant spatial association between subregions of the conjunction results and each neurotransmitter map were examined by comparing the distribution of z-transformed correlations (adjusted for spatial autocorrelation (i.e. adjusting for local gray matter probabilities as estimated from TPM.nii provided with SPM12)) against null distribution using one-sample t-tests (Dukart et al., 2021).

RESULTS

Single meta-analyses

The single meta-analysis on Physiological-Need, i.e. food and water perception during hunger & thirst (combined), revealed consistent activation within the following regions: the bilateral anterior insula, right middle insula, bilateral posterior insula, right claustrum, right putamen, bilateral ACC (B24 and B25), the bilateral caudate head, right caudate body, the left parahippocampal gyrus, the left medial frontal gyrus, right amygdala, right uncus, right mamillary body and right hippocampus (see table 5 and figure 1).

Table 5. Hunger & thirst

Cluster #	x	y	z	ALE	P	Z	Label (Nearest Gray Matter within 5mm)
1	38	10	6	0.031111395	2.5334245E-8	5.448993	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
1	42	4	-10	0.02957781	7.35114E-8	5.2562838	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
1	40	-4	2	0.027335959	3.4063015E-7	4.966653	Right Cerebrum.Sub-lobar.Claustrum.Gray Matter.*
1	46	-12	4	0.02393658	3.1988484E-6	4.51281	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
1	34	-12	6	0.02189089	1.19065735E-5	4.2257686	Right Cerebrum.Sub-lobar.Lentiform Nucleus.Gray Matter.Putamen
1	34	20	-6	0.021366583	1.654994E-5	4.1510224	Right Cerebrum.Sub-lobar.Claustrum.Gray Matter.*
1	28	-8	0	0.019493565	5.3015683E-5	3.8763602	Right Cerebrum.Sub-lobar.Lentiform Nucleus.Gray Matter.Putamen
2	-38	0	-2	0.031286273	2.233257E-8	5.4713836	Left Cerebrum.Sub-lobar.Claustrum.Gray Matter.*
2	-42	-12	10	0.022954691	6.046834E-6	4.375899	Left Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
2	-40	8	12	0.017407449	1.8151532E-4	3.565596	Left Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
3	14	36	-4	0.022025552	1.0891327E-5	4.245793	Right Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 24
3	10	24	0	0.021441728	1.5834556E-5	4.1611233	Right Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head
3	8	20	-12	0.02108268	1.985968E-5	4.1091094	Right Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 25

3	14	10	6	0.018935341	7.378362E-5	3.7951293	Right Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Body
3	10	14	-2	0.0177747	1.4716572E-4	3.620239	Right Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head
4	-26	2	-18	0.024686959	1.9669078E-6	4.61486	Left Cerebrum.Limbic Lobe.Parahippocampal Gyrus.Gray Matter.Brodmann area 34
4	-14	18	-20	0.020973139	2.128351E-5	4.093089	Left Cerebrum.Frontal Lobe.Medial Frontal Gyrus.Gray Matter.Brodmann area 25
4	-14	22	-12	0.017613713	1.6152841E-4	3.5960736	Left Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head
4	-6	22	-10	0.015441236	5.697342E-4	3.2536144	Left Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 24
5	26	6	-16	0.02176692	1.2846617E-5	4.2086267	Right Cerebrum.Sub-lobar.Lentiform Nucleus.Gray Matter.Putamen
5	30	-10	-20	0.02068114	2.5509367E-5	4.0509133	Right Cerebrum.Limbic Lobe.Parahippocampal Gyrus.Gray Matter.Amygdala
5	32	-4	-26	0.019510057	5.2373212E-5	3.8793273	Right Cerebrum.Limbic Lobe.Parahippocampal Gyrus.Gray Matter.Amygdala
5	34	4	-28	0.019396378	5.597474E-5	3.8631182	Right Cerebrum.Limbic Lobe.Uncus.Gray Matter.Brodmann area 28
6	14	-16	0	0.027833832	2.4450569E-7	5.030599	Right Cerebrum.Sub-lobar.Thalamus.Gray Matter.Mammillary Body
7	32	-38	-4	0.021315122	1.7078451E-5	4.1438227	Right Cerebrum.Temporal Lobe.Sub-Gyral.Gray Matter.Hippocampus
7	36	-22	-4	0.02061068	2.6645188E-5	4.0407095	Right Cerebrum.Sub-lobar.Lentiform Nucleus.Gray Matter.Putamen

Coordinates for peak activated clusters in the hunger & thirst condition. We used a cluster-level family-wise error (cFWE) correction (P < 0.05) with a cluster-forming threshold of P < 0.001(uncorrected at the voxel level) (Eklund et al., 2016; Woo et al., 2014), and 1000 permutations. cluster-level family-wise error (cFWE) correction (P < 0.05) with a cluster-forming threshold of P < 0.001(uncorrected at the voxel level) (Eklund et al., 2016; Woo et al., 2014), and 1000 permutations.

The single meta-analysis on Social-Need, i.e. social interaction perception during/after social exclusion, revealed consistent activation within the right anterior insula, bilateral posterior insula, bilateral ACC (B24 and B32), right inferior frontal gyrus, right OFC, left anterior transverse temporal gyrus, and bilateral caudate head (see table 6 and figure 1).

Table 6. Social exclusion

Cluster #	x	y	z	ALE	P	Z	Label (Nearest Gray Matter within 5mm)
1	40	-16	18	0.055232868	3.9646563E-16	8.055735	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
1	48	-10	8	0.021314455	1.0559885E-5	4.252716	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
2	6	38	-6	0.02248284	5.194597E-6	4.408918	Right Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 24
2	2	44	-12	0.021057572	1.2254903E-5	4.219271	Right Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 32
2	-2	36	-16	0.019663427	2.7848786E-5	4.030337	Left Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 32
3	50	32	-8	0.027039705	2.7960468E-7	5.00482	Right Cerebrum.Frontal Lobe.Inferior Frontal Gyrus.Gray Matter.Brodmann area 45
3	38	28	-6	0.020422611	1.7853532E-5	4.1336384	Right Cerebrum.Frontal Lobe.Inferior Frontal Gyrus.Gray Matter.Brodmann area 47
3	38	36	-14	0.016634589	1.7061671E-4	3.5818017	Right Cerebrum.Frontal Lobe.Inferior Frontal Gyrus.Gray Matter.Brodmann area 47
4	-36	-16	20	0.023092672	3.5666646E-6	4.4896793	Left Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
4	-48	-20	12	0.022719312	4.485696E-6	4.4405947	Left Cerebrum.Temporal Lobe.Transverse Temporal Gyrus.Gray Matter.Brodmann area 41
5	10	12	0	0.02310413	3.5442677E-6	4.491021	Right Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head
5	0	12	-6	0.019869018	2.4631125E-5	4.0591025	Left Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head
5	0	18	-4	0.019038977	3.9894312E-5	3.945036	Left Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head

Coordinates for peak activated clusters in the social exclusion condition. We used a cluster-level family-wise error (cFWE) correction (P < 0.05) with a cluster-forming threshold of P < 0.001(uncorrected at the voxel level) (Eklund et al., 2016; Woo et al., 2014), and 1000 permutations. cluster-level family-wise error (cFWE) correction (P < 0.05) with a cluster-forming threshold of P < 0.001(uncorrected at the voxel level) (Eklund et al., 2016; Woo et al., 2014), and 1000 permutations.

Figure 1

Single meta-analyses maps

Maps for activated clusters in each condition: Physiological-Need (green) and Social-Need (blue) and, showing activation patterns for each.

Contrasts meta-analyses

Contrasts meta-analyses results are summarized in tables 7 and 8 and figure 2. Compared to perceiving social interaction during social exclusion, perceiving food or water during hunger or thirst elicited more consistent activation within the bilateral posterior insula, right OFC and the bilateral caudate. Compared to perception of food or water during hunger or thirst, perception of social interaction during social exclusion elicited more consistent activation within the right posterior insula, the right OFC, right inferior frontal gyrus, and left ACC (B32).

Table 7. [Hunger & thirst] minus [Social exclusion]

Cluster #	x	y	z	ALE	P	Z	Label (Nearest Gray Matter within 5mm)

1	40	-16	18	0.055232868	3.9646563E-16	8.055735	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
1	48	-10	8	0.021314455	1.0559885E-5	4.252716	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
2	6	38	-6	0.02248284	5.194597E-6	4.408918	Right Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 24
2	2	44	-12	0.021057572	1.2254903E-5	4.219271	Right Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 32
2	-2	36	-16	0.019663427	2.7848786E-5	4.030337	Left Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 32
3	50	32	-8	0.027039705	2.7960468E-7	5.00482	Right Cerebrum.Frontal Lobe.Inferior Frontal Gyrus.Gray Matter.Brodmann area 45
3	38	28	-6	0.020422611	1.7853532E-5	4.1336384	Right Cerebrum.Frontal Lobe.Inferior Frontal Gyrus.Gray Matter.Brodmann area 47
3	38	36	-14	0.016634589	1.7061671E-4	3.5818017	Right Cerebrum.Frontal Lobe.Inferior Frontal Gyrus.Gray Matter.Brodmann area 47
4	-36	-16	20	0.023092672	3.5666646E-6	4.4896793	Left Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
4	-48	-20	12	0.022719312	4.485696E-6	4.4405947	Left Cerebrum.Temporal Lobe.Transverse Temporal Gyrus.Gray Matter.Brodmann area 41
5	10	12	0	0.02310413	3.5442677E-6	4.491021	Right Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head
5	0	12	-6	0.019869018	2.4631125E-5	4.0591025	Left Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head
5	0	18	-4	0.019038977	3.9894312E-5	3.945036	Left Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head

Coordinates for peak activated clusters in the [Hunger & thirst] minus [Social exclusion] contrast. We used the two cFWE corrected maps with p < .01 (uncorrected at the voxel level), 10,000 permutations (see Eickhoff et al., 2011).

Table 8. [Social exclusion] minus [Hunger & thirst]

Cluster #	x	y	z	P	Z	Label (Nearest Gray Matter within 5mm)
1	42.4	-16.1	20.8	2.0E-4	3.5400841	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
2	48	36	-8	3.0E-4	3.4316144	Right Cerebrum.Frontal Lobe.Middle Frontal Gyrus.Gray Matter.Brodmann area 47
2	44	36	-8	4.0E-4	3.352795	Right Cerebrum.Frontal Lobe.Inferior Frontal Gyrus.Gray Matter.Brodmann area 47
2	50	28	-6	0.0035	2.6968443	Right Cerebrum.Frontal Lobe.Inferior Frontal Gyrus.Gray Matter.*

3	-2	44	-14	0.0026	2.794376	Left Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 32

Coordinates for peak activated clusters in the [Social exclusion] minus [Hunger & thirst] contrast. We used the two cFWE corrected maps with p < .01 (uncorrected at the voxel level), 10,000 permutations (see Eickhoff et al., 2011).

Figure 2

Contrasts maps

In green, clustered thresholded maps for clusters of subtraction {[Physiological-Need] minus [Social-Need]}. In blue, clustered thresholded maps for clusters of subtraction {[Social-Need] minus [Physiological-Need]}.

Conjunction meta-analysis

The intersection between [Physiological-Need] AND [Social-Need] showed overlapping consistent activation in the right posterior insula, right caudate head, and right ACC (B24). (see table 9 and figure 3). It should be noted that the cluster size of the ACC, 8 mm^3, is below what is usually used as minimum, i.e. 10 mm^3.

Table 9. [Hunger & thirst] AND [Social exclusion]

Cluster #	x	y	z	ALE	Label (Nearest Gray Matter within 5mm)
1	48	-10	6	0.019197641	Right Cerebrum.Sub-lobar.Insula.Gray Matter.Brodmann area 13
2	10	14	-2	0.017616214	Right Cerebrum.Sub-lobar.Caudate.Gray Matter.Caudate Head
3	10	38	-6	0.014797952	Right Cerebrum.Limbic Lobe.Anterior Cingulate.Gray Matter.Brodmann area 24

Coordinates for the intersection between s in the [Hunger & thirst] condition AND [Social exclusion] condition. the conjunction was the intersection of the two cFWE thresholded maps.

Figure 3

Conjunction maps

Clustered thresholded maps showing the intersection between activation patterns of [Physiological-Need] AND [Social-Need].

Spatial correlation between the conjunction map and neurotransmitter receptors

Finally, we also looked at the topographical relationship between brain coordinates found in our conjuncion analysis [Physiological-Need] AND [Social-Need] and whole-brain maps of various neurotransmitters (with a focus on serotonin receptors) available in the JuSpace. Because (low) serotonin levels in the brain have been related to sensitivity to food (van Galen et al., 2021) and social exclusion (Preller et al., 2015), our focus was on serotonin receptors. Among serotonin receptors, significant positive spatial correlation between the regions showing consistent activation for physiological and social need and neurotransmitter maps were found for 5HT4 (see figure 4). Although this was the highest correlation with serotonin receptors, it was not the highest of all receptors (see supplementary material for full results). Regarding this, positive correlations were also found for : D1 and D2 dopamine receptors, DAT dopamine transporter, VAchT acetylcholine transporter, SERT serotonin transporter, 5HT4 and 5HT1b serotonin receptors, the Mu opioid receptor, mGluR5 metabotropic glutamate receptor. We found negative correlations for 5HT2a and 5 HT1a serotonin receptors, and CB1 endocannabinoid receptors. This would indicate that dopamine, endogenous opioids, acetylcholine as well as serotonin neurotransmitters are spatially correlated with the intersection network.

Figure 4

Conjunction map correlation with serotonin receptors PET maps

Spatial correlation between the conjunction map and serotonin receptors

Error bars showing the Fisher's z Pearson correlation between the conjunction data on the y axis and the neurotransmitter map based on PET studies on the x axis. The colored dots represent data points and the black ones the mean of the bar which equals the to the single point.

DISCUSSION

Our goal was to investigate possible common and specific brain activation patterns for, on one hand, the processing of physiologically (hunger and thirst) and socially (social exclusion) needed stimuli. To achieve this objective, we used ALE neuroimaging meta-analysis, comparing consistent brain activation patterns during processing of relevant (deprived) stimuli while in physiological vs. social deprivation states. We first carried out separate single meta-analyses on

physiological 'needing' and on social 'needing'. For perception of food and water in hungry and thirsty states, we found consistent activity within the insula, claustrum, putamen, ACC (B24 and B25), caudate head and body, the parahippocampal gyrus, the medial frontal gyrus, amygdala, uncus, mamillary body, and hippocampus . Perception of social interaction during social exclusion revealed consistent activation within the insula, ACC (B24 and B32), inferior frontal gyrus, OFC, anterior transverse temporal gyrus, and caudate head. We then contrasted those maps to identify differences. Compared to social 'needing', physiological 'needing' more consistently elicited activity within the putamen, claustrum, anterior insula, OFC, thalamus, mammillary body and amygdala. Compared to physiological 'needing', social 'needing' more consistently activated the posterior insula, the more lateral part of OFC, inferior frontal gyrus, and ACC. We also intersected the maps of physiological 'needing' and social 'needing' to identify common brain areas between physiological and social 'needing'. In that regard, our results suggest that both processing a physiological and social stimulus we are deprived of (i.e. 'needing') seems to consistently activate parts of the posterior insula, ACC and caudate. Furthermore, we used the spatial pattern of that conjuncion map in order to identify a possible relation between this common network and some neurotransmitter receptors with a focus on serotonergic receptors. Our analysis showed that of all serotonin receptors, the 5HT4 receptor had the highest spatial correlation. In the following paragraphs, we will discuss how these results can help us understand how the brain processes relevant stimuli in a deprived state.

Our single meta-analyses results are in line with the literature. Specifically, previous studies reported activation of insula and ACC for perception of food when hungry (van der Lan et al., 2011; Goldstone, et. al., 2009), water when thirsty (De Araujo et al., 2003), and social interaction when excluded (Mwilambwe & Spreng, 2021). Similar to these, we found consistent activation within the ACC and insula during perception of the physiologically and socially needed stimuli in our single meta-analyses. In that sense, our results show that there is indeed an overlap between processing of physiologically needed stimuli and socially needed stimuli. The contrast [Physiological-Need] minus [Social-Need] revealed activation within the posterior insula, dorsal ACC, pregenual ACC, OFC, and caudate head, whereas the contrast [Social exclusion] minus [Hunger & thirst] did not include the caudate, but also revealed activation within the posterior insula, a more lateral part of the OFC and pregenual ACC. This suggests that neural populations

treating physiologically and/or socially needed stimuli might be spatially closer (or even be the same)i the posterior insula and pregenual ACC. However, for other regions, (such as the OFC) that seem to be activated by both types of need when looking at a macro level, the activations are in different sub-regions when taking a closer look.

States such as hunger and thirst are referred to as homeostatic emotions (Craig, 2003). Despite how they are generated, homeostatic emotions have two important components : the aversive affect, and the affective motivation to terminate that affect (Craig, 2003). More specifically, thalamocortical projections provide both information about (1) the physiological condition of the body in interoceptive cortex at the dorsal and posterior part of the insula ; as well as (2) activation of limbic motor cortex, i.e., the ACC (Craig, 2003). These respectively generate the affective perception and motivation components (Craig, 2003). This is in line with our single meta-analysis results on hunger/food and thirst/water showing consistent activation within the posterior insula and the ACC.

Social exclusion has been said to activate the same dorsal region within the ACC as physical pain (Eisenberg, 2012). However, a recent meta-analysis of social exclusion with the cyberball task found a more ventral part of the ACC (Milambwe & Spreng, 2021). This latter result is similar to our single meta-analysis findings. So, it's possible that social exclusion and physical pain have slightly different brain activity patterns. Also, though social exclusion has been linked to activation of the anterior insula rather than the posterior insula (Eisenberg, 2012), our single meta-analysis on social exclusion found activity in both the posterior insula and in the anterior insula. This is in line with Milambwe & Spreng (2021) recent meta-analysis on social exclusion. However, Vijayakumar and colleagues (2017) did not find the insula in their social exclusion meta-analysis. The difference between our findings and Vijayakumar and colleagues (2017) findings might be related to technical issues or inclusion criteria. Moreover, our study went further than the preceding meta-analyses by investigating the brain activation pattern between perception of social interaction while excluded and that of food/water when hungry/thirsty, enabling us to further assess the need aspect of social excursion in terms of brain patterns.

Some unanswered questions regarding need states might be hinted by looking at the common brain activation pattern map between psychologically and socially needed stimuli. Regarding this, our study shows that consistent brain activity patterns during perception of social interaction while being excluded partly intersects with that of perception of physiologically needed stimuli. This intersection was found within the posterior insula, ventral ACC and caudate which are related to aversive affects, affective motivation and goal directed behavior, respectively (Craig, 2003; Loonen & Ivanova., 2018; Balleine and O'Doherty, 2010). These three regions might be working in the following way: (1) An aversive affect perception of 'needing' is likely processed within the dorsal posterior insula (Craig, 2003), which integrates aversive emotional states and other homeostatic bodily functions (Gerlach et al., 2019) and likely signals a difference between actual and desired interoceptive state (Livneh et al., 2020; Barrett & Simmon, 2015). (2) The ACC contributes in facilitating directional motivation (Craig, 2003) possibly signaling the fulfillment of 'needing' as a requirement and distributing that signal to other brain regions (Weston, 2012) including the caudate (Peak et al., 2019). (3) Activation of the latter might indicate that perceiving needed stimuli leads to goal directed behavior and action choice (Balleine and O'Doherty, 2010; Knutson & Cooper, 2005; Hollon et al., 2014; ito and Doya, 2015; Schwabe & Wolf, 2010), based on the current need (van den Bercken & Cools, 1982). Thus, our method and idea of looking at the common brain areas between psychologically and socially needed stimuli allowed us to further elucidate this mechanism. Following that, our findings suggest that perception of both physiologically and socially needed stimuli embed the affective perception, processed within the posterior insula, and the affective motivation toward the stimuli/event that terminates that need state, processed within the ventral ACC, which directs action choice within the caudate (Peak et al., 2019). Though this was initially suggested for homeostatic/physiological needs and their relevant stimuli (see Craig, 2003), our results lead us to suggest that this might be also true for social need states and social interaction.

As discussed in the previous paragraph, a possible link between social exclusion and hunger/thirst could be that they are both related to some displeasure/aversive state. This might also explain the inconsistency in the relationship between social exclusion and pain (Eisenberg, 2012; Milambwe & Spreng, 2021): social exclusion might be related, not to pain, but to displeasure. The latter is the inverse of pleasure (Cabanac, 2002) and as such includes, but is not

limited to, pain. Indeed, displeasure can also be related to hunger, shortness of breath, disgust, depression, anxiety, fear, etc. (Becker et al., 2019; Berridge & Kringelbach, 2015) and is said to be processed within the pregenual ACC (and mesolimbic and amygdalar circuitry) (Becker et al., 2019), a region that we identified as being commonly recruited during both physiological and social needs.

A recent study by Tomova and colleagues (2020) on the common brain areas between perception of food during hunger and social interaction after isolation only found common activation within the substantia nigra (SN) and ventral tegmental area (VTA). These are dopaminergic regions, and the VTA mesolimbic dopamine has been linked to reward prediction, reward learning and motivation (Hamid et al., 2016). However, in our conjunction analysis, we did not find the VTA/SN nor the ventral striatum which receives VTA dopamine for action invigoration and reward seeking (Li and Daw, 2011; Berridge and Aldrige, 2009; Lex and Hauber, 2008; Hamid et al., 2016; Zhang et al. 2009; Balleine and Killcross, 2006). Though it is a possibility that the difference between our findings and Tomova's is due to the difference between social exclusion and social isolation, other studies on perception of social cues from isolated or lonely participants have not consistently found activity within the mesolimbic dopaminergic VTA or ventral striatum (see Cacioppo et al., 2009; D'agostino et al., 2019). Conversely, Tomova and colleagues (2020) findings might be related to a more pavlovian like dopaminergic conditioned reaction, which might be related to the condition, the task and stimuli used. Indeed, although the task was about (passively) viewing either food or social stimuli (similar to our inclusion criteria), those stimuli were "tailored to each individual's preferred foods and modes of social interaction. During the initial screening, participants were asked to list their top ten favorite foods and social activities." (Tomova et al., 2020). (This was not the case for included studies in the present meta-analysis.) It is possible that these initial preferences facilitated conditioned responses related to reward prediction which are related to dopamine activation in the VTA (Schultz et al., 1997; Schultz, 2015; Montague et al., 1996; Schultz, 1998; Rice et al., 2010). This might be a plausible explanation as other studies and meta-analyses on either thirst, hunger, social exclusion or isolation where participants were passively observing social stimuli have not emphasized dopaminergic activity (ex, van der Lan et al., 2011; Goldstone, et. al., 2009; De Araujo et al., 2003; Mwilambwe-Tshilobo & Spreng, 2021; Vijayakumar et al., 2017; Cacioppo et al., 2009;

D'Agostino et al., 2019). We should also keep in mind that the size of clusters of VTA/SN found in Tomova et al. (2020) might be too small for usual fMRI, and thus could have been activated, but not "captured", in those other studies. Future studies are thus needed to clarify the role of the dopaminergic system during the processing of physiologically and socially needed stimuli. .

Our aim was to go beyond common brain activity patterns between psychologically and socially needed stimuli, by running exploratory analysis to look at how these common brain patterns–posterior insula, caudate and ACC–might be related to serotonergic receptors distribution in the brain. Our results show that at the conjuncion regions between physiological and social 'needing', the serotonin receptor with highest density is the 5HT4 receptor. Our finding is in line with the suggestion that the 5HT4 receptor is a component of a feedback loop from the medial PFC to the dorsal raphe nuclei (DRN) (Rebholz, et al., 2018), specifically the prelimbic and infralimbic subregions (Peyron et al 1998, Lucas et al, 2005), which correspond to Brodmann areas 32 and 25 respectively (Price, 2007); which are part of the ACC (Weston, 2012). The ACC activity is related to both an affective motivation and an update of internal models, i.e. a feedback loop (Craig, 2003; Kolling et al., 2016; Petzschner et al., 2021). Some information from this feedback loop is said to be sent from the ACC to the dorsal raphe nucleus (DRN) (Rebholz et al., 2018; Lucas et al., 2005) which is the largest serotonergic structure in the brain (Liu et al., 2020). Our finding related to the spatial correlation between 5HT4 and the common brain map, leads us to suggest that this feedback loop can allow need states or needed stimuli to influence activity in the DRN through the ACC. That influence can be either inhibitory or excitatory via modulation of GABAergic (DRN inhibition) neurons and CB1 receptors (DRN excitation) in the DRN by the ACC (Lucas et al., 2005; Castello, et al 2018; Geddes et al.2016). Indeed CB1 has been found to be implicated in pleasure for both food (Kirkham, 2009) and social play (Achterberg et al., 2016). This might give some reason why needed stimuli are pleasurable. However we did not find much correlation between the common pattern map and CB1; suggesting that that CB1 actions could be further downstream, closer to the DRN rather than closer to the common brain map found in this study: posterior insula, pregenual ACC and caudate. Furthermore, though our findings differ from those of Tomova regarding the SN/VTA activation; our finding, that the intersection network is correlated to 5HT4 distribution, could partly explain the findings of Tomova and colleagues. VTA\SN is modulated by DRN (Gervais

& Rouillard, 2000), whose activity, as our study and literature suggest, is regulated by ACC via the 5HT4 receptor. In that sense, our results provide some possible explanation on the mechanism underlying Tomova and colleagues (2020) findings: via 5HT4, ACC regulates the serotonergic DRN, which in turn influences VTA/SN dopaminergic activity. In summary, our study suggests that the perception of a stimulus that would alleviate a negative physiological or social state (need) could be linked to brain regions that influence the activity of serotonin neurons.

LIMITS

The main limit of our study is that for physiological needs we only included hunger and thirst, while for social needs we only included social exclusion and not isolation. Nevertheless, it has been argued that both hunger and thirst implicate a similar network, that includes the ACC and insula (Mckinley et al., 2019), and the mechanism of that network is similar to other needs related to temperature, itch, visceral distension, muscle ache, 'air hunger', etc. (Crag, 2003). In the same way, even though social exclusion is different from social isolation (social isolation is more related to meaningful social contacts whereas exclusion is more related to being outcast of, or not able to participate in, some society (Huisman & van Tilburg, 2021)), they have in common the fact that they refer to "lack of ties", either with with society (exclusion) or with other significant persons (isolation) (Huisman & van Tilburg, 2021). Although there has been a debate between whether social exclusion actually causes need related distress (Gerber & Wheeler, 2009; Blackhart et al., 2009), Bernstein and Claypool (2012) found that Cyberball exclusion paradigm, used in this paper, may induce need-like emotional distress (e.g., reduced mood, and lowered self-esteem and other needs). Hence, the use of cyberball in this study is more coherent with a need state (Bernstein & Claypool, 2012). Future studies could assess a larger range of physiologically and socially needed stimuli, including those not included in the present study. Furthermore, one should not forget the limitations of reverse inference (Poldrack, 2006; 2011) in interpreting our results.

CONCLUSION

Our goal was to study the common and specific brain activations during physiological (hunger and thirst) and social (social exclusion/isolation) 'needing' as well as their relationship to the serotonergic system. Our results suggest that regions within the mid-posterior insula, the ACC and the caudate are regions that commonly support processing/perception of both physiologically (hunger and thirst) and socially (exclusion) needed stimuli. So our result lead us to propose that 'needing' whether physiologically or socially is related to (1) am affective perception or response towards the needed stimulus that signal difference between actual and desired state, and which is processed within in the mid-posterior insula ; (2) an affective and directional motivation that requires the termination of the need state, processed within the ACC. (3) This requirement to terminate the need state facilitates goal directed behavior within the caudate. Furthermore, The network of regions at the intersection seem to be related to the distribution of receptors, and among the serotonergic receptors, the 5HT4 seem to have one the highest spatial correlation with that network. We hypothesize that, in need state and/or while processing the needed stimuli, this intersection network, through 5HT4, modulates DRN serotonin activity which signals how beneficial, or not, the current state is. In that sense, in the brain, physiological and social deprivation could lead to low serotonin levels, whereas the onset or presence of physiologically and socially needed stimuli could be related to high serotonin levels.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

ACKNOWLEDGMENTS

The research was supported in part by NSERC Discovery Grant #RGPIN-2018-05698 and UdeM institutional funds.

AUTHOR CONTRIBUTION

Juvénal Bosulu: Designed the study, performed the database search, performed data analysis, interpretation, and wrote the manuscript. **Sébastien Hétu**: Revised the manuscript and provided critical feedbacks. **Yi Luo**: Revised the manuscript and provided critical feedbacks. All authors contributed to and approved the final manuscript version.

DATA AVAILABILITY STATEMENT

All Data are available upon request.

REFERENCES

Achterberg, E. M., van Swieten, M. M., Driel, N. V., Trezza, V., & Vanderschuren, L. J. (2016). Dissociating the role of endocannabinoids in the pleasurable and motivational properties of social play behaviour in rats. *Pharmacological research*, *110*, 151-158.
https://doi.org/10.1016/j.phrs.2016.04.031

Balleine, B. W., & Killcross, S. (2006). Parallel incentive processing: an integrated view of amygdala function. *Trends in neurosciences, 29(5),* 272-279.
https://doi.org/10.1016/j.tins.2006.03.002

Balleine, B. W., & O'doherty, J. P. (2010). Human and rodent homologies in action control: corticostriatal determinants of goal-directed and habitual action. *Neuropsychopharmacology, 35(1),* 48-69.
https://doi.org/10.1038/npp.2009.131

Barrett, L. F., & Simmons, W. K. (2015). Interoceptive predictions in the brain. *Nature reviews neuroscience, 16(7),* 419-429.
https://doi.org/10.1038/nrn3950

Baumeister, R. F., & Leary, M. R. (1995). The need to belong: Desire for interpersonal attachments as a fundamental human motivation. *Psychological Bulletin, 117*(3), 497–529. https://doi.org/10.1037/0033-2909.117.3.497

Becker, C. A., Schmälzle, R., Flaisch, T., Renner, B., & Schupp, H. T. (2015). Thirst and the state-dependent representation of incentive stimulus value in human motive circuitry. Social cognitive and affective neuroscience, 10(12), 1722-1729. https://doi.org/10.1093/scan/nsv063

Becker, C. A., Flaisch, T., Renner, B., & Schupp, H. T. (2017). From thirst to satiety: the anterior mid-cingulate cortex and right posterior insula indicate dynamic changes in incentive value. Frontiers in human neuroscience, 11, 234. https://doi.org/10.3389/fnhum.2017.00234

Becker, S., Bräscher, A. K., Bannister, S., Bensafi, M., Calma-Birling, D., Chan, R. C., ... & Wang, Y. (2019). The role of hedonics in the Human Affectome. *Neuroscience & Biobehavioral Reviews*, *102*, 221-241. https://doi.org/10.1016/j.neubiorev.2019.05.003

Bernstein, M. J., & Claypool, H. M. (2012). Not all social exclusions are created equal: Emotional distress following social exclusion is moderated by exclusion paradigm. *Social Influence*, *7*(2), 113-130. https://doi.org/10.1080/15534510.2012.664326

Berridge, K. C., & Aldridge, J. W. (2009). Decision utility, incentive salience, and cue-triggered "wanting". *Oxford series in social cognition and social neuroscience, 2009,* 509.

Berridge, K. C., & Kringelbach, M. L. (2015). Pleasure systems in the brain. *Neuron*, *86*(3), 646-664. https://doi.org/10.1016/j.neuron.2015.02.018

Bindra, D. (1974). A motivational view of learning, performance, and behavior modification. *Psychological review, 81(3),* 199. https://doi.org/10.1037/h0036330

Blackhart, G. C., Nelson, B. C., Knowles, M. L., & Baumeister, R. F. (2009). Rejection elicits emotional reactions but neither causes immediate distress nor lowers self-esteem: A meta-analytic review of 192 studies on social exclusion. *Personality and Social Psychology Review*, *13*(4), 269-309. https://doi.org/10.1177/1088868309346065

Bolling, D. Z., Pitskel, N. B., Deen, B., Crowley, M. J., McPartland, J. C., Mayes, L. C., & Pelphrey, K. A. (2011). Dissociable brain mechanisms for processing social exclusion and rule violation. NeuroImage, 54(3), 2462-2471.

Bouton, M. E. (2016). *Learning and behavior: A contemporary synthesis* (2nd ed.). Sinauer Associates.

Cabanac, M. (2002). What is emotion?. *Behavioural processes*, *60*(2), 69-83. https://doi.org/10.1016/S0376-6357(02)00078-5

Cacioppo, J. T., Norris, C. J., Decety, J., Monteleone, G., & Nusbaum, H. (2009). In the eye of the beholder: individual differences in perceived social isolation predict regional brain activation to social stimuli. *Journal of cognitive neuroscience*, *21*(1), 83-92. https://doi.org/10.1162/jocn.2009.21007

Castello, J., LeFrancois, B., Flajolet, M., Greengard, P., Friedman, E., & Rebholz, H. (2018). CK2 regulates 5-HT4 receptor signaling and modulates depressive-like behavior. *Molecular psychiatry*, *23*(4), 872-882. https://doi.org/10.1038/mp.2017.240

Chen, J., Papies, E. K., & Barsalou, L. W. (2016). A core eating network and its modulations underlie diverse eating phenomena. Brain and cognition, 110, 20-42. https://doi.org/10.1016/j.bandc.2016.04.004

Chen, E. Y., & Zeffiro, T. A. (2020). Hunger and BMI modulate neural responses to sweet stimuli: fMRI meta-analysis. *International Journal of Obesity, 44(8),* 1636-1652. https://doi.org/10.1038/s41366-020-0608-5

Craig, A. D. (2003). Interoception: the sense of the physiological condition of the body. *Current opinion in neurobiology, 13(4),* 500-505. https://doi.org/10.1016/S0959-4388(03)00090-4

D'Agostino, A. E., Kattan, D., & Canli, T. (2019). An fMRI study of loneliness in younger and older adults. *Social neuroscience*, *14*(2), 136-148. https://doi.org/10.1080/17470919.2018.1445027

Dankoski, E. C., Agster, K. L., Fox, M. E., Moy, S. S., & Wightman, R. M. (2014). Facilitation of serotonin signaling by SSRIs is attenuated by social isolation. *Neuropsychopharmacology*, *39*(13), 2928-2937. https://doi.org/10.1038/npp.2014.162

Dayan, P., & Huys, Q. J. (2009). Serotonin in affective control. *Annual review of neuroscience*, *32*(1), 95-126. https://doi:10.1146/annurev.neuro.051508.135607

De Araujo, I. E., Kringelbach, M. L., Rolls, E. T., & McGlone, F. (2003). Human cortical responses to water in the mouth, and the effects of thirst. *Journal of neurophysiology*, *90*(3), 1865-1876.

Dukart, J., Holiga, S., Rullmann, M., Lanzenberger, R., Hawkins, P. C., Mehta, M. A., ... & Eickhoff, S. B. (2021). *JuSpace: A tool for spatial correlation analyses of magnetic resonance imaging data with nuclear imaging derived neurotransmitter maps, 42(3)*, pp. 555-566). https://doi.org/10.1002/hbm.25244

Egan, G., Silk, T., Zamarripa, F., Williams, J., Federico, P., Cunnington, R., ... & Denton, D. (2003). Neural correlates of the emergence of consciousness of thirst. *Proceedings of the National Academy of Sciences*, *100*(25), 15241-15246. https://doi.org/10.1073/pnas.213665010

Eickhoff, S. B., Laird, A. R., Grefkes, C., Wang, L. E., Zilles, K., & Fox, P. T. (2009). Coordinate-based ALE meta-analysis of neuroimaging data: a random-effects approach based on empirical estimates of spatial uncertainty. *Human brain mapping, 30(9),* 2907. https://doi.org/10.1002/hbm.20718

Eickhoff, S. B., Bzdok, D., Laird, A. R., Kurth, F., & Fox, P. T. (2012). Activation likelihood estimation meta-analysis revisited. *Neuroimage*, *59*(3), 2349-2361. https://doi.org/10.1016/j.neuroimage.2011.09.017

Eisenberger, N. I., Lieberman, M. D., & Williams, K. D. (2003). Does rejection hurt? An fMRI study of social exclusion. *Science, 302*(5643), 290-292. https://DOI: 10.1126/science.10891

Farrell, M. J., Bowala, T. K., Gavrilescu, M., Phillips, P. A., McKinley, M. J., McAllen, R. M., ... & Egan, G. F. (2011). Cortical activation and lamina terminalis functional connectivity during thirst and drinking in humans. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *301*(3), R623-R631.

https://doi.org/10.1152/ajpregu.00817.2010

Führer, D., Zysset, S., & Stumvoll, M. (2008). Brain activity in hunger and satiety: an exploratory visually stimulated FMRI study. Obesity, 16(5), 945-950.

Garavan, H. (2010). Insula and drug cravings. *Brain structure and function, 214(5-6),* 593-601. https://doi.org/10.1038/oby.2008.33

Geddes, S. D., Assadzada, S., Lemelin, D., Sokolovski, A., Bergeron, R., Haj-Dahmane, S., & Béïque, J. C. (2016). Target-specific modulation of the descending prefrontal cortex inputs to the dorsal raphe nucleus by cannabinoids. *Proceedings of the National Academy of Sciences*, *113*(19), 5429-5434. https://doi.org/10.1073/pnas.1522754113

Gehrlach, D. A., Dolensek, N., Klein, A. S., Chowdhury, R. R., Matthys, A., Junghänel, M., Gaitanos, T. N., Podgornik, A., Black, T. D., Vaka, N. R., Conzelmann, K.-K., & Gogolla, N. (2020). Aversive state processing in the posterior insular cortex. *Nature Neuroscience*, 22(9), 1424– 1437. https://doi.org/10.1038/s41593-019-0469-1

Gerber, Jonathan, and Ladd Wheeler. "On being rejected: A meta-analysis of experimental research on rejection." *Perspectives on Psychological Science* 4.5 (2009): 468-488. https://doi.org/10.1111/j.1745-6924.2009.01158.x

Gervais, J., & Rouillard, C. (2000). Dorsal raphe stimulation differentially modulates dopaminergic neurons in the ventral tegmental area and substantia nigra. *Synapse*, *35*(4), 281-291. https://doi.org/10.1002/(SICI)1098-2396(20000315)35:4<281::AID-SYN6>3.0.CO;2-A

Goebel, B. L., & Brown, D. R. (1981). Age differences in motivation related to Maslow's need hierarchy. *Developmental Psychology*, *17*(6), 809.

Goldstone, A. P., Prechtl de Hernandez, C. G., Beaver, J. D., Muhammed, K., Croese, C., Bell, G., ... & Bell, J. D. (2009). Fasting biases brain reward systems towards high‑calorie foods. *European Journal of Neuroscience, 30(8),* 1625-1635. https://doi.org/10.1111/j.1460-9568.2009.06949.x

Hamid, A. A., Pettibone, J. R., Mabrouk, O. S., Hetrick, V. L., Schmidt, R., Vander Weele, C. M., ... & Berke, J. D. (2016). Mesolimbic dopamine signals the value of work. *Nature neuroscience, 19(1),* 117-126. https://doi.org/10.1038/nn.4173

Huisman, M., & van Tilburg, T. G. (2021). Social exclusion and social isolation in later life. In *Handbook of aging and the social sciences* (pp. 99-114). Academic Press. https://doi.org/10.1016/B978-0-12-815970-5.00007-3

Hofstede, G. (1984). *Culture's consequences: International differences in work-related values* (Vol. 5). Sage.

Hollon, N. G., Arnold, M. M., Gan, J. O., Walton, M. E., & Phillips, P. E. (2014). Dopamine-associated cached values are not sufficient as the basis for action selection. Proceedings of the National Academy of Sciences, 111(51), 18357-18362. https://doi.org/10.1073/pnas.1419770111

Hull, C. L. (1943). *Principles of behavior: An introduction to behavior theory.* Appleton-Century.

Inagaki, T. K., Muscatell, K. A., Moieni, M., Dutcher, J. M., Jevtic, I., Irwin, M. R., & Eisenberger, N. I. (2016). Yearning for connection? Loneliness is associated with increased ventral striatum activity to close others. *Social cognitive and affective neuroscience*, *11*(7), 1096-1101. https://doi.org/10.1093/scan/nsv076

Ito, M., & Doya, K. (2015). Distinct neural representation in the dorsolateral, dorsomedial, and ventral parts of the striatum during fixed-and free-choice tasks. *Journal of Neuroscience, 35(8),* 3499-3514.
https://doi.org/10.1523/JNEUROSCI.1962-14.2015

Kirkham, T. C. (2009). Cannabinoids and appetite: food craving and food pleasure. *International Review of Psychiatry*, *21*(2), 163-171.
https://doi.org/10.1080/09540260902782810

Knutson, B., & Cooper, J. C. (2005). Functional magnetic resonance imaging of reward prediction. *Current opinion in neurology, 18(4),* 411-417.
https://doi.org/10.1097/01.wco.0000173463.24758.f6

Kolling, N., Wittmann, M. K., Behrens, T. E. J., Boorman, E. D., Mars, R. B., & Rushworth, M. F. S. (2016). Anterior cingulate cortex and the value of the environment, search, persistence, and model updating. *Nature neuroscience*, *19*(10), 1280.
https://doi.org/10.1038/nn.4382

LaBar, K. S., Gitelman, D. R., Parrish, T. B., Kim, Y. H., Nobre, A. C., & Mesulam, M. (2001). Hunger selectively modulates corticolimbic activation to food stimuli in humans. *Behavioral neuroscience, 115(2)*, 493.
https://doi.org/10.1037/0735-7044.115.2.493

Lancaster, J. L., Tordesillas‑Gutiérrez, D., Martinez, M., Salinas, F., Evans, A., Zilles, K., ... & Fox, P. T. (2007). Bias between MNI and Talairach coordinates analyzed using the ICBM‑152 brain template. *Human brain mapping*, *28*(11), 1194-1205.

https://doi.org/10.1002/hbm.20345

Li, J., & Daw, N. D. (2011). Signals in human striatum are appropriate for policy update rather than value prediction. *Journal of Neuroscience*, *31*(14), 5504-5511. https://doi.org/10.1523/JNEUROSCI.6316-10.2011

Liu, Z., Lin, R., & Luo, M. (2020). Reward contributions to serotonergic functions. *Annu. Rev. Neurosci*, *43*(1), 141-162. https://doi.org/10.1146/annurev-neuro-093019-112252

Livneh, Y., Ramesh, R. N., Burgess, C. R., Levandowski, K. M., Madara, J. C., Fenselau, H., ... & Andermann, M. L. (2017). Homeostatic circuits selectively gate food cue responses in insular cortex. *Nature*, *546*(7660), 611-616. https://doi.org/10.1038/nature22375

Livneh, Y., Sugden, A. U., Madara, J. C., Essner, R. A., Flores, V. I., Sugden, L. A., ... & Andermann, M. L. (2020). Estimation of current and future physiological states in insular cortex. *Neuron, 105(6),* 1094-1111. https://doi.org/10.1016/j.neuron.2019.12.027

Loonen, A. J., & Ivanova, S. A. (2018). Circuits regulating pleasure and happiness: evolution and role in mental disorders. *Acta Neuropsychiatrica*, *30*(1), 29-42. https://doi.org/10.1017/neu.2017.8

Lucas, G., Compan, V., Charnay, Y., Neve, R. L., Nestler, E. J., Bockaert, J., ... & Debonnel, G. (2005). Frontocortical 5-HT4 receptors exert positive feedback on serotonergic activity: viral transfections, subacute and chronic treatments with 5-HT4 agonists. *Biological psychiatry*, *57*(8), 918-925. https://doi.org/10.1016/j.biopsych.2004.12.023

Luo, M., Li, Y., & Zhong, W. (2016). Do dorsal raphe 5-HT neurons encode "beneficialness"?. *Neurobiology of learning and memory*, *135*, 40-49. https://doi.org/10.1016/j.nlm.2016.08.008

MacGregor, D. (1960). *The human side of enterprise* (Vol. 21, No. 166.1960). McGraw‑Hill: New York.

Maner, J. K., DeWall, C. N., Baumeister, R. F., & Schaller, M. (2007). Does social exclusion motivate interpersonal reconnection? Resolving the" porcupine problem.". *Journal of personality and social psychology*, *92*(1), 42.

Maslow, A. H. (1943). A theory of human motivation. *Psychological Review, 50(4)*, 370-96.

Masten, C. L., Eisenberger, N. I., Borofsky, L. A., Pfeifer, J. H., McNealy, K., Mazziotta, J. C., & Dapretto, M. (2009). Neural correlates of social exclusion during adolescence: understanding the distress of peer rejection. Social cognitive and affective neuroscience, 4(2), 143-157.

McKinley, M. J., Denton, D. A., Ryan, P. J., Yao, S. T., Stefanidis, A., & Oldfield, B. J. (2019). From sensory circumventricular organs to cerebral cortex: neural pathways controlling thirst and hunger. *Journal of neuroendocrinology*, *31*(3), e12689. https://doi.org/10.1111/jne.12689

Mohanty, A., Gitelman, D. R., Small, D. M., & Mesulam, M. M. (2008). The spatial attention network interacts with limbic and monoaminergic systems to modulate motivation-induced attention shifts. *Cerebral cortex, 18(11*), 2604-2613. https://doi.org/10.1093/cercor/bhn021

Montague, P. R., Dayan, P., & Sejnowski, T. J. (1996). A framework for mesencephalic dopamine systems based on predictive Hebbian learning. *Journal of neuroscience, 16*(5), 1936-1947. https://doi.org/10.1523/JNEUROSCI.16-05-01936.1996

Mwilambwe-Tshilobo, L., & Spreng, R. N. (2021). Social exclusion reliably engages the default network: A meta-analysis of Cyberball. *NeuroImage*, *227*, 117666.
https://doi.org/10.1016/j.neuroimage.2020.117666

Peak, J., Hart, G., & Balleine, B. W. (2019). From learning to action: the integration of dorsal striatal input and output pathways in instrumental conditioning. *European Journal of Neuroscience*, *49*(5), 658-671.
https://doi.org/10.1111/ejn.13964

Peyron, C., Petit, J. M., Rampon, C., Jouvet, M., & Luppi, P. H. (1997). Forebrain afferents to the rat dorsal raphe nucleus demonstrated by retrograde and anterograde tracing methods. *Neuroscience*, *82*(2), 443-468.
https://doi.org/10.1016/S0306-4522(97)00268-6

Preller, K. H., Pokorny, T., Hock, A., Kraehenmann, R., Stämpfli, P., Seifritz, E., ... & Vollenweider, F. X. (2016). Effects of serotonin 2A/1A receptor stimulation on social exclusion processing. *Proceedings of the National Academy of Sciences*, *113*(18), 5119-5124.
https://doi.org/10.1073/pnas.152418711

Petzschner, F. H., Garfinkel, S. N., Paulus, M. P., Koch, C., & Khalsa, S. S. (2021). Computational models of interoception and body regulation. *Trends in neurosciences, 44(1),* 63-76.
https://doi.org/10.1016/j.tins.2020.09.012

Price, J. L. (2007). Definition of the orbital cortex in relation to specific connections with limbic and visceral structures and other cortical regions. *Annals of the New York Academy of Sciences*, *1121*(1), 54-71.
https://doi.org/10.1196/annals.1401.008

Poldrack, Russell A. "Can cognitive processes be inferred from neuroimaging data?." *Trends in cognitive sciences* 10.2 (2006): 59-63.

https://doi.org/10.1016/j.tics.2005.12.004

Poldrack, Russell A. "Inferring mental states from neuroimaging data: from reverse inference to large-scale decoding." *Neuron* 72.5 (2011): 692-697.
https://doi.org/10.1016/j.neuron.2011.11.001

Rebholz, H., Friedman, E., & Castello, J. (2018). Alterations of expression of the serotonin 5-HT4 receptor in brain disorders. *International Journal of Molecular Sciences*, *19*(11), 3581.
https://doi.org/10.3390/ijms19113581

Rice, M. E., Patel, J. C., & Cragg, S. J. (2011). Dopamine release in the basal ganglia. *Neuroscience*, *198*, 112-137.
https://doi.org/10.1016/j.neuroscience.2011.08.066

Saker, P., Farrell, M. J., Adib, F. R., Egan, G. F., McKinley, M. J., & Denton, D. A. (2014). Regional brain responses associated with drinking water during thirst and after its satiation. *Proceedings of the National Academy of Sciences*, *111*(14), 5379-5384.

Schultz, W. (1998). Predictive reward signal of dopamine neurons. *Journal of neurophysiology, 80(1),* 1-27.
https://doi.org/10.1152/jn.1998.80.1.1

Schultz, W. (2015). Neuronal reward and decision signals: from theories to data. *Physiological reviews, 95(3),* 853-951.
https://doi.org/10.1152/physrev.00023.2014

Schultz, W., Dayan, P., & Montague, P. R. (1997). A neural substrate of prediction and reward. *Science, 275(5306),* 1593-1599.
https://doi.org/10.1126/science.275.5306.1593.

Schwabe, L., & Wolf, O. T. (2010). Learning under stress impairs memory formation. *Neurobiology of learning and memory, 93(2),* 183-188. https://doi.org/10.1016/j.nlm.2009.09.009

Siep, N., Roefs, A., Roebroeck, A., Havermans, R., Bonte, M. L., & Jansen, A. (2009). Hunger is the best spice: an fMRI study of the effects of attention, hunger and calorie content on food reward processing in the amygdala and orbitofrontal cortex. *Behavioural brain research, 198(1),* 149-158. https://doi.org/10.1016/j.bbr.2008.10.035

Sizemore, T. R., Hurley, L. M., & Dacks, A. M. (2020). Serotonergic modulation across sensory modalities. *Journal of Neurophysiology*, *123*(6), 2406-2425. https://doi.org/10.1152/jn.00034.2020

Tay, L., & Diener, E. (2011). Needs and subjective well-being around the world. *Journal of personality and social psychology*, *101*(2), 354.

Toates, F. (1994). Comparing motivational systems—An incentive motivation perspective. In C. R. Legg & D. A. Booth (Eds.), *Appetite: Neural and behavioural bases* (pp. 305–327). Oxford University Press. https://doi.org/10.1093/acprof:oso/9780198547877.003.0013

Tomova, L., Wang, K. L., Thompson, T., Matthews, G. A., Takahashi, A., Tye, K. M., & Saxe, R. (2020). Acute social isolation evokes midbrain craving responses similar to hunger. *Nature Neuroscience*, *23*(12), 1597-1605. https://doi.org/10.1038/s41593-020-00742-z

Tomova, L., Tye, K., & Saxe, R. (2021). The neuroscience of unmet social needs. *Social Neuroscience*, *16*(3), 221-231.

https://doi.org/10.1080/17470919.2019.1694580

Van den Bercken, J. H. L., & Cools, A. R. (1982). Evidence for a role of the caudate nucleus in the sequential organization of behaviour. *Behavioural brain research*, *4*(4), 319-337. https://doi.org/10.1016/0166-4328(82)90058-4

van der Laan, L. N., De Ridder, D. T., Viergever, M. A., & Smeets, P. A. (2011). The first taste is always with the eyes: a meta-analysis on the neural correlates of processing visual food cues. *Neuroimage, 55(1),* 296-303. https://doi.org/10.1016/j.neuroimage.2010.11.055

van Galen, K. A., Ter Horst, K. W., & Serlie, M. J. (2021). Serotonin, food intake, and obesity. *Obesity Reviews*, *22*(7), e13210. https://doi.org/10.1111/obr.13210

Vijayakumar, N., Cheng, T. W., & Pfeifer, J. H. (2017). Neural correlates of social exclusion across ages: A coordinate-based meta-analysis of functional MRI studies. *NeuroImage*, *153*, 359-368. https://doi.org/10.1016/j.neuroimage.2017.02.050

Weston, C. S. (2012). Another major function of the anterior cingulate cortex: the representation of requirements. *Neuroscience & Biobehavioral Reviews*, *36*(1), 90-110. https://doi.org/10.1016/j.neubiorev.2011.04.014

Williams, K. D. (2007). Ostracism. *Annu. Rev. Psychol.*, *58*, 425-452. https://doi.org/10.1146/annurev.psych.58.110405.085641

Williams, K. D., Cheung, C. K., & Choi, W. (2000). Cyberostracism: effects of being ignored over the Internet. *Journal of personality and social psychology*, *79*(5), 748. https://doi.org/10.1037/0022-3514.79.5.748

Zhang, J., Berridge, K. C., Tindell, A. J., Smith, K. S., & Aldridge, J. W. (2009). A neural computational model of incentive salience. *PLoS computational biology, 5(7)*, e1000437. https://doi.org/10.1371/journal.pcbi.1000437