Biodiversity-ecosystem function relationships change in sign and magnitude across the Hill

diversity spectrum

Michael Roswell[1] mroswell@umd.edu **0000-0002-8479-9184**

Tina Harrison[2] tinaharrison09@gmail.com **0000-0001-8735-0289**

Mark A. Genung[2] mark.genung@louisiana.edu **0000-0002-9383-667X**

1 – Department of Entomology, University of Maryland, College Park, MD, USA

2 – Department of Biology, University of Louisiana at Lafayette, Lafayette, LA, USA

Keywords: abundance, biodiversity, diversity profile, ecosystem function, Hill number, rarity

Summary

Motivated by accelerating anthropogenic extinctions, decades of biodiversity-ecosystem function

(BEF) experiments show that ecosystem function declines with species loss from local communities.

Yet, at the local scale, changes in species' total and relative abundances are more common than

species loss. The consensus best biodiversity measures are Hill numbers, which use a scaling

parameter, ℓ, to emphasize rarer versus more common species. Shifting that emphasis captures

distinct, function-relevant biodiversity gradients beyond species richness. Here, we surmised that Hill

numbers that emphasize rare species more than richness may distinguish large, complex, and

presumably higher-functioning assemblages from smaller and simpler ones. In this study, we tested

which values of ℓ produce the strongest BEF relationships in community datasets of ecosystem

functions provided by wild, free-living organisms. We found that ℓ values that emphasized rare species

more than richness most often correlated most strongly with ecosystem functions. As emphasis

shifted to more common species, BEF correlations were often weak and/or negative. We argue that

unconventional Hill diversities that shift emphasis towards rarer species may be useful for describing

biodiversity change, and that employing a wide spectrum of Hill numbers can clarify mechanisms

underlying BEF relationships.

Introduction

A central question in community ecology is, "how will ongoing shifts in biodiversity affect ecosystem function?" In experiments that vary species richness while controlling other community properties, the answer has been clear for some time: ecosystem function has a positive, saturating relationship with species richness [1–3]. There is ongoing interest in "scaling up" research to resolve whether similar patterns hold in natural ecosystems [4,5]. However, richness is not a robust measure of biodiversity in observational data taken from natural ecosystems [6], in large part because most species are rare [7] and likely to be absent from samples. Further, richness often tracks real-world biodiversity gradients poorly, because species composition and abundance can change dramatically with little to no change in observed species richness [8–10]. Therefore, other metrics of biodiversity may provide improved clarity about the real-world linkages between biodiversity and ecosystem function (BEF).

There are both historical and conceptual reasons that BEF research has focused on richness as a measure of biodiversity. Since at least the 1960s, there has been extensive research on how productivity affects species richness [1]. Motivated by intensifying biodiversity loss in the 1980s, declines in richness were (at least implicitly) the global change pattern that seminal BEF studies, with their focus on species loss (e.g., [11]), aimed to understand. This prompted a wave of experiments on how species richness affects ecosystem function [2]. Thus, richness was a natural choice, both because of ecology's long focus on how richness might *respond* to ecosystem functions like productivity, and because of a collective sense that species loss was the correct, or at least most convenient, way to frame anthropogenic changes in biodiversity. Furthermore, there may have been a sense that richness was an expedient measure of functional diversity and redundancy, which were considered the key mechanisms through which biodiversity maintains ecosystem function [12]. However, the choice of richness may not have been based on theoretical expectation that richness, rather than other abundance-weighted diversity measures, best described functionally important biodiversity gradients.

Using species richness as the key biodiversity measure poses methodological problems for BEF research. Species richness is not only sensitive to the extent and depth of sampling, but also to the distribution of relative abundances in the sampled assemblage. To illustrate this, consider the

difficulty of accurately measuring species richness in a community with one hyper-dominant species

and many very rare species, versus measuring richness in a community in which abundance is evenly

distributed. Richness, like other diversity measures, summarizes the distribution of relative

abundances in an assemblage, and when estimated from data, cannot be independent from that

distribution, even if such a measure were desirable [6,13]. However, different diversity measures vary

in the extent to which they emphasize rare vs. common species, with species richness heavily

emphasizing rare species. A unified family of diversity measures, known as "Hill numbers" or "Hill

diversities," summarizes a distribution of relative abundances as the abundance-weighted,

generalized mean rarity [14–16]. Hill numbers are governed by a scaling parameter, ℓ, that scales

species rarity when computing the mean, and higher values of ℓ afford more leverage, or emphasis, to

rare species, while lower values emphasize common species more [16].

The Hill diversity of an assemblage is not a single value, but rather a spectrum that varies

continuously across ℓ [6,14] (figure 1), raising the question of how ecosystem function relates to

biodiversity measures with different emphasis on common vs. rare species. While several recent

studies have compared whether richness ($\ell = 1$), exponentiated Shannon ($\ell = 0$), and inverse Simpson

($\ell = -1$) best predicts ecosystem function [17–20], there has been no examination of how Hill numbers

predict ecosystem function across a wide range of ℓ values. This is a striking knowledge gap because,

although nearly all studies of the relationship between biodiversity and ecosystem function have used

species richness as a measure of diversity [3], other diversity measures could both better describe real

world variation in biodiversity, and also have stronger links to ecosystem functioning. Thus,

biodiversity-function studies may be *underestimating* the importance of biodiversity for function by

not considering Hill diversities with different emphases on rare and common species via different

values of the scaling parameter ℓ.

Despite clear declines in richness at the global scale, local changes in biodiversity are likely better

captured by measuring total abundance and species' relative abundances, for at least three reasons.

First, as already discussed, observed richness is a poor predictor of true richness [21], and good

estimators of true richness based only on species frequencies in samples may never exist [22]. Thus,

even if underlying variation in species richness correlates strongly with, or even drives, ecosystem

function, estimating richness from samples could severely obscure the underlying pattern. Second,

although observed richness does increase with observed abundance, to the extent that abundance

per se drives function, Hill diversities that better reflect abundance (i.e., when $\ell \gg 1$) should be

stronger correlates of function than richness ($\ell = 1$) is. Third, if diversity effects on function are

mediated by positive species interactions [3,23–25], more probable and stronger between equally

abundant species [26,27], Hill diversities that better reflect the probability of interspecific encounter

(e.g. Hill-Simpson diversity, at $\ell = -1$ [28]) may outperform richness as a predictor of function.

Here, we conduct an in-depth examination of Hill numbers across a wide range of values of the scaling

parameter ℓ, and outline the ecological insight that can be gained by knowing which value of ℓ best

predicts ecosystem function. We focus on ecosystem functions that can be expressed as the product

of mean per-capita function and total abundance, which works well for many functions [29,30]. This

provides the opportunity to consider how biodiversity relates to both mean per-capita and total

community function, which are both key targets of BEF research [31–33]. In this endeavor, we use only

simple correlational analyses, omitting other important variables underlying function [34–36].

In this study, we use observational community datasets on three ecosystem functions to ask:

1) Which values of the Hill diversity scaling factor ℓ produce the strongest biodiversity-

109   ecosystem function correlations?

110   2) How do biodiversity-ecosystem function correlations change in sign and strength over a wide

111   range of Hill diversities?

112   3) What is the role of absolute abundance in shaping BEF correlations over the Hill diversity

113   spectrum?

It is not our aim to exhaustively review data in search of a single, best ℓ value for BEF research,

because developing insight does not require that the same ℓ is predictive for all functions. Instead, we

aim to provide a framework that allows researchers interested in biodiversity-function questions to

think about diversity as a spectrum, almost certainly leading to the conclusion that Hill diversities

besides richness can better predict ecosystem function.

Materials and Methods

To find how biodiversity-ecosystem function correlations change with different diversity scaling, we

used previously published community datasets that recorded both the abundance and function of

species at multiple sites. We chose datasets of disparate ecosystem functions and spatial scales:

pollination by wild bee visitors to a landscape-scale array of three plant species (update ref when DOI

is available: Genung et al. 2022, *in press*) [37], reef fish biomass from dive surveys replicated within 32

globally distributed geographic regions [38,39], and above-ground tree biomass in census plots

replicated within four tropical forests [40,41] (Table 1). In each system, total function of a community

can be estimated as the summed contribution across species (or individuals) present in the

community. Pollination was measured as the typical number of pollen grains deposited during a

single visit of a bee taxon to the focal plant species, multiplied, at each site, by bee species's

abundance to calculate total pollen deposition of one species to the focal plant there. Reef fish

biomass was measured by visually estimating individual fish body lengths during dive surveys, which

were then used to calculate biomass using species-specific allometric equations. Tropical tree

biomass was measured by converting observed individual diameter at breast height into biomass

estimates using taxon-specific allometric equations that included information about wood density

[42,43]. In total, we used 39 community datasets, each consisting of one function measured across a

collection of assemblages.

Which value of ℓ produces the strongest biodiversity- ecosystem function correlations?

We compute Hill diversity as a function of species relative abundances, *p_1, p_2, …, p_S,* and a scaling

factor, ℓ, using the formula

144
$$D = \left(\sum_{i=1}^{S} p_i \left(\frac{1}{p_i} \right)^{\ell} \right)^{1/\ell}$$

or its limit as ℓ approaches 0 [16,44] . We used observed species abundances to calculate species

diversities at each site as the Hill diversity along a wide range of ℓ values (from -10 to +10 at intervals

of 0.05) (figure S1). We calculated total function as the sum of species' functions at each site. We

computed the correlation between the natural logarithm of each diversity and the natural logarithm

of total function, across all sites in the community dataset (hereafter, the "BEF correlation"). To

identify the ℓ value that produced the strongest BEF correlation in each community dataset, we

plotted the correlation against the scaling factor ℓ. We identified the single ℓ value with the largest

absolute correlation (i.e., largest R-squared for the relationship between log diversity and log

function).

How do biodiversity-ecosystem function correlations change in sign and strength over a wide range of

Hill diversities?

To determine not only which ℓ value produced the strongest BEF correlation across community

datasets, but also to see how adjusting the Hill diversity scaling parameter affects BEF relationships

more comprehensively, we plotted the BEF correlation against the Hill diversity scaling factor ℓ for

each community dataset. We examined curves to identify patterns in the sign and strength of the BEF

correlation along the spectrum of emphasis on common and rare species.

What is the role of absolute abundance in shaping BEF correlations?

To begin to separate effects of total and relative abundance on BEF correlations across the Hill

diversity spectrum, we looked separately at the relationships between diversity and two

complementary components of total function, namely total abundance and mean per-capita function.

We used the same graphical approach we used to assess the sign and strength of the BEF correlation

across the Hill diversity spectrum. For each community dataset, we found the correlation between the

natural logarithm of Hill diversity at each site and either the natural logarithm of total abundance at

each site), or the natural logarithm of mean per-capita function at each site, and plotted these

correlations against the Hill diversity scaling parameter ℓ. Although on the logarithmic scale,

abundance and mean per-capita function combine additively to create total function, the BEF

correlation does not additively decompose into abundance by biodiversity and per-capita function by

biodiversity correlations, as there is also covariance between abundance and per-capita function.

Nevertheless, by partitioning total function into additive components and examining how each of

these relates to biodiversity gradients across the Hill spectrum, we can better characterize the role of

total abundance in generating patterns in the BEF correlation itself.

Results

Which value of ℓ produces the strongest biodiversity- ecosystem function correlations?

For most datasets, the strongest biodiversity-ecosystem correlations were located at or just above

richness ($\ell = 1$), with a mode at $\ell = 1.5$ (figure 2). A substantial minority (11 of 39 datasets) had

strongest BEF correlations at values of $\ell > 5$, including a peak at $\ell = 10$, the largest value of ℓ we

considered. There were a few outliers: Two tree carbon storage datasets had their strongest BEF

correlations near inverse Simpson ($\ell = -1$) and Shannon ($\ell = 0$) diversities, and a single fish dataset had

a strongest BEF correlation at $\ell = -10$, the smallest value of ℓ we considered (figure 2).

How do biodiversity-ecosystem function correlations change in sign and strength over a wide range of

Hill diversities?

Across all ecosystem functions, we found common patterns in the relationship between the BEF

correlation and the Hill diversity scaling parameter, ℓ. When ℓ is < 1, the diversity-function correlation

is typically weak and may be positive or negative (figure 3). Near $\ell = 1$, the diversity-function

correlation rapidly increases, although a substantial minority of community datasets first show a

sharp negative turn in the relationship near Hill-Simpson and Hill-Shannon diversities (figure 3 b-d).

Across all the datasets we considered, the mean correlation between ln(diversity) and ln(total

function) was not significantly different from zero for either Hill-Simpson or Hill-Shannon diversity

(p>0.28 for two-sided Student's t-test, with no correction for multiple tests). By richness ($\ell = 1$), almost

all datasets showed positive diversity-function correlations, with the mean $R^2 = 0.381$. For most

datasets, the strongest correlations are located near richness, with a mode near $\ell = 1.5$, where the

mean R^2 was 0.445, after which the diversity-function correlation slowly declines as ℓ values continue

to increase (figure 3). A substantial minority of datasets showed continually stronger relationships as ℓ

increased (some profiles in figure 3 b, c), leading to highest R^2 values at or near the maximum ℓ we

considered ($\ell = 10$).

This study was not designed to contrast trends between ecosystem functions, but it is important to

note that the relationship between the BEF correlation, and the emphasis the diversity metric puts on

rare vs. common species (i.e., the value of ℓ), did not appear uniform. For the three bee community

datasets, total pollen deposition and bee diversity were positively correlated at every value of ℓ.

Correlation strength peaked at richness ($\ell = 1$) or just beyond ($\ell = 2$), but remained relatively strong

across all higher values of ℓ (figure 3a). For the 32 reef fish community datasets, total fish biomass and

fish diversity tended to be weakly and often negatively correlated at low ℓ values. Correlation strength

tended to peak either slightly above richness at $\ell = 1.5$, or grow with ℓ for an observed peak near

the maximum value considered ($\ell = 10$). (figure 2, figure 3 b, c). When considering either very high or

very low ℓ values, note that at either end of the Hill number spectrum, diversities rapidly converged

towards their maximum or minimum asymptote. Thus, large changes in the BEF correlation rarely

occurred outside a fairly narrow range between $\ell = -2$ and 2. Finally, the four tropical tree community

datasets showed generally weak correlations. In two tree datasets, BEF correlation strength peaked at

intermediate ℓ values where the BEF correlation was strongly negative (figure 3 d). In another (Barro

Colorado Island) diversity-function correlation was negative even at high ℓ values (figure 3 d, orange

line), but modestly positive for negative ℓ values.

What is the role of absolute abundance in shaping BEF correlations?

As expected, the relationship between diversity and abundance was mostly similar to the relationship

between diversity and function, as total abundance underlies function in our datasets. This can be

seen in the similar shape of the curves showing the correlation between ln(diversity) and either

ln(function) (figure 3) or ln(abundance) (figure 4 a-d), as the sign and strength of correlation typically

moved in similar ways across the ℓ spectrum. In almost all cases, the correlation between ln(diversity)

and ln(abundance) was very strong (and in many cases approached unity), for large, positive values of

the Hill diversity scaling parameter ℓ. As previously remarked, this result is a mathematical

inevitability when datasets contain very rare species/singletons. Additionally, across datasets, we

found that the rise towards the high correlation observed for large ℓ values typically occurred in the

range of ℓ values typically considered by ecologists (-1 to 1), likely reflecting biological and sampling

linkages between abundance and diversity; the correlation frequently saturated once ℓ was greater

than two. While for some community datasets, diversity was largely independent of abundance for

negative ℓ values, we also saw community datasets in which ln(abundance) and ln(diversity) had

modest to strong negative correlation across negative ℓ values. Because Hill diversities typically

change little with ℓ below around -2 [6], this result implies that in these systems, total abundance and

the degree of dominance are positively linked [45].

While the curves in figures 3 a-d and 4 a-d show strong resemblance, for some community datasets

the BEF and diversity-abundance relationships diverge, implying those BEF relationships result from

processes other than abundance. For example, correlations between abundance and diversity were

always strongly positive for the reef fish data for large, positive ℓ values, but in some reef fish

community datasets, correlations between diversity and function were only weakly positive at higher

ℓ values (figure 4 b, c). Such divergences between the diversity-function and diversity-abundance

curves could be due to strong and/or countervailing relationships between mean per-capita function

and diversity, which also showed some overall patterns across community datasets (figure 4 e-h). In

general, Hill diversities with negative ℓ values were positively related to per-capita function,

suggestive of a positive relationship between evenness and mean per-capita function. This pattern

was not ubiquitous, however, with notable exceptions in both tree and bee community datasets

(figure 4 e, h). We found that the correlation between diversity and mean per-capita function often

exhibited a positive peak at intermediate ℓ values, a pattern particularly pronounced in the reef fish

community datasets (figure 4 f, g). Finally, there was a tendency towards a negative correlation

between ln(diversity) and ln(mean per-capita function) at for larger, positive values of ℓ (when

diversity becomes largely synonymous with abundance), though the strength of this relationship was

variable.

Discussion

We explored which Hill diversities best predicted ecosystem function, and were surprised to find that

diversities near richness ($1 < \ell < 2$) often performed the best (figure 2). This result was unexpected

based on three assumptions we described in the introduction: first, observed richness is not a robust

biodiversity measure; second, if abundance drives function, Hill diversities with high ℓ should be more

predictive; third, if positive species interactions (e.g., complementarity [46]) drive function, ℓ values

that emphasize the probability of interspecific encounter ($-1 \le \ell < 1$) should be more predictive. The

disconnect between observed and true richness is not, practically speaking, a resolvable problem and

thus we cannot evaluate how much this first issue is affecting our results [22]. In the following

paragraphs, we first interpret the unexplored meaning of Hill diversity when $\ell > 1$ and then explore the

latter two points, namely: how do Hill diversities near richness outperform Hill diversities with higher ℓ

that better reflect abundance, and why we might have found such low predictive power for Hill

diversities that should capture the effects of (potentially positive) species interactions. To better

ground our discussion, we use an admittedly imprecise simplification and refer to ℓ values as falling

272 within the "evenness range" (-1 ≤ ℓ < 1), the "rare emphasis range" (1 < ℓ < 2, justification follows), or

273 the "abundance range" (ℓ > 2).

275 Before we can fully understand our results, we must interpret Hill diversities with ℓ > 1, which have

276 seen little attention. Traditionally used Hill diversities with ℓ < 1 reach their maximum possible value,

277 species richness, when species all have the same abundance. However, Hill diversities with ℓ > 1,

278 which strongly emphasize rare species, increase with heterogeneity (rather than homogeneity) in

279 species abundance. By emphasizing rare species even *more* than richness does, these diversities may

280 reflect two essential features of community size: the total number of species and the total number of

281 individuals. As these features are not practically separable and both apparently drive function, it may

282 be advantageous to measure them jointly. Arguably, observed richness does this [31], but our results

283 show that ℓ values in the rare emphasis range, which are relatively more sensitive to abundance than

284 richness is, yield measures of diversity that better predict function. Beyond the rare emphasis range,

285 Hill diversity quickly converges on abundance (at least in most observational data). This results in a

286 slight decline in Hill diversity's ability to predict ecosystem function for larger ℓ values (figures 2-3).

288 Hill diversities in the evenness range (-1 ≤ ℓ < 1) should capture the effects of species interactions by

289 emphasizing the probability of interspecific encounter, but these diversities were poor predictors of

290 function. This was unexpected because a consistent BEF finding, including in real-world systems, has

291 been that function increases with evenness [47–50], which increases Hill diversity for ℓ < 1 [51–53].

292 Additionally, in the evenness range, sample Hill diversities have relatively good statistical properties

293 as estimators of true diversity, and asymptotic estimators [54] can further improve the situation,

294 largely avoiding the robustness issues we highlight with species richness. Instead, the observed weak

295 predictive power of Hill diversities with ℓ values in the evenness range is because functions analyzed

296 here are the product of two components, abundance and per-capita function, which each showed

297 different responses to increasing ℓ. Abundance-diversity relationships often mirrored function-

298 diversity correlations (compare figure 3 with the top row of figure 4). In other words, across the ℓ

299 spectrum, Hill diversities had nearly the same relationship with abundance and with function,

300 underlining the necessity of accounting for the role of total abundance in BEF research [55]. However,

301 Hill diversities in the evenness range deviated from this pattern, instead exhibiting often strong,

302 countervailing relationships with abundance and per-capita function (figure 4).

The relationship between diversity and per-capita function differed from the diversity-abundance

relationship, with correlation coefficients for per-capita function generally decreasing with increases

in ℓ, but often showing a positive peak in the evenness range (figure 4). Positive species interactions,

including those that increase per-capita function, are expected to predict total function [46,56]. Our

results partly support these expectations, as Hill diversities in the evenness range, which should track

the probability of interspecific encounter, were positively associated with per-capita function, even as

they tended to be negatively associated with abundance. As we increased ℓ, the correlation between

Hill diversity and per-capita function disappeared near richness ($\ell = 1$), also pushing against

expectations that richness best captures function-relevant biodiversity gradients. In the rare emphasis

and abundance ranges, we typically found a negative correlation between Hill diversity and per-capita

function. This likely reflects spatial constraints and/or fundamental tradeoffs between having many,

smaller-bodied individuals versus fewer larger ones [57,58]. This scenario is particularly easy to

imagine for trees crowding in fixed-area plots, which physically and energetically prohibit arbitrarily

large numbers of the largest trees. Similar energetic and spatial constraints limit the number of very

large fish that might be seen in a single dive. Thus, we suspect that one reason we see a decline in the

correlation between mean per-capita function and diversity with increasing ℓ in the fish and tree

datasets is decreases in per-capita function due to crowding.

Although Hill diversities in the rare emphasis range were most often the best predictor of total

function, Hill diversities with ℓ values in the abundance range also predicted function well, and should

not be discounted. Hill diversities in the abundance range were the *best* predictor of function in a

substantial minority of datasets (Fig. 2), and for nearly all datasets were *strong* predictors of function

(Fig. 3, far right of x-axes). This was expected because of a general link between higher abundance and

higher function [55,59–63]. Even as Hill diversities in the "abundance" range were strong predictors of

function, we also note that abundance can relate to Hill diversity across the full spectrum of ℓ values.

For example, if high-abundance sites tend to be dominated by many individuals of one or a few

species [45], Hill diversities that emphasize the most common species will decrease with abundance.

Thus, we should not expect that strong effects of abundance on function are captured exclusively at

high values of ℓ.

The predictive power of power of Hill diversity changed nonlinearly with increases in ℓ, as multiple

facets of community structure (e.g., richness, abundance, evenness) affect function simultaneously. If

we had found a monotonic strengthening of BEF relationships with increasing ℓ, we would argue that

Hill "diversities" with large positive scaling parameters were simply abundance metrics masquerading

as measures of diversity. Instead, we found, across a variety of regions, taxa, and ecosystem functions,

intermediate, positive ℓ values in the "rare emphasis" range tended to produce the strongest BEF

relationships (Fig. 3). All Hill diversities with positive ℓ values (including richness) tend to increase with

both abundance and richness, which we argue can be a useful property, especially for BEF research.

Because the goal of summarizing species' abundances with diversity metrics is to distill complex,

multivariate information [15], this claim is not radical. In fact, Hill diversities that emphasize rare

species more than richness does can reflect intuitive notions of diversity, which include both high

density and high compositional variation [64]. Our study points to the need for further theoretical

work to explicate the meaning of these seldom-used Hill diversities in the rare emphasis range, and

their linkages to ecosystem function.

By considering Hill diversities over a wide range of ℓ, we place ourselves at odds with the convention

that Hill diversities should be considered only when $\ell \leq 1$ [6,14,15,65,66]. The most compelling

argument for that restricted range of scaling parameters is presented by Patil and Taillie, who argued

that diversity should not decrease when abundance is shifted from more to less abundant species,

including to species with zero abundance, a variation on Dalton's "principle of transfers" [15,67]. This

diversity property does not hold for Hill diversity when $\ell > 1$, which has species richness as its

minimum, occurring in the perfectly even community, and increases (given richness and abundance)

as some species get progressively rarer. A more pragmatic argument comes from Chao et al., who

noted that estimating the relative abundance of rare species is an increasing problem for diversity

measures as ℓ increases; they therefore suggest using only more estimable Hill diversities with $\ell \leq 1$ [6].

However, theoretical work suggests that even richness ($\ell = 1$) is poorly estimated [22], and by this logic

should not be used either. Finally, and most generally, diversity measures have traditionally been

considered separate from abundance/density measures (but see [68,69]), whereas with increasing ℓ

values, observed diversity and observed abundance tend to be more strongly correlated (and in fact

approach a correlation of one in our datasets). Despite these arguments, our results show that Hill

diversities with $\ell > 1$ are meaningful ecological diversity measures, at least in the sense that they

convey more information about function than do more widely-used Hill numbers. Choosing to exclude

these Hill diversities might be desirable for some conceptualizations of diversity, but we are opening

the narrower question of which Hill diversities—with their variable emphasis on richness, abundance,

evenness, and dominance—best predict ecosystem function. In this pursuit, allowing diversity metrics

to highlight absolute abundance is valuable.

As global changes lead to shifting species abundances, ecologists must continue to describe and

predict how these shifts impact ecosystems and the way they function. Yet, understanding the

separate and combined roles of total and relative abundance in mediating ecosystem function

remains a difficult challenge, in large part because total abundance is inextricably linked to diversity

measures. It is mathematically linked for large, positive ℓ values. It is practically constrained by

sampling effects for ℓ closer to 1 (i.e., near species richness). As ℓ becomes negative, Hill diversities

may lose their dependence on total abundance [21]. However, in the majority of community datasets,

we saw at least weak negative correlations between negative-ℓ Hill diversities and observed

abundances, likely due to increasing dominance in more abundant systems [45]. Overall, this suggests

that in observational contexts, simple partitioning of abundance and diversity effects may not be

tractable, at least not in a satisfying manner [31,36]. Since no single-best diversity measure is likely to

emerge for all BEF studies, we encourage researchers to be open-minded towards Hill diversities

across a wide spectrum of ℓ values and their potential links to mechanisms underlying BEF

relationships.

Competing Interests

We have no competing interests.

Acknowledgements

We thank Rachael Winfree for key discussions that sparked this work, Jonathan Dushoff for providing

insight on Hill numbers, and participants in the Hakai Synthesizing Biodiversity Seminar Series for

useful feedback on preliminary results. We gratefully acknowledge all data collectors and data

holders. Data for three tropical forests (Pasoh, Indonesia; Yasuní, Ecuador; and Volcán Barva, Costa

Rica) were originally made available to us by the Tropical Ecology Assessment and Monitoring (TEAM)

Network, a collaboration between Conservation International, the Missouri Botanical Garden, the

Smithsonian Institution, and the Wildlife Conservation Society, and partially funded by these

institutions, the Gordon and Betty Moore Foundation, and other donors.

References

1. Loreau M *et al.* 2001 Biodiversity and Ecosystem Functioning: Current Knowledge and Future

Challenges. *Science (80-.).* **294**, 804–808. (doi:10.1126/science.1064088)

2. Hooper DU *et al.* 2005 Effects of biodiversity on ecosystem functioning: A consensus of current

knowledge. *Ecol. Monogr.* **75**, 3–35. (doi:10.1890/04-0922)

3. Cardinale BJ *et al.* 2012 Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67.

(doi:10.1038/nature11148)

4. Gonzalez A *et al.* 2020 Scaling-up biodiversity-ecosystem functioning research. *Ecol. Lett.* **23**,

757–776. (doi:10.1111/ele.13456)

5. Qiu J, Cardinale BJ. 2020 Scaling up biodiversity–ecosystem function relationships across

space and over time. *Ecology* **101**, 0–2. (doi:10.1002/ecy.3166)

6. Chao A, Gotelli NJ, Hsieh TC, Sander EL, Ma KH, Colwell RK, Ellison AM. 2014 Rarefaction and

extrapolation with Hill numbers: a framework for sampling and estimation in species diversity

studies. *Ecol. Monogr.* **84**, 45–67. (doi:10.1890/13-0133.1)

7. McGill BJ *et al.* 2007 Species abundance distributions: Moving beyond single prediction

theories to integration within an ecological framework. *Ecol. Lett.* **10**, 995–1015.

(doi:10.1111/j.1461-0248.2007.01094.x)

8. Hillebrand H *et al.* 2018 Biodiversity change is uncoupled from species richness trends:

consequences for conservation and monitoring. *J. Appl. Ecol.* **55**, 169–184. (doi:10.1111/1365-

2664.12959)

9. Brose U, Hillebrand H. 2016 Biodiversity and ecosystem functioning in dynamic landscapes.

Philos. Trans. R. Soc. B Biol. Sci. **371**. (doi:10.1098/rstb.2015.0267)

10. Spaak JW *et al.* 2017 Shifts of community composition and population density substantially

affect ecosystem function despite invariant richness. *Ecol. Lett.* **20**, 1315–1324.

(doi:10.1111/ele.12828)

11. Naeem S, Thompson LJ, Lawler SP, Lawton JH, Woodfin RM. 1994 Declining biodiversity can

alter the performance of ecosystems. *Nature* **368**, 734–737. (doi:10.1038/368734a0)

12. Walker BH. 1992 Biodiversity and Ecological Redundancy. *Conserv. Biol.* **6**, 18–23.

(doi:10.1046/j.1523-1739.1992.610018.x)

13.   Chao A, Jost L. 2012 Coverage-based rarefaction and extrapolation: Standardizing samples by

completeness rather than size. *Ecology* **93**, 2533–2547. (doi:10.1890/11-1952.1)

14.   Hill MO. 1973 Diversity and evenness: a unifying notation and its consequences. *Ecology* **54**,

427–432. (doi:10.2307/1934352)

15.   Patil GP, Taillie C. 1982 Diversity as a concept and its measurement. *J. Am. Stat. Assoc.* **77**, 548–

561. (doi:10.1080/01621459.1982.10477845)

16.   Roswell M, Dushoff J, Winfree R. 2021 A conceptual guide to measuring species diversity. *Oikos*

130, 321–338. (doi:10.1111/oik.07202)

17.   Yasuhara M, Doi H, Wei CL, Danovaro R, Myhre SE. 2016 Biodiversity-ecosystem functioning

relationships in long-term time series and palaeoecological records: Deep sea as a test bed.

Philos. Trans. R. Soc. B Biol. Sci. **371**. (doi:10.1098/rstb.2015.0282)

18.   Schweiger AK, Cavender-Bares J, Townsend PA, Hobbie SE, Madritch MD, Wang R, Tilman D,

Gamon JA. 2018 Plant spectral diversity integrates functional and phylogenetic components of

biodiversity and predicts ecosystem function. *Nat. Ecol. Evol.* **2**, 976–982. (doi:10.1038/s41559-

018-0551-1)

19.   Yang JW, Wu W, Chung CC, Chiang KP, Gong GC, Hsieh CH. 2018 Predator and prey biodiversity

relationship and its consequences on marine ecosystem functioning - Interplay between

nanoflagellates and bacterioplankton. *ISME J.* **12**, 1532–1542. (doi:10.1038/s41396-018-0111-3)

20.   Schmidt ML, Biddanda BA, Weinke AD, Chiang E, Januska F, Props R, Denef VJ. 2020

Microhabitats are associated with diversity-productivity relationships in freshwater bacterial

communities. *FEMS Microbiol. Ecol.* **96**, 1–12. (doi:10.1093/FEMSEC/FIAA029)

21.   Chao A, Jost L. 2015 Estimating diversity and entropy profiles via discovery rates of new

species. *Methods Ecol. Evol.* **6**, 873–882. (doi:10.1111/2041-210X.12349)

22.   Haegeman B, Hamelin J, Moriarty J, Neal P, Dushoff J, Weitz JS. 2013 Robust estimation of

microbial diversity in theory and in practice. *ISME J.* **7**, 1092–1101. (doi:10.1038/ismej.2013.10)

23.   Fargione J, Tilman D, Dybzinski R, Lambers JHR, Clark C, Harpole WS, Knops JMH, Reich PB,

Loreau M. 2007 From selection to complementarity: Shifts in the causes of biodiversity-

productivity relationships in a long-term biodiversity experiment. *Proc. R. Soc. B Biol. Sci.* **274**,

871–876. (doi:10.1098/rspb.2006.0351)

24.   Reich PB, Tilman D, Isbell F, Mueller K, Hobbie SE, Flynn DFB, Eisenhauer N. 2012 Impacts of

458    biodiversity loss escalate through time as redundancy fades. *Science (80-.).* **336**, 589–592.

459    (doi:10.1126/science.1217909)

460 25. Tilman D, Isbell F, Cowles JM. 2014 Biodiversity and ecosystem functioning. *Annu. Rev. Ecol.*

461    *Evol. Syst.* **45**, 471–493. (doi:10.1146/annurev-ecolsys-120213-091917)

462 26. Jiang L, Wan S, Li L. 2009 Species diversity and productivity: Why do results of diversity-

463    manipulation experiments differ from natural patterns. *J. Ecol.* **97**, 603–608.

464    (doi:10.1111/j.1365-2745.2009.01503.x)

465 27. Crowder DW, Northfield TD, Strand MR, Snyder WE. 2010 Organic agriculture promotes

466    evenness and natural pest control. *Nature* **466**, 109–112. (doi:10.1038/nature09183)

467 28. Chase JM, Knight TM. 2013 Scale-dependent effect sizes of ecological drivers on biodiversity:

468    Why standardised sampling is not enough. *Ecol. Lett.* **16**, 17–26. (doi:10.1111/ele.12112)

469 29. Balvanera P, Kremen C, Martínez-ramos M. 2005 Applying Community Structure Analysis to

470    Ecosystem Function : Examples from Pollination and Carbon Storage. *Ecol. Appl.* **15**, 360–375.

471 30. Kremen C. 2005 Managing ecosystem services: What do we need to know about their ecology?

472    *Ecol. Lett.* **8**, 468–479. (doi:10.1111/j.1461-0248.2005.00751.x)

473 31. Harrison T, Winfree R, Genung MA. 2022 Price Equations for Understanding the Response of

474    Ecosystem Function to Community Change. *Am. Nat.* **200**, 181–192. (doi:10.1086/720284)

475 32. Carvalho F, Pascoal C, Cássio F, Teixeira A, Sousa R. 2022 Combined per-capita and abundance

476    effects of an invasive species on native invertebrate diversity and a key ecosystem process.

477    *Freshw. Biol.* **67**, 828–841. (doi:10.1111/fwb.13884)

478 33. Ruesink JL, Srivastava DS. 2001 Numerical and per capita responses to species loss:

479    Mechanisms maintaining ecosystem function in a community of stream insect detritivores.

480    *Oikos* **93**, 221–234. (doi:10.1034/j.1600-0706.2001.930206.x)

481 34. Duffy JE, Lefcheck JS, Stuart-Smith RD, Navarrete SA, Edgar GJ. 2016 Biodiversity enhances

482    reef fish biomass and resistance to climate change. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 6230–6235.

483    (doi:10.1073/pnas.1524465113)

484 35. Mori AS, Osono T, Cornelissen JHC, Craine J, Uchida M. 2017 Biodiversity–ecosystem function

485    relationships change through primary succession. *Oikos* **126**, 1637–1649.

486    (doi:10.1111/oik.04345)

487 36. Harte J, Brush M, Newman EA, Umemura K. 2022 An equation of state unifies diversity,

488    productivity, abundance and biomass. *Commun. Biol.* **5**, 874. (doi:10.1038/s42003-022-03817-8)

489   37.   Genung M, Reilly J, Williams N, Buderi A, Gardner J, Winfree R. 2022 Rare and declining bee

490   species are key to consistent pollination of wildflowers and crops across large spatial scales;

491   Dryad, Dataset. (doi:https://doi.org/10.5061/dryad.qnk98sfkc)

492   38.   Lefcheck JS *et al.* 2021 Species richness and identity both determine the biomass of global reef

493   fish communities. *Nat. Commun.* **12**, 1–9. (doi:10.1038/s41467-021-27212-9)

494   39.   Lefcheck JS. 2021 Species richness and identity both determine the biomass of global reef fish

495   communities. *Dataset*. (doi:https://doi.org/10.25573/serc.16847029.v1)

496   40.   Condit R *et al.* 2000 Spatial patterns in the distribution of tropical tree species. *Science* **288**,

497   1414–1418.

498   41.   Cavanaugh KC *et al.* 2014 Carbon storage in tropical forests correlates with taxonomic diversity

499   and functional dominance on a global scale. *Glob. Ecol. Biogeogr.* **23**, 563–573.

500   (doi:10.1111/geb.12143)

501   42.   Zanne AE *et al.* 2009 Data From: Towards a worldwide wood economics spectrum. *Data Dryad*.

502   (doi:https://doi.org/10.5061/dryad.234)

503   43.   Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE. 2009 Towards a worldwide

504   wood economics spectrum. *Ecol. Lett.* **12**, 351–366. (doi:10.1111/j.1461-0248.2009.01285.x)

505   44.   Roswell M, Dushoff J. 2020 MeanRarity: Hill Diversity Estimation and Visualisation.

506   45.   MacDonald ZG, Nielsen SE, Acorn JH. 2017 Negative relationships between species richness

507   and evenness render common diversity indices inadequate for assessing long-term trends in

508   butterfly diversity. *Biodivers. Conserv.* **26**, 617–629. (doi:10.1007/s10531-016-1261-0)

509   46.   Loreau M, Hector A. 2001 Partitioning selection and complementarity in biodiversity

510   experiments. *Nature* **412**, 72–76. (doi:10.1038/35083573)

511   47.   Wilsey BJ, Potvin C. 2000 Biodiversity and ecosystem functioning: Importance of species

512   evenness in an old field. *Ecology* **81**, 887–892. (doi:10.1890/0012-

513   9658(2000)081[0887:BAEFIO]2.0.CO;2)

514   48.   Polley HW, Wilsey BJ, Derner JD. 2003 Do species evenness and plant density influence the

515   magnitude of selection and complementarity effects in annual plant species mixtures? *Ecol.*

516   *Lett.* **6**, 248–256. (doi:10.1046/j.1461-0248.2003.00422.x)

517   49.   Wittebolle L, Marzorati M, Clement L, Balloi A, Daffonchio D, Heylen K, De Vos P, Verstraete W,

518   Boon N. 2009 Initial community evenness favours functionality under selective stress. *Nature*

519   **458**, 623–626. (doi:10.1038/nature07840)

50.   Genung MA, Fox J, Winfree R. 2020 Species loss drives ecosystem function in experiments, but

in nature the importance of species loss depends on dominance. *Glob. Ecol. Biogeogr.* **29**, 1–11.

(doi:10.1111/geb.13137)

51.   Stirling G, Wilsey B. 2001 Empirical Relationships between Species Richness, Evenness, and

Proportional Diversity. *Am. Nat.* **158**, 286–299. (doi:10.1086/321317)

52.   Jost L. 2010 Independence of alpha and beta diversities. *Ecology* **91**, 1969–1974.

(doi:10.1890/09-0368.1)

53.   Chao A, Ricotta C, Aldo P. 2019 Quantifying evenness and linking it to diversity, beta diversity,

and similarity. *Ecology* **100**, 1–15. (doi:10.1002/ecy.2852)

54.   Chao A, Jost L. 2015 Estimating diversity and entropy profiles via discovery rates of new

species. *Methods Ecol. Evol.* **6**, 873–882. (doi:10.1111/2041-21)

55.   Winfree R, Fox JW, Williams NM, Reilly JR, Cariveau DP. 2015 Abundance of common species,

not species richness, drives delivery of a real-world ecosystem service. *Ecol. Lett.* **18**, 626–635.

(doi:10.1111/ele.12424)

56.   Tilman D, Lehman CL, Thomson KT. 1997 Plant diversity and ecosystem productivity:

Theoretical considerations. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 1857–1861.

(doi:10.1073/pnas.94.5.1857)

57.   Yoda K, Kira T, Ogawa H, Hozum K. 1963 Self-thinning in overcrowded pure stands under

cultivated and natural conditions. *J. Inst. Polytech. Osaka City Univerisity. Ser. D, Biol.* , 107–129.

58.   Hughes RN, Griffiths CL. 1988 Self-thinning in barnacles and mussels: the geometry of packing.

Am. Nat. **132**, 484–491. (doi:10.1086/284866)

59.   Dangles O, Malmqvist B. 2004 Species richness-decomposition relationships depend on species

dominance. *Ecol. Lett.* **7**, 395–402. (doi:10.1111/j.1461-0248.2004.00591.x)

60.   Grime JP. 1998 Benefits of plant diversity to ecosystems: immediate, filter and founder effects.

J. Ecol. **86**, 902–910.

61.   Smith MD, Knapp AK. 2003 Dominant species maintain ecosystem function with non-random

species loss. *Ecol. Lett.* **6**, 509–517. (doi:10.1046/j.1461-0248.2003.00454.x)

62.   Vázquez DP, Morris WF, Jordano P. 2005 Interaction frequency as a surrogate for the total effect

of animal mutualists on plants. *Ecol. Lett.* **8**, 1088–1094. (doi:10.1111/j.1461-0248.2005.00810.x)

63.   Gaston KJ. 2011 Common Ecology. *Bioscience* **61**, 354–362. (doi:10.1525/bio.2011.61.5.4)

64.   McGill BJ, Dornelas M, Gotelli NJ, Magurran AE. 2015 Fifteen forms of biodiversity trend in the

anthropocene. *Trends Ecol. Evol.* **30**, 104. (doi:10.1016/j.tree.2014.11.006)

65. Jost L. 2006 Entropy and diversity. *Oikos* **113**, 363–375. (doi:10.1111/j.2006.0030-1299.14714.x)

66. Leinster T, Cobbold CA. 2012 Measuring diversity: the importance of species similarity. *Ecology*
93, 477–489. (doi:10.1890/10-2402.1)

67. Dalton H. 1920 The measurement of the inequality of incomes. *Econ. J.* **30**, 348–361.

68. Ricotta C. 2003 Parametric scaling from species relative abundances to absolute abundances in
the computation of biological diversity: A first proposal using Shannon's entropy. *Acta*
Biotheor. **51**, 181–188. (doi:10.1023/A:1025142106292)

69. Van Strien AJ, Soldaat LL, Gregory RD. 2012 Desirable mathematical properties of indicators for
biodiversity change. *Ecol. Indic.* **14**, 202–208. (doi:10.1016/j.ecolind.2011.07.007)

Table 1. To learn how biodiversity-ecosystem function correlations are affected by different Hill diversity scaling factors, we gathered published, observational community datasets on three ecosystem functions. These were subdivided into a total of 39 community datasets, each including observations of species' identities, abundances, and functions across replicated sites.

Ecosystem function	Citation	Datasets	Replication	Max extent (km)
Rate of wild bee pollen deposition	Genung et al. 2022	Landscape array of 3 plant species	Each plant species present at 25 sites	35
Fish biomass	Lefcheck et al. 2021	32 globally distributed ecoregions (16 temperate, 16 tropical)	11-186 sites (median 59)	17 - 4,677
Above-ground carbon storage	Condit et al. 2000, Cavanaugh et al. 2014	Tree species ID and estimated biomass at four globally distributed tropical forests	50 1-Ha subplots from the 50-Ha BCI census; sets of six 1-Ha plots in three tropical regions.	1 (Condit et al.); 32-681 (Cavanaugh et al.)

592
593
Fig 1. Two hypothetical communities with (a) different species abundance distributions have (b) different diversity profiles. At large negative ell values, each diversity profile converges on the inverse proportional abundance of the one most abundant species in the assemblage (inverse dominance). As ell values are more positive, each diversity profile converges on the inverse proportional abundance of the one least abundant species in the assemblage (equal to total abundance when the least abundant species is a singleton). Because singletons are ubiquitous in observational data, sample Hill diversities converge on observed abundance with increasingly large, positive values of *ell*. In the example, the red community is more even and more abundant, so its diversity is higher compared to the blue community at both ends of the diversity spectrum. However, the blue community has more species, and therefore is more diverse around richness (ell = 1, solid vertical). Other commonly used diversities are inverse Simpson (dotted) and exponentiated Shannon (dashed).

Figure 2. Across 39 observed biodiversity-ecosystem function (BEF) correlations calculated using a wide range of diversity scaling factors, the BEF relationships with the highest r² were typically found using diversities near richness (vertical solid line; modal ℓ = 1.5). The highest BEF r² value for a community dataset was rarely found using diversities that emphasize the relative abundance of common species, including exponentiated Shannon (dashed line) and inverse Simpson (dotted line). Correlations were calculated between log diversity and log ecosystem function at a site (total above-ground carbon in tropical forest plots, rate of pollen grain deposition by wild bees, or total biomass of reef fish encountered in fixed-effort dive surveys in temperate and tropical regions).

647
648
Figure 3. Biodiversity-ecosystem function (BEF) correlations across observed communities in a
study system vary in magnitude and direction, depending on which scaling factor (ell) is used for
calculating species' diversity. Ecosystem services considered here are (a) rate of pollen grain
deposition on one of three flower species by wild bees; total biomass of reef fish encountered in fixed-
effort dive surveys in (b) temperate and (b) tropical global regions; and (d) total above-ground carbon
in tropical forest plots. Correlations are between logged total function at each site, and logged species
diversity at a range of ell values (at 0.5 intervals) emphasizing the relative abundance of common
species' (negative ell values) or rare species (positive ell values). Vertical lines identify correlations at
commonly used diversities: inverse Simpson (dotted), exponential Shannon (dashed), and richness
(solid). Colors visually distinguish different community datasets.

Figure 4. Biodiversity-ecosystem function (BEF) correlation in observational data (figure 3) can be explained by correlation between total abundance and diversity (first row), correlation between mean per-capita function and diversity (second row), and interactions between these two factors (intractable, not shown). With a few exceptions, the abundance-diversity correlations roughly match the BEF correlation across the range of ell values used to calculate species diversities, while per-capita function-diversity correlations show countervailing trends in magnitude and direction. Compare (a, e) wild bee pollination, reef fish biomass in (b, f) temperate and (c, g) tropical regions, and (d, h) tropical forest above-ground biomass with corresponding panels in figure 3. Vertical lines identify correlations at commonly used diversities: inverse Simpson (dotted), exponential Shannon (dashed), and richness (solid). Colors visually distinguish different community datasets.

Figure S1. Community datasets (line color) differed in both the shapes of the diversity profiles (D vs. ell) and the degree to which diversity profiles differed between sites. For example, the grey-blue tree_carbon sites were all 1-Ha subplots from the contiguous BCI 50-Ha forest plot, and diversity profiles were very similar between subplots; by contrast the yellow bee_pollination sites (Floral visitors of *Polemonium reptans*) had variable structure with wide variety in richness (ell = 1, vertical solid lines), inverse dominance (large negative ell), and abundance (large positive ell). Each curve is the diversity profile for a single site; colors indicate a community dataset (set of sites within a region at which a single function was measured).