Biodiversity-ecosystem function relationships change in sign and magnitude across the Hill

diversity spectrum

Michael Roswell[1] mroswell@umd.edu **0000-0002-8479-9184**

Tina Harrison[2] tinaharrison09@gmail.com **0000-0001-8735-0289**

Mark A. Genung[2] mark.genung@louisiana.edu **0000-0002-9383-667X**

1 – Department of Entomology, University of Maryland, College Park, MD, USA

2 – Department of Biology, University of Louisiana at Lafayette, Lafayette, LA, USA

Keywords: abundance, biodiversity, diversity profile, ecosystem function, Hill number, rarity

Summary

Motivated by accelerating anthropogenic extinctions, decades of biodiversity-ecosystem function

(BEF) experiments show that ecosystem function declines with species loss from local communities.

Yet, at the local scale, changes in species' total and relative abundances are more common than

species loss. The consensus best biodiversity measures are Hill numbers, which use a scaling

parameter, ℓ, to emphasize rarer versus more common species. Shifting that emphasis captures

distinct, function-relevant biodiversity gradients beyond species richness. Here, we hypothesized that

Hill numbers that emphasize rare species more than richness does may distinguish large, complex,

and presumably higher-functioning assemblages from smaller and simpler ones. In this study, we

tested which values of ℓ produce the strongest BEF relationships in community datasets of ecosystem

functions provided by wild, free-living organisms. We found that ℓ values that emphasized rare species

more than richness does most often correlated most strongly with ecosystem functions. As emphasis

shifted to more common species, BEF correlations were often weak and/or negative. We argue that

unconventional Hill diversities that shift emphasis towards rarer species may be useful for describing

biodiversity change, and that employing a wide spectrum of Hill numbers can clarify mechanisms

underlying BEF relationships.

Introduction

A central question in community ecology is, "how will ongoing shifts in biodiversity affect ecosystem function?" In experiments that vary species richness while controlling other community properties, the answer has been clear for some time: ecosystem function has a positive, saturating relationship with species richness [1–3]. There is ongoing interest in "scaling up" research to resolve whether similar patterns hold in natural ecosystems [4,5]. However, richness is not a robust measure of biodiversity in observational data taken from natural ecosystems [6], in large part because most species are rare [7] and likely to be absent from samples. Further, richness often tracks real-world biodiversity gradients poorly, because species composition and abundance can change dramatically with little to no change in observed species richness [8–10]. Therefore, other metrics of biodiversity may provide improved clarity about the real-world linkages between biodiversity and ecosystem function (BEF).

There are both historical and conceptual reasons that BEF research has focused on richness as a measure of biodiversity. Since at least the 1960s, there has been extensive research on how productivity affects species richness [1]. Motivated by intensifying biodiversity loss in the 1980s, declines in richness were (at least implicitly) the global change pattern that seminal BEF studies, with their focus on species loss (e.g., [11]), aimed to understand. This prompted a wave of experiments on how species richness affects ecosystem function [2]. Thus, richness was a natural choice, both because of ecology's long focus on how richness might *respond* to ecosystem functions like productivity, and because of a collective sense that species loss was the correct, or at least most convenient, way to frame anthropogenic changes in biodiversity. Furthermore, richness was considered a good proxy for functional diversity and redundancy, which were considered the key mechanisms through which biodiversity maintains ecosystem function [12–14]. However, the choice of richness may not have been based on theoretical expectation that richness, rather than other abundance-weighted diversity measures, best described functionally important biodiversity gradients.

Using species richness as the key biodiversity measure poses methodological problems for BEF research, especially when community properties other than richness vary, as in naturalistic systems. Species richness is not only sensitive to the extent and depth of sampling, but also to the distribution

of relative abundances in the sampled assemblage. To illustrate this, consider the difficulty of

accurately measuring species richness in a community with one hyper-dominant species and many

very rare species, versus measuring richness in a community in which abundance is evenly

distributed. Richness, like other diversity measures, summarizes the distribution of relative

abundances in an assemblage, and when estimated from data, cannot be independent from that

distribution, even if such a measure were desirable [6,15]. However, different diversity measures vary

in the extent to which they emphasize rare vs. common species, with species richness heavily

emphasizing rare species. A unified family of diversity measures, known as "Hill numbers" or "Hill

diversities," summarizes a distribution of relative abundances as the abundance-weighted,

generalized mean rarity [16–18]. Hill numbers are governed by a scaling parameter, ℓ, that scales

species rarity when computing the mean, and higher values of ℓ afford more leverage, or emphasis, to

rare species, while lower values emphasize common species more [18].

The Hill diversity of an assemblage is not a single value, but rather a spectrum that varies

continuously across ℓ [6,16] (Figure 1), raising the question of how ecosystem function relates to

biodiversity measures with different emphasis on common vs. rare species. While several recent

studies have compared whether richness ($\ell = 1$), exponentiated Shannon ($\ell = 0$), or inverse Simpson (ℓ

= -1) best explains ecosystem function [19–22], there has been no examination of how Hill numbers

relate to ecosystem function across a wide range of ℓ values. This is a striking knowledge gap because,

although nearly all studies of the relationship between biodiversity and ecosystem function have used

species richness as a measure of diversity [3], other diversity measures could both better describe real

world variation in biodiversity, and also have stronger links to ecosystem functioning. Furthermore,

Hill diversities with $\ell > 1$, which emphasize rare species even more than richness does, have scarcely

been studied at all, not to mention in relationship to ecosystem function. Thus, biodiversity-function

studies may be *underestimating* the importance of biodiversity for function by not considering Hill

diversities with different emphases on rare and common species via different values of the scaling

parameter ℓ.

Despite clear declines in richness at the global scale, local changes in biodiversity and their

connection to function are likely better captured by measuring total abundance and species' relative

abundances [8,23], for at least three reasons. First, as already discussed, observed richness is a poor

predictor of true richness [24], and good estimators of true richness based only on species frequencies

in samples may never exist [25]. Thus, even if underlying variation in species richness correlates

strongly with, or even drives, ecosystem function, estimating richness from samples could severely

obscure the underlying pattern. Second, although observed richness does increase with observed

abundance, to the extent that abundance *per se* drives function, Hill diversities that better reflect

abundance (i.e., when $\ell \gg 1$) should be stronger correlates of function than richness ($\ell = 1$) is. Third, if

diversity effects on function are mediated by positive species interactions [3,26–28], more probable

and stronger between equally abundant species [29,30], Hill diversities that better reflect the

probability of interspecific encounter (e.g. Hill-Simpson diversity, at $\ell = -1$ [31]) might explain function

better than richness does.

Here, we ask how biodiversity-ecosystem function correlations in observational datasets change in

sign and magnitude across a wide range of values of the scaling parameter ℓ. In natural communities,

classic BEF mechanisms such as selection and complementarity co-occur with other sources of

variation in function, such as abundance, evenness, demography, and environment [32,33].There is no

way to standardize BEF modeling approaches across heterogeneous natural systems [34–36], so

researchers variously use mathematical partitions [37] or regression and path analysis [32,38] to

partially account for subsets of these factors depending on their system knowledge, assumptions, and

preferred study focus. In order to focus attention on the possibilities of a wide diversity spectrum, we

keep analyses simple and general by presenting the overall correlation between total function and

diversity across natural communities, which represents the net effects of many factors. We focus on

ecosystem functions that can be expressed as the product of mean per-capita function and total

abundance, which works well for many functions [39,40]. As a first step towards a more granular

approach, we also present separate correlations between diversity and total abundance [41] and

between diversity and per-capita function, which captures selection effects due to shifts in

community composition, together with complementarity effects on individual-level function. We

explore whether diversity measures other than richness can better explain BEF patterns and

potentially help identify BEF mechanisms in natural systems.

In this study, we use observational community datasets on three ecosystem functions to ask:

1) Which values of the Hill diversity scaling factor ℓ produce the strongest biodiversity-
ecosystem function correlations?
2) How do biodiversity-ecosystem function correlations change in sign and strength over a wide
range of Hill diversities?
3) What is the role of absolute abundance in shaping BEF correlations over the Hill diversity
spectrum?
128

Materials and Methods

To find how biodiversity-ecosystem function correlations change with different diversity scaling, we
used previously published community datasets that recorded both the abundance and function of
species at multiple sites. We chose datasets of disparate ecosystem functions and spatial scales:
pollination by wild bee visitors to a landscape-scale array of three plant species [42,43], reef fish
biomass from dive surveys replicated within 32 globally distributed geographic regions [44,45], and
above-ground tree biomass in census plots replicated within four tropical forests [46,47] (Table 1). In
each system, total function of a community can be estimated as the summed contribution across
species (or individuals) present in the community. Pollination was measured as the product of first,
the typical number of pollen grains deposited during a single visit of a focal bee taxon to the focal
plant species, and second, the number of observed visits to the focal plant species by the focal bee
taxon. Reef fish biomass was measured by visually estimating individual fish body lengths during dive
surveys, which were then used to calculate biomass using species-specific allometric equations.
Tropical tree biomass was measured by converting observed individual diameter at breast height into
biomass estimates using taxon-specific allometric equations that included information about wood
density [48,49]. In total, we used 39 community datasets, each consisting of one function measured
across a collection of assemblages.

Which value of ℓ produces the strongest biodiversity- ecosystem function correlations?
We computed Hill diversity as a function of species relative abundances, *p_1, p_2, …, p_S,* and a
scaling factor, ℓ, using the formula

$$D = \left(\sum_{i=1}^{S} p_i \left(\frac{1}{p_i} \right)^{\ell} \right)^{1/\ell}$$

or its limit as ℓ approaches zero (exponential Shannon entropy) [18,50]. Historically, it has been more

common to express Hill diversity with a scaling parameter "a" or "q", equal to 1-ℓ [6,16,51]. We used

the expression above (with the scaling parameter ℓ, instead), to highlight that this expression is a

specific example of the more general weighted power mean [17,18]. This ℓ formulation's biggest

advantage over the q-formulation is that it clarifies the differences between weights (abundances)

and the rarity scaling controlled by the parameter ℓ [18]. Furthermore, because the equations for

richness, Hill Shannon, and Hill-Simpson diversities straddle $\ell = 0$, this formulation may support the

intuition that the spectrum of Hill diversity measures can extend in either direction (Figure 1), either

further emphasizing common species (as ℓ gets increasingly negative) or further emphasizing rare

ones (as ℓ gets increasingly positive).

We used observed species abundances to calculate species diversities at each site as the Hill diversity

along a wide range of ℓ values (from -10 to +10 at intervals of 0.05) (Figure S1). We calculated total

function as the sum of species' functions at each site. We computed the correlation between the

natural logarithm of each diversity and the natural logarithm of total function, across all sites in the

community dataset (hereafter, the "BEF correlation"). We focused on the logarithms of function, and

later, abundance variables because log(total function) = log(abundance) + log(per capita function),

and we also log-transformed diversity because we anticipated that across the wide array of

ecosystems considered, multiplicative, rather than additive, differences in diversity would be most

comparable. To identify the ℓ value that produced the strongest BEF correlation in each community

dataset, we plotted the correlation against the scaling factor ℓ. We identified the single ℓ value with

the largest absolute correlation (i.e., largest R-squared for the relationship between log(diversity) and

log(function)).

How do biodiversity-ecosystem function correlations change in sign and strength over a wide range of

Hill diversities?

To determine not only which ℓ value produced the strongest BEF correlation across community

datasets, but also to see how adjusting the Hill diversity scaling parameter affects BEF relationships

more comprehensively, we plotted the BEF correlation against the Hill diversity scaling factor ℓ for

each community dataset. We examined curves to identify patterns in the sign and strength of the BEF

correlation along the spectrum of emphasis on common and rare species.

What is the role of absolute abundance in shaping BEF correlations?

To begin to separate effects of total and relative abundance on BEF correlations across the Hill

diversity spectrum, we looked separately at the relationships between diversity and two

complementary components of total function, namely total abundance and mean per-capita function.

We used the same graphical approach we described above, regarding the sign and strength of the BEF

correlation across the Hill diversity spectrum. For each community dataset, we found the correlation

between the natural logarithm of Hill diversity at each site and either the natural logarithm of total

abundance at each site, or the natural logarithm of mean per-capita function at each site, and plotted

these correlations against the Hill diversity scaling parameter ℓ. Although on the logarithmic scale,

abundance and mean per-capita function combine additively to create total function, the BEF

correlation does not additively decompose into abundance by biodiversity and per-capita function by

biodiversity correlations, as there is also covariance between abundance and per-capita function.

Acknowledging this caveat, we suggest that by partitioning total function into additive components

and examining how each of these relates to biodiversity gradients across the Hill spectrum, we can

better characterize the role of total abundance in generating patterns in the BEF correlation itself.

Results

Which value of ℓ produces the strongest biodiversity- ecosystem function correlations?

For most datasets, the strongest biodiversity-ecosystem correlations were located at or just above

richness ($\ell = 1$), with a mode at $\ell = 1.5$ (Figure 2). A substantial minority (11 of 39 datasets) had

strongest BEF correlations at values of $\ell > 5$, including a peak at $\ell = 10$, the largest value of ℓ we

considered. There were a few outliers: Two tree carbon storage datasets had their strongest BEF

correlations near inverse Simpson ($\ell = -1$) and exponential Shannon ($\ell = 0$) diversities, and a single fish

dataset had a strongest BEF correlation at $\ell = -10$, the smallest value of ℓ we considered (Figure 2).

How do biodiversity-ecosystem function correlations change in sign and strength over a wide range of

Hill diversities?

Across all ecosystem functions, we found common patterns in the relationship between the BEF
correlation and the Hill diversity scaling parameter, ℓ. When ℓ was < 1, the diversity-function
correlation was typically weak and could be positive or negative (Figure 3). Near ℓ = 1, the diversity-
function correlation rapidly increased, although a substantial minority of community datasets first
showed a sharp negative turn in the relationship near Hill-Simpson and Hill-Shannon diversities
(Figure 3 b-d). Across all the datasets we considered, the mean correlation between log(diversity) and
log(total function) was not significantly different from zero for either Hill-Simpson or Hill-Shannon
diversity (p>0.28 for two-sided Student's t-test, with no correction for multiple tests). At richness (ℓ =
1), almost all datasets showed positive diversity-function correlations, with the mean R^2 = 0.381. For
most datasets, the strongest correlations were located near richness, with a mode near ℓ = 1.5, where
the mean R^2 was 0.445, after which the diversity-function correlation slowly declined as ℓ values
continued to increase (Figure 3). A substantial minority of datasets showed continually stronger
relationships as ℓ increased (some profiles in Figure 3 b, c), leading to highest R^2 values at or near the
maximum ℓ we considered (ℓ = 10).
225
This study was not designed to contrast trends between ecosystem functions, but it is important to
note that the relationship between the BEF correlation, and the emphasis the diversity metric puts on
rare vs. common species (i.e., the value of ℓ), did not appear uniform across systems. For the three bee
community datasets, total pollen deposition and bee diversity were positively correlated at every
value of ℓ. Correlation strength peaked at richness (ℓ = 1) or just beyond (ℓ = 2), but remained relatively
strong across all higher values of ℓ (Figure 3a). For the 32 reef fish community datasets, total fish
biomass and fish diversity tended to be weakly and often negatively correlated at low ℓ values.
Correlation strength tended to peak either slightly above richness at ℓ = 1.5, or grow with ℓ for an
observed peak near the maximum value considered (ℓ = 10) (Figure 2, Figure 3 b, c). When considering
either very high or very low ℓ values, note that at either end of the Hill number spectrum, diversities
rapidly converged towards their maximum or minimum asymptote. Thus, large changes in the BEF
correlation rarely occurred outside a fairly narrow range between ℓ = -2 and 2. Finally, the four tropical
tree community datasets showed generally weak correlations. In two tree datasets, BEF correlation
strength peaked at intermediate ℓ values where the BEF correlation was strongly negative (Figure 3 d).
In another (Barro Colorado Island) diversity-function correlation was negative even at high ℓ values
(Figure 3 d, orange line), but modestly positive for negative ℓ values.

What is the role of absolute abundance in shaping BEF correlations?

As expected, the relationship between diversity and abundance was mostly similar to the relationship

between diversity and function, as total abundance underlies function in our datasets. This can be

seen in the similar shape of the curves showing the correlation between log(diversity) and either

log(function) (Figure 3) or log(abundance) (Figure 4 a-d), as the sign and strength of correlation

typically moved in similar ways across the ℓ spectrum. In almost all cases, the correlation between

log(diversity) and log(abundance) was very strong (and in many cases approached unity), for large,

positive values of the Hill diversity scaling parameter ℓ. As previously remarked, this result is a

mathematical inevitability when datasets contain very rare species/singletons. Additionally, across

datasets, we found that the rise towards the high correlation observed for large ℓ values typically

occurred in the range of ℓ values typically considered by ecologists (-1 to 1), likely reflecting biological

and sampling linkages between abundance and diversity; the correlation frequently saturated once ℓ

was greater than two. While for some community datasets, diversity was largely independent of

abundance for negative ℓ values, we also saw community datasets in which log(abundance) and

log(diversity) had modest to strong negative correlation across negative ℓ values. Because Hill

diversities typically change little with ℓ below around -2 [6], this result implies that in these systems,

total abundance and the degree of dominance were positively linked [52].

While the curves in Figures 3 a-d and 4 a-d show strong resemblance, for some community datasets

the BEF and diversity-abundance relationships diverged, implying those BEF relationships resulted

from processes other than abundance. For example, correlations between abundance and diversity

were always strongly positive for the reef fish data for large, positive ℓ values, but in some reef fish

community datasets, correlations between diversity and function were only weakly positive at higher

ℓ values (Figure 4 b, c). Such divergences between the diversity-function and diversity-abundance

curves could be due to strong and/or countervailing relationships between mean per-capita function

and diversity, which also showed some overall patterns across community datasets (Figure 4 e-h). In

general, Hill diversities with negative ℓ values were positively related to per-capita function,

suggestive of a positive relationship between evenness and mean per-capita function. This pattern

was not ubiquitous, however, with notable exceptions in both tree and bee community datasets

(Figure 4 e, h). We found that the correlation between diversity and mean per-capita function often

exhibited a positive peak at intermediate ℓ values, a pattern particularly pronounced in the reef fish

community datasets (Figure 4 f, g). Finally, there was a tendency towards a negative correlation

between log(diversity) and log(mean per-capita function) for larger, positive values of ℓ (when

diversity becomes largely synonymous with abundance), though the strength of this relationship was

variable.

Discussion

Diversities near richness ($1 < \ell < 2$) often had strong positive correlations with ecosystem function

(Figure 2), echoing a large body of research emphasizing the importance of richness in BEF

relationships [3]. However, this was not a foregone conclusion, because most empirical BEF work does

not consider other portions of the diversity spectrum [8], and because much of the theoretical and

empirical work is grounded in closed communities where richness has a clear interpretation. We were

surprised by the consistent performance of richness, because of three concerns we described in the

introduction: first, observed richness is not a robust biodiversity measure in observational data;

second, Hill diversities with high ℓ should better explain function in systems with highly variable total

abundance; third, ℓ values that emphasize the probability of interspecific encounter ($-1 \leq \ell < 1$) might

better explain function if species interactions are very important. The disconnect between observed

and true richness is not, practically speaking, a resolvable problem and thus we cannot evaluate how

much this first issue is affecting our results [25]. In the following paragraphs, we explore the latter two

points, namely: how do Hill diversities near richness outperform Hill diversities with higher ℓ that

better reflect abundance, and why we might have found such low explanatory power for Hill

diversities that should capture the effects of (potentially positive) species interactions. Although all

the information about relative abundances is contained in any continuous interval along the Hill

diversity spectrum, at different values of ℓ, different aspects of the abundance distribution are

emphasized. To better ground our discussion, we use an admittedly imprecise simplification, and

refer to ℓ values as falling within the "inverse dominance range" ($\ell < 1$), the "evenness emphasis

range" ($-1 \leq \ell < 1$), the "rare emphasis range" ($1 \leq \ell < 2$, justification follows), or the "abundance

emphasis range" ($\ell > 2$).

As ℓ values increase, empirical diversities values are increasingly dependent on total abundance, since

the abundance of the rarest species is typically at the lower bound set by the detection threshold (for

count data, one). Hill diversities in the "rare emphasis range," like richness itself, are affected by

abundance as well as diversity *per se*. This is reasonably viewed as a sampling nuisance [6,24]. But

viewed in another light, measures that combine abundance, relative abundance, and richness

components could predict function well because, owing partly to their sampling properties, they

describe salient biodiversity gradients. In fact, our results showed that ℓ values in the rare emphasis

range that, compared to richness, are relatively more sensitive to abundance tended to better explain

function. Rare-emphasis Hill diversities increase with richness, abundance, and often also dominance.

As a result, they might vary with the importance of sampling effects whereby higher-functioning

(likely, highly abundant) species are more likely to occur in species-rich assemblages (a kind of

selection effect) [34,37,53]. We found that rare-emphasis Hill diversities tended to explain function

best, with $1 < \ell < 2$ performing better than richness, $\ell = 1$, but because they still reflect compositional

heterogeneity, also better than abundance alone (Figures 2-3).

Hill diversities in the evenness emphasis range ($-1 \leq \ell < 1$) should capture the effects of species

interactions by emphasizing the probability of interspecific encounter, but these diversities explained

function poorly. This was partly unexpected because in several real-world systems, function increases

with evenness [54–57], which increases Hill diversity for $\ell < 1$ [58–60]. Additionally, in the evenness

emphasis range, sample Hill diversities have relatively good statistical properties as estimators of true

diversity, and asymptotic estimators [24] can further improve the situation, largely avoiding the

robustness issues we highlight with species richness. Instead, the observed weak explanatory power

of Hill diversities with ℓ values in the evenness emphasis range is because functions analyzed here are

the product of two components, abundance and per-capita function, which each showed different

responses to increasing ℓ. Abundance-diversity relationships often followed function-diversity

correlations (compare Figure 3 with the top row of Figure 4). In other words, across the ℓ spectrum,

Hill diversities had nearly the same relationship with abundance and with function, underlining the

necessity of accounting for the role of total abundance in BEF research [61]. However, Hill diversities

in the evenness emphasis range deviated from this pattern, instead exhibiting often strong,

countervailing relationships with abundance and per-capita function (Figure 4). This fits with previous

BEF literature, which anticipates a variety of mechanisms linking evenness and ecosystem function,

without a clear prediction for when the net result is positive or negative [62–65].

The relationship between diversity and per-capita function differed from the diversity-abundance

relationship, with correlation coefficients for per-capita function generally decreasing with increases

in ℓ, but often showing a positive peak in the evenness emphasis range (Figure 4). Positive species

interactions, including those that increase per-capita function, are expected to explain total function

[53,66,67]. Our results partly support these expectations, as Hill diversities in the evenness emphasis

range, which should track the probability of interspecific encounter, were positively associated with

per-capita function, even as they tended to be negatively associated with abundance. As we increased

ℓ, the correlation between Hill diversity and per-capita function disappeared near richness ($\ell = 1$), also

pushing against expectations that richness best captures function-relevant biodiversity gradients. In

the rare emphasis and abundance emphasis ranges, we typically found a negative correlation

between Hill diversity and per-capita function. Recent work highlighting the functional contribution of

rare species to ecosystem function led us to suspect the opposite might occur [43,68–70]. The

observed negative correlation likely reflects spatial constraints and/or fundamental tradeoffs

between having many, smaller-bodied individuals versus fewer larger ones [71,72]. This scenario is

particularly easy to imagine for trees crowding in fixed-area plots, which physically and energetically

prohibit arbitrarily large numbers of the largest trees. Similar energetic and spatial constraints limit

the number of very large fish that might be seen in a single dive. Thus, we suspect that one reason we

saw a decline in the correlation between mean per-capita function and diversity with increasing ℓ in

the fish and tree datasets is decreases in per-capita function due to crowding.

Even as Hill diversities in the rare emphasis range most often explained total function best, Hill

diversities with ℓ values in the abundance emphasis range also explained function well, and should

not be discounted. Hill diversities in the abundance emphasis range were the *best* predictor of

function in a substantial minority of datasets (Fig. 2), and for nearly all datasets were *strong* predictors

of function (Fig. 3, far right of x-axes). This was expected because of a general link between higher

abundance and higher function [41,61,73–76]. Even as Hill diversities in the "abundance emphasis"

range related strongly to function, we also note that abundance (like evenness) can relate to Hill

diversity across the full spectrum of ℓ values. For example, if high-abundance sites tend to be

dominated by many individuals of one or a few species [52], Hill diversities that emphasize inverse

dominance will decrease with abundance. Thus, we should not expect that strong effects of

abundance on function are captured exclusively at high values of ℓ.

The explanatory power of Hill diversity changed nonlinearly with increases in ℓ, as multiple facets of

community structure (e.g., richness, abundance, evenness) affect function simultaneously. If we had

found a monotonic strengthening of BEF relationships with increasing ℓ, we would argue that Hill

"diversities" with large positive scaling parameters were simply abundance metrics masquerading as

measures of diversity. Instead, we found, across a variety of regions, taxa, and ecosystem functions,

intermediate, positive ℓ values in the "rare emphasis" range tended to produce the strongest BEF

relationships (Fig. 3). All Hill diversities with positive ℓ values (including richness) tend to increase with

both abundance and richness, which we argue can be a useful property, especially for BEF research.

Because the goal of summarizing species' abundances with diversity metrics is to distill complex,

multivariate information [17], this claim is not radical. In fact, Hill diversities that emphasize rare

species more than richness does can reflect intuitive notions of diversity, which include both high

density and high compositional variation [77]. Our study points to the need for further theoretical

work to explicate the meaning of these seldom-used Hill diversities in the rare emphasis range, and

their linkages to ecosystem function.

By considering Hill diversities over a wide range of ℓ, we place ourselves at odds with the convention

that Hill diversities should be considered only when $\ell \leq 1$ [6,16,17,51,78]. The most compelling

argument for that restricted range of scaling parameters is presented by Patil and Taillie, who argued

that diversity should not decrease when abundance is shifted from more to less abundant species,

including to species with zero abundance, a variation on Dalton's "principle of transfers" [17,79]. This

diversity property does not hold for Hill diversity when $\ell > 1$, which has species richness as its

minimum, occurring in the perfectly even community, and increases (given richness and abundance)

as some species get progressively rarer. A more pragmatic argument comes from Chao et al., who

noted that estimating the relative abundance of rare species is an increasing problem for diversity

measures as ℓ increases; they therefore suggest using only more estimable Hill diversities with $\ell \leq 1$ [6].

However, theoretical work suggests that even richness ($\ell = 1$) is poorly estimated [24,25], and by this

logic should not be used either. Finally, and most generally, diversity measures have traditionally

been considered separate from abundance/density measures (but see [80,81]), whereas with

increasing ℓ values, observed diversity and observed abundance tend to be more strongly correlated

(and in fact approach a correlation of one in our datasets). Despite these arguments, our results show

that Hill diversities with $\ell > 1$ are meaningful ecological diversity measures, at least in the sense that

they convey more information about function than do more widely-used Hill numbers. Choosing to

exclude these Hill diversities might be desirable for some conceptualizations of diversity, but we are

opening the narrower question of which Hill diversities—with their variable emphasis on richness,

abundance, evenness, and dominance—best explain ecosystem function. In this pursuit, allowing

diversity metrics to highlight absolute abundance is valuable.

Our correlational analyses of observational data do not consider confounding variables, which may

obscure links between biodiversity and ecosystem function, across the Hill diversity spectrum.

Experimental and statistical approaches to better account for environmental drivers of diversity

and/or function and more rigorously trace causal pathways (e.g., [38]) will be useful in validating and

extending our findings [82]. One obvious effect of ignoring confounding variables is that the BEF

correlations we found are likely to be low, as the confounding variables add noise that is not

accounted for. Future work linking environmental gradients and other confounding variables to

particular regions of the Hill diversity spectrum (e.g., regions emphasizing inverse dominance,

evenness, rare species, or total abundance) may also clarify mechanisms underlying BEF

relationships.

As global changes lead to shifting species abundances, ecologists must continue to describe and

predict how these shifts impact ecosystems and the way they function. Yet, understanding the

separate and combined roles of total and relative abundance in mediating ecosystem function

remains a difficult challenge, in large part because total abundance is inextricably linked to diversity

measures. It is mathematically linked for large, positive ℓ values. It is practically constrained by

sampling effects for ℓ closer to 1 (i.e., near species richness). As ℓ becomes negative, Hill diversities

may lose their dependence on total abundance [24]. However, in the majority of community datasets,

we saw at least weak negative correlations between negative-ℓ Hill diversities and observed

abundances, likely due to increasing dominance in more abundant systems [52]. Overall, this suggests

that in observational contexts, simple partitioning of abundance and diversity effects may not be

tractable, at least not in a satisfying manner [34,35]. Since no single-best diversity measure is likely to

emerge for all BEF studies, we encourage researchers to be open-minded towards Hill diversities

across a wide spectrum of ℓ values and their potential links to mechanisms underlying BEF

relationships.

Competing Interests

We have no competing interests.

Acknowledgements

We thank Rachael Winfree for key discussions that sparked this work, Jonathan Dushoff for providing

insight on Hill numbers, and participants in the Hakai Synthesizing Biodiversity Seminar Series for

useful feedback on preliminary results. Mary O'Connor and two anonymous reviewers provided

helpful feedback on an earlier draft of this manuscript. We gratefully acknowledge all data collectors

and data holders. Data for three tropical forests (Pasoh, Indonesia; Yasuní, Ecuador; and Volcán Barva,

Costa Rica) were originally made available to us by the Tropical Ecology Assessment and Monitoring

(TEAM) Network, a collaboration between Conservation International, the Missouri Botanical Garden,

the Smithsonian Institution, and the Wildlife Conservation Society, and partially funded by these

institutions, the Gordon and Betty Moore Foundation, and other donors.

References

1.   Loreau M *et al.* 2001 Biodiversity and Ecosystem Functioning: Current Knowledge and Future

Challenges. *Science (80-.).* **294**, 804–808. (doi:10.1126/science.1064088)

2.   Hooper DU *et al.* 2005 Effects of biodiversity on ecosystem functioning: A consensus of current

knowledge. *Ecol. Monogr.* **75**, 3–35. (doi:10.1890/04-0922)

3.   Cardinale BJ *et al.* 2012 Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67.

(doi:10.1038/nature11148)

4.   Gonzalez A *et al.* 2020 Scaling-up biodiversity-ecosystem functioning research. *Ecol. Lett.* **23**,

757–776. (doi:10.1111/ele.13456)

5.   Qiu J, Cardinale BJ. 2020 Scaling up biodiversity–ecosystem function relationships across

space and over time. *Ecology* **101**, 0–2. (doi:10.1002/ecy.3166)

6.   Chao A, Gotelli NJ, Hsieh TC, Sander EL, Ma KH, Colwell RK, Ellison AM. 2014 Rarefaction and

extrapolation with Hill numbers: a framework for sampling and estimation in species diversity

studies. *Ecol. Monogr.* **84**, 45–67. (doi:10.1890/13-0133.1)

7. McGill BJ *et al.* 2007 Species abundance distributions: Moving beyond single prediction

theories to integration within an ecological framework. *Ecol. Lett.* **10**, 995–1015.

(doi:10.1111/j.1461-0248.2007.01094.x)

8. Hillebrand H *et al.* 2018 Biodiversity change is uncoupled from species richness trends:

consequences for conservation and monitoring. *J. Appl. Ecol.* **55**, 169–184. (doi:10.1111/1365-

2664.12959)

9. Brose U, Hillebrand H. 2016 Biodiversity and ecosystem functioning in dynamic landscapes.

Philos. Trans. R. Soc. B Biol. Sci. **371**. (doi:10.1098/rstb.2015.0267)

10. Spaak JW *et al.* 2017 Shifts of community composition and population density substantially

affect ecosystem function despite invariant richness. *Ecol. Lett.* **20**, 1315–1324.

(doi:10.1111/ele.12828)

11. Naeem S, Thompson LJ, Lawler SP, Lawton JH, Woodfin RM. 1994 Declining biodiversity can

alter the performance of ecosystems. *Nature* **368**, 734–737. (doi:10.1038/368734a0)

12. Walker BH. 1992 Biodiversity and Ecological Redundancy. *Conserv. Biol.* **6**, 18–23.

(doi:10.1046/j.1523-1739.1992.610018.x)

13. Naeem S. 2002 Disentangling the impacts of diversity on ecosystem functioning in

combinatorial experiments. *Ecology* **83**, 2925–2935. (doi:10.1890/0012-

9658(2002)083[2925:DTIODO]2.0.CO;2)

14. Tilman D, Downing JA. 1994 Biodiversity and stability in grasslands. *Nature* **367**, 363–365.

15. Chao A, Jost L. 2012 Coverage-based rarefaction and extrapolation: Standardizing samples by

completeness rather than size. *Ecology* **93**, 2533–2547. (doi:10.1890/11-1952.1)

16. Hill MO. 1973 Diversity and evenness: a unifying notation and its consequences. *Ecology* **54**,

427–432. (doi:10.2307/1934352)

17. Patil GP, Taillie C. 1982 Diversity as a concept and its measurement. *J. Am. Stat. Assoc.* **77**, 548–

561. (doi:10.1080/01621459.1982.10477845)

18. Roswell M, Dushoff J, Winfree R. 2021 A conceptual guide to measuring species diversity. *Oikos*

130, 321–338. (doi:10.1111/oik.07202)

19. Yasuhara M, Doi H, Wei CL, Danovaro R, Myhre SE. 2016 Biodiversity-ecosystem functioning

relationships in long-term time series and palaeoecological records: Deep sea as a test bed.

Philos. Trans. R. Soc. B Biol. Sci. **371**. (doi:10.1098/rstb.2015.0282)

20. Schweiger AK, Cavender-Bares J, Townsend PA, Hobbie SE, Madritch MD, Wang R, Tilman D,

Gamon JA. 2018 Plant spectral diversity integrates functional and phylogenetic components of

biodiversity and predicts ecosystem function. *Nat. Ecol. Evol.* **2**, 976–982. (doi:10.1038/s41559-

018-0551-1)

21.   Yang JW, Wu W, Chung CC, Chiang KP, Gong GC, Hsieh CH. 2018 Predator and prey biodiversity

relationship and its consequences on marine ecosystem functioning - Interplay between

nanoflagellates and bacterioplankton. *ISME J.* **12**, 1532–1542. (doi:10.1038/s41396-018-0111-3)

22.   Schmidt ML, Biddanda BA, Weinke AD, Chiang E, Januska F, Props R, Denef VJ. 2020

Microhabitats are associated with diversity-productivity relationships in freshwater bacterial

communities. *FEMS Microbiol. Ecol.* **96**, 1–12. (doi:10.1093/FEMSEC/FIAA029)

23.   Dornelas M, Gotelli NJ, McGill B, Shimadzu H, Moyes F, Sievers C, Magurran AE. 2014

Assemblage time series reveal biodiversity change but not systematic loss. *Science (80-.).* **344**,

296–299. (doi:10.1126/science.1248484)

24.   Chao A, Jost L. 2015 Estimating diversity and entropy profiles via discovery rates of new

species. *Methods Ecol. Evol.* **6**, 873–882. (doi:10.1111/2041-210X.12349)

25.   Haegeman B, Hamelin J, Moriarty J, Neal P, Dushoff J, Weitz JS. 2013 Robust estimation of

microbial diversity in theory and in practice. *ISME J.* **7**, 1092–1101. (doi:10.1038/ismej.2013.10)

26.   Fargione J, Tilman D, Dybzinski R, Lambers JHR, Clark C, Harpole WS, Knops JMH, Reich PB,

Loreau M. 2007 From selection to complementarity: Shifts in the causes of biodiversity-

productivity relationships in a long-term biodiversity experiment. *Proc. R. Soc. B Biol. Sci.* **274**,

871–876. (doi:10.1098/rspb.2006.0351)

27.   Reich PB, Tilman D, Isbell F, Mueller K, Hobbie SE, Flynn DFB, Eisenhauer N. 2012 Impacts of

biodiversity loss escalate through time as redundancy fades. *Science (80-.).* **336**, 589–592.

(doi:10.1126/science.1217909)

28.   Tilman D, Isbell F, Cowles JM. 2014 Biodiversity and ecosystem functioning. *Annu. Rev. Ecol.*

Evol. Syst. **45**, 471–493. (doi:10.1146/annurev-ecolsys-120213-091917)

29.   Jiang L, Wan S, Li L. 2009 Species diversity and productivity: Why do results of diversity-

manipulation experiments differ from natural patterns. *J. Ecol.* **97**, 603–608.

(doi:10.1111/j.1365-2745.2009.01503.x)

30.   Crowder DW, Northfield TD, Strand MR, Snyder WE. 2010 Organic agriculture promotes

evenness and natural pest control. *Nature* **466**, 109–112. (doi:10.1038/nature09183)

31.   Chase JM, Knight TM. 2013 Scale-dependent effect sizes of ecological drivers on biodiversity:

Why standardised sampling is not enough. *Ecol. Lett.* **16**, 17–26. (doi:10.1111/ele.12112)

32.   Duffy JE, Lefcheck JS, Stuart-Smith RD, Navarrete SA, Edgar GJ. 2016 Biodiversity enhances

reef fish biomass and resistance to climate change. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 6230–6235.

(doi:10.1073/pnas.1524465113)

33.   Mori AS, Osono T, Cornelissen JHC, Craine J, Uchida M. 2017 Biodiversity–ecosystem function

relationships change through primary succession. *Oikos* **126**, 1637–1649.

(doi:10.1111/oik.04345)

34.   Harrison T, Winfree R, Genung MA. 2022 Price equations for understanding the response of

ecosystem function to community change. *Am. Nat.* **200**, 181–192. (doi:10.1086/720284)

35.   Harte J, Brush M, Newman EA, Umemura K. 2022 An equation of state unifies diversity,

productivity, abundance and biomass. *Commun. Biol.* **5**, 874. (doi:10.1038/s42003-022-03817-8)

36.   Hagan JG, Vanschoenwinkel B, Gamfeldt L. 2021 We should not necessarily expect positive

relationships between biodiversity and ecosystem functioning in observational field data. *Ecol.*

Lett. **24**, 2537–2548. (doi:10.1111/ele.13874)

37.   Fox JW. 2006 Using the Price Equation to partition the effects of biodiversity loss on ecosystem

function. *Ecology* **87**, 2687–2696. (doi:10.1890/0012-9658(2006)87[2687:UTPETP]2.0.CO;2)

38.   Grace JB *et al.* 2016 Integrative modelling reveals mechanisms linking productivity and plant

species richness. *Nature* **529**, 390–393. (doi:10.1038/nature16524)

39.   Balvanera P, Kremen C, Martínez-ramos M. 2005 Applying Community Structure Analysis to

Ecosystem Function : Examples from Pollination and Carbon Storage. *Ecol. Appl.* **15**, 360–375.

40.   Kremen C. 2005 Managing ecosystem services: What do we need to know about their ecology?

Ecol. Lett. **8**, 468–479. (doi:10.1111/j.1461-0248.2005.00751.x)

41.   Grime JP. 1998 Benefits of plant diversity to ecosystems: immediate, filter and founder effects.

J. Ecol. **86**, 902–910.

42.   Genung M, Reilly J, Williams N, Buderi A, Gardner J, Winfree R. 2022 Rare and declining bee

species are key to consistent pollination of wildflowers and crops across large spatial scales;

Dryad, Dataset. (doi:https://doi.org/10.5061/dryad.qnk98sfkc)

43.   Genung MA, Reilly J, Williams NM, Buderi A, Gardner J, Winfree R. 2022 Rare and declining bee

species are key to consistent pollination of wildflowers and crops across large spatial scales.

Ecology , 0–2. (doi:10.1002/ecy.3899)

44.   Lefcheck JS *et al.* 2021 Species richness and identity both determine the biomass of global reef

fish communities. *Nat. Commun.* **12**, 1–9. (doi:10.1038/s41467-021-27212-9)

45.   Lefcheck JS. 2021 Species richness and identity both determine the biomass of global reef fish

communities. *Dataset*. (doi:https://doi.org/10.25573/serc.16847029.v1)

46.   Condit R *et al.* 2000 Spatial patterns in the distribution of tropical tree species. *Science* **288**,

1414–1418.

47.   Cavanaugh KC *et al.* 2014 Carbon storage in tropical forests correlates with taxonomic diversity

and functional dominance on a global scale. *Glob. Ecol. Biogeogr.* **23**, 563–573.

(doi:10.1111/geb.12143)

48.   Zanne AE *et al.* 2009 Data From: Towards a worldwide wood economics spectrum. *Data Dryad*.

(doi:https://doi.org/10.5061/dryad.234)

49.   Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE. 2009 Towards a worldwide

wood economics spectrum. *Ecol. Lett.* **12**, 351–366. (doi:10.1111/j.1461-0248.2009.01285.x)

50.   Roswell M, Dushoff J. 2020 MeanRarity: Hill Diversity Estimation and Visualisation.

51.   Jost L. 2006 Entropy and diversity. *Oikos* **113**, 363–375. (doi:10.1111/j.2006.0030-1299.14714.x)

52.   MacDonald ZG, Nielsen SE, Acorn JH. 2017 Negative relationships between species richness

and evenness render common diversity indices inadequate for assessing long-term trends in

butterfly diversity. *Biodivers. Conserv.* **26**, 617–629. (doi:10.1007/s10531-016-1261-0)

53.   Loreau M, Hector A. 2001 Partitioning selection and complementarity in biodiversity

experiments. *Nature* **412**, 72–76. (doi:10.1038/35083573)

54.   Wilsey BJ, Potvin C. 2000 Biodiversity and ecosystem functioning: Importance of species

evenness in an old field. *Ecology* **81**, 887–892. (doi:10.1890/0012-

9658(2000)081[0887:BAEFIO]2.0.CO;2)

55.   Polley HW, Wilsey BJ, Derner JD. 2003 Do species evenness and plant density influence the

magnitude of selection and complementarity effects in annual plant species mixtures? *Ecol.*

Lett. **6**, 248–256. (doi:10.1046/j.1461-0248.2003.00422.x)

56.   Wittebolle L, Marzorati M, Clement L, Balloi A, Daffonchio D, Heylen K, De Vos P, Verstraete W,

Boon N. 2009 Initial community evenness favours functionality under selective stress. *Nature*

458, 623–626. (doi:10.1038/nature07840)

57.   Genung MA, Fox J, Winfree R. 2020 Species loss drives ecosystem function in experiments, but

in nature the importance of species loss depends on dominance. *Glob. Ecol. Biogeogr.* **29**, 1–11.

(doi:10.1111/geb.13137)

58.   Stirling G, Wilsey B. 2001 Empirical Relationships between Species Richness, Evenness, and

Proportional Diversity. *Am. Nat.* **158**, 286–299. (doi:10.1086/321317)

59.   Jost L. 2010 Independence of alpha and beta diversities. *Ecology* **91**, 1969–1974.

(doi:10.1890/09-0368.1)

60.   Chao A, Ricotta C, Aldo P. 2019 Quantifying evenness and linking it to diversity, beta diversity,

and similarity. *Ecology* **100**, 1–15. (doi:10.1002/ecy.2852)

61.   Winfree R, Fox JW, Williams NM, Reilly JR, Cariveau DP. 2015 Abundance of common species,

not species richness, drives delivery of a real-world ecosystem service. *Ecol. Lett.* **18**, 626–635.

(doi:10.1111/ele.12424)

62.   Stavert JR, Bartomeus I, Beggs JR, Gaskett AC, Pattemore DE. 2019 Plant species dominance

increases pollination complementarity and plant reproductive function. *Ecology* **100**, 1–16.

(doi:10.1002/ecy.2749)

63.   Clare DS, Culhane F, Robinson LA. 2022 Secondary production increases with species richness

but decreases with species evenness of benthic invertebrates. *Oikos* , 1–10.

(doi:10.1111/oik.08629)

64.   Filstrup CT, King KBS, McCullough IM. 2019 Evenness effects mask richness effects on

ecosystem functioning at macro-scales in lakes. *Ecol. Lett.* **22**, 2120–2129.

(doi:10.1111/ele.13407)

65.   Wang XY, Ge Y, Gao S, Chen T, Wang J, Yu FH. 2021 Evenness alters the positive effect of species

richness on community drought resistance via changing complementarity. *Ecol. Indic.* **133**,

108464. (doi:10.1016/j.ecolind.2021.108464)

66.   Tilman D, Lehman CL, Thomson KT. 1997 Plant diversity and ecosystem productivity:

Theoretical considerations. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 1857–1861.

(doi:10.1073/pnas.94.5.1857)

67.   Kirwan L *et al.* 2007 Evenness drives consistent diversity effects in intensive grassland systems

across 28 European sites. *J. Ecol.* **95**, 530–539. (doi:10.1111/j.1365-2745.2007.01225.x)

68.   Dee LE, Cowles J, Isbell F, Pau S, Gaines SD, Reich PB. 2019 When Do Ecosystem Services

Depend on Rare Species? *Trends Ecol. Evol.* **34**, 746–758. (doi:10.1016/j.tree.2019.03.010)

69.   Jousset A *et al.* 2017 Where less may be more: How the rare biosphere pulls ecosystems strings.

ISME J. **11**, 853–862. (doi:10.1038/ismej.2016.174)

70.   Lyons KG, Brigham CA, Traut BH, Schwartz MW. 2005 Rare species and ecosystem functioning.

613   *Conserv. Biol.* **19**, 1019–1024. (doi:10.1111/j.1523-1739.2005.00106.x)

614 71. Yoda K, Kira T, Ogawa H, Hozum K. 1963 Self-thinning in overcrowded pure stands under

615   cultivated and natural conditions. *J. Inst. Polytech. Osaka City Univerisity. Ser. D, Biol.* , 107–129.

616 72. Hughes RN, Griffiths CL. 1988 Self-thinning in barnacles and mussels: the geometry of packing.

617   *Am. Nat.* **132**, 484–491. (doi:10.1086/284866)

618 73. Dangles O, Malmqvist B. 2004 Species richness-decomposition relationships depend on species

619   dominance. *Ecol. Lett.* **7**, 395–402. (doi:10.1111/j.1461-0248.2004.00591.x)

620 74. Smith MD, Knapp AK. 2003 Dominant species maintain ecosystem function with non-random

621   species loss. *Ecol. Lett.* **6**, 509–517. (doi:10.1046/j.1461-0248.2003.00454.x)

622 75. Vázquez DP, Morris WF, Jordano P. 2005 Interaction frequency as a surrogate for the total effect

623   of animal mutualists on plants. *Ecol. Lett.* **8**, 1088–1094. (doi:10.1111/j.1461-0248.2005.00810.x)

624 76. Gaston KJ. 2011 Common Ecology. *Bioscience* **61**, 354–362. (doi:10.1525/bio.2011.61.5.4)

625 77. McGill BJ, Dornelas M, Gotelli NJ, Magurran AE. 2015 Fifteen forms of biodiversity trend in the

626   anthropocene. *Trends Ecol. Evol.* **30**, 104. (doi:10.1016/j.tree.2014.11.006)

627 78. Leinster T, Cobbold CA. 2012 Measuring diversity: the importance of species similarity. *Ecology*

628   **93**, 477–489. (doi:10.1890/10-2402.1)

629 79. Dalton H. 1920 The measurement of the inequality of incomes. *Econ. J.* **30**, 348–361.

630 80. Ricotta C. 2003 Parametric scaling from species relative abundances to absolute abundances in

631   the computation of biological diversity: A first proposal using Shannon's entropy. *Acta*

632   *Biotheor.* **51**, 181–188. (doi:10.1023/A:1025142106292)

633 81. Van Strien AJ, Soldaat LL, Gregory RD. 2012 Desirable mathematical properties of indicators for

634   biodiversity change. *Ecol. Indic.* **14**, 202–208. (doi:10.1016/j.ecolind.2011.07.007)

635 82. Simpson EH. 1951 The interpetation of interactions in contingency tables. *J. R. Stat. Soc. Ser. B*

636   *Stat. Methodol.* **13**, 238–241.

639
640
641
642

Table 1. To learn how biodiversity-ecosystem function correlations are affected by different Hill diversity scaling factors, we gathered published, observational community datasets on three ecosystem functions. These were subdivided into a total of 39 community datasets, each including observations of species' identities, abundances, and functions across replicated sites.

Ecosystem function	Citation	Datasets	Replication	Max extent (km)
Rate of wild bee pollen deposition	Genung et al. 2022	Landscape array of 3 plant species	Each plant species present at 25 sites	35
Fish biomass	Lefcheck et al. 2021	32 globally distributed ecoregions (16 temperate, 16 tropical)	11-186 sites (median 59)	17 - 4,677
Above-ground carbon storage	Condit et al. 2000, Cavanaugh et al. 2014	Tree species ID and estimated biomass at four globally distributed tropical forests	50 1-Ha subplots from the 50-Ha BCI census; sets of six 1-Ha plots in three tropical regions.	1 (Condit et al.); 32-681 (Cavanaugh et al.)

668
669
Figure 1. Two hypothetical communities with (a) different species abundance distributions have
(b) different diversity profiles. At large negative ℓ values, each diversity profile converges on the
inverse proportional abundance of the one most abundant species in the assemblage (inverse
dominance). As ℓ values are more positive, each diversity profile converges on the inverse
proportional abundance of the one least abundant species in the assemblage (equal to total
abundance when the least abundant species is a singleton). Because singletons are ubiquitous in
observational data, sample Hill diversities converge on observed abundance with increasingly large,
positive values of ℓ. In the example, the red community is more even and more abundant, so its
diversity is higher compared to the blue community at both ends of the diversity spectrum. However,
the blue community has more species, and therefore is more diverse around richness (ℓ = 1, solid
vertical). Other commonly used diversities are inverse Simpson (dotted, ℓ = -1) and exponentiated
Shannon (dashed, ℓ = 0). In the diversity literature, it has been less common to explore the right side of
this spectrum (i.e., ℓ > 1)

Figure 2. Across 39 observed biodiversity-ecosystem function (BEF) correlations calculated using a wide range of diversity scaling factors, the BEF relationships with the highest r² were typically found using diversities near richness (vertical solid line; modal ℓ = 1.5). The highest BEF r² value for a community dataset was rarely found using diversities that emphasize the relative abundance of common species, including exponentiated Shannon (dashed line) and inverse Simpson (dotted line). Correlations were calculated between log(diversity) and log(ecosystem) function at a site (total above-ground carbon in tropical forest plots, rate of pollen grain deposition by wild bees, or total biomass of reef fish encountered in fixed-effort dive surveys in temperate and tropical regions).

724
725
Figure 3. Biodiversity-ecosystem function (BEF) correlations across observed communities in a
study system vary in magnitude and direction, depending on which scaling factor "ℓ" is used for
calculating species' diversity. Ecosystem services considered here are (a) rate of pollen grain
deposition on one of three flower species by wild bees; total biomass of reef fish encountered in fixed-
effort dive surveys in (b) temperate and (b) tropical global regions; and (d) total above-ground carbon
in tropical forest plots. Correlations are between logged total function at each site, and logged species
diversity at a range of ℓ values (at 0.05 intervals) emphasizing the relative abundance of common
species' (negative ℓ values) or rare species (positive ℓ values). Vertical lines identify correlations at
commonly used diversities: inverse Simpson (dotted), exponential Shannon (dashed), and richness
(solid). Colors visually distinguish different community datasets.

736
737
Figure 4. Biodiversity-ecosystem function (BEF) correlation in observational data (Figure 3) can
be explained by correlation between total abundance and diversity (first row), correlation
between mean per-capita function and diversity (second row), and interactions between these
two factors (intractable, not shown). With a few exceptions, the abundance-diversity correlations
roughly match the BEF correlation across the range of ℓ values used to calculate species diversities,
while per-capita function-diversity correlations show countervailing trends in magnitude and
direction. Compare (a, e) wild bee pollination, reef fish biomass in (b, f) temperate and (c, g) tropical
regions, and (d, h) tropical forest above-ground biomass with corresponding panels in Figure 3.
Vertical lines identify correlations at commonly used diversities: inverse Simpson (dotted),
exponential Shannon (dashed), and richness (solid). Colors visually distinguish different community
datasets.

754
Figure S1. Community datasets (line color) differed in both the shapes of the diversity profiles
(Hill diversity vs. ℓ) and the degree to which diversity profiles differed between sites. For
example, the grey-blue tree_carbon sites were all 1-Ha subplots from the contiguous BCI 50-Ha forest
plot, and diversity profiles were very similar between subplots; by contrast the yellow bee_pollination
sites (Floral visitors of *Polemonium reptans*) had variable structure with wide variety in richness (ℓ = 1,
vertical solid lines), inverse dominance (large negative ℓ), and abundance (large positive ℓ). Each curve
is the diversity profile for a single site; colors indicate a community dataset (set of sites within a region
at which a single function was measured).