# The unique synaptic circuitry of specialized olfactory glomeruli

## in *Drosophila melanogaster*

Lydia Gruber[1], Rafael Cantera[2], Markus William Pleijzier[3], Michael Steinert[4], Thomas Pertsch[4], Bill S. Hansson[1#], Jürgen Rybak [1#]

1 Max Planck Institute for Chemical Ecology, Department of Evolutionary Neuroethology, 07745 Jena, Germany

2 Instituto de Investigaciones Biológicas Clemente Estable, Departamento de Biología del Neurodesarrollo, 11600 Montevideo, Uruguay

3 Neurobiology Division, MRC Laboratory of Molecular Biology, Cambridge, CB2 0QH, England, United Kingdom

4 Institute of Applied Physics, Abbe Center of Photonics, Friedrich Schiller University Jena, Albert Einstein Strasse 15, 07745 Jena, Germany

# contributed equally

*Correspondence: Jürgen Rybak (jrybak@ice.mpg.de)

**Keywords**: olfactory circuitry, connectome*, Drosophila melanogaster*, FIB-SEM, antennal lobe, glomerulus

## ABSTRACT

In the *Drosophila* olfactory system most odorants are encoded in the antennal lobe in a combinatory way, activating several glomerular circuits. However, odorants of particular ecological role for the fly are encoded through activation of a single specialized olfactory pathway. Comparative analyses of densely reconstructed connectomes of one broadly tuned glomerulus (DL5) and one narrowly tuned glomerulus (DA2) gained detailed insight into the variations of synaptic circuitries of glomeruli with different computational tasks. Our approach combined laser-branding of glomeruli of interest with volume based focused ion beam-scanning electron microscopy (FIB-SEM) to enable precise targeting and analysis of the two glomeruli. We discovered differences in their neuronal innervation, synaptic composition and specific circuit diagrams of their major cell types: olfactory sensory neurons (OSNs), uniglomerular projection neurons (uPNs) and multiglomerular neurons (MGNs). By comparing our data with a previously mapped narrowly tuned glomerulus (VA1v), we identified putative generic features of narrowly tuned glomerular circuits, including higher density of neuronal fibers and synapses, lower degree of OSN lateralization, stronger axo-axonic connections between OSNs, dendro-dendritic connections between many uPNs, and lower degree of presynaptic inhibition on OSN axons. In addition, this work revealed that the dendrites of the single uPN in DL5 contain a substantial amount of autapses interconnecting distant regions of the dendritic tree**.** The comparative analysis of glomeruli allows to formulate synaptic motifs implemented in olfactory circuits with different computational demands.

## INTRODUCTION

Olfaction is an anatomically shallow sensory system. In mammals and invertebrates just one synapse separates the sensory periphery from the central brain (Dolan *et al.*, 2018; Liang *et al.*, 2010; Owald *et al.*, 2015; Shepherd, 2011; Su *et al.*, 2009). In the olfactory system of *Drosophila melanogaster*, the first relay station of synaptic transmission is the antennal lobe (AL), which has a circuit architecture homologous to that of the vertebrate olfactory bulb (Boeckh *et al.*, 1990; Sachse *et al.*, 2021; Shepherd *et al.*, 2021). The fly AL consists of approximately 58 spherical compartments, called glomeruli, which can be distinguished by size, shape and location (Bates *et al.*, 2020; Gao *et al.*, 2000; Grabe *et al.*, 2015; Laissue *et al.*, 1999; Vosshall *et al.*, 2000). Each glomerulus receives stereotypic input from axon terminals of olfactory sensory neurons (OSNs), which have their cell bodies and dendrites located in the antennae or maxillary palps (Benton *et al.*, 2006; de Bruyne *et al.*, 1999; de Bruyne *et al.*, 2001; Hallem *et al.*, 2004; Shanbhag *et al.*, 1999). All the OSNs innervating a given glomerulus express a typical repertoire of ligand-gated chemoreceptors (Benton *et al.*, 2006; Couto *et al.*, 2005; Fishilevich *et al.*, 2005), which represent a wide range of specifications, binding either a single, few, or many distinct chemicals (Hallem *et al.*, 2006; Hallem *et al.*, 2004; Knaden *et al.*, 2012; Münch *et al.*, 2016; Seki *et al.*, 2017; Wicher *et al.*, 2021).

Most OSNs project bilaterally to the corresponding glomeruli in the left and right AL (Gaudry *et al.*, 2013; Tobin *et al.*, 2017). In the AL, OSNs convey odor signals to excitatory uniglomerular projection neurons (uPNs), which branch only within a single glomerulus, or to inhibitory multiglomerular PNs (mPNs) and inhibitory or excitatory local interneurons (LNs) (Ai *et al.*, 2013; Bates *et al.*, 2020; Cuntz *et al.*, 2007; Kazama *et al.*, 2008; Kazama *et al.*, 2009; Kreher *et al.*, 2008; Masse *et al.*, 2009; Ng *et al.*, 2002; Tanaka *et al.*, 2012; Wilson, 2013). LNs innervate each several glomeruli and are the key modulatory neurons in the AL (Chou *et al.*, 2010; Seki *et al.*, 2010). The highly converging OSNs-to-PN signal transmission (Chen *et al.*, 2005; Jeanne *et al.*, 2015; Masse *et al.*, 2009) is lateralized, activating ipsilateral uPNs more strongly than contralateral ones (Agarwal *et al.*, 2011; Gaudry *et al.*, 2013; Tobin *et al.*, 2017). From the AL, uPNs and mPNs relay processed signal information to higher brain centers (Bates *et al.*, 2020; Fiala, 2007; Galizia, 2014; Guven-Ozkan *et al.*, 2014; Jefferis *et al.*, 2007; Keene *et al.*, 2007; Norgate *et al.*, 2006; Strutz *et al.*, 2014).

The stereotypic activity pattern elicited by distinct odorants encodes the odor space, represented in a so-called odotopic map of the AL according to the glomerular activation by distinct chemical classes. (Couto *et al.*, 2005; Grabe *et al.*, 2018; Grabe *et al.*, 2015; Knaden *et al.*, 2014; Laissue *et al.*, 2008). Some odorants induce a fixed innate behavior (aversion or attraction), activating characteristically specific glomeruli (Gao *et al.*, 2015; Grabe *et al.*, 2018; Knaden *et al.*, 2014; Knaden *et al.*, 2012; Semmelhack *et al.*, 2009). The encoding of hedonic valence already at the level of the AL is important for a fast odor coding. Most odorants are encoded in a combinatorial manner in the fly AL by activating multiple OSNs types expressing broadly tuned

receptors and their glomerular circuits, including broadly tuned uPNs (de Bruyne *et al.*, 2001; Galizia, 2014; Masse *et al.*, 2009; Sachse *et al.*, 2016; Seki *et al.*, 2017; Silbering *et al.*, 2007; Silbering *et al.*, 2008; Szyszka *et al.*, 2015). Certain chemoreceptors and their downstream glomerular circuits, however, have evolved a very high specificity and sensitivity to single or very few chemicals (Andersson *et al.*, 2015; Haverkamp *et al.*, 2018; Keesey *et al.*, 2021). These narrowly tuned glomerular circuits often belong to dedicated olfactory pathways, called "labeled lines", which process information regarding single odorants of particular importance for reproduction and survival (Datta *et al.*, 2008; Dweck *et al.*, 2015; Gao *et al.*, 2015; Kurtovic *et al.*, 2007; Stensmyr *et al.*, 2012). An extreme example is the DA2 glomerulus, which responds exclusively to geosmin, an ecologically relevant chemical that alerts flies to the presence of harmful microbes, causing the fly to avoid laying eggs at these locations (Stensmyr et al., 2012). This dedicated olfactory pathway and its receptor sequence is conserved throughout evolution (Keesey *et al.*, 2021; Keesey *et al.*, 2019). Another example is glomerulus VA1v, which responds to methyl laurate, a pheromone that induces a strongly attractive response in female flies leading to aggregation behavior (Dweck *et al.*, 2015). DL5, on the other hand, is an example of a broadly tuned glomerulus, innervated by OSNs activated by several odorants, like E2-hexenal and benzaldehyde (Knaden *et al.*, 2012; Mohamed *et al.*, 2019b; Münch *et al.*, 2016; Seki *et al.*, 2017). This functional diversity suggests differences in neuronal composition and synaptic connectivity between broadly and narrowly tuned glomeruli.

A survey of neuronal composition across glomeruli revealed great variation in the numbers of the different types of neurons innervating narrowly and broadly tuned glomeruli (Grabe *et al.*, 2016). In general, narrowly tuned glomeruli are innervated by more uPNs and fewer LNs compared to more broadly tuned glomeruli (Chou *et al.*, 2010; Grabe *et al.*, 2016). In addition, narrowly tuned OSNs receive less global LN inhibition than broadly tuned ones (Grabe *et al.*, 2020; Hong *et al.*, 2015; Schlegel *et al.*, 2021). For example, in female flies, the narrowly tuned glomerulus DA2 contains dendrites of 6-8 uPNs, whereas the broadly tuned glomerulus DL5 houses only 1 or 2 uPNs and has a higher number of innervating LNs. Interestingly, both glomeruli are innervated by the same number of OSNs (Grabe *et al.*, 2016).

Little is known, however, about the microarchitecture of the synaptic circuitry in distinct glomeruli and, in particular, about ultrastructural differences between narrowly vs. broadly tuned glomerular circuits. Electron microscopy (EM) allows volume imaging with dense reconstruction of fine neurite branches and synapses in brain tissue at nanometer resolution, necessary to map synapses (Briggman *et al.*, 2006; Cardona *et al.*, 2009; Helmstaedter, 2013; Meinertzhagen, 2018; Rybak, 2013). The first ultrastructural insights into the synaptic connectivity of *Drosophila* olfactory glomeruli were obtained by studies based on serial section transmission EM (ssTEM) (Rybak, 2016; Tobin *et al.*, 2017). Rybak *et al*. (2016) showed that all three basic classes of AL neurons make synapses with each other, while Tobin *et al*. (2017) revealed that the differences in number of innervating uPNs between the left and right DM6 glomeruli are compensated by differences in synaptic strength. With focused ion beam-scanning electron microscopy (FIB-SEM; (Knott *et al.*,

2008)) a complete reconstruction of all neurons in the narrowly tuned, pheromone processing
glomerulus VA1v was obtained (Horne *et al.*, 2018). Recent technological innovations in ssTEM,
FIB-SEM and automated neuron reconstruction have made connectome datasets of the adult
*Drosophila* central nervous system available (Li, P. H. *et al.*, 2020; Saalfeld *et al.*, 2009; Scheffer *et*
*al.*, 2020; Zheng *et al.*, 2018) and provided complete circuit descriptions of several brain centres
(Auer *et al.*, 2020; Bates *et al.*, 2020; Coates *et al.*, 2020; Dolan *et al.*, 2019; Felsenberg *et al.*, 2018;
Hulse *et al.*, 2021; Huoviala *et al.*, 2020; Li, F. *et al.*, 2020; Marin *et al.*, 2020; Otto *et al.*, 2020;
Schlegel *et al.*, 2021).

In an attempt to find answers to how highly specialized olfactory glomerular circuits of
dedicated olfactory pathways differ in their signal integration from broadly tuned glomerular
circuits, we compared the microarchitecture and synaptic circuitry of a narrowly and a broadly
tuned glomerulus (DA2 and DL5). By using a correlative workflow combining transgenic markers
with FIB-SEM, in order to identify these glomeruli, we reconstructed OSNs, uPNs and
multiglomerular neurons (MGNs), mapped all associated synapses and compared the circuit
organization of both glomeruli.

**RESULTS**

**Volume-based electron microscopy of two different olfactory glomeruli**

To compare the synaptic circuitries of two olfactory glomeruli known to belong to either
narrowly or broadly tuned glomerular types in *Drosophila melanogaster*, we mapped all synapses
of glomeruli DA2 (right AL) and DL5 (left AL) in a single female fly (**Figure 1A-B**) with the aid of FIB-
SEM. A partial reconstruction of a second DA2 in another fly was used to measure neuronal volume
(see Methods). The reconstructions were based on high resolution (4x4x20 nm) datasets (**Figure 1**;
**Figure 1 – video 1),** thus allowing reconstruction of the finest neuronal branches (~20 nm diameter;
**Figure 1C-D**) as well as mapping chemical synapses (example in **Figure 1E**) in the two volumes of
interest (VOI). To restrict the imaging volume to the target VOIs, we employed a correlative
approach for the first time for a *Drosophila* EM volume reconstruction. Glomeruli DA2 and DL5
were identified by their size, shape and location in brains of transgenic flies (*Orco-GAL4; UAS-*
*GCaMP6s*) using the glomerular map of (Grabe *et al.*, 2015). The flies expressed the green
fluorescent protein GCaMP6 coupled with calmodulin and M13 (a peptide sequence from myosin
light-chain kinase; **Figure 1A-B**). Subsequently, the identified glomeruli were marked by laser
branding using a two-photon laser (Bishop *et al.*, 2011). These fiducial marks were apparent under
both light (Figure 1A-B) and electron microscopy **(Figure 1C-D)** and facilitated the delimitation of
the VOIs during FIB-SEM scanning. We produced two complete FIB-SEM datasets: one for
glomerulus DA2 and one for DL5 (pure imaging time for both glomeruli: ~60 h) and a partial dataset
for DA2 in a second fly.

**Skeleton based neuron reconstruction and synapse identification**

We reconstructed all neurons within the two VOIs (example neuron: **Figure 1F**) and mapped
all their synaptic connections using an iterative skeleton-based reconstruction approach, similar to
previously reported procedures (Berck *et al.*, 2016; Schneider-Mizell *et al.*, 2016; Zheng *et al.*,
2017) with the aid of the web-based neuron reconstruction software CATMAID
(http://www.catmaid.org; RRID:SCR_006278; (Cardona *et al.*, 2009; Schneider-Mizell *et al.*, 2016);
**Figure 1 – video 1**). Synapses were identified by their presynaptic transmitter release site, which
in *Drosophila* is composed of a presynaptic density called a T-bar, surrounded by synaptic vesicles
and apposed postsynaptic elements (**Figure 1E**), as previously described (Fröhlich, 1985; Huang *et*
*al.*, 2018; Li, P. H. *et al.*, 2020; Rybak *et al.*, 2016; Trujillo-Cenoz, 1969). All synapses observed in
our FIB-SEM data sets were polyadic, i.e. each presynaptic site connected to multiple postsynaptic
sites (See example in **Figure 1E**), a feature of insect brain synapses (Hartenstein, 2016; Malun *et*
*al.*, 1993; Meinertzhagen *et al.*, 1991; Prokop *et al.*, 2006; Rybak *et al.*, 2016). Some synapses had
up to 16 postsynaptic sites (**Figure 2 – figure supplement 1B**), i.e. one T-bar and 16 single synaptic
profiles (i.e. sixteen 1:1 single output-input connections). Short neuronal fragments (<10 μm),

which could not be connected to any neuronal fiber were designated as "orphans". These
fragments represented 4% of the total length of all traced neuronal fibers in DA2 and 6% in DL5
and contained about ~12% of all synaptic contacts in both glomeruli.
168
**Glomerular neurons: classification, description and inventory**

Previous descriptions of the ultrastructural characteristics of the AL in *Drosophila* helped to
classify AL neurons into 3 main classes (**Figure 2A**) Olfactory sensory neurons (OSNs),
uniglomerular projection neurons (uPNs) and multiglomerular neurons (MGNs; cells that
interconnect multiple glomeruli). MGNs are further subdivided into multiglomerular projection
neurons (mPNs) and local interneurons (LNs) (Berck *et al.*, 2016; Gruber *et al.*, 2018; Horne *et al.*,
2018; Li, P. H. *et al.*, 2020; Rybak *et al.*, 2016; Schlegel *et al.*, 2021; Zheng *et al.*, 2017). Most of the
neuronal profiles within the MGN neuron class belong probably to inhibitory local neurons, as this
cell type is the most numerous and broadly arborizing of the multiglomerular cell types in the
antennal lobe (Chou *et al.*, 2010; Lin *et al.*, 2012). In addition, we observed a few neuronal fibers
with an electron-dense and vesicle-rich cytosol, which we interpreted to be either peptidergic
neurons (Eckstein *et al.*, 2020; Nässel *et al.*, 2006) or the contralaterally projecting, serotonin-
immunoreactive deutocerebral (CSD) neuron, (Coates *et al.*, 2020; Dacks *et al.*, 2006; Eckstein *et
al.*, 2020; Goyal *et al.*, 2013; Zheng *et al.*, 2017). Except for these neuronal fibers containing
abundant electron-dense vesicles, all other neuronal fibers were assigned to either OSNs, uPNs or
MGNs based on their morphology (**Figure 2A, B;** see Methods).
OSNs formed large, elongated synaptic boutons (**Figure 2A**), had the largest volume/length
ratio of all three neuron classes (**Figure 2 – figure supplement 1A**) and displayed the lowest degree
of branching intensity of all neurons in both glomeruli (**Figure 2B**). In agreement with what had
been observed in other glomeruli (Rybak *et al.*, 2016), the majority of output synapses made by
OSN terminals were triads (1:3) and tetrads (1:4). The T-bars of OSN synapses exhibited a large
variation in size and some of them were large enough to accommodate 16 postsynaptic contacts
(**Figure 2 – figure supplement 1B**). The frequency of large T-bars was much higher in OSNs than in
other neuron classes with an average polyadicity (average number of postsynaptic sites at each T-
bar) of 6 (1:6; (**Table 1,** row 14). As OSNs had the greatest T-bar and output density along their
axons (**Table 1**, row 10-11) they also displayed the largest synaptic ratios (both for the T-bars/input
sites and output sites/input sites) of all neuron classes (**Table 1,** row 12-13), which was in line with
previous observations (Rybak *et al.*, 2016).
The uPNs exhibited the highest degree of branching intensity of the three neuron classes in
both glomeruli (**Figure 2A-B**). They showed numerous very fine apical branches that frequently
connected multiple times via spines to the same presynaptic site, leading to an entangled 3D shape
typical for this neuron class (**Figure 2A**) (Rybak *et al.*, 2016; Schlegel *et al.*, 2021; Tobin *et al.*, 2017).
uPNs had the smallest volume/length ratio of all neuron classes (for the DA2: **Figure 2 - supplement**

**1A**). In addition to having many fine branches, uPN dendrites also had enlarged regions with almost
no cytosol that were packed with large mitochondrial profiles extending over considerable
distances. These enlarged profiles showed a larger degree of mitochondria fission (dividing and
segregating mitochondrion organelles; personal observation) than the other neuron classes with
rather round and compact mitochondria (**Figure 2A**; FIB-SEM image; see data availability). Seven
uPNs were found in DA2, confirming light microscopy findings (Grabe *et al.*, 2016). Two of them
(PN#1, PN#2; see data availability) branched broadly and innervated the full glomerulus, and
received more synaptic input than the other 5 uPNs (PN#3-#7; **see Table S3**), which branched
exclusively in sub-regions of the glomerulus, with partial overlap. In addition to abundant clear
small vesicles (~20 nm in diameter) (Bates *et al.*, 2020; Strutz *et al.*, 2014; Yasuyama *et al.*, 2003),
uPN dendrites also displayed small electron-dense vesicles, as previously reported for PN axon
terminals in the mushroom body calyx (Butcher *et al.*, 2012; Yang *et al.*, 2022). These electron-
dense vesicles are packed with different types of neuropeptides that act as neuromodulators or
co-transmitters (Croset *et al.*, 2018; Eckstein *et al.*, 2020; Gondré-Lewis *et al.*, 2012; Li *et al.*, 2017).
In both glomeruli, uPNs had the highest neuronal synaptic input density and the lowest T-bar and
output density of the three neuron classes (**Table 1,** row 9-11; DA2 and DL5 differences: see next
section). The synaptic ratios (T-bars/input sites and output sites/input sites) were much lower for
uPNs than for the other neuron classes (**Table 1,** row 12-13). The majority of uPN dendritic output
synapses (feedback synapses) were tetrads in both glomeruli, with an average polyadicity of
around 5 (lower than in OSNs; (**Figure 2 – supplement 1; Table 1,** row 14).

The majority of the neuronal fibers in both glomeruli belonged to MGNs (**Figure 2A**). MGNs
exhibited variable morphology and ultrastructure, as expected, but shared also some
ultrastructural features. Their synaptic boutons were formed by thin fibers, thus the volume/length
ratio of MGNs was lower than that of OSNs but greater than that of uPNs (**Figure 2 – figure
supplement 1A**). A similar relationship was found for the number of output sites and the T-bar
density along MGN fibers, which were smaller than in OSNs but larger than in uPNs (**Table 1,** row
10-11). In contrast, branching intensity in MGNs was larger than in OSNs but smaller than in uPNs
(**Figure 2B**). The synaptic ratio of output-to-input sites was around one (**Table 1**, row 12-13). MGNs
had the lowest polyadicity (~3) of the three neuron classes (**Table 1**, row 14) and their synapses
were mainly triads (**Figure 2 – supplement 1D**). Interestingly, besides the abundant clear small
vesicles (~20 nm in diameter), some MGNs had small electron-dense vesicles, most likely housing
the neuropeptide sNPF (Nässel *et al.*, 2008).

**DA2 is more densely innervated and has a higher synapse density than DL5**

In our FIB-SEM datasets the volume of glomerulus DA2 was 45% smaller than that of
glomerulus DL5 (1500 µm$^3$ vs. 2640 µm$^3$), which is in agreement with measurements based on light
microscopy (DA2 = 1600 µm$^3$, DL5 = 2900 µm$^3$ (Grabe *et al.*, 2016). We also confirmed that a similar

number of OSNs (44-46 OSNs) innervated both glomeruli (**Figure 2C**), and that each glomerulus
received OSN innervation from both the ipsilateral and contralateral antennae (Grabe *et al.*, 2016;
Vosshall *et al.*, 2000). Also in agreement with (Grabe *et al.*, 2016), the DA2 glomerulus was
innervated by 7 uPNs whereas DL5 had a single uPN (**Figure 2C**). MGN cell numbers could not be
counted in our study due to their multiglomerular morphology, which also prevented us from
tracing MGN fibers to their soma due to our partial volume acquisition (see Methods).

To investigate differences between DA2 and DL5 we turned our attention to their glomerular
innervation and synaptic composition. We measured the total length (sum in μm) of all neuronal
fibers of each neuron class within the DA2 and DL5 (**Figure 2C; Table 1**, row 1). In addition, we
counted all T-bars and their output sites (1:1 synaptic contacts) as well as all postsynaptic sites
(input sites) for all neuron fibers together and for each neuron class individually (**Table 1, row 2-
4**). We counted in total ~ 14 000 synaptic contacts and 2648 T-bars in DA2 and ~ 17 000 contacts
and 3387 T-bars in DL5 (**Figure 2C, Table 1, row 4**). Most of these synapses were triads and tetrads
(**Figure 2 – figure supplement 1B-D**). In order to compare DA2 and DL5 we normalized neuronal
length and synaptic numbers to glomerular volume. We then analyzed (**1**) the innervation density,
i.e., the length of neuronal fibers per glomerular volume ($μm/μm^3$) and (**2**) the glomerular synaptic
density (T-bar #, output site or input site $#/μm^3$). Data are reported in total for all neuronal fibers
of each neuron class (**Table 1, row 5-8**) and as an average for neuronal fibers of the respective
neuron class **(Figure 3)**. In addition, we compared (**3**) the average polyadicity for each neuron class
**(Figure 3)** and **(4)** the average neuronal synaptic density (T-bar, output and input site density along
each neuronal fiber) (#/μm) (**Figure 3 – figure supplement 1B**).

We observed that the average neuron innervation density of OSNs was 20% higher in DA2
than in DL5 (**Figure 3A**)**, Table S1**). Also the glomerular synaptic density of input sites, output sites
and T-bars along OSNs was significantly higher in DA2 than in DL5 (**Figure 3A**). OSNs in DA2 formed
therefore more input sites, and much more T-bars and output sites per glomerular volume than in
DL5 (**Table 1**, row 7-8; relative differences: **Table S1**). In contrast, the density of input sites
distributed along the length of OSN fibers was similar in DA2 and DL5, whereas T-bar and output
site density along the OSN axons was significantly higher in DA2 (**Figure 3 – figure supplement 1A**).
267

We then asked if the DA2 glomerulus, due to its higher number of uPNs, also had a higher
uPN innervation density and synaptic density of its postsynaptic sites and/or presynaptic sites
compared to the DL5 glomerulus, which contains a single uPN. In the DA2, the fibers of the 7 uPNs
had almost the same total length as the fibers of the single uPN in the more voluminous DL5 (4652
μm in DA2 vs. 5015 μm in DL5; **Table 1**, row 1). The DA2 uPNs had in addition a similar total number
of input sites than the single uPN in DL5 (3887 vs. 3955; **Table 1**, row 2). As such, in DA2 the total
innervation density of its 7 uPNs was higher as compared to the innervation density of the single
uPN in DL5 (**Table 1**, row 5), even though the average innervation density of DA2-uPNs was lower
(**Figure 3B**). The total glomerular input density of all uPNs was higher in DA2 as compared to DL5

(**Table 1**, row 6). On the other hand, the total glomerular synaptic density of the T-bars and output
sites was similar in DA2 and DL5 (**Table 1**, rows 7-8). In line with these results, the neuronal density
of T-bars and output sites was less in the DA2 uPNs compared to the DL5 single uPN, whereas the
neuronal density of input sites was similar (**Figure 3 – figure supplement 1B; Table 1**, row 9-10).
This caused almost twice as high synaptic ratios (T-bars-to-inputs and outputs-to-inputs) in the DL5
uPN relative to DA2 uPNs (**Table 1**; row 12-13).

We then hypothesized that DA2 will have a lower innervation density of MGNs (mainly LNs)
than DL5 as it had been reported that DL5 is innervated by fewer LNs (Chou *et al.*, 2010; Grabe *et*
*al.*, 2016). However, we observed the opposite: the innervation density of MGNs was significantly
higher in DA2 than in DL5 (**Figure 3C**), with slightly higher total innervation density (**Table 1,** row
5**).** Interestingly, only the glomerular input density was significantly higher for DA2 MGNs
compared to that found in DL5, not the glomerular synaptic density of output sites or of the T-bars
(**Figure 3C**). However, the total glomerular synaptic density of input sites, output sites and T-bars
were still higher in DA2 than in DL5 **(Table 1,** rows 6-8)**.** Synaptic densities along the MGN fibers
were similar in DA2 and DL5 (**Figure 3 – supplement 1**).

In summary, the DA2 glomerulus is more densely innervated than DL5 and has a more
densely packed neuropil with more synaptic contacts relative to the DL5. The DA2 has a
significantly higher innervation density and higher density of T-bars, output and input sites per
volume (**Figure 3D**, **Table 1,** row 5-8)**.** The degree of synapse polyadicity is also significantly higher
in DA2 than in DL5 (**Figure 3D, Table 1,** row 14) due to a shift to higher polyadicity among OSN
(**Figure 3A**) and MGN synapses (**Figure 3C**). OSNs show the strongest shift in polyadicity, with
tetrads being the most abundant synapse type in DA2 whereas triads are the most abundant in
DL5 OSNs (**Figure 2 – figure supplement 1B).**

**Lateralization of OSN glomerular connectivity**

In *Drosophila melanogaster*, the majority of olfactory glomeruli receive bilateral OSN input
(Silbering *et al.*, 2011; Stocker *et al.*, 1990; Stocker *et al.*, 1983; Vosshall *et al.*, 2000) see scheme
in **Figure 4A**). Recent studies have shown that ipsi- and contralateral OSNs are asymmetric in their
synaptic connectivity to other neurons in the majority of the glomeruli (Schlegel *et al.*, 2021; Tobin
*et al.*, 2017) and that ipsi- and contralateral OSNs activate uPNs in an asymmetric way (Gaudry *et*
*al.*, 2013; Tobin *et al.*, 2017). However, not all glomeruli appear to have the same degree of
lateralized OSN connectivity (Schlegel *et al.*, 2021). At least for one narrowly tuned glomerulus
(DA1), there is functional evidence that in female flies its uPNs are evenly activated by either ipsi-
or contralateral antennal stimulation (Agarwal *et al.*, 2011). We hypothesized that this lack of
lateralization could be a feature of other narrowly tuned glomeruli.

Ipsi- and contralateral OSNs in DA2 and DL5 were identified based on the location and
trajectory of their axons (**Figure 4B**). In both glomeruli, ipsilateral OSN terminals were longer than

their contralateral counterparts within the VOI, while polyadicity was stronger in contralateral
axons. Synaptic density was not consistently higher or lower in ipsilateral OSNs compared to
contralateral ones in DA2 and DL5 (**Figure 4 – figure supplement 1**).

We observed that the synaptic output of ipsi- vs. contralateral OSNs was asymmetric, with
significant differences in the ipsi- and contralateral OSN output to either uPNs, OSNs or MGNs
(**Figure 4C**, DA2 and DL5). In agreement with previous observations in other glomeruli (Schlegel *et*
*al.*, 2021), the output fraction to uPNs and OSNs was greater in ipsilateral OSNs than in
contralateral ones (**Figure 4C**, DA2 and DL5). Vice versa, the OSN output to MGNs was greater in
the contralateral glomerulus than in the ipsilateral side (**Figure 4C**, DA2 and DL5). However, the
differences between the medians and means were smaller in DA2 than in DL5 (**Figure 4C**;
differences between means: see data availability).

Our finding of less lateralized connections in the DA2 (**Figure 4C**, DA2 and DL5) was also
observed in another narrowly tuned glomerulus (VA1v; Dweck et al., 2015) for which connectome
data is available (Horne *et al.*, 2018). In VA1v, the OSN output to uPNs and MGNs was significantly
asymmetric in the same manner as in DA2 and DL5, i.e. with greater ipsilateral OSN output fractions
to uPNs and OSNs and greater contralateral OSN output fraction to MGNs (**Figure 4C**). Asymmetry
in the VA1v OSN output fractions was even less distinct than in DA2 (regarding both the difference
between the median and the mean; **Figure 4C** and data availability). In VA1v, the OSN output
fraction to OSNs was similar in ipsi- and contralateral OSNs (**Figure 4C**). In addition, the OSN input,
from either sister OSNs or MGNs, was asymmetric in DL5 but not in the narrowly tuned glomeruli
(**Figure 4D**). The inputs from uPNs to ipsi- or contralateral OSNs were not compared due to their
low numbers.

In summary, our data add to the knowledge of lateralized connectivity within olfactory
glomeruli and supports the hypothesis that narrowly tuned glomeruli have a lower degree of
lateralization of OSN connectivity compared to broadly tuned glomeruli.

**Glomeruli DA2 and DL5 differ in several features of their circuitry**

Next, we asked whether the synaptic circuitries of DA2 and DL5 differ from each other. We
counted each synaptic contact (**Table S2 and S3**) and categorized the distinct connection motifs
according to the neuron class to which the output and input neuron belonged (**Figure 5A; Table**
**S2).** Each connection motif (for example OSN>uPN, i.e., the OSN-to-uPN feedforward connection)
was then assessed for its relative synaptic strength, i.e. how many synaptic contacts of this
particular connection motif were found compared to the total number of synaptic contacts within
the respective circuitry (**Figure 5A-D**; see Methods).

We found that neurons from each class made synaptic contacts with each other in DA2 and
DL5, as previously reported for other glomeruli (Berck *et al.*, 2016; Horne *et al.*, 2018; Rybak *et al.*,
2016; Schlegel *et al.*, 2021; Tobin *et al.*, 2017). In both DA2 and DL5, OSNs provided the strongest

relative synaptic output, i.e. 49% of all synaptic connections in DA2 and 43% in DL5 were formed
by OSNs (**Figure 5B-C**). Thus, even though DA2 and DL5 had similar numbers of OSNs (44 and 46,
respectively), those in DA2 provided a stronger circuit output (14% stronger; **Table S2**) than those
in DL5 (**Figure 5B-C**). In both glomeruli the main OSN output partners were MGNs and uPNs, i.e.
27% of all circuitry connections in DA2 and 24% in DL5 were OSN>MGN connections and 20% in
DA2 and 18% in DL5 were OSN>uPN connections (**Figure 5B-C**). In DA2, interestingly, each of the 7
uPNs received input from almost all OSNs and so could maintain a high degree of convergent signal
transmission (**Table S3**). In contrast, OSNs received the lowest relative input of all neuron classes
in DA2 and DL5 (7% and 8% respectively; **Figure 5B-C**). In line with previous observations in other
glomeruli (Horne *et al.*, 2018; Schlegel *et al.*, 2021), OSNs also made abundant axo-axonic synapses
with sister OSNs (2.6% in DA2 and 1.5% in DL5; **Figure 5B-C**). Thus, the relative synaptic strength
of the OSN>OSN connection was 70% stronger in DA2 than in DL5 (**Figure 5B-C; Table S2**).
The uPNs in both glomeruli had the weakest relative output of all neuron classes within their
circuitry, and this was even weaker (38%) in DA2 (**Figure 5B-C; Table S2**). In contrast, the relative
synaptic input onto uPNs was greater in DA2 than in DL5 (33% vs. 28%, respectively; **Figure 5B-C;**
16% stronger in DA2; **Table S2**), which is in line with our finding that in DA2, the uPNs provide more
input sites per unit of glomerular volume than in the DL5 (**Figure 3B-C**). In both glomeruli, the
feedback connections from uPNs (depicted in **Figure 5A**), were almost exclusively directed towards
MGNs, as previously reported for the broadly tuned DM6 and the narrowly tuned glomerulus VA1v
(Horne *et al.*, 2018; Tobin *et al.*, 2017). However, the relative synaptic strength of the uPN>MGN
connection was 40% weaker in DA2 than in DL5 (uPN>MGN: 10% in DA2 and 17% in DL5). Only a
few cases of uPN>OSN synaptic connections were observed (a total of 16 in DA2 and 26 in DL5)
representing a synaptic strength of 0.1% in DA2 and 0.2% in DL5 (**Table S2**). Finally, uPNs in DA2
also made 71 reciprocal synaptic connections (representing a synaptic strength of 0.6%; **Table S2**;
**Figure 5B**), consistent with electrophysiological evidence for reciprocal synaptic interactions
between sister uPNs (Kazama *et al.*, 2009). The single uPN of the DL5 had 54 dendro-dendritic
synapses (representing 0.4% of all DL5 synaptic contacts; **Figure 5C**), which were exclusively
autapses, i.e. synapses formed by a neuron onto itself. Dendritic uPN autapses exist also in DA2-
uPNs, but they were few: we observed only 14 autaptic uPN-uPN connections in DA2, which were
mainly located at the two longest uPN dendrites (for further analysis of autapses see next section).
MGNs received the strongest input in both glomeruli (60% of the total input in DA2 and 64%
in DL5; **Figure 5B-C**). This is in line with the observation that MGNs provided the majority of all
traced neuronal fibers in each glomerulus and had the highest innervation density of all neuron
classes; **Table 1**). The relative output strength of MGNs was similar in both glomerular circuits
(~40% of the total output in each glomerulus; **Figure 5B-C**). MGNs made many reciprocal synapses
to each other, accounting for 23% of all synapses in both glomeruli (**Figure 5B-C**). The relative
synaptic strength between MGN>uPN was stronger in DA2 (12%) than DL5 (10%) (**Figure 5B-C;**

**Table S2**). The MGN>OSN feedback connection was relatively weak in both glomeruli (5% in DA2
vs. 6% in DL5; **Figure 5B-C**) but weaker (25%) in DA2 than in DL5 (**Table S2**).

We then looked at the fractional output and input of each neuron class (Figure 5E', E''). In
both glomeruli OSNs had a similar proportion of their synaptic output onto uPNs (40%-41%), onto
MGNs (55% in both) and onto sister OSNs (4%-5%) (**Figure 5E'**). From the uPNs perspective, over
93%-96% of their recurrent synaptic output was directed to MGNs in both DA2 and DL5, and few
synapses were directed onto OSNs (~1% of the uPN output; **Figure 5E'**). The uPN>uPN output
fraction of the 7 uPNs in DA2 (reciprocal synapses) was twice the uPN output fraction (autaptic) of
the single uPN dendrite in DL5 (6% vs. 3%; **Figure 5E'**). MGNs formed synaptic output mainly to
other MGNs (58%-59% of the total MGN output in DA2 and DL5). Among MGNs we found also rare
cases of autapses. The MGN>uPN output fraction was greater in DA2 (30%) than in DL5 (25%),
whereas the MGN>OSN output fraction was smaller in DA2 (12%) than in DL5 (16%; **Figure 5E'**).

Turning to the input fractions of each neuron class, we found that in both glomeruli, OSNs
received most of their input from MGNs (>50%). In DA2 the input fraction onto OSNs (MGN>OSN)
was smaller than in DL5 (63% vs. 78%; **Figure 5E''**). In contrast, the OSN input fraction from sister
OSNs was greater in DA2 (35% vs. 20%; **Figure 5E''**). In both glomeruli, the OSNs received only weak
uPN input (2%) (**Figure 5E''**). The input fractions onto the 7 uPNs, formed by uPNs, MGNs and OSNs,
in the DA2 and the single uPN in DL5 were similar (**Figure 5E''**). Most uPN input was delivered by
OSNs (~62% in both glomeruli) and less from MGNs (~36%). The uPN input fraction from other
uPNs in DA2 or the autaptic input from the single uPN in DL5 was small (2%; **Figure 5E''**). In DA2
the MGNs received a smaller fraction of uPN feedback input than in DL5 (17% vs. 26%; **Figure 5E''**)
but a greater OSN input fraction (45% vs. 38%; **Figure 5E''**). The fraction of MGN>MGN input was
similar in both glomeruli.

To further explore whether the differences in circuitry between DA2 and DL5 reported here
might represent features characteristic of narrowly tuned glomeruli, we analyzed connectome
data from another narrowly tuned glomerulus (VA1v; (Horne *et al.*, 2018). We calculated the
relative synaptic strength between OSNs (n=107), uPNs (n=5) and MGNs (n=74) in the VA1v (**Figure
5D; Table S2**). We found that the two narrowly tuned glomeruli shared five circuit features that
were different from the broadly tuned glomerulus DL5: (**1**), OSNs in VA1v, as reported above for
DA2, displayed a stronger relative feedforward output to uPNs (22%) and to MGNs (32%) (**Figure
5D**). The uPNs and MGNs in VA1v, received a larger fraction of OSN input than in DL5 (**Figure 5E''**).
(**2**), the OSN>OSN synaptic output was four times stronger (6%) than in DL5 (1.5%; **Figure 5B-D,
Table S1**). This was also reflected in the OSN output fraction to sister OSNs (10%), which in VA1v
was more than twice that of DL5 (4%; **Figure 5E'**) and in the much greater OSN input fraction (38%)
to OSNs in the VA1v than in DL5 (20%; **Figure 5E''**). (**3**), in the VA1v the uPN>uPN relative synaptic
output was more than twice that of DL5 (1% vs. 0.4% in DL5; **Figure 5D**), which is in accordance
with a much greater uPN output fraction to uPNs (14%) in VA1v than in DL5 (3%) (**Figure 5E'**). (**4**),
as observed before in DA2, VA1v uPNs had fewer feedback synapses onto MGNs than in DL5

(relative synaptic strength of uPN>MGN connection: 6% vs. 17%; **Figure 5C-D**), also reflected in a
smaller output fraction from uPNs to MGNs in VA1v than in DL5 (81% vs. 96%; **Figure 5E'**). In
agreement, the MGN input fraction from uPNs in VA1v was much smaller than in DL5 (10% vs. 26%;
**Figure 5E''**). (**5**), OSNs in VA1v received a smaller MGN input fraction than DL5 OSNs (60% vs. 78%;
**Figure 5E''**).

Besides relative differences (stronger or weaker) in DA2 and VA1v connection motifs
compared to DL5, two connection motifs were stronger in DA2 and DL5 than in VA1v: **(1)** the
MGN>uPN connection showed a synaptic strength of 12% and 10% in DA2 and DL5 vs. 8% in VA1v
(**Figure 5B-D, Table S2**). In agreement with this, the MGN output fraction to uPNs (**Figure 5E'**, MGN
output) and the MGN input fraction in uPNs was greater in DA2 and DL5 than in VA1v (**Figure 5E''**,
uPN input). **(2)**, the relative synaptic strength in MGN>MGN motifs was similar between DA2 and
DL5 (23%; **Figure 5B-C**), but weaker in VA1v (17%; **Figure 5D, Table S2**). This was also reflected in
a smaller MGN output and input fraction from or to MGNs (**Figure 5E' and E''**).

In summary, the two narrowly tuned glomerular circuits studied here shared five circuit
features when compared with the broadly tuned glomerular circuit (all glomerular circuit features
in DA2, DL5 and VA1v are shown in **Figure 6A**). These features were (1) a stronger OSN>uPN and
OSN>MGN connection, (2) a much stronger axo-axonic communication between sister OSNs, (3) a
stronger dendro-dendritic connection between uPN dendrites, (4) less feedback from uPNs to
MGNs and (5) less feedback from MGNs to OSNs (**Figure 6B**).

**Autapses in the large DL5 uPN connect distant regions of its dendritic tree**

Autapses (synapses made by a neuron upon itself) have seldomly been reported in the
*Drosophila* central nervous system (Horne *et al.*, 2018; Takemura *et al.*, 2015). In the DA2
glomerulus we found few autapses in uPNs and MGNs (**Figure 5C; Figure 7A**). In the dendritic tree
of the single DL5 uPN, on the other hand, three observers registered 54 autaptic connections
independently (see Methods). This represents 3% of the output connections of this neuron and
0.4% of all synaptic contacts in the whole glomerulus (**Figure 7A**; **Figure 5C; E'**). We found that
these autapses were not distributed evenly along the dendritic tree of the DL5 uPN. Some dendritic
branches received several autaptic inputs, whereas other had no autaptic input (**Figure 7A**) and we
hypothesized that these autapses could connect distant parts of this very large dendritic tree. We
thus analyzed the exact location and distribution of their presynaptic and postsynaptic sites (**Figure**
**7A**) We discovered a difference in the distribution of the pre- and postsynaptic elements of DL5
autapses. Whilst their presynaptic T-bars were evenly distributed at basal (strahler order: 5) and
distal regions (strahler order: 1-4), 95% of their postsynaptic sites were located in the most distal
region (strahler order 2-1; **Figure 7B-C**). We also calculated the geodesic distance (i.e., along-the-
arbor distance) from pre- and post-synaptic sites to the basal root node, which is the node point
where the uPN enters the glomerulus and is equivalent to the closest point to the soma in our

reconstruction. The geodesic distance to the basal root node from the presynaptic site was
significantly shorter than for postsynaptic sites (**Figure 7 – figure supplement 1B**). The pre- and
postsynaptic sites of each autapse were either close to each other along the dendritic tree, or
distant from each other (see examples in the dendrogram depicted in **Figure 7D**). Thus, the
geodesic distance between pre- and postsynaptic sites, (see scheme in **Figure 7E**), as well as the
number of branching points between pre- and postsynaptic partners, were bimodally distributed
(**Figure 7F-G**). Autapses that connected distant dendritic branches were more frequent than those
that connected close dendritic branches (**Figure 7E-G**). In summary, we found abundant autapses
within the uPN dendrite of DL5 and they were unevenly distributed, with many output sites located
in a few sub-branches connecting distal dendritic regions.

**DISCUSSION**

We hypothesized that specialized, narrowly tuned olfactory glomeruli differ in their
ultrastructure and microcircuitry from broadly tuned glomeruli. By comparing data obtained with
dense reconstructions of two narrowly tuned olfactory glomeruli with that of a broadly tuned
glomerulus in *Drosophila melanogaster,* we found prominent features of narrowly tuned glomeruli
involving synaptic composition, lateralization of sensory input and synaptic circuitry.

**Glomerular circuit analysis: a correlative approach**

The small size of olfactory glomeruli in *Drosophila* gave us the opportunity to reconstruct
and analyze the dense connectome of entire glomeruli with volume-based electron microscopy in
a reasonable time period. Here we developed a correlative workflow that combines transgenic
neuron labeling with near-infra-red-laser-branding for precise volume targeting. We then used FIB-
SEM (Bishop *et al.*, 2011) to resolve glomerular networks at the synaptic level. A similar procedure
was used recently to investigate single cellular organelles (Ronchi *et al.*, 2021). An advantage of
this approach is that it facilitates localization of the volume of interest with high precision and
consequently limits to a minimum the volume to be scanned and reconstructed. At the same time,
the limitation in volume is a drawback of our workflow, as it was impossible to reconstruct neurons
back to their soma. This fact prevented the identification of individual neurons as in other
connectome studies (Bates *et al.*, 2020; Berck *et al.*, 2016; Eichler *et al.*, 2017; Horne *et al.*, 2018;
Scheffer *et al.*, 2020; Schlegel *et al.*, 2021; Xu *et al.*, 2020; Zheng *et al.*, 2018).

We provide data on innervation and synapse density of olfactory sensory neurons (OSNs),
uniglomerular projection neurons (uPNs) and multiglomerular neurons (MGNs) in the *Drosophila*
antennal lobe (AL). We observed a higher innervation density of all neuron types but mainly by
uPNs and MGNs and in parallel higher density of synaptic contacts along OSN terminals in the
narrowly tuned DA2 compared with DL5. These results suggest that narrowly tuned glomeruli have
a more densely packed neuropil, with more numerous synaptic connections in the feedforward
motifs OSN>uPN and OSN>MGN. Overall, our observations on synapse density were comparable
with previous reports (Horne *et al.*, 2018; Mosca *et al.*, 2014; Rybak *et al.*, 2016).

**Specific features of narrowly tuned glomerular circuits**

Our analysis revealed circuit features in the narrowly tuned glomerulus DA2 and VA1v
that might be adaptations specific of such dedicated glomerular circuits. Nevertheless, future
studies, analyzing precise numbers of synaptic connections in more individuals, combined with
physiological studies and computational models are required to test this hypothesis.

**The OSN>uPN feedforward connection is stronger in narrowly tuned glomeruli**

Presynaptic OSN terminals provide the major input to uPNs in insect olfactory glomeruli
(Chen *et al.*, 2005; Hansson *et al.*, 2000; Horne *et al.*, 2018; Kazama *et al.*, 2008; Lei *et al.*, 2010;
Rybak *et al.*, 2018; Schlegel *et al.*, 2021; Tobin *et al.*, 2017). Here we showed that this connection
is stronger in DA2 and VA1v than in DL5 (**Figure 5 and 6**). A strong OSNs>uPN synaptic connection
will drive non-linear signal amplification, which improves signal detection at low odor
concentrations (Bhandawat *et al.*, 2007; Kazama *et al.*, 2008; Masse *et al.*, 2009; Ng *et al.*, 2002).
A larger number of synapses of this type could be an adaptation to improve this amplification
effect, as shown by artificial increase of synaptic sites in the AL  (Acebes *et al.*, 2001) and in lateral
horn dendrites (Liu *et al.*, 2022).

Each of the 7 uPNs in DA2 received convergent synaptic input from almost all DA2-OSNs.
This is in agreement with reports on the narrowly tuned glomeruli DA1 and VA1v (Agarwal *et al.*,
2011; Horne *et al.*, 2018; Jeanne *et al.*, 2015) and for broadly tuned glomeruli (Chen *et al.*, 2005;
Kazama *et al.*, 2009; Masse *et al.*, 2009; Tobin *et al.*, 2017; Vosshall *et al.*, 2000). High OSN>uPN
convergence is the main driver of highly correlated activity among uPNs in pheromone coding
glomeruli in flies as well as moths (Kazama *et al.*, 2009; Rospars *et al.*, 2014). High convergence in
the lateral horn improves signal transmission from uPNs to lateral horn neurons without sacrificing
speed (Huoviala *et al.*, 2020; Jeanne *et al.*, 2015). In the mushroom body calyces, however, the
high degree of convergence is only pursued for DA2 uPNs, which converge onto few Kenyon cells,
whereas VA1v uPNs synapse randomly onto many dispersed Kenyon cells (Caron 2013; Zheng
2020; Li 2020), indicating diverse signal integration in the mushroom body.

From our study, we hypothesize that in narrowly tuned glomerular circuits, which have
more uPNs, the maintained strong OSN>uPN convergence, improve signal transmission accuracy.
Secondly, a stronger OSN>uPN connection might compensate for the lack of OSN>uPN signal
transmission sites in the case of odorants activating OSNs in a single glomerulus.

**Reciprocal connections between sister OSNs and sister uPNs are stronger in narrowly tuned**
**glomeruli**

532
533

The reciprocal OSN-OSN synapse is generally stronger in narrowly tuned glomeruli DA1, DL3
and DL4, compared to broadly tuned glomeruli DL5, DM6, DM3 and DM4 (Dweck *et al.*, 2015;
Ebrahim *et al.*, 2015; Grabe *et al.*, 2016; Knaden *et al.*, 2012; Schlegel *et al.*, 2021; Seki *et al.*, 2017;
Suh *et al.*, 2004; Tobin *et al.*, 2017). A high degree of axo-axonic synapses between sister OSNs was
also found in VA1v (Horne *et al.*, 2018; Schlegel *et al.*, 2021)and DA2 but not in the DL5 (this study).
Hence, we suggest that a strong OSN-OSN connection is a characteristic feature of the synaptic
circuitry of narrowly tuned olfactory glomeruli. Axo-axonic connections have also been reported
between gustatory and mechanosensory neurons in *Drosophila* larvae (Miroschnikow *et al.*, 2018)
and in the olfactory epithelium and the olfactory bulb of vertebrates (Hirata, 1964; Shepherd *et*

*al.*, 2021). In vertebrates, axo-axonic synapses between excitatory sensory neurons are involved in
correlated transmitter release (Cover *et al.*, 2021), reminiscent of correlated uPN activity due to
reciprocal synaptic and electric coupling in the *Drosophila* AL and LH (Huoviala *et al.*, 2020; Kazama
*et al.*, 2009). A strong OSN-OSN connection also has the potential to increase the correlation of
OSN spiking events and therefore facilitate a robust OSN signal (de la Rocha *et al.*, 2007).

Reciprocal dendro-dendritic synapses between sister uPNs are reported here for the DA2
have been reported previously also for glomeruli DM6, DM4, VA7 and VA1v (Horne *et al.*, 2018;
Kazama *et al.*, 2009; Rybak *et al.*, 2016; Tobin *et al.*, 2017). These types of synapses enhance uPN
signal correlation (Kazama *et al.*, 2009), as reported for mitral and tufted cells of the vertebrate
olfactory bulb, the circuit equivalent to PNs of insect ALs (Christie *et al.*, 2005; McTavish *et al.*,
2012; Shepherd *et al.*, 2021). In *Drosophila* multiple uPNs could induce correlated PN
depolarization events, which improve the signal-to-noise-ratio of PN signal transmission (Chen *et*
*al.*, 2005; Jeanne *et al.*, 2015; Kazama *et al.*, 2009).

In summary, our data give evidence that reciprocal OSN-OSN and uPN-uPN connections are
a prominent feature of the synaptic circuit of narrowly tuned glomeruli. We suggest that those
reciprocal OSN-OSN and uPN-uPN connections support correlation of neuronal activity and
therefore boosts signal induced depolarization events. This will in turn enhance the signal-to-noise
ratio (accuracy) and transmission probability of weak and/or irregular odorant input, increasing
processing speed.

**Less lateralization in the OSN bilateral connectivity in narrowly tuned glomeruli**

In *Drosophila*, most OSN axons project bilaterally and form synapses in their corresponding
glomerulus on both the left and right brain hemispheres (Couto *et al.*, 2005; Kazama *et al.*, 2009;
Schlegel *et al.*, 2021; Silbering *et al.*, 2011; Stocker *et al.*, 1990; Tobin *et al.*, 2017; Vosshall *et al.*,
2000). This is rarely observed in other insects and absent in vertebrates (Anton *et al.*, 2003; Dalal
*et al.*, 2020; Galizia *et al.*, 1998; Hansson *et al.*, 2000; Masson *et al.*, 1990; Parthasarathy *et al.*,
2013; Stocker *et al.*, 1983). In the mammalian olfactory system, bilateral comparison of olfactory
input only occurs in higher brain centers (Dalal *et al.*, 2020). In flies, bilateral sensory input enables
them to discriminate odor sources of different spatial origin through bilateral comparison of
olfactory stimulation (Borst *et al.*, 1982; Duistermars *et al.*, 2009; Gaudry *et al.*, 2013; Mohamed
*et al.*, 2019a; Taisz *et al.*, 2022). Asymmetric OSN connectivity, shown for many olfactory OSNs
(Schlegel *et al.*, 2021; Tobin *et al.*, 2017) seems to be the origin of a bilateral contrast in the uPN
response (Agarwal *et al.*, 2011; Gaudry *et al.*, 2013; Taisz *et al.*, 2022; Tobin *et al.*, 2017), and is
most likely the key to precise odor source localization (Taisz *et al.*, 2022). Bilateral comparison is
also used in the lateral horn (a higher olfactory brain center in *Drosophila*) for odorant position
coding (Mohamed *et al.*, 2019a). However, not all glomeruli are similar in the magnitude of
bilateral asymmetry with respect to their OSN connectivity (Schlegel *et al.*, 2021) or their uPN
responses (Agarwal *et al.*, 2011).

In agreement with observations in other olfactory glomeruli (Schlegel *et al.*, 2021; Tobin *et*
*al.*, 2017), we found that glomeruli DL5, DA2 and VA1v (data from: (Horne *et al.*, 2018) have
ipsilaterally asymmetric OSN synaptic output to excitatory uPNs and sister OSNs and contralaterally
an enhanced OSN>MGN output (**Figure 4**). We believe that, in agreement with a recent study,
these asymmetric connections determine a strong left-right-contrast in the uPN response, akin to
a "winner-takes-all" principle (Taisz *et al.*, 2022).

We also observed that the degree of bilateral OSN asymmetry in DA2 and VA1v was much
weaker than in DL5 (**Figure 4**). Weakly lateralized OSN connectivity is perhaps insufficient to induce
an adequate bilateral contrast necessary for odor source localization. Recent work supports this
idea by showing the importance of the interplay of asymmetric OSN signaling and LN inhibition to
enhance the bilateral contrast of uPN activity and to facilitate navigation (Taisz *et al.*, 2022).

Why do these narrowly tuned glomeruli have weaker bilateral contrast than broadly tuned
glomeruli? The answer could lie in the ecological significance of the individual odorants. Geosmin,
encoded by glomerulus DA2 (Stensmyr *et al.*, 2012), and the pheromone methyl laurate, encoded
by glomerulus VA1v (Dweck *et al.*, 2015), act at short distances, mainly when the fly is walking and
not flying. Perhaps, the behavioral response to geosmin or methyl laurate does not need a precise
odor source location. On the other hand, food odor detection at a distance, which happens mainly
at flying conditions, needs continuous processing of odor position and body alignment to navigate
towards the odor source (Demir *et al.*, 2020; Thoma *et al.*, 2015). The bilateral OSN projection onto
uPNs in DA2 and VA1v potentially has a distinct function other than odor position coding and could,
via the enhancement of the effect of convergence of OSN>uPN signal transmission, enhance odor
signal amplification (Bhandawat *et al.*, 2007; Jeanne *et al.*, 2015; Kazama *et al.*, 2009; Masse *et al.*,
2009)

**Distinct synaptic integration of local modulatory neurons in narrowly tuned glomeruli**

MGNs are composed of multiglomerular projection neurons (mPNs) that project directly to
the LH (Bates *et al.*, 2020; Jefferis *et al.*, 2007; Strutz *et al.*, 2014) and inhibitory and excitatory local
interneurons (LNs) that interconnect the AL glomeruli (Chou *et al.*, 2010; Liu *et al.*, 2013; Masse *et*
*al.*, 2009; Okada *et al.*, 2009; Seki *et al.*, 2010). Since LNs are the most numerous and broadly
arborizing of the multiglomerular cell types in the AL (Chou *et al.*, 2010; Lin *et al.*, 2012), we focus
our discussion on these. Multiglomerular LNs are crucial for the modulation of the OSN>uPN signal
transmission (Chou *et al.*, 2010; Galizia, 2014; Masse *et al.*, 2009; Seki *et al.*, 2010; Szyszka *et al.*,
2015).

Previous observations have shown that glomeruli DA2 and VA1v have a lower number of
innervating LNs (Chou *et al.*, 2010; Grabe *et al.*, 2016) and receive less global interglomerular LN
inhibition than broadly tuned glomeruli (Hong *et al.*, 2015). We therefore assumed that DA2 or
VA1v would have a lower LN innervation density and less LN synaptic integration in their circuitry.
However, we did not observe a general lower synaptic integration in DA2 (**Figure 5**) and found a

greater MGN innervation density, and a higher density of input sites than in DL5. VA1v MGNs on
the other hand received less synaptic input and provided less output in its glomerular circuit than
MGNs in DL5.

Taking a closer look at particular synaptic connection motifs of MGNs we saw that narrowly
tuned glomeruli had a relatively weak uPN>MGN feedback (**Figure 6**). uPN feedback onto LNs and
their reciprocal connection (LN>uPN) were reported in *Drosophila* and other insects, such as honey
bees, cockroaches and moths, but their function is still poorly understood (Boeckh *et al.*, 1993;
Sachse *et al.*, 2002; Sun *et al.*, 1997). In the honey bee reciprocal dendro-dendritic synapses
between excitatory and inhibitory neurons enhance signal contrast and the reliability of true signal
representations throughout the AL (Sachse *et al.*, 2002; Yokoi *et al.*, 1995). Here we could not
differentiate the LN types involved in the uPN>MGN synaptic motif. However, the prevailing
uPN>LN synapses involve mainly widespread pan-glomerular LNs in the adult (Horne et al., 2018)
and larval AL (Berck et al., 2016), which are important for combinatorial coding (Galizia, 2014;
Sachse *et al.*, 2016). Thus, weaker uPN>MGN feedback in the narrowly tuned DA2 and VA1v circuits
might be a compensatory mechanism to lower the computational demand of interglomerular
communication for odor identity coding.

We also observed that OSNs received less MGN in the narrowly tuned DA2 and VA1v than
in the DL5, suggesting that the OSNs in DA2 and VA1v receive relatively weak presynaptic
inhibition. Pan-glomerular GABAergic LNs induce presynaptic inhibition at OSN presynaptic site
(Berck *et al.*, 2016; Schlegel *et al.*, 2021). These inhibitory LNs are drivers of balanced glomerular
gain control and are a key player for odor identity coding, balancing incoming and alternating odor
intensities (Asahina *et al.*, 2009; Galizia, 2014; Hong *et al.*, 2015; Olsen *et al.*, 2008; Root *et al.*,
2008; Sachse *et al.*, 2016; Silbering *et al.*, 2008; Szyszka *et al.*, 2015; Wang, 2012). Our data support
these observations and provide an argument for why narrowly tuned OSNs receive much lower
inhibition during AL stimulation with odorants activating other OSN populations (Hong *et al.*, 2015).
Even though DA2 and VA1v might receive less interglomerular inhibition, their OSN>MGN output
is still strong, in agreement with studies showing that throughout the AL, global lateral inhibition
mediated by LNs scales with general OSN activation (Hong *et al.*, 2015; Olsen *et al.*, 2008).

In summary, narrowly tuned circuits are probably influenced more strongly by
intraglomerular than by interglomerular modulation. Narrowly tuned circuits perhaps have greater
computational capacities in intraglomerular modulation of signal transmission, which could be
important for example for PN fine-tuning and response adjustment (Assisi *et al.*, 2012; Ng *et al.*,
2002).

Above we discussed putative generic features of narrowly tuned glomerular circuits.
Besides these circuit features, we found a strong MGN>MGN connection in the aversive glomerular
circuits DA2 and DL5 in contrast to a much weaker MGN>MGN connection in the attractive
glomerulus VA1v (Dweck *et al.*, 2015; Knaden *et al.*, 2014; Knaden *et al.*, 2012; Mohamed *et al.*,
2019b; Stensmyr *et al.*, 2012). Why do aversive olfactory circuits have a stronger MGN>MGN

connection than attractive circuits? In the larval *Drosophila* AL, reciprocal LN>LN synapses induce
disinhibition induced by a strong connection between the pan-glomerular LNs and a bilateral
projecting LN, the Keystone LN, which synapses strongly onto pan-glomerular LNs and selectively
onto OSNs, which are activated by attractive food odors. This is thought to be a key feature to
switch from homogenous to heterogeneous presynaptic inhibition and therefore to a selective gain
control enhancing contrast between attractive and aversive odor activation (Berck *et al.*, 2016).
Such balanced inhibitory systems could also be present in the adult *Drosophila* AL, reflected in the
strong LN>LN connection in DA2 and DL5. Disinhibition of interglomerular presynaptic inhibition in
aversive glomeruli circuits might be important for the fly to stay vigilant to aversive odors, while
perceiving attractive cues, for example during feeding conditions so that a fast switch in behavior
can be initiated if necessary.
666

**Autaptic connection within the dendritic tree of a single uPN**

We observed autapses along the large dendritic tree of the single DL5-uPN. To our
knowledge, this is the first report of bulk dendro-dendritic autapses in the *Drosophila* olfactory
system, indicating a cell-type specific occurrence of autapses in the DL5-uPN as reported for other
cell types in the optic lobe (Takemura et al., 2015). Autapses are also reported to be present at
different frequencies in different types of neurons in the mammalian brain (Bacci *et al.*, 2006;
Bekkers, 1998; Bekkers, 2009; Ikeda *et al.*, 2006; Saada *et al.*, 2009; Tamás *et al.*, 1997; Van der
Loos *et al.*, 1972). In *Drosophila,* most uPNs are cholinergic (Croset *et al.*, 2018; Kazama *et al.*, 2008;
Tanaka *et al.*, 2012; Yasuyama *et al.*, 2003; Yasuyama *et al.*, 1999) and the DL5-uPN autapses
reported here might activate either nicotinic or muscarinic acetylcholine postsynaptic receptors.
Muscarinic acetylcholine receptors have an inhibitory effect in the Kenyon cells of the mushroom
body (Bielopolski *et al.*, 2019), but mediate excitation in the AL (Rozenfeld *et al.*, 2019).
What could be the function of these autaptic feedback loops within the DL5-uPN dendritic
tree? Recent studies in vertebrates show that excitatory autapses enhance neuron bursting and
excitability (Guo *et al.*, 2016; Wiles *et al.*, 2017; Yin *et al.*, 2018). Autaptic inhibitory connections
have been implicated in circuit synchronization, spike-timing precision, self-stabilization of
neuronal circuits and feedback inhibition (Bacci *et al.*, 2006; Bekkers, 1998; Ikeda *et al.*; Saada *et*
*al.*, 2009; Tamás *et al.*, 1997; Van der Loos *et al.*, 1972).
Autapses in the DL5 uPN form mainly long-distance feedback loops, connecting distinct
dendritic subtrees and the basal dendrite region (closer to the soma) with distal branches. This
spatial segregation is similar to the distribution of non-autaptic pre- and postsynaptic sites in
*Drosophila* uPNs, where presynapses are located more frequently at basal dendrites than
postsynapses (Rybak *et al.*, 2016) and other insects, such as *Periplaneta americana* and moths (Lei
*et al.*, 2010; Malun, 1991; Sun *et al.*, 1997). Dendro-dendritic autaptic feedback loops connecting
basal to distal branches and distinct dendritic subtrees of a large dendritic tree might facilitate

activity correlation between distant dendritic subunits, as described for non-autaptic, reciprocal
uPN>uPN connections (Kazama *et al.*, 2009). This could be important in a large compartmentalized
dendrite that receives inhomogeneous excitation by several OSNs at distinct dendritic sites, in
order to enhance synchronized depolarization events along the dendrite, supporting signal
integration (Graubard *et al.*, 1980; Tran-Van-Minh *et al.*, 2015). Clustered autapses could mediate
local signal input amplification for distinct dendritic subunits (Kumar *et al.*, 2018; Liu *et al.*, 2022).
Autaptic contacts, finally, could be able to shift the uPN membrane depolarization towards the
spiking threshold, and enhance the firing probability during activation.

In conclusion, we provide a comprehensive comparative analysis of the ultrastructure and
synaptic circuitry of two functionally diverse olfactory glomeruli with distinct computational
demands, processing either single odorant information in a dedicated olfactory pathway (DA2) or
input regarding several odorants and taking part in combinatory coding across distributed
glomeruli (DL5). Our work provides an opportunity to gain insight into variations in network
architecture and provides fundamental knowledge for future understanding of glomerular
processing. By comparing our data with those from another narrowly tuned glomerulus (VA1v), we
distilled prominent circuit features that suggest that narrowly tuned glomerular circuits encode
odor signals with a weaker left-right-contrast, improved accuracy, stronger signal amplification and
stronger intraglomerular signal modulation relative to broadly tuned glomeruli. Our findings reveal
the existence of autapses in olfactory glomeruli and indicate that dendro-dendritic autapses play
an important role in dendritic signal integration.

## MATERIAL AND METHODS:

### Fly line and fly rearing

Flies of the genotype *Orco-GAL4; UAS-GCaMP6s* (Vosshall *et al.,* 2000) were obtained from the Bloomington *Drosophila* Stock Center (https://bdsc.indiana.edu) and reared on standard *Drosophila* food at 25°C and 70% humidity on a 12 h:12 h day:night cycle. Seven-days old female flies were used. In these flies, Orco-positive olfactory sensory cells emit green fluorescence, making possible to identify individual glomeruli.

### Brain dissection and fixation for Focus Ion Beam Microscopy-Scanning electron microscopy (FIB-SEM)

Two 7-day old female flies were anesthetized with nitric oxide (with Sleeper TAS; INJECT+MATIC, Switzerland) and decapitated with forceps. Heads were dipped for one minute in 0.05% Triton X-100 in 0.1M Sorensen's phosphate buffer, pH 7.3 and transferred to a droplet of freshly prepared ice-cooled fixative (2.5% glutaraldehyde and 2.0% paraformaldehyde in 0.1M Sørensen's phosphate buffer, pH 7.3; as in (Karnovsky, 1965). The proboscis was removed and the back of the head was opened to improve fixative penetration. After 5-10 minutes, the brain was dissected out of the head capsule and post-fixed for two hours on ice. Fixation was stopped by rinsing the brain several times in ice-cooled 0.1M Sørensen's phosphate buffer, pH 7.3 (after (Rybak *et al.,* 2016)).

### Laser branding of glomeruli for identification during FIB-SEM microscopy

To identify the glomeruli of interest at the ultrastructural level and to limit to a minimum the volume of tissue to be scanned with FIB-SEM, near-infrared laser branding (NIRB,(Bishop *et al.,* 2011). Glomeruli of interest were first located with light microscopy in brains of *Orco-GAL4; UAS-GCaMP6s* flies using a confocal microscope (ZEISS LSM 710 NLO, Carl Zeiss, Jena, Germany), a 40x water immersion objective (W Plan-Apochromat 40x/1.0 DIC VIS-IR, Carl Zeiss, Jena, Germany), a laser wavelength of 925 nm at 30% laser power and ZEN software (Carl Zeiss, Jena, Germany). Once glomeruli DA2 or DL5 were identified by means of location, shape and size the volume of interest (VOI) was tagged with fiducial marks ("laser-branded") close to the borders of the glomerulus (**Figure 1A-B**), using an infrared Chameleon Ultra diode-pumped laser (Coherent, Santa Clara, USA)

at wavelength 800 nm and at 75-90% of laser power). Two laser scan rounds were performed for

each induced fiducial brand. DA2 (right AL) and DL5 (left AL) were laser-branded in the same fly. A

second glomerulus DA2 was marked in the right AL of another fly.

**Transmission Electron Microscopy**

Brains were rinsed with 2.5% sodium-cacodylate buffer and post-fixed in 1% osmium tetroxide and

1% potassium ferrocyanide in cacodylate buffer for 2 hours. After rinsing with cacodylate buffer

the brains were dehydrated with a graded acetone series (30%-100% acetone), including an

additional *en bloc* staining step in-between, in which the brains were incubated in 1% uranyl

acetate in 50% acetone for 30 minutes in the dark, and gradually infiltrated with Araldite (glycerol-

based aromatic epoxy resins; Serva, Germany). In the final step, the tissue was embedded in pure

resin and left in a 60°C incubator to polymerize for 48h. Resin blocks were trimmed with a Reichert

UltraTrim microtome (Leica, Deer Park, USA) and the fiducial laser marks were then located in

semi-thin sections. To check tissue quality before performing high-resolution volume-based

electron microscopy, serial sections 50 nm in thickness were cut with a diamond knife (Ultra 45°,

Diatome, Switzerland) on a Reichert Ultracut S ultramicrotome (Leica, Deer Park, Germany),

collected on single slot grids (2 x 1 mm), and imaged with a JEM 1400 electron microscope (Jeol,

Freising, Germany) operated at 80 kV. Digital micrographs were obtained with a Gatan Orius SC

1000 CCD camera (Gatan Orius SC 1000; Gatan, Pleasanton, USA) controlled with the Gatan

Microscopy Suite software Vers. 2.31.734.0.

.

**Focused Ion Beam-Scanning Electron Microscopy (FIB-SEM)**

Before serial Focused Ion Beam milling and Scanning Electron Microscopy imaging (FIB-SEM; (Knott

*et al.*, 2008; Xu *et al.*, 2017), the surface of the trimmed block was coated with a conductive carbon

layer to prevent charging artifacts. A FEI Helios NanoLab G3 UC (FEI, Hilsboro, USA) was used for

FIB-SEM process. The laser marks used to landmark the VOI were visible across the surface of the

block. The VOI surface was protected via a local deposition of platinum using a gas injection system

for subsequent ion and electron beam deposition. The material surrounding the VOI at the front

and the side was removed to reduce re-deposition of material during FIB-SEM. Serial images across

the entire VOI were generated by repeated cycles of milling slices orthogonal to the block surface

via FIB and imaging via SEM the newly exposed surface. The tissue was milled with a focused beam

of gallium ions using FEI's Tomahawk ion column (accelerating voltage: 30 kV, beam current: 790

pA, milling steps: 20 nm). FEI's Elstar electron column was used to create the backscattered

electron contrast images using an In-Column Detector (accelerating voltage. 3kV; 1.6 nA; dwell

time: 10 µs). The DA2 and DL5 volumes in the first fly were imaged with a resolution of 4.9 x 4.9 x

20 nm$^3$/vox (DA2: 769 images with 4096 x 3536 pix; DL5: 976 images with 5218 x 3303 pix). The

volume of a second DA2 in a second fly was imaged with a resolution of 4.4 x 4.4 x 20 nm$^3$/vox (571

images with 4096 x 3536 pix). The milling/imaging cycles were controlled with the Auto Slice and

View 4.0 software (FEI, Hilsboro, USA).

**Image alignment, 3D reconstruction and segmentation**

FIB-SEM image stacks were aligned by maximizing the Pearson correlation coefficient of the central

part of two consecutive images using template matching from the openCV library

(https://opencv.org). Dense reconstructions of the glomeruli were produced by manually tracing

all neuronal fibers and by annotating all synapses within the two glomeruli, using a skeleton-based

reconstruction procedure similar to previous approaches (Berck *et al.*, 2016; Schneider-Mizell *et

al.*, 2016; Zheng *et al.*, 2017). Up to five independent tracers and two reviewers participated in an

iterative reconstruction process using the web-based reconstruction software CATMAID

(http://www.catmaid.org; RRID:SCR_006278; (Saalfeld *et al.*, 2009; Schneider-Mizell *et al.*, 2016);

**Figure 1D, Figure 1 – video 1**), performing a dense reconstruction of synaptic neuropil. In a second

fly, neurons of a DA2 glomerulus were manually reconstructed with the volume-based

reconstruction method TrakEM2 (Cardona *et al.*, 2012), an ImageJ (Fiji) plugin

(https://imagej.net/TrakEM2).

**Neuron visualization**

Reconstructed neurons were visualized using CATMAID 3D visualization (http://www.catmaid.org

and  using Blender 3D, an open-source 3D software (https://www.blender.org/; **Figure 7 – figure

supplement 1**). Neuron data from CATMAID were imported and shaded by Strahler order using an

existing CATMAID plugin for Blender (https://github.com/schlegelp/CATMAID-to-Blender; Schlegel

et al., 2016). Volume-based reconstructions were visualized as surface shapes in CATMAID

imported from TrakEM2 (https://imagej.net/TrakEM2).

**Glomerular border definition**

The definition of the boundary between olfactory glomeruli was based on the combination of

several structural features: the spatial position of pre- and postsynaptic elements along OSN axons,

the position of the majority of uPN postsynaptic sites, the faint glial leaflets scattered at the

periphery of the glomerulus, and the fiducial laser marks (**Figure 1B, D**).

**Neuron identification**

Neuronal fibers were assigned to one of three pre-defined neuron classes: OSNs, uPNs, and MGNs.

The classification was based on their 3D shape (**Figure 2A**), their branching intensity (**Figure 2B**),

the average diameter of their fibers (neuronal profiles: **Figure 2A -** FIB-SEM image; exemplary

volume-based reconstruction), the ratio of T-bars-to-input sites and the size of their T-bars, which

were either "small" (few postsynaptic connections) or "large" (many postsynaptic connections

**Figure 2 – supplement 1B-D)**. In addition, several intracellular features helped to classify neuron

classes: the shape and appearance of mitochondria, the size and electron density of vesicles and

the amount of synaptic spinules (small filopodia-like invaginations of neighboring cells (**Figure 2A**

**-** FIB-SEM image; (Gruber *et al.*, 2018). OSNs and uPNs could be counted, due to their uniglomerular

character, by means of the identification of the axons (OSNs) or main dendrites (uPNs) entering

the glomerulus. The number of MGNs could not be counted because of their pan-glomerular

projection patterns in the AL. Ipsi- and contralateral OSNs in DA2 and DL5 were identified based

on the trajectory of axonal fibers and their entry location in each glomerulus, (example neurons:

**Figure 4B)**. Ipsilateral OSNs reach the glomerulus from the ipsilateral antennal nerve and leave the

glomerulus towards the antennal lobe commissure (ALC: (Tanaka *et al.*, 2012). Contralateral OSNs

reach the glomerulus projecting from the ALC.

**Data analysis**

With the aid of the web-based software CATMAID ([http://www.catmaid.org](http://www.catmaid.org)) the following

properties were quantified: glomerular volume, neuronal fiber length (in µm), number of fiber

branching points, number of synaptic input and output sites and T-bars (see data availability). In a

second fly, the volume of neurons in DA2 was measured (**Figure 2 – supplement 1A**) with the aid

of TrakEM2 (Cardona *et al.*, 2012), an ImageJ (Fiji) plugin ([https://imagej.net/TrakEM2](https://imagej.net/TrakEM2)). The

following calculations were performed:

1. $Innervation\ density = \frac{total\ neuron\ length\ (\mu m)}{glomerular\ volume\ (\mu m^3)}$ ,

835           a. calculated as a ratio: (1) the sum of all neuronal fibers of each neuron class or (2) all

836           together (**Table 1**) or (3) for each neuron individually (**Figure 3**)

837       2. $Glomerular\ synaptic\ density = \frac{\#\ of\ synaptic\ inputs,-\ outputs\ or\ T-bars}{glomerular\ volume\ (\mu m^3)}$ ,

838           a. calculated as a ratio: (1) the sum of all neuronal fibers of each neuron class or (2) all

839           together (**Table 1**) or (3) for each neuron individually (**Figure 3**)

3. $Neuronal\ synaptic\ density = \frac{\#\ of\ synaptic\ inputs-,outputs\ or\ T-bars}{neuronal\ fiber\ length\ (\mu m)}$ (**Table 1; Figure 3 –**

**figure supplement 1**)

4. $Synaptic\ ratios = \frac{\#\ of\ T-bars\ or\ outputs}{inputs}$ (represents the average for each neuron class;

**Table 1**)

5. $Polyadicity = \frac{\#\ of\ outputs}{T-bars}$ (represents the average number of postsynaptic sites at a T-bar

of each neuron class; **Table 1 and Figure 1E**)

6. Relative differences $= \frac{respective\ value\ target\ glomerulus - value\ source\ glomerulus}{source\ glomerulus} \times 100$

(**Table S1; Table S2**)

7. Relative synaptic strength $= \frac{\#\ of\ synaptic\ contacts\ from\ neuron\ class\ A\ to\ B}{\#\ all\ synaptic\ contacts\ in\ corresponding\ glomerulus}$ (**Table  S1;**

**Table S2**)

8. Fraction of output $= \frac{\#\ of\ outputs\ of\ neuron\ class\ A\ directed\ to\ neuron\ class\ B}{total\ \#\ of\ outputs\ of\ neuron\ class\ A} \times 100$

9. Fraction of input $= \frac{\#\ of\ inputs\ from\ neuron\ class\ A\ from\ class\ B}{total\ \#\ of\ inputs\ of\ neuron\ class\ A} \times 100$

Graphs were made with the programming language R and RStudio (R Core Team, 2018) using the

packages 'ggplot2' and 'reshape' (see data availability) or with Python (see data availability). EM

and fluorescence images were visualized with ImageJ (Fiji) (http://fiji.sc/; (Schindelin *et al.*, 2012)

(and All figures were compiled with Adobe Illustrator CS5 software (Adobe Inc.).

Statistical analysis was performed with R Studio (R Studio Team, 2016) using the packages

'ggsignif'. Differences between samples DA2 and DL5 or between ipsilateral and contralateral OSNs

were tested for significance with a two-sided student's t-test if sample size was normally

distributed, or with Wilcoxon two sample test if the data was not normally distributed (noted in

figure legend). Data is in all cases represented as mean + standard deviation.

### Analysis of autapses

The location of autapses, the measurement of their geodesic distance (distance along the neuronal

dendrite) and the number of branching points from point A (presynaptic site) to B (postsynaptic

profile) was analyzed with Python using the package 'neuroboom'

https://github.com/markuspleijzier/neuroboom (see also data availability).

### Data availability

Datasets will be available through the public CATMAID instance:

https://catmaid.ice.mpg.de/catmaid_2020.02.15/. Neurons are named according to their neuron

classification. The neuroboom Python package was used for dendrogram analysis, available at

https://github.com/markuspleijzier/neuroboom and https://pypi.org/project/neuroboom/.

Further newly generated source codes can be made available upon request. Source data files are

provided (see source data files).

**Acknowledgments**

The authors are most grateful to Katrin Buder for the support with electron microscopy sample

preparation, and Veit Grabe for advice on two-photon imaging. Great thanks also to Albert Cardona

for discussion on synaptic networks, him, and Tom Kazimiers (Kazmos GmbH) for instruction in the

use of CATMAID. The neuronal reconstructions were conducted with the outstanding support

of Damilola E. Akinyemi**,** Eckard E. Schumann, and Michael Adewoye. We thank Martin Nawrot and

Magdalena Springer for constructive comments and discussions about autapses. The work was

supported by Roland Kilper and Ute Müller (aura optics, Jena), the European Regional

Development Fund, by funds from the DFG (grant no. 430592330), in the Priority Program

'Evolutionary Optimisation of Neuronal Processing' (DFG-SPP 2205) and by the Max Planck Society.

**REFERENCES**

Acebes, A., & Ferrus, A. (2001). Increasing the Number of Synapses Modifies Olfactory Perception
in Drosophila. *Journal of Neuroscience, 21*(16), 6264–6273.
Agarwal, G., & Isacoff, E. (2011). Specializations of a pheromonal glomerulus in the Drosophila
olfactory system. *Journal of Neurophysiology, 105*(4), 1711-1721.
doi:10.1152/jn.00591.2010
Ai, H., & Hagio, H. (2013). Morphological analysis of the primary center receiving spatial
information transferred by the waggle dance of honeybees. *Journal of Comparative
Neurology, 521*(11), 2570-2584. doi:10.1002/cne.23299
Andersson, M. N., Löfstedt, C., & Newcomb, R. D. (2015). Insect olfaction and the evolution of
receptor tuning. *Frontiers in Ecology and Evolution, 3*. doi:10.3389/fevo.2015.00053
Anton, S., van Loon, J. J., Meijerink, J., Smid, H. M., Takken, W., & Rospars, J. P. (2003). Central
projections of olfactory receptor neurons from single antennal and palpal sensilla in
mosquitoes. *Arthropod Structure and Development, 32*(4), 319-327.
doi:10.1016/j.asd.2003.09.002
Asahina, K., Louis, M., Piccinotti, S., & Vosshall, L. B. (2009). A circuit supporting concentration-
invariant odor perception in Drosophila. *Journal of Biology, 8*(1), 9. doi:10.1186/jbiol108
Assisi, C., Stopfer, M., & Bazhenov, M. (2012). Excitatory Local Interneurons Enhance Tuning of
Sensory Information. *PLoS Computational Biology, 8*(7), e1002563.
doi:10.1371/journal.pcbi.1002563
Auer, T. O., Khallaf, M. A., Silbering, A. F., Zappia, G., Ellis, K., Álvarez-Ocaña, R., . . . Benton, R.
(2020). Olfactory receptor and circuit evolution promote host specialization. *Nature*.
doi:10.1038/s41586-020-2073-7
Bacci, A., & Huguenard, J. R. (2006). Enhancement of spike-timing precision by autaptic
transmission in neocortical inhibitory interneurons. *Neuron, 49*(1), 119-130.
doi:10.1016/j.neuron.2005.12.014
Bates, A. S., Schlegel, P., Roberts, R. J. V., Drummond, N., Tamimi, I. F. M., Turnbull, R., . . .
Jefferis, G. (2020). Complete Connectomic Reconstruction of Olfactory Projection Neurons
in the Fly Brain. *Current Biology, 30*(16), 3183-3199 e3186. doi:10.1016/j.cub.2020.06.042
Bekkers, J. M. (1998). Neurophysiology: are autapses prodigal synapses? *Current Biology, 8*(2),
R52-55. doi:10.1016/s0960-9822(98)70033-8
Bekkers, J. M. (2009). Synaptic Transmission: Excitatory Autapses Find a Function? *Current
Biology, 19*(7), R296-R298. doi:http://dx.doi.org/10.1016/j.cub.2009.02.010
Benton, R., Sachse, S., Michnick, S. W., & Vosshall, L. B. (2006). Atypical Membrane Topology and
Heteromeric Function of *Drosophila* Odorant Receptors In Vivo. *PLoS Biology, 4*(2), 240-
257. doi:10.1371/journal.pbio.0040020
Berck, M. E., Khandelwal, A., Claus, L., Hernandez-Nunez, L., Si, G., Tabone, C. J., . . . Cardona, A.
(2016). The wiring diagram of a glomerular olfactory system. *Elife, 5*, e14859.
doi:10.7554/eLife.14859
Bhandawat, V., Olsen, S. R., Gouwens, N. W., Schlief, M. L., & Wilson, R. I. (2007). Sensory
processing in the *Drosophila* antennal lobe increases reliability and separability of
ensemble odor representations. *Nature Neuroscience, 10*(11), 1474-1482.
doi:10.1038/nn1976

Bielopolski, N., Amin, H., Apostolopoulou, A. A., Rozenfeld, E., Lerner, H., Huetteroth, W., . . .
Parnas, M. (2019). Inhibitory muscarinic acetylcholine receptors enhance aversive
olfactory learning in adult Drosophila. *Elife, 8*. doi:10.7554/eLife.48264
Bishop, D., Nikic, I., Brinkoetter, M., Knecht, S., Potz, S., Kerschensteiner, M., & Misgeld, T. (2011).
Near-infrared branding efficiently correlates light and electron microscopy. *Nat Meth,*
*8*(7), 568-570. doi:10.1038/nmeth.1622
Boeckh, J., Distler, P., Ernst, K. D., Hösl, M., & Malun, D. (1990). Olfactory bulb and antennal lobe.
In D. Schild (Ed.), *NATO ASI Series, Vol. H39: Chemosensory Information Processing* (pp.
201-227). Berlin, Heidelberg: Springer Verl.
Boeckh, J., & Tolbert, L. P. (1993). Synaptic organization and development of the antennal lobe in
insects. *Microscopy Research and Technique, 24*(3), 260-280.
doi:10.1002/jemt.1070240305
Borst, A., & Heisenberg, M. (1982). Osmotropotaxis inDrosophila melanogaster. *Journal of*
*comparative Physiology, 147*(4), 479-484. doi:10.1007/BF00612013
Briggman, K. L., & Denk, W. (2006). Towards neural circuit reconstruction with volume electron
microscopy techniques. *Current Opinion in Neurobiology, 16*(5), 562-570.
doi:10.1016/j.conb.2006.08.010
Butcher, N. J., Friedrich, A. B., Lu, Z., Tanimoto, H., & Meinertzhagen, I. A. (2012). Different
classes of input and output neurons reveal new features in microglomeruli of the adult
*Drosophila* mushroom body calyx. *Journal of Comparative Neurology, 520*(10), 2185-2201.
doi:10.1002/cne.23037
Cardona, A., Saalfeld, S., Schindelin, J., Arganda-Carreras, I., Preibisch, S., Longair, M., . . .
Douglas, R. J. (2012). TrakEM2 Software for Neural Circuit Reconstruction. *PLoS ONE, 7*(6),
e38011. doi:10.1371/journal.pone.0038011
Cardona, A., Saalfeld, S., Tomancak, P., & Hartenstein, V. (2009). *Drosophila* brain development:
closing the gap between a macroarchitectural and microarchitectural approach. *Cold*
*Spring Harb Symp Quant Biol, 74*, 235-248. doi:10.1101/sqb.2009.74.037
Chen, W. R., & Shepherd, G. M. (2005). The olfactory glomerulus: A cortical module with specific
functions. *Journal of Neurocytology, 34*, 353–360.
Chou, Y. H., Spletter, M. L., Yaksi, E., Leong, J. C., Wilson, R. I., & Luo, L. (2010). Diversity and
wiring variability of olfactory local interneurons in the *Drosophila* antennal lobe. *Nature*
*Neuroscience, 13*(4), 439-449. doi:10.1038/nn.2489
Christie, J. M., Bark, C., Hormuzdi, S. G., Helbig, I., Monyer, H., & Westbrook, G. L. (2005).
Connexin36 mediates spike synchrony in olfactory bulb glomeruli. *Neuron, 46*(5), 761-772.
doi:10.1016/j.neuron.2005.04.030
Coates, K. E., Calle-Schuler, S. A., Helmick, L. M., Knotts, V. L., Martik, B. N., Salman, F., . . . Dacks,
A. M. (2020). The Wiring Logic of an Identified Serotonergic Neuron That Spans Sensory
Networks. *The Journal of Neuroscience, 40*(33), 6309-6327. doi:10.1523/jneurosci.0552-
20.2020
Couto, A., Alenius, M., & Dickson, B. J. (2005). Molecular, anatomical, and functional organization
of the *Drosophila* olfactory system. *Current Biology, 15*(17), 1535-1547.
doi:10.1016/j.cub.2005.07.034
Cover, K. K., & Mathur, B. N. (2021). Axo-axonic synapses: Diversity in neural circuit function.
*Journal of Comparative Neurology, 529*(9), 2391-2401. doi:10.1002/cne.25087

Croset, V., Treiber, C. D., & Waddell, S. (2018). Cellular diversity in the Drosophila midbrain
974           revealed by single-cell transcriptomics. *Elife, 7*. doi:10.7554/eLife.34550

975    Cuntz, H., Borst, A., & Segev, I. (2007). Optimization principles of dendritic structure. *Theor Biol*
976           *Med Model, 4*, 21. doi:10.1186/1742-4682-4-21

977    Dacks, A. M., Christensen, T. A., & Hildebrand, J. G. (2006). Phylogeny of a serotonin-
978           immunoreactive neuron in the primary olfactory center of the insect brain. *Journal of*
979           *Comparative Neurology, 498*(6), 727-746. doi:10.1002/cne.21076

980    Dalal, T., Gupta, N., & Haddad, R. (2020). Bilateral and unilateral odor processing and odor
981           perception. *Commun Biol, 3*(1), 150. doi:10.1038/s42003-020-0876-6

982    Datta, S. R., Vasconcelos, M. L., Ruta, V., Luo, S., Wong, A., Demir, E., . . . Axel, R. (2008). The
983           *Drosophila* pheromone cVA activates a sexually dimorphic neural circuit. *Nature,*
984           *452*(7186), 473-477. doi:10.1038/nature06808

985    de Bruyne, M., Clyne, P. J., & Carlson, J. R. (1999). Odor coding in a model olfactory organ: the
986           *Drosophila* maxillary palp. *Journal of Neuroscience, 19*(11), 4520-4532.

987    de Bruyne, M., Foster, K., & Carlson, J. R. (2001). Odor Coding in the *Drosophila* Antenna. *Neuron,*
988           *30*(2), 537-552. doi:10.1016/S0896-6273(01)00289-6

989    de la Rocha, J., Doiron, B., Shea-Brown, E., Josić, K., & Reyes, A. (2007). Correlation between
990           neural spike trains increases with firing rate. *Nature, 448*(7155), 802-806.
991           doi:10.1038/nature06028

992    Demir, M., Kadakia, N., Anderson, H. D., Clark, D. A., & Emonet, T. (2020). Walking Drosophila
993           navigate complex plumes using stochastic decisions biased by the timing of odor
994           encounters. *Elife, 9*. doi:10.7554/eLife.57524

995    Dolan, M.-J., Frechter, S., Bates, A. S., Dan, C., Huoviala, P., Roberts, R. J. V., . . . Jefferis, G. S. X. E.
996           (2019). Neurogenetic dissection of the Drosophila lateral horn reveals major outputs,
997           diverse behavioural functions, and interactions with the mushroom body. *Elife, 8*, e43079.
998           doi:10.7554/eLife.43079

Dolan, M. J., Belliart-Guérin, G., Bates, A. S., Frechter, S., Lampin-Saint-Amaux, A., Aso, Y., . . .
Jefferis, G. (2018). Communication from Learned to Innate Olfactory Processing Centers Is
Required for Memory Retrieval in Drosophila. *Neuron, 100*(3), 651-668.e658.
doi:10.1016/j.neuron.2018.08.037

Duistermars, B. J., Chow, D. M., & Frye, M. A. (2009). Flies require bilateral sensory input to track
odor gradients in flight. *Current Biology, 19*(15), 1301-1307.

Dweck, H. K. M., Ebrahim, S. A. M., Thoma, M., Mohamed, A. A. M., Keesey, I. W., Trona, F., . . .
Hansson, B. S. (2015). Pheromones mediating copulation and attraction in *Drosophila*.
*Proceedings of the National Academy of Sciences, 112*(21), E2829-E2835.
doi:10.1073/pnas.1504527112

Ebrahim, S. A. M., Dweck, H. K. M., Stökl, J., Hofferberth, J. E., Trona, F., Weniger, K., . . . Knaden,
M. (2015). *Drosophila* Avoids Parasitoids by Sensing Their Semiochemicals via a Dedicated
Olfactory Circuit. *PLoS Biology, 13*(12), e1002318. doi:10.1371/journal.pbio.1002318

Eckstein, N., Bates, A. S., Du, M., Hartenstein, V., Jefferis, G. S. X. E., & Funke, J. (2020).
Neurotransmitter Classification from Electron Microscopy Images at Synaptic Sites in
Drosophila. *bioRxiv*, 2020.2006.2012.148775. doi:10.1101/2020.06.12.148775

Eichler, K., Litwin-Kumar, A., Li, F., Park, Y., Andrade, I., Schneider-Mizell, C. M., . . . Cardona, A. (2017). The Complete Connectome Of A Learning And Memory Center In An Insect Brain. *bioRxiv*. doi:10.1101/141762

Felsenberg, J., Jacob, P. F., Walker, T., Barnstedt, O., Edmondson-Stait, A. J., Pleijzier, M. W., . . . Waddell, S. (2018). Integration of Parallel Opposing Memories Underlies Memory Extinction. *Cell, 175*(3), 709-722.e715. doi:https://doi.org/10.1016/j.cell.2018.08.021

Fiala, A. (2007). Olfaction and olfactory learning in Drosophila: recent progress. *Current Opinion in Neurobiology, 17*(6), 720-726. doi:10.1016/j.conb.2007.11.009

Fishilevich, E., & Vosshall, L. B. (2005). Genetic and functional subdivision of the *Drosophila* antennal lobe. *Current Biology, 15*(17), 1548-1553. doi:10.1016/j.cub.2005.07.066

Fröhlich, A. (1985). Freeze-Fracture Study of an Invertebrate Multiple-Contact Synapse: The Fly Photoreceptor Tetrad. *Journal of Comparative Neurology, 241*(3), 311-326. doi:10.1002/cne.902410306

Galizia, C. G. (2014). Olfactory coding in the insect brain: data and conjectures. *European Journal of Neuroscience, 39*(11), n/a-n/a. doi:10.1111/ejn.12558

Galizia, C. G., Nagler, K., Holldobler, B., & Menzel, R. (1998). Odour coding is bilaterally symmetrical in the antennal lobes of honeybees (Apis mellifera). *European Journal of Neuroscience, 10*(9), 2964-2974.

Gao, Q., Yuan, B., & Chess, A. (2000). Convergent projections of Drosophila olfactory neurons to specific glomeruli in the antennal lobe. *Nature*.

Gao, X. J., Clandinin, T. R., & Luo, L. (2015). Extremely Sparse Olfactory Inputs Are Sufficient to Mediate Innate Aversion in *Drosophila PLoS ONE, 10*(4), e0125986. doi:10.1371/journal.pone.0125986

Gaudry, Q., Hong, E. J., Kain, J., de Bivort, B. L., & Wilson, R. I. (2013). Asymmetric neurotransmitter release enables rapid odour lateralization in *Drosophila*. *Nature, 493*(7432), 424-428. doi:10.1038/nature11747

Gondré-Lewis, M. C., Park, J. J., & Loh, Y. P. (2012). Chapter Two - Cellular Mechanisms for the Biogenesis and Transport of Synaptic and Dense-Core Vesicles. In K. W. Jeon (Ed.), *International Review of Cell and Molecular Biology* (Vol. 299, pp. 27-115): Academic Press.

Goyal, R. K., & Chaudhury, A. (2013). Structure activity relationship of synaptic and junctional neurotransmission. *Autonomic Neuroscience: Basic and Clinical, 176*(1), 11-31. doi:10.1016/j.autneu.2013.02.012

Grabe, V., Baschwitz, A., Dweck, Hany K. M., Lavista-Llanos, S., Hansson, Bill S., & Sachse, S. (2016). Elucidating the Neuronal Architecture of Olfactory Glomeruli in the *Drosophila* Antennal Lobe. *Cell Reports, 16*(12), 3401-3413. doi:10.1016/j.celrep.2016.08.063

Grabe, V., & Sachse, S. (2018). Fundamental principles of the olfactory code. *BioSystems, 164*, 94-101. doi:10.1016/j.biosystems.2017.10.010

Grabe, V., Schubert, M., Strube-Bloss, M., Reinert, A., Trautheim, S., Lavista-Llanos, S., . . . Sachse, S. (2020). Odor-Induced Multi-Level Inhibitory Maps in Drosophila. *eNeuro, 7*(1). doi:10.1523/eneuro.0213-19.2019

Grabe, V., Strutz, A., Baschwitz, A., Hansson, B. S., & Sachse, S. (2015). Digital in vivo 3D atlas of the antennal lobe of *Drosophila melanogaster*. *Journal of Comparative Neurology, 523*(3), 530-544. doi:10.1002/cne.23697

Graubard, K., Raper, J. A., & Hartline, D. K. (1980). Graded synaptic transmission between spiking
neurons. *Proc Natl Acad Sci U S A, 77*(6), 3733-3735. doi:10.1073/pnas.77.6.3733
Gruber, L., Rybak, J., Hansson, B. S., & Cantera, R. (2018). Synaptic Spinules in the Olfactory
Circuit of Drosophila melanogaster. *Front Cell Neurosci, 12*(86), 86.
doi:10.3389/fncel.2018.00086
Guo, D., Wu, S., Chen, M., Perc, M., Zhang, Y., Ma, J., . . . Yao, D. (2016). Regulation of Irregular
Neuronal Firing by Autaptic Transmission. *Sci Rep, 6*, 26096. doi:10.1038/srep26096
Guven-Ozkan, T., & Davis, R. L. (2014). Functional neuroanatomy of *Drosophila* olfactory memory
formation. *Learning & Memory, 21*(10), 519-526. doi:10.1101/lm.034363.114
Hallem, E. A., & Carlson, J. R. (2006). Coding of odors by a receptor repertoire. *Cell, 125*(1), 143-
160. doi:10.1016/j.cell.2006.01.050
Hallem, E. A., Ho, M. G., & Carlson, J. R. (2004). The molecular basis of odor coding in the
Drosophila antenna. *Cell, 117*(7), 965-979. doi:10.1016/j.cell.2004.05.012
Hansson, B. S., & Anton, S. (2000). Function and morphology of the antennal lobe: new
developments. *Annual Review of Entomology, 45*, 203-231.
doi:10.1146/annurev.ento.45.1.203
Hartenstein, V. (2016). The Central Nervous System of Invertebrates. In S. V. Shepherd (Ed.), *The
Wiley Handbook of Evolutionary Neuroscience* (pp. 173-235).
Haverkamp, A., Hansson, B. S., & Knaden, M. (2018). Combinatorial Codes and Labeled Lines:
How Insects Use Olfactory Cues to Find and Judge Food, Mates, and Oviposition Sites in
Complex Environments. *Front Physiol, 9*, 49. doi:10.3389/fphys.2018.00049
Helmstaedter, M. (2013). Cellular-resolution connectomics: challenges of dense neural circuit
reconstruction. *Nature Methods, 10*(6), 501-507. doi:10.1038/nmeth.2476
Hirata, Y. (1964). Some observations on the fine structure of the synapses in the olfactory bulb of
the mouse, with particular reference to the atypical synaptic configurations. *Archivum
histologicum japonicum, 24*(3), 293-302.
Hong, E. J., & Wilson, R. I. (2015). Simultaneous encoding of odors by channels with diverse
sensitivity to inhibition. *Neuron, 85*(3), 573-589. doi:10.1016/j.neuron.2014.12.040
Horne, J. A., Langille, C., McLin, S., Wiederman, M., Lu, Z., Xu, C. S., . . . Meinertzhagen, I. A.
(2018). A resource for the *Drosophila* antennal lobe provided by the connectome of
glomerulus VA1v. *Elife, 7*, e37550. doi:10.7554/eLife.37550
Huang, G. B., Scheffer, L. K., & Plaza, S. M. (2018). Fully-Automatic Synapse Prediction and
Validation on a Large Data Set. *Frontiers in Neural Circuits, 12*(87).
doi:10.3389/fncir.2018.00087
Hulse, B. K., Haberkern, H., Franconville, R., Turner-Evans, D. B., Takemura, S.-y., Wolff, T., . . .
Jayaraman, V. (2021). A connectome of the Drosophila central complex reveals network
motifs suitable for flexible navigation and context-dependent action selection. *Elife, 10*,
e66039. doi:10.7554/eLife.66039
Huoviala, P., Dolan, M.-J., Love, F. M., Myers, P., Frechter, S., Namiki, S., . . . Jefferis, G. S. X. E.
(2020). Neural circuit basis of aversive odour processing in *Drosophila* from sensory input
to descending output. 394403. doi:10.1101/394403 %J bioRxiv
Ikeda, K., & Bekkers, J. M. Autapses. *Current Biology, 16*(9), R308. doi:10.1016/j.cub.2006.03.085
Ikeda, K., & Bekkers, J. M. (2006). Autapses. *Current Biology, 16*(9), R308.
doi:10.1016/j.cub.2006.03.085

Jeanne, James M., & Wilson, Rachel I. (2015). Convergence, Divergence, and Reconvergence in a
Feedforward Network Improves Neural Speed and Accuracy. *Neuron, 88*(5), 1014-1026.
doi:10.1016/j.neuron.2015.10.018
Jefferis, G. S., Potter, C. J., Chan, A. M., Marin, E. C., Rohlfing, T., Maurer, C. R., Jr., & Luo, L.
(2007). Comprehensive maps of *Drosophila* higher olfactory centers: spatially segregated
fruit and pheromone representation. *Cell, 128*(6), 1187-1203.
doi:10.1016/j.cell.2007.01.040
Karnovsky, M. J. (1965). A formaldehyde-glutaraldehyde fixative of high osmolarity for use in
electron microscopy. *Journal of Cellular Biology, 27*, 137.
Kazama, H., & Wilson, R. I. (2008). Homeostatic Matching and Nonlinear Amplification at
Identified Central Synapses. *Neuron, 58*(3), 401-413.
doi:http://dx.doi.org/10.1016/j.neuron.2008.02.030
Kazama, H., & Wilson, R. I. (2009). Origins of correlated activity in an olfactory circuit. *Nature*
*Neuroscience, 12*(9), 1136-1144. doi:10.1038/nn.2376
Keene, A. C., & Waddell, S. (2007). Drosophila olfactory memory: single genes to complex neural
circuits. *Nature Reviews: Neuroscience, 8*(5), 341-354.
Keesey, I. W., & Hansson, B. S. (2021). 10 - The neuroethology of labeled lines in insect olfactory
systems. In G. J. Blomquist & R. G. Vogt (Eds.), *Insect Pheromone Biochemistry and*
*Molecular Biology (Second Edition)* (pp. 285-327). London: Academic Press.
Keesey, I. W., Zhang, J., Depetris-Chauvin, A., Obiero, G. F., Knaden, M., & Hansson, B. S. (2019).
Evolution of a pest: towards the complete neuroethology of <em>Drosophila
suzukii</em> and the subgenus <em>Sophophora</em>. *bioRxiv*, 717322.
doi:10.1101/717322
Knaden, M., & Hansson, B. S. (2014). Mapping odor valence in the brain of flies and mice. *Current*
*Opinion in Neurobiology, 24*(1), 34-38. doi:10.1016/j.conb.2013.08.010
Knaden, M., Strutz, A., Ahsan, J., Sachse, S., & Hansson, B. S. (2012). Spatial representation of
odorant valence in an insect brain. *Cell Reports, 1*, 392-399.
doi:10.1016/j.celrep.2012.03.002
Knott, G., Marchman, H., & Lich, B. (2008). Serial Section Scanning Electron Microscopy of Adult
Brain Tissue Using Focused Ion Beam Milling. *Journal of Neuroscience, 28*(12), 2964-2959.
doi:10.1523/JNEUROSCI.3189-07.2008
Kreher, S. A., Mathew, D., Kim, J., & Carlson, J. R. (2008). Translation of sensory input into
behavioral output via an olfactory system. *Neuron, 59*(1), 110-124.
doi:10.1016/j.neuron.2008.06.010
Kumar, A., Schiff, O., Barkai, E., Mel, B. W., Poleg-Polsky, A., & Schiller, J. (2018). NMDA spikes
mediate amplification of inputs in the rat piriform cortex. *Elife, 7*, e38446.
doi:10.7554/eLife.38446
Kurtovic, A., Widmer, A., & Dickson, B. J. (2007). A single class of olfactory neurons mediates
behavioural responses to a Drosophila sex pheromone. *Nature, 446*(7135), 542-546.
doi:10.1038/nature05672
Laissue, P. P., Reiter, C., Hiesinger, P. R., Halter, S., Fischbach, K. F., & Stocker, R. F. (1999). Three-
dimensional reconstruction of the antennal lobe in *Drosophila melanogaster*. *The Journal*
*of Comparative Neurology, 405*(4), 543-552. doi:10.1002/(sici)1096-
9861(19990322)405:4<543::aid-cne7>3.0.co;2-a

Laissue, P. P., & Vosshall, L. B. (2008). The Olfactory Sensory Map in *Drosophila*. In G. M. Technau (Ed.), *Brain Development in Drosophila melanogaster* (pp. 102-114 ). New York: Springer.

Lei, H., Oland, L. A., Riffell, J. A., Beyerlein, A., & Hildebrand, J. G. (2010). Microcircuits for Olfactory Information Processing in the Antennal Lobe of *Manduca sexta*. In G. M. Shepherd & S. Grillner (Eds.), *Handbook of Brain Microcircuits* (pp. 417-426). New York: Oxford University Press.

Li, F., Lindsey, J. W., Marin, E. C., Otto, N., Dreher, M., Dempsey, G., . . . Rubin, G. M. (2020). The connectome of the adult Drosophila mushroom body provides insights into function. *Elife, 9*. doi:10.7554/eLife.62576

Li, H., Horns, F., Wu, B., Xie, Q., Li, J., Li, T., . . . Luo, L. (2017). Classifying Drosophila Olfactory Projection Neuron Subtypes by Single-Cell RNA Sequencing. *Cell, 171*(5), 1206-1220.e1222. doi:https://doi.org/10.1016/j.cell.2017.10.019

Li, P. H., Lindsey, L. F., Januszewski, M., Zheng, Z., Bates, A. S., Taisz, I., . . . Jain, V. (2020). Automated Reconstruction of a Serial-Section EM *Drosophila* Brain with Flood-Filling Networks and Local Realignment. *bioRxiv*, 605634. doi:10.1101/605634

Liang, L., & Luo, L. (2010). The olfactory circuit of the fruit fly *Drosophila melanogaster*. *Sci China Life Sci, 53*(4), 472-484. doi:10.1007/s11427-010-0099-z

Lin, S., Kao, C.-F., Yu, H.-H., Huang, Y., & Lee, T. (2012). Lineage analysis of Drosophila lateral antennal lobe neurons reveals notch-dependent binary temporal fate decisions. *PLoS Biology, 10*(11), e1001425-e1001425. doi:10.1371/journal.pbio.1001425

Liu, T. X., Davoudian, P. A., Lizbinski, K. M., & Jeanne, J. M. (2022). Connectomic features underlying diverse synaptic connection strengths and subcellular computation. *Current Biology, 32*(3), 559-569.e555. doi:https://doi.org/10.1016/j.cub.2021.11.056

Liu, W. W., & Wilson, R. I. (2013). Glutamate is an inhibitory neurotransmitter in the *Drosophila* olfactory system. *Proceedings of the National Academy of Sciences, 110*(25), 10294–10299. doi:10.1073/pnas.1220560110

Malun, D. (1991). Inventory and distribution of synapses of identified uniglomerular projection neurons in the antennal lobe of Periplaneta americana. *Journal of Comparative Neurology, 305*(2), 348-360. doi:10.1002/cne.903050215

Malun, D., Waldow, U., Kraus, D., & Boeckh, J. (1993). Connections between the Deutocerebrum and the Protocerebrum, and Neuroanatomy of Several Classes of Deutocerebral Projection Neurons in the Brain of Male Periplaneta-Americana. *Journal of Comparative Neurology, 329*(2), 143-162.

Marin, E. C., Büld, L., Theiss, M., Sarkissian, T., Roberts, R. J. V., Turnbull, R., . . . Jefferis, G. (2020). Connectomics Analysis Reveals First-, Second-, and Third-Order Thermosensory and Hygrosensory Neurons in the Adult Drosophila Brain. *Current Biology, 30*(16), 3167-3182.e3164. doi:10.1016/j.cub.2020.06.028

Masse, N. Y., Turner, G. C., & Jefferis, G. S. (2009). Olfactory information processing in Drosophila. *Current Biology, 19*(16), R700-713. doi:10.1016/j.cub.2009.06.026

Masson, C., & Mustaparta, H. (1990). Chemical information processing in the olfactory system of insects. *Physiological Review, 70(1)*, 199-245.

McTavish, T., Migliore, M., Shepherd, G., & Hines, M. (2012). Mitral cell spike synchrony modulated by dendrodendritic synapse location. *Frontiers in Computational Neuroscience, 6*. doi:10.3389/fncom.2012.00003

Meinertzhagen, I. A. (2018). Of what use is connectomics? A personal perspective on the
*Drosophila* connectome. *Journal of Experimental Biology, 221*(10), jeb164954.
doi:10.1242/jeb.164954 %J The Journal of Experimental Biology

Meinertzhagen, I. A., & O'Neil, S. D. (1991). Synaptic organization of columnar elements in the
lamina of the wild type in Drosophila melanogaster. *Journal of Comparative Neurology,*
*305*, 232-263.

Miroschnikow, A., Schlegel, P., Schoofs, A., Hueckesfeld, S., Li, F., Schneider-Mizell, C. M., . . .
Pankratz, M. J. (2018). Convergence of monosynaptic and polysynaptic sensory paths onto
common motor outputs in a Drosophila feeding connectome. *Elife, 7*, e40247.
doi:10.7554/eLife.40247

Mohamed, A. A. M., Hansson, B. S., & Sachse, S. (2019a). Third-Order Neurons in the Lateral Horn
Enhance Bilateral Contrast of Odor Inputs Through Contralateral Inhibition in Drosophila.
*Front Physiol, 10*, 851. doi:10.3389/fphys.2019.00851

Mohamed, A. A. M., Retzke, T., Das Chakraborty, S., Fabian, B., Hansson, B. S., Knaden, M., &
Sachse, S. (2019b). Odor mixtures of opposing valence unveil inter-glomerular crosstalk in
the Drosophila antennal lobe. *Nat Commun, 10*(1), 1201. doi:10.1038/s41467-019-09069-
1

Mosca, T. J., & Luo, L. (2014). Synaptic organization of the *Drosophila* antennal lobe and its
regulation by the Teneurins. *Elife, 3*(3), e03726. doi:10.7554/eLife.03726

Münch, D., & Galizia, C. G. (2016). DoOR 2.0 - Comprehensive Mapping of Drosophila
melanogaster Odorant Responses. *Scientific Reports, 6*, 21841. doi:10.1038/srep21841
http://www.nature.com/articles/srep21841#supplementary-information

Nässel, D. R., Enell, L. E., Santos, J. G., Wegener, C., & Johard, H. A. (2008). A large population of
diverse neurons in the Drosophila central nervous system expresses short neuropeptide F,
suggesting multiple distributed peptide functions. *BMC Neuroscience, 9*, 90.
doi:10.1186/1471-2202-9-90

Nässel, D. R., & Homberg, U. (2006). Neuropeptides in interneurons of the insect brain. *Cell and*
*Tissue Research, 326*(1), 1-24. doi:10.1007/s00441-006-0210-8

Ng, M., Roorda, R. D., Lima, S. Q., Zemelman, B. V., Morcillo, P., & Miesenböck, G. (2002).
Transmission of Olfactory Information between Three Populations of Neurons in the
Antennal Lobe of the Fly. *Neuron, 36*(3), 463-474. doi:10.1016/s0896-6273(02)00975-3

Norgate, M., Lee, E., Southon, A., Farlow, A., Batterham, P., Camakaris, J., & Burke, R. (2006).
Essential Roles in Development and Pigmentation for the *Drosophila* Copper Transporter
DmATP7. *Molecular Biology of the Cell, 17*(1), 475-484. doi:10.1091/mbc.E05-06-0492

Okada, R., Awasaki, T., & Ito, K. (2009). Gamma-aminobutyric acid (GABA)-mediated neural
connections in the *Drosophila* antennal lobe. *Journal of Comparative Neurology, 514*(1),
74-91. doi:10.1002/cne.21971

Olsen, S. R., & Wilson, R. I. (2008). Lateral presynaptic inhibition mediates gain control in an
olfactory circuit. *Nature, 452*(7190), 956-960. doi:10.1038/nature06864

Otto, N., Pleijzier, M. W., Morgan, I. C., Edmondson-Stait, A. J., Heinz, K. J., Stark, I., . . . Waddell,
S. (2020). Input Connectivity Reveals Additional Heterogeneity of Dopaminergic
Reinforcement in Drosophila. *Current Biology, 30*(16), 3200-3211.e3208.
doi:https://doi.org/10.1016/j.cub.2020.05.077

Owald, D., & Waddell, S. (2015). Olfactory learning skews mushroom body output pathways to
steer behavioral choice in Drosophila. *Current Opinion in Neurobiology, 35*, 178-184.
doi:10.1016/j.conb.2015.10.002
Parthasarathy, K., & Bhalla, U. S. (2013). Laterality and symmetry in rat olfactory behavior and in
physiology of olfactory input. *Journal of Neuroscience, 33*(13), 5750-5760.
doi:10.1523/jneurosci.1781-12.2013
Prokop, A., & Meinertzhagen, I. A. (2006). Development and structure of synaptic contacts in
Drosophila. *Seminars in Cell & Developmental Biology, 17*(1), 20-30.
doi:10.1016/j.semcdb.2005.11.010
Ronchi, P., Mizzon, G., Machado, P., D'Imprima, E., Best, B. T., Cassella, L., . . . Schwab, Y. (2021).
High-precision targeting workflow for volume electron microscopy. *Journal of Cell Biology,*
*220*(9). doi:10.1083/jcb.202104069
Root, C. M., Masuyama, K., Green, D. S., Enell, L. E., Nassel, D. R., Lee, C. H., & Wang, J. W. (2008).
A presynaptic gain control mechanism fine-tunes olfactory behavior. *Neuron, 59*(2), 311-
321. doi:10.1016/j.neuron.2008.07.003
Rospars, J.-P., Grémiaux, A., Jarriault, D., Chaffiol, A., Monsempes, C., Deisig, N., . . . Martinez, D.
(2014). Heterogeneity and Convergence of Olfactory First-Order Neurons Account for the
High Speed and Sensitivity of Second-Order Neurons. *PLoS Computational Biology, 10*(12),
e1003975. doi:10.1371/journal.pcbi.1003975
Rozenfeld, E., Lerner, H., & Parnas, M. (2019). Muscarinic Modulation of Antennal Lobe
GABAergic Local Neurons Shapes Odor Coding and Behavior. *Cell Rep, 29*(10), 3253-
3265.e3254. doi:10.1016/j.celrep.2019.10.125
Rybak, J. (2013). Exploring Brain Connectivity in Insect Model Systems of Learning and Memory.
In R. Menzel & P. Benjamin (Eds.), *Invertebrate Learning and Memory* (pp. 26-40). San
Diego: Academic Press.
Rybak, J. (2016). Perspective-Brain atlases for studying neuronal circuitry in arthropods. In A.
Schmidt-Rhaesa, S. Harzsch, & G. Purschke (Eds.), *Structure and Evolution of Invertebrate*
*Nervous Systems* (Vol. 1). New York: Oxford University Press.
Rybak, J., & Hansson, B. S. (2018). Olfactory Microcircuits in *Drosophila Melanogaster*. In G. M.
Shepherd & S. Grillner (Eds.), *Handbook of Brain Microcircuits* (2nd ed., pp. 361-367).
Oxford, UK: Oxford University Press.
Rybak, J., Talarico, G., Ruiz, S., Arnold, C., Cantera, R., & Hansson, B. S. (2016). Synaptic circuitry
of identified neurons in the antennal lobe of *Drosophila melanogaster*. *Journal of*
*Comparative Neurology, 524*(9), 1920-1956. doi:10.1002/cne.23966
Saada, R., Miller, N., Hurwitz, I., & Susswein, A. J. (2009). Autaptic excitation elicits persistent
activity and a plateau potential in a neuron of known behavioral function. *Current Biology,*
*19*(6), 479-484. doi:10.1016/j.cub.2009.01.060
Saalfeld, S., Cardona, A., Hartenstein, V., & Tomančák, P. (2009). CATMAID: collaborative
annotation toolkit for massive amounts of image data. *Bioinformatics, 25*(15), 1984-1986.
doi:10.1093/bioinformatics/btp266
Sachse, S., & Galizia, C. G. (2002). Role of inhibition for temporal and spatial odor representation
in olfactory output neurons: a calcium imaging study. *Journal of Neurophysiology, 87*(2),
1106-1117.

Sachse, S., & Hansson, B. S. (2016). Research spotlight: Olfactory coding in *Drosophila*
*melanogaster*. In A. Schmidt-Rhaesa, S. Harzsch, & G. Purschke (Eds.), *Structure and*
*Evolution of Invertebrate Nervous Systems* (pp. 640-645). Oxford: Oxford University Press.
Sachse, S., & Manzini, I. (2021). Editorial for the special issue "Olfactory Coding and Circuitries".
*Cell and Tissue Research, 383*(1), 1-6. doi:10.1007/s00441-020-03389-1
Scheffer, L. K., Xu, C. S., Januszewski, M., Lu, Z., Takemura, S.-y., Hayworth, K. J., . . . Plaza, S. M.
(2020). A connectome and analysis of the adult Drosophila central brain. *Elife, 9*, e57443.
doi:10.7554/eLife.57443
Schlegel, P., Bates, A. S., Stürner, T., Jagannathan, S. R., Drummond, N., Hsu, J., . . . Jefferis, G. S.
X. E. (2021). Information flow, cell types and stereotypy in a full olfactory connectome.
*Elife, 10*, e66018. doi:10.7554/eLife.66018
Schneider-Mizell, C. M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M. F., . . . Cardona, A.
(2016). Quantitative neuroanatomy for connectomics in *Drosophila*. *Elife, 5*, e12059.
doi:10.7554/eLife.12059
Seki, Y., Dweck, H. K. M., Rybak, J., Wicher, D., Sachse, S., & Hansson, B. S. (2017). Olfactory
coding from the periphery to higher brain centers in the Drosophila brain. *BMC Biology,*
*15*(1), 56. doi:10.1186/s12915-017-0389-z
Seki, Y., Rybak, J., Wicher, D., Sachse, S., & Hansson, B. S. (2010). Physiological and morphological
characterization of local interneurons in the *Drosophila* antennal lobe. *Journal of*
*Neurophysiology, 104*(2), 1007-1019. doi:jn.00249.2010 [pii]
10.1152/jn.00249.2010
Semmelhack, J. L., & Wang, J. W. (2009). Select Drosophila glomeruli mediate innate olfactory
attraction and aversion. *Nature, 459*(7244), 218-223.
Shanbhag, S. R., Muller, B., & Steinbrecht, R. A. (1999). Atlas of olfactory organs of *Drosophila*
*melanogaster* - 1. Types, external organization, innervation and distribution of olfactory
sensilla. *International Journal of Insect Morphology & Embryology, 28*(4), 377-397. doi:Doi
10.1016/S0020-7322(99)00039-2
Shepherd, G. M. (2011). The Olfactory Bulb: A Simple System in the Mammalian Brain
*Comprehensive Physiology*: John Wiley & Sons, Inc.
Shepherd, G. M., Rowe, T. B., & Greer, C. A. (2021). An Evolutionary Microcircuit Approach to the
Neural Basis of High Dimensional Sensory Processing in Olfaction. *Frontiers in Cellular*
*Neuroscience, 15*. doi:10.3389/fncel.2021.658480
Silbering, A. F., & Galizia, C. G. (2007). Processing of odor mixtures in the Drosophila antennal
lobe reveals both global inhibition and glomerulus-specific interactions. *Journal of*
*Neuroscience, 27*(44), 11966-11977. doi:10.1523/jneurosci.3099-07.2007
Silbering, A. F., Okada, R., Ito, K., & Galizia, C. G. (2008). Olfactory information processing in the
*Drosophila* antennal lobe: anything goes? *Journal of Neuroscience, 28*(49), 13075-13087.
doi:10.1523/JNEUROSCI.2973-08.2008
Silbering, A. F., Rytz, R., Grosjean, Y., Abuin, L., Ramdya, P., Jefferis, G. S., & Benton, R. (2011).
Complementary Function and Integrated Wiring of the Evolutionarily Distinct *Drosophila*
Olfactory Subsystems. *Journal of Neuroscience, 31*(38), 13357-13375.
doi:10.1523/JNEUROSCI.2360-11.2011

Stensmyr, M. C., Dweck, H. K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., . . . Hansson, B. S. (2012). A conserved dedicated olfactory circuit for detecting harmful microbes in *Drosophila*. *Cell, 151*(6), 1345-1357. doi:10.1016/j.cell.2012.09.046

Stocker, R. F., Lienhard, M. C., Borst, A., & Fischbach, K. F. (1990). Neuronal architecture of the antennal lobe in Drosophila melanogaster. *Cell and Tissue Research, 262*(1), 9-34.

Stocker, R. F., Singh, R. N., Schorderet, M., & Siddiqi, O. (1983). Projection patterns of different types of antennal sensilla in the antennal glomeruli of Drosophila melanogaster. *Cell and Tissue Research, 232*(2), 237-248. doi:10.1007/bf00213783

Strutz, A., Soelter, J., Baschwitz, A., Farhan, A., Grabe, V., Rybak, J., . . . Sachse, S. (2014). Decoding odor quality and intensity in the *Drosophila* brain. *Elife, 3*, e04147. doi:10.7554/eLife.04147

Su, C.-Y., Menuz, K., & Carlson, J. R. (2009). Olfactory Perception: Receptors, Cells, and Circuits. *Cell, 139*(1), 45-59. doi:10.1016/j.cell.2009.09.015

Suh, G. S. B., Wong, A. M., Hergarden, A. C., Wang, J. W., Simon, A. F., Benzer, S., . . . Anderson, D. J. (2004). A single population of olfactory sensory neurons mediates an innate avoidance behaviour in *Drosophila*. *Nature, 431*(7010), 854-859. doi:http://www.nature.com/nature/journal/v431/n7010/suppinfo/nature02980_S1.html

Sun, X. J., Tolbert, L. P., & Hildebrand, J. G. (1997). Synaptic organization of the uniglomerular projection neurons of the antennal lobe of the moth *Manduca sexta*: a laser scanning confocal and electron microscopic study. *Journal of Comparative Neurology, 379*(1), 2-20.

Szyszka, P., & Galizia, C. G. (2015). Olfaction in Insects. In R. L. Doty (Ed.), *Handbook of Olfaction and Gustation* (3 ed., pp. 531-546): John Wiley & Sons, Inc.

Taisz, I., Donà, E., Münch, D., Bailey, S. N., Morris, W. J., Meechan, K. I., . . . Galili, D. S. (2022). Generating parallel representations of position and identity in the olfactory system. *bioRxiv*, 2022.2005.2013.491877. doi:10.1101/2022.05.13.491877

Takemura, S.-y., Xu, C. S., Lu, Z., Rivlin, P. K., Parag, T., Olbris, D. J., . . . Scheffer, L. K. (2015). Synaptic circuits and their variations within different columns in the visual system of Drosophila. *Proceedings of the National Academy of Sciences, 112*(44), 13711-13716. doi:10.1073/pnas.1509820112

Tamás, G., Buhl, E. H., & Somogyi, P. (1997). Massive Autaptic Self-Innervation of GABAergic Neurons in Cat Visual Cortex. *The Journal of Neuroscience, 17*(16), 6352-6364. doi:10.1523/jneurosci.17-16-06352.1997

Tanaka, N. K., Endo, K., & Ito, K. (2012). The organization of antennal lobe-associated neurons in the adult *Drosophila* melanogaster brain. *Journal of Comparative Neurology, 520*(18), 4067-4130. doi:10.1002/cne.23142

Thoma, M., Hansson, B. S., & Knaden, M. (2015). High-resolution Quantification of Odor-guided Behavior in Drosophila melanogaster Using the Flywalk Paradigm. *J Vis Exp*(106), e53394. doi:10.3791/53394

Tobin, W. F., Wilson, R. I., & Lee, W.-C. A. (2017). Wiring variations that enable and constrain neural computation in a sensory microcircuit. *Elife, 6*, e24838. doi:10.7554/eLife.24838

Tran-Van-Minh, A., Cazé, R. D., Abrahamsson, T., Cathala, L., Gutkin, B. S., & DiGregorio, D. A. (2015). Contribution of sublinear and supralinear dendritic integration to neuronal computations. *Frontiers in Cellular Neuroscience, 9*. doi:10.3389/fncel.2015.00067

Trujillo-Cenoz, O. (1969). Some Aspects of the Structural Organization of the Medulla in Muscoid
Flies l. *J Ultrastructural Research, 27*, 533-553.
Van der Loos, H., & Glaser, E. M. (1972). Autapses in neocortex cerebri: synapses between a
pyramidal cell's axon and its own dendrites. *Brain Research, 48*, 355-360.
doi:10.1016/0006-8993(72)90189-8
Vosshall, L. B., Wong, A. M., & Axel, R. (2000). An olfactory sensory map in the fly brain. *Cell,
102*(2), 147-159. doi:10.1016/S0092-8674(00)00021-0
Wang, J. W. (2012). Presynaptic modulation of early olfactory processing in *Drosophila*.
*Developmental Neurobiology, 72*(1), 87-99. doi:10.1002/dneu.20936
Wicher, D., & Miazzi, F. (2021). Functional properties of insect olfactory receptors: ionotropic
receptors and odorant receptors. *Cell and Tissue Research, 383*(1), 7-19.
doi:10.1007/s00441-020-03363-x
Wiles, L., Gu, S., Pasqualetti, F., Parvesse, B., Gabrieli, D., Bassett, D. S., & Meaney, D. F. (2017).
Autaptic Connections Shift Network Excitability and Bursting. *Sci Rep, 7*, 44006.
doi:10.1038/srep44006
Wilson, R. I. (2013). Early Olfactory Processing in Drosophila: Mechanisms and Principles. *Annual
Review of Neuroscience, 36*, 217-241. doi:10.1146/annurev-neuro-062111-150533
Xu, C. S., Hayworth, K. J., Lu, Z., Grob, P., Hassan, A. M., García-Cerdán, J. G., . . . Hess, H. F.
(2017). Enhanced FIB-SEM systems for large-volume 3D imaging. *Elife, 6*, e25916.
doi:10.7554/eLife.25916
Xu, C. S., Januszewski, M., Lu, Z., Takemura, S.-y., Hayworth, K. J., Huang, G., . . . Plaza, S. M.
(2020). A Connectome of the Adult *Drosophila* Central Brain. 2020.2001.2021.911859.
doi:10.1101/2020.01.21.911859 %J bioRxiv
Yang, K., Liu, T., Wang, Z., Liu, J., Shen, Y., Pan, X., . . . Zhang, K. (2022). Classifying Drosophila
olfactory projection neuron boutons by quantitative analysis of electron microscopic
reconstruction. *iScience, 25*(5), 104180. doi:https://doi.org/10.1016/j.isci.2022.104180
Yasuyama, K., Meinertzhagen, I. A., & Schurmann, F. W. (2003). Synaptic connections of
cholinergic antennal lobe relay neurons innervating the lateral horn neuropile in the brain
of *Drosophila melanogaster*. *Journal of Comparative Neurology, 466*(3), 299-315.
doi:10.1002/cne.10867
Yasuyama, K., & Salvaterra, P. M. (1999). Localization of choline acetyltransferase-expressing
neurons in *Drosophila* nervous system. *Microscopy Research and Technique, 45*(2), 65-79.
Yin, L., Zheng, R., Ke, W., He, Q., Zhang, Y., Li, J., . . . Shu, Y. (2018). Autapses enhance bursting
and coincidence detection in neocortical pyramidal cells. *Nature Communications, 9*(1),
4890. doi:10.1038/s41467-018-07317-4
Yokoi, M., Mori, K., & Nakanishi, S. (1995). Refinement of odor molecule tuning by
dendrodendritic synaptic inhibition in the olfactory bulb. *Proceedings of the National
Academy of Sciences, 92*(8), 3371-3375. doi:doi:10.1073/pnas.92.8.3371
Zheng, Z., Lauritzen, J. S., Perlman, E., Robinson, C. G., Nichols, M., Milkie, D., . . . Bock, D. D.
(2017). A Complete Electron Microscopy Volume Of The Brain Of Adult Drosophila
melanogaster. *bioRxiv*. doi:10.1101/140905
Zheng, Z., Lauritzen, J. S., Perlman, E., Robinson, C. G., Nichols, M., Milkie, D., . . . Bock, D. D.
(2018). A Complete Electron Microscopy Volume of the Brain of Adult Drosophila
melanogaster. *Cell, 174*(3), 730-743.e722. doi:https://doi.org/10.1016/j.cell.2018.06.019

 **FIGURES**

 **Figure 1: A correlative approach to analyze the ultrastructure of identified olfactory glomeruli**

**A-B:** Two-photon laser scans of the antennal lobes in *Orco-Gal4; UAS-GCaMP6s* flies where Orco-
positive olfactory sensory neurons (OSNs) in the glomerular neuropils were labeled by GCaMP
(green fluorescence). Glomeruli DA2 (A) and DL5 (B) are encircled. Schematics show their relative
position in the antennal lobe. Once the glomeruli of interest were identified, glomerular borders
were marked with fiducial marks (white triangles) via laser branding, which enabled their
identification at the ultrastructural level. **C-D**: Representative images of the same glomeruli (DA2
in **C** and DL5 in **D**) obtained with focused-ion-beam electron microscopy (FIB-SEM), showing their
ultrastructure. Asterisks indicate the main neurite of uniglomerular projection neurons entering
the glomerulus. White triangle shows a 2-photon laser mark (see also **A** and **B**). **E**: FIB-SEM image
of a polyadic synapse: the presynaptic site (red arrowhead) is composed of a T-bar shaped
presynaptic density surrounded by small vesicles and is opposed by several postsynaptic profiles
(cyan dots). Scheme of a tetrad synapse: a presynaptic site with its T-bar (red arrowhead) forms
four output connections (arrows) with four postsynaptic input sites (cyan dots). **F:** A skeleton-based
reconstruction of an OSN axon terminal (green line) with presynaptic (red dots) and postsynaptic
sites (cyan dots). The dark grey shading surrounding the OSN trace represents the volume-based
reconstruction of the same neuron. Tracing and reconstruction were performed within the FIB-
SEM dataset (light grey area).
1424

**Figure 2: Neuron classification and neuronal composition of the DA2 and DL5 glomeruli**
**A**: Example FIB-SEM images (left column), volumetric neuronal reconstructions (middle column),
and skeleton-based neuron traces (right column) of a representative example of each neuron class:
OSNs (green), uniglomerular projection neurons (uPNs, red) and multiglomerular neurons (MGNs,
blue). The ultrastructure of neurons, including T-bars (black triangle), mitochondria (asterisks) and
spinules (white triangle) are indicated. Exemplar volumetric reconstructions (middle column) show
the general morphology of each neuron class. Presynapses and postsynapses are indicated with
red and cyan dots on the skeleton traces (right column). **B:** Average branching intensity (branching
points per μm of neuronal-fiber length) of each neuron class OSNs, uPNs and MGNs in DA2 and
DL5. Data represent mean+ standard deviation (error bars). Data points represent single values.
Means were compared using Wilcoxon two-sample test. No significant differences of branching
points/μm in OSNs or MGNs between glomeruli were found (significance was not tested for uPNs
due to the presence of a single uPN in DL5). **C**: Schematic summary indicating, for each glomerulus,
its volume (in $\mu m^3$), the number of neurons of each class (MGNs were not counted), the total fiber
length of all neurons for each neuron class and the total number of single synaptic contacts for
each glomerulus.
1441

**Table 1. Glomerular innervation and synaptic composition**
Quantitative neuronal data comparing glomeruli DA2 and DL5, detailing glomerular innervation
and synaptic properties for each neuronal class: OSNs (green), uPNs (red) and MGNs (blue) and the
sum of all of them. **Row 1**: Total length of all neurons of each neuron class and total length for all
neurons in each glomerulus. **Row 2-4**: Synaptic counts: input sites (inputs), output sites (outputs)

and T-bars (T-bars). **Row 5**: Innervation density: total neuron length (μm; row 1)/glomerular
volume (μm$^3$); glomerular volume: DA2=1500 μm$^3$ and DL5=2600 μm$^3$ (see Figure 1C). **Row 6-8**:
Total synaptic density per unit of glomerular volume (μm$^3$): sum of all input sites (inputs), output
sites (outputs) and T-bars of each neuron class or of all neurons/glomerular volume. **Row 9-11**:
Average synaptic density along neuronal fibers (illustrated also in Figure 3 – supplement 1):
number of inputs, outputs or T-bars/neuron length (μm). **Row 12-13:** Average synaptic ratios: the
ratio of T-bars-to-inputs or outputs-to-inputs. **Row 14:** Polyadicity: the average number of
postsynaptic sites at each T-bar in DA2 and DL5. The ratios in rows 12-14 were calculated based on
synaptic counts normalized to neuron length (rows 9-11). The color shading highlights values that
have a relative difference greater than 20% (see relative differences Table S1) between DA2 and
DL5. Dark shades highlights values that are greater in DA2 than in DL5 (green (OSNs), red (uPNs),
blue (MGNs)) and light colors highlight values that are less in DA2 than in DL5.
1459
**Figure 3: Innervation density and synaptic density in DA2 and DL5**
**A-E**: The average glomerular innervation density of OSNs (**A**), uPNs (**B**), MGNs (**C**) and collectively
of all glomerular neurons (**D**); the average synaptic density of input sites (inputs), output sites
(outputs) and T-bars and the average polyadicity. Innervation density: length (μm) of each
neuronal fiber normalized to one μm$^3$ of glomerular (glom.) volume. Synaptic density: number of
input sites, output sites or T-bars of each neuronal fiber normalized to one μm$^3$ of glomerular
volume. Polyadicity: average number of single output sites per T-bar in each neuronal fiber. Data
for DA2 shown in dark colors and for DL5 in light colors. Number of neurons in DA2: OSNs (green)
n= 44; uPNs (red) n= 7; MGNs (blue) n=180; all neurons n=231, in DL5: OSNs n=46; uPN n=1; MGNs
n=221; all neurons n=268. Data represent mean + standard deviation (error bars). Data points
represent single values. Means were compared using either Student's t-test (OSNs) or Wilcoxon
two-sample test (MGNs and all neurons). uPNs were not compared, since the DL5 has only one.
Significance value: p>0.05 (not significant, no star), p≤0.05 (*), p≤0.01 (**), p≤0.001 (***). Values
are provided at data availability; polyadicity values are listed in Table 1, row 14.
1474
**Figure 4: Lateralization of OSN terminals in the antennal lobes**
**A**: Illustration of an ipsilateral (dark green) and a contralateral (light green) OSN with dendrites in
the corresponding antennae and their axonal projections to the ipsilateral olfactory glomerulus in
the antennal lobe (AL) (dashed rectangle). **B**: Exemplary skeleton traces of an ipsilateral (dark
green) and a contralateral (light green) OSN terminal inside glomerulus DA2. The ipsilateral OSN
axons reach the glomerulus via the ipsilateral antennal nerve (arrow down) and leave the
glomerulus towards the AL commissure (arrow up) while OSN axons originating at the contralateral
antenna reach the glomerulus via the AL commissure. Red dots: presynapses; blue dots:
postsynapses. **C**: Boxplots showing the fraction of synaptic output to uPNs (in red), - to OSNs (in
green) or - toMGNs (in blue), , for the ipsilateral OSNs (dark green boxplot) and contralateral OSNs
(light green), respectively, in the DA2, DL5 and VA1v glomeruli (Horne *et al.*, 2018). **D**: Boxplots
showing the fraction of synaptic input of the same ipsilateral and contralateral OSNs that they
receive from OSNs and MGNs. Connection polarity is indicated by arrows in the schematic neuronal
drawings on the left of each plot. Dots represent single values. Means were compared using

Student's T-test. Significance value: p>0.05 (not significant, no star)), p≤0.05 (*), p≤0.01 (**),
p≤0.001 (***). Mean and Median values are provided at data availability.
1491

**Figure 5: Strength of synaptic connections between neuron classes in the circuitry of DA2, DL5**
**and VA1v.**
**A:** Schematic representation of principal connection motifs between the neuron classes OSNs
(green), uPNs (red) and MGNs (blue). The synaptic flow directed towards uPNs is a feedforward
and that directed towards OSNs or from uPNs to MGNs defined as a feedback connection (arrows).
**B-D**: Alluvial diagrams of the glomerular circuitry in DA2 (**B**), DL5 (**C**) and VA1v (**D**). Each diagram
shows the relative synaptic strength calculated as the proportion of 1:1 single synaptic contacts
between each neuron class in relation to the total number of synaptic contacts in their respective
glomerulus. The synaptic strength between each neuron class, given as percentage, is indicated by
the thickness of the lines. The proportions (as percentage) of output (left side) or input (right side)
are illustrated by colored rectangles to the left or right of each alluvial diagram. The total number
of synaptic contacts is indicated below the diagrams. Percentages of the relative synaptic strength
and synaptic counts are listed in the supplementary Table S1. **E:** Stacked bar charts depict output
(E') and input (E'') fractions (given as percentages) of each neuron class: OSNs (green), uPNs (red),
MGNs (blue), schematically illustrated next to the bar charts respectively, to each of the other
neuron classes for glomeruli DA2, DL5 and VA1v. Fractions are color-coded according to the neuron
class of the respective connecting partner.
1509

**Figure 6: Differences in connectivity strength in glomeruli DA2, DL5 and VA1v**
**A**: Schematic representation of synaptic connection motifs (arrows) between OSNs (green), uPNs
(red), and MGNs (blue) in glomeruli DA2, DL5 and VA1v. The number of neurons of each class or
truncated neuronal fibers (in brackets) is noted in the corresponding circle. **B**: Schematics of
connection motifs (left) that are jointly stronger or weaker in DA2 and VA1v than in DL5. The
relative differences (as percentage) between DA2 and DL5 as well as VA1v and DL5 are illustrated
as arrows up (stronger) or arrows down (weaker) according to their intensity (see legend at the
bottom) from the perspective of the target glomerulus (defined in the table header). The values of
relative differences are listed in the Table S2.
1519

**Figure 7: Distribution of pre- and postsynaptic partners of autapses in the uPN dendrite of the**
**DL5**
**A**: Distribution of autaptic presynaptic (red dots) and postsynaptic sites (cyan dots) mapped in a
dendrogram of the dendrite of the single uPN in glomerulus DL5. The basal root node (black dot)
represents the entry site of the uPN dendrite into the glomerulus (closest point to its soma).
Clustering of autaptic input sites along some branches are encircled. **B**: Simplified representation
of the uPN's dendrogram illustrating the distinct strahler orders, at distal branches (1-4) and at
basal branches (5-8); see legend on the right). **C**: Distribution of autaptic presynaptic (left) and
postsynaptic input sites (right) along the dendrite, as proportions at each corresponding strahler
order (color coded). Note that autaptic postsynaptic sites are located almost exclusively at the
most distal dendritic branches. **D**: Dendrogram of the DL5-uPN showing the distribution of
presynaptic sites (triangles) and postsynaptic sites (circles) of selected autapses (indicated by same
color). Distant pairs of pre- and postsynapses (long geodesic distance) are indicated by numbers

whereas closely attached synaptic sites (short geodesic distance) are encircled and labelled with letters. **E**: Schematic of the dendrogram illustrating the location of the presynaptic (red dot) and postsynaptic (cyan dot) sites of a single autapse, the geodesic distance between them, i.e. the distance along the dendrite (µm), and the number of branching points (orange dots) between the pre- and postsynaptic components of the same autapse. **F**: Number of autapses with distinct geodesic distances between their pre- and postsynapses (illustrated in E). **G**: Number of autapses with the number of branch points between their pre- and postsynapses counted along the uPN dendrite (illustrated in E).

Acebes, A., & Ferrus, A. (2001). Increasing the Number of Synapses Modifies Olfactory Perception in Drosophila. *Journal of Neuroscience, 21*(16), 6264–6273.

Agarwal, G., & Isacoff, E. (2011). Specializations of a pheromonal glomerulus in the Drosophila olfactory system. *Journal of Neurophysiology, 105*(4), 1711-1721. doi:10.1152/jn.00591.2010

Ai, H., & Hagio, H. (2013). Morphological analysis of the primary center receiving spatial information transferred by the waggle dance of honeybees. *Journal of Comparative Neurology, 521*(11), 2570-2584. doi:10.1002/cne.23299

Andersson, M. N., Löfstedt, C., & Newcomb, R. D. (2015). Insect olfaction and the evolution of receptor tuning. *Frontiers in Ecology and Evolution, 3*. doi:10.3389/fevo.2015.00053

Anton, S., van Loon, J. J., Meijerink, J., Smid, H. M., Takken, W., & Rospars, J. P. (2003). Central projections of olfactory receptor neurons from single antennal and palpal sensilla in mosquitoes. *Arthropod Structure and Development, 32*(4), 319-327. doi:10.1016/j.asd.2003.09.002

Asahina, K., Louis, M., Piccinotti, S., & Vosshall, L. B. (2009). A circuit supporting concentration-invariant odor perception in Drosophila. *Journal of Biology, 8*(1), 9. doi:10.1186/jbiol108

Assisi, C., Stopfer, M., & Bazhenov, M. (2012). Excitatory Local Interneurons Enhance Tuning of Sensory Information. *PLoS Computational Biology, 8*(7), e1002563. doi:10.1371/journal.pcbi.1002563

Auer, T. O., Khallaf, M. A., Silbering, A. F., Zappia, G., Ellis, K., Álvarez-Ocaña, R., . . . Benton, R. (2020). Olfactory receptor and circuit evolution promote host specialization. *Nature*. doi:10.1038/s41586-020-2073-7

Bacci, A., & Huguenard, J. R. (2006). Enhancement of spike-timing precision by autaptic transmission in neocortical inhibitory interneurons. *Neuron, 49*(1), 119-130. doi:10.1016/j.neuron.2005.12.014

Bates, A. S., Schlegel, P., Roberts, R. J. V., Drummond, N., Tamimi, I. F. M., Turnbull, R., . . . Jefferis, G. (2020). Complete Connectomic Reconstruction of Olfactory Projection Neurons in the Fly Brain. *Current Biology, 30*(16), 3183-3199 e3186. doi:10.1016/j.cub.2020.06.042

Bekkers, J. M. (1998). Neurophysiology: are autapses prodigal synapses? *Current Biology, 8*(2), R52-55. doi:10.1016/s0960-9822(98)70033-8

Bekkers, J. M. (2009). Synaptic Transmission: Excitatory Autapses Find a Function? *Current Biology, 19*(7), R296-R298. doi:http://dx.doi.org/10.1016/j.cub.2009.02.010

Benton, R., Sachse, S., Michnick, S. W., & Vosshall, L. B. (2006). Atypical Membrane Topology and Heteromeric Function of *Drosophila* Odorant Receptors In Vivo. *PLoS Biology, 4*(2), 240-257. doi:10.1371/journal.pbio.0040020

Berck, M. E., Khandelwal, A., Claus, L., Hernandez-Nunez, L., Si, G., Tabone, C. J., . . . Cardona, A. (2016). The wiring diagram of a glomerular olfactory system. *Elife, 5*, e14859. doi:10.7554/eLife.14859

Bhandawat, V., Olsen, S. R., Gouwens, N. W., Schlief, M. L., & Wilson, R. I. (2007). Sensory processing in the *Drosophila* antennal lobe increases reliability and separability of ensemble odor representations. *Nature Neuroscience, 10*(11), 1474-1482. doi:10.1038/nn1976

Bielopolski, N., Amin, H., Apostolopoulou, A. A., Rozenfeld, E., Lerner, H., Huetteroth, W., . . . Parnas, M. (2019). Inhibitory muscarinic acetylcholine receptors enhance aversive olfactory learning in adult Drosophila. *Elife, 8*. doi:10.7554/eLife.48264

Bishop, D., Nikic, I., Brinkoetter, M., Knecht, S., Potz, S., Kerschensteiner, M., & Misgeld, T. (2011). Near-infrared branding efficiently correlates light and electron microscopy. *Nat Meth, 8*(7), 568-570. doi:10.1038/nmeth.1622

Boeckh, J., Distler, P., Ernst, K. D., Hösl, M., & Malun, D. (1990). Olfactory bulb and antennal lobe. In D. Schild (Ed.), *NATO ASI Series, Vol. H39: Chemosensory Information Processing* (pp. 201-227). Berlin, Heidelberg: Springer Verl.

Boeckh, J., & Tolbert, L. P. (1993). Synaptic organization and development of the antennal lobe in insects. *Microscopy Research and Technique, 24*(3), 260-280. doi:10.1002/jemt.1070240305

Borst, A., & Heisenberg, M. (1982). Osmotropotaxis inDrosophila melanogaster. *Journal of comparative Physiology, 147*(4), 479-484. doi:10.1007/BF00612013

Briggman, K. L., & Denk, W. (2006). Towards neural circuit reconstruction with volume electron microscopy techniques. *Current Opinion in Neurobiology, 16*(5), 562-570. doi:10.1016/j.conb.2006.08.010

Butcher, N. J., Friedrich, A. B., Lu, Z., Tanimoto, H., & Meinertzhagen, I. A. (2012). Different classes of input and output neurons reveal new features in microglomeruli of the adult *Drosophila* mushroom body calyx. *Journal of Comparative Neurology, 520*(10), 2185-2201. doi:10.1002/cne.23037

Cardona, A., Saalfeld, S., Schindelin, J., Arganda-Carreras, I., Preibisch, S., Longair, M., . . . Douglas, R. J. (2012). TrakEM2 Software for Neural Circuit Reconstruction. *PLoS ONE, 7*(6), e38011. doi:10.1371/journal.pone.0038011

Cardona, A., Saalfeld, S., Tomancak, P., & Hartenstein, V. (2009). *Drosophila* brain development: closing the gap between a macroarchitectural and microarchitectural approach. *Cold Spring Harb Symp Quant Biol, 74*, 235-248. doi:10.1101/sqb.2009.74.037

Chen, W. R., & Shepherd, G. M. (2005). The olfactory glomerulus: A cortical module with specific functions. *Journal of Neurocytology, 34*, 353–360.

Chou, Y. H., Spletter, M. L., Yaksi, E., Leong, J. C., Wilson, R. I., & Luo, L. (2010). Diversity and wiring variability of olfactory local interneurons in the *Drosophila* antennal lobe. *Nature Neuroscience, 13*(4), 439-449. doi:10.1038/nn.2489

Christie, J. M., Bark, C., Hormuzdi, S. G., Helbig, I., Monyer, H., & Westbrook, G. L. (2005). Connexin36 mediates spike synchrony in olfactory bulb glomeruli. *Neuron, 46*(5), 761-772. doi:10.1016/j.neuron.2005.04.030

Coates, K. E., Calle-Schuler, S. A., Helmick, L. M., Knotts, V. L., Martik, B. N., Salman, F., . . . Dacks, A. M. (2020). The Wiring Logic of an Identified Serotonergic Neuron That Spans Sensory

Networks. *The Journal of Neuroscience, 40*(33), 6309-6327. doi:10.1523/jneurosci.0552-20.2020

Couto, A., Alenius, M., & Dickson, B. J. (2005). Molecular, anatomical, and functional organization of the *Drosophila* olfactory system. *Current Biology, 15*(17), 1535-1547. doi:10.1016/j.cub.2005.07.034

Cover, K. K., & Mathur, B. N. (2021). Axo-axonic synapses: Diversity in neural circuit function. *Journal of Comparative Neurology, 529*(9), 2391-2401. doi:10.1002/cne.25087

Croset, V., Treiber, C. D., & Waddell, S. (2018). Cellular diversity in the Drosophila midbrain revealed by single-cell transcriptomics. *Elife, 7*. doi:10.7554/eLife.34550

Cuntz, H., Borst, A., & Segev, I. (2007). Optimization principles of dendritic structure. *Theor Biol Med Model, 4*, 21. doi:10.1186/1742-4682-4-21

Dacks, A. M., Christensen, T. A., & Hildebrand, J. G. (2006). Phylogeny of a serotonin-immunoreactive neuron in the primary olfactory center of the insect brain. *Journal of Comparative Neurology, 498*(6), 727-746. doi:10.1002/cne.21076

Dalal, T., Gupta, N., & Haddad, R. (2020). Bilateral and unilateral odor processing and odor perception. *Commun Biol, 3*(1), 150. doi:10.1038/s42003-020-0876-6

Datta, S. R., Vasconcelos, M. L., Ruta, V., Luo, S., Wong, A., Demir, E., . . . Axel, R. (2008). The *Drosophila* pheromone cVA activates a sexually dimorphic neural circuit. *Nature, 452*(7186), 473-477. doi:10.1038/nature06808

de Bruyne, M., Clyne, P. J., & Carlson, J. R. (1999). Odor coding in a model olfactory organ: the *Drosophila* maxillary palp. *Journal of Neuroscience, 19*(11), 4520-4532.

de Bruyne, M., Foster, K., & Carlson, J. R. (2001). Odor Coding in the *Drosophila* Antenna. *Neuron, 30*(2), 537-552. doi:10.1016/S0896-6273(01)00289-6

de la Rocha, J., Doiron, B., Shea-Brown, E., Josić, K., & Reyes, A. (2007). Correlation between neural spike trains increases with firing rate. *Nature, 448*(7155), 802-806. doi:10.1038/nature06028

Demir, M., Kadakia, N., Anderson, H. D., Clark, D. A., & Emonet, T. (2020). Walking Drosophila navigate complex plumes using stochastic decisions biased by the timing of odor encounters. *Elife, 9*. doi:10.7554/eLife.57524

Dolan, M.-J., Frechter, S., Bates, A. S., Dan, C., Huoviala, P., Roberts, R. J. V., . . . Jefferis, G. S. X. E. (2019). Neurogenetic dissection of the Drosophila lateral horn reveals major outputs, diverse behavioural functions, and interactions with the mushroom body. *Elife, 8*, e43079. doi:10.7554/eLife.43079

Dolan, M. J., Belliart-Guérin, G., Bates, A. S., Frechter, S., Lampin-Saint-Amaux, A., Aso, Y., . . . Jefferis, G. (2018). Communication from Learned to Innate Olfactory Processing Centers Is Required for Memory Retrieval in Drosophila. *Neuron, 100*(3), 651-668.e658. doi:10.1016/j.neuron.2018.08.037

Duistermars, B. J., Chow, D. M., & Frye, M. A. (2009). Flies require bilateral sensory input to track odor gradients in flight. *Current Biology, 19*(15), 1301-1307.

Dweck, H. K. M., Ebrahim, S. A. M., Thoma, M., Mohamed, A. A. M., Keesey, I. W., Trona, F., . . . Hansson, B. S. (2015). Pheromones mediating copulation and attraction in *Drosophila*. *Proceedings of the National Academy of Sciences, 112*(21), E2829-E2835. doi:10.1073/pnas.1504527112

Ebrahim, S. A. M., Dweck, H. K. M., Stökl, J., Hofferberth, J. E., Trona, F., Weniger, K., . . . Knaden,
M. (2015). *Drosophila* Avoids Parasitoids by Sensing Their Semiochemicals via a Dedicated
Olfactory Circuit. *PLoS Biology, 13*(12), e1002318. doi:10.1371/journal.pbio.1002318
Eckstein, N., Bates, A. S., Du, M., Hartenstein, V., Jefferis, G. S. X. E., & Funke, J. (2020).
Neurotransmitter Classification from Electron Microscopy Images at Synaptic Sites in
Drosophila. *bioRxiv*, 2020.2006.2012.148775. doi:10.1101/2020.06.12.148775
Eichler, K., Litwin-Kumar, A., Li, F., Park, Y., Andrade, I., Schneider-Mizell, C. M., . . . Cardona, A.
(2017). The Complete Connectome Of A Learning And Memory Center In An Insect Brain.
*bioRxiv*. doi:10.1101/141762
Felsenberg, J., Jacob, P. F., Walker, T., Barnstedt, O., Edmondson-Stait, A. J., Pleijzier, M. W., . . .
Waddell, S. (2018). Integration of Parallel Opposing Memories Underlies Memory
Extinction. *Cell, 175*(3), 709-722.e715. doi:https://doi.org/10.1016/j.cell.2018.08.021
Fiala, A. (2007). Olfaction and olfactory learning in Drosophila: recent progress. *Current Opinion
in Neurobiology, 17*(6), 720-726. doi:10.1016/j.conb.2007.11.009
Fishilevich, E., & Vosshall, L. B. (2005). Genetic and functional subdivision of the *Drosophila*
antennal lobe. *Current Biology, 15*(17), 1548-1553. doi:10.1016/j.cub.2005.07.066
Fröhlich, A. (1985). Freeze-Fracture Study of an Invertebrate Multiple-Contact Synapse: The Fly
Photoreceptor Tetrad. *Journal of Comparative Neurology, 241*(3), 311-326.
doi:10.1002/cne.902410306
Galizia, C. G. (2014). Olfactory coding in the insect brain: data and conjectures. *European Journal
of Neuroscience, 39*(11), n/a-n/a. doi:10.1111/ejn.12558
Galizia, C. G., Nagler, K., Holldobler, B., & Menzel, R. (1998). Odour coding is bilaterally
symmetrical in the antennal lobes of honeybees (Apis mellifera). *European Journal of
Neuroscience, 10*(9), 2964-2974.
Gao, Q., Yuan, B., & Chess, A. (2000). Convergent projections of Drosophila olfactory neurons to
specific glomeruli in the antennal lobe. *Nature*.
Gao, X. J., Clandinin, T. R., & Luo, L. (2015). Extremely Sparse Olfactory Inputs Are Sufficient to
Mediate Innate Aversion in *Drosophila PLoS ONE, 10*(4), e0125986.
doi:10.1371/journal.pone.0125986
Gaudry, Q., Hong, E. J., Kain, J., de Bivort, B. L., & Wilson, R. I. (2013). Asymmetric
neurotransmitter release enables rapid odour lateralization in *Drosophila*. *Nature,
493*(7432), 424-428. doi:10.1038/nature11747
Gondré-Lewis, M. C., Park, J. J., & Loh, Y. P. (2012). Chapter Two - Cellular Mechanisms for the
Biogenesis and Transport of Synaptic and Dense-Core Vesicles. In K. W. Jeon (Ed.),
*International Review of Cell and Molecular Biology* (Vol. 299, pp. 27-115): Academic Press.
Goyal, R. K., & Chaudhury, A. (2013). Structure activity relationship of synaptic and junctional
neurotransmission. *Autonomic Neuroscience: Basic and Clinical, 176*(1), 11-31.
doi:10.1016/j.autneu.2013.02.012
Grabe, V., Baschwitz, A., Dweck, Hany K. M., Lavista-Llanos, S., Hansson, Bill S., & Sachse, S.
(2016). Elucidating the Neuronal Architecture of Olfactory Glomeruli in the *Drosophila*
Antennal Lobe. *Cell Reports, 16*(12), 3401-3413. doi:10.1016/j.celrep.2016.08.063
Grabe, V., & Sachse, S. (2018). Fundamental principles of the olfactory code. *BioSystems, 164*, 94-
101. doi:10.1016/j.biosystems.2017.10.010

Grabe, V., Schubert, M., Strube-Bloss, M., Reinert, A., Trautheim, S., Lavista-Llanos, S., . . . Sachse, S. (2020). Odor-Induced Multi-Level Inhibitory Maps in Drosophila. *eNeuro, 7*(1). doi:10.1523/eneuro.0213-19.2019

Grabe, V., Strutz, A., Baschwitz, A., Hansson, B. S., & Sachse, S. (2015). Digital in vivo 3D atlas of the antennal lobe of *Drosophila melanogaster. Journal of Comparative Neurology, 523*(3), 530-544. doi:10.1002/cne.23697

Graubard, K., Raper, J. A., & Hartline, D. K. (1980). Graded synaptic transmission between spiking neurons. *Proc Natl Acad Sci U S A, 77*(6), 3733-3735. doi:10.1073/pnas.77.6.3733

Gruber, L., Rybak, J., Hansson, B. S., & Cantera, R. (2018). Synaptic Spinules in the Olfactory Circuit of Drosophila melanogaster. *Front Cell Neurosci, 12*(86), 86. doi:10.3389/fncel.2018.00086

Guo, D., Wu, S., Chen, M., Perc, M., Zhang, Y., Ma, J., . . . Yao, D. (2016). Regulation of Irregular Neuronal Firing by Autaptic Transmission. *Sci Rep, 6*, 26096. doi:10.1038/srep26096

Guven-Ozkan, T., & Davis, R. L. (2014). Functional neuroanatomy of *Drosophila* olfactory memory formation. *Learning & Memory, 21*(10), 519-526. doi:10.1101/lm.034363.114

Hallem, E. A., & Carlson, J. R. (2006). Coding of odors by a receptor repertoire. *Cell, 125*(1), 143-160. doi:10.1016/j.cell.2006.01.050

Hallem, E. A., Ho, M. G., & Carlson, J. R. (2004). The molecular basis of odor coding in the Drosophila antenna. *Cell, 117*(7), 965-979. doi:10.1016/j.cell.2004.05.012

Hansson, B. S., & Anton, S. (2000). Function and morphology of the antennal lobe: new developments. *Annual Review of Entomology, 45*, 203-231. doi:10.1146/annurev.ento.45.1.203

Hartenstein, V. (2016). The Central Nervous System of Invertebrates. In S. V. Shepherd (Ed.), *The Wiley Handbook of Evolutionary Neuroscience* (pp. 173-235).

Haverkamp, A., Hansson, B. S., & Knaden, M. (2018). Combinatorial Codes and Labeled Lines: How Insects Use Olfactory Cues to Find and Judge Food, Mates, and Oviposition Sites in Complex Environments. *Front Physiol, 9*, 49. doi:10.3389/fphys.2018.00049

Helmstaedter, M. (2013). Cellular-resolution connectomics: challenges of dense neural circuit reconstruction. *Nature Methods, 10*(6), 501-507. doi:10.1038/nmeth.2476

Hirata, Y. (1964). Some observations on the fine structure of the synapses in the olfactory bulb of the mouse, with particular reference to the atypical synaptic configurations. *Archivum histologicum japonicum, 24*(3), 293-302.

Hong, E. J., & Wilson, R. I. (2015). Simultaneous encoding of odors by channels with diverse sensitivity to inhibition. *Neuron, 85*(3), 573-589. doi:10.1016/j.neuron.2014.12.040

Horne, J. A., Langille, C., McLin, S., Wiederman, M., Lu, Z., Xu, C. S., . . . Meinertzhagen, I. A. (2018). A resource for the *Drosophila* antennal lobe provided by the connectome of glomerulus VA1v. *Elife, 7*, e37550. doi:10.7554/eLife.37550

Huang, G. B., Scheffer, L. K., & Plaza, S. M. (2018). Fully-Automatic Synapse Prediction and Validation on a Large Data Set. *Frontiers in Neural Circuits, 12*(87). doi:10.3389/fncir.2018.00087

Hulse, B. K., Haberkern, H., Franconville, R., Turner-Evans, D. B., Takemura, S.-y., Wolff, T., . . . Jayaraman, V. (2021). A connectome of the Drosophila central complex reveals network motifs suitable for flexible navigation and context-dependent action selection. *Elife, 10*, e66039. doi:10.7554/eLife.66039

Huoviala, P., Dolan, M.-J., Love, F. M., Myers, P., Frechter, S., Namiki, S., . . . Jefferis, G. S. X. E.
(2020). Neural circuit basis of aversive odour processing in *Drosophila* from sensory input
to descending output. 394403. doi:10.1101/394403 %J bioRxiv
Ikeda, K., & Bekkers, J. M. Autapses. *Current Biology, 16*(9), R308. doi:10.1016/j.cub.2006.03.085
Ikeda, K., & Bekkers, J. M. (2006). Autapses. *Current Biology, 16*(9), R308.
doi:10.1016/j.cub.2006.03.085
Jeanne, James M., & Wilson, Rachel I. (2015). Convergence, Divergence, and Reconvergence in a
Feedforward Network Improves Neural Speed and Accuracy. *Neuron, 88*(5), 1014-1026.
doi:10.1016/j.neuron.2015.10.018
Jefferis, G. S., Potter, C. J., Chan, A. M., Marin, E. C., Rohlfing, T., Maurer, C. R., Jr., & Luo, L.
(2007). Comprehensive maps of *Drosophila* higher olfactory centers: spatially segregated
fruit and pheromone representation. *Cell, 128*(6), 1187-1203.
doi:10.1016/j.cell.2007.01.040
Karnovsky, M. J. (1965). A formaldehyde-glutaraldehyde fixative of high osmolarity for use in
electron microscopy. *Journal of Cellular Biology, 27*, 137.
Kazama, H., & Wilson, R. I. (2008). Homeostatic Matching and Nonlinear Amplification at
Identified Central Synapses. *Neuron, 58*(3), 401-413.
doi:http://dx.doi.org/10.1016/j.neuron.2008.02.030
Kazama, H., & Wilson, R. I. (2009). Origins of correlated activity in an olfactory circuit. *Nature*
*Neuroscience, 12*(9), 1136-1144. doi:10.1038/nn.2376
Keene, A. C., & Waddell, S. (2007). Drosophila olfactory memory: single genes to complex neural
circuits. *Nature Reviews: Neuroscience, 8*(5), 341-354.
Keesey, I. W., & Hansson, B. S. (2021). 10 - The neuroethology of labeled lines in insect olfactory
systems. In G. J. Blomquist & R. G. Vogt (Eds.), *Insect Pheromone Biochemistry and*
*Molecular Biology (Second Edition)* (pp. 285-327). London: Academic Press.
Keesey, I. W., Zhang, J., Depetris-Chauvin, A., Obiero, G. F., Knaden, M., & Hansson, B. S. (2019).
Evolution of a pest: towards the complete neuroethology of <em>Drosophila
suzukii</em> and the subgenus <em>Sophophora</em>. *bioRxiv*, 717322.
doi:10.1101/717322
Knaden, M., & Hansson, B. S. (2014). Mapping odor valence in the brain of flies and mice. *Current*
*Opinion in Neurobiology, 24*(1), 34-38. doi:10.1016/j.conb.2013.08.010
Knaden, M., Strutz, A., Ahsan, J., Sachse, S., & Hansson, B. S. (2012). Spatial representation of
odorant valence in an insect brain. *Cell Reports, 1*, 392-399.
doi:10.1016/j.celrep.2012.03.002
Knott, G., Marchman, H., & Lich, B. (2008). Serial Section Scanning Electron Microscopy of Adult
Brain Tissue Using Focused Ion Beam Milling. *Journal of Neuroscience, 28*(12), 2964-2959.
doi:10.1523/JNEUROSCI.3189-07.2008
Kreher, S. A., Mathew, D., Kim, J., & Carlson, J. R. (2008). Translation of sensory input into
behavioral output via an olfactory system. *Neuron, 59*(1), 110-124.
doi:10.1016/j.neuron.2008.06.010
Kumar, A., Schiff, O., Barkai, E., Mel, B. W., Poleg-Polsky, A., & Schiller, J. (2018). NMDA spikes
mediate amplification of inputs in the rat piriform cortex. *Elife, 7*, e38446.
doi:10.7554/eLife.38446

Kurtovic, A., Widmer, A., & Dickson, B. J. (2007). A single class of olfactory neurons mediates
behavioural responses to a Drosophila sex pheromone. *Nature, 446*(7135), 542-546.
doi:10.1038/nature05672

Laissue, P. P., Reiter, C., Hiesinger, P. R., Halter, S., Fischbach, K. F., & Stocker, R. F. (1999). Three-
dimensional reconstruction of the antennal lobe in *Drosophila melanogaster*. *The Journal
of Comparative Neurology, 405*(4), 543-552. doi:10.1002/(sici)1096-
9861(19990322)405:4<543::aid-cne7>3.0.co;2-a

Laissue, P. P., & Vosshall, L. B. (2008). The Olfactory Sensory Map in *Drosophila*. In G. M. Technau
(Ed.), *Brain Development in Drosophila melanogaster* (pp. 102-114 ). New York: Springer.

Lei, H., Oland, L. A., Riffell, J. A., Beyerlein, A., & Hildebrand, J. G. (2010). Microcircuits for
Olfactory Information Processing in the Antennal Lobe of *Manduca sexta*. In G. M.
Shepherd & S. Grillner (Eds.), *Handbook of Brain Microcircuits* (pp. 417-426). New York:
Oxford University Press.

Li, F., Lindsey, J. W., Marin, E. C., Otto, N., Dreher, M., Dempsey, G., . . . Rubin, G. M. (2020). The
connectome of the adult Drosophila mushroom body provides insights into function. *Elife,
9*. doi:10.7554/eLife.62576

Li, H., Horns, F., Wu, B., Xie, Q., Li, J., Li, T., . . . Luo, L. (2017). Classifying Drosophila Olfactory
Projection Neuron Subtypes by Single-Cell RNA Sequencing. *Cell, 171*(5), 1206-
1220.e1222. doi:https://doi.org/10.1016/j.cell.2017.10.019

Li, P. H., Lindsey, L. F., Januszewski, M., Zheng, Z., Bates, A. S., Taisz, I., . . . Jain, V. (2020).
Automated Reconstruction of a Serial-Section EM *Drosophila* Brain with Flood-Filling
Networks and Local Realignment. *bioRxiv*, 605634. doi:10.1101/605634

Liang, L., & Luo, L. (2010). The olfactory circuit of the fruit fly *Drosophila melanogaster*. *Sci China
Life Sci, 53*(4), 472-484. doi:10.1007/s11427-010-0099-z

Lin, S., Kao, C.-F., Yu, H.-H., Huang, Y., & Lee, T. (2012). Lineage analysis of Drosophila lateral
antennal lobe neurons reveals notch-dependent binary temporal fate decisions. *PLoS
Biology, 10*(11), e1001425-e1001425. doi:10.1371/journal.pbio.1001425

Liu, T. X., Davoudian, P. A., Lizbinski, K. M., & Jeanne, J. M. (2022). Connectomic features
underlying diverse synaptic connection strengths and subcellular computation. *Current
Biology, 32*(3), 559-569.e555. doi:https://doi.org/10.1016/j.cub.2021.11.056

Liu, W. W., & Wilson, R. I. (2013). Glutamate is an inhibitory neurotransmitter in the *Drosophila
olfactory system. *Proceedings of the National Academy of Sciences, 110*(25), 10294–
10299. doi:10.1073/pnas.1220560110

Malun, D. (1991). Inventory and distribution of synapses of identified uniglomerular projection
neurons in the antennal lobe of Periplaneta americana. *Journal of Comparative
Neurology, 305*(2), 348-360. doi:10.1002/cne.903050215

Malun, D., Waldow, U., Kraus, D., & Boeckh, J. (1993). Connections between the Deutocerebrum
and the Protocerebrum, and Neuroanatomy of Several Classes of Deutocerebral
Projection Neurons in the Brain of Male Periplaneta-Americana. *Journal of Comparative
Neurology, 329*(2), 143-162.

Marin, E. C., Büld, L., Theiss, M., Sarkissian, T., Roberts, R. J. V., Turnbull, R., . . . Jefferis, G. (2020).
Connectomics Analysis Reveals First-, Second-, and Third-Order Thermosensory and
Hygrosensory Neurons in the Adult Drosophila Brain. *Current Biology, 30*(16), 3167-
3182.e3164. doi:10.1016/j.cub.2020.06.028

Masse, N. Y., Turner, G. C., & Jefferis, G. S. (2009). Olfactory information processing in Drosophila. *Current Biology, 19*(16), R700-713. doi:10.1016/j.cub.2009.06.026

Masson, C., & Mustaparta, H. (1990). Chemical information processing in the olfactory system of insects. *Physiological Review, 70(1)*, 199-245.

McTavish, T., Migliore, M., Shepherd, G., & Hines, M. (2012). Mitral cell spike synchrony modulated by dendrodendritic synapse location. *Frontiers in Computational Neuroscience, 6*. doi:10.3389/fncom.2012.00003

Meinertzhagen, I. A. (2018). Of what use is connectomics? A personal perspective on the *Drosophila* connectome. *Journal of Experimental Biology, 221*(10), jeb164954. doi:10.1242/jeb.164954 %J The Journal of Experimental Biology

Meinertzhagen, I. A., & O'Neil, S. D. (1991). Synaptic organization of columnar elements in the lamina of the wild type in Drosophila melanogaster. *Journal of Comparative Neurology, 305*, 232-263.

Miroschnikow, A., Schlegel, P., Schoofs, A., Hueckesfeld, S., Li, F., Schneider-Mizell, C. M., . . . Pankratz, M. J. (2018). Convergence of monosynaptic and polysynaptic sensory paths onto common motor outputs in a Drosophila feeding connectome. *Elife, 7*, e40247. doi:10.7554/eLife.40247

Mohamed, A. A. M., Hansson, B. S., & Sachse, S. (2019a). Third-Order Neurons in the Lateral Horn Enhance Bilateral Contrast of Odor Inputs Through Contralateral Inhibition in Drosophila. *Front Physiol, 10*, 851. doi:10.3389/fphys.2019.00851

Mohamed, A. A. M., Retzke, T., Das Chakraborty, S., Fabian, B., Hansson, B. S., Knaden, M., & Sachse, S. (2019b). Odor mixtures of opposing valence unveil inter-glomerular crosstalk in the Drosophila antennal lobe. *Nat Commun, 10*(1), 1201. doi:10.1038/s41467-019-09069-1

Mosca, T. J., & Luo, L. (2014). Synaptic organization of the *Drosophila* antennal lobe and its regulation by the Teneurins. *Elife, 3*(3), e03726. doi:10.7554/eLife.03726

Münch, D., & Galizia, C. G. (2016). DoOR 2.0 - Comprehensive Mapping of Drosophila melanogaster Odorant Responses. *Scientific Reports, 6*, 21841. doi:10.1038/srep21841 http://www.nature.com/articles/srep21841#supplementary-information

Nässel, D. R., Enell, L. E., Santos, J. G., Wegener, C., & Johard, H. A. (2008). A large population of diverse neurons in the Drosophila central nervous system expresses short neuropeptide F, suggesting multiple distributed peptide functions. *BMC Neuroscience, 9*, 90. doi:10.1186/1471-2202-9-90

Nässel, D. R., & Homberg, U. (2006). Neuropeptides in interneurons of the insect brain. *Cell and Tissue Research, 326*(1), 1-24. doi:10.1007/s00441-006-0210-8

Ng, M., Roorda, R. D., Lima, S. Q., Zemelman, B. V., Morcillo, P., & Miesenböck, G. (2002). Transmission of Olfactory Information between Three Populations of Neurons in the Antennal Lobe of the Fly. *Neuron, 36*(3), 463-474. doi:10.1016/s0896-6273(02)00975-3

Norgate, M., Lee, E., Southon, A., Farlow, A., Batterham, P., Camakaris, J., & Burke, R. (2006). Essential Roles in Development and Pigmentation for the *Drosophila* Copper Transporter DmATP7. *Molecular Biology of the Cell, 17*(1), 475-484. doi:10.1091/mbc.E05-06-0492

Okada, R., Awasaki, T., & Ito, K. (2009). Gamma-aminobutyric acid (GABA)-mediated neural connections in the *Drosophila* antennal lobe. *Journal of Comparative Neurology, 514*(1), 74-91. doi:10.1002/cne.21971

Olsen, S. R., & Wilson, R. I. (2008). Lateral presynaptic inhibition mediates gain control in an
olfactory circuit. *Nature, 452*(7190), 956-960. doi:10.1038/nature06864
Otto, N., Pleijzier, M. W., Morgan, I. C., Edmondson-Stait, A. J., Heinz, K. J., Stark, I., . . . Waddell,
S. (2020). Input Connectivity Reveals Additional Heterogeneity of Dopaminergic
Reinforcement in Drosophila. *Current Biology, 30*(16), 3200-3211.e3208.
doi:https://doi.org/10.1016/j.cub.2020.05.077
Owald, D., & Waddell, S. (2015). Olfactory learning skews mushroom body output pathways to
steer behavioral choice in Drosophila. *Current Opinion in Neurobiology, 35*, 178-184.
doi:10.1016/j.conb.2015.10.002
Parthasarathy, K., & Bhalla, U. S. (2013). Laterality and symmetry in rat olfactory behavior and in
physiology of olfactory input. *Journal of Neuroscience, 33*(13), 5750-5760.
doi:10.1523/jneurosci.1781-12.2013
Prokop, A., & Meinertzhagen, I. A. (2006). Development and structure of synaptic contacts in
Drosophila. *Seminars in Cell & Developmental Biology, 17*(1), 20-30.
doi:10.1016/j.semcdb.2005.11.010
Ronchi, P., Mizzon, G., Machado, P., D'Imprima, E., Best, B. T., Cassella, L., . . . Schwab, Y. (2021).
High-precision targeting workflow for volume electron microscopy. *Journal of Cell Biology,*
*220*(9). doi:10.1083/jcb.202104069
Root, C. M., Masuyama, K., Green, D. S., Enell, L. E., Nassel, D. R., Lee, C. H., & Wang, J. W. (2008).
A presynaptic gain control mechanism fine-tunes olfactory behavior. *Neuron, 59*(2), 311-
321. doi:10.1016/j.neuron.2008.07.003
Rospars, J.-P., Grémiaux, A., Jarriault, D., Chaffiol, A., Monsempes, C., Deisig, N., . . . Martinez, D.
(2014). Heterogeneity and Convergence of Olfactory First-Order Neurons Account for the
High Speed and Sensitivity of Second-Order Neurons. *PLoS Computational Biology, 10*(12),
e1003975. doi:10.1371/journal.pcbi.1003975
Rozenfeld, E., Lerner, H., & Parnas, M. (2019). Muscarinic Modulation of Antennal Lobe
GABAergic Local Neurons Shapes Odor Coding and Behavior. *Cell Rep, 29*(10), 3253-
3265.e3254. doi:10.1016/j.celrep.2019.10.125
Rybak, J. (2013). Exploring Brain Connectivity in Insect Model Systems of Learning and Memory.
In R. Menzel & P. Benjamin (Eds.), *Invertebrate Learning and Memory* (pp. 26-40). San
Diego: Academic Press.
Rybak, J. (2016). Perspective-Brain atlases for studying neuronal circuitry in arthropods. In A.
Schmidt-Rhaesa, S. Harzsch, & G. Purschke (Eds.), *Structure and Evolution of Invertebrate*
*Nervous Systems* (Vol. 1). New York: Oxford University Press.
Rybak, J., & Hansson, B. S. (2018). Olfactory Microcircuits in *Drosophila Melanogaster*. In G. M.
Shepherd & S. Grillner (Eds.), *Handbook of Brain Microcircuits* (2nd ed., pp. 361-367).
Oxford, UK: Oxford University Press.
Rybak, J., Talarico, G., Ruiz, S., Arnold, C., Cantera, R., & Hansson, B. S. (2016). Synaptic circuitry
of identified neurons in the antennal lobe of *Drosophila melanogaster*. *Journal of*
*Comparative Neurology, 524*(9), 1920-1956. doi:10.1002/cne.23966
Saada, R., Miller, N., Hurwitz, I., & Susswein, A. J. (2009). Autaptic excitation elicits persistent
activity and a plateau potential in a neuron of known behavioral function. *Current Biology,*
*19*(6), 479-484. doi:10.1016/j.cub.2009.01.060

Saalfeld, S., Cardona, A., Hartenstein, V., & Tomančák, P. (2009). CATMAID: collaborative
annotation toolkit for massive amounts of image data. *Bioinformatics, 25*(15), 1984-1986.
doi:10.1093/bioinformatics/btp266
Sachse, S., & Galizia, C. G. (2002). Role of inhibition for temporal and spatial odor representation
in olfactory output neurons: a calcium imaging study. *Journal of Neurophysiology, 87*(2),
1106-1117.
Sachse, S., & Hansson, B. S. (2016). Research spotlight: Olfactory coding in *Drosophila*
*melanogaster*. In A. Schmidt-Rhaesa, S. Harzsch, & G. Purschke (Eds.), *Structure and*
*Evolution of Invertebrate Nervous Systems* (pp. 640-645). Oxford: Oxford University Press.
Sachse, S., & Manzini, I. (2021). Editorial for the special issue "Olfactory Coding and Circuitries".
*Cell and Tissue Research, 383*(1), 1-6. doi:10.1007/s00441-020-03389-1
Scheffer, L. K., Xu, C. S., Januszewski, M., Lu, Z., Takemura, S.-y., Hayworth, K. J., . . . Plaza, S. M.
(2020). A connectome and analysis of the adult Drosophila central brain. *Elife, 9*, e57443.
doi:10.7554/eLife.57443
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . . Cardona, A.
(2012). Fiji: an open-source platform for biological-image analysis. *Nat Meth, 9*(7), 676-
682.
doi:[http://www.nature.com/nmeth/journal/v9/n7/abs/nmeth.2019.html#supplementary](http://www.nature.com/nmeth/journal/v9/n7/abs/nmeth.2019.html#supplementary)
-information
Schlegel, P., Bates, A. S., Stürner, T., Jagannathan, S. R., Drummond, N., Hsu, J., . . . Jefferis, G. S.
X. E. (2021). Information flow, cell types and stereotypy in a full olfactory connectome.
*Elife, 10*, e66018. doi:10.7554/eLife.66018
Schneider-Mizell, C. M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M. F., . . . Cardona, A.
(2016). Quantitative neuroanatomy for connectomics in *Drosophila*. *Elife, 5*, e12059.
doi:10.7554/eLife.12059
Seki, Y., Dweck, H. K. M., Rybak, J., Wicher, D., Sachse, S., & Hansson, B. S. (2017). Olfactory
coding from the periphery to higher brain centers in the Drosophila brain. *BMC Biology,*
*15*(1), 56. doi:10.1186/s12915-017-0389-z
Seki, Y., Rybak, J., Wicher, D., Sachse, S., & Hansson, B. S. (2010). Physiological and morphological
characterization of local interneurons in the *Drosophila* antennal lobe. *Journal of*
*Neurophysiology, 104*(2), 1007-1019. doi:jn.00249.2010 [pii]
10.1152/jn.00249.2010
Semmelhack, J. L., & Wang, J. W. (2009). Select Drosophila glomeruli mediate innate olfactory
attraction and aversion. *Nature, 459*(7244), 218-223.
Shanbhag, S. R., Muller, B., & Steinbrecht, R. A. (1999). Atlas of olfactory organs of *Drosophila*
*melanogaster* - 1. Types, external organization, innervation and distribution of olfactory
sensilla. *International Journal of Insect Morphology & Embryology, 28*(4), 377-397. doi:Doi
10.1016/S0020-7322(99)00039-2
Shepherd, G. M. (2011). The Olfactory Bulb: A Simple System in the Mammalian Brain
*Comprehensive Physiology*: John Wiley & Sons, Inc.
Shepherd, G. M., Rowe, T. B., & Greer, C. A. (2021). An Evolutionary Microcircuit Approach to the
Neural Basis of High Dimensional Sensory Processing in Olfaction. *Frontiers in Cellular*
*Neuroscience, 15*. doi:10.3389/fncel.2021.658480

Silbering, A. F., & Galizia, C. G. (2007). Processing of odor mixtures in the Drosophila antennal
lobe reveals both global inhibition and glomerulus-specific interactions. *Journal of*
*Neuroscience, 27*(44), 11966-11977. doi:10.1523/jneurosci.3099-07.2007
Silbering, A. F., Okada, R., Ito, K., & Galizia, C. G. (2008). Olfactory information processing in the
*Drosophila* antennal lobe: anything goes? *Journal of Neuroscience, 28*(49), 13075-13087.
doi:10.1523/JNEUROSCI.2973-08.2008
Silbering, A. F., Rytz, R., Grosjean, Y., Abuin, L., Ramdya, P., Jefferis, G. S., & Benton, R. (2011).
Complementary Function and Integrated Wiring of the Evolutionarily Distinct *Drosophila*
Olfactory Subsystems. *Journal of Neuroscience, 31*(38), 13357-13375.
doi:10.1523/JNEUROSCI.2360-11.2011
Stensmyr, M. C., Dweck, H. K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., . . . Hansson, B. S.
(2012). A conserved dedicated olfactory circuit for detecting harmful microbes in
*Drosophila*. *Cell, 151*(6), 1345-1357. doi:10.1016/j.cell.2012.09.046
Stocker, R. F., Lienhard, M. C., Borst, A., & Fischbach, K. F. (1990). Neuronal architecture of the
antennal lobe in Drosophila melanogaster. *Cell and Tissue Research, 262*(1), 9-34.
Stocker, R. F., Singh, R. N., Schorderet, M., & Siddiqi, O. (1983). Projection patterns of different
types of antennal sensilla in the antennal glomeruli of Drosophila melanogaster. *Cell and*
*Tissue Research, 232*(2), 237-248. doi:10.1007/bf00213783
Strutz, A., Soelter, J., Baschwitz, A., Farhan, A., Grabe, V., Rybak, J., . . . Sachse, S. (2014).
Decoding odor quality and intensity in the *Drosophila* brain. *Elife, 3*, e04147.
doi:10.7554/eLife.04147
Su, C.-Y., Menuz, K., & Carlson, J. R. (2009). Olfactory Perception: Receptors, Cells, and Circuits.
*Cell, 139*(1), 45-59. doi:10.1016/j.cell.2009.09.015
Suh, G. S. B., Wong, A. M., Hergarden, A. C., Wang, J. W., Simon, A. F., Benzer, S., . . . Anderson, D.
J. (2004). A single population of olfactory sensory neurons mediates an innate avoidance
behaviour in *Drosophila*. *Nature, 431*(7010), 854-859.
doi:http://www.nature.com/nature/journal/v431/n7010/suppinfo/nature02980_S1.html
Sun, X. J., Tolbert, L. P., & Hildebrand, J. G. (1997). Synaptic organization of the uniglomerular
projection neurons of the antennal lobe of the moth *Manduca sexta*: a laser scanning
confocal and electron microscopic study. *Journal of Comparative Neurology, 379*(1), 2-20.
Szyszka, P., & Galizia, C. G. (2015). Olfaction in Insects. In R. L. Doty (Ed.), *Handbook of Olfaction*
*and Gustation* (3 ed., pp. 531-546): John Wiley & Sons, Inc.
Taisz, I., Donà, E., Münch, D., Bailey, S. N., Morris, W. J., Meechan, K. I., . . . Galili, D. S. (2022).
Generating parallel representations of position and identity in the olfactory system.
*bioRxiv*, 2022.2005.2013.491877. doi:10.1101/2022.05.13.491877
Takemura, S.-y., Xu, C. S., Lu, Z., Rivlin, P. K., Parag, T., Olbris, D. J., . . . Scheffer, L. K. (2015).
Synaptic circuits and their variations within different columns in the visual system of
Drosophila. *Proceedings of the National Academy of Sciences, 112*(44), 13711-13716.
doi:10.1073/pnas.1509820112
Tamás, G., Buhl, E. H., & Somogyi, P. (1997). Massive Autaptic Self-Innervation of GABAergic
Neurons in Cat Visual Cortex. *The Journal of Neuroscience, 17*(16), 6352-6364.
doi:10.1523/jneurosci.17-16-06352.1997

Tanaka, N. K., Endo, K., & Ito, K. (2012). The organization of antennal lobe-associated neurons in the adult *Drosophila* melanogaster brain. *Journal of Comparative Neurology, 520*(18), 4067-4130. doi:10.1002/cne.23142

Thoma, M., Hansson, B. S., & Knaden, M. (2015). High-resolution Quantification of Odor-guided Behavior in Drosophila melanogaster Using the Flywalk Paradigm. *J Vis Exp*(106), e53394. doi:10.3791/53394

Tobin, W. F., Wilson, R. I., & Lee, W.-C. A. (2017). Wiring variations that enable and constrain neural computation in a sensory microcircuit. *Elife, 6*, e24838. doi:10.7554/eLife.24838

Tran-Van-Minh, A., Cazé, R. D., Abrahamsson, T., Cathala, L., Gutkin, B. S., & DiGregorio, D. A. (2015). Contribution of sublinear and supralinear dendritic integration to neuronal computations. *Frontiers in Cellular Neuroscience, 9*. doi:10.3389/fncel.2015.00067

Trujillo-Cenoz, O. (1969). Some Aspects of the Structural Organization of the Medulla in Muscoid Flies l. *J Ultrastructural Research, 27*, 533-553.

Van der Loos, H., & Glaser, E. M. (1972). Autapses in neocortex cerebri: synapses between a pyramidal cell's axon and its own dendrites. *Brain Research, 48*, 355-360. doi:10.1016/0006-8993(72)90189-8

Vosshall, L. B., Wong, A. M., & Axel, R. (2000). An olfactory sensory map in the fly brain. *Cell, 102*(2), 147-159. doi:10.1016/S0092-8674(00)00021-0

Wang, J. W. (2012). Presynaptic modulation of early olfactory processing in *Drosophila*. *Developmental Neurobiology, 72*(1), 87-99. doi:10.1002/dneu.20936

Wicher, D., & Miazzi, F. (2021). Functional properties of insect olfactory receptors: ionotropic receptors and odorant receptors. *Cell and Tissue Research, 383*(1), 7-19. doi:10.1007/s00441-020-03363-x

Wiles, L., Gu, S., Pasqualetti, F., Parvesse, B., Gabrieli, D., Bassett, D. S., & Meaney, D. F. (2017). Autaptic Connections Shift Network Excitability and Bursting. *Sci Rep, 7*, 44006. doi:10.1038/srep44006

Wilson, R. I. (2013). Early Olfactory Processing in Drosophila: Mechanisms and Principles. *Annual Review of Neuroscience, 36*, 217-241. doi:10.1146/annurev-neuro-062111-150533

Xu, C. S., Hayworth, K. J., Lu, Z., Grob, P., Hassan, A. M., García-Cerdán, J. G., . . . Hess, H. F. (2017). Enhanced FIB-SEM systems for large-volume 3D imaging. *Elife, 6*, e25916. doi:10.7554/eLife.25916

Xu, C. S., Januszewski, M., Lu, Z., Takemura, S.-y., Hayworth, K. J., Huang, G., . . . Plaza, S. M. (2020). A Connectome of the Adult *Drosophila* Central Brain. 2020.2001.2021.911859. doi:10.1101/2020.01.21.911859 %J bioRxiv

Yang, K., Liu, T., Wang, Z., Liu, J., Shen, Y., Pan, X., . . . Zhang, K. (2022). Classifying Drosophila olfactory projection neuron boutons by quantitative analysis of electron microscopic reconstruction. *iScience, 25*(5), 104180. doi:https://doi.org/10.1016/j.isci.2022.104180

Yasuyama, K., Meinertzhagen, I. A., & Schurmann, F. W. (2003). Synaptic connections of cholinergic antennal lobe relay neurons innervating the lateral horn neuropile in the brain of *Drosophila melanogaster*. *Journal of Comparative Neurology, 466*(3), 299-315. doi:10.1002/cne.10867

Yasuyama, K., & Salvaterra, P. M. (1999). Localization of choline acetyltransferase-expressing neurons in *Drosophila* nervous system. *Microscopy Research and Technique, 45*(2), 65-79.

2053     Yin, L., Zheng, R., Ke, W., He, Q., Zhang, Y., Li, J., . . . Shu, Y. (2018). Autapses enhance bursting
2054          and coincidence detection in neocortical pyramidal cells. *Nature Communications, 9*(1),
2055          4890. doi:10.1038/s41467-018-07317-4
2056     Yokoi, M., Mori, K., & Nakanishi, S. (1995). Refinement of odor molecule tuning by
2057          dendrodendritic synaptic inhibition in the olfactory bulb. *Proceedings of the National*
2058          *Academy of Sciences, 92*(8), 3371-3375. doi:doi:10.1073/pnas.92.8.3371
2059     Zheng, Z., Lauritzen, J. S., Perlman, E., Robinson, C. G., Nichols, M., Milkie, D., . . . Bock, D. D.
2060          (2017). A Complete Electron Microscopy Volume Of The Brain Of Adult Drosophila
2061          melanogaster. *bioRxiv*. doi:10.1101/140905
2062     Zheng, Z., Lauritzen, J. S., Perlman, E., Robinson, C. G., Nichols, M., Milkie, D., . . . Bock, D. D.
2063          (2018). A Complete Electron Microscopy Volume of the Brain of Adult Drosophila
2064          melanogaster. *Cell, 174*(3), 730-743.e722. doi:https://doi.org/10.1016/j.cell.2018.06.019
2065

2066

2067    **Figure 1**

2068
2069   **Figure 2**

| Row | Values | Unit | OSNs | | uPNs | | MGNs | | all neurons | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | DA2 | DL5 | DA2 | DL5 | DA2 | DL5 | DA2 | DL5 |
| 1 | Total neuronal length | μm | 2012 | 2727 | 4652 | 5015 | 10705 | 14411 | 17370 | 22153 |
| 2 | | input | 868 | 1083 | 3887 | 3955 | 7229 | 9018 | 11984 | 14056 |
| 3 | Total synaptic counts | output | 6671 | 6828 | 1624 | 3108 | 5659 | 6749 | 13954 | 16685 |
| 4 | | T-bars | 1063 | 1213 | 322 | 602 | 1263 | 1572 | 2648 | 3387 |
| 5 | Total innervation density (sum of length of all neuronal fibers/glomerular volume) | μm/μm$^3$ | 1.26 | 1.05 | 2.91 | 1.93 | 6.69 | 5.54 | 10.86 | 8.52 |
| 6 | | inputs/μm$^3$ | 0.54 | 0.42 | 2.43 | 1.52 | 4.52 | 3.47 | 7.49 | 5.41 |
| 7 | Total glomerular synaptic density (total synaptic counts/glomerular volume) | outputs/μm$^3$ | 4.17 | 2.63 | 1.02 | 1.20 | 3.54 | 2.60 | 8.72 | 6.42 |
| 8 | | T-bars/μm$^3$ | 0.66 | 0.47 | 0.20 | 0.23 | 0.79 | 0.60 | 1.66 | 1.30 |
| 9 | | inputs/μm | 0.42 | 0.39 | 0.83 | 0.79 | 0.62 | 0.59 | 0.59 | 0.56 |
| 10 | Neuronal synaptic density (synaptic counts/neuronal length) | outputs/μm | 3.37 | 2.62 | 0.33 | 0.62 | 0.52 | 0.51 | 1.06 | 0.87 |
| 11 | | T-bars/μm | 0.53 | 0.46 | 0.07 | 0.12 | 0.12 | 0.12 | 0.19 | 0.18 |
| 12 | Synaptic ratio | T-bars/inputs | 1.31 | 1.27 | 0.08 | 0.15 | 0.23 | 0.24 | 0.43 | 0.42 |
| 13 | | outputs/inputs | 8.29 | 7.29 | 0.40 | 0.79 | 1.04 | 1.11 | 2.40 | 2.17 |
| 14 | Polyadicity | outputs/T-bars | 6.35 | 5.70 | 4.95 | 5.16 | 3.22 | 2.64 | 3.88 | 3.17 |

2070    **Table 1**

2071
2072    **Figure 3**

2073
2074      **Figure 4**

2075
2076 **Figure 5**

**A**

**B**

**Figure 6**

 **Figure 7**

**SUPPLEMENTARY FIGURES**

**Figure 1 -- Video 1: FIB-SEM dataset of a DA2 glomerulus with highlighted uPN reconstruction**
(see extra file).
The video shows a full FIB-SEM scan of a DA2 glomerulus at pixel resolution 4x4x20 nm, with the
neuron trace of a single uniglomerular projection neuron (uPN#2) highlighted in yellow.

**Figure 2 – figure supplement 1: Neuronal volume and polyadicity**

**A**: Ratio between neuronal fiber volume and length in glomerulus DA2. Data represent mean +
standard deviation (error bars) (OSNs n=30; uPNs n =5; MGNs n = 15). **B-D**: Frequency of T-bars
associated with a number of postsynaptic contacts (Polyadicity) in OSNs (**B**), uPNs (**C**) and MGNs
(**D**) in DA2 (dark shade) and DL5 (light shade)

**Supplementary Table S1: Relative differences of innervation and synaptic composition**
**between glomeruli DA2 and DL5**

The Table lists the relative differences between DA2 and DL5 (see Methods for calculations).
Values that are at least 20% greater in DA2 than in DL5 are highlighted in dark shades and values
that are at least 20% less in DA2 than in DL5 are highlighted in light shades.

**Figure 3 – figure supplement 1: Synaptic density along neuronal fibers in DA2 and DL5**

Counts of synaptic inputs, synaptic outputs and T-bars normalized to 1 µm of neuronal length
along OSN, uPN or MGN fibers and collectively for all neurons within glomeruli DA2 (dark colors)
and DL5 (light colors). DA2: OSNs (green) n= 44; uPNs (red) n= 7; MGNs (blue) n=180; all neurons
n=231. DL5: OSNs n=46; uPN n=1; MGNs n=221; all neurons n=268. Data represent mean +
standard deviation (error bars). Data points represent single values. Means are compared using
either Student's T-test (in OSNs) or Wilcoxon two-sample test (in MGNs and all neurons). The
uPNs of the DA2 are not compared to the single uPN of the DL5. Significance value: $p>0.05$ (not
significant, no star), $p≤0.05$ (*), $p≤0.01$ (**), $p≤0.001$ (***). Values are listed in Table 1, row 9-11.

**Figure 4 – figure supplement 1: Properties of ipsi- and contralateral OSNs.**

**A:** Boxplots for total neuronal-fiber length and synaptic density (inputs, outputs, T-bars per unit
of neuronal fiber length) of ipsilateral (dark green) and contralateral OSN terminals (light green).
Dots represent single values. Means were compared using Stu
2110

dent's T-test. Significance value: p≤0.05 (*), p≤0.01 (**), p≤0.001 (***).

**Supplementary Table S2**: **Synaptic connectivity and relative differences between DA2, DL5 and**
**VA1v**

Synapse counts and synaptic strength of each connection type in DA2, DL5 and VA1v. Three
comparisons are shown: DA2 compared with DL5 (top table), VA1v with DL5 (middle) and VA1v
with DA2 (bottom). The relative synaptic strength (rel syn strength) of each connection type is
listed on the left side and the relative differences (rel differences) are listed on the right side.

**Supplementary Table S3: Connectivity of single neurons in DA2** (see extra file)

**Supplementary Table S4: Connectivity of single neurons in DL5** (see extra file)

**Figure 7 – figure supplement 1: Distribution of synapses and autapses along the DL5 uPN**
**dendrite in DL5**
**A**: 3D-reconstruction of the dendritic tree of the single uPN in glomerulus DL5 shown as skeleton
trace with artificial thickness according to Strahler order. Autapses are shown as cyan dots. The
entry site of the uPN main dendrite into the glomerulus (point closest to the soma) is the basal
root node. **B**: Number of autaptic presynaptic (red) and postsynaptic sites (cyan) according to
their geodesic distance to the basal root node point (indicated with a black circle in A). **C**:
Proportional distribution of all presynapses and postsynapses (excluding autaptic connections) in
the DL5 uPN at each strahler order (see legend inset). Note the high proportion of postsynaptic
sites on most distal dendritic branches.

2130
2131 **Figure 2 – figure supplement 1**

| Row | Values | Unit | OSNs | uPNs | MGNs | all neurons |
|---|---|---|---|---|---|---|
| | | | \multicolumn Relative differences between DA2 and DL5 | | | |
| 1 | **Total neuronal length** | µm | -26 | -7 | -26 | -22 |
| 2 | | input | -20 | -2 | -20 | -15 |
| 3 | **Total synaptic counts** | output | -2 | -48 | -16 | -16 |
| 4 | | T-bars | -12 | -47 | -20 | -22 |
| 5 | **Total innervation density (sum of length of all neuronal fibers/glomerular volume)** | µm/µm³ | 20 | 51 | 21 | 27 |
| 6 | | inputs/µm³ | 30 | 60 | 30 | 39 |
| 7 | **Total glomerular synaptic density (total synaptic counts/glomerular volume)** | outputs/µm³ | 59 | -15 | 36 | 36 |
| 8 | | T-bars/µm³ | 42 | -13 | 31 | 27 |
| 9 | | inputs/µm | 8 | 5 | 5 | 6 |
| 10 | **Neuronal synaptic density (synaptic counts/neuronal length)** | outputs/µm | 29 | -47 | 2 | 21 |
| 11 | | T-bars/µm | 16 | -45 | -5 | 7 |
| 12 | **Synaptic ratio** | T-bars/inputs | 3 | -47 | -6 | 2 |
| 13 | | outputs/inputs | 14 | -50 | -6 | 11 |
| 14 | **Polyadicity** | outputs/T-bars | 11 | -4 | 22 | 22 |

2132    **Supplementary Table S1**

2133
2134 **Figure 3 – figure supplement 1**

2135
2136 **Figure 4 – figure supplement 1**

| neuron class | connection motif | DA2 | rel syn strength (%) | rel out classes (%) | rel in classes (%) | DL5 | rel syn strength (%) | rel out classes (%) | rel in classes (%) | rel difference (%) | rel out difference (%) | rel in difference (%) |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| OSNs | OSNs>uPNs | 2365 | 19.99 | | | 2354 | 17.76 | | | 12.52 | | |
| | OSNs>MGNs | 3186 | 26.92 | 49.48 | 7.37 | 3201 | 24.15 | 43.44 | 7.89 | 11.47 | 13.93 | -6.63 |
| | OSNs>OSNs | 305 | 2.58 | | | 202 | 1.52 | | | 69.11 | | |
| uPNs | uPNs>OSNs | 16 | 0.14 | | | 26 | 0.20 | | | -31.08 | | |
| | uPNs>MGNs | 1205 | 10.18 | 10.92 | 32.62 | 2240 | 16.90 | 17.50 | 28.07 | -39.75 | -37.63 | 16.18 |
| | uPNs>uPNs | 71 | 0.60 | | | 54 | 0.41 | | | 47.26 | | |
| MGNs | MGNs>OSNs | 551 | 4.66 | | | 818 | 6.17 | | | -24.56 | | |
| | MGNs>uPNs | 1424 | 12.03 | 39.60 | 60.01 | 1313 | 9.91 | 39.06 | 64.03 | 21.47 | 1.38 | -6.28 |
| | MGNs>MGNs | 2711 | 22.91 | | | 3046 | 22.98 | | | -0.32 | | |
| | SUM | 11833 | 100 | 100 | 100 | 13254 | 100 | 100 | 100 | | | |

2137 **Supplementary Table S2**

| neuron class | connection motif | VA1v | rel syn strength (%) | rel out classes (%) | rel in classes (%) | DL5 | rel syn strength (%) | rel out classes (%) | rel in classes (%) | rel difference (%) | rel out difference (%) | rel in difference (%) |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| **OSNs** | **OSNs>uPNs** | 7226 | 22.26 | | | 2354 | 17.76 | | | 25.35 | | |
| | **OSNs>MGNs** | 10295 | 31.72 | 59.84 | 15.51 | 3201 | 24.15 | 43.44 | 7.89 | 31.33 | 37.76 | 96.53 |
| | **OSNs>OSNs** | 1901 | 5.86 | | | 202 | 1.52 | | | 284.30 | | |
| **uPNs** | **uPNs>OSNs** | 117 | 0.36 | | | 26 | 0.20 | | | 83.76 | | |
| | **uPNs>MGNs** | 1801 | 5.55 | 6.83 | 30.67 | 2240 | 16.90 | 17.50 | 28.07 | -67.17 | -60.96 | 9.26 |
| | **uPNs>uPNs** | 300 | 0.92 | | | 54 | 0.41 | | | 126.86 | | |
| **MGNs** | **MGNs>OSNs** | 3016 | 9.29 | | | 818 | 6.17 | | | 50.56 | | |
| | **MGNs>uPNs** | 2430 | 7.49 | 33.33 | 53.82 | 1313 | 9.91 | 39.06 | 64.03 | -24.42 | -14.68 | -15.96 |
| | **MGNs>MGNs** | 5371 | 16.55 | | | 3046 | 22.98 | | | -27.99 | | |
| | **SUM** | 32457 | 100 | 100 | 100 | 13254 | 100 | 100 | 100 | | | |

**Supplementary Table S3**

2140

| neuron class | connection motif | VA1v | rel syn strength (%) | rel out classes (%) | rel in classes (%) | DA2 | rel syn strength (%) | rel out classes (%) | rel in classes (%) | rel difference (%) | rel out difference (%) | rel in difference (%) |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| **OSNs** | **OSNs>uPNs** | 7226 | 22.26 | | | 2365 | 19.99 | | | 11.40 | | |
| | **OSNs>MGNs** | 10295 | 31,.2 | 59.84 | 15.51 | 3186 | 26.92 | 49.48 | 7.37 | 17.82 | 20.92 | 110.48 |
| | **OSNs>OSNs** | 1901 | 5.86 | | | 305 | 2.58 | | | 127.25 | | |
| **uPNs** | **uPNs>OSNs** | 117 | 0.36 | | | 16 | 0.14 | | | 166.60 | | |
| | **uPNs>MGNs** | 1801 | 5.55 | 6.83 | 30.67 | 1205 | 10.18 | 10.92 | 32.62 | -45.51 | -37.41 | -5.96 |
| | **uPNs>uPNs** | 300 | 0.92 | | | 71 | 0.60 | | | 54.06 | | |
| **MGNs** | **MGNs>OSNs** | 3016 | 9.29 | | | 551 | 4.66 | | | 99.57 | | |
| | **MGNs>uPNs** | 2430 | 7.49 | 33.33 | 53.82 | 1424 | 12.03 | 39.60 | 60.01 | -37.78 | -15.84 | -10.33 |
| | **MGNs>MGNs** | 5371 | 16.55 | | | 2711 | 22.91 | | | -27.76 | | |
| | **SUM** | 32457 | 100 | 100 | 100 | 11833 | 100 | 100 | 100 | | | |

2141

2142

| color code | rel syn strength (%) | <5 | 5-10 | 10-15 | 15-20 | >20 |
|---|---|---|---|---|---|---|
| | rel difference (%) | <10 | 10-20 | 20-50 | 50-100 | >100 |

2144 **Supplementary Table S4**

**A**

**B**

**C**

Synapse distribution along dendrite

2145
2146    **Figure 7 – figure supplement 1**