1   **Phylogeographic breaks and how to find them: An empirical attempt at separating vicariance**
2   **from isolation by distance in a lizard with restricted dispersal.**
3
4   **Running title:** *Acanthodactylus erythrurus* phylogeography
5
6

7   Loïs Rancilhac[1,2,3*$], Aurélien Miralles[4*], Philippe Geniez[5], Daniel Mendez-Aranda[6], Menad
8   Beddek[1,7], José Carlos Brito[8,9,10], Raphaël Leblois[11], Pierre-André Crochet[1]
9
1 CEFE, CNRS, Univ Montpellier, EPHE, IRD, Montpellier, France
2 Zoologisches Institut, Technische Universität Braunschweig, Mendelssohnstr. 4, 38106,
Braunschweig, Germany
3 Animal Ecology, Institute of Ecology and Genetics, Evolutionary Biology Centre, Uppsala
University, Uppsala, Sweden
4 Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle,
CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France
5 CEFE, Univ Montpellier, CNRS, EPHE-PSL University, IRD, Biogéographie et Ecologie des
Vertébrés, Montpellier, France
6 Max Delbrueck Center for Molecular Medicine in the Helmholtz Association, Robert-Roessle-Str.
10, 13092, Berlin, Germany
7 Naturalia Environnement, Site Agroparc, Avignon, France
8 CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório
Associado, Campus de Vairão, Universidade do Porto, Vairão, Portugal
9 BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão,
Vairão, Portugal
10 Department of Biology, Faculty of Sciences, University of Porto, Porto, Portugal
11 CBGP, INRAE, CIRAD, IRD, Institut Agro, Montpellier SupAgro, Univ. Montpellier,
Montferrier-sur-Lez Cedex, France
29
30   ***These authors contributed equally to this work**
31   **$Corresponding author: Loïs Rancilhac, loisrancilhac@gmail.com**

Abstract

Aim

Discontinuity in the distribution of genetic diversity (often based on mtDNA) is usually interpreted as evidence for phylogeographic breaks, underlying vicariant units. However, a misleading signal of phylogeographic break can arise in the absence of barrier to gene flow, under mechanisms of isolation by distance (IBD). How and under which conditions phylogeographic breaks can be reliably differentiated from populations evolving under IBD remain unclear. Here, we use multi-locus sequence data from a widely distributed lizard species to address these questions in an empirical setting.

Location

Morocco

Taxon

Spiny-footed lizard (*Acanthodactylus erythrurus*), Squamata: Lacertidae

Methods

Using 325 samples from 40 localities, we identified genetic discontinuities within *A. erythrurus* based on a mitochondrial fragment and nine nuclear markers independently. Using the nuclear markers, we then applied linear regression models to investigate whether genetic divergence could be explained by geographical distances alone, or barriers to gene flow (real phylogeographic breaks).

Result

A. erythrurus is characterized by an important mitochondrial diversity, with 11 strongly supported phylogeographic lineages with a crown age of 6 Mya. Nuclear markers, however, yielded weak phylogenetic support for these lineages. Using clustering methods based on genotypes at nine nuclear loci, we identified phylogeographic clusters that were partly discordant with the mtDNA lineages. Tests of IBD delimited at least four groups of populations separated by barriers to gene flow, but unambiguous separation of vicariance from IBD remained challenging in several cases.

Main conclusions

The genetic diversity of *A. erythrurus* originates from a mix of IBD and vicariance, which were difficult to distinguish, and resulted in similar levels of mitochondrial differentiation. These results highlight that phylogeographic breaks inferred from mitochondrial data should be further investigated using multi-locus data and explicit testing to rule out alternative processes generating discontinuities in mitochondrial diversity, including IBD. We identified four groups of populations within *A. erythrurus*, separated by barriers to gene flow, but even using nine independent nuclear makers the power of our approach was limited, and further investigation using genome-wide data will be required to resolve the phylogeographic history of this species.

Keywords: *Acanthodactylus*, Isolation by Distance, Morocco, Phylogeography, Vicariance, Admixture, Cytonuclear discordance, Phylogeographic breaks

Introduction

One of the main goals of phylogeography is to identify discontinuities in the distribution of genetic diversity across conspecific populations – phylogeographic breaks – and investigate the biotic and abiotic factors underlying them (Kumar & Kumar, 2018). Typically, phylogeographic breaks are interpreted as evidence for the existence of evolutionary lineages that diverged through vicariance events (Kuo & Avise, 2005). As such, they have been widely used to investigate biogeography and evolutionary biology questions, including the impact of geological events and landscape features on populations connectivity (e.g., Leaché et al., 2020), and their respective roles in promoting divergence and speciation (Huang, 2020). Furthermore, phylogeographic breaks are of interest for other fields of study, such as taxonomy and conservation biology (Braby et al., 2012; Shaffer et al., 2015). However, correct identification of phylogeographic breaks remains a non-trivial problem (Audzijonyte & Vrijenhoek, 2010).

Historically, phylogeographic studies have heavily relied on uniparentally inherited markers, such as mitochondrial DNA (mtDNA), to infer patterns of genetic diversity (Avise et al., 1987). However, while being characterized by a number of advantages (e.g., high mutation rate, lower effective population size, established standardized sequencing protocols), these markers often yield misleading signals of phylogeographic breaks (or the absence thereof) due to a number of processes (reviewed in Toews & Brelsford, 2012). Among them, different selective pressures (Irwin, 2012), introgression (e.g., Zieliński et al., 2013) and sex-biased dispersal (Palumbi & Baker, 1994) are well documented, while incomplete lineage sorting is not expected to generate predictable, biogeographically discordant patterns (Funk & Omland, 2003; Toews & Brelsford, 2012). Conversely, breaks in mitochondrial genetic diversity can also appear in continuous populations solely as a result of isolation by distance (IBD; Irwin, 2002), especially in the presence of sampling gaps (Audzijonyte & Vrijenhoek, 2010), and mtDNA is often poorly suited to detect IBD patterns (Teske et al., 2018). Additionally, mtDNA gene trees often differ from the population tree, mainly due to introgression and incomplete lineage sorting (ILS), hence phylogeographic breaks identified from mtDNA might not reflect actual populations splits (e.g., Dufresnes, Nicieza et al., 2020; Velo-Antón et al., 2018).

The limitations of mitochondrial phylogeography are usually overcome by considering multiple unlinked nuclear loci, which enable more accurate inferences (Brito & Edwards, 2009), particularly in the framework of the coalescent theory (Knowles, 2009). This approach reduces the likelihood of detecting spurious phylogeographic breaks, as independent loci are not expected to support concordant breaks in the absence of barriers to gene flow (Avise & Ball, 1990; Kuo & Avise, 2005). However, populations evolving along a single geographical gradient (i.e., under IBD) might still be apparently divided into distinct lineages, for example in the presence of large

sampling gaps (Schwartz & McKelvey, 2009; Tolley et al., 2022; Wiemers & Fiedler, 2007). In that

context, it appears crucial to test explicitly whether apparent phylogeographic breaks (e.g.,

reciprocally monophyletic groups in phylogenetic trees) delimit vicariant units separated by barriers

to gene flow or have been generated by alternative processes such as IBD. A commonly used

approach is to study the correlation between genetic and geographical distances among populations

or individuals in a statistical framework (e.g., Ramachandran et al., 2005), to determine whether the

null hypothesis that all populations differentiate along the same IBD gradient can be rejected, which

would indicate the presence of a barrier to gene flow (e.g., Binks et al., 2019; Hausdorf & Hennig,

2020; Goudarzi et al., 2019). Other popular approaches include the estimation of migration rates

over geographical surfaces, with low migration rates highlighting barriers to gene flow (Petkova et

al., 2016). However, in practice, the implementation of such tests is often limited in

phylogeographic studies by sparse sampling schemes and the low number of genetic markers

available. Here, we investigate the extent to which vicariant units can safely be identified from

populations evolving along IBD gradients, using multi-locus data of a genetically diverse North

African lizard.

Compared to other regions of the Mediterranean Basin biodiversity hotspot, fewer

phylogeographic studies have focused on North African communities (Beddek et al., 2018;

Husemann et al., 2014). Among these, many studies have revealed important intra-specific genetic

diversity in various taxa (see Husemann et al., 2014 for a review), in relation with the range of

landscapes and bioclimatic regions and the complex biogeographic history that characterize this

area. Consequently, important gaps still hamper our knowledge of the taxonomic diversity of North

African taxa and the evolutionary processes which created it (Beddek et al., 2018; Brito et al., 2014;

Ficetola et al., 2013). Phylogeographic studies of North African taxa are needed to fill the

aforementioned knowledge gaps, by documenting the extent and geographic distribution of intra-

specific genetic diversity, and identifying unrecognized species-level lineages. Secondly,

biodiversity hotspots such as North Africa are prime areas to study the ecological and evolutionary

processes shaping the distribution of genetic diversity and underlying lineages diversification at a

global scale (e.g., Carnaval et al., 2009; Demos et al. 2014, Durrant et al. 2015). In that context,

regionally distributed (as opposed to narrow-ranged) taxa are particularly valuable models, as they

are likely to inform on a larger variety of processes.

In this study, we focus on the Red-tailed Spiny-footed lizard *Acanthodactylus erythrurus*

(Squamata: Lacertidae), a species widely distributed across the Maghreb and the Iberian Peninsula.

This species is characterized by pronounced morphological variations that led to the recognition of

several morphological subspecies or species (Salvador, 1982; Bons & Geniez, 1995), which inhabit

distinct habitat types (from coastal sand dunes to high altitude xerophytic vegetation through open

forests and semi-arid steppes). Several molecular studies (Beddek et al., 2018; Fonseca et al., 2009;

Miralles et al., 2020) have revealed an important and previously unsuspected intraspecific genetic

diversity, leading to the description of two new species, and emphasizing the need for a more in-

depth taxonomic revision of this group. Notably, these studies highlight important disagreements

between the accepted taxonomy and molecular clades. One of the main questions raised by these

findings is whether the morphological and ecological diversification of this group has been

produced by vicariance and lineage diversification, or by local adaptation in the face of historical

gene flow. However, most studies (except Miralles et al., 2020) have relied only on mtDNA,

offering an incomplete picture of the historical processes underlying *A. erythrurus*' genetic

diversity. Here, we complement the multi-locus data produced by Miralles et al. (2020) with new

sequences, focusing on Moroccan populations with a comprehensive sampling, to clarify whether

multiple vicariant units explain the genetic diversity in the *A. erythrurus* complex (referred to as *A.*

erythrurus here).

170   The goals of this study are 1) to identify candidate phylogeographic breaks among

Moroccan populations of *A. erythrurus* using both mtDNA and nuclear DNA (nDNA) markers; 2)

to determine whether these candidate breaks delimit vicariant units or could result from IBD, and 3)

integrate these processes into a phylogeographic scenario to explain *A. erythrurus*' present genetic

diversity. Across the range of *A. erythrurus,* we decided to focus on Moroccan populations to

mitigate the effect of sampling gaps, as we only had sparse sampling in other countries. We

employed a two-step approach to untangle the processes at play: first, we analyzed mitochondrial

and nuclear (nine gene sequence fragments) data independently to identify phylogeographic

lineages. In both cases, phylogeographic lineages were defined as the smallest genetically

homogeneous geographical units. Secondly, we used the nuclear dataset to test whether the

described lineages corresponded to vicariant units or could result from IBD alone. Through this

approach, we aim at providing a more comprehensive picture of the phylogeographic history of

Moroccan *A. erythrurus*, but also empirically assess our ability to identify vicariant units using

multi-locus data in a genetically variable species.

Material & Methods

1. Sampling and DNA sequencing

189   We used a multi-locus dataset of 325 samples from the Moroccan populations of *A.*

erythrurus. Among these, 128 samples have been previously used by Miralles et al. (2020) in their

revision of the whole *A. erythrurus* complex using samples from its whole range (Algeria, Morocco,

Portugal, Spain, Tunisia). The remaining 197 have never been analyzed in any publication so far. In

the present study, we focus on the Moroccan populations of *A. erythrurus* because they offer the

most complete sampling. These 325 samples represent the Moroccan localities of the "Ibero-

Moroccan" (IM) clade defined by Miralles et al. (2020), but with significantly more samples per

locality. The other Moroccan populations historically assigned to *A. erythrurus* (Western and

Eastern High Atlas clades of Miralles et al., 2020) represent distinct species described as such by

Miralles et al. (2020). They are isolated from the IM clade by strong barriers to gene flow and were

thus not considered here. Ten additional samples from other clades and species were also included

to be used as outgroups in phylogenetic analyses, as described below. Details on the localities and

number of samples for the present study are given in Figure 1 and Table S1.

All samples have been sequenced for DNA fragments from one mitochondrial gene (776 bp

of the NADH-dehydrogenase subunit 4 [ND4] gene) and nine nuclear protein-coding genes:

Recombination Activating Gene 2 (RAG2, 321 pb); Oocyte maturation factor (C-mos, 369 pb),

Ornithine Decarboxylase (OD, 396 pb), Phosphogluconase Dehydrogenase intron 7 (PgD7, 420 pb),

Melanocortin Receptor 1 (MC1R, 630 pb), Pyruvate Kinase Muscle (PKM, 351 pb*),* Nuclear

transcription factor Y gamma (NFYC, 666 pb), Acetylcholinergic receptor M4 (ACM4, 390 pb) and

RNA fingerprint protein 35 (R35, 588 pb). Detailed sampling and laboratory protocols are given in

Miralles et al. (2020). All sequences were submitted to Genbank, and individual accession numbers

are available in Appendix 1.

Finally, for the ingroup samples, individual genotypes were inferred at each nuclear locus using the

software PHASE (Stephens et al., 2001) implemented in DNAsp v. 5 (Librado & Rozas, 2009). The

samples were grouped based on their mitochondrial lineages (cf. Results) to account for the

assumption that all samples are drawn from a single population at Hardy-Weinberg equilibrium

(Stephens et al., 2001). Such approach could be problematic if individuals from separated

populations shared closely related mtDNA haplotypes (e.g., in the case of mitochondrial

introgression between two sampled populations). However, preliminary analyses confirmed that

samples sharing similar mtDNA haplotypes were always closely related in the nDNA data. PHASE

was run on each group separately, with 5,000 sampled steps and 10,000 burnin-steps in the MCMC.

Uncertain phases (i.e., Posterior Probability < 0.90) were treated as unresolved sites using PHASE's

default settings.

2. Mitochondrial phylogeography

As a first step to identify the lineages used as basic units for the subsequent analyses, we

conducted phylogenetic inference based on 776 bp of the ND4 mitochondrial gene. In addition to all

our Moroccan samples, two samples from Portugal and Spain were also included to allow

estimation of the divergence times based on the opening of the Gibraltar straight (more details

below). Finally, the tree was rooted by including representatives of each of the following lineages (for the lineages nomenclature, we follow Miralles et al., 2020): *A. savignyi* (N=1), *A. margaritae* (N=1), *A. lacrymae* (N=2), the AT lineage from eastern Algeria and Tunisia (including *A. blanci,* N=2), *A. montanus* (N=2) and the CA lineage from central Algeria (N=2). In total, 331 samples were included in the final alignment (ND4 sequencing failed for four ingroup samples).

Maximum-likelihood (ML) phylogenetic inference was performed using IQTREE v. 1.6.8 (Chernomor et al., 2016; Nguyen et al., 2015). The best substitution model was selected using ModelFinder (Kalyaanamoorthy et al., 2017) as implemented in IQTREE, and the branches support was assessed using the SH-like approximate likelihood test (aLRT) with 5,000 pseudoreplicates. Based on this tree, we defined phylogeographic lineages corresponding to the smallest geographical units supported as monophyletic groups (in practice, with an aLRT support > 90). Finally, we estimated divergence times through two approaches. First, we calibrated the divergence between the Iberian populations and their closest Moroccan relatives based on the opening of the Strait of Gibraltar (between 5.3 and 5.5 Million years ago [Mya]). However, as more recent colonization events across the Strait of Gibraltar have been found in reptiles and amphibians (e.g., Carranza et al., 2006; Gutiérrez-Rodríguez et al., 2017), we also calibrated the tree based on a fixed mutation rate for the ND4 gene. Details on the procedure can be found in Supplementary materials S1.

3. Nuclear phylogeography

Preliminary analyses revealed that the phylogeographic signal in the nuclear data was much weaker compared to that of the mitochondrial dataset, with much higher levels of allele sharing among populations and lower nodes support in phylogenies. Our initial approach to infer genetic clusters and admixture among them was to use the software STRUCTURE (Pritchard et al., 2000). However, we failed to reach proper convergence of the MCMC chains and to identify the optimal number of clusters (K). Furthermore, for a single K value, independent runs yielded very different, yet equally likely, clustering solutions (results not included). These solutions suggested clearly separated groups (i.e., high Q values), but their composition varied across runs, and did not highlight a clear geographic signal. Likewise, phylogenetic reconstructions of the concatenated data failed to converge toward a consistent topology. In that context, it was difficult to apply a strict criterion to delimit phylogeographic lineages from the nuclear dataset. Thus, we relied on congruent results from several analyses, especially multivariate analyses of the genotypes at the nine nuclear loci, as described below.

3.1 Phylogenetic analyses

We first performed ML phylogenetic inference using the concatenation of the nine nuclear

loci. Best substitution models and partition schemes were selected on a per-locus partition, and the

analyses were run in IQTREE v. 1.6.8 with the same settings as for the mitochondrial analyses. Five

independent ML searches were performed.

In addition, we also perform Neighbour-Joining (NJ) inference based on between-populations (i.e.,

sampling localities) pairwise distance matrices for the ingroup samples only. This analysis was

repeated twice, firstly using average p-distances calculated from the concatenation of the nine

markers, secondly using Nei's distances computed from the genotypes at the nine loci. In the first

case, both distances computation and tree inference were performed in MEGA 11 (Tamura et al.,

2021); in the second case Nei's distances were calculated using adegenet v. 2.1.5 (Jombart, 2008)

and NJ inference performed with ape v. 5.6 (Paradis & Schliep, 2019).

Because our attempts to infer a phylogenetic tree from the nuclear markers yielded poorly

supported and variable topologies, in which the position of the Iberian samples is different from that

of the mitochondrial tree (see Results and Figure S5), we did not attempt to calibrate the nuclear

phylogeny. Furthermore, we spent a substantial amount of time trying to get a species tree from the

nine nDNA markers using *BEAST (Drummond et al., 2012) but did not manage to reach proper

convergence and obtained variable topologies across runs.

3.2 Multivariate analyses

As an alternative to phylogenetic analyses, we used Multiple Correspondence Analyses

(MCA) to detect phylogeographic breaks based on the nuclear dataset. MCAs were calculated from

the genotypes at the 9 nuclear loci, using the software Genetix (Belkhir, 2004). We used an iterative

approach to define phylogeographic lineages: the analysis was first run on a dataset including all

Moroccan populations (same sampling as for the phylogenetic analyses but without outgroups and

Iberian samples); the results were visually inspected and groups of populations that were clearly

differentiated from the rest were identified as phylogeographic lineages. These populations were

then removed from the dataset, and the analysis was run again on the resulting subset. This process

was repeated until no clear structure could be identified in the results (in total, five iterations were

needed, see Results).

4. Tests of Isolation by Distance and vicariance

Population structure analyses based on both mtDNA and nDNA data supported several

genetic groups in Moroccan *A. erythrurus*. While the structure inferred from both types of markers

was mostly concordant, there were a few notable differences. For further analyses, we considered

the nDNA genetic groups (summarized in the following, cf. Table 1 and Figure 2 for more details),

which minimized the number of clusters represented by a single locality. Indeed, the employed

statistical test relies on calculating a regression on within-group geographic distances, which is not

possible for single localities. More precisely, we defined four inland groups: the Rif Mountains (Rif,

including N-Rif and S-Rif), the Middle-Atlas Mountains (Mid-Atl, including Mid-Atl1 and Mid-

Atl2), the surroundings of Marrakech (Mara), and the population of Sidi-Yahia (S-Yah); as well as

six groups distributed along the Atlantic coast, most of which consist in single localities, except for

the northernmost (N-Atl) and southernmost (S-Atl) groups. To determine whether this pattern of

genetic diversity reflects true phylogeographic breaks, we investigated whether these genetic groups

differentiate along similar geographical gradients (i.e., IBD gradients). The populations were

divided into two groups, analyzed separately for practical purposes: 1) the inland lineages and 2) the

lineages distributed along the Atlantic coast. All scripts and data files used to run these analyses are

available at https://github.com/rancilhac/IBD_vs_Vicariance

(http://doi.org/10.5281/zenodo.7142595).

4.1 Differentiation between inland lineages

To assess whether the Rif, Mid-Atl and Mara groups are vicariant units, we analyzed the

correlation between genetic and geographical distances in pairs of geographically adjacent lineages,

using the statistical framework of Hausdorf & Hennig (2020). This approach relies on the null

hypothesis that the two considered groups of population differentiate along the same IBD gradient.

Testing this relies on a two-steps approach: firstly, we tested that the within-groups distances of

both lineages followed similar regressions (hypothesis *H01* of Hausdorf & Hennig, 2020). In case

this was verified, we then tested whether regressions fitted on between-groups distances only and

all within-group distances are similar (hypothesis *H02* of Hausdorf & Hennig, 2020). A rejection of

that hypothesis (i.e. within-groups and between-groups distances do not fit the same regression line)

would support a barrier to gene flow between the tested groups. However, this test is not

appropriate if *H01* is not verified. In such case, we tested whether between-groups distances follow

the same regression as the within-group distances of one of the two lineages (hypothesis *H03* of

Hausdorf & Hennig 2020). A verification of that hypothesis is incompatible with a barrier to gene

flow. When testing *H03*, we performed two tests, *H031* and *H032,* in which the within-group

distances used for the comparison are respectively those of group 1 and group 2. A failure to reject

both *H031* and *H032* would indicate inconclusive results deserving further investigation (Hausdorf

& Hennig, 2020). In all cases, the significance of the difference between regression lines is assessed

using a non-parametric jackknife test, as described in Hausdorf & Hennig (2020). We used this

approach to test the following combinations of lineages, corresponding to the inland lineages and
adjacent coastal lineages: Rif + Atl-N, Rif + Mid-Atl, Mid-Atl + Mara, Mid-Atl + (S-Atl, O-Gha, S-
Bou), Mara + S-Atl.

The case of S-Yah is more complicated to address, because this lineage is represented by a single
locality, thus the tests described in the previous section are not applicable. Considering the results of
previous analyses (cf. Results, Figure 2), we hypothesize that S-Yah is closely related to the Mid-
Atl group. To test whether S-Yah differentiate within the same IBD gradient as Mid-Atl, we tested
whether regressions within (Mid-Atl + S-Yah) and within Mid-Atl alone were similar. If this
hypothesis is verified, we interpret it as an indication for an absence of barrier to gene flow between
Mid-Atl and S-Yah.

All regression tests were performed using the prabclus R package (Hausdorf & Hennig,
2020) based on pairwise distances between individuals. In the specific case of S-Yah, the *regdistdiff*
function had to be adapted in a custom R script to match our needs. Geographic distance matrices
were calculated from coordinates using prabclus, and log transformed. Genetic distance matrices
were computed from the genotype data at the nine nuclear loci using two metrics: the $\hat{a}$ statistic (an
equivalent of $Fst/(1-Fst)$ calculated between individuals), calculated in Genepop (Rousset, 2008),
and the shared allele distance (s), calculated in prabclus. Each analysis was repeated twice, using
both measures of genetic distance. We decided not to use mitochondrial distances because empirical
evidence suggest that mitochondrial markers are not suited to detect IBD gradients (Audzijonyte &
Vrijenhoek, 2010; Teske et al., 2018). Furthermore, male-biased dispersal has been identified in
most Squamates studied to date (Ferchaud et al. 2015), meaning that maternally inherited markers
might not accurately represent dispersal patterns in *A. erythrurus*.

4.2 Isolation by distance along the Atlantic coast

Because several of the genetic groups along the Atlantic coast are represented by single
localities, testing for the presence of barriers to gene flow between each group pair is not
straightforward, as explained in the case of S-Yah (above). Thus, rather than testing each putative
phylogeographic breaks, we explored two hypotheses. First, considering that N-Atl is very distinct
from other coastal populations on multivariate analyses and population trees (where it appears to be
more related to the Rif, cf. Results), we hypothesize that there is a barrier to gene flow between N-
Atl and the population adjacent to the south, M-Bou. Testing this hypothesis is relatively
straightforward as the framework of Hausdorf & Hennig 2020 applies here, by testing (S-Atl, O-
Gha, S-Bou, Blad-Mehd, M-Bou) and N-Atl. However, it is also possible that there are more

barriers to gene flow along the coast, or none. To gain more insight into this, we first calculated a
regression between genetic and logarithm of geographical distances including all coastal lineages
(to the exclusion of N-Atl in case the previous test showed that it does not evolve along the same
IBD gradient as the southernmost populations). We then successively removed lineages and re-
calculated the regression. In case removing populations did not significantly change the regression,
we interpret this as an indication that all coastal lineages differentiate along a single IBD gradient.
All tests were performed in the same set up as detailed in the previous section.
379
380 **Results**
381
382 *1. Mitochondrial DNA supports deep phylogeographic lineages in Moroccan* **Acanthodactylus**
383 **erythrurus**
384
Using 776 bp of the ND4 mitochondrial gene, we recovered a relatively well supported
phylogenetic tree (Figure 1, Figure S1), fully consistent with the analyses of the same data by
Miralles et al. (2020). Particularly, the monophyly of the "Ibero-Moroccan" (IM) clade is fully
supported (aLRT = 100), and this group is further divided into 11 phylogeographic lineages with
aLRT support >90 (Figure 1, Table 1), namely (from north to south): 1) the northern Atlantic coast
(N-Atl, from Souk-Tnine to Larache); 2) the northern part of the Rif mountains (N-Rif, from Martil
to Ketama); 3) the southern part of the Rif (S-Rif, Tazzeka and the North of Ifrane); 4) the coastal
population of Moulay-Bousselham (M-Bou); 5) the coastal populations of Blad-Dendoun and
Mehdia (Blad-Mehd); 6) the inner-most parts of the Middle-Atlas mountains (Mid-Atl1, around
Azrou and Ifrane); 7) the rest of the Middle-Atlas mountains (Mid-Atl2, from Oulmès to
Ouaourioud and Boumia); 8) the population of Sidi-Yahia (S-Yah); 9) the coastal population of
Sidi-Boumaine (S-Bou); 10) the populations surrounding Marrakech (Mara; including Sidi-Chiker
and Timlite) and 11) the Southern Atlantic coast (S-Atl; from Ouled-Ghanem south to Essaouira).
All lineages are geographically isolated by sampling gaps, except for S-Rif and Mid-Atl1, which
come into contact north of Ifrane (Middle-Atlas). Of the three samples from this locality, two
cluster with the former and one with the latter (see Doniol-Valcroze et al. 2022 and Figure S1; the
correspondence between samples and localities is given in Appendix 1).
The 11 lineages cluster into three main clades: i) Mara, S-Atl, S-Bou; ii) Rif N, Rif S (as
well as the Iberian samples) and iii) N-Atl, M-Bou, Blad-Mehd, Mid-Atl1, Mid-Atl2. The aLRT
support for these clades ranges from 77 to 100, but the relationships among them remain
unresolved. Finally, S-Yah harbors a remarkably divergent mitochondrial lineage which is basal to
all other Moroccan mitochondrial clades of *A. erythrurus*.

Figure 1. *A) Simplified phylogenetic tree obtained from the maximum likelihood analysis of 776 bp*
of the ND4 mitochondrial fragment. The outgroups used for rooting (10 samples of closely related
A. erythrurus *lineages and five other* Acanthodactylus *species) were removed from the figure and*
terminal groups were pruned for improved graphical resolution (for full tree, see Figure S1).
Numbers at the nodes show aLRT support Numbers on terminal branches show the aLRT support of
collapsed terminal clades. Color boxes show the 11 identified phylogeographic lineages. B)
Neighbour-Joining population tree inferred from the pairwise Nei's distances between populations,
calculated from the genotypes at nine nuclear loci, with colors referring to the mitochondrial
lineages. C) Map of Morocco showing the sampled localities, with colors and texts showing the
mitochondrial lineages they belong to. Localities are numbered as in Table 1, with coordinates and
sample sizes given in Table S1.

The time-tree analyses based on the end of the Messinian crisis (Figure S2) inferred an

origin of the "IM" clade at 5.99 Mya (95% credibility interval [5.36 – 7.07]). Inferences with a

fixed mutation rate (Figures S3 & S4) yielded slightly older dates (6.05 and 7.12 Mya based on the

mutation rate of *Podarcis* and *Gallotia* respectively), but the 95% confidence intervals were much

wider. Overall, the three calibration approaches gave similar dates. However, these results should be

taken with care since the analyses failed to reach proper convergence (see Supplementary materials

S1 for more details).

Table 1. *Summary of the phylogeographic lineages delimited using 776 bp of the ND4*

mitochondrial gene, and genotypes at nine nuclear loci. Numbers between brackets in the fourth

column refer to the localities numbering on Figure 1C and Table S1. The full sample list for each

locality is given in Appendix 1.

Mitochondrial lineages	Nuclear lineages	Geographic range	Localities
N-Atl	N-Atl	Northernmost Atlantic coast	Larache, Souk-Khemis, Souk-Tnine [1-3]
N-Rif	Rif	Northern part of the Rif mountains	Ketama, Martil, Bab-Taza [4-6]
S-Rif		Northern parts of Middle-Atlas	Tazzeka, Ifrane North [7-8]
M-Bou	M-Bou	Single locality on the North of the Atlantic coast south of Larache	Moulay-Bousselam [9]
Blad-Mehd	Blad-Mehd	Two neighbour localities on the Atlantic coast just north of Rabat	Blad dendoun, Mehdia [10-11]
Mid-Atl1	Mid-Atl	North-eastern parts of the middle-Atlas	Ifrane South, Azrou area [12-13]
Mid-Atl2		Southern and western parts of the middle-Atlas	Oulmès area, Boumia, Col Zad, Ouaourioud, Oued Zem [14-18]
S-Yah	S-Yah	Single locality just south of Rabat	Sidi-Yahia [19]
S-Bou	S-Bou	Single locality on the Atlantic coast south of Casablanca	Sidi-Boumaine [20]
Mara	Mara	Populations surrounding Marrakech, in southern Morocco	Marakech area, Sidi Chiker, Timlite [21-23]
S-Atl	O-Gha	Single locality on the Atlantic coast north of Essaouira	Ouled Ghanem [24]
	S-Atl	Populations of the southernmost Atlantic coast	Essaouira area, Souira [25-26]

2. Multi-locus nuclear phylogeography

The ML phylogenetic analysis of the concatenation of the nine nuclear markers yielded a

much less resolved topology compared to that of the mtDNA data (Figure S5). While the

monophyly of the IM clade is strongly supported (aLRT = 99), most of the mitochondrial lineages

are mixed and do not form reciprocally monophyletic groups. However, the following

mitochondrial lineages are also supported as monophyletic in the nuclear phylogeny: Mara (aLRT =

97); S-Bou (aLRT = 81); and a clade composed of M-Bou and N-Atl (aLRT = 90) within which the

former is paraphyletic but the latter is monophyletic (aLRT = 91). Finally, a clade comprising all N-

Rif and S-Rif samples to the exclusion of one from the contact zone with Mid-Atl1 is relatively well

supported (aLRT = 75), but also includes one sample from Mid-Atl1.

Multiple Correspondence Analyses (MCA) of the genotypes at the nine nuclear loci

recovered a phylogeographic structure partially concordant with that of the mitochondrial data

(Figure 2). With this approach, we could identify 10 phylogeographic lineages (from north to south,

the names refer to those used for the mitochondrial lineages, see also Table 1): 1) N-Atl; 2) the Rif

mountains (Rif, mitochondrial lineages N-Rif & S-Rif; including all three samples from IFRA-N);

3) M-Bou; 4) Blad-Mehd; 5) all the Middle Atlas populations (Mid-Atl, mitochondrial lineages

Mid-Atl1 & Mid-Atl2); 6) S-Yah; 7) S-Bou; 8) Mara; 9) the coastal locality of Ouled-Ghanem (O-

Gha, grouped within the S-Atl lineage in the mitochondrial data); and 10) S-Atl (to the exception of

O-Gha). Overall, these groups are mostly overlapping with the mitochondrial lineages, but are less

straightforward to delimit due to the weak phylogenetic resolution of the nuclear markers. The

locality O-Gha appears as an exception, being clustered with the other southern Atlantic populations

in the mitochondrial data but quite divergent in the nuclear data (Figure 2C). Interestingly, S-Yah

appears here closely related to Mid-Atl populations, from which it could be distinguished only at

the finer scale (Figure 2E), contrasting with its high mitochondrial divergence.

459
460
Figure 2. *Results of Multiple Correspondence Analyses calculated from the genotypes at the nine*
nuclear genes. The analysis was repeated five times iteratively with subsets of populations to detect
finer phylogeographic structure. Colors and names correspond to the mitochondrial lineages. In A,
the black ellipse highlights three samples from the contact zone between Rif and Mid-Atl. Blad-
Mehd is highlighted twice in D and E to illustrate its intermediate position between the Southern
Atlantic and Middle-Atlas populations.
467
Population NJ trees inferred from both p-distances (Figure S4) and Nei distances (Figure
1B) supported a structure similar to that of the MCA. The Rif, N-Atl and Mara groups are separated

from the others by relatively long branches (although Mara is closer to the other populations than to

Rif and N-Atl), while populations from the southern Atlantic coast, Mid-Atl and S-Yah appear

closely related. Both genetic distances did not yield completely concordant topologies, especially

regarding the position of M-Bou, which is alternatively placed close to N-Atl or S-Atl. On both

trees, Blad-Mehd has an intermediate position, suggesting an effect of admixture between several

genetic groups.

3. Isolation by Distance tests

IBD tests for pairs of inland lineages based on both $\hat{a}$ and s genetic distances were overall

congruent (Tables 2 & 3). Figures 3 & 4 display the regression lines based on $\hat{a}$, while those based

on s are provided in Figures S7 & S8.

Starting from the north, all tests congruently supported a barrier to gene flow between N-Atl

and the Rif (Fig. 3A). This was also the case when comparing Rif and Mid-Atl (Fig. 3B), to the

exception of one test: based on $\hat{a}$ genetic distances, we couldn't reject the null hypothesis *H032* that

the between-groups regression differed from the within-group regression of Rif, although this

difference was significative when considering the within-group regression of Mid-Atl (*H031*). The

results for Mid-Atl and Mara (Fig. 3C) were also ambiguous: based on the s genetic distance, we

found that the between-groups distances were different from the within-group distances of Mara,

but other tests (i.e., *H02* based on $\hat{a}$ and *H031* based on s) were non-significant. On the contrary,

tests between Mara and S-Atl (Fig. 3D) clearly supported a barrier to gene flow. Finally, tests based

on both $\hat{a}$ and s unambiguously supported that S-Atl and Mid-Atl (Fig. 3E) differentiate along the

same IBD gradient. While the differentiation between S-Yah and Mid-Atl could not be tested in the

same framework because S-Yah is represented by a single locality, we were not able to distinguish

regression lines calculated on the distances within Mid-Atl alone or Mid-Atl + S-Yah (Fig 3F),

which we interpret as an indication that they evolve along the same IBD gradient.

Figure 3. *Linear regressions between genetic (â) and geographical distances in pairs of*

geographically adjacent inland lineages. Black and red dots show within-group distances for both

lineages, and corresponding dashed lines represent linear regression fitted to them. Blue crosses

and solid lines show between-group distances. Between-group distances are not represented on the

last graph as the test was adapted to account for zero-only geographical distances in S-Yah.

On the Atlantic coast, tests based on both $â$ and s genetic distances suggested a barrier to

gene flow between N-Atl and the other coastal populations (Figure 4A, Table 2). However, these

results remain inconclusive, as the regression on between-groups distances was significantly

different from that on the within-group distances of N-Atl, but not the within-group distances of the

rest of the coastal populations. Our tests further suggested that all coastal populations south of N-

Atl differentiate along a single IBD gradient (Figure 4B).

508
509 ***Figure 4.*** *Isolation by distance tests along the Atlantic coast based on the â genetic distance. A)*
510 *Comparison between N-Atl (red) and the rest of the coastal populations (black). Blue cross show*
511 *between-groups distances and lines represent the respective regressions. B) Regressions on within-*
512 *group distances of several combinations of coastal populations South of N-Atl.*
513
514

		H01		*H02*		*H031*		*H032*	
Group 1	**Group 2**	$\hat{a}$	*s*	$\hat{a}$	*s*	$\hat{a}$	*s*	$\hat{a}$	*s*
Inland lineages									
N-Atl	Rif	0.35	0.40	5.69e-6	6.58e-6	-	-	-	-
Rif	Mid-Atl	2.96e-4	0.08	-	3.93e-7	2.17e-6	-	0.19	-
Mid-Atl	Mara	0.07	3.94e-5	0.55	-	-	0.91	-	0.025
Mara	S-Atl	1.25e-5	3.6e-13	-	-	4.14e-8	0.006	6.20e-8	6.48e-14
S-Atl	Mid-Atl	0.0005	0.0006	-	-	0.50	0.49	0.73	0.74
Coastal lineages									
N-Atl	(M-Bou + Bad-Mehd + S-Bou + O-Gha + S-Atl)	6.05e-20	1.33e-15	-	-	7.94e-13	3.02e-5	1.00	1.00

		p-value	
Group 1	**Group2**	$\hat{a}$	*s*
Inland lineages			
Mid-Atl	S-Yah	0.39	0.71
Coastal lineages			
(M-Bou + Blad-Mehd + S-Bou + O-Gha + S-Atl)	(Blad-Mehd + S-Bou + O-Gha + S-Atl)	0.12	0.28

(M-Bou + Blad-Mehd + S-Bou + O-Gha + S-Atl)	(S-Bou + O-Gha + S-Atl)	1.00	0.95
(M-Bou + Blad-Mehd + S-Bou + O-Gha + S-Atl)	(O-Gha + S-Atl)	1.00	0.38
(M-Bou + Blad-Mehd + S-Bou + O-Gha + S-Atl)	S-Atl	0.14	0.80
(Blad-Mehd + S-Bou + O-Gha + S-Atl)	(S-Bou + O-Gha + S-Atl)	0.77	1.00
(Blad-Mehd + S-Bou + O-Gha + S-Atl)	(O-Gha + S-Atl)	1.00	0.32
(Blad-Mehd + S-Bou + O-Gha + S-Atl)	S-Atl	0.88	0.54
(S-Bou + O-Gha + S-Atl)	(O-Gha + S-Atl)	1.00	0.91
(S-Bou + O-Gha + S-Atl)	S-Atl	0.41	1.00
(O-Gha + S-Atl)	S-Atl	0.63	1.00

Discussion

1. Mitochondrial and nuclear markers yield contrasting phylogeographic signals in Moroccan **Acanthodactylus erythrurus**

As already highlighted by Miralles et al. (2020), our analyses of 776 bp of the ND4 mtDNA gene supported an important genetic diversity in the Moroccan populations of *A. erythrurus*, which can be divided into several deep lineages. Our analyses recovered an origin of the Ibero-Moroccan clade ("IM" clade) around 6 Mya, a relatively old age for a species-level lineage in comparison with calibrations in *Podarcis* wall lizards (Salvi et al., 2021) and other Lacertids (Garcia-Porta et al., 2019; Joger et al., 2007; Tamar et al., 2016). Furthermore, the overall congruence between calibrations based on the end of the Messinian crisis and on fixed mutation rates seem to confirm that Iberian populations have been isolated since the re-opening of the Strait of Gibraltar, around 5.4

Mya. While the deepest nodes within the IM clade are unresolved, possibly as a result of rapid

diversification, or the limited informativeness of the sequence used, we could identify 11 strongly

supported (aLRT > 90) clades corresponding to geographical units (subsequently referred to as

mtDNA lineages) among the Moroccan populations. While some of these mtDNA lineages include

several populations distributed across relatively large areas, many well differentiated monophyletic

groups corresponded to single localities, especially along the Atlantic coast. It is remarkable that in

some cases geographically distant populations share closely related haplotypes (e.g., in the Mid-

Atl2 and S-Atl lineages), while in other instances localities that are close to each other cluster in

distinct lineages (e.g., S-Rif and Mid-Atl1 lineages meeting in Ifrane, and the localities in the

Northern half of the Atlantic coast).

This pattern could suggest the existence of many phylogeographic breaks across Morocco,

resulting from allopatric diversification following gene flow disruptions. Especially, in the Middle-

Atlas, two highly differentiated mitochondrial lineages (S-Rif and Mid-Atl1) enter in contact with

limited apparent admixture, a pattern consistent with allopatric divergence followed by secondary

contact (Avise et al., 1987). Alternatively, this pattern could also result from fine-scale barriers to

gene flow, including divergent ecological preferences between populations.

Contrastingly, our analyses of nine nuclear loci provided little support to the mtDNA

lineages. Several "state-of-the-art" approaches, including phylogeny and STRUCTURE clustering,

failed to recover consistent lineages, seemingly because of high levels of allele sharing among

populations. This result was surprising, as we expected that populations evolving allopatrically for

several millions of years would accumulate genome-wide mutations reflecting the mitochondrial

structure. The high levels of allele sharing could thus indicate ongoing nuclear gene flow between

lineages, or an important retention of ancestral polymorphisms due to large effective population

sizes (Pamilo & Nei, 1988). Given the matrilineal inheritance of the mitochondria, male-biased

dispersal could exacerbate differences in structure with nuclear markers (Palumbi & Baker, 1994).

Finally, the poor performance of STRUCTURE to infer genetic clusters could be due to the

presence of IBD, which breaks the model's assumptions (Perez et al., 2018), resulting in over-

estimating the number of genetic clusters in the data (Frantz et al., 2009; Schwartz & McKelvey,

2009; Serre & Pääbo, 2004).

Contrary to other approaches based on more constrained models (i.e., phylogenetic trees,

STRUCTURE), multivariate analyses of individual genotypes at the nine nuclear loci supported a

phylogeographic structure somewhat consistent with the mtDNA lineages. One important difference

was the position of the individuals from Sidi-Yahia (S-Yah, just south of Rabat), which harbors a

very distinctive mitochondrial lineage, branching as sister to all the other populations. However,

nuclear data suggest that these individuals are closely related to populations from the middle-Atlas

and the Atlantic coast. Two scenarios can be considered to explain this discrepancy: firstly, the

population of Sidi-Yahia could be closely related to its neighbours (i.e., Atlantic coast, Middle-

Atlas), with a mitochondrial genome introgressed from a distinct genetic pool, absent from our

analyses. Alternatively, Sidi-Yahia could be the remnant of an older lineage whose nuclear genome

got swamped by that of invading populations from the middle-Atlas or Atlantic coast. Either way, it

seems that this pattern is the legacy of an unsampled, possibly extinct *Acanthodactylus* lineage,

whose mitochondrial genome persisted to that day. The existence of "ghost mitochondrial lineages"

is well-documented in animals (for examples in reptiles, see Renoult et al., 2009; Salvi et al., 2017;

Schultze et al., 2020), and their persistence can result from various mechanisms, including local

adaptation (Carpio et al., 2021). Future studies of *A. erythrurus* based on genome-wide data might

be useful to understand the processes underlying such discordance.

582   At a larger scale, in the absence of a well resolved nDNA tree, it is difficult to assess the

extent of cytonuclear discordances in *A. erythrurus*. However, genetic groups identified from

mtDNA and nDNA are mostly concordant, the main difference residing in the lower phylogenetic

resolution yielded by nDNA markers. We hypothesize that this pattern could result from two

scenarios: 1) the mitochondrial lineages reflect phylogeographic breaks, and the poor resolution of

nDNA markers is the result of ancestral polymorphism retention; or 2) the mitochondrial lineages

are largely connected by nuclear gene flow. We tried to untangle these processes by studying the

distribution of genetic differentiation along geographical gradients.

2. The conundrum of separating vicariance from isolation by distance

Geographical and genetic distances were overall strongly correlated in our data, highlighting the

role of IBD in generating genetic diversity in *A. erythrurus*. However, IBD alone did not explain the

observed phylogeographic patterns. Using linear regressions, we could identify at least four

geographic groups separated by barriers to gene flow (summarized in Figure 5, discussed in-depth

in the next section), within which we could not reject the null hypothesis that populations

differentiate through IBD. It is interesting to note that our results suggest that similar levels of

mitochondrial divergence can arise from both IBD and allopatric divergence (and even allopatric

speciation, as highlighted at the Rif/Middle-Atlas transition by Doniol-Valcroze et al., 2022). In that

context, interpreting mitochondrial splits as evidence for phylogeographic breaks is potentially

misleading and biased toward an over-estimation of the number of phylogeographic lineages. While

contrasting mitochondrial and nuclear phylogeographic patterns can help determining the extent of

mitochondrial diversity that can be attributed to vicariance, this issue remains very challenging in

the absence of clear signal in nuclear markers. This problem is especially relevant as mitochondrial

markers are still widely used to infer biogeographic patterns and species boundaries. In many cases,

multi-locus studies employ only a single or few nuclear markers, whose signal gets eclipsed by that

of the more variable mtDNA fragments. Hence, we argue that a thorough interpretation of mtDNA

phylogeographic patterns should rely on a comparison with independently analyzed nuclear

markers.

As we show here, employing statistical, spatially explicit methods to detect discontinuities

in genetic diversity can yield valuable insights on the presence of phylogeographic breaks.

However, our analyses often yielded inconclusive results, even in a case where we expected the

signal for vicariance to be clear (i.e., the contact zone between Rif and Middle-Atlas, as inferred

from the low admixture limited to a single locality, despite geographic distances being < 1km),

suggesting low power of these methods even with nine independent markers. We observed a large

variance in genetic distances within populations, making our tests more prone to stochastic errors.

Employing more informative data (typically genome-wide markers) would likely reduce this

variance, yielding more power to distinguish regression lines.

More importantly, we identify the sampling scheme as a significant factor driving the results

of IBD tests. For example, it seems that the inconclusive results for the Rif/Middle-Atlas tests could

be due to the asymmetric sampling around the contact zone, with only a few samples available in

the former group, and a large sampling gap between the contact zone and the core area. In-depth

analyses of this transition by Doniol-Valcroze et al. (2022) also identified sampling gaps as a

limiting factor, even when considering genome-wide data. Likewise, our IBD tests for coastal

populations were very limited because of "lineages" represented by single localities, which made

thorough comparisons of within-groups and between-groups distances impossible. This issue could

be mitigated by sampling many localities, especially to fill gaps between identified genetic groups.

Finally, the IBD model used assumes that genetic and geographic distances follow a linear

relationship (Hausdorf & Hennig, 2020). Because most genetic distance metrics are comprised

between 0 and 1, this assumption is likely to be broken when genetic distances are high. Especially,

intra- and between-groups regressions can be difficult to distinguish when intra-group distances are

close to 1 (Hausdorf & Hennig, 2020). However, the linear relationship is expected at least in one

case, when considering log-transformed geographic distances and $Fst/(1-Fst)$ as an estimate of

genetic distance (and, by extension, the $\hat{a}$ genetic distance for individuals; Rousset, 1997). In our

case, intra-group distances were very variable and sometimes very large, especially in S-Atl and

Mid-Atl, which might result in false negatives or inconclusive results. Considering more markers

(e.g., genome-wide SNPs) might improve this issue by reducing the variability across individuals

and avoiding very high distances. Furthermore, violations of the linearity assumption might explain

the differences in results between the $\hat{a}$ and s genetic distances.

To conclude, our results demonstrate that identifying phylogeographic breaks is far from a trivial

issue and should not rely on mitochondrial markers alone. It would be interesting to further study

the factors underlying the concordance of mitochondrial breaks with vicariance events, especially

the effects of dispersal abilities and population sizes.

3. Mechanisms of diversification in the **Acanthodactylus erythrurus** *group*

Small terrestrial vertebrates, including lizards, often have low dispersal abilities, which sometimes

result in strong phylogeographic patterns (e.g., Alfonso Silva et al., 2017; Potter et al., 2019).

Hence, it is not surprising to find a high genetic diversity associated with strong IBD gradients in *A.*

erythrurus. However, low dispersal alone could not explain the phylogeographic pattern observed in

A. erythrurus across Morocco, and we found at least four groups separated by barriers to gene flow

(Figure 5): the northern Atlantic Coast, the Rif, the surroundings of Marrakech, and a large group

including the remaining the Atlantic coast, the Middle-Atlas and Sidi-Yahia. However, several of

the inferred barriers to gene flow were not unambiguously supported, highlighting a need for further

investigations. Surprisingly, this was the case for Rif and Middle-Atlas, two groups that show very

little to no traces of admixture at their contact zone, near Ifrane, where individuals belonging to

either group occur 1 km apart. An in-depth analysis of this contact zone based on genome-wide data

by Doniol-Valcroze et al. (2022) showed a very steep transition, demonstrating that the Middle-

Atlas and Rif are strongly differentiated groups with restricted admixture, consistent with the

existence of intrinsic barriers to gene flow (Dufresnes et al., 2020; see Doniol-Valcroze et al. (2022)

for a more in-depth discussion of the Rif/Middle-Atlas transition). This result supports the

hypothesis that ancestral polymorphism retention in our nine loci dataset, rather than widespread

gene flow, blurred the phylogeographic signal, to the point that two lineages at an advanced stage of

speciation were sometimes difficult to separate. Both the northern Atlantic and Marrakech plains

populations showed patterns of divergence akin that of the Rif (i.e., monophyly in the nDNA tree,

clear divergence in the MCAs), and might be candidates for deep intra-specific - or even species-

level - lineages. Future studies employing a larger number of molecular markers will hopefully help

clarifying how these groups are related, and whether they are linked by extensive gene flow or not.

Figure 5. *Summary of the processes underlying populations differentiation in Moroccan*
Acanthodactylus erythrurus. *The red star highlights the contact zone between the Rif and Middle-*
Atlas lineages, that is inconclusive in our analysis but is identified as a species-level barrier in
Doniol-Valcroze et al. (2022).

Although the application of our approach was limited by lineages represented by single localities,

our results support that populations from the Middle-Atlas, Atlantic coast to the exclusion of the

three northernmost localities, and Sidi-Yahia are all part of the same IBD gradient, without barriers

to gene flow detected. Alternatively, it is also possible that the Middle-Atlas and Atlantic coast

represent two formerly allopatric lineages that resumed high levels of gene flow upon secondary

contact, as supported by the intermediate position of some coastal populations (especially Blad-

Dendoun and Mehdia) in both MCAs and population trees. In the absence of explicit measures of

admixture, such scenario would be difficult to distinguish from strict IBD. Interestingly, these

populations are separated into two morphologically well differentiated subspecies, *A. e. atlanticus*

(Middle-Atlas) and *A. e. lineomaculatus* (Atlantic coast), the latter having even be proposed as a

distinct species (Bons & Geniez, 1995). Our results suggest that this morphological variation could

have arisen, or been maintained, in the face of gene flow, maybe as a result of local adaptation.

Contrary to what we observed in *A. erythrurus*, phylogeographic studies of reptile species

widespread in Western and Northern Morocco mostly report low levels of genetic diversity within

Morocco (e.g., El-Farhati et al., 2021; Faria & Harris, 2020; Nicolas et al., 2018; Rosado et al.,

2017; but see Klesser et al., 2021 and Veríssimo et al., 2016 for more complex patterns). Some of

the phylogeographic breaks we identified here correspond with large rivers (e.g., Oued Loukos

between N-Atl and the other costal Atlantic populations), others with ecological transitions (soft

coastal grounds versus hard inland substrates between S-Atl and Mara) but most do not align with obvious landscape or ecological breaks. It is possible that a substantial fraction of the lineages we identified correspond to different species (as for the Rif and Middle-Atlas lineages; Doniol-Valcroze et al., 2022), and that their divergences are maintained by intrinsic barriers to gene flow, similarly to the various species of the wall lizards (*Podarcis*) that currently inhabit the Iberian Peninsula (Caeiro-Dias et al., 2018, 2021a,b). Further sampling around contact zones and analyses of gene flow across these contact zones with genomic markers will be needed to partition the effects of intrinsic (e.g., mate recognition, selection against hybrids) and extrinsic (geography and ecology) barriers to gene flow in *A. erythrurus* in North Africa.

Whatever the current mechanisms underlying phylogeographic breaks in the species in Morocco, it is hard to pinpoint precisely which processes generated such a diversity over a relatively small spatial scale. It is possible that this pattern is the legacy of populations fragmentation caused by climatic variations during the Pleistocene, when periods of high aridity occurred (Le Houérou, 1997). *Acanthodactylus erythrurus* populations, being associated with Mediterranean and mesic habitats (Fonseca et al., 2009), might have repeatedly retracted to cooler and more humid places during arid periods, such as mountain ranges or coasts (similar to "refugia-within-refugia", Gómez & Lunt, 2007). Our results highlight that the recent biogeographic history of taxa within Morocco, and the Maghreb in general, is still poorly understood, and call for more comparative phylogeography studies in this area, especially relying on multi-locus data sets.

Acknowledgements

The authors are grateful to Raouaa Fathalla and Patricia Sourrouille for their help with laboratory work, and to Christian Hennig for his helpful input on linear models. Part of the analyses were carried out by L.R. using resources in project 2022/22-653 provided by the Swedish National Infrastructure for Computing (SNIC) at UPPMAX, partially funded by the Swedish Research Council through grant agreement no. 2018-05973. Part of this work was carried out by using the resources of the national INRAe MIGALE (Migale bioinformatics Facility, https://doi.org/10.15454/1.5572390655343293E12) and GENOTOUL (Bioinfo Genotoul, https://doi.org/10.15454/1.5572369328961167E12) bioinformatics HPC platforms, as well as the local Montpellier Bioinformatics Biodiversity (MBB, supported by the LabEx CeMEB ANR-10-LABX-04-01) and CBGP HPC platform services. This research was conducted in the scope of the LIA "Biodiversity and Evolution".

Funding

R.L. was supported by the Agence Nationale de la Recherche (GENOSPACE ANR-16-CE02-0008 and INTROSPEC ANR-19-CE02-0011), and by recurrent funding from INRAE.

Conflict of interest statement

737
738 The authors state that there are no conflicts of interest regarding this research.
739
740 **Authors contributions**
741

748 **References**
749

750 Afonso Silva, A. C., Bragg, J. G., Potter, S., Fernandes, C., Coelho, M. M., & Moritz, C. (2017).
751 Tropical specialist vs. climate generalist: diversification and demographic history of sister species
752 of *Carlia* skinks from northwestern Australia. *Molecular Ecology*, *26*(15), 4045–4058.
753
754 Audzijonyte, A., & Vrijenhoek, R. C. (2010). When gaps really are gaps: statistical phylogeography
755 of hydrothermal vent invertebrates. *Evolution 64*(8), 2369–2384.
756
757 Avise, J. C., Arnold, J., Ball, R. M., Bermingham, E., Lamb, T., Neigel, J. E., Reeb, A. C. & Saun-
758 ders, N. C. (1987). Intraspecific phylogeography: the mitochondrial DNA bridge between popula-
759 tion genetics and systematics. *Annual review of ecology and systematics*, *18*(1), 489–522.
760
761 Avise J. C., Ball R. M. Jr (1990). Principles of genealogical concordance in species concepts and
762 biological taxonomy. *Oxford surveys in evolutionary biology, 7,* 45–67
763
764 Beddek, M., Zenboudji-Beddek, S., Geniez, P., Fathalla, R., Sourouille, P., Arnal, V., Dellaoui, B.,
765 Koudache, F., Telailia, S., Peyre, O., & Crochet, P. A. (2018). Comparative phylogeography of am-
766 phibians and reptiles in Algeria suggests common causes for the east-west phylogeographic breaks
767 in the Maghreb. *PloS one*, *13*(8), e0201218.
768
769 Binks, R. M., Byrne, M., McMahon, K., Pitt, G., Murray, K., & Evans, R. D. (2019). Habitat dis-
770 continuities form strong barriers to gene flow among mangrove populations, despite the capacity for
771 long-distance dispersal. *Diversity and Distributions*, *25*(2), 298–309.
772
773 Bons, J., & Geniez, P. (1995). Contribution to the systematics of the lizard *Acanthodactylus*
774 *erythrurus* (Sauria, Lacertidae) in Morocco. *Herpetological Journal, 5,* 271–271.
775
776 Braby, M. F., Eastwood, R., & Murray, N. (2012). The subspecies concept in butterflies: has its ap-
777 plication in taxonomy and conservation biology outlived its usefulness? *Biological Journal of the*
778 *Linnean Society, 106*(4), 699–716.
779
780 Brito, P. H., & Edwards, S. V. (2009). Multilocus phylogeography and phylogenetics using se-
781 quence-based markers. *Genetica, 135*(3), 439–455.
782
783 Brito, J. C., Godinho, R., Martínez-Freiría, F., Pleguezuelos, J. M., Rebelo, H., Santos, X., ... &
784 Carranza, S. (2014). Unravelling biodiversity, evolution and threats to conservation in the Sahara-
785 Sahel. *Biological Reviews*, *89*(1), 215–231.
786

Caeiro-Dias, G., Luís, C., Pinho, C., Crochet, P. A., Sillero, N., & Kaliontzopoulou, A. (2018). Lack of congruence of genetic and niche divergence in *Podarcis hispanicus* complex. *Journal of Zoological Systematics and Evolutionary Research*, *56*(4), 479–492.

Caeiro-Dias, G., Rocha, S., Couto, A., Pereira, C., Brelsford, A., Crochet, P. A., & Pinho, C. (2021a). Nuclear phylogenies and genomics of a contact zone establish the species rank of *Podarcis lusitanicus* (Squamata, Lacertidae). *Molecular Phylogenetics and Evolution*, *164*, 107270.

Caeiro-Dias, G., Brelsford, A., Kaliontzopoulou, A., Meneses-Ribeiro, M., Crochet, P. A., & Pinho, C. (2021b). Variable levels of introgression between the endangered Podarcis carbonelli and highly divergent congeneric species. *Heredity*, *126*(3), 463–476.

Carnaval, A. C., Hickerson, M. J., Haddad, C. F., Rodrigues, M. T., & Moritz, C. (2009). Stability predicts genetic diversity in the Brazilian Atlantic foDurrantrest hotspot. *Science*, *323*(5915), 785–789.

Carpio, A. J., Melo-Ferreira, J., Martinez-Haro, M., Farelo, L., Campos, R., Alves, P. C., & Acevedo, P. (2021). Iberian hares with anciently introgressed mitochondrial DNA express a marginal environmental niche. *Journal of Biogeography*, *48*(9), 2328–2336.

Dávalos, L. M., & Russell, A. L. (2014). Sex-biased dispersal produces high error rates in mitochondrial distance-based and tree-based species delimitation. *Journal of Mammalogy*, *95*(4), 781–791.

Demos, T. C., Peterhans, J. C. K., Agwanda, B., & Hickerson, M. J. (2014). Uncovering cryptic diversity and refugial persistence among small mammal lineages across the Eastern Afromontane biodiversity hotspot. *Molecular Phylogenetics and Evolution*, *71*, 41–54.

Doniol-Valcroze, P., Rancilhac, L., Brito, J. C., Miralles, A., Geniez, P., Benoit, L., Loiseau, A., Leblois, R., Dufresnes, C., & Crochet, P.-A. (2022). Combining RADseq and contact zone analysis to decipher cryptic diversification in reptiles: Insights from *Acanthodactylus erythrurus* (Reptilia: Lacertidae). *BioRxiv*, 2022.09.30.510260. https://doi.org/10.1101/2022.09.30.510260

Drummond, A. J., Xie, W., & Heled, J. (2012). Bayesian inference of species trees from multilocus data using* BEAST. *Molecular Biology and Evolution*, *29*, 1969–1973.

Dufresnes, C., Nicieza, A. G., Litvinchuk, S. N., Rodrigues, N., Jeffries, D. L., Vences, M., Perrin, N., & Martínez-Solano, Í. (2020). Are glacial refugia hotspots of speciation and cytonuclear discordances? Answers from the genomic phylogeography of Spanish common frogs. *Molecular Ecology*, *29*(5), 986–1000.

Dufresnes, C., Pribille, M., Alard, B., Gonçalves, H., Amat, F., Crochet, P. A., Dubey, S., Perrin, N., Fumagalli, L., Vences, M., & Martínez-Solano, I. (2020). Integrating hybrid zone analyses in species delimitation: lessons from two anuran radiations of the Western Mediterranean. *Heredity*, *124*(3), 423–438.

Durrant, H. M., Barrett, N. S., Edgar, G. J., Coleman, M. A., & Burridge, C. P. (2015). Shallow phylogeographic histories of key species in a biodiversity hotspot. *Phycologia*, *54*(6), 556-565.

El-Farhati, H., Khaldi, M., Ribas, A., Hizem, M. W., Nouira, S., & Nicolas, V. (2021). Evolutionary history of the two North African hedgehogs (Mammalia: Erinaceidae) *Atelerix algirus* and

Paraechinus aethiopicus based on phylogeography and species distribution modelling. *Vertebrate Zoology*, *71*, 799.

Ferchaud, A. L., Eudeline, R., Arnal, V., Cheylan, M., Pottier, G., Leblois, R., & Crochet, P. A. (2015). Congruent signals of population history but radically different patterns of genetic diversity between mitochondrial and nuclear markers in a mountain lizard. *Molecular Ecology*, *24*(1), 192–207.

Faria, J. F., & Harris, D. J. (2020). Phylogeography of *Psammodromus algirus* (Linnaeus, 1758) in Morocco. *Herpetology Notes*, *13*, 1055–1061.

Fonseca, M. M., Brito, J. C., Paulo, O. S., Carretero, M. A., & Harris, D. J. (2009). Systematic and phylogeographical assessment of the *Acanthodactylus erythrurus* group (Reptilia: Lacertidae) based on phylogenetic analyses of mitochondrial and nuclear DNA. *Molecular Phylogenetics and Evolution*, *51*(2), 131–142.

Ficetola, G. F., Bonardi, A., Sindaco, R., & Padoa-Schioppa, E. (2013). Estimating patterns of reptile biodiversity in remote regions. *Journal of Biogeography*, *40*(6), 1202–1211.

Frantz, A. C., Cellina, S., Krier, A., Schley, L., & Burke, T. (2009). Using spatial Bayesian methods to determine the genetic structure of a continuously distributed population: clusters or isolation by distance? *Journal of Applied Ecology*, *46*(2), 493–505.

Funk, D. J., & Omland, K. E. (2003). Species-level paraphyly and polyphyly: frequency, causes, and consequences, with insights from animal mitochondrial DNA. *Annual Review of Ecology, Evolution, and Systematics*, 397–423.

Garcia-Porta, J., Irisarri, I., Kirchner, M., Rodríguez, A., Kirchhof, S., Brown, J. L., MacLeod, A., Turner, A. P., Ahmadzadeh, F., Albaladejo, G., Crnobrnja-Isailovic, J., De la Riva, I., Fawzi, A., Galan, P., Göçmen, B., Harris, D. J., Jiménez-Robles, O., Joger, U., Jovanović Glavaš, O., Karış, M., Koziel, G., Künzel, S., Lyra, M., Miles, D., Nogales, M., Anıl Oğuz, M., Pafilis, P., Rancilhac, L., Rodriguez Concepcion, B., Sanchez, E., Salvi, D., Slimani, T., S'khifa, A., Turk Qashqaei, A., Žagar, A., Lemmon, A., Lemmon, E. M., Carretero, M. A., Carranza, S., Philippe, H., Sinervo, B., Müller, J., Vences, M., & Valero, K. C. W. (2019). Environmental temperatures shape thermal physiology as well as diversification and genome-wide substitution rates in lizards. *Nature communications*, *10*(1), 1–12.

Gómez, A., & Lunt, D. H. (2007). Refugia within refugia: patterns of phylogeographic concordance in the Iberian Peninsula. In *Phylogeography of southern European refugia* (pp. 155–188). Springer, Dordrecht.

Goudarzi, F., Hemami, M. R., Rancilhac, L., Malekian, M., Fakheran, S., Elmer, K. R., & Steinfartz, S. (2019). Geographic separation and genetic differentiation of populations are not coupled with niche differentiation in threatened Kaiser's spotted newt (*Neurergus kaiseri*). *Scientific Reports*, *9*(1), 1–12.

Hausdorf, B., & Hennig, C. (2020). Species delimitation and geography. *Molecular Ecology Resources*, *20*(4), 950–960.

Huang, J. P. (2020). Is population subdivision different from speciation? From phylogeography to species delimitation. *Ecology and Evolution*, *10*(14), 6890–6896.

Husemann, M., Schmitt, T., Zachos, F. E., Ulrich, W., & Habel, J. C. (2014). Palaearctic biogeogra-
phy revisited: evidence for the existence of a North African refugium for Western Palaearctic biota.
Journal of Biogeography, *41*(1), 81–94.

Irwin, D. E. (2002). Phylogeographic breaks without geographic barriers to gene flow. *Evolution*,
56(12), 2383–2394.

Irwin, D. E. (2012). Local adaptation along smooth ecological gradients causes phylogeographic
breaks and phenotypic clustering. *The American Naturalist*, *180*(1), 35–49.

Joger, U., Fritz, U., Guicking, D., Kalyabina-Hauf, S., Nagy, Z. T., & Wink, M. (2007). Phylogeog-
raphy of western Palaearctic reptiles–Spatial and temporal speciation patterns. *Zoologischer Anzei-*
ger-A Journal of Comparative Zoology, *246*(4), 293–313.

Jombart, T. (2008). Adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics, 24(11), 1403–1405.

Klesser, R., Husemann, M., Schmitt, T., Sousa, P., Moussi, A., & Habel, J. C. (2021). Molecular
biogeography of the Mediterranean *Buthus* species complex (Scorpiones: Buthidae) at its southern
Palaearctic margin. *Biological Journal of the Linnean Society, 133*(1), 166–178.

Knowles, L. L. (2009). Statistical phylogeography. *Annual Review of Ecology, Evolution, and Sys-*
tematics, *40*, 593–612.

Kumar, R., & Kumar, V. (2018). A review of phylogeography: biotic and abiotic factors. *Geology,*
Ecology, and Landscapes, *2*(4), 268–274.

Kuo, C. H., & Avise, J. C. (2005). Phylogeographic breaks in low-dispersal species: the emergence
of concordance across gene trees. *Genetica*, *124*(2), 179–186.

Leaché, A. D., Oaks, J. R., Ofori-Boateng, C., & Fujita, M. K. (2020). Comparative phylogeogra-
phy of West African amphibians and reptiles. *Evolution*, *74*(4), 716–724.

Le Houérou, H. N. (1997). Climate, flora and fauna changes in the Sahara over the past 500 million
years. *Journal of Arid Environments*, *37*(4), 619–647.

Librado, P., & Rozas, J. (2009). DnaSP v5: a software for comprehensive analysis of DNA poly-
morphism data. *Bioinformatics*, *25*(11), 1451–1452.

Miralles, A., Geniez, P., Beddek, M., Aranda, D. M., Brito, J. C., Leblois, R., & Crochet, P. A.
(2020). Morphology and multilocus phylogeny of the Spiny-footed Lizard (*Acanthodactylus*
erythrurus) complex reveal two new mountain species from the Moroccan Atlas. *Zootaxa*, *4747*(2).

Muir, G., & Schloetterer, C. (2005). Evidence for shared ancestral polymorphism rather than recur-
rent gene flow at microsatellite loci differentiating two hybridizing oaks (*Quercus* spp.). *Molecular*
ecology, 14(2), 549–561.

Nicolas, V., Mataame, A., Crochet, P. A., Geniez, P., Fahd, S., & Ohler, A. (2018). Phylogeography
and ecological niche modeling unravel the evolutionary history of the African green toad, *Bufotes*
boulengeri boulengeri (Amphibia: Bufonidae), through the Quaternary. *Journal of Zoological Sys-*
tematics and Evolutionary Research, *56*(1), 102–116.

Palumbi, S. R., & Baker, C. S. (1994). Contrasting population structure from nuclear intron se-
quences and mtDNA of humpback whales. *Molecular Biology and Evolution*, *11*(3), 426–435.
945

946 Pamilo, P., & Nei, M. (1988). Relationships between gene trees and species trees. *Molecular biol-
947 ogy and evolution*, *5*(5), 568-583.
948

949 Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolution-
950 ary analyses in R. *Bioinformatics*, *35*(3), 526–528.
951

952 Perez, M. F., Franco, F. F., Bombonato, J. R., Bonatelli, I. A., Khan, G., Romeiro-Brito, M., Fegies,
953 A. C., Ribeiro, P. M., Silva, G. A. R., & Moraes, E. M. (2018). Assessing population structure in
954 the face of isolation by distance: are we neglecting the problem? *Diversity and Distributions*,
955 *24*(12), 1883–1889.
956

958 Peters, J. L., Bolender, K. A., & Pearce, J. M. (2012). Behavioural vs. molecular sources of conflict
959 between nuclear and mitochondrial DNA: the role of male-biased dispersal in a Holarctic sea duck.
960 *Molecular Ecology, 21*(14), 3562–3575.
961

962 Petkova, D., Novembre, J., & Stephens, M. (2016). Visualizing spatial population structure with es-
963 timated effective migration surfaces. *Nature genetics*, *48*(1), 94–100.
964

965 Potter, S., Afonso Silva, A. C., Bragg, J. G., Catalano, S. R., Donnellan, S., Doughty, P., Scott, M.
966 L., & Moritz, C. (2019). Contrasting scales of local persistence between monsoonal and arid biomes
967 in closely related, low-dispersal vertebrates. *Journal of Biogeography, 46*(11), 2506–2519.
968

969 Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multi-
970 locus genotype data. *Genetics*, *155*(2), 945–959.
971

972 Ramachandran, S., Deshpande, O., Roseman, C. C., Rosenberg, N. A., Feldman, M. W., & Cavalli-
973 Sforza, L. L. (2005). Support from the relationship of genetic and geographic distance in human
974 populations for a serial founder effect originating in Africa. *Proceedings of the National Academy
975 of Sciences*, *102*(44), 15942–15947.
976

977 Renoult, J. P., Geniez, P., Bacquet, P., Benoit, L., & Crochet, P. A. (2009). Morphology and nuclear
978 markers reveal extensive mitochondrial introgressions in the Iberian Wall Lizard species complex.
979 *Molecular ecology*, *18*(20), 4298–4315.
980

981 Rosado, D., Rato, C., Salvi, D., & Harris, D. J. (2017). Evolutionary history of the Morocco lizard-
982 fingered geckos of the *Saurodactylus brosseti* complex. *Evolutionary Biology*, *44*(3), 386–400.
983

984 Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics under isola-
985 tion by distance. *Genetics, 145*(4), 1219–1228.
986

987 Salvador, A. (1982) A revision of the lizards of the genus *Acanthodactylus* (Sauria: Lacertidae).
988 Bonner Zoologisches Monographien, Nr. 16, *Zoologisches Forschungsinstitut und Museum Alexan-
989 der Koenig*, 1–167.
990

Salvi, D., Pinho, C., & Harris, D. J. (2017). Digging up the roots of an insular hotspot of genetic di-
versity: decoupled mito-nuclear histories in the evolution of the Corsican-Sardinian endemic lizard
Podarcis tiliguerta. *BMC evolutionary biology*, *17*(1), 1–22.

Salvi, D., Pinho, C., Mendes, J., & Harris, D. J. (2021). Fossil-calibrated time tree of *Podarcis* wall
lizards provides limited support for biogeographic calibration models. *Molecular Phylogenetics and*
Evolution, 161, 107169.

Serre, D., & Pääbo, S. (2004). Evidence for gradients of human genetic diversity within and among
continents. *Genome research*, *14*(9), 1679–1685.

1001

1002 Shaffer, H. B., Gidiş, M., McCartney-Melstad, E., Neal, K. M., Oyamaguchi, H. M., Tellez, M., &
1003 Toffelmier, E. M. (2015). Conservation genetics and genomics of amphibians and reptiles. *Annual*
1004 *Review of Animal Biosciences*, *3*(1), 113–138.

1005

1006 Schultze, N., Spitzweg, C., Corti, C., Delaugerre, M., Di Nicola, M. R., Geniez, P., Lapini, L.,
1007 Liuzzi, C., Lunghi, E., Novarini, N., Picariello, O., Razzetti, E., Sperone, E., Stellati, L., Vignoli,
1008 L., Asztalos, M., Kindler, C., Vamberger, M., & Fritz, U. (2020). Mitochondrial ghost lineages blur
1009 phylogeography and taxonomy of *Natrix helvetica* and *N. natrix* in Italy and Corsica. *Zoologica*
1010 *Scripta, 49*(4), 395–411.

1011

1012 Schwartz, M. K., & McKelvey, K. S. (2009). Why sampling scheme matters: the effect of sampling
1013 scheme on landscape genetic results. *Conservation Genetics*, *10*(2), 441.

1014

1015 Tamar, K., Carranza, S., Sindaco, R., Moravec, J., Trape, J. F., & Meiri, S. (2016). Out of Africa:
1016 phylogeny and biogeography of the widespread genus *Acanthodactylus* (Reptilia: Lacertidae). *Mo-*
1017 *lecular Phylogenetics and evolution*, *103*, 6–18.

1018

1019 Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: molecular evolutionary genetics analysis
1020 version 11. *Molecular biology and evolution*, *38*(7), 3022–3027.

1021

1022 Teske, P. R., Golla, T. R., Sandoval-Castillo, J., Emami-Khoyi, A., Van der Lingen, C. D., Von der
1023 Heyden, S., ... & Beheregaray, L. B. (2018). Mitochondrial DNA is unsuitable to test for isolation
1024 by distance. *Scientific reports*, *8*(1), 1–9.

1025

1026 Toews, D. P., & Brelsford, A. (2012). The biogeography of mitochondrial and nuclear discordance
1027 in animals. *Molecular ecology, 21*(16), 3907–3930.

1028

1029 Tolley, K. A., Telford, N. S., Taft, J. M., Bates, M. F., Conradie, W., Makhubo, B. G., & Alexan-
1030 der, G. J. (2022). Taxonomic inflation due to inadequate sampling: are girdled lizards (*Cordylus mi-*
1031 *nor* species complex) from the Great Karoo one and the same? *Biological Journal of the Linnean*
1032 *Society*, *135*(1), 1–24.

1033

1034 Velo-Antón, G., Martínez-Freiría, F., Pereira, P., Crochet, P. A., & Brito, J. C. (2018). Living on
1035 the edge: Ecological and genetic connectivity of the spiny-footed lizard, *Acanthodactylus aureus*,
1036 confirms the Atlantic Sahara desert as a biogeographic corridor and centre of lineage diversifica-
1037 tion. *Journal of Biogeography*, *45*(5), 1031–1042.

1038

1039 Veríssimo, J., Znari, M., Stuckas, H., Fritz, U., Pereira, P., Teixeira, J., Arculeo, M., Marrone, F.,
1040 Sacco, F., Naimi, M., Kehlmaier, C., & Velo-Antón, G. (2016). Pleistocene diversification in Mo-
1041 rocco and recent demographic expansion in the Mediterranean pond turtle *Mauremys leprosa*. *Bio-*
1042 *logical Journal of the Linnean Society*, *119*(4), 943–959.

1043

1044   Vignoli, L., Vuerich, V., & Bologna, M. A. (2012). Experimental study of dispersal behaviour in a
1045   wall lizard species (*Podarcis sicula*) (Sauria Lacertidae). *Ethology Ecology & Evolution, 24*(3),
1046   244–256.
1047
1048   Wiemers, M., & Fiedler, K. (2007). Does the DNA barcoding gap exist? A case study in blue but-
1049   terflies (Lepidoptera: Lycaenidae). *Frontiers in zoology, 4*(1), 1–16.
1050
1051   Zieliński, P., Nadachowska-Brzyska, K., Wielstra, B., Szkotak, R., Covaciu-Marcov, S. D.,
1052   Cogălniceanu, D., & Babik, W. (2013). No evidence for nuclear introgression despite complete mt
1053   DNA replacement in the Carpathian newt (*Lissotriton montandoni*). *Molecular Ecology, 22*(7),
1054   1884–1903.