Neural representations of the content and production of human vocalization

Vera A. Voigtlaender[1,2,3,4], Florian Sandhaeger[1,2,3,4], David Hawellek[1,2,3,6], Steffen R. Hage[2,5] & Markus Siegel[1,2,3]

[1] Department of Neural Dynamics and Magnetoencephalography, Hertie Institute for Clinical Brain Research, University of Tübingen, Germany
[2] Centre for Integrative Neuroscience, University of Tübingen, Germany
[3] MEG Center, University of Tübingen, Germany
[4] Graduate Training Centre of Neuroscience, International Max Planck Research School, University of Tübingen, Germany
[5] Dept. of Otolaryngology-Head and Neck Surgery, Hearing Research Centre, University of Tübingen, Germany
[6] Roche, Pharmaceutical Research and Early Development, Roche Innovation Center Basel, Basel, Switzerland

Correspondence: vera.voigtlaender@uni-tuebingen.de & markus.siegel@uni-tuebingen.de

Abstract

Speech, as the spoken form of language, is fundamental for human communication. The phenomenon of covert inner speech implies a functional independence of speech content and motor production. However, it remains unclear how a flexible mapping between speech content and production is achieved on the neural level. To address this, we recorded magnetoencephalography (MEG) in humans performing a rule-based vocalization task. On each trial, vocalization content (one of two vowels) and production form (overt or covert) were instructed independently. Using multivariate pattern analysis, we found robust neural information about vocalization content and production, mostly originating from speech areas of the left hemisphere. Production signals dynamically transformed upon presentation of the content cue, whereas content signals remained largely stable throughout the trial. In sum, our results show dissociable neural representations of vocalization content and production in the human brain and provide new insights into the neural dynamics underlying human vocalization.

Vocal behavior is an essential component of human communication. Particularly speech, the

spoken form of language, is a highly sophisticated skill exclusive to humans. Thereby, we can

encode information not only in sound (overt speech), but also in thought (covert speech).

These different speech forms imply a functional independence of speech content and motor

production. However, it remains an open question how content and production are represented

neuronally and how the brain achieves a flexible mapping between the two.

In speech, two levels need to be distinguished. The lexical level, which refers to entire words,

and the sub-lexical level, which refers to parts of words, such as phonemes or syllables [1]. On

the lexical level, neural activity underlying overt and covert speech is similar, with differences

largely accounted for by a higher degree of executive motor control during overt speech [2–5].

Broca's area was found to act as a supra-modal hub, exhibiting language-specific activation

independent of the production form [6]. These findings suggest that there is a neural

representation of speech content which is to some degree independent of its motor production.

While this independence is intuitive on the lexical level, it is less clear on the sub-lexical level,

where content may be expected to be tightly bound to the motor production. Still, initial

evidence suggests such production independent representations of sub-lexical entities like

syllables and phonemes [7,8].

In sum, past work suggests that neural activity underlying lexical and sub-lexical vocalizations

represents both speech content and motor production. However, it remains unclear whether

content representations generalize across different motor plans, and to what extent it is

possible to dissociate these aspects. Furthermore, the dynamic interplay between emerging

representations of content and motor production is not known, as most previous studies used

neural data with inherently poor temporal resolution.

Here, we therefore aimed, first, to independently manipulate and decode neuronal content and

motor components of human vocalization, and second, to investigate their dynamic interplay

across time. We recorded MEG while subjects performed a novel rule-based vocalization task

dissociating the content and motor aspects of sub-lexical speech. Content (one of two vowels)

and production (overt or covert) were instructed sequentially and in random order. Multivariate

pattern analysis (MVPA) of time-resolved MEG data allowed us to characterize the format,

overlap and temporal dynamics of neural content and production representations.

With this joint approach, we were able to read out content and motor information several

seconds before speech onset. The strength of neural information correlated with the degree of

motor involvement and the two representations overlapped when isolated. The production

representation transformed once the content was known, whereas the content representation

remained stable until the onset of vocalization.

Results

The components of vocalization can be decoded independently

We recorded MEG while subjects performed a rule-based vocalization task. Participants had

to overtly vocalize or covertly imagine the vocalization of two different vowels. During each

trial, content (/u/ / /ə/) and production (vocalized/ imagined) were instructed sequentially with

visual cues (Fig. 1A). Each cue lasted 100 ms and was followed by a 2 s delay. At the end of

the trial, a brief dimming of the fixation point served as a go-cue for the onset of vocalization

or imagination. The order of instruction was randomized, as was the assignment of the

instructed content or production to the visual cues (Fig. 1B). Participants performed the correct

production type (vocalized vs. imagined) in 97.98 % of the trials. In case of vocalized trials, the

correct vowel (/u/ vs /ə/), was performed in 100 % of the trials. We checked vocalized trials for

their onset latency (possible for 37 sessions). In 98.48 % of the trials, the vocal onset was after

the go cue, as instructed. The mean vocal latency of these trials was 0.58 s (+/- 0.12 SD).

Fig. 1 | Rule-based vocalization task. Participants imagined or vocalized different contents (phonemes /u/ and /ə/). **a**, Production and the content were instructed successively with visual cues, according to the respective rule of the trial block. **b**, Rules for four trial blocks per recording session. In two blocks the content was instructed first, in the other two blocks the production was instructed first.

For each subject and recording session, we computed neural information about content and production. We applied cross-validated MANOVA on preprocessed single-trial MEG data from all sensors (cvMANOVA; see Methods) [9,10]. As an extension of the commonly used cross-validated Mahalanobis distance, cvMANOVA allows for the simultaneous quantification of the variability in neural data due to several variables of interest. The resulting measure of neural information can be interpreted analogously to classifier performance from multivariate decoding analyses. To enable robust statistical tests, we averaged information in the time windows between cue 1 and 2 (delay 1), as well as between cue 2 and the go cue (delay 2). In both cases, we excluded the first 250 ms after cue offset to avoid a possible confound due to sensory activity evoked by the cues (see Methods).

We observed significant neural information about both variables (Fig. 2). For both orders of cue presentation, we found information about the variables shortly after their respective instruction. When content was instructed first, there was significant content information in delay 1 (p = 0.003; corrected) and significant information about content and production in delay 2 (content: $p = 3.4 \times 10^{-4}$, production: $p = 1.6 \times 10^{-9}$; corrected). Conversely, when production was instructed first, content information was only present in delay 2, whereas production

information was highly significant in both delays (content: p_{del2} = 7.8x10^{-5}; production: p_{del1} =

2.8x10^{-6}, p_{del2} = 8.5x10^{-9}; corrected). Thus, both the content of a vocalization and its production

form were represented neuronally, several seconds before the actual execution.

Fig. 2 | Neural information about content and production. a, Information in trials with content instructed first.
b, Information in trials with production instructed first. Shaded regions and error bars indicate SEM. Bar plots show
average information in delay 1 and 2. Asterisks indicate significance (n = 24, p < 0.05 corrected; t-test, one-tailed).

The components of vocalization are modulated by effort

Both experimental dimensions entailed differences in motor effort. Imagined vowels lacked

actual vocalization, just as /ə/, as a non-articulated vowel, lacked the strong articulation of /u/.

Therefore, we wanted to investigate whether the neural information about content was equally

strong for both production types and whether the neural information about production type was

identical for both vowels. To test this, we repeated the analysis after splitting the data based

on both variables. In other words, content was decoded separately for both production types,

and production was decoded separately for both vowels (Fig. 3).

Fig. 3 | Neural information about content and production in split conditions. Shaded regions and error bars indicate SEM. Bar plots show averaged information of delay 1 and 2. Asterisks indicate above bars significant information (n = 24, p < 0.05 corrected; t-test, one-tailed). Horizontal lines with asterisks on top indicate a significant difference (n = 24, p < 0.05 corrected; paired t-test, two-tailed).

In all conditions, information about both variables could be decoded after their respective instruction. When content was instructed first, production information was significant for both vowels, in delay 2 (/ə/: $p = 7.8\times10^{-9}$, /u/: $p = 6\times10^{-9}$; corrected). When production was instructed first, it could be read out in both delays and for both vowels (/ə/: $p_{del1} = 4.1\times10^{-6}$, $p_{del2} = 4.8\times10^{-7}$; /u/: $p_{del1} = 1.4\times10^{-4}$, $p_{del2} = 1.6\times10^{-9}$; corrected). In the second delay, production information was higher in /u/ than in /ə/ trials, but only significantly so when content was instructed first ($p = 0.026$, paired t-test; corrected).

Content information was significant in both delays and production types when content was instructed first (imagined: $p_{del1} = 0.044$, $p_{del2} = 0.044$; vocalized: $p_{del1} = 0.004$, $p_{del2} = 2.7\times10^{-5}$; corrected). When production was instructed first, content information was only significant in the second delay, again in both production types (imagined: $p = 0.007$, vocalized: $p = 1.5\times10^{-4}$; corrected). Content information was higher during vocalized than imagined trials, in both orders and all relevant delays. This difference was significant in the second delay when content was instructed first ($p = 0.007$, paired t-test; corrected).

133 In sum, both production and content information were present in all individual conditions and,

134 furthermore, higher in those conditions with stronger motor involvement.

Fig. 4 | Spatial dynamics of neural information about content production. a, Production information in both orders. **b**, Content information in both orders. Left and top view of the cortical distribution of information. Information was averaged in 500 ms intervals, except for the intervals directly after the cues, for which the first 250 ms were excluded.

The components of vocalization are represented in cortical speech areas

We found that vocalization content and production were represented in cortical areas typically

associated with speech. To characterize the cortical distribution of content- and production

information, we repeated the cvMANOVA-analysis on the source level using a searchlight

approach. We then averaged neural information within four 500ms time windows per delay.

Again, for the 1st time window after each cue, the initial 250ms were excluded. This analysis

revealed spatially stable representations of both variables (Fig. 4). Descriptively, the highest

level of production information was found in Broca's area and other motor cortices. Content

was represented in the same areas, but additionally in temporal cortices.

Fig. 5 | Lateralization index of content and production information. The lateralization index (left – right hemisphere information) was computed for both delays and both orders. Asterisks indicate significance (n = 24, p < 0.05 corrected; t-test, one-tailed).

For higher-order language processes neural activity is known to be left-lateralized, which is debated for speech on the sub-lexical level [11,12]. Our source patterns suggested a clear left-lateralization. We tested this by computing a lateralization index for both variables (information in the left hemisphere - right hemisphere, Fig. 5). Both variables were left-lateralized in all delays after the respective instruction. This was significant for content information in the first delay when it was instructed first (p = 0.037; corrected) and in the second delay when production was instructed first (p = 0.006; corrected). Production information was significantly left-lateralized in both delays when production was instructed first (p_{del1} = 0.048, p_{del2} = 0.009; corrected) and in the second delay when it was instructed second (p = 0.003; corrected).

Taken together, source analysis showed that both content and production had stable left-lateralized representations on the cortical level. This did not only identify the origin of neural information within well-known speech-associated areas, but also excluded confounds due to inherently non-lateralized effects of visual cues or electromagnetic artifacts.

The components of vocalization have different representational formats

The searchlight analysis indicated a spatial stability of the coarse cortical distribution of content and production information across time. Nonetheless, the representational format, i.e. the fine-grained cortical pattern underlying each representation may be dynamic. To test if neural representations of content and production transformed across time, we decoded both variables on the sensor level across time [13] (Fig. 6). By using each pair of time-points for training and testing the cvMANOVA, we could assess to what extent neural representations were

173 overlapping across timepoints. Stable or dynamic representations would yield high or low

174 cross-time decoding, respectively. We performed this analysis both on the content-first

175 condition and on the production-first condition, as well as training on all timepoints of one

176 condition and testing on those of the other (Fig. 6, mixed orders).

Fig. 6 | Cross-time information about content and production. a, Cross-time production information. **b**, Cross-
time content information. Bar plots show the averaged cross-temporal information for delay 1, 2 and 1/2, as well as
the expected values for stable representations. The diagonals ± 100 ms were excluded from calculations. Asterisks
indicate significant information (n = 24, p < 0.05 corrected; t-test, one-tailed) and horizontal bars with asterisks on
top indicate significantly smaller cross-information than the expected value (n = 24, p < 0.05 corrected; paired t-
test, one-tailed).

For statistical testing, we averaged neural cross-information per variable and delay (Fig. 6, bar

plots). Cross-temporal Information within delay 1 was averaged over all conditions in which the

relevant cue was instructed first, as information could only be present in these conditions. To

estimate cross-temporal generalization between delays, we averaged an off-diagonal time-

window over those conditions where the relevant cue was presented first, as well as those

mixed-order conditions where the relevant cue was presented first in one condition but second

in the other. To estimate cross-temporal information within delay 2, we averaged data from all

cue orders. Briefly, this ensured that neural information about content or production was

available at all training and test timepoints included in the statistical analysis, allowing us to

meaningfully address the question whether representations were shared between training and

test timepoints.

Within both delay 1 and 2, there was significant cross-temporal information about both

variables (production: p_{del1} = 6.3x10^{-5}, p_{del2} = 2.9x10^{-9}; content: p_{del1} = 0.027, p_{del2} = 1.1x10^{-5};

corrected). There was also significant cross-temporal information between delays 1 and 2 for

both variables (production: p = 6.3x10^{-5}; content: p = 0.012; corrected).

To quantitatively interpret these results in relation to the information available at each individual

timepoint, we computed an estimate of the expected cross-information in case of perfectly

stable representational formats but potentially different information magnitudes over time [10].

This expected cross-information allowed us to test whether representations significantly

differed between timepoints. The cross-temporal stability of production representations was

lower than expected in all time windows, and significantly so between delay 1 and 2 and within

delay 2 ($p_{del1/2}$ = 2x10^{-4}, p_{del2} = 7.3x10^{-6}; corrected). In contrast, cross-temporal content

information was never significantly smaller than expected for a temporally stable

representation. Thus, while we found evidence for a partially dynamic representation of

production type, this was not the case for content, which appeared stable over time.

To what extent are the neural representations of content and production similar? Our cross-

time decoding analysis showed that both representations evolve differently across time. We

took this as an indication that these representations are not identical. To rigorously estimate

the extent of representational overlap, we implemented a cross-variable analysis, training the

algorithm on one variable and testing it on the other. Again, we computed the expected cross-

information under the assumption of identical representations and compared it with the

observed cross-information. We computed cross-variable information in both cue orders, but

also in mixed cue orders where the relevant cue was either first or second (Fig. 7).

Fig. 7 | Cross-variable information of content and production. Cross-information in both orders and mixed orders with the relevant cue first or second. **a**, Cross-information in trials with content instructed first. **b**, Cross-information across trials with relevant cue first. **c**, Cross-information across trials with relevant cue second. **d**, Cross-information in trials with production instructed first. Solid lines indicate cross-information and dashed lines indicate the respective expected values for identical representations. Bar plots show the averaged cross-information and the averaged expected values. Asterisks indicate significant cross-information (n = 24, p < 0.05 corrected; t-test, one-tailed) and horizontal bars with asterisks on top indicate significantly smaller cross-information than the expected value (n = 24, p < 0.05 corrected; paired t-test, one-tailed).

We found significant cross-information only in delay 1 in the order with the relevant cue first (p = 0.046; corrected). In the other orders, as information about both variables could only be present after the second cue, no cross-information was expected in delay 1. While we observed a small amount of cross-information in delay 2 of the mixed orders, this was not significant. On the other hand, cross-information was significantly smaller than its expected value in the second delay of all orders (content first: p = 0.004; both first: p = 0.02; both second: p = 0.01; production first: p = 0.002; corrected). Taken together, these results show that content and production representations overlap but are not identical. In our data, isolated content representations (before knowledge of the production type) cannot be distinguished from isolated production representations (before knowledge of the content). Thus, content and production representations are indistinguishable as long as the respective other variable is unknown to the participant. However, as soon as both aspects become available, their representations show strong differences.

Information about the components of vocalization remains stable across sessions

Electromagnetic artifacts and other, usually visually driven confounders, often pollute the data in speech studies [14]. Our source level analysis provided evidence that content- and production information was indeed speech related, originating from well-known speech-associated areas. In addition, we implemented a control analysis to test if a possible visual cue confound had an impact on our results. Because the order of trial blocks was reversed in each participant's second recording session, it was possible to cancel out the visual cue effect by decoding across sessions. To this end, we trained the cvMANOVA on one session and tested on the other.

Fig. 8 | Cross-session decoding of content and production. Information was averaged for both orders and delays. Asterisks indicate significant information (n = 24, p < 0.05 corrected; t-test, one-tailed).

Decoding across sessions was expected to be more challenging than decoding within session, as the signal-to-noise ratio was impacted by additional variability due to head movement between the sessions. Nevertheless, we found significant information about both content and production in all relevant delays. There was significant content information in both delays if it was instructed first ($p_{del1} = 0.004$, $p_{del2} = 5.1 \times 10^{-5}$; corrected) and in delay 2 if it was instructed second ($p = 0.006$; corrected). Production information was significant in delay 2 if content was instructed first ($p = 3.4 \times 10^{-8}$; corrected) and in both delays if production was instructed first ($p_{del1} = 4.8 \times 10^{-5}$, $p_{del2} = 3.4 \times 10^{-6}$; corrected). Thus, content- and production information were stable across recording sessions, and therefore also not driven by a visual, cue-related confound due to the sequential order of rule blocks within each session.

Discussion

Our results shed new light on the neural mechanisms underlying the flexible mapping between two components of human speech: its content and motor production. Combining MEG and MVPA, we could decode robust neural information about these components directly after their respective instruction, which was up to several seconds preceding the onset of the vocalization. Thus, our factorial task design allowed us to dissociate content and production in the pre-execution phase, where key processing stages take place [15–18] and electromagnetic artifacts of the motor production itself are ruled out [14].

We found significant information about each component directly after the first cue, when the respective other instruction was still missing. The early content information suggests that content can be represented independently of a specific motor plan, which falls in line with implications from previous studies [2,6–8,12,15,19]. Therefore, the actual motor production is not necessary to form a neural representation of content. However, this does not imply that the content representation is completely independent from motor planning.

Content information was present for both production types, as was production information for both vowels. However, content information was higher when vocalized and production information was higher for the vowel /u/, implying a dependence of information strength on the degree of motor involvement. As previous studies have shown, overt and covert speech differ in terms of executive motor control, including M1 recruitment for the efference copy [2,5], which could account for the higher content information in vocalized trials. In addition, the stronger phonological code retrieval and encoding in overt speech could also contribute to the observed difference of content information between the production types [5]. The two vowels also differ in motor involvement, as /ə/ is a non-articulated innate-like vowel, whereas /u/ is strongly articulated and learned [20]. Therefore, the stronger motor involvement could account for higher production information in the vowel /u/.

The vocal motor network comprises many cortical and subcortical areas (Hage and Nieder,

2016; Jürgens, 2002; Schulz et al., 2005). As expected, we located representations of content

and production in frontal and central cortex consistent with well-known speech-associated

areas. In addition, content representation extended to temporal cortices, which may be due to

efference copies in sensory regions [23–25]. Although one may not expect the higher-order

language network to be recruited in our paradigm, we found neural representations of both

content and production to be stronger in the left hemisphere. This could either indicate that

language capacities beyond low-level speech were recruited or that low-level speech

processes can already be lateralized under specific circumstances. Independent of these

alternatives, our finding of lateralization in an early pre-execution phase suggests that the

uncovered neural representations were indeed speech or language specific and did not reflect

general working memory processes.

Our cross-decoding approach provided new insights into the temporal dynamics and

similarities between the neural representations of vocalization components. During the first

delay, when information about only one of the components was available to the subjects, both

representations were temporally stable and indistinguishable from each other. Thus, while

there may be subtle differences that MEG is insensitive to, our results suggest an overlap of

neural representations of content and production. One possible explanation for such an

overlap could be an effort effect where conditions with a higher degree of motor involvement

elicit higher neural activity than those with a lesser degree. Concretely, |u vs. ə| and |vocalized

vs. imagined| could both correspond to contrasts of |high effort vs. low effort|. This effect could

reflect priming motor signals preceding vocalizations [18,26]. Alternatively, it could also reflect the

firing patterns of one or more speech specific neural populations that encode several content-

and production-related features. Here, effort could drive either the firing rates of individual

neurons or the number of recruited cells. Further invasive research is required to determine

whether the same population, or spatially close and therefore indistinguishable neurons are

modulated by both content- and production-related effort.

In the second delay, cross-information between content and production was similar as in the first delay. However, as both content- and production information were much higher, the representations were now clearly distinguishable. Moreover, cross-temporal decoding between both delays revealed that the content representation remained stable over time, whereas the production representation transformed once the content was known. Consequently, the divergence of the representations in the second delay was likely driven by the transformation of production. Taken together, this implies that the production representation during the second delay was a combination of the initial format during the first delay and an additional component. This additional component, building up once the content was known, may reflect the specific motor program used to prepare the articulation of the respective vowel.

In sum, our results show that, when isolated, both representations overlap and correlate with the degree of motor involvement. This overlap of the representations could be caused by pre-motor-like activity that is modulated by effort, but independent of the actual execution. While the content representation remains stable, the production representation changes once the content is known, likely reflecting the addition of a specific motor program.

While natural speech can functionally be decomposed into content and motor production, there is little evidence for a content dimension on the neural level and even less about its dynamic interplay with the motor production. Most implications for a neural content dimension come from studies focusing on the lexical [3,6,24] or the sub lexical syllable level [2,7,23]. Temporal dynamics have so far only been studied on the lexical level [15,16]. Yet, the elementary building blocks of speech are phonemes, and to our knowledge all previous research on this level is related to motor production [12,19,27,28]. Our results uncover a neural content dimension for phonemes that was present independently of motor production and could therefore allow for a generalization between production forms. These results accord well with the larger body of work on the higher sub lexical and lexical levels.

339 Our findings set the stage for future research, to investigate how neural codes of isolated

340 phonemes and their motor production translate to those embedded in speech. Our combined

341 approach of MEG, MVPA and a rule-based paradigm, provides a fruitful framework for that,

342 thus, opening a new window for non-invasive speech research in health and disease.

Methods

Subjects

345 24 healthy humans with normal or corrected-to-normal vision participated in the study (14

346 male; 21 right-handed; mean age: 29 years; 5 years SD). All participants gave written informed

347 consent before participation and received monetary reward afterwards. The study was

348 conducted in accordance with the declaration of Helsinki and approved by the ethics committee

349 of the University of Tübingen.

Behavioral Task and Stimuli

Participants performed a rule-based vocalization task. In each trial, one of two vowels (/u/ or

/ə/ had to be either overtly or covertly vocalized. Vowel and production type were instructed

sequentially by visual cues. The corresponding rule, i.e. the assignment of the visual cues to

their instructed content, changed across recording blocks and was indicated before the

beginning of each block.

Participants self-paced the trials using closed-loop eye movement control. Each trial started

with an initiation phase of 1000 ms, during which a white fixation spot (diameter: 0.1° of the

visual angle) appeared at the center of the screen. Once fixation was acquired, the first visual

cue (a forward or backward white slash, length: 2°, width: 0.25° of the visual angle) appeared

for 100 ms, instructing either content or production. Then, the fixation spot appeared again for

a delay period of 2000 ms. The second visual cue appeared for 100 ms, instructing the

respective missing variable, followed by a second delay period displaying the fixation spot.

Dimming the spot for 100 ms served as the go-cue for the participants' response. After the go-

cue, the fixation spot remained on screen for 1500 ms, which gave time for the response of the participants. The intertrial interval was 1000 ms long, indicated by dimming of the fixation spot. If fixation was broken after the onset of the first visual cue, the trial was aborted, which was indicated by a color change of the fixation spot to red for 500 ms. The cue configuration of the aborted trial was repeated at a random position later within this block.

Within each recording block, the order of instruction and the meaning of each instruction cue was fixed. However, in half of the blocks, the content was instructed first, whereas in the other half of the blocks, the production was instructed first. Moreover, the assignment of each visual cue to its meaning (content or production) was different in half of the blocks for each order. These four different rules led to four blocks of trials. Each block contained 80 trials, with 20 per condition (1: /u/ vocalized, 2: /u/ imagined, 3: /ə/ vocalized, 4: /ə/ imagined). The order of the conditions was randomized per block. In total, there were 16 different conditions, including the different assignments of the visual cues to their instructed content. The 24 possible orders of the four blocks were randomly assigned to the participants.

Before the experiment, participants were instructed on how to articulate the vowels correctly. Thereby, we made sure that /u/ was clearly articulated, whereas /ə/ was produced with minimal involvement of the vocal tract. We also made sure that there was no involuntary articulatory movement visible for imagined vowels. During the experiment, the participants memorized the respective rule before each block. The rule was presented on the screen for as long as necessary. Eight training trials preceded each block to ensure correct performance. If necessary, the rule was shown again during the block while the sequence of trials was paused. All participants performed two MEG sessions with 320 trials each. For each participant, the order of blocks from the first session was inversed for the second session. During both sessions, the performance of the participants was monitored with a microphone and a camera. After the experiments, each trial was labeled for production type and in case of vocalized trials for vowels.

Data Acquisition

We recorded MEG (Omega 2000, CTF Systems Inc., Port Coquitlam, Canada) with 273 sensors at a sampling rate of 2342.75 Hz in a magnetically shielded chamber. Participants sat upright with a screen at 65 cm viewing distance. Stimuli were projected onto the screen by an LCD projector (Sanyo PLC-XP41, Moriguchi, Japan) with a refresh rate of 60 Hz. The projection was the only source of light in the chamber. Continuous head movement was monitored with three coils attached to fiducial points. In two participants, head movement could not be measured due to technical issues. Eye movements were recorded using an infrared eye-tracker (EyeLink CL-OC, SR Research Ltd., Canada) at a sampling rate of 1000 Hz. For labeling of trials and vocal onset detection in vocalized trials, the participants' responses were recorded with a microphone integrated into the MEG System, at a sampling rate of 2343.75 Hz. An additional microphone (MD 419, Sennheiser electronic GmbH & Co KG, Germany) with a sampling rate of 44.1 kHz was used to record a higher quality audio trace for the labelling of production types and vowels. In a separate session, we acquired structural T1-weighted MRIs (3 Tesla MAGNETOM, Siemens Healthcare GmbH, Germany) for source reconstruction based on each participants individual's anatomy (resolution: 1 mm^3, MPRAGE).

Data Preprocessing

Technically caused channel jumps were detected and corrected and time lags between digital triggers and actual stimulus presentation were corrected based on a photodiode signal. For one subject, we excluded a noisy channel from the analysis. We low-pass-filtered the MEG data at 30 Hz (6th order, zero-phase Butterworth IIR filter) and down-sampled to 300 Hz. Each trial was baseline-corrected using the 500 ms preceding the onset of the first visual cue. To reduce electromagnetic artifacts, we ran independent component analysis (ICA) on the data. To ensure convergence of the ICA algorithm, the data was high-pass-filtered at 0.05 Hz. However, we applied the resulting unmixing matrix to the original data without any high-pass filter, and removed artefact components like heart beats and other small muscle activities. Vocal onset detection in vocalized trials was possible for 37 sessions. Audio traces were

missing due to technical issues in the remaining session. However, performance was closely

monitored during the recordings and immediately corrected, if participants vocalized before the

go-cue appeared. The available audio traces were smoothed with a median-based sliding

window model (window size 42.66ms). A participant-wise threshold served for onset detection.

For that, the root mean square (rms) of a 1000 ms period of one of the imagined trials was

calculated. The respective standard deviation was multiplied by eight and added to the rms. In

case of low vocalization amplitudes due to very soft voices of a few participants, the threshold

was manually adjusted by slightly decreasing it until the onsets could reliably be detected. The

vocalization was correctly performed, with an onset after the go-cue, in 99% of the trials.

For all subsequent analyses, only correct trials were used. Those were trials with the correct

production type, and, in case of vocalized trials, with the correct vowel and a vocal onset after

the go-cue. For the nine sessions with a missing audio trace, only the correct vowel was

considered.

Cross-validated MANOVA

We estimated the amount of neural information about the variables of interest in the

MEG data with cross-validated MANOVA [9,10]. We performed 20 repetitions of

cvMANOVA for each session with five-fold cross-validation. All folds and repetitions

were subsequently averaged. We first estimated a noise covariance matrix using trials

from all conditions. Next, we estimated contrasts of beta weights of each condition in

a cross-validation fold's training set, which accounted for the "training" of the model.

The "testing" was done by estimating contrasts of beta weights in the respective fold's

test set. The dot product of these contrasts, normalized by the noise covariance,

served as an estimate of the true pattern distinctness:

440
$$D = trace\left(\frac{1}{n} B'_{train} C_{train} C_{train}^{-1} X'_{test} X_{test} C_{train} C_{test}^{-1} B_{test} \Sigma^{-1}\right)$$

441 where Σ^{-1} is the inverted noise covariance matrix, C_{train} is the contrast vector the model is

442 trained on, C_{test} is the test contrast vector and X_{test} is the design matrix indicating the unique

443 condition of each trial in the test set. B_{train} and B_{test} contain the regression parameters of a

444 multivariate general linear model:

445
$$B_{train} = X_{train}^{-1} Y_{train}$$

447
$$B_{test} = X_{test}^{-1} Y_{test}$$

448 where Y_{train} and Y_{test} are the training and test data sets. The inverted noise covariance matrix

449 was estimated with the mean of the time window from cue 1 offset to go cue onset:

450
$$B_{train_{tw}} = X_{train}^{-1} Y_{train_{tw}}$$

451
$$\Xi = Y_{train_{tw}} - X_{train} B_{train_{tw}}$$

452
$$\Sigma^{-1} = (fE - p - 1) \cdot (\Xi' \Xi)^{-1}$$

with fE as the degrees of freedom and p as the number of sources.

Technically, cvMANOVA is a multivariate information based cross-validated encoding

approach. However, it shares many similarities with common multivariate decoding methods

[29]. The measure of neural information about the variables of interest can, theoretically, also be

used to decode these variables on individual trials. Therefore, we refer to our results as

decoding results.

For all sensor level analyses, a subset of 137 approximately equally distributed sensors were

included. This was to ensure a sufficient number of trials in relation to the degrees of freedom

of the dataset.

Cross-decoding

With cvMANOVA we were able to decode across conditions, by training and testing on different

time points, variables, and levels of the variables. Therefore, the contrast vectors C_{train} and C_{test}

were constructed to only contain the respective conditions to be trained or tested on. With this

approach, we decoded across variables by using content (/u/ vs. /∂/) for the construction of the

training contrast C_{train} and production type (vocalized vs. imagined) for the construction of the

test contrast C_{test}. To estimate content information for both production types separately, we

constructed C_{train} based on trials including both production types, but C_{test} based on trials with

only one production type, respectively. The same principle was applied for separately decoding

the production type from both vowels. For decoding across time, we used the regression

parameters B_{train} from one time point and B_{test} from another. We applied this to all pairs of time

points. For cross-session decoding, we implemented a two-fold cross-validation such that trials

from session 1 and 2 served as training and test sets alternately.

Expected cross-decoding

To estimate a benchmark for the overlap between representations, we computed the expected

cross-information [10]. As the maximally possible amount of shared information between

contexts depends on the information available in each individual context, the strength of the

shared representation must be compared to the strength of both representations. Therefore,

we estimated the expected cross-decoding

$$E_{12} = \sqrt{|D_1 D_2|} \cdot sign(D_1) \cdot sign(D_2)$$

where D_1 and D_2 denote the pattern distinctness in the two contexts. If the representations

were identical, the cross-decoding D_{12} would approach E_{12}. Conclusively, cross-decoding

values smaller than E_{12} indicate that the representations are not identical and, therefore, not

fully overlapping.

Source estimation

We generated individual single-shell head models [30] based on each subjects' structural T1-

weighted MRI. Using linear spatial filtering [31], we estimated three-dimensional MEG source

activity at 457 equally spaced locations ~7 mm beneath the skull. For searchlight analysis, we

used the three dipole directions of each source and the respective immediate neighbors. The

lateralization index (LI) was computed by averaging the searchlight results for each

hemisphere and subtracting right from left. For cross-session decoding, the three orientations

were added and a subset of 229 equally spaced sources was used for decoding.

Statistical Analysis

Neural information was averaged within two different time windows (delay 1: from 250ms after

cue 1 offset to cue 2 onset; delay 2: from 250ms after cue 2 offset to go-cue onset). For testing

the significance of neural information and cross-information to be larger than 0 we employed

one-tailed one-sample t-tests. One-tailed paired t-tests were applied for testing cross-time and

cross-variable information to be smaller than the expected cross-information. For the

comparison of content information in both production types and vice versa, we used two-tailed

paired t-tests. All p-values were FDR corrected for the number of tested time intervals [32].

Potential cue confound

We identified the source of a potential confound: due to the blocked design, in combination

with possible non-stationarities in the dataset, content- and production information could

theoretically be influenced by representations of the physical cue itself, even though the cue

appearance and its meaning were counterbalanced. Briefly, if noise led to independent shifts

of the neural activity patterns of both cue options, this could mistakenly be identified as

information about the variable indicated by the cue. To make sure our results were not

dependent on this potential confound, we took two measures. First, we excluded the 250 ms

after cue onset in both time windows used for statistical analysis, as this was the time period

that would likely be affected. Secondly, we performed cross-session decoding to confirm that

content- and production information were present if the possible cue confound was accounted

for.

Visualization

For all line plots, data were smoothed with a 100 ms Hanning window (full width at half

maximum).

Software

All analyses were performed using the Fieldtrip toolbox [33] and custom code in MATLAB

(R2014a & 2017b).

Data availability

The preprocessed data are available from the authors upon reasonable request.

Code availability

The analysis code is available from the authors upon reasonable request.

524 Acknowledgements

525 We thank Gabriele Walker-Dietrich, Jürgen Dax and Christoph Braun for help with data

526 acquisition. This research was supported by the European Research Council (ERC) StG

527 335880 (M.S.) and CoG 864491 (M.S.) and by the Evangelisches Studienwerk Villigst (V.V.).

528 Author Contributions

529 Conceptualization: M.S., V.A.V., F.S., D.H., S.R.H.; investigation: V.A.V.; formal analysis:

530 V.A.V., F.S.; resources: F.S., D.H., M.S.; writing – original draft preparation: V.A.V.; writing –

531 review and editing: F.S., M.S.; supervision: M.S., F.S.; funding acquisition: M.S., V.A.V.

532 Competing Interests

533 All authors declare no competing interests.

534 References

1. Vitevitch, M. S. The influence of sublexical and lexical representations on the processing

of spoken words in English. *Clin. Linguist. Phon.* **17**, 487–499 (2003).

2. Whitford, T. J. *et al.* Neurophysiological evidence of efference copies to inner speech.

eLife **6**, e28197 (2017).

3. Perrone-Bertolotti, M., Rapin, L., Lachaux, J.-P., Baciu, M. & Lœvenbruck, H. What is that

little voice inside my head? Inner speech phenomenology, its role in cognitive

performance, and its relation to self-monitoring. *Behav. Brain Res.* **261**, 220–239 (2014).

4. Corley, M., Brocklehurst, P. H. & Moat, H. S. Error biases in inner and overt speech:

Evidence from tongue twisters. *J. Exp. Psychol. Learn. Mem. Cogn.* **37**, 162–175 (2011).

5. Stephan, F., Saalbach, H. & Rossi, S. The Brain Differentially Prepares Inner and Overt

Speech Production: Electrophysiological and Vascular Evidence. *Brain Sci.* **10**, 148

(2020).

6. Trettenbrein, P. C., Papitto, G., Friederici, A. D. & Zaccarella, E. Functional neuroanatomy

of language without speech: An ALE meta-analysis of sign language. *Hum. Brain Mapp.*

42, 699–712 (2021).

7. Zhang, W., Liu, Y., Wang, X. & Tian, X. The dynamic and task-dependent representational

transformation between the motor and sensory systems during speech production. *Cogn.*

Neurosci. **11**, 194–204 (2020).

8. Oppenheim, G. M. & Dell, G. S. Motor movement matters: The flexible abstractness of

inner speech. *Mem. Cognit.* **38**, 1147–1160 (2010).

9. Allefeld, C. & Haynes, J.-D. Searchlight-based multi-voxel pattern analysis of fMRI by

cross-validated MANOVA. *NeuroImage* **89**, 345–357 (2014).

10. Sandhaeger, F., Omejc, N., Pape, A.-A. & Siegel, M. *Abstract neuronal choice signals*

during embodied decisions. http://biorxiv.org/lookup/doi/10.1101/2020.10.02.323832

(2020) doi:10.1101/2020.10.02.323832.

11. Cogan, G. B. *et al.* Sensory–motor transformations for speech occur bilaterally. *Nature*

507, 94–98 (2014).

12. Peeva, M. G. *et al.* Distinct representations of phonemes, syllables, and supra-syllabic

sequences in the speech production network. *NeuroImage* **50**, 626–638 (2010).

13. King, J.-R. & Dehaene, S. Characterizing the dynamics of mental representations: the

temporal generalization method. *Trends Cogn. Sci.* **18**, 203–210 (2014).

14. Ewald, A., Aristei, S., Nolte, G. & Rahman, R. A. Brain Oscillations and Functional

Connectivity during Overt Language Production. *Front. Psychol.* **3**, (2012).

15. Flinker, A. *et al.* Redefining the role of Broca's area in speech. *Proc. Natl. Acad. Sci.* **112**,

2871–2875 (2015).

16. Long, M. A. *et al.* Functional Segregation of Cortical Regions Underlying Speech Timing

and Articulation. *Neuron* **89**, 1187–1193 (2016).

17. Salmelin, R., Schnitzler, A., Schmitz, F. & Freund, H.-J. Single word reading in

developmental stutterers and fluent speakers. *Brain* **123**, 1184–1202 (2000).

18. Schulz, G. M., Varga, M., Jeffires, K., Ludlow, C. L. & Braun, A. R. Functional

Neuroanatomy of Human Vocalization: An H215O PET Study. *Cereb. Cortex* **15**, 1835–

1847 (2005).

19. Papoutsi, M. *et al.* From Phonemes to Articulatory Codes: An fMRI Study of the Role of

Broca's Area in Speech Production. *Cereb. Cortex* **19**, 2156–2165 (2009).

20. Irwin, O. C. Infant Speech: Development of Vowel Sounds. *J. Speech Hear. Disord.* **13**,

31–34 (1948).

21. Jürgens, U. Neural pathways underlying vocal control. *Neurosci. Biobehav. Rev.* 24

(2002).

22. Hage, S. R. & Nieder, A. Dual Neural Network Model for the Evolution of Speech and

Language. *Trends Neurosci.* **39**, 813–829 (2016).

23. Tian, X. Mental imagery of speech and movement implicates the dynamics of internal

forward models. *Front. Psychol.* **1**, (2010).

24. Shergill, S. S. *et al.* Modulation of activity in temporal cortex during generation of inner

speech. *Hum. Brain Mapp.* **16**, 219–227 (2002).

25. Houde, J. F., Nagarajan, S. S., Sekihara, K. & Merzenich, M. M. Modulation of the

Auditory Cortex during Speech: An MEG Study. *J. Cogn. Neurosci.* **14**, 1125–1138

(2002).

26. Gavrilov, N., Hage, S. R. & Nieder, A. Functional Specialization of the Primate Frontal

Lobe during Cognitive Control of Vocalizations. *Cell Rep.* **21**, 2393–2406 (2017).

27. Conant, D. F., Bouchard, K. E., Leonard, M. K. & Chang, E. F. Human Sensorimotor

Cortex Control of Directly Measured Vocal Tract Movements during Vowel Production. *J.*

Neurosci. **38**, 2955–2966 (2018).

28. Correia, J. M., Caballero-Gaudes, C., Guediche, S. & Carreiras, M. Phonatory and

articulatory representations of speech production in cortical and subcortical fMRI

responses. *Sci. Rep.* **10**, 4529 (2020).

29. Hebart, M. N. & Baker, C. I. Deconstructing multivariate decoding for the study of brain

function. *NeuroImage* **180**, 4–18 (2018).

30. Nolte, G. The magnetic lead field theorem in the quasi-static approximation and its use

for magnetoencephalography forward calculation in realistic volume conductors. *Phys.*

Med. Biol. **48**, 3637–3652 (2003).

31. Van Veen, B. D., Van Drongelen, W., Yuchtman, M. & Suzuki, A. Localization of brain

electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Trans.*

Biomed. Eng. **44**, 867–880 (1997).

32. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300

(1995).

33. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open Source Software

for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Comput.*

Intell. Neurosci. **2011**, 1–9 (2011).