Changes in a new type of genomic accordion may open the pallets to increased

monkeypox transmissibility

Sara Monzón[1#], Sarai Varona[1#], Anabel Negredo[2,3#], Juan Angel Patiño-Galindo[4], Santiago Vidal-Freire[4], Angel Zaballos[5], Eva Orviz[6], Oskar Ayerdi[6], Ana Muñoz-García[6], Alberto Delgado-Iribarren[6], Vicente Estrada[3,6], Cristina García[2], Francisca Molero[2], Patricia Sánchez[2,3], Montserrat Torres[2], Ana Vázquez[2,7], Juan-Carlos Galán[7,8], Ignacio Torres[9], Manuel Causse del Río[10], Laura Merino[11], Marcos López[12], Alicia Galar[13], Laura Cardeñoso[14], Almudena Gutiérrez[15], Cristina Loras[16], Isabel Escribano[17], Marta Elena Alvarez-Argüelles[18], Leticia del Río[19], María Simón[20], MªAngeles Meléndez[21], Juan Camacho[2], Laura Herrero[2], Pilar Jiménez Sancho[5], Maria Luisa Navarro-Rico[5], Jens H. Kuhn[22], Mariano Sanchez-Lockhart[23], Nicholas Di Paola[23], Jeffrey R. Kugelman[23], Elaina Giannetti[4], Susana Guerra[4,24,25], Adolfo García-Sastre[4,24,26,27,28], Gustavo Palacios[4,24&*], Isabel Cuesta[1&], and Maripaz P. Sánchez-Seco[2,3&]

[1]Unidad de Bioinformática, Unidades Centrales Científico Técnicas, Instituto de Salud Carlos III, 28029 Madrid, Spain

[2]Centro Nacional de Microbiología, Instituto de Salud Carlos III, 28029 Madrid, Spain

[3]Centro de Investigación Biomédica en Red de Enfermedades Infecciosas (CIBERINFEC), Instituto de Salud Carlos III, 28029 Madrid, Spain

[4]Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[5]Unidad de Genómica, Unidades Centrales Científico Técnicas, Instituto de Salud Carlos III, 28029 Madrid, Spain

[6]Centro Sanitario Sandoval, Hospital Clínico San Carlos, 28040 Madrid, Spain

[7]Centro de Investigación Biomédica en Red de Epidemiología y Salud Pública (CIBERESP),

Instituto de Salud Carlos III, 28029 Madrid, Spain

[8]Servicio de Microbiología, Hospital Universitario Ramón y Cajal, Instituto Ramón y Cajal

de Investigación Sanitaria (IRYCIS), 28034 Madrid, Spain

[9]Servicio de Microbiología, Hospital Clínico Universitario, Instituto de Investigación

INCLIVA, 46010 Valencia, Spain

[10]Unidad de Microbiología, Hospital Universitario Reina Sofía, Instituto Maimónides de

Investigación Biomédica de Córdoba, 14004 Córdoba, Spain

[11]Unidad Clínico de Enfermedades Infecciosas, Microbiología y Medicina Preventiva,

Hospital Universitario Virgen del Rocío, 41013 Sevilla, Spain

[12]Servicio de Microbiología y Parasitología, Hospital Universitario Puerta de Hierro

Majadahonda, 28222 Madrid, Spain

[13]Servicio de Microbiología Clínico y Enfermedades Infecciosas, Hospital General

Universitario Gregorio Marañón, 28007 Madrid, Spain

[14]Servicio de Microbiología, Instituto de Investigación Sanitaria, Hospital Universitario de la

Princesa, 28006 Madrid, Spain

[15]Servicio de Microbiología y Parasitología Clínica, Hospital Universitario La Paz, 28046

Madrid, Spain

[16]Servicio de Microbiología, Hospital General y Universitario, 13005 Ciudad Real, Spain

[17]Hospital General Universitario Dr. Balmis, 03010 Alicante, Spain

[18]Servicio de Microbiología, Hospital Universitario Central, 33006 Asturias, Spain

[19]Hospital Quironsalud Torrevieja, 03184 Alicante, Spain

[20]Servicio de Microbiología, Hospital Central de la Defensa "Gómez Ulla", 28947 Madrid,

Spain

[21]Servicio de Microbiología y Parasitología, Hospital Universitario 12 de Octubre, 28041

Madrid, Spain

[22]Integrated Research Facility at Fort Detrick, National Institute of Allergy and Infectious

Diseases, National Institutes of Health, Fort Detrick, Frederick, MD 21702, USA

[23]United States Army Research Institute for Infectious Disease, Fort Detrick, Frederick, MD

21702, USA

[24]Global Health Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai,

New York, NY 10029, USA

[25]Departmento de Medicina Preventiva, Salud Publica y Microbiología, Universidad

Autónoma de Madrid, 28029 Madrid, Spain

[26]Department of Medicine, Division of Infectious Diseases, Icahn School of Medicine at

Mount Sinai, New York, NY 10029, USA

[27]The Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY

10029, USA

[28]Department of Pathology, Molecular and Cell-Based Medicine, Icahn School of Medicine

at Mount Sinai, New York, NY 10029, USA

66
67 [#]These authors contributed equally to this work.

68
[&]These senior authors contributed equally to this work.

Correspondence

[*]Gustavo Palacios: gustavo.palacios@mssm.edu

SUMMARY

The currently expanding monkeypox epidemic is caused by a subclade IIb descendant of a

monkeypox virus (MPXV) lineage traced back to Nigeria in 1971. In contrast to monkeypox

cases caused by clade I and subclade IIa MPXV, the prognosis of current cases is generally

favorable, but person-to-person transmission is much more efficient. MPXV evolution is

driven by selective pressure from hosts and loss of virus–host interacting genes. However,

there is no satisfactory genetic explanation using single-nucleotide polymorphisms (SNPs)

for the observed increased MPXV transmissibility. We hypothesized that key genomic

changes may occur in the genome's low-complexity regions (LCRs), which are highly

challenging to sequence and have been dismissed as uninformative. Using a combination of

highly sensitive techniques, we determined a first high-quality MPXV genome sequence of a

representative of the current epidemic with LCRs resolved at unprecedented accuracy. This

effort revealed significant variation in short-tandem repeats within LCRs. We demonstrate

that LCR entropy in the MPXV genome is significantly higher than that of SNPs and that

LCRs are not randomly distributed. *In silico* analyses indicate that expression, translation,

stability, or function of MPXV orthologous poxvirus genes (OPGs) 153, 204, and 208 could

be affected in a manner consistent with the established "genomic accordion" evolutionary

strategies of orthopoxviruses. Consequently, we posit that genomic studies focusing on

phenotypic MPXV clade-/subclade-/lineage-/strain differences should change their focus to

the study of LCR variability instead of SNP variability.

Keywords

Chordopoxvirinae, genomic accordion, low-complexity region, molecular epidemiology,

monkeypox, MPX, MPXV, OPG, orthopoxvirus, *Poxviridae*, short tandem repeat, STR,

transmission, genomics, bioinformatics, computational biology

INTRODUCTION

Monkeypox virus (MPXV) is a double-stranded DNA virus that belongs to genus

Orthopoxvirus (varidnavirian *Nucleocytoviricota*: *Poxviridae*: *Chordopoxvirinae*) along with

other human viruses, such as vaccinia virus (VACV) and variola virus (VARV) (International

Committee on Taxonomy of Viruses, 2022). First encountered in 1958 in crab-eating

macaques imported to Belgium (Magnus et al., 2009), MPXV has caused sporadic disease

outbreaks in humans since the 1970s in Eastern, Middle, and Western Africa, totaling

approximately 25,000 cases (case fatality rate 1–10%) (Beer and Rao, 2019), and also

sporadic disease outbreaks among wild monkeys and apes (Patrono et al., 2020; Radonić et

al., 2014). Exposure to MPXV animal reservoirs, in particular rope squirrels and sun

squirrels, is a significant risk factor of human infections (Khodakevich et al., 1988).

The human disease caused by MPXV is designated as monkeypox in the World

Health Organization (WHO) International Classification of Diseases, Eleventh Revision

(ICD-11; code 1E71) (World Health Organization, 2022a). Phylogenetically, historic MPXV

isolates cluster into two clades (Likos et al., 2005), designated I and II (Happi et al., 2022;

World Health Organization, 2022b). Clade I viruses are considered more virulent and

transmissible than clade II viruses (Damon, 2011; Likos *et al.*, 2005; World Health

Organization, 2022b).

Since May 2022, multiple European countries have reported a continuously increasing

number of MPXV infections and associated disease, including clusters of cases associated

with potential superspreading events in Belgium, Spain, and the United Kingdom (UK). As of

September 30, 2022, a total of 68,428 cases had been reported in 106

countries/territories/areas in all six WHO regions. Of these, 67,739 were in 99 countries that

have not reported MPXV infections prior to 2022 (Centers for Disease Control and

Prevention, 2022). This rapid increase in infections prompted WHO to declare this epidemic

a Public Health Emergency of International Concern (PHEIC) (Nuzzo et al., 2022). The

viruses of the 2022 epidemic belong to subclade IIb (Antinori et al., 2022; Nextstrain, 2022;

Vivancos et al., 2022), a line of descent of MPXV that had been circulating in Nigeria, likely

since 1971 (Faye et al., 2018).

The clinical presentation monkeypox caused by clade I or IIa includes fever,

headache, lymphadenopathy, and/or malaise, followed by a characteristic rash that progresses

centrifugally from maculopapules via vesicles and pustules to crusts that may occur on the

face, body, mucous membranes, palms of the hands, and soles of the feet (Ježek et al., 1987).

The clinical presentation of the current MPXV subclade IIb infections diverges from classical

monkeypox by having a good prognosis, self-limiting but infectious skin lesions (typically

emerging at and restricted to the genital, perineal/perianal, and/or peri-oral areas) before the

development of fever, lymphadenopathy, and malaise. Generalized disease usually manifests

with a rash that has not been widely observed in the current outbreak. Human-to-human

transmission is substantially higher in subclade IIb MPXV-associated outbreaks than in clade

I and clade IIa MPXV (Bunge et al., 2022; Otu et al., 2022; Thornhill et al., 2022; Ulaeto et

al., 2022; Vusirikala et al., 2022). The R_0 for MXPV IIb among men who have sex with men

(MSM) is higher than 1. Transmission may be catalyzed by decreasing protection associated

with the VARV/smallpox vaccination campaign that ended in 1980 (Grant et al., 2020;

Rimoin et al., 2010). Furthermore, a change in transmission route may be the cause of the

difference in clinical presentation and pathogenesis as was shown in animal models

(Reynolds et al., 2006).

Orthopoxvirus infections are classified as systemic or localized illnesses. Localized

usually means that signs are restricted to the site of MPXV entry, as described in the 2022

outbreak. The involved orthopoxvirus, its route of entry, and the immune status of the host

are usually the only determinants of generalized or localized infection. Different mechanisms

of virion entry and egress, as well as virus-encoded host factors, are the main biological

determinants (Liu et al., 2019; McFadden, 2005; Moss, 2006; 2016; Roberts and Smith,

2008). Changes in the genome of the current MPXV variant, such as gene loss (Kugelman et

al., 2014), may explain both trends.

The MPXV genome is a linear, ≈197-kb-long double-stranded DNA with covalently

closed hairpin ends. The genome's densely packed orthologous poxvirus genes (OPGs)

(Senkevich et al., 2021) are distributed over a central conserved region ("core") and flanking

terminal regions, each of which ends in identical but oppositely oriented ≈6.4-kb-long

terminal inverted repetitions (ITRs). Roughly 193 open reading frames (ORFs) encode

proteins with ≥60 amino-acid residues. "Housekeeping" proteins involved in MPXV

transcription, replication, and virion assembly are encoded by OPGs located in the central

conserved region, whereas proteins involved in host range and pathogenesis are mostly

encoded by OPGs located in the terminal regions. Like all orthopoxvirus genomes, the

MPXV genome contains numerous tandem repeats in the ITRs as well as nucleotide

homopolymers all over the genome (Moss and Smith, 2021; Shchelkunov et al., 2002; Wittek

and Moss, 1980). However, we also observed other similar structures through the MPXV

genome in the form of short tandem repeats (STRs). Moreover, initial observations appear to

indicate that these STRs (which may consist of dinucleotide, trinucleotide, or more complex

palindromic repeats) are localized in areas where more variation is observed suggesting a

crucial role in the MXPV biology and evolution.

Orthopoxviruses rapidly acquire higher fitness by massive gene amplification

(genome expansion) when encountering severe bottlenecks *in vitro*. This amplification, akin

to gene reduplication in organismal evolution, enables gene copies to accumulate mutations,

potentially resulting in new protein variants that can overcome the bottlenecks. Subsequent

gene copy reduction (genome contraction) offsets the costs associated with increasing

genome length, thereby retaining the adaptive mutations (Elde et al., 2012). Orthopoxviruses

also rapidly adapt to selective pressures by single-nucleotide insertions (genome expansion)

or deletions (genome contractions) within poly-A or poly-T stretches, resulting in easily

reversible gene-inactivating or re-activating frameshifts (Senkevich et al., 2020). These

rhythmic genome expansions and contractions are referred to as "genomic accordions" at the

gene and base level (Elde *et al.*, 2012). Given the overall conservation of STRs in

orthopoxvirus genomes, we hypothesized that their variation could be a third type of genomic

accordion and that, overall, this type of adaptation (which we designate here as low-

complexity regions [LCRs]), rather than single-nucleotide polymorphisms (SNPs), could be

the key to understanding the unusual epidemiology of 2022 subclade IIb MPXV.

RESULTS

***De novo* assembly of subclade II lineage B.1 monkeypox virus (MPXV) genome**

sequence 353R

Using a template-based mapping approach, shotgun metagenomic short-read-based

sequencing of nucleic acids in vesicular lesion swabs from Spanish monkeypox patients

resulted in the determination of 48 MPXV consensus genome sequences with at least 10X

read depth. A median of 39,697,742 high-quality reads per swab (maximum = 111,030,976;

minimum = 7,780,032) were obtained using a NovaSeq 6000 Illumina sequencer. Although

98.12% of the reads were assigned as being of human origin, a median of 74,085 MPXV

reads (maximum = 27,516,891; minimum = 30,854) sufficed to cover >99% of the genome

(**Table S1**).

194   Read mapping indicated that, as expected, LCRs of the MPXV genome were mostly

unresolved. More importantly, those results were biased by the MPXV genome sequence

used for mapping (subclade II lineage A MPXV isolate MPXV-M5312_HM12_Rivers):

LCRs were resolved by the used reference-mapping software tools "following" the sequence

provided in the reference genome instead of reporting the actual sequence (**Figure S1A**). To

determine the actual LCR sequence, we explored different assembly strategies generally used

for resolving eukaryotic genomes, which mostly combine different sequencing technologies.

To increase the chances of success, we applied these technologies to a monkeypox patient

sample with a high proportion of high-quality viral reads (swab 353R). The *de novo* assembly

obtained from NovaSeq (2x150-bp pair-ended reads), MiSeq (2x300-bp pair-ended reads),

and Nanopore sequencing generated 3, 2, and 1 contigs belonging to MPXV, covering 97%,

97%, and 101% of the MPXV-M5312_HM12_Rivers sequence, respectively (**Figure 1**).

Characterization and validation of non-randomly distributed low-complexity regions

(LCRs) in monkeypox virus (MPXV) genome sequence 353R

We applied a systematic approach for LCR discovery to the MPXV 353R sequence that

resulted in the identification of 21 LCRs (13 STRs, 8 homopolymers; **Tables 1** and **S2**). Two

pairs of LCRs (1/4 and 10/11) are located in the ITRs and are identical copies in reverse-

complementary form. Consequently, we moved forward with 19 LCRs.

In general, LCRs were resolved using the assembly obtained from single-molecule

sequencing and further validated using short-read sequencing since most sequences were 13–

67 bp long and therefore were covered by reads from each side or flanking region without

mismatches (**Figure S1B**; **File S1)**. All LCRs (except pair 1/4 and 3) were validated this way.

LCR pair 1/4 (256 bp) and LCR3 (468 bp) were only resolved with single-molecule

sequencing reads due to their lengths (**Table 2**).

LCR3 contains a complex tandem repeat with the sequence ATAT [ACATTATAT]$_n$.

Our analysis indicated n=52. No publicly available MPXV genome sequence contains a

tandem repeat of similar length. However, applying the analysis to 35 publicly available

MPXV National Center for Biotechnology Information (NCBI) Sequence Read Archive

(SRA) datasets of single-molecule raw reads allowed us to resolve the LCR3 of some (**Figure**

2A). Fifteen datasets revealed supporting long reads that include both LCR3 flanking regions.

Interestingly, four subclade IIb lineage B.1 MPXV sequences associated with the 2022

monkeypox epidemic and available in SRA have n=54–62 repeats in LCR3. Their number of

repeats separate these sequences from 2018–2019 subclade IIb lineage A sequences that have

n=12–42 repeats in LCR3, indicating LCR3 as a region of genomic instability and high

variability.

LCR pair 1/4 contains a tandem repeat with the sequence

[AACTAACTTATGACTT]$_n$. Our analysis indicated n=16. Instead, the sequences of the

subclade IIb lineage B.1 MPXV isolate MPXV_USA_2022_MA001 and lineage A reference

isolate MPXV-M5312_HM12_Rivers LCR pair 1/4 have n=8 (**Table 3**). Inclusion of the

NCBI SRA datasets into the analysis confirmed the n=16 value (**Figure 2B**). In addition, the

analysis revealed subclade IIb lineage-specific repeat differences in LCR pair 1/4. Lineage

A.1 viruses are polymorphic, having 14 repeats (n=1 sequences), 16 (n=3), 17 (n=7), and 19

(n=1); lineage A.2 viruses have n=23–26 repeats; lineage A viruses have n=32, 43, 53, or 71

repeats. In contrast, lineage B1 viruses consistently have n=16. While LCR3 appears to have

"increased" in length since the spillover, LCR pair 1/4 appears to be decreasing in length,

thus behaving like an "accordion" over time.

The subclade II lineage B.1 353R and MPXV_USA_2022_MA001 genome sequences

have the same 67 SNPs called against the subclade II lineage A reference isolate MPXV-

M5312_HM12_Rivers genome sequence. Additionally, the 353R sequence has two

additional paired SNPs in the left and right ITRs (5,595G→A; 191,615C→T compared with

the MPXV-M5312_HM12_Rivers sequence) that result in the introduction of a stop codon in

OPG015. We observed this variation in only two other patients among our sample set. The

353R and MPXV_USA_2022_MA001 sequences also differ by two number of insertion (ins)

or deletion (del) of bases (indels) at positions 133,077 and 173,273, respectively, which

correspond to differences in LCR2 and LCR5, respectively. As a result of the resolution

LCRs, sequence 353R differs by 1,342 bp in genome length compared with the

MPXV_USA_2022_MA001 sequence and 1,338 bp compared with the MPXV-

M5312_HM12_Rivers sequence. Most of the variation is due to differences in the length of

LCR pair 1/4 and LCR3, along with minor length differences in LCR2, LCR5, and LCR pair

10/11 (**Table 3**). In general, the number of repeats (n) found with the hybrid assembly

approach we used here doubled the length of LCRs.

Based on the higher resolution of the MPXV 353R genome sequence, in particular

regarding LCRs, we propose this sequence as the new MPXV high-quality genome (HQG)

reference sequence according to the sequence quality standards defined in (Ladner et al.,

2014).

Low-complexity regions (LCRs) are non-randomly distributed in the monkeypox virus

(MPXV) genome

We compared the distribution of LCRs between the different major functional protein OPG

groups following the classification of (Senkevich *et al.*, 2021). Differences between

functional groups were statistically significant (Kruskal–Wallis test, χ^2 p-value <0.001).

Pairwise analysis demonstrated that the functional group "core" (orthopoxvirus genomic

central conserved region) includes LCRs at a significantly lower frequency (multiple

pairwise-comparison Wilcoxon test) than functional groups "ANK/PRANC" (false discovery

rate [FDR]-corrected p-value <0.0001), "Bcl-2 domain" (corrected p-value = 0.04), and

"accessory" (FDR-corrected p-value <0.0001) (**Figure 3A**). These analyses indicate that

LCRs in orthopoxvirus genomes are non-randomly distributed and that there is a significant

purifying selection force against introducing LCRs in central conserved region areas.

Next, we compared the degree of diversity among the 21 identified LCRs with the

observed SNP variability that had been the focus of the field. In the 353R HQG sequence,

LCRs 2, 5, 7, 10, 11, and 21 had intra-host genetic diversity, with entropy values that ranged

from 0.18 (LCR7) to 1.66 (LCR2), with an average of 0.81 and a standard deviation (SD) of

0.64 among them (**Table 2**). Only five nucleotide positions (1,285; 6,412; 88,807; 133,894;

and 145,431) had intra-host genetic diversity at the SNP level. The entropy values ranged

from 0.17 (position 133,894) to 0.69 (position 6,412), with an average of 0.38 and a SD=0.21

among them. Interestingly, a student's t-test revealed a significantly higher level of diversity

in LCRs than in SNPs (p-value = 0.021; **Figure 3B**).

Then, we characterized, collected, and compared the allele frequencies for all LCRs

from all dataset samples (**Table S3**) applying the filters described above. Our analyses

revealed that the average inter-sample Euclidean distances at LCRs ranged from 0.05

(LCR21) to 0.73 (LCR2) (**Figure 3C**). We found statistically significant differences between

LCRs (Kruskal–Wallis χ^2 p-value <0.001). More specifically, multiple pairwise comparison

Wilcoxon test results showed that all LCRs have significantly different levels of inter-sample

distances (FDR-corrected p-values < 0.001), except in case of the LCR10 versus LCR11

(FDR-corrected p-value = 0.48) and LCR2 versus LCR5 (FDR-corrected p-value = 0.25)

(**Figure 3C**). Average distances in SNPs were 0.0018–0.4168. Our randomization tests

revealed that all LCRs have a significantly higher level of inter-sample diversity than the

SNPs (all FDR-corrected p-values <0.05) (**Figure 3C**). These analyses uncovered that most

of the variability in the orthopoxvirus genome is located in LCR. Consequently, we posit that

studies focusing on phenotypic MPXV (and likely other orthopoxvirus) clade-/subclade-

/lineage-/strain differences due to genomic sequence variation should change their focus to

the study of LCR variability instead of SNP variability.

Low-complexity regions (LCRs) might be more phylogenetically informative than

single-nucleotide polymorphisms (SNPs) for inter-host sequence analysis

Analysis of only two monkeypox patient samples (353R and 349R) resulted in sufficient

sequence coverage information to enable allele frequency comparison in most LCRs (**Figure**

4A). Their side-by-side comparison revealed differences in allele frequency in some of them

(LCR2, LCR5, and LCR pair 10/11) (**Figure 4B**). The sequence coverage achieved with the

remaining samples only enabled to unequivocally resolve an LCR subset (i.e., covering both

flanking regions: LCRs 2, 5, 7, 8, 9, and pair 10/11). LCR8 and LCR9 were identical across

the entire sample set. However, LCR7 and LCR pair 10/11 had considerable intra-host

variation, as well as differences in the preponderant allele (LCR pair 10/11) between samples

(**Figure 4C**).

Phylogenetic (**Figure S2A**) and haplotype network (**Figure S2B**) analyses of the

sequences determined from the monkeypox patient samples yielded limited information

regarding the outbreak. Most sequences were highly similar and are, therefore, part of the

basal ancestral MPXV subclade IIb lineage B1 node. Some sequences formed supported

clusters: Sequences clustered into groups (**Table S4**):

- group 1 (lineage B.1): sequences from patients 395, 399, and 441 and Floridian

MPXV isolate MPXV_USA_2022_FL002 (GenBank #ON676704);

- group 2 (lineage B.1): sequences from patients 347, 352R, 353R, 416, and Spanish

MPXV isolate MPXV/ES0001/HUGTiP/2022 (GenBank #ON622718); all share a

stop-codon mutation in OPG015;

•   group 3 (lineage B.1.3): sequences from patients 22,369 formed with Slovenian

320   MPXV isolate SLO (GenBank #ON609725.2), French isolate

321   MPXV_FR_HCL0001_2022 (GenBank #ON622722), and 38 other sequences

322   worldwide; defined by NBT03_gp174 mutation G190,660A, resulting in an R84K

323   amino-acid residue change;

324   •   group 4 (lineage B.1): sequences from patients 417 and 2,437;

325   •   group 5 (lineage B.1.1): sequences from patients 698; 1,300; 2,388; 2,428; German

326   MPXV isolate MPXV/Germany/2022/RKI01 (GenBank #ON637938.1) and 97 other

327   sequences worldwide; defined by OPG094 mutation G74,360A resulting in a R194H

328   amino-acid residue change); and

•   group 6 (lineage B.1): sequences from patients 2,309 and 2,317.

Only one epidemiological link among group samples was identified; patients 395 and

399 came from sexual partners who attended events in Portugal and Spain.

In summary, at least at this time in the monkeypox epidemic, there appears to be

limited value in full-genome SNP characterization for transmission analysis.

Conservation and variation in proteins encoded by orthologous poxvirus gene (OPG)

and codon usage analysis in OPG low-complexity regions (LCRs)

Analyses showed that LCRs were associated with intra-host and inter-sample variation.

Although most LCRs that showed variability in our sample set (pair 1/4, 2, 3, 5, 7, pair 10/11,

and 21), three (3, 7, and 21) are located in regions that, considering orthopoxvirus

evolutionary history, are associated with virulence or transmission (Chen et al., 2005;

Kastenmayer et al., 2014). Noteworthy, three of the 21 highly repetitive areas identified in

our intra-host variation analysis (those of LCRs 5, 6, and 7) are located in a defined central

conserved region of the orthopoxvirus genome between positions 130,000 and 138,000

(**Figure 1**). This region contains OPG152 (which is truncated in the MPXV genome),

OPG153 (directly affected by LCR7), and OPG154. LCR7 is the only STR that is located at

the center of a functional ORF). In contrast, LCR3 and LCR21 are situated in the

promoter/start area, potentially modifying the ORF start site. The repeat area of LCR7

encodes a poly-D homopolymer in a nonstructured region of OPG153 (**Figure 5A**). The

changes we uncovered result in the insertion of two isoleucyls. This change resembles the

primary structure found in clade I viruses. In contrast, pre-2017 African subclade IIa viruses

lack such insertions.

Another region of potential functional impact is the area between 170,000 to 180,000

that includes LCRs 19, 20, 2, 21 and 3) (**Figure 1**). The LCR3 repeat [CATTATATA]$_n$ is

located 21 bp upstream of the putative translation start site of OPG208. Importantly, a

methionyl codon is located immediately upstream of LCR3. The usage of this start codon

would result in the introduction of an Ile-Ile-Tyr repeat. This codon has a medium to low

probability of being used as start codon in the cognate mRNA (T base in position -3),

compared to a "strong" Kozak sequence of the downstream putative start codon.

Nevertheless, LCR3 remained in-frame in all clade II MPXV samples, indicating selective

pressure to maintain the possibility of alternative start translation (**Figure 5B**). Interestingly,

LCR3 is not located in-frame in most clade I viruses. This may be significant because

OPG208 is a member of a set of genes most likely responsible for increased virulence of

clade I MPXV compared to classical (pre-epidemic) clade IIa MPXV (Chen *et al.*, 2005). The

LCR3 tandem repeat [CATTATATA]$_n$ is present with n=52, 54, and 62 copies in epidemic

subclade IIb lineage B viruses (**Figure 2A**), whereas it is n=7, 37, and 27 in subclade IIa

MPXV isolates Sierra Leone (GenBank #AY741551), MPXV-WRAIR7-61 (GenBank

#AY603973), and MPXV-COP-58 (GenBank #AY753185) sequences, respectively (Chen *et*

al., 2005), as well as n=16 for clade I MPXV isolate Zaire-96-I-16 (GenBank #AF380138).

Interestingly, all publicly available subclade II lineage A single-molecule long-read sequence

data imply a repeat n<40 (**Figure 2A**). Alternatively, the LCR3 repeat sequence could also

alter promoter function. Interestingly, the repeat sequence also introduces codons with a low

usage ratio that are not be optimized for expression in primates. The codon triplet ATA,

which encodes isoleucyl, has a rare codon usage of 0.17 (Stothard, 2000) (**Figure 5B**).

Similarly, the STR downstream of LCR21 introduces a methionyl codon upstream of

the putative start codon for OPG204 (**Figure 5C**). Kozak sequence analysis revealed a

medium to high probability for translation compared with the putative start codon (**Figure

5C**).

The remaining LCRs (2, pair 4/1, and pair 10/11) are located downstream of known

ORFs; thus, their variation is less likely to be associated with a change in phenotype.

DISCUSSION

MPXV subclade IIb traces back to a human MPXV infection that likely occurred after spillover from a local animal reservoir in Ihie, Abia State, Nigeria, in 1971. An additional 10 human infections with MPXV of this lineage were detected through 1978, when this lineage seemed to have disappeared. However, in 2017, it reemerged in Yenagoa, Bayelsa State, Nigeria (Faye et al., 2018). Since then, hundreds of monkeypox cases have been reported and MPXV belonging to the IIb lineage has been sampled in several countries. But, there were no secondary cases prior to the 2022 epidemic (Cohen-Gihon et al., 2020; Ng et al., 2019; Vaughan et al., 2018; Yong et al., 2020).

Subclade IIb viruses cause monkeypox that presents differently than the classical disease caused by clade I and subclade IIa viruses, i.e., subclade IIb infections are associated with higher prevalence among adults rather than adolescents, are predominant in the MSM community, and are efficiently transmitted human-to-human from localized infectious skin lesions rather than requiring disseminated infection (Bunge et al., 2022; Otu et al., 2022; Thornhill et al., 2022; Ulaeto et al., 2022; Vusirikala et al., 2022).

Comparative genomics demonstrated obvious relationships between orthopoxvirus genotype and phenotype, driven by selective pressure from hosts (Baroudy and Moss, 1982; Chen et al., 2005; Esposito et al., 2006; Gubser et al., 2004; Hendrickson et al., 2010; Kugelman et al., 2014; Shchelkunov, 2012; Shchelkunov et al., 2001). Consequently, it was expected that increased MPXV genotype IIb human-to-human transmission would go hand in hand with genotypic changes. However, since orthopoxviruses genomes are organized in redundant ways (Bratke and McLysaght, 2008; Elde et al., 2012; McLysaght et al., 2003; Senkevich et al., 2020), genotypic changes were expected to be modulating, rather than radical.

Thus far, genomic characterization of the 2022 epidemic has focused on describing its

evolutionary history and tracking MPXV introductions into western countries. The 2022

MPXV cluster diverges from its predecessor viruses by an average of 50 SNPs. Of these, the

majority (n=24) are non-synonymous mutations with a second minority subset of

synonymous mutations (n=18) and a few intergenic differences (n=4) (Isidro et al., 2022). A

strong mutational bias was mainly attributed to the potential action of apolipoprotein B

mRNA-editing catalytic polypeptide-like 3 (APOBEC3) enzymes (O'Tool and Rambaut,

2022). Genetic variation, including deletion of immunomodulatory genes, also occurred

(Jones et al., 2022). MPXV sublineages mostly represent very small variations, usually

characterized by one or two SNP differences to basal nodes in phylogenetic analyses

(Nextstrain, 2022). Four of the MPXV genome sequences determined in this study can be

assigned to global lineage B.1.1, one sequence could be assigned to global lineage B.1.3, and

the remaining new sequences belong to lineage B1. We detected additional clusters that were

also defined by few SNPs, but only in one case could we identify an epidemiological link.

Thus, there appears to be a limited relationship between SNPs and epidemiology, which

might hint to sequencing errors. Consequently, MPXV genomic epidemiology might need a

change of focus.

Our analyses located considerable MPXV genomic variability in areas previously

considered of poor informative value, i.e., in LCRs. Because LCR entropy is significantly

higher than that of SNPs and LCRs are not randomly distributed in defined coding areas in

the genome, person-to-person transmission-associated changes were observed in the

immunomodulatory region (Kugelman et al., 2014), and genomic accordions are a rapid path

for adaptation of orthopoxvirus during serial passaging (Elde et al., 2012; Senkevich et al.,

2020), we posit that LCR changes might be associated with MPXV transmissibility

differences over time.

Eight LCRs had evident signs of intra-host and inter-sample variation (pair 1/4, 2, 3,

5, 6, 7, pair 10/11, 21). Five of them (5, 6, 7, 3, and 21) were co-located in two areas of the

MPXV genome: base pairs 130,000–135,000 (5, 6, and 7) which are in the central conserved

region of the OPXV genome in which most "housekeeping" genes are located; and base pairs

170,000–180,000 (3 and 21), which are located in the immunomodulatory area (**Figure 1**).

Three of those LCRs are located inside the putative translated regions of genes OPG153,

OPG204, and OPG208. Changes in OPG204 and OPG208 are located near the N-terminal

region and might involve modulating the expression of translation. The changes observed in

OPG153 stand out as they are located inside a region that is under high selective pressure for

transmission in a "housekeeping" orthologous poxvirus gene, which is involved in virion

attachment and egress (Senkevich *et al.*, 2021). Thus, we urge that these areas be scrutinized

for changes that might affect the MPXV interactome.

The OPG153 repeat results in a poly-Asp amino-acid homopolymer string (**Figure**

5A); the LCR repeat in OPG208 results in an Ile-Ile-Tyr repeat (**Figure 5B**); and the N-

terminal domain variation in OPG204 results in a Met-Lys repeat (**Figure 5C**). Self-

association guided by stretches of single amino-acid residue repeats may lead to the

formation of aggregates (Oma et al., 2004). Many human diseases are associated with

detrimental effects of homopolymers (Gatchel and Zoghbi, 2005; Shoubridge et al., 2007).

Expansion or contraction of the repeat may increase self-attraction and trigger disease. These

homopolymers also regulate the activity of transcription factors (Gemayel et al., 2015) or

direct proteins to different cellular compartments (Oma *et al.*, 2004; Salichs et al., 2009).

Modulation of ORF translation via MPXV LCR-like repeats also has been described for

various microbes. For instance, the functionality of baker's yeast proteins Flo1p and Flo11p

is proportionally modulated by the repeat length of their N-terminal regions. A relatively

small change in the number of tandem repeats is crucial for yeast adaptation to a new

environment (Fidalgo et al., 2006; Verstrepen et al., 2005). Another example is glutamic

acid-rich protein (GARP) of plasmodia, which contains repetitive sequences that direct the

protein to the periphery of the infected erythrocyte. At least nine other exported plasmodium

proteins target the periphery of the erythrocyte using this strategy. Interestingly, the lengths

of the tandem repeats vary among plasmodium strains (Davies et al., 2017; Davies et al.,

2016).

Protein translation rates are in part regulated by the availability of mRNA codon-

cognate aminoacylated tRNAs. Homopolymers composed of codons for rare tRNAs directly

diminish translation via tRNA depletion. The non-optimal tyrosyl codons in MPXV LCRs 3

and 21 suggest such translation modulation for OPG208 and OPG204, respectively. The

protein encoded by OPG208, B19R (Cop-K2L, SPI-1) is a serine protease inhibitor-like

protein that functions as an inhibitor of apoptosis in VACV-infected cells (Kettle et al., 1995;

Kotwal and Moss, 1989), which could prevent VACV proliferation and protect nearby cells

(Brooks et al., 1995; Jorgensen et al., 2017). Consequently, MPXV B19R is now considered a

potential MPXV virulence marker (Chen et al., 2005). The protein encoded by OPG204,

B16R, is a secreted decoy receptor for interferon type I (Colamonici et al., 1995; Hernáez et

al., 2018). We did not observe any repeat number changes OPG204-associated LCR21, but

SNPs in clade I, subclade IIa, and subclade IIb result in alternative translational start sites

followed by a, suggesting these SNPs could have direct effects on OPG204 translation.

The most intriguing finding from our dataset involves LCR7 in OPG153. The

OPG153 expression product (A26L) attaches orthopoxvirus particles to laminin (Chiu et al.,

2007) and regulates orthopoxvirus particle egress (Howard et al., 2008; Kastenmayer et al.,

2014; Liu et al., 2014; Ulaeto et al., 1996), thereby modulating key steps in the virus

lifecycle. OPG153 is unique as it is the central conserved region gene that has been "lost" the

most times during orthopoxvirus evolution (Senkevich et al., 2021). Inactivation of OPG153

genes by frameshift mutations occurs rapidly in experimental orthopoxvirus evolution models

(Senkevich *et al.*, 2020), resulting increased virus replication levels, changes in particle

morphogenesis, decreased particle-to-PFU ratios, and differences in pathogenesis

(Kastenmayer *et al.*, 2014; Senkevich *et al.*, 2020). Finally, A26L is the main target of the

host antibody response to orthopoxvirus infection (Keasey et al., 2010; Pugh et al., 2016).

Thus, any genomic change that modulates OPG153 is likely of significance. LCR7 encodes a

poly-D non-structured region that is conserved among orthopoxviruses A26Ls; however, its

length is highly variable. In mammals, poly-D stretches appear to provide functionality to

asporin, a small leucin rich repeat proteoglycan that also possesses a unique stretch of

aspartyls at its N terminus (Henry et al., 2001), associated with calcium-binding (Zhu et al.,

2018). Interestingly, orthopoxviruses that form a dense protein matrix within the cytoplasm

called A-type inclusions (ATIs), such as MPXV, generically have very long poly-D stretches

in this region, whereas orthopoxviruses that do not form ATIs, such as VACV and VARV,

have reduced their LCR7 poly-D stretches to four Ds. Even among MPXV clades, patterns

are observable. Subclade IIa viruses have an extended 21 amino-acid residue poly-D stretch,

whereas clade I and subclade IIb viruses have poly-D stretches with two inserted isoleucyls.

Intriguingly, both insertions result from the incorporation of the same "ATCATA" nucleotide

insertion in the "GAT" repetitive stretch.

In summary, our findings expand the concept of genome accordions as a simple and

recurrent mechanism of adaptation on a genomic scale in orthopoxvirus evolution. A

consequence of this broadening is the recognition that MPXV genome LCRs might hold the

key to improved understanding of current monkeypox epidemiology and clinical

presentation. A new standardized approach to generate and analyze sequencing data via

prioritizing LCR characterization and subsequent functional mechanism-of-action studies are

warranted.

MATERIALS AND METHODS

Study population

This study includes confirmed human monkeypox cases diagnosed from May 18 to July 14, 2022, at the Centro Nacional de Microbiología (CNM), Instituto de Salud Carlos III, Madrid, Spain. The study was performed as part of the public health response to the current monkeypox epidemic by the Spanish Ministry of Health. Sample information is listed in **Tables S1** and **S5**.

The samples used in this work were obtained in the context of the Microbiological Surveillance and Diagnosis Program for the Monkeypox Outbreak of the Centro Nacional de Microbiología, Instituto de Salud Carlos III. The study was based on routine testing, did not involve any additional sampling or tests and stored RNA extracts were used, so specific ethical approval was not required for this study. All sequenced viruses corresponded to those to patients that gave consent to be analyzed for diagnosis or surveillance purposes.

Study sample processing

Swabs of vesicular lesions from study patients in viral transport media were sent refrigerated to CNM. Nucleic acids were extracted at CNM using either QIAamp MinElute Virus Spin (DNA) or QIAamp Viral RNA Mini kits (Qiagen, Germantown, MD, USA) according to the manufacturer's recommendations. Inactivation of samples was conducted in a certified class II biological safety cabinet in a biosafety level (BSL) 2 laboratory using BSL-3 best practices with appropriate personal protective equipment.

Monkeypox virus (MPXV) laboratory confirmation

MPXV detection by PCR in a sample was considered laboratory confirmation and resulted in inclusion of the swab in the study. A previously described orthopoxvirus-generic real-time

PCR (qPCR) was used for screening (Fedele et al., 2006). A previously described

conventional validated nested PCR targeting OPG002 (encoding a TFN receptor) was used

for results confirmation (Sánchez-Seco et al., 2006).

MPXV genome sequencing

Sequencing libraries were prepared with a tagmentation-based Illumina DNA Prep kit

(Illumina, San Diego, CA, USA) and run in a NovaSeq 6000 SP Reagent Kit (Illumina) flow

cell using 2x150 paired-end sequencing. To improve assembly quality, the library from swab

353R, an unpassaged vesicular fluid from a confirmed case, was also run in a MiSeq Reagent

Kit v3 (Illumina) flow cell using 2 x 300 paired-end sequencing. Additionally, sample 353R

was also analyzed by single-molecule methods using Nanopore sequencing (Oxford

Nanopore Technologies, Oxford, UK). For Nanopore sequencing, 210 ng of DNA was

extracted from swab 353R and used to prepare a sequence library with a Rapid Sequencing

Kit (Oxford Nanopore Technologies); the library was analyzed in an FLO-MIN106D (Oxford

Nanopore Technologies) flow cell for 25 h. The process rendered 1.12 Gb of filter-passed

bases.

Bioinformatics

De novo **assembly and annotation of subclade II lineage B.1 MPXV genome sequence**

353R. Due to the high yield of MPXV genomic material in a preparatory run, swab 353R was

selected as source material for the determination of an MPXV high-quality genome (HQG)

sequence. Single-molecule long-sequencing reads were preprocessed using Porechop

v0.3.2pre (Wick et al., 2017) with default parameters. Reads were *de novo* assembled using

Flye v2.9-b1768 (Kolmogorov et al., 2019) in single-molecule sequencing raw read mode

with default parameters, resulting in one MPXV contig of 198,254 bp. Short 2x150

sequencing reads were mapped with Bowtie2 v2.4.4 (Langmead and Salzberg, 2012) against

the selected contig, and resulting BAM files were used to correct the assembly using Pilon

v1.24 (Walker et al., 2014). At this intermediate step, this corrected sequence was used as a

reference in the nf-core/viralrecon v2.4.1 pipeline (Patel et al., 2022) for mapping and

consensus generation with short sequencing reads. The allele frequency threshold of 0.5 was

used for including variant positions in the corrected contig.

Short MiSeq 2x300 and NovaSeq 2x150 sequencing reads were also assembled *de*

novo using the nf-core/viralrecon v2.4.1 pipeline, written in Nextflow (Di Tommaso et al.,

2017) in collaboration between the nf-core community (Ewels et al., 2020) and the Unidad de

Bioinformática, Instituto de Salud Carlos III, Madrid, Spain (https://github.com/BU-ISCIII).

FASTQ files containing raw reads were quality controlled using FASTQC v0.11.9 (Andrews,

2010). Raw reads were trimmed using fastp v0.23.2 (Chen et al., 2018). The sliding-window

quality-filtering approach was performed, scanning the read with a 4-base-wide sliding

window and cutting 3' and 5' base ends when average quality per base dropped below a

Qphred33 of 20. Reads shorter than 50 bp and reads with more than 10% read quality under

Qphred 20 were removed. Host genome reads were removed via kmer-based mapping of the

trimmed reads against the human genome reference sequence GRCh38

(https://www.ncbi.nlm.nih.gov/data-hub/genome/GCF_000001405.26/) using Kraken 2

v2.1.2 (Wood et al., 2019). The remaining non-host reads were assembled using SPADES

v3.15.3 (Antipov et al., 2016; Prjibelski et al., 2020) in rnaviral mode. A fully ordered MPXV

genome sequence was generated using ABACAS v1.3.1 (Assefa et al., 2009), based on the

MPXV isolate MPXV_USA_2022_MA001 (Nextstrain subclade IIb lineage B.1) sequence

(GenBank #ON563414.3) (Gigante et al., 2022). The independently obtained *de novo*

assemblies and reference-based consensus genomes obtained from swab 353R were aligned

using MAFFT v7.475 ([Katoh et al., 2019](#)) and visually inspected for variation using Jalview

v2.11.0 ([Waterhouse et al., 2009](#)).

Systematic identification of low-complexity regions in orthopoxvirus genomes. Detection of

short tandem repeats (STRs) in the HQG sequence and other orthopoxvirus genomes was

performed with Tandem repeats finder ([Benson, 1999](#)), using default parameters. Briefly, the

algorithm works without the need to specify either the pattern or its length. Tandem repeats

are identified considering percent identity and frequency of insertion (ins) or deletion (del) of

bases (indels) between adjacent pattern copies, using statistically based recognition criteria.

Since Tandem repeats finder does not detect single-nucleotide repeats, we developed an R

script to systematically identify homopolymers of at least 9 nucleotide residues in all

available orthopoxvirus genome sequences. STRs and homopolymers were annotated as low-

complexity regions (LCRs).

Curation of low-complexity regions in the MPXV high-quality virus genome sequence. We

curated LCRs in the HQG sequence using a modified version of STRsearch ([Wang et al.,](#)

[2020](#)). Once provided with identifying flanking regions, STRsearch performed a profile

analysis of STRs in massively parallel sequencing data. To ensure high-quality

characterization of LCR alleles, we modified the script ([https://github.com/BU-](https://github.com/BU-ISCIII/MPXstreveal)

[ISCIII/MPXstreveal](https://github.com/BU-ISCIII/MPXstreveal)) to complement reverse reads that map against the reverse genome

strand according to their BAM flags. In addition, output was modified to add information

later accessed by a custom Python script to select only reads containing both LCR flanking

regions. All LCRs in the HQG sequence were manually validated using STRsearch results

and *de novo* assemblies obtained from all sequencing approaches. When an LCR was only

resolved by single-molecule long-sequencing technologies (LCR pair 1/4 and LCR3), we also

analyzed publicly available data by downloading all single-molecule long-sequencing data

from the National Center for Biotechnology Information (NCBI) Sequence Read Archive

(SRA) (https://www.ncbi.nlm.nih.gov/sra) as of August 10, 2022, and analyzed the data

according to **File S1**.

Final MPXV high-quality virus genome sequence assembly. The consensus genome

constructed with the nf-core/viralrecon v2.4.1 pipeline using the corrected *de novo* contig as

stated above, along with the resulting curated and validated consensus LCRs, were used to

build the final HQG reference sequence using a custom Python script. The resulting HQG is

available from the European Nucleotide Archive (#OX044336.2).

Generation of MPXV high-quality virus genome reference-based consensus sequence for

all other samples. For the remaining specimens, sequencing reads were analyzed for MPXV

genome sequence determination using the nf-core/viralrecon v2.4.1 pipeline. Trimmed reads

were mapped with Bowtie2 v2.4.4 against the HQG sequence and the sequence of subclade II

lineage A MPXV isolate M5312_HM12_Rivers (GenBank #MT903340.1) (Mauldin et al.,

2022). Picard v2.26.10 (The Broad Institute, 2018) and SAMtools v1.14 (Li et al., 2009) were

used to generate MPXV genome mapping statistics. iVar v1.3.1 (Grubaugh et al., 2019),

which calls for low-frequency and high-frequency variants, was used for variant calling.

Variants with an allele frequency higher than 75% were kept to be included in the consensus

genome sequence. BCFtools v1.14 (Danecek et al., 2021) was used to obtain the MPXV

genome sequence consensus with filtered variants and masked genomic regions with

coverage values lower than 10X. All variants, included or not, in the consensus genome

sequence, were annotated using SnpEff v5.0e (Cingolani et al., 2012b), and SnpSift v4.3

(Cingolani et al., 2012a). Final summary reports were created using MultiQC v.1.11 (Ewels

et al., 2016). Consensus genome sequences were analyzed with Nextclade v2.4.1

(Aksamentov et al., 2021) using the "MPXV (All clades)" dataset (timestamp 2022-08-

19T12:00:00Z). Raw reads and consensus genomes are available from the European

Nucleotide Archive (#ERS12168855–ERS12168865, #ERS12168867, #ERS12168868,

#ERS13490510–ERS13490543).

Intra-host and inter-host allele frequency analyses. Intra-host genetic entropy (defined as -

sum(Xi*log(Xi)), in which Xi denotes each of the allele frequencies in a position) was

calculated according to the single-nucleotide polymorphisms (SNP) frequencies of each

position along the genome using nf-core/viralrecon v2.4.1 pipeline results. Similarly, genetic

entropy for each LCR was calculated considering the frequencies of repeat lengths.

LCR intra-host and inter-host variations in the sample set were analyzed using the

modified version of STRsearch. As a filter for quality for this analysis, STRsearch results

(**Table S5**) were filtered, keeping alleles with at least 10 reads spanning the region and allele

frequency above 0.03. Quality control and allele frequency graphs were created using a

customized R script.

Pairwise genetic distances between samples were calculated as Euclidean distances

(defined as /X-Y/=sqrt(sum(xi-yi)^2), in which xi and yi are the allele frequencies of sample

X and Y at a given position, respectively), thus accounting for the major and minor alleles at

each analyzed position. Distances were calculated individually for each variable LCR (STRs

2, 5, 7, 10, 11, and 21) and for each of all 5,422 SNPs showing inter-sample variability

(compared to MPXV-M5312_HM12_Rivers). The distributions of inter-sample distances

were compared between LCRs using a Kruskal–Wallis test (χ^2 *p*-values) followed by multiple

pairwise-comparison between groups (Wilcoxon test), with *p*-values subjected to the false

discovery rate (FDR) correction. A randomization test was used to test whether inter-sample

variability in LCRs is higher than that in SNPs: first, the average Euclidean distance for each

LCR and each SNP position was calculated; then, the average value of each LCR was

compared to a random sample of 1,000 values from the distribution of mean distances from

the SNPs along the genome. The *p*-value was calculated from the percentage of times that the

mean of the LCR was higher than the randomly taken values from the SNPs.

Phylogenetic analysis of the MPXV central conserved region. Variant calling and SNP

matrix generation was performed using Snippy v4.4.5 (Seeman, 2015), including sequence

samples and representative MPXV genome sequences downloaded from GenBank (**Table**

S5). The SNP matrix with both invariant and variant sites was used for phylogenetic analysis

using IQ-Tree 2 v. 2.1.4-beta (Minh et al., 2020) via predicted model K3Pu+F+I and 1,000

bootstrap replicates. A phylogenetic tree was visualized and annotated using iTOL v6.5.8

(Letunic and Bork, 2021). The SNP matrix was also used for generating the haplotype

network using PopArt v1.7 (Bandelt et al., 1999).

Selected MPXV ORF analysis. Representative orthopoxvirus genomes (Senkevich *et al.*,

2021) were downloaded from GenBank together with the consensus genome sequences from

the specimens analyzed in this study (**Table S5**). MPXV genomes were assigned to clades

and lineages according to the most recent nomenclature recommendations according to

Nextstrain (Nextstrain, 2022) using Nextclade v2.4.1. Annotations from RefSeq

#NC_063383.1 (subclade II lineage A MPXV virus isolate MPXV-M5312_HM12_Rivers)

GFF file were transferred to all FASTA genome sequences using Liftoff v1.6.3 (Shumate and

Salzberg, 2021). OPG153 was extracted using AGAT v0.9.1 (Dainat et al., 2022) and multi-

FASTA files were generated for each group and gene. OPG204 and OPG208 alternative

annotation start site ORFs were re-annotated in Geneious Prime (Biomatters, San Diego, CA,

USA), and extracted as new alignments. We used MUSCLE v3.8.1551 for aligning each

multi-FASTA file and Jalview v2.11.0 for inspecting and editing the alignments. Finally,

MetaLogo v1.1.2 (Chen et al., 2022) was used for creating and aligning the sequence logos

for each orthopoxvirus group of the OPG153/LCR7, OPG204/LCR21, and OPG208/LCR3

areas.

Comparison of LCR frequencies in protein functional groups. The potential biological

impact of LCRs was evaluated by mapping the frequency and location of STRs and

homopolymers in the orthopoxvirus genome and considering the biological function of the

affected genes. The frequency of inclusion of LCRs between distinct functional groups of

genes was compared as previously described (Senkevich *et al.*, 2021). Orthopoxviruses

(n=231, Akhmeta virus [AKMV]: n=6 sequences; alaskapox virus [AKPV]: n=1; cowpox

virus [CPXV]: n=82; ectromelia virus [ECTV]: n=5; MPXV: n=62; VACV: n=18; VARV:

n=57) include 216 functionally annotated OPGs classified in 5 categories ("Housekeeping

genes/Core" ANK/PRANC family, Bcl-2 domain family, PIE family, and "Accessory/Other"

[e.g., virus-host interacting genes]). The frequency was calculated after normalizing count

numbers with the sample size of the OPG alignment. Statistical analysis of the significance of

differences was performed by applying a Kruskal–Wallis test (χ^2 *p*-values) followed by a

non-parametric multiple pairwise comparison between groups (Wilcoxon test), with *p*-values

subjected to FDR correction.

ACKNOWLEDGMENTS

We would like to thank the work of the Rapid Response Unit of the National Center for

Microbiology, especially MªJosé Buitrago, and Cristobal Belda, ISCIII General Director. We

also thank Anya Crane (Integrated Research Facility at Fort Detrick, National Institute of

Allergy and Infectious Diseases, National Institutes of Health) for critically editing the

manuscript and Jiro Wada (Integrated Research Facility at Fort Detrick, National Institute of

Allergy and Infectious Diseases, National Institutes of Health) for helping with figure

preparation.

The work for this study at Instituto de Salud Carlos III was partially funded by Acción

Estratégica "Impacto clínico y microbiológico del brote por el virus de la viruela del mono en

pacientes en España (2022): proyecto multicéntrico MONKPOX-ESP22" (CIBERINFEC).

The work for this study at the Department of Microbiology, Icahn School of Medicine at

Mount Sinai as part of Global Health Emerging Pathogen Institute activities was funded by

institutional funds (G.P.) from the Department of Microbiology, Icahn School of Medicine at

Mount Sinai in support of Global Health Emerging Pathogen Institute activities.

This work was also supported in part through Laulima Government Solutions, LLC,

prime contract with the U.S. National Institute of Allergy and Infectious Diseases (NIAID)

under Contract No. HHSN272201800013C. J.H.K. performed this work as an employee of

Tunnell Government Services (TGS), a subcontractor of Laulima Government Solutions,

LLC, under Contract No. HHSN272201800013C.

Opinions, interpretations, conclusions, and recommendations are those of the authors

and are not necessarily endorsed by the U.S. Army.

The views and conclusions contained in this document are those of the authors and

should not be interpreted as necessarily representing the official policies, either expressed or

implied, of the U.S. Department of Health and Human Services or of the institutions and

companies affiliated with the authors, nor does mention of trade names, commercial products,

or organizations imply endorsement by the U.S. Government.

AUTHOR CONTRIBUTIONS

Conceptualization, S. M., S.V., A. N., I. J., M.S.L., A.G.S., I.C., M.S.S., G.P.

Methodology, S.M., S.V., A.N., J.A.P.G., S.V.F., A.Z., J. H. K., M.S.L., N.D., J.R.K., E.G.,

S.G., G.P.

Investigation, S.M., S.V., A.N., J.A.P.G., S.V.F., A.Z., E.O., O.A., A.M.G., A.D.I., V.E.,

C.G., F.M., P.S., M.T., A.V., J.C.G., I.T., M.C.R., L.M., M.L., A.G., L.C., A.G., J.C., L.H.,

P.J.S., M.L.N.R., I.J., M.E.A.A., C.L., L.R., I.E., M.S., M.A.M., J.H.K., M.S.L., N.D.P.,

J.R.K., E.G., S.G., A.G.S., I.C., M.S.S., G.P.

Formal analysis, S.M., S.V., A.N., J.A.P.G., S.V.F., J.H.K., M.S.L., N.D.P., J.R.K., E.G.,

I.C., M.S.S, G.P.

Writing – original draft, S.M., S.V., A.N., G.P.

Writing – review & editing, S. M., S.V., A. N., A.G.S., I.C., M.S.S., G.P.

Visualization, S.M., S.V., A.N., G.P.

Supervision, A.G.S., I.C., M.S.S., G.P.

Resources, A.G.S., I.C., M.S.S., G.P.

Funding Acquisition, A.G.S., I.C., M.S.S., G.P.

DECLARATION OF INTERESTS

A.G.-S. has consulting agreements for the following companies involving cash and/or stock:

Castlevax, Amovir, Vivaldi Biosciences, Contrafect, 7Hills Pharma, Avimex, Vaxalto,

Pagoda, Accurius, Esperovax, Farmak, Applied Biological Laboratories, Pharmamar, Paratus,

CureLab Oncology, CureLab Veterinary, Synairgen, and Pfizer, outside of the reported work.

A.G.-S. has been an invited speaker in meeting events organized by Seqirus, Janssen, Abbott,

and Astrazeneca. A.G.-S. is inventor on patents and patent applications on the use of

antivirals and vaccines for the treatment and prevention of virus infections and cancer, owned

by the Icahn School of Medicine at Mount Sinai, New York, outside of the reported work.

The authors declare no competing interests.

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Gustavo Palacios (gustavo.palacios@mssm.edu).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

All scripts and codes used for this study can be found at github repository:

(https://github.com/BU-ISCIII/MPXstreveal).

 FIGURE LEGENDS

Figure 1. *De novo* **assembly of subclade II lineage B.1 monkeypox virus (MPXV)**

genome sequence 353R

A visual representation of the fully annotated MPXV isolate 353R genome (based on the

subclade II lineage A reference isolate MPXV-M5312_HM12_Rivers genome sequence

annotation). Shown are (from the outside to the inside): high-quality genome (HQG) hybrid

assembly (wide outer light blue ring); sequencing coverage distribution graph (thin ragged

line [green: $\geq$10,000x, 99.42%; black: 1,000x–10,000, 0.28%; orange: <1,000x-10, 0.1%;

red: <10-0, 0.2%]); orthologous poxvirus gene (OPG) annotations according to the

standardized nomenclature (Senkevich *et al.*, 2021) (lettering and shaded boxes [orange:

ANK/PRANC (N-terminal ankyrin protein with PRANC domain) inverted terminal repetition

[ITR] regions; gold: Bcl-2 domain; blue: BTB/Kelch domains; green: housekeeping; purple:

other; pink: TNFR and/or PIE domains); contigs from NovaSeq, MiSeq, and Nanopore

sequencing (wide inner gray rings). Additionally, radial lines and shading that originate in the

center and reach outward on the white background indicate low-complexity regions (LCRs;

royal blue) and areas with more change (light blue).

Figure 2. Characterization and validation of non-randomly distributed low-complexity

regions (LCRs) in monkeypox virus (MPXV) genome sequence 353R

(A) LCR3 sequence validation using MPXV 353R Nanopore sequencing data and 15

additional raw data sequencing reads downloaded from the National Center for

Biotechnology Information (NCBI) Sequence Read Archive (SRA). (B) LCR pair 1/4

sequence validation using MPXV 353R Nanopore sequencing data and 20 additional raw data

sequencing reads downloaded from NCBI SRA. Detailed information on the represented

materials, along with originator and epidemiological data, is provided in **Table S6**.

Figure 3. Low-complexity regions (LCRs) are non-randomly distributed in the

monkeypox virus (MPXV) genome

(A) Frequencies (mean +/- standard error) at which LCRs occur in orthologous poxvirus

genes (OPGs) of different functional groups. Shown are functional classes in which pairwise

comparisons had a significantly different frequency than other groups (multiple pairwise

Wilcoxon test, false discovery rate [FDR] corrected p-value <0.05). (B) Entropy value

distribution for short tandem repeats (STRs) in LCRs (left) and single-nucleotide

polymorphisms (SNPs; right). (C) Distributions of the pairwise inter-sample Euclidean

distances for each STR in LCRs (**Table S3**). SNPs in the boxplot represent the distribution of

average Euclidean distances of each variable position along the MPXV genome.

Figure 4. Low-complexity regions (LCRs) might be more phylogenetically informative

than single-nucleotide polymorphisms (SNPs) for inter-host sequence analysis

Monkeypox virus (MPXV) population genomics within the biological specimen (intra-host)

and across different specimens (inter-host). (A) Panel shading indicates the number of reads

supporting each LCR for each sample. Only paired reads that include a perfect match to both

flanking regions were counted; the gradient shows the maximum value in black and the

minimum value (n=1) in the lightest blue. Samples without coverage are indicated in gray.

(B) Comparison of LCR allele frequency for samples 353R and 349R. Only LCRs with at

least 10 supporting paired reads including both flanking regions were counted; only alleles

with a frequency of 0.03 or higher were considered. The gradient shows the maximum value

in black and the minimum value (n=0.03) in the lightest blue. (C) Comparison of LCR allele

frequency in all samples for LCRs with good coverage (7, pair 10/11, 12, 13, 14, 19, 20, and

21). Only LCRs with at least 10 supporting paired reads including both flanking regions were

counted; only alleles with a frequency of 0.03 or higher were considered. The gradient shows

the maximum value in black and the minimum value (n=0.03) in the lightest blue.

Figure 5. Conservation and variation in proteins encoded by orthologous poxvirus gene

(OPG) and codon usage analysis in OPG low-complexity regions (LCRs)

MetaLogo visualization of conserved and varying amino-acid residues in OPG-encoded

proteins among monkeypox virus (MPXV) clade I, subclade IIa, and subclade IIb, with

homologous and nonhomologous sites highlighted. (A) OPG153/ LCR7-derived variability.

CMLV, camelpox virus; VACV, vaccinia virus; VARV, variola virus; CPXV, cowpox virus;

MPXV, monkeypox virus. (B) OPG204/LCR21-derived variability; and (C) OPG208/LCR3-

derived variability. Visualizations created at Biorender.com and Geneious version 2022.2

created by Biomatters.

 TABLE LEGENDS

 Table 1. Low-complexity regions (LCRs) in monkeypox virus (MPXV) genome sequence 353R

Short tandem repeats (STRs) are described using nucleotide base-pair coordinates with reference to

the high-quality genome (HQG) sequence (ENA Accession #OX044336). Listed are the number of

repeat units, description of the sequence (with $_n$ = number of repeats for this particular genome),

identification of the nearest annotated orthologous poxvirus gene (OPG), type of LCR (STR or

homopolymer), position of the LCR to the nearest gene, and distance of the LCR to the nearest gene.

OPG notations follow the standardized nomenclature (Senkevich *et al.*, 2021); vaccinia virus

(VACV) Copenhagen strain and classical VACV gene notations are shown in addition to enable

comparisons.

Table 2. Low-complexity region (LCR) validation and entropy-level intra-host analysis in

monkeypox virus (MPXV) genome sequence 353R

Listed are the type and number of supporting reads for each LCR. Definitions of quality: Yes, LCR is

found entirely in the assembly in one contig; no, LCR is not assembled with the reported method. All

LCRs with entropy levels above 0.15 are shaded in gray. OPG, orthologous poxvirus gene.

Table 3. Comparison of monkeypox virus (MPXV) genome sequence 353R with reference

sequences

LCR repetitions for each genome are indicated. Discrepant number (n) of LCR repeats are shaded in

gray. indels, number of insertion (ins) or deletion (del) of bases; LCR, low-complexity region; SNP,

single-nucleotide polymorphism.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. Examples for read mapping artefacts and correction in monkeypox virus (MPXV)

genome low-complexity regions (LCRs)

(A) LCR2 alignment highlighting differences compared with various consensus sequences. (B) LCR7

alignment highlighting differences of results obtained using three sequencing platforms compared to

the subclade II lineage A monkeypox reference isolate MPXV-M5312_HM12_Rivers sequence.

Figure S2. Phylogenetic analysis of monkeypox virus (MPXV)

(A) Phylogenetic maximum-likelihood (ML) tree showing monkeypox virus (MPXV) subclade IIb

single-nucleotide polymorphism (SNP) clustering. Bootstrap supports >60 are indicated by labels

with their number of supports. (B) Haplotype network showing SNP differences among samples

included in the phylogenetic tree. Details on groups can be found in **Table S4**.

Figure S3. Conservation and variation in proteins encoded by orthologous poxvirus gene

(OPG) 208. (A) MetaLogo visualization of conserved and varying amino-acid residues in OPG-

encoded proteins among monkeypox virus (MPXV) clade I, subclade IIa, and subclade IIb, with

homologous and nonhomologous sites highlighted; (B) Entropy heatmap; (C) Entropy analysis by

site.

Figure S4. Conservation and variation in proteins encoded by orthologous poxvirus gene

(OPG) 153.

(A) Entropy heatmap. CMLV, camelpox virus; VARV, variola virus; VACV, vaccinia virus; CPXV,

cowpox virus; MPXV, monkeypox virus. (B) . MPXV, monkeypox virus; CPXV, cowpox virus;

VACV, vaccinia virus; VARV, variola virus; CMLV, camelpox virus.

SUPPLEMENTAL TABLE LEGENDS

Table S1. NovaSeq sequencing quality control values

Table S2. Low-complexity regions (LCRs) in the monkeypox virus (MPXV) high-quality

genome (HQG) sequence 353R

Listed are annotated positions (according to the reference MPXV-M5312_HM12_Rivers isolate

genome sequence), sequence, and flanking regions of each area, as described previously (Phillips et

al., 2018). ID, identification; STR, short tandem repeats.

Table S3. Short tandem repeats (STRs) in low-complexity regions (LCRs)

Detailed information on STRs of the entire dataset used for analysis in **Figure 2**.

Table S4. Phylogenetic analysis

Detailed information on groups depicted in **Figure S2**. SNP, single nucleotide polymorphism.

Table S5. Genomes used in the study

GI, genome identifier; NA, not applicable

Table S6. Characterization and validation of non-randomly distributed low-complexity regions

(LCRs) in the monkeypox virus (MPXV) 353R genome sequence

Detailed information on the represented materials, along with originator and epidemiological data,

used for analysis in **Figure 2**. SRA, Sequence Read Archive (SRA); ID, identification; QC, quality

control.

SUPPLEMENTAL FILE LEGENDS

File S1. Analysis parameters for validation of short tandem repeats (STRs) in National Center

for Biotechnology Information (NCBI) Sequence Read Archive (SRA)

REFERENCES

Aksamentov, I., Roemer, C., Hodcroft, E.B., and Neher, R.A. (2021). Nextclade: clade assignment,

mutation calling and quality control for viral genomes. J Open Source Softw *6*, 3773.

10.21105/joss.03773.

Andrews, S. (2010). FASTQC: a quality control tool for high throughput sequence data.

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Antinori, A., Mazzotta, V., Vita, S., Carletti, F., Tacconi, D., Lapini, L.E., D'Abramo, A., Cicalini,

S., Lapa, D., Pittalis, S., et al. (2022). Epidemiological, clinical and virological characteristics of four

cases of monkeypox support transmission through sexual contact, Italy, May 2022. Euro Surveill *27*,

2200421. 10.2807/1560-7917.ES.2022.27.22.2200421.

Antipov, D., Korobeynikov, A., McLean, J.S., and Pevzner, P.A. (2016). hybridSPAdes: an

algorithm for hybrid assembly of short and long reads. Bioinformatics *32*, 1009-1015.

10.1093/bioinformatics/btv688.

Assefa, S., Keane, T.M., Otto, T.D., Newbold, C., and Berriman, M. (2009). ABACAS: algorithm-

based automatic contiguation of assembled sequences. Bioinformatics *25*, 1968-1969.

10.1093/bioinformatics/btp347.

Bandelt, H.-J., Forster, P., and Röhl, A. (1999). Median-joining networks for inferring intraspecific

phylogenies. Mol Biol Evol *16*, 37-48. 10.1093/oxfordjournals.molbev.a026036.

Baroudy, B.M., and Moss, B. (1982). Sequence homologies of diverse length tandem repetitions near

ends of vaccinia virus genome suggest unequal crossing over. Nucleic Acids Res *10*, 5673-5679.

10.1093/nar/10.18.5673.

Beer, E.M., and Rao, V.B. (2019). A systematic review of the epidemiology of human monkeypox

outbreaks and implications for outbreak strategy. PLoS Negl Trop Dis *13*, e0007791.

10.1371/journal.pntd.0007791.

Benson, G. (1999). Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res

27, 573-580. 10.1093/nar/27.2.573.

Bratke, K.A., and McLysaght, A. (2008). Identification of multiple independent horizontal gene

transfers into poxviruses using a comparative genomics approach. BMC Evol Biol *8*, 67.

10.1186/1471-2148-8-67.

Brooks, M.A., Ali, A.N., Turner, P.C., and Moyer, R.W. (1995). A rabbitpox virus serpin gene

controls host range by inhibiting apoptosis in restrictive cells. J Virol *69*, 7688-7698.

10.1128/JVI.69.12.7688-7698.1995.

Bunge, E.M., Hoet, B., Chen, L., Lienert, F., Weidenthaler, H., Baer, L.R., and Steffen, R. (2022).

The changing epidemiology of human monkeypox—A potential threat? A systematic review. PLoS

Negl Trop Dis *16*, e0010141. 10.1371/journal.pntd.0010141.

Centers for Disease Control and Prevention (2022). 2022 Monkeypox Outbreak Global Map.

https://www.cdc.gov/poxvirus/monkeypox/response/2022/world-map.html.

Chen, N., Li, G., Liszewski, M.K., Atkinson, J.P., Jahrling, P.B., Feng, Z., Schriewer, J., Buck, C.,

Wang, C., Lefkowitz, E.J., et al. (2005). Virulence differences between monkeypox virus isolates

from West Africa and the Congo basin. Virology *340*, 46-63. 10.1016/j.virol.2005.05.030.

Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics *34*, i884-i890. 10.1093/bioinformatics/bty560.

Chen, Y., He, Z., Men, Y., Dong, G., Hu, S., and Ying, X. (2022). MetaLogo: a heterogeneity-aware

sequence logo generator and aligner. Brief Bioinform *23*, 1-7. 10.1093/bib/bbab591.

Chiu, W.-L., Lin, C.-L., Yang, M.-H., Tzou, D.-L., and Chang, W. (2007). Vaccinia virus 4c (A26L)

protein on intracellular mature virus binds to the extracellular cellular matrix laminin. J Virol *81*,

2149-2157. 10.1128/JVI.02302-06.

Cingolani, P., Patel, V.M., Coon, M., Nguyen, T., Land, S.J., Ruden, D.M., and Lu, X. (2012a).

Using *Drosophila melanogaster* as a model for genotoxic chemical mutational studies with a new

program, SnpSift. Front Genet *3*, 35. 10.3389/fgene.2012.00035.

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X., and

Ruden, D.M. (2012b). A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w^{1118}; *iso*-2; *iso*-3.

Fly (Austin) *6*, 80-92. 10.4161/fly.19695.

Cohen-Gihon, I., Israeli, O., Shifman, O., Erez, N., Melamed, S., Paran, N., Beth-Din, A., and Zvi, A.

(2020). Identification and whole-genome sequencing of a monkeypox virus strain isolated in Israel.

Microbiol Resour Announc *9*, e01524-01519. 10.1128/MRA.01524-19.

Colamonici, O.R., Domanski, P., Sweitzer, S.M., Larner, A., and Buller, R.M. (1995). Vaccinia virus

B18R gene encodes a type I interferon-binding protein that blocks interferon α transmembrane

signaling. J Biol Chem *270*, 15974-15978. 10.1074/jbc.270.27.15974.

Dainat, J., Hereñú, D., LucileSol, and pascal-git (2022). NBISweden/AGAT: AGAT-v0.8.1.

https://zenodo.org/record/5834795#.YzJOCzTMJyE. 10.5281/zenodo.5834795.

Damon, I.K. (2011). Status of human monkeypox: clinical disease, epidemiology and research.

Vaccine *29 Suppl 4*, D54-59. 10.1016/j.vaccine.2011.04.014.

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane,

T., McCarthy, S.A., Davies, R.M., and Li, H. (2021). Twelve years of SAMtools and BCFtools.

GigaScience *10*, giab008. 10.1093/gigascience/giab008.

Davies, H.M., Nofal, S.D., McLaughlin, E.J., and Osborne, A.R. (2017). Repetitive sequences in

malaria parasite proteins. FEMS Microbiol Rev *41*, 923-940. 10.1093/femsre/fux046.

Davies, H.M., Thalassinos, K., and Osborne, A.R. (2016). Expansion of lysine-rich repeats in

Plasmodium proteins generates novel localization sequences that target the periphery of the host

erythrocyte. J Biol Chem *291*, 26188-26207. 10.1074/jbc.M116.761213.

Di Tommaso, P., Chatzou, M., Floden, E.W., Barja, P.P., Palumbo, E., and Notredame, C. (2017).

Nextflow enables reproducible computational workflows. Nat Biotechnol *35*, 316-319.

10.1038/nbt.3820.

Elde, N.C., Child, S.J., Eickbush, M.T., Kitzman, J.O., Rogers, K.S., Shendure, J., Geballe, A.P., and

Malik, H.S. (2012). Poxviruses deploy genomic accordions to adapt rapidly against host antiviral

defenses. Cell *150*, 831-841. 10.1016/j.cell.2012.05.049.

Esposito, J.J., Sammons, S.A., Frace, A.M., Osborne, J.D., Olsen-Rasmussen, M., Zhang, M., Govil,

D., Damon, I.K., Kline, R., Laker, M., et al. (2006). Genome sequence diversity and clues to the

evolution of variola (smallpox) virus. Science *313*, 807-812. 10.1126/science.1125134.

Ewels, P., Magnusson, M., Lundin, S., and Käller, M. (2016). MultiQC: summarize analysis results

for multiple tools and samples in a single report. Bioinformatics *32*, 3047-3048.

10.1093/bioinformatics/btw354.

Ewels, P.A., Peltzer, A., Fillinger, S., Patel, H., Alneberg, J., Wilm, A., Garcia, M.U., Di Tommaso,

P., and Nahnsen, S. (2020). The nf-core framework for community-curated bioinformatics pipelines.

Nat Biotechnol *38*, 276-278. 10.1038/s41587-020-0439-x.

Faye, O., Pratt, C.B., Faye, M., Fall, G., Chitty, J.A., Diagne, M.M., Wiley, M.R., Yinka-Ogunleye,

A.F., Aruna, S., Etebu, E.N., et al. (2018). Genomic characterisation of human monkeypox virus in

Nigeria. Lancet Infect Dis *18*, 246. 10.1016/S1473-3099(18)30043-4.

Fedele, C.G., Negredo, A., Molero, F., Sánchez-Seco, M.P., and Tenorio, A. (2006). Use of internally

controlled real-time genome amplification for detection of variola virus and other orthopoxviruses

infecting humans. J Clin Microbiol *44*, 4464-4470. 10.1128/JCM.00276-06.

Fidalgo, M., Barrales, R.R., Ibeas, J.I., and Jimenez, J. (2006). Adaptive evolution by mutations in

the *FLO11* gene. Proc Natl Acad Sci U S A *103*, 11228-11233. 10.1073/pnas.0601713103.

Gatchel, J.R., and Zoghbi, H.Y. (2005). Diseases of unstable repeat expansion: mechanisms and

common principles. Nat Rev Genet *6*, 743-755. 10.1038/nrg1691.

Gemayel, R., Chavali, S., Pougach, K., Legendre, M., Zhu, B., Boeynaems, S., van der Zande, E.,

Gevaert, K., Rousseau, F., Schymkowitz, J., et al. (2015). Variable glutamine-rich repeats modulate

transcription factor activity. Mol Cell *59*, 615-627. 10.1016/j.molcel.2015.07.003.

Gigante, C.M., Korber, B., Seabolt, M.H., Wilkins, K., Davidson, W., Rao, A.K., Zhao, H., Hughes,

C.M., Minhaj, F., Waltenburg, M.A., et al. (2022). Multiple lineages of *Monkeypox virus* detected in

the United States, 2021- 2022. bioRxiv, 2022.2006.2010.495526. 10.1101/2022.06.10.495526.

Grant, R., Nguyen, L.L., and Breban, R. (2020). Modelling human-to-human transmission of

monkeypox. Bull World Health Organ *98*, 638-640. 10.2471/blt.19.242347.

Grubaugh, N.D., Gangavarapu, K., Quick, J., Matteson, N.L., De Jesus, J.G., Main, B.J., Tan, A.L.,

Paul, L.M., Brackney, D.E., Grewal, S., et al. (2019). An amplicon-based sequencing framework for

accurately measuring intrahost virus diversity using PrimalSeq and iVar. Genome Biol *20*, 8.

10.1186/s13059-018-1618-7.

Gubser, C., Hué, S., Kellam, P., and Smith, G.L. (2004). Poxvirus genomes: a phylogenetic analysis.

J Gen Virol *85*, 105-117. 10.1099/vir.0.19565-0.

Happi, C., Adetifa, I., Mbala, P., Njouom, R., Nakoune, E., Happi, A., Ndodo, N., Ayansola, O.,

Mboowa, G., Bedford, T., et al. (2022). Urgent need for a non-discriminatory and non-stigmatizing

nomenclature for monkeypox virus. PLoS Biol *20*, e3001769. 10.1371/journal.pbio.3001769.

Hendrickson, R.C., Wang, C., Hatcher, E.L., and Lefkowitz, E.J. (2010). Orthopoxvirus genome

evolution: the role of gene loss. Viruses *2*, 1933-1967. 10.3390/v2091933.

Henry, S.P., Takanosu, M., Boyd, T.C., Mayne, P.M., Eberspaecher, H., Zhou, W., de Crombrugghe,

B., Höök, M., and Mayne, R. (2001). Expression pattern and gene characterization of *Asporin*: a

newly discovered member of the leucine-rich repeat protein family. J Biol Chem *276*, 12212-12221.

10.1074/jbc.M011290200.

Hernáez, B., Alonso-Lobo, J.M., Montanuy, I., Fischer, C., Sauer, S., Sigal, L., Sevilla, N., and Alcamí, A. (2018). A virus-encoded type I interferon decoy receptor enables evasion of host immunity through cell-surface binding. Nat Commun *9*, 5440. 10.1038/s41467-018-07772-z.

Howard, A.R., Senkevich, T.G., and Moss, B. (2008). Vaccinia virus A26 and A27 proteins form a stable complex tethered to mature virions by association with the A17 transmembrane protein. J Virol *82*, 12384-12391. 10.1128/JVI.01524-08.

International Committee on Taxonomy of Viruses (2022). Current ICTV Taxonomy Release. Current ICTV Taxonomy Release. https://ictv.global/taxonomy.

Isidro, J., Borges, V., Pinto, M., Sobral, D., Santos, J.D., Nunes, A., Mixão, V., Ferreira, R., Santos, D., Duarte, S., et al. (2022). Phylogenomic characterization and signs of microevolution in the 2022 multi-country outbreak of monkeypox virus. Nat Med *28*, 1569-1572. 10.1038/s41591-022-01907-y.

Jones, T.C., Schneider, J., Mühlemann, B., Veith, T., Beheim-Schwarzbach, J., Tesch, J., Schmidt, M.L., Walper, F., Bleicker, T., Isner, C., et al. (2022). Genetic variability, including gene duplication and deletion, in early sequences from the 2022 European monkeypox outbreak. bioRxiv, 2022.2007.2023.501239. 10.1101/2022.07.23.501239.

Jorgensen, I., Rayamajhi, M., and Miao, E.A. (2017). Programmed cell death as a defence against infection. Nat Rev Immunol *17*, 151-164. 10.1038/nri.2016.147.

Kastenmayer, R.J., Maruri-Avidal, L., Americo, J.L., Earl, P.L., Weisberg, A.S., and Moss, B. (2014). Elimination of A-type inclusion formation enhances cowpox virus replication in mice: implications for orthopoxvirus evolution. Virology *452-453*, 59-66. 10.1016/j.virol.2013.12.030.

Katoh, K., Rozewicki, J., and Yamada, K.D. (2019). MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. Brief Bioinform *20*, 1160-1166. 10.1093/bib/bbx108.

Keasey, S., Pugh, C., Tikhonov, A., Chen, G., Schweitzer, B., Nalca, A., and Ulrich, R.G. (2010). Proteomic basis of the antibody response to monkeypox virus infection examined in cynomolgus

macaques and a comparison to human smallpox vaccination. PLoS One *5*, e15547.

10.1371/journal.pone.0015547.

Kettle, S., Blake, N.W., Law, K.M., and Smith, G.L. (1995). Vaccinia virus serpins B13R (SPI-2)

and B22R (SPI-1) encode M_r 38.5 and 40K, intracellular polypeptides that do not affect virus

virulence in a murine intranasal model. Virology *206*, 136-147. 10.1016/s0042-6822(95)80028-x.

Khodakevich, L., Ježek, Z., and Messinger, D. (1988). Monkeypox virus: ecology and public health

significance. Bull World Health Organ *66*, 747-752.

Kolmogorov, M., Yuan, J., Lin, Y., and Pevzner, P.A. (2019). Assembly of long, error-prone reads

using repeat graphs. Nat Biotechnol *37*, 540-546. 10.1038/s41587-019-0072-8.

Kotwal, G.J., and Moss, B. (1989). Vaccinia virus encodes two proteins that are structurally related

to members of the plasma serine protease inhibitor superfamily. J Virol *63*, 600-606.

10.1128/JVI.63.2.600-606.1989.

Kugelman, J.R., Johnston, S.C., Mulembakani, P.M., Kisalu, N., Lee, M.S., Koroleva, G., McCarthy,

S.E., Gestole, M.C., Wolfe, N.D., Fair, J.N., et al. (2014). Genomic variability of monkeypox virus

among humans, Democratic Republic of the Congo. Emerg Infect Dis *20*, 232-239.

10.3201/eid2002.130118.

Ladner, J.T., Beitzel, B., Chain, P.S.G., Davenport, M.G., Donaldson, E.F., Frieman, M., Kugelman,

J.R., Kuhn, J.H., O'Rear, J., Sabeti, P.C., et al. (2014). Standards for sequencing viral genomes in the

era of high-throughput sequencing. mBio *5*, e01360-01314. 10.1128/mBio.01360-14.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods *9*,

357-359. 10.1038/nmeth.1923.

Letunic, I., and Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic

tree display and annotation. Nucleic Acids Res *49*, W293-W296. 10.1093/nar/gkab301.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence

Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079.

10.1093/bioinformatics/btp352.

Likos, A.M., Sammons, S.A., Olson, V.A., Frace, A.M., Li, Y., Olsen-Rasmussen, M., Davidson, W.,

Galloway, R., Khristova, M.L., Reynolds, M.G., et al. (2005). A tale of two clades: monkeypox

viruses. J Gen Virol *86*, 2661-2672. 10.1099/vir.0.81215-0.

Liu, L., Cooper, T., Howley, P.M., and Hayball, J.D. (2014). From crescent to mature virion:

vaccinia virus assembly and maturation. Viruses *6*, 3787-3808. 10.3390/v6103787.

Liu, R., Mendez-Rios, J.D., Peng, C., Xiao, W., Weisberg, A.S., Wyatt, L.S., and Moss, B. (2019).

SPI-1 is a missing host-range factor required for replication of the attenuated modified vaccinia

Ankara (MVA) vaccine vector in human cells. PLoS Pathog *15*, e1007710.

10.1371/journal.ppat.1007710.

Magnus, P.v., Andersen, E.K., Petersen, K.B., and Birch-Andersen, A. (2009). A pox-like disease in

cynomolgus monkeys. Acta Pathol Microbiol Scand *46*, 156-176. 10.1111/j.1699-

0463.1959.tb00328.x.

Mauldin, M.R., McCollum, A.M., Nakazawa, Y.J., Mandra, A., Whitehouse, E.R., Davidson, W.,

Zhao, H., Gao, J., Li, Y., Doty, J., et al. (2022). Exportation of monkeypox virus from the African

continent. J Infect Dis *225*, 1367-1376. 10.1093/infdis/jiaa559.

McFadden, G. (2005). Poxvirus tropism. Nat Rev Microbiol *3*, 201-213. 10.1038/nrmicro1099.

McLysaght, A., Baldi, P.F., and Gaut, B.S. (2003). Extensive gene gain associated with adaptive

evolution of poxviruses. Proc Natl Acad Sci U S A *100*, 15655-15660. 10.1073/pnas.2136653100.

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., and

Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic inference in the

genomic era. Mol Biol Evol *37*, 1530-1534. 10.1093/molbev/msaa015.

Moss, B. (2006). Poxvirus entry and membrane fusion. Virology *344*, 48-54.

10.1016/j.virol.2005.09.037.

Moss, B. (2016). Membrane fusion during poxvirus entry. Semin Cell Dev Biol *60*, 89-96.

10.1016/j.semcdb.2016.07.015.

Moss, B., and Smith, G.L. (2021). *Poxviridae*: the viruses and their replication. In Fields virology.

Volume 2: DNA viruses P.M. Howley, D.M. Knipe, B.A. Damania, and J.I. Cohen, eds. (Wolters

Kluwer), pp. 573-613.

Nextstrain (2022). Genomic epidemiology of monkeypox virus.

https://nextstrain.org/monkeypox/hmpxv1.

Ng, O.T., Lee, V., Marimuthu, K., Vasoo, S., Chan, G., Lin, R.T.P., and Leo, Y.S. (2019). A case of

imported monkeypox in Singapore. Lancet Infect Dis *19*, 1166. 10.1016/S1473-3099(19)30537-7.

Nuzzo, J.B., Borio, L.L., and Gostin, L.O. (2022). The WHO declaration of monkeypox as a Global

Public Health Emergency. JAMA *328*, 615-617. 10.1001/jama.2022.12513.

O'Tool, Á., and Rambaut, A. (2022). Initial observations about putative APOBEC3 deaminase

editing driving short-term evolution of MPXV since 2017. https://virological.org/t/initial-

observations-about-putative-apobec3-deaminase-editing-driving-short-term-evolution-of-mpxv-

since-2017/830.

Oma, Y., Kino, Y., Sasagawa, N., and Ishiura, S. (2004). Intracellular localization of homopolymeric

amino acid-containing proteins expressed in mammalian cells. J Biol Chem *279*, 21217-21222.

10.1074/jbc.M309887200.

Otu, A., Ebenso, B., Walley, J., Barceló, J.M., and Ochu, C.L. (2022). Global human monkeypox

outbreak: atypical presentation demanding urgent public health action. Lancet Microbe *3*, e554-e555.

10.1016/S2666-5247(22)00153-7.

Patel, H., Varona, S., Monzón, S., Espinosa-Carrasco, J., Heuer, M.L., Underwood, A., Gabernet, G.,

nf-core bot, Ewels, P., MiguelJulia, et al. (2022). nf-core/viralrecon: nf-core/viralrecon v2.4.1 -

Plastered Magnesium Marmoset. https://zenodo.org/record/6320980#.YzJOTzTMJyE.

10.5281/zenodo.6320980.

Patrono, L.V., Pléh, K., Samuni, L., Ulrich, M., Röthemeier, C., Sachse, A., Muschter, S., Nitsche,

A., Couacy-Hymann, E., Boesch, C., et al. (2020). Monkeypox virus emergence in wild chimpanzees

reveals distinct clinical outcomes and viral diversity. Nat Microbiol *5*, 955-965. 10.1038/s41564-020-

0706-0.

Phillips, C., Gettings, K.B., King, J.L., Ballard, D., Bodner, M., Borsuk, L., and Parson, W. (2018).

"The devil's in the detail": release of an expanded, enhanced and dynamically revised forensic STR

Sequence Guide. Forensic Sci Int Genet *34*, 162-169. 10.1016/j.fsigen.2018.02.017.

Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A., and Korobeynikov, A. (2020). Using SPAdes

De Novo Assembler. Curr Protoc Bioinformatics *70*, e102. 10.1002/cpbi.102.

Pugh, C., Brown, E.S., Quinn, X., Korman, L., Dyas, B.K., Ulrich, R.G., and Pittman, P.R. (2016).

Povidone iodine ointment application to the vaccination site does not alter immunoglobulin G

antibody response to smallpox vaccine. Viral Immunol *29*, 361-366. 10.1089/vim.2016.0025.

Radonić, A., Metzger, S., Dabrowski, P.W., Couacy-Hymann, E., Schuenadel, L., Kurth, A., Mätz-

Rensing, K., Boesch, C., Leendertz, F.H., and Nitsche, A. (2014). Fatal monkeypox in wild-living

sooty mangabey, Côte d'Ivoire, 2012. Emerg Infect Dis *20*, 1009-1011. 10.3201/eid2006.13-1329.

Reynolds, M.G., Yorita, K.L., Kuehnert, M.J., Davidson, W.B., Huhn, G.D., Holman, R.C., and

Damon, I.K. (2006). Clinical manifestations of human monkeypox influenced by route of infection. J

Infect Dis *194*, 773-780. 10.1086/505880.

Rimoin, A.W., Mulembakani, P.M., Johnston, S.C., Lloyd Smith, J.O., Kisalu, N.K., Kinkela, T.L.,

Blumberg, S., Thomassen, H.A., Pike, B.L., Fair, J.N., et al. (2010). Major increase in human

monkeypox incidence 30 years after smallpox vaccination campaigns cease in the Democratic

Republic of Congo. Proc Natl Acad Sci U S A *107*, 16262-16267. 10.1073/pnas.1005769107.

Roberts, K.L., and Smith, G.L. (2008). Vaccinia virus morphogenesis and dissemination. Trends

Microbiol *16*, 472-479. 10.1016/j.tim.2008.07.009.

Salichs, E., Ledda, A., Mularoni, L., Albà, M.M., and de la Luna, S. (2009). Genome-wide analysis

of histidine repeats reveals their role in the localization of human proteins to the nuclear speckles

compartment. PLoS Genet *5*, e1000397. 10.1371/journal.pgen.1000397.

Sánchez-Seco, M.P., Hernández, L., Eiros, J.M., Negredo, A., Fedele, G., and Tenorio, A. (2006).

Detection and identification of orthopoxviruses using a generic nested PCR followed by sequencing.

Br J Biomed Sci *63*, 79-85. 10.1080/09674845.2006.11732725.

Seeman, T. (2015). Snippy: rapid haploid variant calling and core genome alignment.

https://github.com/tseemann/snippy.

Senkevich, T.G., Yutin, N., Wolf, Y.I., Koonin, E.V., and Moss, B. (2021). Ancient gene capture and

recent gene loss shape the evolution of orthopoxvirus-host interaction genes. mBio *12*, e0149521.

10.1128/mBio.01495-21.

Senkevich, T.G., Zhivkoplias, E.K., Weisberg, A.S., and Moss, B. (2020). Inactivation of genes by

frameshift mutations provides rapid adaptation of an attenuated vaccinia virus. J Virol *94*, e01053-

01020. 10.1128/JVI.01053-20.

Shchelkunov, S.N. (2012). Orthopoxvirus genes that mediate disease virulence and host tropism. Adv

Virol *2012*, 524743. 10.1155/2012/524743.

Shchelkunov, S.N., Totmenin, A.V., Babkin, I.V., Safronov, P.F., Ryazankina, O.I., Petrov, N.A.,

Gutorov, V.V., Uvarova, E.A., Mikheev, M.V., Sisler, J.R., et al. (2001). Human monkeypox and

smallpox viruses: genomic comparison. FEBS Lett *509*, 66-70. 10.1016/s0014-5793(01)03144-1.

Shchelkunov, S.N., Totmenin, A.V., Safronov, P.F., Mikheev, M.V., Gutorov, V.V., Ryazankina,

O.I., Petrov, N.A., Babkin, I.V., Uvarova, E.A., Sandakhchiev, L.S., et al. (2002). Analysis of the

monkeypox virus genome. Virology *297*, 172-194. 10.1006/viro.2002.1446.

Shoubridge, C., Cloosterman, D., Parkinson-Lawerence, E., Brooks, D., and Gécz, J. (2007).

Molecular pathology of expanded polyalanine tract mutations in the Aristaless-related homeobox

gene. Genomics *90*, 59-71. 10.1016/j.ygeno.2007.03.005.

Shumate, A., and Salzberg, S.L. (2021). Liftoff: accurate mapping of gene annotations.

Bioinformatics *37*, 1639-1643. 10.1093/bioinformatics/btaa1016.

Stothard, P. (2000). The sequence manipulation suite: JavaScript programs for analyzing and

formatting protein and DNA sequences. Biotechniques *28*, 1102, 1104. 10.2144/00286ir01.

The Broad Institute (2018). Picard coomand-line tools. https://broadinstitute.github.io/picard/.

Thornhill, J.P., Barkati, S., Walmsley, S., Rockstroh, J., Antinori, A., Harrison, L.B., Palich, R., Nori,

A., Reeves, I., Habibi, M.S., et al. (2022). Monkeypox virus infection in humans across 16 countries

— April–June 2022. N Engl J Med *387*, 679-691. 10.1056/NEJMoa2207323.

Ulaeto, D., Grosenbach, D., and Hruby, D.E. (1996). The vaccinia virus 4c and A-type inclusion

proteins are specific markers for the intracellular mature virus particle. J Virol *70*, 3372-3377.

10.1128/JVI.70.6.3372-3377.1996.

Ulaeto, D.O., Dunning, J., and Carroll, M.W. (2022). Evolutionary implications of human

transmission of monkeypox: the importance of sequencing multiple lesions. Lancet Microbe *3*, e639-

e640. 10.1016/S2666-5247(22)00194-X.

Vaughan, A., Aarons, E., Astbury, J., Balasegaram, S., Beadsworth, M., Beck, C.R., Chand, M.,

O'Connor, C., Dunning, J., Ghebrehewet, S., et al. (2018). Two cases of monkeypox imported to the

United Kingdom, September 2018. Euro Surveill *23*, 1800509. 10.2807/1560-

7917.ES.2018.23.38.1800509.

Verstrepen, K.J., Jansen, A., Lewitter, F., and Fink, G.R. (2005). Intragenic tandem repeats generate

functional variability. Nat Genet *37*, 986-990. 10.1038/ng1618.

Vivancos, R., Anderson, C., Blomquist, P., Balasegaram, S., Bell, A., Bishop, L., Brown, C.S.,

Chow, Y., Edeghere, O., Florence, I., et al. (2022). Community transmission of monkeypox in the

United Kingdom, April to May 2022. Euro Surveill *27*, 2200422. 10.2807/1560-

7917.ES.2022.27.22.2200422.

Vusirikala, A., Charles, H., Balasegaram, S., Macdonald, N., Kumar, D., Barker-Burnside, C.,

Cumiskey, K., Dickinson, M., Watson, M., Olufon, O., et al. (2022). Epidemiology of early

monkeypox virus transmission in sexual networks of gay and bisexual men, England, 2022. Emerg

Infect Dis *28*, 2082-2086. 10.3201/eid2810.220960.

Walker, B.J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.A., Zeng, Q.,

Wortman, J., Young, S.K., and Earl, A.M. (2014). Pilon: an integrated tool for comprehensive

microbial variant detection and genome assembly improvement. PLoS One *9*, e112963.

10.1371/journal.pone.0112963.

Wang, D., Tao, R., Li, Z., Pan, D., Wang, Z., Li, C., and Shi, Y. (2020). STRsearch: a new pipeline

for targeted profiling of short tandem repeats in massively parallel sequencing data. Hereditas *157*, 8.

10.1186/s41065-020-00120-6.

Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., and Barton, G.J. (2009). Jalview

Version 2—a multiple sequence alignment editor and analysis workbench. Bioinformatics *25*, 1189-

1191. 10.1093/bioinformatics/btp033.

Wick, R.R., Judd, L.M., Gorrie, C.L., and Holt, K.E. (2017). Completing bacterial genome

assemblies with multiplex MinION sequencing. Microb Genom *3*, e000132. 10.1099/mgen.0.000132.

Wittek, R., and Moss, B. (1980). Tandem repeats within the inverted terminal repetition of vaccinia

virus DNA. Cell *21*, 277-284. 10.1016/0092-8674(80)90135-x.

Wood, D.E., Lu, J., and Langmead, B. (2019). Improved metagenomic analysis with Kraken 2.

Genome Biol *20*, 257. 10.1186/s13059-019-1891-0.

World Health Organization (2022a). ICD-11. International Classification of Diseases 11th Revision.

https://icd.who.int/en/.

World Health Organization (2022b). Monkeypox: experts give virus variants new names.

https://www.who.int/news/item/12-08-2022-monkeypox--experts-give-virus-variants-new-names.

1208 Yong, S.E.F., Ng, O.T., Ho, Z.J.M., Mak, T.M., Marimuthu, K., Vasoo, S., Yeo, T.W., Ng, Y.K.,

1209 Cui, L., Ferdous, Z., et al. (2020). Imported monkeypox, Singapore. Emerg Infect Dis *26*, 1826-1830.

1210 10.3201/eid2608.191387.

1211 Zhu, X., Jiang, L., Lu, Y., Wang, C., Zhou, S., Wang, H., and Tian, T. (2018). Association of aspartic

1212 acid repeat polymorphism in the asporin gene with osteoarthritis of knee, hip, and hand: a PRISMA-

1213 compliant meta-analysis. Medicine (Baltimore) *97*, e0200. 10.1097/MD.0000000000010200.

1214

Name	Location start[a]	Location end[b]	Repeat unit[c]	Pattern[d]	Nearest OPG[e]	Type of LCR[f]	Relative position to the OPG[g]	Distance in bp[h]	Copenhagen notation[i]	Vaccinia virus (VACV) notation[j]	Comments
LCR1	5,369	5,624	16	[AACTAACTTATGACTT]n	OPG003 (ITR)	STR	Downstream	72	Cop-C19L	NA	
LCR1					OPG015 (ITR)	STR	Upstream	35	CPXV-017	NA	
LCR2	174,063	174,112	2	[ATAT]n	NA	STR	Downstream	46	Cop-B16R	B14R	
LCR3	179,872	180,345	9	ATAT [ACATTATAT]n	OPG208	STR	ATG Start/Promoter	21	Cop-K2L	B19R	SPI-1 apoptosis inhibition
LCR4	193,504	193,759	16	[AAGTCATAAGTTAGTT]n	OPG003 (ITR)	STR	Downstream	72	Cop-C19L	NA	
LCR4					OPG015 (LITR)	STR	Upstream	35	CPXV-017	NA	
LCR5	133,895	133,918	1	[T]n	MPXVgp137	homopolymer	Upstream	889	Cop-A25L	A27L	Fragmented gene area
LCR6	133,980	133,989	10	[CAATCTTTCT]n	MPXVgp137	STR	Upstream	818	Cop-A25L	A27L	
LCR7	137,319	137,375	3	[ATC]n	OPG153	STR	Inside ORF	NA	Cop-A28L	A26L	Attachment MVs/laminin
LCR8	147,655	147,718	5+7	[ATATTTT]n [ATTTT]n [ATATTTT]n [ATTTT]n [ATATTTT]n [ATTTT]n [ATATTTT]n	OPG171	STR	Upstream	75	Cop-A42R	A42R	
					OPG170	STR	Upstream	70	Cop-A41L	A41L	
LCR9	151,350	151,417	9	[TATGAAG]n [GATATGAT]n [GATATGATG]n [GATATGAT]n	OPG176	STR	Upstream	12	Cop-A46R	A47R	
LCR10	197,830	197,842	1	[T]n	OPG001 (ITR)	homopolymer	Downstream	225	NA	NA	
LCR11	1,286	1,298	1	[T]n	OPG001 (ITR)	homopolymer	Downstream	225	NA	NA	
LCR12	29,326	29,364	1	[A]n	OPG044	homopolymer	Inside ORF	NA	Cop-K7R	B15R	C-terminal position
LCR13	76,896	76,904	1	[T]n	OPG097	homopolymer	Inside ORF	NA	Cop-L3L/L4R	L3L/L4R	
LCR14	81,658	81,666	1	[T]n	OPG104	homopolymer	Inside ORF	NA	Cop-J5L	L5L	Essential for viral replication
LCR15	140,911	140,977	9	[ATAACAATT]n [ATAATTGTT]n [ATAATAATT]n [ATAATTGTT]n	OPG159	STR	Inside ORF	NA	Cop-A31L	A33L	PKR inhibitor candidate? / C-terminal position
LCR16	153,457	153,465	1	[A]n	OPG180	homopolymer	Upstream	15	Cop-A50R	A50R	
LCR17	163,979	164,003	4	[TAAC]n	OPG188	STR	Downstream	82	Cop-B2R	B4R	
LCR18	166,865	166,920	7	[AATAATT]n	OPG190	STR	Downstream	15	Cop-B5R	B6R	
LCR19	170,508	170,563	6	[GATACA]n	OPG197	STR	Inside ORF	NA	Cop-B11R	B11R	Hypothetical protein
LCR20	172,868	172,876	1	[T]n	OPG199	homopolymer	Downstream	56	Cop-K2L	SPI-2/B12R	
LCR21	175,299	175,357	6	[GATGAA]n	OPG204	STR	ATG Start/Promoter	NA	Cop-B19R	B16R	Alternative ATG repeat start

a   Nucleotide base coordinate in reference HQG (Genbank# OX044336)

b   Nucleotide base coordinate in reference HQG (Genbank# OX044336)

c   Number of repeat units in the HQG (Genbank# OX044336)

d   Description of the pattern of the LCR in representative MPXV, in which n is the number of repeats for this particular genome

e   Identification according to Senkevich et al. of nearest identified gene. New Notation

f   Type of LCR: short tandem repeats or homopolymer

g   Position of the LCR to the neareast gene

h   Distance of the LCR to the nearest gene

i   Notation of the gene in the VACV Copenhagen strain

j   Notation of the gene in the VACV Western Reserve strain

Name	Repeat Unit	Pattern HQ	Number of	Nearest Gene	Variation	Type of	Entropy	Resolved	Nanopore	MiSeq	NovaSeq	#	#
LCR4	16	TAGTCATAAGTTAGTT [AAGTCATAAGTTAGTT]$_{15}$	16	OPG003 (ITR)	NR	Length	NA	No$^{\&}$	Yes	No	No	NA	NA
LCR3	9	ATAT [ACATTATAT]$_{52}$	52	OPG208	Yes	Length	NA	Yes	Yes	No	No	NA	NA
LCR1	16	[AACTAACTTATGACTT]$_{15}$ AACTAACTTATGACTA	16	OPG003 (ITR)	NR	Length	NA	No$^{\&}$	Yes	No	No	NA	NA
LCR2	2	[AT]$_{25}$	25	NA	Yes	Length	1.66	No	Yes	Yes	Yes	768	90
LCR5	1	[T]$_{24}$	24	OPG152	Yes	Length	1.535	Yes	No	No	Yes	NA	112
LCR10	1	[T]$_{13}$	13	OPG001 (ITR)	Yes	Length	0.63	No$^{\&}$	Yes	No	No	6,561	11,945
LCR11	1	[T]$_{13}$	13	OPG001 (ITR)	Yes	Length	0.627	No$^{\&}$	Yes	Yes	Yes	6,448	11,589
LCR21	6	[GATGAA]$_4$ GATGA	4.5	OPG204	Yes	Mutation	0.207	Yes	Yes	Yes	Yes	6,578	6,661
LCR7	3	[ATC]$_{14}$ TATGAT [ATC]$_3$	19	OPG153	Yes	Length	0.181	Yes	Yes	Yes	Yes	4,541	6,607
LCR9	9	[TATGAAG]$_1$ [GATATGAT]$_1$ [GATATGATG]$_5$ [GATATGAT]$_1$	8	OPG176	No	NA	0	Yes	Yes	Yes	Yes	5,208	5,737
LCR8	5+7	[ATATTTT]$_1$ [ATTTT]$_1$ [ATATTTT]$_3$ [ATTTT]$_1$ [ATATTTT]$_2$ [ATTTT]$_1$ [ATATTTT]$_1$	10	OPG171	No	NA	0	Yes	Yes	Yes	Yes	6,581	6,790
LCR6	10	[CAATCTTTCT]$_1$	1	OPG152	Yes*	NA	0	No*	Yes	Yes	Yes	4,884	12,930
LCR20	1	[T]$_9$	9	OPG199	No	NA	0	Yes	Yes	Yes	Yes	10,106	13,315
LCR19	6	GATTCA [GATACA]$_8$ GAT	9.3	OPG197	No	NA	0	Yes	Yes	Yes	yes	4,119	4,685
LCR18	7	[AATAATT]$_3$ AATAA	3	OPG190	No	NA	0	Yes	Yes	Yes	Yes	9,755	11,838
LCR17	4	[TAAC]$_6$ T	6.1	OPG188	No	NA	0	Yes	Yes	Yes	Yes	7,388	9,474
LCR16	1	[A]$_9$	9	OPG180	No	NA	0	Yes	Yes	Yes	Yes	10,340	16,044
LCR15	9	[ATAACAATT]$_4$ [ATATTGTT]$_1$ [ATAATAATT]$_1$ [ATAATTGTT]$_1$	7	OPG159	No	NA	0	Yes	Yes	Yes	Yes	7,067	6,569
LCR14	1	[T]$_9$	9	OPG104	No	NA	0	Yes	Yes	Yes	Yes	7,819	12,521
LCR13	1	[T]$_9$	9	OPG097/098	No	NA	0	Yes	Yes	Yes	Yes	7,480	12,126
LCR12	1	[A]$_9$	9	OPG044	No	NA	0	Yes	Yes	Yes	Yes	9,789	13,592

*   LCR6 is a 10-bp repeat that was reported early in the outbreak as an insertion (https://virological.org/t/first-german-genome-sequence-of-monkeypox-virus-associated-to-multi-country-outbreak-in-may-2022/812). In our dataset, we have not seen any variation in this area.

$^{\&}$ LCR pairs 1/4 and 10/11 are located in ITRs. Given that no read covering this area reached a unique are outside of the ITRs, we cannot technically state that we solved the repeat. Nonethless, the ITRs should be identical based on the know poxvirid replication mode.

	MPXV-M5312_HM12_Rivers	MPXV_USA_2022_MA001	353R
Genome length	197209	197205	198547
SNPs*	NA	67	69
Indels*	NA	10del 7ins	11del 6 ins
Homopolymeric sites**	408	405	399
Unique SNPs	NA	0	2
LCR characterization			
LCR pair 1/4	8	8	16
LCR2	22	24	25
LCR3	18	16	52
LCR5	25	28	24
LCR6	2	1	1
LCR7	19	17.6	17.6
LCR8	10	10	10
LCR9	8	6	6
LCR pair 10/11	17	14	13
LCR12	9	9	9
LCR13	9	9	9
LCR14	9	9	9
LCR15	7	7	7
LCR16	9	9	9
LCR17	6.1	6.1	6.1
LCR18	3.5	3.5	3.5
LCR19	9.3	9.3	9.3
LCR20	9	9	9
LCR21	4.5	4.5	4.5
*SNPs and indels vs MPXV-M5312_HM12_Rivers			
** Homopolymers >8 nt in length			

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5