Spatial summation of pain is associated with pain expectations: Results from home-based paradigm

Jakub Nastaj[1], Jacek Skalski[1], Aleksandra Budzisz[1], Tibor M. Szikszay[2], Sylwia Swoboda[1], Weronika Kowalska[1],

Daria Nowak[1], Edyta Zbroja[1], Natalia Kruszyna[1], Marta Jakubińska[1], Dominika Grygny[1], Romuald Polczyk[3], Andrzej

Małecki[1], Kerstin Luedtke[2], & Wacław M. Adamczyk[1]*

[1]Laboratory of Pain Research, Institute of Physiotherapy and Health Science, The Jerzy Kukuczka Academy of

Physical Education, Katowice, Poland

[2]Institute of Health Sciences, Department of Physiotherapy, Pain & Exercise Research Lübeck, University of Lübeck,

Lübeck, Germany

[3]Institute of Psychology, Jagiellonian University, Kraków, Poland

*Corresponding author:

E-mail: w.adamczyk@awf.katowice.pl

Number of figures: 5

Number of tables: 3

Number of pages: 22

Number of Supplementary Materials: 8

Conflict of interest: None.

Abstract

The purpose of this study was to reproduce the previously observed spatial summation of pain effect (SSp) using non-laboratory procedures and commercial equipment. An additional aim was to explore the association between expectations and SSp. The Cold Pressor Task (CPT) was used to induce SSp. Healthy participants (N=68) immersed their non-dominant hands (divided into 5 segments) into cold water (Cold Pressor Task). Two conditions were used 1) gradual hand immersion (ascending condition) and 2) gradual hand withdrawal (descending condition). Pain intensity was measured on a Visual Analogue Scale (VAS). The influence of psychological factors, such as the volunteer's expectations of pain intensity, on the actual perception of pain were also measured on a VAS. Results showed significant SSp ($\chi^2(4) = 116.90$, $p < 0.001$), reproduced with non-laboratory equipment in a home-based set-up. Furthermore, two novel findings were observed: i) there was a significant correlation between expectations and perceived pain, indicating a link between pain expectations and SSp, ii) spatial summation increased with the increase in duration exposure to the noxious stimulus (Wald $\chi^2(2) = 157.5$, $p < 0.001$). This study suggests that SSp is shaped by a mixture of excitatory and inhibitory mechanisms and can be influenced by the temporal summation of the nociceptive system. Moreover, this study proposes a new feasible way to induce SSp using a home-based set-up.

Keywords: Spatial summation, noxious cold, cold pressor task, COVID-19, expectations, expectancy

1. Introduction

Spatial summation of pain (SSp) may be characterized as an increase of pain intensity when body area [1–3] or distance [4,5] between stimulated areas are enlarged [3]. There is also evidence that pain thresholds decrease as the area of noxious stimulation increases [6–8]. However, studies show that when the area of noxious stimulation or the distance between stimulated areas exceeds 30 cm (in case of cold pain) [9] or 10 cm (in the case of heat pain) [10] SSp does not occur. Several possible mechanisms have been proposed to contribute to the SSp effect, e.g., local stimulus integration [11], neural recruitment [5,11], lateral inhibition [12], or diffuse noxious inhibitory control [3]. An interesting observation is that there is no exponential or linear increase in pain intensity during a linear increase in the stimulated body area [1] or in the distance between the stimuli [1,4]. Instead, an increase in pain seems to be characterized by a relatively quick saturation when the area of the body exposed to noxious stimulation is expanded in a linear fashion, further suggesting that inhibitory processes are involved in SSp.

The cold pressor task (CPT) is widely used in studies on nociception [2,13–18] and SSp [2,16–21]. In CPT, a body part is immersed in a cold water serving as a noxious stimulus. The low temperature activates nociceptive fibers [18,22,23] through low sensitive ion channels TRPM8 [24] and TRPA1 [25] leading to increasing pain of mild to moderate intensity [26]. Although this method does not allow adequate standardization when there are anthropometric differences, it is easy and commonly used and has been successfully employed in pain studies. A similar method using hot water instead of cold, was also used by Marchand & Arsenault [3] to study SSp. Their results showed that SSp was observed only in the condition using a gradual decrease of the stimulated area but did not occur in a progressively increased stimulated area. Moreover, the perceived pain was less intense during the decreasing compared to the increasing condition.

One of the factors that can be related to SSp are expectations. Indeed, there is evidence that pain can be influenced by expectations [27,28]. Studies on stimulus expectations in pain showed that even short-term predictive cues could have effects on pain perception [29,30]. The role of expected (predicted) value on pain perception could be explained using the predictive coding framework [31,32], wherein the prior distribution of predicted values leads to the shift of posterior distribution following the integration with the collected data, here nociceptive information. Interestingly, despite widely documented effect of expectations on pain perception, any relationship between expectations and SSp has not been studied (or published) yet.

Both publication bias, based on phenomenon of higher probability of publishing statistically significant findings than nonsignificant findings [33], and lack of replication studies are affecting the validity of scientific research [34]. This problem was particularly addressed by researchers in the field of psychology, leading to "crisis of

confidence" in previous findings [35], but it is also found in other fields like medicine [36], economy, [37] or genetic research [38]. To address this is in SSp research we decided to replicate this effect and adopt the methodology from the study of Marchand & Arsenault [3]. Based on previous studies on the CPT paradigm using non-laboratory equipment without water circulation [39,40], the authors decided to additionally test if it is possible to conduct and replicate, for a first time, SSp outside the laboratory setting [21]. This novel yet adopted methodology could enable the conducting of similar replications and preliminary studies for researchers who do not have dedicated equipment for CPT in their laboratories, or it could be used in exceptional situations when conducting research in laboratories is hampered [41,42]. Moreover, the proposed method could be very useful for clinical purposes if there is lack of dedicated CPT equipment.

Furthermore, the current experiment aimed to investigate the potential association between pain-related expectations and SSp by examining participants' predictions regarding pain using an SSp paradigm. It was hypothesized that participants can predict pain produced by noxious stimulation of different sizes.

2. Materials and methods

2.1 General information

The study design was based on the experiment by Marchand & Arsenault [3]. The study was designed as a within-subject experiment. Each participant took part in two consecutive experimental conditions conducted during which they immersed their hand in a cold-water tank. The study was conducted in a home environment, i.e., each examiner performed an experiment within their households (see details below). The home setting differed between examiners. Mainly living rooms, bedrooms and kitchens were chosen to build up the setup. Settings were chosen such that the subject could stay without distractions such as music or the sounds of other household members. Examiners followed an intensive training for the precise data collection and screening procedures. Each examiner was involved in all phases of study preparation. Firstly, they took part in a series of online meetings during which they were introduced to the design of the experiment. Secondly, they received training on the preparation of the CPT (see below). During training, water temperatures achieved in their home environment were measured (water temperature in CPT before adding 6 foil ice-cube packs and after adding ice cubes every 10 minutes for 2h). Thirdly, examiners conducted a pilot study, which was preceded by training in the laboratory. Each of the examiners was individually trained how to follow each step of the study SOP (standard operating procedures) and was supervised by JN. Each examiner conducted a pilot assessment of one single individual in their home environment. To investigate if the CPT method was consistent among examiners, measures of temperatures were compared (**see Supplementary Material 1**). Finally, before conducting main experiment, examiners were demonstrating all procedures to JN, who assessed them in terms of

quality and ensured feedback if any violations from SOP were detected. The study was approved by the Bioethics Committee of Academy of Physical Education in Katowice (1-2021/02-25-21) and was conducted in accordance with the Declaration of Helsinki [43]: Every participant provided written informed consent. The study was pre-registered at the OSF platform (*https://osf.io/kjbdz*). Protocol deviations are described in **Supplementary Material 2.**

2.2 Study population

Examiners (n=13) consisted of members of the Laboratory of Pain Research at the Academy of Physical Education in Katowice or laboratory collaborators. Only healthy participants aged 19-65 years could take part in the study. A thorough interview was conducted using a screening questionnaire which allowed to enroll healthy participants [44]. Participants were excluded if they had a current trauma or wounds in the non-dominant hand, had COVID-19 disease (at the study date or in the past), suffered from acute pain on the study date or within 24 hours preceding the study, took psychoactive substances or medications on the study date, were diagnosed with a disease related to cold temperature intolerance (e.g., Raynaud syndrome, cryoglobulinemia, cold urticaria, etc.), had experienced in the past a pathological reaction to cold temperature (e.g., excessive edema or redness, blisters, etc.). Additionally, due to the fact that the experiment was conducted outside the laboratory and to avoid any adverse events, rigorous exclusion criteria were applied **(see Supplementary Material 3)**. Apart from those derived from previous experiments on cold pressor task (see [45] for example) additional exclusion criteria were obtained from the literature regarding cryotherapy [46,47]. Lastly, if there was any concern about the health condition, the decision to participate was made by a medical doctor who was a member of the research team (AM).

2.3 Equipment for Cold Pressor Task

Because the experiment was designed to be performed in a non-laboratory setting, commercially available equipment and tools were used. Transparent plastic rectangular containers ($36.5 \times 27.5 \times 22$cm) filled with cold tap water (until 15cm height of the container) were used for the Cold Pressor Task (CPT). To obtain the desired water temperature of approximately 5°C, 6 foil ice-cube packs were used (a total of 144 ice cubes) for each experimental condition. An electronic ($\pm0.1\square$) thermometer was attached to the plastic box [21] to monitor the temperature. This non-laboratory version of the CPT was previously validated and led to comparable SSp induced via laboratory-based cold pressor with water circulation [21].

2.4 Experimental design

Before the first immersion of the hand, participants were instructed about the test procedure and prepared for the measurements. Subsequently, the participants were asked about their general fear of pain and fear of pain caused by the cold temperature by using a 0-10 Verbal Rating Scale, where 0 was defined as "no fear of pain" and 10 as "the

greatest fear of pain you can imagine". The non-dominant hand was then divided into 5 segments using the anatomical points of the hand (**Figure 1**): (1) first segment - from the fingertips to the distal interphalangeal joint of the third finger; (2) second segment - from the distal interphalangeal joint to the proximal interphalangeal joint of the third finger; (3) third segment - from the distal interphalangeal joint of the third finger to the metacarpophalangeal joint; (4) fourth segment and (5) fifth segment - the distance between the flexion line of the metacarpophalangeal joint of the third finger and the beginning of the flexion line of the metacarpophalangeal joint of the thumb are divided into two parts. Segment lines were marked on the skin with a pen. Length and width of each segment was measured using flexible measuring tape before the main phase of experiment. The size of areas exposed to cold stimulation is presented in **Figure 2**. Before and during the main phase of experiment, examiners recorded the water temperature at multiple time points as well as the room temperature before each experimental condition. The amount of ice added during the experiment was also recorded. Experiment began when the water temperature had decreased to 4.5 - 5.5□. Two experimental conditions, ascending and descending, were used in the study for all participants. In the ascending condition (**see Figure 1**), participants started from immersion of just 1 segment and sequentially they immersed a greater number of segments finishing with all 5 segments immersed. In the descending condition the order was reversed, i.e., they first immersed all 5 segments and finished with immersion of just 1. In both conditions there was a 5 minutes break between immersion of the next segment to ensure that the skin temperature reached a baseline level [19]. The order of testing (ascending vs. descending condition) was random. The interval between each condition was one hour. During that break, participants were asked to complete the SEWL (subjective experience of workload) questionnaire which measures physical activity level [48,49]. Before and after each experimental condition (ascending, descending), cold pain thresholds (PT_{COLD}) were tested on the examined limb: an ice cube was placed on the palmar surface of the participants' hand and the time (in seconds) until the first pain sensation occurred, was recorded [50]. This procedure was used to control for sensitivity and was an integral part of the screening procedure. At the end, subjects were asked to provide demographic information and guess the purpose of the study. No one of participants knew the correct purpose of the experiment. A familiarization phase was not embedded within the study design to not bias expectations regarding pain from cold water immersion. However, measuring PT_{COLD} prior to the main data collection, allowed participants to familiarize with the cold sensations without inducing expectations. A detailed presentation of the study flow is presented in **Supplementary Material 4**.

2.5 Trial design

Each immersion/trial lasted 60 seconds (regardless of the number of segments involved) and inter-trial intervals were set at 5 minutes [3,20]. Each single trial started with the question about expected pain intensity for this

trial. Participants were told and shown a figure demonstrating the area of the hand which will be exposed to water immersion. Participants were asked to mark a point on the Visual Analogue Scale (VAS) which best reflected their expected pain intensity. The scale for expectations had anchors of 0 (no pain) to 100 (worst pain imaginable). First measurement of expectations (prior to any immersion) started from descending condition segments 5/5 (segments 1-5). Next, participants were instructed to immerse their hand until the line which separated a given number of segments **(Figure 1)**. They were explained and shown to stabilize their thumb to avoid its accidental stimulation. Participants were prompted to rate their pain intensity on the VAS scale (same as for expectations) on the following time points: after 10, 30 and 50s. Participants were blinded towards their previous ratings as these were covered and remained inaccessible during the study. While participants provided ratings, examiners recorded the temperature of the water in a room where assessments were run. After removing their hand, participants were instructed to rest their hand in their axilla for approximately 4 minutes and 30 seconds. This allowed the skin temperature of the hand to return to the baseline level before the next immersion.

2.6 Data extraction and statistical analysis

The main analyses were conducted in the following stages: In the first stage, the effect of stimulation area on pain intensity was investigated using a General Estimated Equations (GEE) model with three within-subject conditions treated as "factors": "condition" (ascending, descending), "segment" (1, 2, 3, 4, 5 segments) "time" (10s, 30s, 50s). The same analysis was repeated with pooled temperature set as a covariate. Polynomial contrast analysis was performed to check the pattern (nonlinear, linear) of pain increase between immersions. Furthermore, in case of significant main and/or interaction effects, t-tests contrasts were performed to describe reported effects.

Next, two sets of expectation ratings (measured before vs. after) were compared within the same statistical model to explore potential differences. The effect of stimulation area on pain expectations was tested using (GEE) with two within-subject factors: "condition" (ascending, descending) and "segment" (1, 2, 3, 4, 5 segments). Next, we analyzed differences in the actual sizes of areas of stimulation (approximated via anthropometric measurements) by using repeated measures ANOVA and size of immersed segments (1,1+2, 1+2+3, 1+2+3+4, 1+2+3+4+5) as dependent variable. Greenhouse-Geisser correction was applied if sphericity assumption was violated.

Lastly, the correlation between pain intensity and pain expectations was conducted using Pearson rank coefficients, scatter plots and regression lines fitted to the data. Correlation between SSp (difference between pain in immersion of 5 segments versus 1 segment) and physical activity (subjective experience of workload [SEWL] score), age, Δarea (difference in size between all 5 segments and 1 segment), and temperature of the water were also tested. The comparison between water temperature and PPTs before and after the experiment was conducted using paired

Student t tests. The α level was set as 0.05 and Sidak correction was employed to control type-I error rate. Unless otherwise indicated figures and tables include descriptive statistics based on raw data. All statistical analyses were performed using IBM Statistical Package for Social Science (SPSS version 25, Armonk, NY, USA).

3. RESULTS

A total number of 68 participants (29 males; 39 females; age: 18-57, mean: 31.08) completed the experiment. Further descriptive characteristics are shown in **Table 1**. The mean temperature of the cold water was 5.13°C during the ascending condition and 5.12°C during the descending condition. In the ascending condition ($t_{[66]} = -0.5$, $p = 0.63$) as well as during the descending condition ($t_{[66]} = 1.2$, $p = 0.22$) no significant differences (before condition vs. after condition) were shown for the pain thresholds, indicating that pain sensitivity was stable over the course of experiment **(Supplementary Material 5)**.

3.1 Primary analyses

Means and standard deviations for pain are presented in **Supplementary Material 6**. The GEE showed a significant main effect for the factor "segment" (Wald $\chi^2(4) = 116.90$, $p < 0.001$), indicating a significant SSp effect. Pairwise post-hoc comparisons showed significant differences in pain between the immersions of different number of hand segments. SSp occurred between segment 1 and segments 1+2 (Mean difference (MD): -4.1; 95% CI -6.84, -1.36), segment 1 and segments 1+2+3 (MD: -9.88; 95% CI -13.66, -6.09), segment 1 and segments 1+2+3+4 (MD: -17.17, 95% CI -22.01, -12.33) as well as between segment 1 and segments 1+2+3+4+5 (MD: -23.72, 95% CI -29.95, -17.49). Furthermore, a linear relationship of these areas could be reported as indicated by the polynomial contrast ($p < 0.001$).

Furthermore, a significant main effect was found for the factor "time" (Wald $\chi^2(2) = 157.45$, $p < 0.001$), indicating that (likely) temporal summation occurred during immersion. Pairwise comparisons showed a significant difference between the first (10s) and second (30s) as well as between the first and third (50s) (last) pain measurement ($p < 0.001$). No significant effect of "condition" (Wald $\chi^2(1) = 1.07$, $p = 0.30$) was found.

Significant two-way interactions were found between the factors "condition" × "segment" (Wald $\chi^2(4) = 18.57$, $p = 0.001$), "condition" × "time" (Wald $\chi^2(2) = 8.31$, $p = 0.02$) and "segment" × "time" (Wald $\chi^2(8) = 80.80$, $p < 0.001$). Pairwise comparisons following these effects revealed that there was a significant difference in pain between the ascending and descending condition but only for the immersion of segment 1 **(Figure 3)** ($p = 0.016$). Likewise, for the two-way "segment" × "time" interaction, pairwise comparisons showed that spatial summation was significant

regardless of the timepoint of measurement, however, largest effects were observed for the last pain measurement (50s) ($p < 0.001$).

A significant three-way, i.e., "condition" × "area" × "time" interaction was shown (Wald $\chi^2(8) = 17.49$, $p = 0.03$), indicating that different spatial summation trajectories were observed across the two conditions (ascending vs. descending) in respect to timepoint of measurement (10, 30, 50s). Exploration of this interaction with the pairwise comparisons showed that significant differences between the ascending and descending condition occurred for the immersion of just segment 1 only during the third (50s) timepoint of measurement ($p = 0.017$) **(Figure 4)**, but not for the second (30s) ($p = 1.00$) and first (10s) ($p = 0.23$). Adding "temperature" as a covariate had only a marginal effect on the three-way interaction (Wald $\chi^2(8) = 15.3$, $p = 0.054$) and had no influence on other statistical results.

Descriptive statistics for expectations are depicted in **Figure 3** and **Table 2**. No significant difference was observed between expectations measured before the first immersion (unbiased) and expectations measured during the assessment (Wald $\chi^2(1) = 1.52$, $p = 0.22$). Thus, for clarity reasons, expectations measured during the assessment are presented below whereas measures taken prior to assessment are in **Supplementary Material 7**. GEE on expectations data did not show significant main effect for the factor "condition" (Wald $\chi^2(1) = 3.04$, $p = 0.08$), but for the factor "segment" (Wald $\chi^2(4) = 111.89$, $p < 0.001$). Pairwise comparisons showed significant differences in expected pain between immersions of different numbers of hand segments (all p values < 0.001). Namely, the expected pain level was lower for a single segment compared to two (MD): -4.95; 95% CI -7.58, -2.32), three (MD: -13.00; 95% CI -17.60, -8.40), four (MD: -19.38, 95% CI -24.92, -13.84) as well as five segments (MD: -23.38, 95% CI -29.82, -16.95). The "segment" × "condition" interaction was also significant (Wald $\chi^2(4) = 27.30$, $p < 0.001$), indicating that the pattern of increase in expected pain level was different across two conditions **(Figure 3)**. Pearson coefficients ($r = 0.53 - 0.81$) were statistically significant for all correlations between expected pain and actual pain **(Table 3 and Figure 5)**.

3.2 Results of exploratory analyses

Exploratory correlations revealed no significant relationship between the magnitude of SSp and physical activity ($r = -0.04$, $p = 0.72$), maximal increment in the stimulated area (total stimulation area for 5 segments minus only 1 segment, $r = -0.13$, $p = 0.35$), or mean water temperature ($r = -0.11$, $p = 0.36$), age ($r = -0.03$, $p = 0.82$). Further, the repeated measures ANOVA of the size of immersed segments showed a significant effect for "size", indicating that the area of stimulation increased from trial to trial ($F_{(1.08, 72.17)} = 135.17$, $p < 0.001$, $\eta^2_p = 0.70$).

15   Polynomial contrasts showed that the increase in size of the stimulated area could be explained by both a linear and an

16   exponential function ($F_{(1,67)} = 35.47$, $p < 0.001$, $\eta^2_p = 0.35$).

4. Discussion

The main aim of this study was to reproduce the SSp effect, reported previously using an ascending/descending paradigm [2,3,20] and noxious cold stimulation [2,20]. The second aim was to introduce a novel methodology to study SSp in non-laboratory setting. The last aim was to use this novel methodology to investigate the associations between expectations and SSp.

Current results confirmed that SSp, as previously shown with noxious cold [20], and heat [3] stimulation of the whole upper extremity, can be reproduced in a home-based setting. This finding implies that the proposed "adapted" paradigm is feasible for conducting bedside testing in clinical and non-laboratory environments. Furthermore, our results contribute to the mechanistic understanding of SSp by showing that spatial summation might be predicted from subjects' pain expectations. Moreover, SSp is strongly influenced by temporal summation. Additionally, differential effects of the order of immersions indicate that -to some degree- SSp is shaped by descending pain inhibition.

4.1 Reproducibility of spatial-related effects

The current study aimed to reproduce the effect previously shown by Marchand & Arsenault [3], yet using a modified methodology. In the mentioned experiment, participants' upper extremities were divided into 8 segments, such that the first segment included only the fingertips and the last segment included the entire arm (from fingertips to axilla). They used three experimental sessions: increasing session (immersion from fingertip to the axilla), decreasing session (immersion from the axilla to fingertip) and whole-arm-first + increasing session (first immersion to the axilla and next from fingertip to the axilla). Authors not only observed a significant SSp effect, as pain was on average higher when a larger area was stimulated, but also an interaction between the sequence of immersions and the size of the stimulated area. This interaction indicated that the same size of stimulated body area (segments) was perceived differently in the ascending (immersion from fingertip to the axilla) compared to the descending (from axilla to fingertips) condition. Their findings demonstrated that SSp was only detected in the decreasing session and whole-arm-first + increasing session and did not take place in the increasing session condition. The authors proposed an explanation that in the increasing condition, both facilitatory and inhibitory mechanisms were gradually being recruited. The results from the third condition (whole-arm-first + increasing session) appear to support their suggestion. If the gradual recruitment of inhibitory mechanisms is responsible for the lack of correlation between the stimulated area and reported pain during the increasing session, a positive relationship between the stimulated area and pain should be reestablished in a whole-arm-first + increasing session whereas starting with the whole arm immersion would stimulate the activation of inhibitory mechanisms at the beginning of the session. In our study, this interaction

was replicated, although it was prominent when only the immersion of the first segment (fingertips) was compared. In line with the previous observation, this interaction can be a manifestation of a robust activation of the descending pain inhibitory system in the descending condition.

Interestingly, SSp was reproduced in our experiment in a cohort of 68 healthy individuals, despite introduced changes to original methodology [3,20,51]. Firstly, only the hand was used and divided into 5 segments. Secondly, here we used noxious cold stimulation. Thirdly, the study was conducted with home-based equipment outside of the laboratory. The latter aspect supports the robustness of observed findings. Despite larger variability and random noise cause by different CPT temperatures within the individual households, SSp was stable. Results were replicated even when controlling for water temperature.

4.2 Physiological mechanisms of SSp

The mechanisms of SSp are not fully understood, however, it seems that both facilitatory and inhibitory processes interact during summation. Inhibitory processes could be inferred from an interaction between the sequence and the stimulated body area (segments). It was observed that when the stimulation started from the largest area, pain provoked by immersion of segment 1 was slightly lower than the analogue stimulation in the ascending sequence. That discrepancy can be explained by the fact that in the descending sequence, inhibitory mechanisms are activated to their maximum and persist during subsequent immersions, thereby resulting in lower pain during the immersion of segment 1, which is in line with a previous study [3]. Moreover, a study on rats which showed that an increase in stimulation area from 1.9 to 18cm^2 gradually decreased the frequency of convergent neuron discharges in intact, yet not spinally transected animals [52]. Furthermore, the increase in pain in either sequence was disproportional. A 5 times larger area does not multiply the reported pain intensity by the same value, which is in line with previous SSp studies [1,3,20].

As for facilitatory processes, it appears that they could be modulated by other factors. Temporal summation has not yet been considered in SSp induced via CPT. Our results suggest that the duration of the stimulus strongly affects the summation pattern. The curve representing the summation trajectory was steeper when the last measurements were considered (50s), indicating that SSp can be partially mediated by the temporal and/or spatial summation of nociceptors innervating deeper layers of soft tissue [53] and possible veins [54].

Another feature of SSp is the pattern of pain increase which was explained by a linear and a non-linear equation to the same extent. In a recent experiment with electrical stimuli, the pain increase followed a logarithmic curve when the size of the stimulated area increased in a linear fashion [1]. It can be hypothesized that in the current study, the increase in the stimulus area was rather exponential **(Figure 4)**, which could result in a less efficient

inhibition and thus a more linear increase in pain. However, this requires further research as a linear fit only negligibly less fit to our data.

4.3 Spatial summation using cold pain

Additionally to the topical administration of cold stimuli using a thermode [9,55], SSp was previously reproduced by CPT in 6 experiments [16]; five of these showed significantly higher pain during the immersion of a larger area of the body [16,17,20,21]. A first attempt by Wolf & Hardy [18], was not successful. The authors compared pain provoked by the immersion of one finger to pain provoked by whole hand immersion and did not observe SSp. However, the study sample was small (n=2), and results could be explained by individual differences in SSp which are known to be large: lateral inhibition [12] may paradoxically lead to lower pain in a larger stimulated area [12,19]. In a study by Westcott et al. [17], SSp was confirmed in 40 individuals by demonstrating more intense pain during full-hand immersion compared to the immersion of one finger into water of 0°C. In one study, SSp was provoked with a temperature comparable to that used in the current experiment (4.7°C) [16]. Participants withdrew their hand faster during stimulation of the whole hand compared to partial immersion. Julien et al. [20] divided the upper extremity into 8 segments and performed ascending and descending immersions with a water temperature of 12°C, which was necessary to allow patients with chronic pain to tolerate the stimuli. They induced robust SSp and its disruption in the group of patients suffering from chronic pain. Two recent studies from our group confirmed the existence of SSp during hand immersion by dividing the hand into 5 segments [21], or 2 segments (ulnar and radial side) [19].

4.4 Spatial summation in non-laboratory setting

Assessment of spatial summation outside the laboratory has never been investigated apart from our pilot study [21]. The proposed methodology was inspired by difficulties with the data collection during the COVID-19 pandemic. However, proposed methodology exceeds beyond pandemic use. For instance, it could be used to conduct replications and preliminary studies by researchers who do not have specialized equipment for CPT in their laboratories. In fact, experimental pain research often requires expensive equipment allowing for e.g. a control of the temperature of a stimulus during the experiment [26]. This methodology can also be helpful for assessing participants with CPT who cannot come to the laboratory e.g., due to their health condition. Furthermore, it can be discussed that the current study is a step forward by moving the laboratory-based pain research into field studies. This approach could be also very useful in clinical context when there is lack of specialized equipment for CPT or there are difficulties to transport CPT equipment to patients who have musculoskeletal dysfunctions.

The methodology employed here was recently proposed in a preliminary validation study, conducted on 9 volunteers. This previous publication validated home-based CPT against laboratory-based equipment with a constant temperature and circulation of the water [21]. The pilot study showed that the SSp trajectory was comparable in the laboratory vs. the non-laboratory paradigm, although the average temperature of the water might have been higher and more variable in the home set-up. In another study by McIntyre et al. [56] healthy participants were trained (online) to self-administer CPT and showed that 97% of participants did not report issues with the test procedure of CPT. These findings together with the current results suggest, that the complex assessment of pain modulation can be used safely in the home environment. Nonetheless, it is important to note a number of difficulties in applying such a methodology. The first one is maintaining the target temperature by adjusting the volume of CPT (water and ice) which must be preceded by testing. It is advisable to compare the resulting temperature with the available specialized CPT, as the authors have done [21]. Another is the problem with standardizing the environment in which the assessment is conducted. The environment outside the lab does not provide as much control over the environmental stimuli as laboratory setting does. For this reason, research under such conditions can be more challenging and require very strict procedure of conduct.

4.5 Expectations and spatial summation

Expectations have not yet been considered in SSp. Correlations reported in the current study were positive and significant in all cases, indicating that participants expected more pain before immersion of the larger area of their body. Participants could correctly adjust their expectations after each immersion and correctly predict that larger area of immersion causes more intense pain in ascending condition. The same results were observed for descending condition as participants correctly predict decrease in pain during subsequent immersions of smaller area. This results could be explained by predictive coding [31,57] as an increase or decrease in pain intensity during consecutive immersions was in line with participants expectations.

In contrast to SSp, expectations -so far- were considered in conditioned pain modulation (CPM), a pain-inhibiting-pain effect. In a typical CPM experiment, participants are exposed to a test stimulus after the pre-exposure to a conditioning stimulus (CS). In one study [58], a high level of expected pain hampered the magnitude of the CPM effect. However, in a study by France et al. [59], expectations were matching the level of pain after application of the CS. Namely, those participants who expected lower reductions in CPM, experienced a more robust inhibitory effect. In another study, expectations were manipulated, showing that expectation -if influenced by verbal suggestion- lead to a reduced CPM effect - but only in females [60]. As both CPM and SSp are pain modulation paradigms testing the spatial aspects of pain, it is reasonable to assume that expectations also shape SSp.

4.6 Limitations and future directions

There are two main limitations to this study. First is the lack of standardization of the environmental factors that result from conducting this study in outside the laboratory settings. Examiners conducted the study in their households, resulting in significant differences in the characteristics of the selected rooms and environmental stimuli. The second is that the study was conducted by multiple examiners, differing in factors such as age and gender that can affect the subjects. Another is that the examiners varied significantly in the number of subjects assessed which comes as the result of restrictions during COVID-19 pandemic. Unfortunately, a direct test of the influence of the examiners on the result was not possible, because of the large number of examiners. Each examiner tested very few participants; some only one participant. Therefore, including examiners as a factor in the ANOVA was not feasible. Numbers of participants per each examiner are included in **Supplementary Material 8**. Future attempts to adapt methodologies using non-laboratory equipment should emphasize standardizing the environment in which measurements take place to control environmental stimuli. As the results showed, subjects' expectations correlate with the pain they experience. Also, test subjects are able to predict the intensity of pain. Further research should focus on the manipulation of expectations to see if the subjects' level of expectations affects SSp and is able to modulate it.

4.7 Conclusions

This study successfully replicated the SSp using CPT. It proposes a new method to induce SSp using a non-laboratory setup. Also, three novel aspects emerged from this study. Firstly, SSp can be assessed outside of the laboratory, providing a new tool for experiments outside of the laboratory in e.g., clinical settings. Secondly, it seems that SSp can be shaped by a mixture of excitatory and inhibitory mechanisms and is influenced by the temporal summation of the nociceptive system. Lastly, spatial summation seems to be strictly linked with expectations, but future studies that directly modulate expectations are needed to investigate their influence on SSp, and physiological measurements should also be taken.

5. ACKNOWLEDGEMENTS

The authors declare no conflict of interest. The authors would like to thank Magdalena Grygiel and Daniel Adamczyk for help with conducting the experiment.

6. REFERENCESS

1. Adamczyk WM, Manthey L, Domeier C, Szikszay TM, Luedtke K. Nonlinear increase of pain in distance-based and area-based spatial summation. Pain. 2021 Jun;162(6):1771–80.

2. Julien N, Marchand S. Endogenous pain inhibitory systems activated by spatial summation are opioid-mediated. Neurosci Lett. 2006 Jul;401(3):256–60.

3. Marchand S, Arsenault P. Spatial summation for pain perception: interaction of inhibitory and excitatory mechanisms. Pain. 2002 Feb;95(3):201–6.

4. Adamczyk WM, Szikszay TM, Kung T, Carvalho GF, Luedtke K. Not as "blurred" as expected? Acuity and spatial summation in the pain system. Pain. 2021 Mar;162(3):794–802.

5. Quevedo AS, Coghill RC. Filling-In, Spatial Summation, and Radiation of Pain: Evidence for a Neural Population Code in the Nociceptive System. J Neurophysiol. 2009 Dec;102(6):3544–53.

6. Defrin R, Ronat A, Ravid A, Peretz C. Spatial summation of pressure pain: effect of body region. Pain. 2003 Dec;106(3):471–80.

7. Lautenbacher, Jesper Nielsen, Thim S. Spatial summation of heat pain in males and females. Somatosens Mot Res. 2001 Jan;18(2):101–5.

8. Defrin R, Urca G. Spatial summation of heat pain: a reassessment. Pain. 1996 Jul;66(1):23–9.

9. Defrin R, Sheraizin A, Malichi L, Shachen O. Spatial summation and spatial discrimination of cold pain: Effect of spatial configuration and skin type. Pain. 2011 Dec;152(12):2739–45.

10. Defrin R, Givon R, Raz N, Urca G. Spatial summation and spatial discrimination of pain sensation. Pain. 2006 Dec;126(1):123–31.

11. Price DD, McHaffie JG, Larson MA. Spatial summation of heat-induced pain: influence of stimulus area and spatial separation of stimuli on perceived pain sensation intensity and unpleasantness. J Neurophysiol. 1989 Dec 1;62(6):1270–9.

12. Quevedo AS, Mørch CD, Andersen OK, Coghill RC. Lateral inhibition during nociceptive processing. Pain. 2017 Jun;158(6):1046–52.

13. Freeman R, Barabasz A, Barabasz M, Warner D. Hypnosis and Distraction Differ in Their Effects on Cold Pressor Pain. Am J Clin Hypn. 2000 Oct;43(2):137–48.

14. Johnson MI, Ashton CH, Bousfield DR, Thompson JW. Analgesic effects of different frequencies of transcutaneous electrical nerve stimulation on cold-induced pain in normal subjects. Pain. 1989 Nov;39(2):231–6.

15. Keogh E, Herdenfeldt M. Gender, coping and the perception of pain. Pain. 2002 Jun;97(3):195–201.

16. Martikainen IK, Närhi MV, Pertovaara A. Spatial integration of cold pressor pain sensation in humans. Neurosci Lett. 2004 May;361(1–3):140–3.

17. Westcott TB, Huesz L, Boswell D, Herold P. Several Variables of Importance in the Use of the Cold Pressor as a Noxious Stimulus in Behavioral Research. Percept Mot Skills. 1977 Apr;44(2):401–2.

18. Wolf S, Hardy JD. STUDIES ON PAIN. OBSERVATIONS ON PAIN DUE TO LOCAL COOLING AND ON FACTORS INVOLVED IN THE "COLD PRESSOR" EFFECT. J Clin Invest. 1941 Sep 1;20(5):521–33.

19. Adamczyk WM, Katra M, Szikszay TM, Peugh J, King CD, Luedtke K, Coghill RC. Spatial Tuning in Nociceptive Processing Is Driven by Attention. J Pain. 2023 Jun;24(6):1116-1125

20. Julien N, Goffaux P, Arsenault P, Marchand S. Widespread pain in fibromyalgia is related to a deficit of endogenous pain inhibition. Pain. 2005 Mar;114(1):295–302.

21. Skalski J, Nastaj J, Swoboda S, Budzisz A, Zbroja E, Małecki A, et al. Nielaboratoryjna adaptacja badania przestrzennego sumowania bólu w dobie pandemii COVID-19. BÓL. 2021;21:1–7.

22. Campero M, Serra J, Ochoa JL. C-polymodal nociceptors activated by noxious low temperature in human skin. J Physiol. 1996 Dec 1;497(2):565–72.

23. Middleton SJ, Barry AM, Comini M, Li Y, Ray PR, Shiers S, et al. Studying human nociceptors: from fundamentals to clinic. Brain. 2021 Jun 22;144(5):1312–35.

24. McKemy DD, Neuhausser WM, Julius D. Identification of a cold receptor reveals a general role for TRP channels in thermosensation. Nature. 2002 Mar;416(6876):52–8.

25. Kozyreva TV, Kozaruk VP, Meyta ES. Skin TRPA1 ion channel participates in thermoregulatory response to cold. Comparison with the effect of TRPM8. J Therm Biol. 2019 Aug;84:208–13.

26. Vonbaeyer C, Piira T, Chambers C, Trapanotto M, Zeltzer L. Guidelines for the cold pressor task as an experimental pain stimulus for use with children. J Pain. 2005 Apr;6(4):218–27.

27. Atlas LY, Wager TD. How expectations shape pain. Neurosci Lett. 2012 Jun;520(2):140–8.

28. Fields HL. How expectations influence pain. Pain. 2018 Sep;159(1):S3–10.

29. Atlas LY, Bolger N, Lindquist MA, Wager TD. Brain Mediators of Predictive Cue Effects on Perceived Pain. J Neurosci. 2010 Sep 29;30(39):12964–77.

30. Koyama T, McHaffie JG, Laurienti PJ, Coghill RC. The subjective experience of pain: Where expectations become reality. Proc Natl Acad Sci. 2005 Sep 6;102(36):12950–5.

31. Song Y, Yao M, Kemprecos H, Byrne A, Xiao Z, Zhang Q, et al. Predictive coding models for pain perception. J Comput Neurosci. 2021 May;49(2):107–27.

32. Büchel C, Geuter S, Sprenger C, Eippert F. Placebo Analgesia: A Predictive Coding Perspective. Neuron. 2014 Mar;81(6):1223–39.

33. Greenwald AG. Consequences of prejudice against the null hypothesis. Psychol Bull. 1975 Jan;82(1):1–20.

34. Nuijten MB. Practical tools and strategies for researchers to increase replicability. Dev Med Child Neurol. 2019 May;61(5):535–9.

35. Pashler H, Wagenmakers E. Editors' Introduction to the Special Section on Replicability in Psychological Science: A Crisis of Confidence? Perspect Psychol Sci. 2012 Nov;7(6):528–30.

36. Nosek BA, Errington TM. Making sense of replications. eLife. 2017 Jan 19;6:e23383.

37. Ioannidis JPA, Stanley TD, Doucouliagos H. The Power of Bias in Economics Research. Econ J. 2017 Oct 1;127(605):F236–65.

38. Ioannidis JPA. Calibration of credibility of agnostic genome-wide associations. Am J Med Genet B Neuropsychiatr Genet. 2008 Sep 5;147B(6):964–72.

39. Uman LS, Stewart SH, Watt MC, Johnston A. Differences in High and Low Anxiety Sensitive Women's Responses to a Laboratory Based Cold Pressor Task. Cogn Behav Ther. 2006 Dec;35(4):189–97.

40. Vigil JM, Rowell LN, Alcock J, Maestes R. Laboratory Personnel Gender and Cold Pressor Apparatus Affect Subjective Pain Reports. Pain Res Manag. 2014;19(1):e13–8.

41. Miki Y, Chubachi N, Imamura F, Yaegashi N, Ito K. Impact of COVID-19 restrictions on the research environment and motivation of researchers in Japan. Prog Disaster Sci. 2020 Dec;8:100128.

42. Sohrabi C, Mathew G, Franchi T, Kerwan A, Griffin M, Soleil C Del Mundo J, et al. Impact of the coronavirus (COVID-19) pandemic on scientific research and implications for clinical academic training – A review. Int J Surg. 2021 Feb;86:57–63.

43. World Medical Association. World Medical Association Declaration of Helsinki. Ethical principles for medical research involving human subjects. Bull World Health Organ. 2001;79(4):373–4.

44. Coghill RC, Yarnitsky D. Healthy and normal? The need for clear reporting and flexible criteria for defining control participants in quantitative sensory testing studies. Pain. 2015 Nov;156(11):2117–8.

45. Liu X, Wang S, Chang S, Chen W, Si M. Effect of Brief Mindfulness Intervention on Tolerance and Distress of Pain Induced by Cold-Pressor Task: Brief Mindfulness Intervention on Tolerance Pain. Stress Health. 2013 Aug;29(3):199–204.

46. Lombardi G, Ziemann E, Banfi G. Whole-Body Cryotherapy in Athletes: From Therapy to Stimulation. An Updated Review of the Literature. Front Physiol. 2017 May 2;8:258.

47. Swenson C, Swärd L, Karlsson J. Cryotherapy in sports medicine. Scand J Med Sci Sports. 2007 Jan 30;6(4):193–200.

48. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. Am J Clin Nutr. 1982 Nov 1;36(5):936–42.

49. Knapik A, Saulicz E, Kuszewski M, Plinta R. An Analysis of Relations Between a Self-Assessment of Health and Active Life-Style. Med Sport. 2009 Mar 1;13(1):17–21.

50. Tilley P, Bisset L. The Reliability and Validity of Using Ice to Measure Cold Pain Threshold. BioMed Res Int. 2017;2017:1–6.

51. de Souza JB, Potvin S, Goffaux P, Charest J, Marchand S. The Deficit of Pain Inhibition in Fibromyalgia Is More Pronounced in Patients With Comorbid Depressive Symptoms. Clin J Pain. 2009 Feb;25(2):123–7.

52. Bouhassira D, Gall O, Chitour D, Bars DL. Dorsal horn convergent neurones: negative feedback triggered by spatial summation of nociceptive afferents. Pain. 1995 Aug;62(2):195–200.

53. Campero M, Bostock H. Unmyelinated afferents in human skin and their responsiveness to low temperature. Neurosci Lett. 2010 Feb;470(3):188–92.

54. Klement W, Arndt JO. The role of nociceptors of cutaneous veins in the mediation of cold pain in man. J Physiol. 1992 Apr 1;449(1):73–83.

55. Machet-Pietropaoli H, Chery-Croze S. Spatial summation of thermal pain in human beings. Sens Processes. 1979;3(2):183–7.

56. McIntyre MH, 23andMe Research Team, Kless A, Hein P, Field M, Tung JY. Validity of the cold pressor test and pain sensitivity questionnaire via online self-administration. Pan F, editor. PLOS ONE. 2020 Apr 16;15(4):e0231697.

57. Wiech K. Deconstructing the sensation of pain: The influence of cognitive processes on pain perception. Science. 2016 Nov 4;354(6312):584–7.

58. Traxler J, Hanssen MM, Lautenbacher S, Ottawa F, Peters ML. General versus pain specific cognitions: Pain catastrophizing but not optimism influences conditioned pain modulation. Eur J Pain. 2019 Jan;23(1):150–9.

59. France CR, Burns JW, Gupta RK, Buvanendran A, Chont M, Schuster E, et al. Expectancy Effects on Conditioned Pain Modulation Are Not Influenced by Naloxone or Morphine. Ann Behav Med. 2016 Aug;50(4):497–505.

60. Bjørkedal E, Flaten. Expectations of increased and decreased pain explain the effect of conditioned pain modulation in females. J Pain Res. 2012 Aug;289.

    7. Figures and Tables

Figure 1. Experimental procedure. Example of the "ascending" condition with the hand divided into 5 segments: In

the ascending condition, nociceptive stimulation started from a small area of the hand (fingertips) and increased in

subsequent immersions. In the descending condition, the order was reversed: nociceptive stimulation started with the

whole hand (segments 1 to 5) and decreased in subsequent immersions.

Figure 2. Hand areas exposed to noxious cold stimulation. Exponential (red) growth negligibly better ($R^2 = 0.48$ vs.

0.47) than linear (blue) fit the increase ($p < 0.001$) of the area of stimulation.

Figure 3. Significant interactions between segment(s) immersions and condition. Left: Estimated marginal mean

pain ratings during immersion of each number of segments for the two conditions (ascending, descending). Right:

Mean expectations ratings prior to the first immersion of each number of segments for the two conditions. *Significant

difference between the ascending and descending condition of the segment 1 ($p = 0.015$). **Significant difference in

expected pain level ($p < 0.01$) between the ascending and descending condition.

Figure 4. Temporal summation drives the interaction between condition (ascending, descending) and spatial summation of pain. Figure shows mean pain ratings collected at 10s (3a), at 30s (3b), at 50s (3c). Note that the

difference in pain between ascending and descending condition becomes significant (p = 0.017) after 50s of immersion.

Figure 5. Associations between predicted pain and actual pain of participants. Associations between mean pain ratings during immersion for each number of segments in the two conditions and expectations measured during immersions in cold water. Left: ascending condition, right: descending condition. Significant correlation coefficients: *p < 0.05, ***p < 0.05.

Table 1. Descriptive statistics

Variable	Mean (SD)
Age (years)	31.08 (12.17)
Height (cm)	173.83 (9.50)
Body mass (kg)	68.23 (14.75)
Fear of Pain General (NRS 0-10)	4.38 (2.15)
Fear of Pain Cold (NRS 0-10)	3.36 (2.19)
SEWL	10.31 (1.27)
Variable	**N**
Sex	F = 39 (57%) ; M = 29 (43%)
Handedness	R = 67 (98.5%) ; L = 1 (1.5%)

F- Female, M -Male, R - right, L - left. NRS - numeric rating scale, SEWL - subjective experience of workload questionnaire, SD – standard deviation

Table 2. Pain-related expectations within spatial summation paradigm

	Ascending				**Descending**			
Segment	**Mean**	**SD**	**M**	**IQR**	**Mean**	**SD**	**M**	**IQR**
1/5	19.5	18.22	9	22.5	17.52	15.22	7	15
2/5	21.57	16.53	16	16	23.47	17.57	14	25.5
3/5	25.05	16.83	21	25.5	31.09	19.35	30	34.5
4/5	32.64	19.26	26.5	32.5	36.61	20.37	38	35.5
5/5	40.41	24.22	32.5	47.5	37.85	21.87	37	39

1/5 – Segment 1, 2/5 – Segments 1 to 2, 3/5- Segments 1 to 3, 4/5 – Segments 1 to 4, 5/5- Segments 1 to 5. SD, standard deviations. M, median. IQR, interquartile range.

Table 3. Correlations between expectation and perceived pain

Condition	**Immersed segments**				
	1/5	**2/5**	**3/5**	**4/5**	**5/5**

Descending	0.85^{***}	0.76^{***}	0.79^{*}	0.82^{*}	0.65^{*}
Ascending	0.73^{***}	0.71^{***}	0.82^{***}	0.74^{***}	0.85^{***}

Pearson`s correlation coefficients: $^{*}p < 0.001$