Evaluation of neural regulation and microglial responses to brain injury in larval zebrafish exposed to perfluorooctane sulfonate

Shannon E. Paquette[1][#], Nathan R. Martin[1][#], April Rodd[1], Katherine E. Manz[2], Manuel Camarillo[1], Eden Allen[1], Hannah I. Weller[3], Kurt Pennell[2], and Jessica S. Plavicki[1]

[1]Department of Pathology & Laboratory Medicine, Brown University, Providence, RI, USA

[2]School of Engineering, Brown University, Providence, RI, USA

[3]Department of Ecology, Evolution, and Organismal Biology, Brown University, Providence, RI, USA

[#]These authors contributed equally

Corresponding Author

Jessica S. Plavicki, PhD

Department of Pathology & Laboratory Medicine
Brown University
70 Ship Street
Providence, RI 02903
Email: jessica_plavicki@brown.edu

The authors have no conflicts of interest to declare.

ABSTRACT

Background: Per- and polyfluoroalkyl substances (PFAS) are biopersistent pollutants that have become global contaminants as a result of their diverse applications in commerce and industry. While some *in vitro* and epidemiological studies have explored the neurotoxic potential of perfluorooctane sulfonate (PFOS), a prevalent PFAS congener, it is unknown how developmental exposure to PFOS affects neuronal communication and other developmentally critical neural cell types, including microglia.

Objectives: We sought to determine the extent to which PFOS exposure disrupts brain health, neuronal activity, and neuron-microglia communication during brain development. In addition, while PFOS impairs humoral immunity, its impact on innate immune cells, including resident microglia, is unclear. As such, we aimed to determine whether microglia are cellular targets of PFOS and, if so, whether disrupted microglial development and/or function could contribute to or is influenced by PFOS-induced neural dysfunction.

Methods: Zebrafish were chronically exposed to either control solution (0.1% DMSO), 7 µM PFOS, 14 µM PFOS, 28 µM PFOS, or 64 µM perfluorooctanoic acid (PFOA). We used *in vivo* imaging and gene expression analysis to assess microglial populations in the developing brain and to determine shifts in microglial state. We functionally challenged microglial using a larval brain injury model and, to assess the neuronal signaling environment, performed functional neuroimaging experiments utilizing the photoconvertible calcium indicator CaMPARI. These studies were paired with optogenetic manipulations of neurons and microglia, an untargeted metabolome wide association study (MWAS), and larval swim behavior assessments.

Results: Developmental PFOS exposure resulted in a shift away from the homeostatic microglia state, as determined by functional and morphological changes, as well as transcriptional upregulation of the microglia activation gene *p2ry12*. PFOS-induced effects on microglia state exacerbated microglia responses to brain injury in the absence of increased cell death or inflammation. PFOS exposure also heightened neural calcium activity, and optogenetic silencing of neurons or microglia independently was sufficient to normalize microglial responses to injury. An untargeted MWAS of larval brains revealed PFOS-exposed larvae had neurochemical signatures of excitatory-inhibitory imbalance. Behaviorally, PFOS-exposed larvae also exhibited anxiety-like thigmotaxis. To test whether the neuronal and microglial phenotypes were specific to

PFOS, we exposed embryos to PFOA, a known immunotoxic PFAS. PFOA did not alter thigmotaxis,

neuronal activity, or microglial responses, further supporting a role for neuronal activity as a critical modifier

of microglial function following PFOS exposure.

Discussion: Together, this study provides the first detailed account of the effects of PFOS exposure on

neural cell types in the developing brain *in vivo* and adds neuronal hyperactivity as an important endpoint

to assess when studying the impact of toxicant exposures on microglia function.

INTRODUCTION

Pollution poses a global threat to environmental, ecologic, and economic health and stability. As

international geologic committees consider whether we have now entered the "Anthropocene Age," it is of

increasing importance to understand the broad impacts of human-made pollutants. It is also worth

emphasizing that the burden of pollution is disproportionately shared, with minority and low-income

communities being more susceptible to the physical consequences of insufficient regulatory laws, industrial

waste disposal, and air pollution[1-3]. If we are to protect our health, society, and environment, more research

efforts are needed to address the costs of our past and present environmental negligence.

One major class of chemicals of increasing concern are per- and polyfluoroalkyl substances (PFAS). While

PFAS were only introduced in industrial manufacturing in the mid-20th century, the expansive and

international utilization of PFAS has led to near universal exposure in humans[4, 5]. PFAS congeners are

found in fire-fighting foams, commercial household products such as water-repellant fabrics, food

packaging, and non-stick finishes, and are used in a variety of applications in the aerospace, aviation, and

automotive industries[6]. Structurally, PFAS consist of fully or polyfluorinated aliphatic substances of varying

carbon chain length and head groups, with longer chain length tending to be associated with increased

toxicity[7-9]. The strength of carbon-fluorine bonds lends most PFAS to be biopersistent, bioaccumulative,

and resistant to degradation[10]. As such, PFAS have acquired the disconcerting moniker "forever

chemicals."

Of the 4,700+ PFAS congeners[6], one of the most environmentally prevalent and extensively studied is

perfluorooctane sulfonate (PFOS)[10]. PFOS has an 8-carbon chain with a sulfonic acid head group and

adversely impacts the functioning and health of several major organ systems[11, 12], and is considered a

metabolic, endocrine, and immune disruptor[13-19]. *In utero* and developmental exposure to PFOS has

particularly concerning consequences on adaptive immunity. Grandjean et al. demonstrated in humans that

elevated PFOS levels in infancy and early childhood significantly attenuate adaptive immune responses,

curb antibody production, and limit vaccine efficacy[20, 21]. Such studies reinforce the necessity of

understanding the impact of pollution on population health, especially considering the need to vaccinate

individuals against emerging or evolving pathogens. PFOS-induced humoral immune suppression[14, 22]

prompted the National Toxicology Program to classify PFOS as a "presumed immune hazard to humans"

in 2016[23]. However, the effects of PFOS on innate immunity are still largely inconclusive, with some groups

finding that PFOS dampens innate cell infiltration, gene expression, or activity[24-26], and others describing

heightened inflammation or immune function[27-29]. Even less understood is the tissue-specific impact of

PFOS on local immune populations, namely tissue-resident macrophages. Tissue-resident macrophages

are largely yolk-sac or fetal liver-derived, heterogenous, and most notably carry out discrete, non-canonical

tissue-specific functions essential for development and homeostasis[30]. Determining whether tissue-

resident macrophages are PFAS targets, especially during development, is essential for our understanding

of long-term consequences of macrophage dysregulation following PFAS exposure.

In this work, we focus on microglia, the resident immune population of the central nervous system (CNS),

as a potential PFOS target. Beyond effector immune responses, microglia have a highly dynamic and

diverse repertoire of homeostatic functions, including developmental pruning of extra-numerary

synapses[31, 32], facilitating synaptogenesis and maturation[33], and regulation of neural plasticity and

dendritic spine density through frequent synaptic monitoring[34-36]. As sentinels of the CNS, these persistent,

self-renewing cells are also highly sensitive and rapidly adaptable to any changes in their environment.

Microglial dysfunction has been well documented in disease models of various neuropathological states,

including Down Syndrome[37], Alzheimer's disease[38], and epilepsy[39]. Due to the critical regulatory roles

of microglia in neural health and development, there has been a concerted effort to clarify the mechanisms

of bidirectional crosstalk between neurons and microglia[35]. However, the impact of environmental

exposures on neuron-microglia interactions during developmentally sensitive periods of life, including

gestation, has been largely overlooked. Some epidemiological studies suggested correlations between

developmental PFOS exposure and ADHD incidence[40, 41], while others found no such relationship[42-45].

Meanwhile in mice, a single neonatal PFOS exposure was sufficient to alter proteins required for neuronal

growth and synaptogenesis, as well as cause spontaneous behavior and hyperactivity in adults[46, 47].

Developmental exposure to PFOS was also shown to cause hyperactive locomotor activity in larval

zebrafish[48-50]. *In vitro* experiments using cultured neurons revealed that PFOS interacts with inositol 1,4,5-

triphosphate receptors (IP_3Rs) and ryanodine receptors (RyRs), leading to the release of intracellular

calcium stores, which suggests a potential role for calcium in PFOS-induced neurotoxicity[51]. However, it

is still not known whether the induced locomotor activity in zebrafish is attributed to neuronal hyperactivity

or skeletal muscle calcium utilization[52]. It is also not known if potential changes in the neuronal environment

following PFOS exposure can impact microglia function, and vice versa.

Herein, we utilized the versatility, transparency, and *ex vivo* development of the larval zebrafish model to

determine how chronic PFOS exposure impacts microglial and neuronal function *in vivo* during

development. We employed a brain injury model, functional neuroimaging, larval behavioral assays,

optogenetic modulation of cell-specific membrane potential, a metabolome wide association study (MWAS),

and a microglia knockout model in larval zebrafish to provide the first *in vivo* account of PFOS-induced

disruption of microglial and neuronal activity. We also exposed larvae to a non-excitatory PFAS congener,

perfluorooctanoic acid (PFOA), to determine whether structurally similar compounds can have distinct

effects on CNS health and neural cell function.

MATERIALS & METHODS

Zebrafish husbandry

Zebrafish (*Danio rerio*) maintenance and experimental procedures were approved by the Brown University

Institutional Animal Care and Use Committee (IACUC; 19-12-0003) adhering to the National Institute of

Health's "Guide for the Care and Use of Laboratory Animals." Zebrafish colonies were maintained in an

aquatic housing system (Aquaneering Inc., San Diego, CA) maintaining water temperature (28.5 ± 2°C),

filtration, and purification, automatic pH and conductivity stabilization, and ultraviolet (UV) irradiation

disinfection. Adult and larval zebrafish were sustained in a 14:10 hour light-dark cycle (Westerfield 2000).

Adult zebrafish were fed once a day with GEMMA Micro (Skretting, Westbrook, ME).

Adult zebrafish (varying ages >3 months post-fertilization) were placed into 1.7 L sloped spawning tanks

(Techniplast, USA) 15-18 hours prior to mating. Sexes were separated by a transparent partition. Within 2

hours of light cycle onset, the partition was removed, and zebrafish were allowed to spawn for 1 hour.

Embryos were collected in fresh egg water (60 mg/L Instant Ocean Sea Salts; Aquarium Systems, Mentor,

OH) and placed into 100 mm non-treated culture petri dishes (CytoOne, Cat. No. CC7672-3394). Embryonic

and larval zebrafish were maintained at 28.5 ± 1°C in an incubator (Powers Scientific Inc., Pipersville, PA)

up to 120 hours post-fertilization (hpf).

Zebrafish lines

The following zebrafish lines were used in this study, either independently or in combination. To visualize

green fluorescent macrophages, we used *Tg(mpeg1:EGFP)*[53] zebrafish. To modulate cellular membrane

potential by inducing chloride ion influx, we used the optogenetic line halorhodopsin, which is a light-gated

chloride pump that is sensitive to 589 nm light[54, 55]. This third-generation opsin exhibits reliable membrane

trafficking for uniform surface expression, is resistant to inactivation, and has step-like, potent photocurrents

that are stable over long timescales[55]. Halorhodopsin was under the control of a Gal4-regulated upstream

activating sequence (UAS) for cell-specific control, *Tg(UAS:eNpHR3.0-mCherry)*[54, 55]. To detect calcium

activity in neurons *in vivo,* we used a transgenic line with neuron-specific expression of the calcium sensor

CaMPARI (Calcium-Modulated Photoactivatable Ratiometric Integrator) under the pan-neuronal promoter

elavl3 (*Tg(elavl3:CaMPARI(W391F+V398L))jf9*)[56]*.* CaMPARI is a permanent photoconvertible calcium

sensor that undergoes allosteric chromophore modulation from green-to-red in response to ultraviolet light,

but only upon simultaneous binding of free intracellular calcium[56]. To achieve neuron-specific expression

of a transgene of interest, we used a neuron-driven Gal4 line, *Tg(elavl3:Gal4-VP16)*[57]. Likewise, to achieve

macrophage-specific   expression   of   a   transgene,   we   used   a   macrophage-driven   Gal4,

Tg(mpeg1:Gal4FF)gl25[53]. To visualize neurons and macrophages simultaneously, we used a

photoconvertible pan-neuronal transgenic line in which the neurons express a photoconvertible protein,

Kaede, that converts from green-to-red in response to blue light, *Tg(HuC:Kaede)*[58]. To visualize

macrophages in the red channel as a control experiment for the optogenetic stimulation study, we used a

transgenic line with UAS-driven nitroreductase fused to mCherry, *Tg(UAS:nfsB-mCherry)*[59] and crossed

these fish with the macrophage-Gal4 line. To determine whether or not PFOS-induced microglial

dysfunction during early larval development influenced neuronal excitation and behavior, we used the

embryonic macrophage mutant line targeting the gene *irf8* (TALE-NT 2.0; frameshift mutation of the st96

allele)[60]. Irf8 mutants lack all embryonic-derived macrophage populations, including microglia[60]. This line

was kindly provided to us by Dr. William S. Talbot of Stanford University.

Perfluorooctanesulfoinc acid (PFOS) and perfluorooctanoic acid (PFOA) exposure

Perfluorooctane sulfonic acid (PFOS; Sigma-Aldrich, Cat. No. A-5040) and perfluorooctanoic acid (PFOA;

Sigma-Aldrich, Cat. No. 171468) were prepared by dissolving the powdered compounds in 100% DMSO.

PFOS and PFOA stock concentrations were verified using LC-HRMS (certified reference standards and

internal standards were purchased from Wellington, Ontario, Canada). Upon verification of the stock

concentration, we found that the PFOS solution contained the following impurities: 4409.49 ng/L of PFHpS,

4588.56 ng/L of PFHxS, and 7493.40 ng/L of PFNS. PFOS and PFOA stock solutions were diluted by a

factor of 5000x in a mixture 1:1 methanol:water and 2 mM ammonium acetate to accommodate the

detection range of the LC-HRMS. LC/MS grade water and methanol were purchased from Honeywell

(Muskegon, MI 49442). Ammonium acetate solution (5 M) was purchased from Millipore Sigma (Burlington,

MA 01803).

Timed spawns of relevant transgenic zebrafish crosses were performed for 1 hr. Embryos were collected

and screened for embryo quality at 4 hpf. Healthy embryos were placed in 24-well plates at a density of 3

embryos per well. Prior to treatment, PFAS compounds were diluted in egg water to the final concentration

of interest. Egg water containing 0.1% DMSO was used as vehicle control. Embryos were dosed with 2 mL

of diluted PFOS solution or vehicle control per well at 4 hpf. The 24-well plates were sealed with parafilm

to limit evaporative loss and placed in an incubator (28.5 ± 1°C). Embryos were dechorionated at 24 hpf

and statically exposed until the experimental timepoint of interest. PFOS body burden was assessed at 48

hpf and 72 hpf as described in the section "Targeted Analysis of PFOS Using Liquid Chromatography High

Resolution Mass Spectrometry" below.

We selected a range of PFOS concentrations based on previously published developmental toxicology

research that provided important information regarding mortality and gross morphological changes in

response to exposure in larval zebrafish at similar timepoints[61, 62]. The survival rate following exposure to

the three highest concentrations of PFOS used in the study (14 µM PFOS, 28 µM PFOS, and 56 µM PFOS)

was determined between 24 hpf (1 dpf) and 120 hpf (5 dpf). At 120 hpf, there was no significant difference

in larval survival in the 14 µM PFOS-exposed group. However, significant reductions in survival were first

detected at 96 hpf in the 28 µM and 56 µM PFOS-exposed groups (Table S1). As such, larvae exposed to

7 µM and 14 µM PFOS were assessed until 5 dpf, while larvae exposed to 28 µM or 56 µM PFOS were

assessed until 3 dpf.

Brain injury model

At 3 dpf, larval zebrafish were anesthetized in 0.02% tricaine-s solution (Syndel, Ferndale, WA) and restrained dorsal-side-up in 2% agarose (Fisher Scientific, Cat. No. BP160-100). Larvae were injured anteriorly at the right telencephalon with a 9 um OD pulled glass capillary needle. This method was an adaptation for larval fish[63]. For time-lapse imaging, larvae were immediately mounted dorsally on a 35 mm glass bottom microwell dish (MatTek, Part No. P35G-1.5-14-C) in 2% low-melting agarose (Fisher Scientific, Cat. No. BP160-100) surrounded by egg water. Multi-slice projection images of the forebrain and optic tectum at various timepoints post-injury, as well as time-lapse videos composed of 10-minute imaging intervals, were captured using a Zeiss LSM 880 confocal microscope at 20x magnification. Area of microglia response was measured using Zen Blue (Zeiss).

Microglial cell quantification

Adult transgenic zebrafish expressing *Tg(HuC:Kaede)* and *Tg(mpeg1:EGFP)* were crossed to generate *Tg(HuC:Kaede; mpeg1:EGFP)* fish, which had pan-neuronal expression of the photoconvertible protein kaede as well as green macrophages. Embryos were exposed to PFOS as described. At 3 dpf, larvae were fixed in 4% paraformaldehyde (PFA, Sigma-Aldrich, Cat. No. P6148) for 18-24 hours at 4°C. Post-fixation, larvae were washed 3 times in PBS-T (phosphate buffered solution + 0.6% Triton-X 100) (Sigma-Aldrich, Cat. No. X100). PBS-T was removed and VECTASHIELD Mounting Media (Vector Labs, Cat. No. H-1000) was added to the samples. Samples incubated for 5 minutes at room temperature, then were gently mixed and placed in -20°C until imaging. The larvae were removed from VECTASHIELD and mounted in 2% low-melting agarose (Fisher Scientific, Cat. No. BP160-100) in 35mm glass bottom microwell dishes (MatTek, Part No. P35G-1.5-14-C). Prior to image acquisition, the kaede fluorophore was photoconverted from green to red fluorescence using 405 nm light (about 1-minute exposure) on the Zeiss LSM 880 confocal microscope. Microglia were defined as macrophages (mpeg1+ cells) amongst and in contact with differentiated neurons (HuC+ cells) in the forebrain, optic tectum, and hindbrain. The entirety of the zebrafish brain was imaged at 20x magnification (2 tile panels with 10% overlap). Microglia were counted by panning through confocal z-stacks using FIJI/ImageJ. Immune cells in contact with HuC+ differentiated

neurons were counted as parenchymal cells (i.e. microglia). Surface cells in contact with differentiated

neurons that failed to extend into the parenchyma were defined as non-parenchymal cells (i.e.

macrophage). Location and proximity of mpeg1+ cells were confirmed using z-stack orthogonal views.

Microglial morphology quantification following PFOS exposure

Microglial morphology was performed on high-resolution confocal micrographs of the optic tectum in fixed

transgenic zebrafish larvae expressing either *Tg(mpeg1:EGFP), Tg(mpeg1:Gal4FF;UAS:eNpHR3.0-*

mCherry), or *Tg(mpeg1:GalFF;UAS:nfsb-mCherry).* Individual microglia were isolated based on their

specific z-stack range, to prevent analysis of overlapping cells, using FIJI/ImageJ. Z-slices were also

processed to rescale intensity (compensating for loss of intensity in deeper tissue) and modestly

smoothened using Gaussian blur with a sigma radius of 1 to better define cellular extensions and reduce

signal noise. Each cell was individually outlined, and morphological parameters measured with CellProfiler

software (version 3.1.9) or FIJI/ImageJ. Area, perimeter, and cell shape (perimeter-to-area ratio) were

quantified.

Quantitative RT-PCR

Larval zebrafish heads were collected at 3 dpf. To collect, larvae were placed on ice for 4 minutes to

immobilize swim activity but prevent pain or distress during collection[64]. The heads were removed by

cutting at the base of the hindbrain at a 45° angle, limiting collection of the heart and pericardium. Heads

were pooled (n = 10) and flash frozen in liquid nitrogen. Four independent experimental replicates were

collected per treatment group. RNA isolation and purification was carried out using the Rneasy Plus Kit

(QIAGEN). cDNA synthesis was achieved using the SuperScript IV Reverse Transcriptase First-Strand

Synthesis System kit (Invitrogen, Cat. No. 18091050). qRT-PCR for genes of interest was performed with

7.5 ng/uL of cDNA using the ViiA7 Real Time PCR System (Applied Biosystems). Gene targets were

detected by using either pre-designed TaqMan probes (Thermo Fisher Scientific) or custom primers with

Power Track SYBR Green Master Mix (Thermo Fisher, Cat. No. A46012). See list of probes and primers

for qRT-PCR and genotyping in Table S2.

Genotyping *irf8* mutants

irf8[st96/st96] were generated by Shiau et al. via TALEN-targeting[60]. After swim behavior testing or CaMPARI

imaging, whole larvae were placed directly in 50 uL of DNA lysis buffer (10 mM Tris pH 8.4 (Fisher, Cat.

No. BP152-500), 50 mM KCl (Sigma, Cat. No. P9333), 1.5 mM MgCl2 (Sigma, Cat. No. M0250), 0.30%

Tween-20 (Fisher, Cat. No. BP337), 0.30% Igepal/NP-40 (Sigma, Cat. No. 56741)) to be used for

genotyping. Larvae were incubated in DNA lysis buffer for 20 minutes at 94°C, cooled to 55°C, then 5 uL

of 10 mg/mL of proteinase K (Millipore Sigma, Cat. No. 70663-4) was added. All temperature specific

reactions were carried out in the ProFlex PCR System (Applied Biosystems). Samples were then incubated

for 1 hour at 55°C, followed by 20 minutes at 94°C, and held at 4°C until the PCR reaction. The PCR

reaction was performed with 2 uL of DNA and 18 uL of master mix (diH2O, 10x Taq Buffer (Thermo

Scientific, Cat. No. EP0404), 2.5 mM dNTP (Thermo Scientific, Cat. No. R1121), 25 mM MgCl2 (Thermo

Scientific, Cat. No. AB0359), 10 uM forward primer (IDT), 10 uM reverse primer (IDT), DMSO (Millipore

Sigma, Cat. No. D8418), and Taq DNA polymerase (Thermo Scientific, Cat. No. EP0404)). The PCR

reaction conditions were as follows: 1 minute at 95°C; 35 cycles of 30 seconds at 95°C, 45 seconds at

68°C, and 30 seconds at 72°C; then 7 minutes at 72°C; and hold at 4°C. The *irf8* exon 1 fragment was

amplified using primers listed in Table S2. As this *irf8* mutant has a frameshift mutation at an AvaI site, the

PCR product was used for restriction digest with AvaI (New England BioLabs, Cat. No. R0152L) to identify

the presence of the mutation. For the restriction digest reaction, 10 uL of PCR product and 10 uL of digest

master mix (diH2O, rCutSmart Buffer, and AvaI) were combined and run on the thermocycler for 20 minutes

at 37°C and cooled to 4°C. The digest reaction product was run in a 2% agarose gel containing RedSafe

nucleic acid staining solution (Bulldog Bio Inc, Cat. No. 21141). The gel results were visualized using the

Bio-Rad Gel Doc XR+.

Acridine orange staining

Cell death was determined using the vital dye acridine orange (AO) in live zebrafish embryos. AO is a cell

permeable stain that selectively intercalates uncoiled nucleic acid present in apoptotic cells[65]. Zebrafish

embryos were injured at 3 dpf and placed in 5 ug/mL acridine orange (ThermoFisher, Cat. No. A1301)

diluted in egg water for 20 minutes. Zebrafish were then washed in egg water for 15 minutes, refreshing

the solution every 5 minutes. Live zebrafish were mounted in 2% agarose and immediately imaged on a

Zeiss LSM 880 confocal microscope. For 1 hour-post injury (hpi) analysis, staining took place from 40

minutes to 1 hpi. For 4 hpi analysis, staining took place from 3 hours and 40 minutes to 4 hpi.

CaMPARI photoconversion and image acquisition

Tg[elavl3:CaMPARI(W391F+V398L)]^{jf9} embryos were screened and the brightest (highest expressing)

were selected for toxicant exposure and subsequent photoconversion. At either 3 dpf or 5 dpf, CaMPARI-

positive larvae were individually placed into a modified 1-well dish (15 mm diameter) containing PFOS or

0.1% DMSO. The dish was placed onto a constructed pedestal within a DanioVision Observation Chamber

(Noldus, Wageningen, The Netherlands) adapted with optogenetics components (Prizmatix, Southland,

MI). Use of the pedestal decreased working distance from the LED light source to the free-swimming larva,

concordantly increasing light intensity. Cells expressing CaMPARI with high calcium content will

photoconvert (green-to-red) in the presence of 405 nm light. Photoconversion was performed by exposing

zebrafish to 405 nm (135 mW/cm^2) wavelength light for 1 minute. Live zebrafish larvae were then

anesthetized in 0.02% tricaine-s (MS-222) and mounted in 2% low-melting agarose in 35mm glass bottom

microwell dishes. Confocal z-stacks were acquired on a Zeiss LSM 880 confocal microscope and maximum

intensity projections generated in Zen Black (Zeiss) representing a global snapshot of neural activity. Image

parameters were set during acquisition of the first image and maintained for the duration of each

experiment.

CaMPARI analyses (R/G ratio calculations)

CaMPARI photoconversion from green-to-red during exposure to 405 nm wavelength light was used to

determine neuronal calcium levels following PFOS exposure at 3 dpf or 5 dpf. To assess the level of

photoconversion, maximum intensity projections were imported into FIJI/ImageJ. Brain regions of interest

including forebrain (FB), habenula (H), optic tectum (OT), cerebellum (Ce), hindbrain (HB), and whole brain

(WB) were manually selected and integrated density of red and green was measured. Measurements were

blank corrected by selecting 3 separate background regions of each image (Equation 1). The ratios of

corrected integrated density were then used to determine R/G ratio for each respective image (Equation 2).

R/G ratios were then normalized to the average control R/G ratio within a particular experiment (Equation

3).

297   <u>Equation 1: Corrected Integrated Density</u> -- $C = d - (\Sigma(b1 + b2 + b3)) \div 3$

298   b = background integrated density

299   d = integrated density of region of interest

301   <u>Equation 2: R/G Ratio</u> -- $R = CRed \div CGreen$

303   <u>Equation 3: Normalized R/G Ratio</u> -- $Normalized\ R/G\ Ratio = R \div \underline{X}$

304            $\underline{X}$= mean of control R/G ratios (for each experimental day)

Pentylenetetrazol (PTZ) exposure

For validation of CaMPARI functionality, 3 dpf *Tg(elavl3:CaMPARI(W391F+V398L))*[jf9] larvae were exposed

to 10 mM pentylenetetrazol (PTZ) for 10 minutes. PTZ was washed out 3 times with egg water immediately

followed by CaMPARI photoconversion, as described. For assessment of microglia response to injury

following PTZ exposure, larvae were exposed to control solution, 28 µM PFOS, or 5 mM PTZ at 3 dpf.

Larvae were injured in the right telencephalon, as described, and placed back in their original dosing

solutions for 4 hours. At 4 hpi, larvae were mounted in 2% agarose in a 35mm glass bottom imaging dish

and multi-slice projection images of the forebrain and optic tectum were captured using a Zeiss LSM 880

confocal microscope at 20x magnification. Area of microglia response was measured using Zen Blue

(Zeiss).

Light/dark behavioral assay

317 Behavioral assessments were performed in 24-well plates (ThermoFisher, Cat. No. 144530) using a

318 DanioVision Observation Chamber with EthoVision XT live-tracking software (Noldus, Wageningen, The

319 Netherlands). Larval zebrafish were exposed to either control solution, 7 µM, 14 µM, or 28 µM PFOS at 4

320 hpf, as described above. Approximately 18-24 hours prior to behavioral assessment, larvae were

321 transferred into individual wells of the same dosing solution. All dosing solutions were prepared and plated

322 on the same day as the initial larval exposure to maintain "chronic" exposure conditions, rather than

323 introducing the larvae to a fresh dosing solution. All light/dark behavioral assays were performed between

324 8 AM and 12 PM to prevent behavioral changes attributable to circadian differences. Briefly, zebrafish were

325 placed into the DanioVision observation chamber and acclimated during a 15-minute dark cycle, followed

326 by a 5-minute light cycle (Light 1), a 5-minute dark cycle (Dark 1), another 5-minute light cycle (Light 2),

327 and completed with a 15-minute dark cycle (Dark 2). Behavioral experiments lasted a total of 45 minutes.

328 Total distance moved (mm) during light/dark cycles was quantified using the EthoVision XT software as a

329 measure for hyperactivity. Anxiety-like behavior following PFOS exposure was assessed by monitoring

330 larval well location (center vs. edge) over the course of the assay. Defined center and edge regions each

331 constituted 50% of total well area. Edge versus center preference was quantified using EthoVision XT.

332 Center avoidance data was normalized relative to controls for each experimental replicate.

333 **Optogenetic manipulation in Noldus behavioral unit**

334 To stimulate the optogenetic channel halorhodopsin in the transgenic lines *Tg(elavl3:Gal4;UAS:eNpHR3.0-*

335 *mCherry)* and *Tg(mpeg1:Gal4ff;UAS:eNpHR3.0-mCherry),* screened larvae were placed a DanioVision

336 observation chamber (Noldus, Wageningen, The Netherlands) outfitted with a 570 nm wavelength laser

337 (Prizmatix, Southland, MI). Halorhodopsin's peak excitation wavelength is 589 nm, though because its

338 excitation spectrum is broad, halorhodopsin is also responsive to 570 nm. Halorhodopsin[+] neurons or

339 microglia were stimulated with 570 nm light for 4 hours then immediately imaged for cell morphology or

340 injury response.

Targeted analysis of PFOS using liquid chromatography high resolution mass spectrometry (LC-HRMS)

Body burden was assessed at 48 hpf and 72 hpf to determine the level of uptake of PFOS in the larvae, as well as to determine the extent to which the detected PFAS impurities (PFNS, PFHpS, PFHxS) were present in the larvae. Body burden results can be found in Table S3. Chemicals: Certified PFOS, PFNS, PFHpS, and PFHxS and their respective isotope labelled reference standards were purchased from Wellington Laboratories (Ontario, Canada). LC/MS grade water and methanol were purchased from Honeywell (Muskegon, MI 49442). Ammonium acetate solution (5 M) was purchased from Millipore Sigma (Burlington, MA 01803).

Sample Extraction: Three technical replicates of ten larvae were pooled, flash frozen in liquid nitrogen, and stored in a -80°C freezer until extracted and defrosted at 20°C. 1 mL methanol was added to the centrifuge tube containing the embryos. The samples were sonicated for 90 minutes, vortex mixed 1 minute, and allowed to reach equilibrium for 3 hours at 20°C. Samples were then centrifuged at 3,000 rpm for 10 minutes. The following was added to an LC analysis vial: 50 µL of the methanol extract, 10 µL labelled PFOS internal standard, and 440 µL of a mixture containing 1:1 methanol:water and 2 mM ammonium acetate.

Targeted analysis of PFOS was performed using a Thermo Liquid Chromatography (LC) Orbitrap Q Exactive HF-X mass spectrometer (MS) equipped with a Thermo Vanquish UHPLC system. Mobile phase A contained 2 mM ammonium acetate in water and mobile phase B contained 2 mM ammonium acetate in methanol. 20 µl of each sample extract was injected in triplicate and separated in a Thermo Hypersil Gold Vanquish C18 column (50 mm X 2.1 mm x 1.9 µm) at a constant temperature of 60°C. PFOS was eluted from the column at a constant flow rate of 0.4 mL/min using a mobile phase gradient as follows: equilibration with 10% B for 1 minute, followed by a gradient ramp from 10% B to 95% B over 4 minutes and held for 2 minutes, and back to 10 % B over 1 minute and held for 2 minutes (total run time 9 minutes, data collected from 0.6 to 8.5 minutes. The MS was operated in full scan dd-MS^2 mode (70 NCE) with an inclusion list for PFOS. Ionization was performed in negative mode with an ionization window of 1.0 m/z, sheath gas flow

rate of 40, auxiliary gas flow rate of 10, sweep gas flow rate of 2, spray voltage of 2.7 kV, 310°C capillary

temperature, funnel RF level of 35, and 320°C auxiliary gas heater temperature. Ions were further

fragmented in the HCD collision cell filled with N2 (produced by a Peak Scientific Nitrogen Generator,

Genius NM32LA). For the full-scan, the Orbitrap was operated with a resolution of 120,000, automatic gain

control (AGC) of 3e6, and maximum dwell time of 100ms. For dd-MS2, the Orbitrap was operated with a

resolution of 15,000, AGC of 2e5, and maximum dwell time of 400ms.

Four ions were monitored for PFOS, including 498.9302 m/z (quantifying), 79.9573 m/z (confirming),

98.9556 m/z (confirming), and 82.9607 m/z (confirming). The retention time of PFOS was 5.18 min.

Quantification was performed in TraceFinder 5.0 General with external seven-point calibration curve

prepared by serial dilutions of the calibration standard. Limits of detection (LOD) were determined by

injecting a calibration 7 standard seven times and using Equation 4:

$\underline{\text{Equation 4: Limits of Detection (LOD)}}$ -- $LOD = (3 * s) \div m$

where s is the sample standard deviation and m is the calibration curve slope. The resulting PFOS LOD

was 33.87 ppt.

Untargeted metabolome wide association study (MWAS)

High resolution metabolomics: The sample extracts were re-analyzed using LC-HRMS to collect untargeted

metabolomics data. A 10 µL volume was injected in triplicate onto on the Thermo LC-Orbitrap system

described above. Two chromatography separation methods were used, normal and reverse-phase. The

normal-phase LC was performed with a HILIC column (Thermo Syncronis HILIC 50 mm X 2.1 mm x 3 µm)

at a constant temperature of 25⁰C. Mobile phase A contained 2 mM ammonium acetate in acetonitrile and

mobile phase B contained 2 mM aqueous ammonium acetate. Metabolites were eluted from the column at

a constant flow rate of 0.2 mL/minute using a solvent gradient as follows: equilibrate with 10% B for 1

minute, increase to 65% B for 9 minutes and hold for 3 minutes, decrease to 10% over 1 minute and hold

for 1 minute. The reverse-phase LC was performed with a C18 column (Thermo Hypersil Gold Vanquish,

50 mm X 2.1 mm x 1.9 µm) at a constant temperature of 60⁰C. Mobile phase A contained 2 mM aqueous

ammonium acetate and mobile phase B contained 2 mM ammonium acetate in acetonitrile. Metabolites

were eluted from the column at a constant flow rate of 0.5 mL/minute using a mobile phase gradient as

follows: equilibration with 2.5% B for 1 minute, increase to 100% B over 11 minutes and held for 2 minutes,

and back to 2.5% B over 1 minute and held for 1.5 minutes (total run time 16.5 minutes, data were collected

from 0.05 to 12.5 minutes). For both normal and reverse-phase LC, the MS was operated in full scan mode

with 120,000 resolution, automatic gain control of 3×10^6, and maximum dwell time of 100 ms. Electrospray

ionization was conducted in positive mode for normal-phase and negative mode for reverse phase LC.

Ionization was performed at a sheath gas flow of 40 units, auxiliary gas flow of 10 units, sweep gas flow of

2 units, spray voltage of 3.5 kV, 310ºC capillary temperature, funnel radio frequency (RF) level of 35, and

320ºC auxiliary gas heater temperature.

<u>Metabolomics data analysis</u>: Data files were converted from *.raw files to *.cdf files using Xcalibur file

Converter, and then processed in R packages apLCMS[66] and xMSanalyzer[67] to produce m/z feature

tables. Feature intensities were normalized by log2 transformation. Association of metabolite feature with

PFOS exposure was assessed using a t-test ($P < 0.05$) in comparison to the control samples. The *P*-values

were adjusted using Benjamini and Hochberg with a false discovery rate (FDR) ≤ 20% to control for Type I

errors in multiple comparisons. Significant metabolites were analyzed for pathway enrichment using

MetaboAnalystR[68] using the zebrafish mummichog curated model, which includes the KEGG, BiGG, and

Edinburgh maps. All metabolomics data analysis was performed in R (version 4.0.2).

Statistical analyses and reproducibility

Each experiment was carried out in at least three independent experimental replicates. An experimental

replicate was considered a cohort of zebrafish that were spawned on separate days and, when applicable,

dosed with separate freshly prepared dosing solutions. Dosing groups for each experimental replicate were

composed of siblings, such that sibling controls could be compared to dosed siblings. For each graph,

datasets were tested for normality using a Shapiro-Wilk test and by plotting a QQ plot. If the data were

normally distributed, parametric tests were performed, when applicable. If the data were not normally

distributed, non-parametric tests were performed, when applicable. When comparing two independent

groups, unpaired parametric t-tests with Welch's correction were performed if the data were normally

distributed, or Mann Whitney U tests (non-parametric) were performed if the data was not normally

distributed. In cases where there were 3 or more groups with one independent variable and the data were

normally distributed, a Welch ANOVA with post-hoc Dunnett's T3 multiple comparisons test was performed.

In cases where there were 3 or more groups with one independent variable and the data were not normally

distributed, a Kruskal-Wallis test (non-parametric ANOVA) with Dunn's test for multiple comparisons was

performed. In cases where there were 3 or more groups with two independent variables and the mean of

each group was compared to the mean of every other group, a two-way ANOVA with a post-hoc Tukey

multiple comparisons test was performed. In cases where there were 3 or more groups with two

independent variables and the mean of two selected groups were selected for comparison, a two-way

ANOVA with a post-hoc Sidak multiple comparisons test was performed. All statistical analyses were

performed using GraphPad Prism 9 (Dotmatics, San Diego, CA).

Data availability

All data is available in the main text and supplementary materials. All targeted and untargeted metabolomics

files will be deposited into Metabolomics Workbench.

RESULTS

Microglial morphology and response to brain injury following chronic PFOS exposure

Whether PFOS affects microglia development and function *in vivo* is not yet known. Given the critical

developmental and modulatory roles of microglia in the CNS, we sought to determine whether embryonic

exposure to PFOS had a direct effect on microglia colonization of the developing brain. Transgenic

zebrafish embryos with macrophage-specific GFP expression under the *mpeg1* promoter

(*Tg(mpeg1:EGFP)*)[53] were chronically exposed to either a control solution (0.1% DMSO) or 28 µM PFOS

from 4 hours post-fertilization (hpf) until 3 days post-fertilization (dpf) (Figure 1A). Using liquid

chromatography with high-resolution mass spectrometry (LC-HRMS), we validated our exposure solutions

and determined that 28 µM PFOS resulted in a total body burden of 70.46 ± 2.72 ng/embryo at 3 dpf (Table

S3). Next, we examined microglial number and morphology at 3 dpf, a time point at which no significant

differences in mortality were observed across the tested concentration range (Table S1). Non-parenchymal,

parenchymal, and total microglia number was not affected at 3 dpf between the control and 28 µM PFOS-

exposed larvae (Figure S1A-C). While microglia number was unchanged, cell morphology was significantly

different. PFOS-exposed microglia acquired a less ramified and more amoeboid cell shape (Figure 1D-E''

vs 1I-J''). This corresponded to a significant reduction in cell area (Figure 1N), cell perimeter (Figure 1O)

and the acquisition of a more circular cell shape as calculated by a decrease in the perimeter-to-area ratio

(Figure 1P). Relative mRNA analysis for the gene *p2ry12*, a G-protein coupled receptor for ATP directly

involved in microglia activation and migration behavior[69-71], was also significantly upregulated in the brains

of PFOS-exposed larvae (Figure 1Q), further supporting that PFOS exposure was affecting microglial state.

To determine whether the phenotypic and transcriptional changes were accompanied by functional

changes, we tested whether pollutant-induced microglia activation affected the ability of microglia to

respond to minor brain injury using an established zebrafish injury model[72, 73]. At 3 dpf, larvae were injured

at the right hemisphere of the telencephalon using a pulled glass needle (OD 9 um; Figure 1B) and microglia

recruitment to the injury site was monitored for the first 4.5 hours post-injury (hpi; Figure 1C). PFOS-

exposed larvae had significantly more microglia recruit to the injury site in the first 4.5 hpi compared to

sibling controls (Figure 1R; Supplemental videos 1 versus 2). Additionally, the area of microglial response

in exposed larvae was significantly expanded at 4 hpi (Figure 1G vs 1L; Figure 1S), which persisted up to

12 hpi (Figure 1H vs 1M; Figure 1T). A major function of microglia following brain injury is the removal of

damaged cells and cellular debris to prevent excessive inflammation, which is deleterious to brain health[74].

Using the nucleic acid stain acridine orange in live larvae, we asked whether an increased incidence of cell

death was driving the microglia response to injury in pollutant-exposed larvae; however, PFOS-exposure

did not result in more cell death in the whole brain nor at the injury site at 1 or 4 hpi (Figure S1D-H).

Moreover, relative mRNA expression of the proinflammatory cytokines *il1β*, *tnfα*, and *il6* were unchanged

in larval brains following exposure (Figure S1I).

Microglia morphology and response to brain injury following optogenetic reversal of

microglia activation state in PFOS-exposed larvae

To determine the extent to which microglial state contributed to the heightened microglial response to injury,

we next asked whether PFOS-induced changes in microglia state could be reversed *in vivo*, and if this

reversal of state was sufficient to normalize the injury response. Since the homeostatic microglia membrane

potential is largely maintained by chloride channel currents[75], we utilized transgenic zebrafish expressing

the   light-gated   chloride   pump   halorhodopsin   (eNpHR3.0)   specifically   in   microglia

(*Tg(mpeg1:Gal4FF;UAS:eNpHR3.0-mCherry)*) (Figure 2A). Larvae with halorhodopsin+ microglia were

exposed to 28 μM PFOS, injured at 3 dpf, and exposed to 570 nm light for 4 hours (Figure 2B,C) to induce

the halorhodopsin construct specifically in mpeg+ cells. We confirmed the PFOS-induced changes in

microglial morphology in unstimulated PFOS-exposed larvae (Figure 2D-E'') and compared the microglia

morphology of these unstimulated larvae to PFOS-exposed larvae stimulated with 570 nm light.

Halorhodopsin stimulation of PFOS-exposed microglia produced a significantly more ramified morphology

(Figure 2H-I''). Additionally, while stimulation had no effect on microglia area (Figure 2L), it significantly

increased the cell perimeter (Figure 2M) and perimeter-to-area ratio (Figure 2N), supporting the transition

from an amoeboid to a ramified state. We also confirmed 570 nm stimulation alone had no impact on

microglia morphology by stimulating and assessing microglia from PFOS-exposed larvae lacking

halorhodopsin expression, *Tg(mpeg1:Gal4FF;UAS:nfsb-mCherry)* (Figure S2). We next tested whether

optogenetic modulation of microglia state was sufficient to rescue their responses to brain injury. Indeed,

while unstimulated larvae had exacerbated microglial responses to injury (Figure 2F vs 2G), stimulation of

halorhodopsin in PFOS-exposed microglia significantly reduced the microglia response (Figure 2G vs 2K;

area of response quantified in Figure 2O). These data suggest that modulation of microglia activity state

alone is sufficient to normalize their functions following toxicant exposure.

Assessment of global and regional neuronal network activity in PFOS-exposed larvae

using CaMPARI

Given that microglia are highly responsive to their microenvironments, microglia hyperresponsiveness in

PFOS-exposed larvae may be influenced by changes in neuronal communication. While evidence suggests

PFOS exposure can impact developing and adult brain health, the direct effects of exposure on neuronal

network activity, whole brain metabolome, and neurotransmitter release are not well understood. To

understand the effect of PFOS on global brain activity *in vivo*, we assessed neuronal calcium activity by

driving the fluorescent calcium sensor CaMPARI (Calcium-Modulated Photoactivatable Ratiometric

Integrator) under the pan-neuronal promoter *elavl3* (*Tg(elavl3:CaMPARI(W391F+V398L))^{jf9}*). CaMPARI is

a permanent photoconvertible calcium sensor that undergoes allosteric chromophore modulation from

green-to-red in response to ultraviolet light, but only upon simultaneous binding of free intracellular

calcium[56]. Therefore, inactive neurons at the time of photoconversion are green, while active neurons

convert to red (Figure 3A-D). In this study, free swimming larvae were subjected to 135 mW/cm^2 of 405 nm

light for 1 minute inside our Noldus behavioral unit (Figure S3A-D). We validated our CaMPARI

photoconversion, imaging, and analysis pipeline by exposing larvae to pentylenetetrazol (PTZ), a GABAA

receptor antagonist known to cause neuronal hyperactivity and seizures in zebrafish[76] (Figure S3E-H

versus 3E'-H'). Indeed, 10 mM PTZ led to a significant ratio-metric increase of neuronal intracellular calcium

in the optic tectum (+22.2%) and a 16.7% increase in whole brain activity (Figure S3J). To determine

whether PFOS exposure affected brain activity, we applied this validated pipeline to 3 dpf larvae exposed

to 28 μM PFOS, the same concentration used for the injury model. PFOS exposure notably heightened

neuronal activity in the forebrain (FB; +11.4%), optic tectum (OT; +14.1%), cerebellum (Ce; +17.9%),

hindbrain (HB; +14.6%), and whole brain collectively (WB; +15.7%) (Figure 3G,I). We also examined

whether lower concentrations of PFOS were able to induce differences in neuronal activity. While there

were no significant differences in network activity at 3 dpf following 7 μM PFOS (Figure 3F,H), this group

had notably and significantly heightened regional and global activity at 5 dpf (Figure 3K,M; FB, +7.1%; OT,

+31.2%; Ce, +24.0%; HB, +15.7%; WB, +20.8%), as did the 14 µM exposed group (Figure 3L,N; FB,

+14.0%; OT, +52.6%; Ce, +51.8%; HB, +34.3%; WB, +39.6%). We also assessed gross brain morphology

of PFOS-exposed larvae and found that while whole brain area was slightly reduced at 3 dpf in the 7 µM (-

2.8%) and 14 µM (-2.7%) groups, brain area was unchanged by 5 dpf, suggesting that the effects of PFOS

on morphology are nominal and temporary (Figure S4).

To better understand the neurochemical changes in the PFOS exposed brain, we performed an untargeted

metabolome wide association study (MWAS) on isolated heads of 3 dpf control and 28 µM PFOS exposed

larvae. Several features were significantly upregulated or downregulated between the control and exposed

groups (Figure S5). Significantly different metabolites from the MWAS were further analyzed for pathway

enrichment using MetaboAnalystR[68]. Several of the significantly enriched pathways, including glutamate

metabolism, aspartate and asparagine metabolism, tyrosine metabolism, and phosphoinositide

metabolism, are involved in neuronal excitation, catecholamine synthesis, and neurotransmitter release,

uptake, and recycling (Figure S5C; Table S4).

We next sought to determine whether the neurochemical imbalances were predictors of abnormal

behavioral activity. To first determine whether our exposure paradigm replicates previously reported

behavioral hyperactivity, we exposed embryos to either 7 µM, 14 µM, or 28 µM PFOS at 4 hpf, and

performed light/dark behavioral assays at 3 dpf, 4 dpf, and 5 dpf (Figure S6A,B). Consistent with previous

locomotor assays at 6 dpf[48], PFOS-exposed larvae were significantly more responsive to light changes,

designated by elevated distance traveled within the well at 3 dpf following 28 µM exposure (Figure S6E-H)

and 5 dpf following 7 µM and 14 µM exposure (Figure S6N-R). In addition to distance traveled, behavior

videos were analyzed to assess the frequency of center avoidance within the wells of a 24-well plate. Similar

to the mammalian open-field test, center avoidance in a well is an indication of zebrafish anxiety-like

behavior, while willingness to cross the center suggests a reduced anxiety-like behavior[77]. As such, we

examined the time each larva spent along the well's edge (anxiety-like) versus center (exploratory)

throughout each light/dark behavioral assay (Figure 4A,B). PFOS exposure resulted in notable and

significant reductions in time spent in the center in all dosed groups at 3 and 5 dpf (Figure 4C-E).

Interestingly, while the 7 µM PFOS group were not significantly more active during the light on cycle at 5

dpf (Figure S6P), they only exhibited center-avoidance during the light on cycle (Figure S7G,H). Likewise,

while 14 µM PFOS-exposed larvae exhibited significant center-avoidance behavior at 3 dpf, they did not

have increased swim activity at 3 dpf (Figure S6H). Further, the 14 µM PFOS group exhibited center-

avoidance behavior in both the light and dark cycles at 5 dpf (Figure S7I,J), though were only significantly

more active when the light was on, and not when the light was off (Figure S6P,Q). This suggests that

anxiety-like behaviors are separable from swim hyperactivity and supports the need to more thoroughly

understand the independent behavioral changes associated with PFOS exposure. Since anxiety-, fear-,

and aversive-like behaviors are regulated by the medial habenula in zebrafish[78, 79], we assessed neuron

activity at the habenula using CaMPARI. PFOS exposure did not result in a difference in habenular activity

at 3 dpf, and only 14 µM PFOS-exposed larvae showed a significant difference in habenular activity at 5

dpf (Figure S8).

Evaluation of light/dark response in microglia mutants exposed to PFOS

Neuron-microglia bi-directional signaling enables the reciprocal regulation of microglia and neuronal

functions, including microglial modulation of neuronal activity and circuit refinement[35, 80]. Considering the

observed shift in microglial state following PFOS exposure, we sought to determine whether or not PFOS-

induced microglial dysfunction during early larval development influenced neuronal excitation and behavior.

To do this, we exposed PFOS to zebrafish with a null mutation for *irf8* (*irf8*$^{st96/st96}$), a gene required for

macrophage formation during primitive and transient definitive hematopoiesis[60]. *Irf8* mutant larvae lack the

earliest embryonic-derived macrophage populations, including microglia[60]. Control-treated *irf8* mutants

had a slight reduction in swim behavior throughout the recording (Figure 5A; gray versus yellow traces),

though were significantly less reactive to the light-to-dark transitions than control-treated wildtype larvae

(Figure 5B; 96.8 $\pm$ 8.3 mm moved versus 71.5 $\pm$ 6.8 mm moved). At 5 dpf, both wildtype and *irf8* mutant

larvae exposed to 8 µM PFOS were significantly more active than control-treated larvae (Figure 5A; Figure

S9E-G). However, unlike the control-treated *irf8* mutants, PFOS-exposed mutants did not display a trend

toward reduced swim activity during the light/dark behavioral assay (Figure 5A; blue versus red traces).

PFOS-exposed mutants also exhibited the most dramatic behavioral response to the light-to-dark

transitions, especially during the second dark cycle (Figure 5B; "Minute 15"). In addition, while wildtype

control larvae and wildtype PFOS-exposed larvae had a 14.9% and 23.8% reduction in swim activity by the

end of the assay ("Minute 30"), respectively, PFOS-exposed *irf8* mutants had a 47.6% reduction in activity

(Figure 5B). These data suggest that microglia may attempt to modulate the PFOS-exposed brain, such

that the absence of microglia may further promote PFOS-induced behavioral reactivity and altered patterns

of activity post-dark transitions. Heightened reactivity may also result in behaviors that impair swim ability,

such as seizing or convulsions. PFOS exposure also results in increases in neuronal-driven CaMPARI in

microglia-deficient larvae, but this was not significant between genotypes (Figure 5C).

Microglia response to brain injury following optogenetic modulation of neuronal activity

Given the bi-directional communication between microglia and neurons and that microglial state can be

modulated by neural activity, we asked whether elevated neural activity alone was sufficient to modify

microglial responses to injury. To test this hypothesis, 3 dpf larvae were treated with 5 mM PTZ to increase

neuronal activity, then were injured in the right telencephalon as described. PTZ-induced neuronal

hyperactivity significantly increased the microglia response at 4 hpi, similar to that of PFOS-exposed larvae

(Figure S10). We next asked whether inhibiting neuronal activity in PFOS-exposed larvae using

optogenetics could normalize the observed hyperresponsive microglial phenotype. Larvae expressing

Tg(elavl3:Gal4;cryaa:RFP;UAS:eNpHR3.0;mpeg1:EGFP), which have pan-neuronal expression of

halorhodopsin as well as GFP-labeled macrophages, were exposed to either a control solution or 28 µM

PFOS at 4 hpf. After brain injury at 3 dpf, larvae were either left unstimulated or stimulated for 4 hours with

570 nm light to silence neuronal activity. Microglia of unstimulated PFOS-exposed larvae were 34.5% more

responsive to injury (Figure 6E,F,I). Conversely, neuronal hyperpolarization rescued microglia

hyperresponsiveness, resulting in a 6.6% decrease in microglia response in stimulated PFOS-exposed

larvae compared to stimulated controls (Figure 6G,H,I), further supporting that pollutant-induced changes

in the neural signaling environment significantly influences microglia behavior, independent of cell death or

inflammation.

Microglia response to brain injury and neuronal network activity following chronic exposure to PFOA, a known immunotoxic congener

To further assess if microglia hyperresponsiveness following PFOS exposure is a result of neuronal

hyperactivity, we exposed larval zebrafish to a structurally similar PFAS congener, perfluorooctanoic acid

(PFOA). Whereas PFOS has a sulfonic acid head group, PFOA is an 8-carbon PFAS with a carboxylic acid

head group. Following the same exposure paradigm used for PFOS exposure (Figure 1A), zebrafish

embryos were dosed with either a control solution (0.1% DMSO) or 64 µM PFOA at 4 hpf. Exposure to 64

µM PFOA did not result in differing swim behavior during light/dark behavioral assays at 5 dpf (Figure S11A-

D). While PFOA-exposed larvae had shorter body length, there were no observable gross morphological

effects on the spine that would affect swimming ability (Figure S12). Using neuronally-driven CaMPARI,

PFOA exposure did not result in any regional or global differences in neuronal network activity in 5 dpf

larvae (Figure 7A-C), unlike the PFOS-exposed groups at this timepoint (Figure 3). Given that PFOA did

not result in neuronal hyperactivity, we asked whether PFOA exposure affected microglial responses to

brain injury. Indeed, PFOA exposure did not result in differing microglial responses to brain injury (Figure

7D-F). Together, these data further support that neuronal hyperactivity is a key driver of microglia response,

and that structurally similar PFAS congeners can have distinct impacts on the CNS.

DISCUSSION

For more than two decades, researchers have revealed the many biologic, ecologic, and

environmental ramifications of PFOS toxicity, as well as the toxicity profiles for a subset of other 'forever

chemical' congeners. Here, we used a combination of *in vivo* imaging of cellular behavior, functional

neuroimaging, optogenetic modulation, and behavioral assays to address the impact of PFOS exposure on

microglia-neuron interactions and activity in larval zebrafish. Our data demonstrated that developmental

PFOS exposure modulated microglia state and induced hyperactivity in response to injury independent of

cell death or inflammation. Since the homeostatic microglia membrane potential is largely maintained by

chloride channel currents[75], we asked whether activating the optogenetic chloride channel halorhodopsin

(eNpHR3.0) could rescue microglia hyperresponsiveness. Indeed, electrical modulation of microglia was

sufficient to revert microglia from an amoeboid to a homeostatic morphology and normalize their responses

to injury, suggesting electrical dysfunction as a previously underappreciated pathway worth interrogating

when studying microglial state and immunotoxicity. PFOS-exposed larvae also exhibited global and

regional increases in neuronal activity and anxiety-like behaviors, a previously unidentified

neurodevelopmental phenotype of PFOS exposure in zebrafish, with only nominal and temporary impacts

on regional brain morphology. We demonstrated that PFOS-induced neuronal hyperactivity was a key

mediator of microglia reactivity and that optogenetic silencing of neurons was sufficient to normalize

microglia responses to injury. Further, exposure to PFOA did not result in neuronal hyperactivity nor

microglia hyperresponsiveness. Together, this study provides the first detailed account of the effects of

PFOS exposure on the developing brain *in vivo* and adds neuronal hyperactivity as an important endpoint

to assess when studying the impact of toxicant exposures on microglia function.

Zebrafish have long been considered excellent models to study innate immune development and

function[81, 82], including to understand neuro-immune interactions[83]. While the immunotoxic impact of

PFOS on adaptive immunity has been well documented[84], the effects on the innate immune arm are less

understood and, at times, contradictory. Studies showing either innate immune activation or suppression

following PFOS exposure are likely attributed to varying exposure paradigms, PFOS concentration, animal

model, age, or cell line used[24, 26, 28, 85, 86]. The innate immune system is highly sensitive to both endogenous

and xenobiotic stimuli and reacts rapidly, creating a signaling cascade that informs all downstream immune

functions[87]. Macrophages in particular have the ability to dynamically polarize from a homeostatic state to

be pro- or anti-inflammatory depending on the environmental needs. As antigen-presenting cells,

macrophages also have the vital role of instructing adaptive immune cells on their responses[87]. The

dependency of the adaptive immune system on the innate emphasizes the need to clarify the immunotoxic

mechanisms of pollutants like PFOS.

645   This work constitutes the first *in vivo* analysis of microglia, the resident immune population of the

CNS, following PFAS exposure. Previous *in vitro* studies using immortalized microglia cell lines suggested

PFOS exposure decreased microglial viability, mitochondrial stability, and increased ROS production *in*

vitro in a concentration-dependent manner[88, 89]. We found that PFOS exposure altered the microglia state,

demonstrated by their morphological transition from ramified to amoeboid shaped, as well as by the

upregulation of the microglia activation gene *p2ry12,* a purinergic receptor that responds to ATP. It was

previously demonstrated that microglia response to injury was mediated by glutamate-evoked calcium

waves and ATP release from the injury site[73]; therefore, ATP released by the injury site and/or through

high neuronal activity could be contributing to the microglial hyperresponsiveness in PFOS-exposed larvae.

It is still unclear to what extent PFOS is acting directly on the microglia or indirectly through neuronal

activation. Microglia activation and injury responses were attenuated by both microglia and neuronal

optogenetic silencing, suggesting that the neuronal environment has a significant influence on the microglia.

Of note, while larvae with neuron-driven halorhodopsin had a 34.5% higher microglial response to injury

following PFOS exposure, the response was not as significant as it was in other genetic backgrounds ($P =$

0.0879). This is likely due to the fact that optogenetic channels and pumps are highly sensitive, and while

we reduced potential exposure to ambient light, leakiness of these opsins may contribute to low level ion

exchange. More work on the baseline permeability of optogenetic channels in a controlled and cell-specific

context is needed.

663   While we are providing the first demonstration of microglial dysfunction in response to toxicant

exposure, a shift away from a homeostatic microglial state has been well documented in disease models

of various neuropathological states. In addition, pharmacological manipulation of microglia state or

prevention of microglia depletion was shown to lead to cognitive and functional improvements in some

neurological disease models[37, 38, 90]. However, it is worth noting that modulation of microglia state is not

inherently pathological. For example, microglia depletion following stroke significantly increased infarct size,

neuronal cell death, and caused calcium overload[91]. Microglia depletion during acute seizures also

exacerbated excitotoxicity and seizure sensitivity[92]. Additionally, the regional specificity of microglia state

in a mouse model of chronic stress was considered a protective and/or adaptive response[93]. Lastly, it is

important to note that the dosing paradigm used in this study has been shown to impact other organ

systems, such as the pancreas[61]. Therefore, one cannot dismiss the potential for systemic effects

contributing to the neurochemical imbalances or altered microglial state. While our timelapse imaging

videos, cell death and inflammation data, and microglia quantifications do not suggest macrophages are

recruited from the periphery to the brain injury site, more extensive analysis using cell lineage tracing from

the periphery would need to be performed to confirm this. However, the potential for the periphery to impact

brain health further emphasizes the benefit of understanding systems-level interactions when studying

PFAS exposure.

Due to the developmental and homeostatic roles microglia have on maintaining the

excitatory/inhibitory balance[94], as well as the situation-specific adverse effects that changes in microglia

state may have on the CNS, we asked whether modulation of microglial state following PFOS exposure

was contributing to the behavioral or neuronal hyperactivity. Using microglia-deficient zebrafish with a

mutation in the gene *irf8*, we found that control-treated *irf8* mutants had a slight but consistent reduction in

swim behavior that was not seen in the PFOS-treated mutants. Additionally, PFOS-exposed mutants had

the greatest response to the light-dark transition, suggesting that microglia may actually attempt to repress

PFOS-induced behavioral hyperactivity. While microglia loss did not affect baseline neuronal calcium

signaling, it is worth noting that microglia loss is distinct from microglial dysfunction, and thus does not rule

out the contribution of microglia to neuronal hyperactivity in this context. For example, signals derived from

dysfunctional microglial could impact neuronal firing and perhaps swim behavioral responses, which would

not be recapitulated in a model of microglial loss. In addition, microglia increasingly accumulate in synaptic

regions between 7 and 28 dpf in zebrafish[95], suggesting that any significant death-dependent or activity-

dependent neuronal pruning may occur developmentally later than the timepoints investigated in this study.

Lastly, the concentrations of PFOS used in this study might increase neuronal activity too substantially or

even irreversibly, such that electrical modulation or loss of microglia is not sufficient to influence the activity

in a measurable way.

To our knowledge, this is the first report of *in vivo* neuronal hyperactivity caused by embryonic

PFOS exposure. We show that embryonic PFOS exposure resulted in higher intracellular calcium

concentrations across multiple brain regions at various concentrations (7 μM, 14 μM, and 28 μM) and time

points (3 dpf & 5 dpf). Both gestational and adult PFOS exposures have been shown to impact calcium

dependent signaling molecules important for memory, including Ca^{2+}/calmodulin dependent kinase II

(CaMKII) and cAMP response element-binding protein (CREB) in rats[96], suggesting that disrupted calcium

signaling is an important mediator of PFOS-induced neurotoxicity. *In vitro* studies also linked PFOS toxicity

to disrupted calcium homeostasis[51, 97]. While the underlying mechanisms remain unknown, possible

explanations include increased influx through activation of L-type Ca^{2+} channels[97] and release of

intracellular calcium stores through interaction with ryanodine and inositol 1,4,5-trisphosphate receptors[51].

Intracellular calcium excess in neurons promotes excitotoxicity and can cause brain damage leading to

various neurological and neurodegenerative disorders[98, 99], emphasizing the need to further understand

the influence of PFAS on neuronal function. PFAS compounds have also been shown to activate

peroxisome proliferator-activated receptors (PPARs) *in vitro* as well as in zebrafish[100, 101]. PPAR-γ in

particular is expressed in neurons, microglia, and astrocytes and mediates inflammatory responses in the

CNS[102]. Interrogating PPAR activation in the CNS following PFOS exposure could also provide important

information about the specific pathways impacted by exposure.

The susceptibility of the developing human brain to PFOS remains a contentious point due to

conflicting data. While some studies have demonstrated significant correlations between developmental

PFOS exposure and ADHD incidence[40, 41], others found no such relationship[42-45, 103]. In the mouse brain,

PFOS concentrations have been shown to increase over time and lead to tonic convulsions, despite a lack

of morphological phenotypes[104]. The heightened regional and global network activity demonstrated in this

study warrants an assessment of convulsive phenotypes and seizure activity. While outside the scope of

this study, determining the threshold and timescale at which neuronal hyperactivity increases incidence of

convulsion would be interesting to pursue.

Similar to previous reports[48-50], we observed hyperactive swim behavior in PFOS-exposed larvae

during light/dark behavioral assays. However, we are the first to report PFOS-induced hyperactivity during

the light and dark phases prior to 6 dpf. Hyperactive behavioral changes corresponded well with increased

intracellular neuronal calcium concentrations. Of note, increased anxiety-like behaviors were separable

from increased swim activity at 3 dpf and 5 dpf and was also specific to light or dark cycles depending on

PFOS concentration. This suggests that there may be further disruptions in neuronal communication

beyond just increased calcium concentrations influencing swim activity. In-depth regional brain activity

analyses, possibly through the integration of developed analytical pipelines[105], could provide further insight

into how PFOS induced neural activation is linked to behavioral abnormalities in the larval zebrafish.

We identify increased anxiety-like behaviors as a novel phenotype of PFOS neurotoxicity in the

larval zebrafish using an adapted open-field test model[106-108]. Anxiety is an associated symptom of several

neurobehavioral disorders linked to PFAS exposures, including autism spectrum disorder and ADHD[45, 109-

111]. PFOS-induced center avoidance has been observed in mice exposed during adulthood[112], but

developmental anxiety-like behavior has not been previously reported. Of note, previous published

research reporting PFOS-induced larval hyperactivity utilized 96-well plates for higher throughput[48, 50];

however, smaller well sizes may not be conducive to conducting anxiety-like behavioral analyses. We

therefore conducted behavioral experiments using 24-well plates, which allowed us to define regions within

the well to quantify where fish spent their time swimming. Because dysregulation of habenular activity has

been associated with increased anxiety, depression, and fear across species, including the zebrafish[78, 79,

113], we assessed whether neuronal calcium was affected in the habenula following PFOS exposure. While

only the 14 μM PFOS group showed neuronal dysregulation at the habenula, all groups displayed anxiety-

like behaviors. This suggests that the habenula may be but one region dictating anxiety-like responses in

larval zebrafish or that temporal differences in habenular activity, not captured using CaMPARI, contribute

to anxiety-like responses.

Not only does this study provide the first *in vivo* assessment of neuronal calcium activity following

PFOS exposure, but it is also the first to evaluate PFOA exposure in this context. The chemical structure

of PFOA is similar to PFOS, differing only by the presence of a carboxylic acid head group rather than

sulfonic acid, respectively. PFOA has been shown to cause significant health effects, including

immunotoxicity, elevated cholesterol, dysregulated liver metabolism, kidney dysfunction, thyroid disease,

among many others[114]. Previous reports using larval zebrafish have found that PFOA concentrations

ranging from 4.4 µM to 80 µM did not result in a significant difference in 6 dpf swim behavior during the

light/dark behavioral assays[48]. At the concentrations and timepoints tested here, PFOA exposure did not

result in heightened larval swim behavior or neuronal network activity. However, others have reported that

5 dpf larvae exposed to 1 µM PFOA[115] as well as 14 dpf larvae exposed to 2 µM PFOA[50] did result in

swim hyperactivity. In addition, while neonatal exposure to PFOA in mice has been shown to result in

abnormal expression of proteins important for brain growth and neuron function[46], PFOA exposure of rat

primary cortical neurons did not affect spontaneous neuronal activity or burst duration[116]. Further analysis

is needed to clarify the extent to which PFOA, and other shorter chain length or carboxylic group PFAS,

could be neurotoxic or contribute to neural dysregulation. Nevertheless, that PFOA-exposed larvae had

normal microglia responses to injury, compared to those exposed to PFOS, further highlights the role of

neuronal activity on microglia function.

This study provides the first *in vivo* analysis of how developmental PFOS exposure disrupts larval

brain health and function. It also highlights the relevance of understanding pollution-induced effects on

innate immune cells, in both a non-canonical developmental and homeostatic context, as well as

considering the long-term consequences of potential antigen-presentation dysfunction. In summary, the

complexity of neural cell communication, especially during the sensitive period of brain development,

emphasizes the importance of studying pollutant exposure in non-isolated systems.

ACKNOWLEDGMENTS

We thank Dr. William Talbot for sharing their $irf8^{st96/st96}$ mutant line with us. We also thank Rachel Cyr for excellent care and oversight of our zebrafish facility. Thank you to other members of the Plavicki lab for their discussions in input during various stages of this study. Lastly, thank you to the undergraduate trainees that were involved in the early coordination of this work, Rekha Dhillon-Richardson and Ratna Patel. This work was supported by a NIEHS Outstanding New Environmental Scientist (ONES) award, and cardiopulmonary vascular COBRE Phase II (2PG20GM103652) awarded to J.S.P. In addition, this work was supported by the Ruth L. Kirschstein Predoctoral Individual National Research Service Award (NRSA; F31HL156460) by the NHLBI awarded to S.E.P. S.E.P and N.R.M. were previously supported by the Brown University Environmental Pathology Training Grant (T32ES007272-26) from NIEHS. The Thermo LC-Orbitrap MS was partially funded by NSF Major Research Instrumentation (MRI) award CBET-1919870 to K.P. (PI) and J.S.P (Co-I).

REFERENCES

1. Bullard RD. Overcoming Racism in Environmental Decisionmaking. Environment: Science and Policy for Sustainable Development. 1994;36(4):10-44. Epub 08 Jul 2010. doi:https://doi.org/10.1080/00139157.1994.9929997.

2. Muller C, Sampson RJ, Winter AS. Environmental Inequality: The Social Causes and Consequences of Lead Exposure. Annual Review of Sociology. 2018;44:263-282. doi:https://doi.org/10.1146/annurev-soc-073117-041222.

3. Liu J, Clark LP, Bechle MJ, Hajat A, Kim SY, Robinson AL, Sheppard L, Szpiro AA, Marshall JD. Disparities in Air Pollution Exposure in the United States by Race/Ethnicity and Income, 1990-2010. Environ Health Perspect. 2021 Dec;129(12):127005. Epub 2021/12/16. doi:10.1289/EHP8584. Cited in: Pubmed; PMID 34908495.

4. Kannan K, Corsolini S, Falandysz J, Fillmann G, Kumar KS, Loganathan BG, Mohd MA, Olivero J, Van Wouwe N, Yang JH, Aldoust KM. Perfluorooctanesulfonate and related fluorochemicals in human blood from several countries. Environ Sci Technol. 2004 Sep 1;38(17):4489-95. Epub 2004/10/06. doi:10.1021/es0493446. Cited in: Pubmed; PMID 15461154.

5. Paul AG, Jones KC, Sweetman AJ. A first global production, emission, and environmental inventory for perfluorooctane sulfonate. Environ Sci Technol. 2009 Jan 15;43(2):386-92. Epub 2009/02/26. doi:10.1021/es802216n. Cited in: Pubmed; PMID 19238969.

6. Group. OUGPCP. Portal on Per and Poly Fluorinated Chemicals. 2020. Available from: https://www.oecd.org/chemicalsafety/portal-perfluorinated-chemicals/.

7. Buck RC, Franklin J, Berger U, Conder JM, Cousins IT, de Voogt P, Jensen AA, Kannan K, Mabury SA, van Leeuwen SP. Perfluoroalkyl and polyfluoroalkyl substances in the environment: terminology, classification, and origins. Integr Environ Assess Manag. 2011 Oct;7(4):513-41. Epub 2011/07/28. doi:10.1002/ieam.258. Cited in: Pubmed; PMID 21793199.

8. Gomis MI, Vestergren R, Borg D, Cousins IT. Comparing the toxic potency in vivo of long-chain perfluoroalkyl acids and fluorinated alternatives. Environ Int. 2018 Apr;113:1-9. Epub 20180128. doi:10.1016/j.envint.2018.01.011. Cited in: Pubmed; PMID 29421396.

9. Chambers WS, Hopkins JG, Richards SM. A Review of Per- and Polyfluorinated Alkyl Substance Impairment of Reproduction. Front Toxicol. 2021;3:732436. The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a

819 potential conflict of interest. Epub 20211122. doi:10.3389/ftox.2021.732436. Cited in: Pubmed; PMID
820 35295153.

821
822 10. Langenbach B, Wilson M. Per- and Polyfluoroalkyl Substances (PFAS): Significance and
823 Considerations within the Regulatory Framework of the USA. Int J Environ Res Public Health. 2021 Oct
824 23;18(21). Epub 2021/11/14. doi:10.3390/ijerph182111142. Cited in: Pubmed; PMID 34769660.

825
826 11. Sunderland EM, Hu XC, Dassuncao C, Tokranov AK, Wagner CC, Allen JG. A review of the pathways of
827 human exposure to poly- and perfluoroalkyl substances (PFASs) and present understanding of health
828 effects. J Expo Sci Environ Epidemiol. 2019 Mar;29(2):131-147. Epub 2018/11/25. doi:10.1038/s41370-
829 018-0094-1. Cited in: Pubmed; PMID 30470793.

830
831 12. Zeng Z, Song B, Xiao R, Zeng G, Gong J, Chen M, Xu P, Zhang P, Shen M, Yi H. Assessing the human
832 health risks of perfluorooctane sulfonate by in vivo and in vitro studies. Environ Int. 2019 May;126:598-
833 610. Epub 2019/03/12. doi:10.1016/j.envint.2019.03.002. Cited in: Pubmed; PMID 30856447.

834
835 13. DeWitt JC, Copeland CB, Luebke RW. Suppression of humoral immunity by perfluorooctanoic acid is
836 independent of elevated serum corticosterone concentration in mice. Toxicol Sci. 2009 May;109(1):106-
837 12. Epub 2009/02/26. doi:10.1093/toxsci/kfp040. Cited in: Pubmed; PMID 19240040.

838
839 14. DeWitt JC, Peden-Adams MM, Keller JM, Germolec DR. Immunotoxicity of perfluorinated
840 compounds: recent developments. Toxicol Pathol. 2012;40(2):300-11. Epub 2011/11/24.
841 doi:10.1177/0192623311428473. Cited in: Pubmed; PMID 22109712.

842
843 15. DeWitt JC, Shnyra A, Badr MZ, Loveless SE, Hoban D, Frame SR, Cunard R, Anderson SE, Meade BJ,
844 Peden-Adams MM, Luebke RW, Luster MI. Immunotoxicity of perfluorooctanoic acid and
845 perfluorooctane sulfonate and the role of peroxisome proliferator-activated receptor alpha. Crit Rev
846 Toxicol. 2009;39(1):76-94. Epub 2008/09/20. doi:10.1080/10408440802209804. Cited in: Pubmed; PMID
847 18802816.

848
849 16. DeWitt JC, Williams WC, Creech NJ, Luebke RW. Suppression of antigen-specific antibody responses
850 in mice exposed to perfluorooctanoic acid: Role of PPARalpha and T- and B-cell targeting. J
851 Immunotoxicol. 2016;13(1):38-45. Epub 2015/01/17. doi:10.3109/1547691X.2014.996682. Cited in:
852 Pubmed; PMID 25594567.

853
17. Stein CR, McGovern KJ, Pajak AM, Maglione PJ, Wolff MS. Perfluoroalkyl and polyfluoroalkyl
substances and indicators of immune function in children aged 12-19 y: National Health and Nutrition
Examination Survey. Pediatr Res. 2016 Feb;79(2):348-57. Epub 2015/10/23. doi:10.1038/pr.2015.213.
Cited in: Pubmed; PMID 26492286.

18. Braun JM. Early-life exposure to EDCs: role in childhood obesity and neurodevelopment. Nat Rev
Endocrinol. 2017 Mar;13(3):161-173. Epub 2016/11/20. doi:10.1038/nrendo.2016.186. Cited in:
Pubmed; PMID 27857130.

862
863 19. Chang ET, Adami HO, Boffetta P, Wedner HJ, Mandel JS. A critical review of perfluorooctanoate and
864 perfluorooctanesulfonate exposure and immunological health conditions in humans. Crit Rev Toxicol.
865 2016;46(4):279-331. Epub 2016/01/14. doi:10.3109/10408444.2015.1122573. Cited in: Pubmed; PMID
866 26761418.

867
868 20. Grandjean P, Andersen EW, Budtz-Jorgensen E, Nielsen F, Molbak K, Weihe P, Heilmann C. Serum
869 vaccine antibody concentrations in children exposed to perfluorinated compounds. JAMA. 2012 Jan
870 25;307(4):391-7. Epub 2012/01/26. doi:10.1001/jama.2011.2034. Cited in: Pubmed; PMID 22274686.

871
872 21. Grandjean P, Heilmann C, Weihe P, Nielsen F, Mogensen UB, Timmermann A, Budtz-Jorgensen E.
873 Estimated exposures to perfluorinated compounds in infancy predict attenuated vaccine antibody
874 concentrations at age 5-years. J Immunotoxicol. 2017 Dec;14(1):188-195. Epub 2017/08/15.
875 doi:10.1080/1547691X.2017.1360968. Cited in: Pubmed; PMID 28805477.

876
877 22. Peden-Adams MM, Keller JM, Eudaly JG, Berger J, Gilkeson GS, Keil DE. Suppression of humoral
878 immunity in mice following exposure to perfluorooctane sulfonate. Toxicol Sci. 2008 Jul;104(1):144-54.
879 Epub 2008/03/25. doi:10.1093/toxsci/kfn059. Cited in: Pubmed; PMID 18359764.

880
881 23. NTP. Monograph on Immunotoxicity Associated with Exposure to Perfluorooctanoic acid (PFOA) and
882 perfluorooctane sulfonate (PFOS). 2016. Available from:
883 https://ntp.niehs.nih.gov/ntp/ohat/pfoa_pfos/pfoa_pfosmonograph_508.pdf.

884
885 24. Castano-Ortiz JM, Jaspers VLB, Waugh CA. PFOS mediates immunomodulation in an avian cell line
886 that can be mitigated via a virus infection. BMC Vet Res. 2019 Jun 25;15(1):214. Epub 2019/06/27.
887 doi:10.1186/s12917-019-1953-2. Cited in: Pubmed; PMID 31238913.

888
889 25. Keil DE, Mehlmann T, Butterworth L, Peden-Adams MM. Gestational exposure to perfluorooctane
890 sulfonate suppresses immune function in B6C3F1 mice. Toxicol Sci. 2008 May;103(1):77-85. Epub
891 2008/02/07. doi:10.1093/toxsci/kfn015. Cited in: Pubmed; PMID 18252804.

892
26. Ryu MH, Jha A, Ojo OO, Mahood TH, Basu S, Detillieux KA, Nikoobakht N, Wong CS, Loewen M,
Becker AB, Halayko AJ. Chronic exposure to perfluorinated compounds: Impact on airway
hyperresponsiveness and inflammation. Am J Physiol Lung Cell Mol Physiol. 2014 Nov 15;307(10):L765-
74. Epub 2014/09/14. doi:10.1152/ajplung.00100.2014. Cited in: Pubmed; PMID 25217661.

27. Dong GH, Zhang YH, Zheng L, Liang ZF, Jin YH, He QC. Subchronic effects of perfluorooctanesulfonate
exposure on inflammation in adult male C57BL/6 mice. Environ Toxicol. 2012 May;27(5):285-96. Epub
2010/08/26. doi:10.1002/tox.20642. Cited in: Pubmed; PMID 20737580.

28. Qazi MR, Bogdanska J, Butenhoff JL, Nelson BD, DePierre JW, Abedi-Valugerdi M. High-dose, short-
term exposure of mice to perfluorooctanesulfonate (PFOS) or perfluorooctanoate (PFOA) affects the
number of circulating neutrophils differently, but enhances the inflammatory responses of macrophages
to lipopolysaccharide (LPS) in a similar fashion. Toxicology. 2009 Aug 21;262(3):207-14. Epub
2009/06/23. doi:10.1016/j.tox.2009.06.010. Cited in: Pubmed; PMID 19540903.

29. Wang LQ, Liu T, Yang S, Sun L, Zhao ZY, Li LY, She YC, Zheng YY, Ye XY, Bao Q, Dong GH, Li CW, Cui J.
Perfluoroalkyl substance pollutants activate the innate immune system through the AIM2
inflammasome. Nat Commun. 2021 May 18;12(1):2915. Epub 2021/05/20. doi:10.1038/s41467-021-
23201-0. Cited in: Pubmed; PMID 34006824.

30. Davies LC, Jenkins SJ, Allen JE, Taylor PR. Tissue-resident macrophages. Nat Immunol. 2013
Oct;14(10):986-95. Epub 2013/09/21. doi:10.1038/ni.2705. Cited in: Pubmed; PMID 24048120.

31. Paolicelli RC, Bolasco G, Pagani F, Maggi L, Scianni M, Panzanelli P, Giustetto M, Ferreira TA, Guiducci
E, Dumas L, Ragozzino D, Gross CT. Synaptic pruning by microglia is necessary for normal brain
development. Science. 2011 Sep;333(6048):1456-8. eng. Epub 2011/07/21.
doi:10.1126/science.1202529. Cited in: Pubmed; PMID 21778362.

32. Schafer DP, Lehrman EK, Kautzman AG, Koyama R, Mardinly AR, Yamasaki R, Ransohoff RM,
Greenberg ME, Barres BA, Stevens B. Microglia sculpt postnatal neural circuits in an activity and
complement-dependent manner. Neuron. 2012 May 24;74(4):691-705. Epub 2012/05/29.
doi:10.1016/j.neuron.2012.03.026. Cited in: Pubmed; PMID 22632727.

33. Hoshiko M, Arnoux I, Avignone E, Yamamoto N, Audinat E. Deficiency of the microglial receptor
CX3CR1 impairs postnatal functional development of thalamocortical synapses in the barrel cortex. J
Neurosci. 2012 Oct;32(43):15106-11. eng. doi:10.1523/JNEUROSCI.1167-12.2012. Cited in: Pubmed;
PMID 23100431.

34. Parkhurst CN, Yang G, Ninan I, Savas JN, Yates JR, 3rd, Lafaille JJ, Hempstead BL, Littman DR, Gan
WB. Microglia promote learning-dependent synapse formation through brain-derived neurotrophic
factor. Cell. 2013 Dec 19;155(7):1596-609. Epub 2013/12/24. doi:10.1016/j.cell.2013.11.030. Cited in:
Pubmed; PMID 24360280.

35. Szepesi Z, Manouchehrian O, Bachiller S, Deierborg T. Bidirectional Microglia-Neuron
Communication in Health and Disease. Front Cell Neurosci. 2018;12:323. Epub 2018/10/16.
doi:10.3389/fncel.2018.00323. Cited in: Pubmed; PMID 30319362.

36. Wake H, Moorhouse AJ, Jinno S, Kohsaka S, Nabekura J. Resting microglia directly monitor the
functional state of synapses in vivo and determine the fate of ischemic terminals. J Neurosci. 2009 Apr
1;29(13):3974-80. Epub 2009/04/03. doi:10.1523/JNEUROSCI.4363-08.2009. Cited in: Pubmed; PMID
19339593.

37. Pinto B, Morelli G, Rastogi M, Savardi A, Fumagalli A, Petretto A, Bartolucci M, Varea E, Catelani T,
Contestabile A, Perlini LE, Cancedda L. Rescuing Over-activated Microglia Restores Cognitive
Performance in Juvenile Animals of the Dp(16) Mouse Model of Down Syndrome. Neuron. 2020 Dec
9;108(5):887-904 e12. Epub 2020/10/08. doi:10.1016/j.neuron.2020.09.010. Cited in: Pubmed; PMID
33027640.

38. Mancuso R, Fryatt G, Cleal M, Obst J, Pipi E, Monzon-Sandoval J, Ribe E, Winchester L, Webber C,
Nevado A, Jacobs T, Austin N, Theunis C, Grauwen K, Daniela Ruiz E, Mudher A, Vicente-Rodriguez M,
Parker CA, Simmons C, Cash D, Richardson J, Consortium N, Jones DNC, Lovestone S, Gomez-Nicola D,
Perry VH. CSF1R inhibitor JNJ-40346527 attenuates microglial proliferation and neurodegeneration in
P301S mice. Brain. 2019 Oct 1;142(10):3243-3264. Epub 2019/09/11. doi:10.1093/brain/awz241. Cited
in: Pubmed; PMID 31504240.

39. Hiragi T, Ikegaya Y, Koyama R. Microglia after Seizures and in Epilepsy. Cells. 2018 Mar 28;7(4). Epub
2018/03/31. doi:10.3390/cells7040026. Cited in: Pubmed; PMID 29597334.

40. Hoffman K, Webster TF, Weisskopf MG, Weinberg J, Vieira VM. Exposure to polyfluoroalkyl
chemicals and attention deficit/hyperactivity disorder in U.S. children 12-15 years of age. Environ Health
Perspect. 2010 Dec;118(12):1762-7. Epub 2010/06/17. doi:10.1289/ehp.1001898. Cited in: Pubmed;
PMID 20551004.

41. Lenters V, Iszatt N, Forns J, Cechova E, Kocan A, Legler J, Leonards P, Stigum H, Eggesbo M. Early-life
exposure to persistent organic pollutants (OCPs, PBDEs, PCBs, PFASs) and attention-deficit/hyperactivity
disorder: A multi-pollutant analysis of a Norwegian birth cohort. Environ Int. 2019 Apr;125:33-42. Epub
2019/02/01. doi:10.1016/j.envint.2019.01.020. Cited in: Pubmed; PMID 30703609.

42. Fei C, McLaughlin JK, Lipworth L, Olsen J. Prenatal exposure to perfluorooctanoate (PFOA) and
perfluorooctanesulfonate (PFOS) and maternally reported developmental milestones in infancy. Environ
Health Perspect. 2008 Oct;116(10):1391-5. Epub 2008/10/23. doi:10.1289/ehp.11277. Cited in: Pubmed;
PMID 18941583.

43. Lien GW, Huang CC, Shiu JS, Chen MH, Hsieh WS, Guo YL, Chen PC. Perfluoroalkyl substances in cord
blood and attention deficit/hyperactivity disorder symptoms in seven-year-old children. Chemosphere.
2016 Aug;156:118-127. Epub 2016/05/14. doi:10.1016/j.chemosphere.2016.04.102. Cited in: Pubmed;
PMID 27174824.

44. Liew Z, Ritz B, von Ehrenstein OS, Bech BH, Nohr EA, Fei C, Bossi R, Henriksen TB, Bonefeld-Jorgensen
EC, Olsen J. Attention deficit/hyperactivity disorder and childhood autism in association with prenatal
exposure to perfluoroalkyl substances: a nested case-control study in the Danish National Birth Cohort.
Environ Health Perspect. 2015 Apr;123(4):367-73. Epub 2015/01/24. doi:10.1289/ehp.1408412. Cited in:
Pubmed; PMID 25616253.

45. Ode A, Kallen K, Gustafsson P, Rylander L, Jonsson BA, Olofsson P, Ivarsson SA, Lindh CH, Rignell-
Hydbom A. Fetal exposure to perfluorinated compounds and attention deficit hyperactivity disorder in
childhood. PLoS One. 2014;9(4):e95891. Epub 2014/04/25. doi:10.1371/journal.pone.0095891. Cited in:
Pubmed; PMID 24760015.

46. Johansson N, Eriksson P, Viberg H. Neonatal exposure to PFOS and PFOA in mice results in changes in
proteins which are important for neuronal growth and synaptogenesis in the developing brain. Toxicol
Sci. 2009 Apr;108(2):412-8. Epub 2009/02/13. doi:10.1093/toxsci/kfp029. Cited in: Pubmed; PMID
19211617.

47. Johansson N, Fredriksson A, Eriksson P. Neonatal exposure to perfluorooctane sulfonate (PFOS) and
perfluorooctanoic acid (PFOA) causes neurobehavioural defects in adult mice. Neurotoxicology. 2008
Jan;29(1):160-9. Epub 2007/12/08. doi:10.1016/j.neuro.2007.10.008. Cited in: Pubmed; PMID
18063051.

1001

1002   48. Gaballah S, Swank A, Sobus JR, Howey XM, Schmid J, Catron T, McCord J, Hines E, Strynar M, Tal T.
1003   Evaluation of Developmental Toxicity, Developmental Neurotoxicity, and Tissue Dose in Zebrafish
1004   Exposed to GenX and Other PFAS. Environ Health Perspect. 2020 Apr;128(4):47005. Epub 2020/04/10.
1005   doi:10.1289/ehp5843. Cited in: Pubmed; PMID 32271623.

1006

1007   49. Huang H, Huang C, Wang L, Ye X, Bai C, Simonich MT, Tanguay RL, Dong Q. Toxicity, uptake kinetics
1008   and behavior assessment in zebrafish embryos following exposure to perfluorooctanesulphonicacid
1009   (PFOS). Aquat Toxicol. 2010 Jun 10;98(2):139-47. Epub 2010/02/23. doi:10.1016/j.aquatox.2010.02.003.
1010   Cited in: Pubmed; PMID 20171748.

1011

1012   50. Jantzen CE, Annunziato KM, Cooper KR. Behavioral, morphometric, and gene expression effects in
1013   adult zebrafish (Danio rerio) embryonically exposed to PFOA, PFOS, and PFNA. Aquat Toxicol. 2016
1014   Nov;180:123-130. Epub 2016/10/07. doi:10.1016/j.aquatox.2016.09.011. Cited in: Pubmed; PMID
1015   27710860.

1016

1017 51. Liu X, Jin Y, Liu W, Wang F, Hao S. Possible mechanism of perfluorooctane sulfonate and
1018 perfluorooctanoate on the release of calcium ion from calcium stores in primary cultures of rat
1019 hippocampal neurons. Toxicol In Vitro. 2011 Oct;25(7):1294-301. Epub 2011/05/18.
1020 doi:10.1016/j.tiv.2011.04.016. Cited in: Pubmed; PMID 21575708.

1021

1022 52. Christou M, Fraser TWK, Berg V, Ropstad E, Kamstra JH. Calcium signaling as a possible mechanism
1023 behind increased locomotor response in zebrafish larvae exposed to a human relevant persistent
1024 organic pollutant mixture or PFOS. Environ Res. 2020 Aug;187:109702. Epub 2020/06/01.
1025 doi:10.1016/j.envres.2020.109702. Cited in: Pubmed; PMID 32474314.

1026

1027 53. Ellett F, Pase L, Hayman JW, Andrianopoulos A, Lieschke GJ. mpeg1 promoter transgenes direct
1028 macrophage-lineage expression in zebrafish. Blood. 2011;117(4). doi:10.1182/blood-2010-10-314120.

1029

1030 54. Arrenberg AB, Del Bene F, Baier H. Optical control of zebrafish behavior with halorhodopsin. Proc
1031 Natl Acad Sci U S A. 2009 Oct 20;106(42):17968-73. Epub 2009/10/07. doi:10.1073/pnas.0906252106.
1032 Cited in: Pubmed; PMID 19805086.

1033

1034 55. Gradinaru V, Zhang F, Ramakrishnan C, Mattis J, Prakash R, Diester I, Goshen I, Thompson KR,
1035 Deisseroth K. Molecular and cellular approaches for diversifying and extending optogenetics. Cell. 2010
1036 Apr 2;141(1):154-165. Epub 2010/03/23. doi:10.1016/j.cell.2010.02.037. Cited in: Pubmed; PMID
1037 20303157.

1038

1039 56. Fosque BF, Sun Y, Dana H, Yang CT, Ohyama T, Tadross MR, Patel R, Zlatic M, Kim DS, Ahrens MB,
1040 Jayaraman V, Looger LL, Schreiter ER. Neural circuits. Labeling of active neural circuits in vivo with
1041 designed calcium integrators. Science. 2015 Feb 13;347(6223):755-60. Epub 2015/02/14.
1042 doi:10.1126/science.1260922. Cited in: Pubmed; PMID 25678659.

1043

1044 57. Kimura Y, Satou C, Higashijima S. V2a and V2b neurons are generated by the final divisions of pair-
1045 producing progenitors in the zebrafish spinal cord. Development. 2008 Sep;135(18):3001-5. eng. Epub
1046 20080806. doi:10.1242/dev.024802. Cited in: Pubmed; PMID 18684740.

1047

1048 58. Sato T, Takahoko M, Okamoto H. HuC:Kaede, a useful tool to label neural morphologies in networks
1049 in vivo. Genesis. 2006 Mar;44(3):136-42. Epub 2006/02/24. doi:10.1002/gene.20196. Cited in: Pubmed;
1050 PMID 16496337.

1051

1052 59. Davison JM, Akitake CM, Goll MG, Rhee JM, Gosse N, Baier H, Halpern ME, Leach SD, Parsons MJ.
1053 Transactivation from Gal4-VP16 transgenic insertions for tissue-specific cell labeling and ablation in
1054 zebrafish. Dev Biol. 2007 Apr 15;304(2):811-24. Epub 2007/03/06. doi:10.1016/j.ydbio.2007.01.033.
1055 Cited in: Pubmed; PMID 17335798.

1056

1057 60. Shiau CE, Kaufman Z, Meireles AM, Talbot WS. Differential requirement for irf8 in formation of
1058 embryonic and adult macrophages in zebrafish. PLoS One. 2015;10(1):e0117513. Epub 2015/01/24.
1059 doi:10.1371/journal.pone.0117513. Cited in: Pubmed; PMID 25615614.

1060

1061 61. Sant KE, Jacobs HM, Borofski KA, Moss JB, Timme-Laragy AR. Embryonic exposures to
1062 perfluorooctanesulfonic acid (PFOS) disrupt pancreatic organogenesis in the zebrafish, Danio rerio.
1063 Environ Pollut. 2017 Jan;220(Pt B):807-817. Epub 20161031. doi:10.1016/j.envpol.2016.10.057. Cited in:
1064 Pubmed; PMID 27810111.

1065

1066 62. Chen J, Tanguay RL, Tal TL, Gai Z, Ma X, Bai C, Tilton SC, Jin D, Yang D, Huang C, Dong Q. Early life
1067 perfluorooctanesulphonic acid (PFOS) exposure impairs zebrafish organogenesis. Aquat Toxicol. 2014
1068 May;150:124-32. Epub 20140312. doi:10.1016/j.aquatox.2014.03.005. Cited in: Pubmed; PMID
1069 24667235.

1070

1071 63. Kishimoto N, Shimizu K, Sawamoto K. Neuronal regeneration in a zebrafish model of adult brain
1072 injury. Dis Model Mech. 2012 Mar;5(2):200-9. Epub 2011/10/27. doi:10.1242/dmm.007336. Cited in:
1073 Pubmed; PMID 22028327.

1074

1075 64. Wallace CK, Bright LA, Marx JO, Andersen RP, Mullins MC, Carty AJ. Effectiveness of Rapid Cooling as
1076 a Method of Euthanasia for Young Zebrafish (Danio rerio). J Am Assoc Lab Anim Sci. 2018 Jan 1;57(1):58-
1077 63. Epub 2018/02/07. Cited in: Pubmed; PMID 29402353.

1078

1079 65. Delic J, Coppey J, Magdelenat H, Coppey-Moisan M. Impossibility of acridine orange intercalation in
1080 nuclear DNA of the living cell. Exp Cell Res. 1991 May;194(1):147-53. eng. doi:10.1016/0014-
1081 4827(91)90144-j. Cited in: Pubmed; PMID 2015848.

1082

1083 66. Yu T, Park Y, Johnson JM, Jones DP. apLCMS--adaptive processing of high-resolution LC/MS data.
1084 Bioinformatics. 2009 Aug 1;25(15):1930-6. Epub 2009/05/06. doi:10.1093/bioinformatics/btp291. Cited
1085 in: Pubmed; PMID 19414529.

1086

1087 67. Uppal K, Soltow QA, Strobel FH, Pittard WS, Gernert KM, Yu T, Jones DP. xMSanalyzer: automated
1088 pipeline for improved feature detection and downstream analysis of large-scale, non-targeted
1089 metabolomics data. BMC Bioinformatics. 2013 Jan 16;14:15. Epub 2013/01/18. doi:10.1186/1471-2105-
1090 14-15. Cited in: Pubmed; PMID 23323971.

1091

1092 68. Chong J, Xia J. MetaboAnalystR: an R package for flexible and reproducible analysis of metabolomics
1093 data. Bioinformatics. 2018 Dec 15;34(24):4313-4314. Epub 2018/06/30.
1094 doi:10.1093/bioinformatics/bty528. Cited in: Pubmed; PMID 29955821.

1095

1096   69. Badimon A, Strasburger HJ, Ayata P, Chen X, Nair A, Ikegami A, Hwang P, Chan AT, Graves SM, Uweru
1097   JO, Ledderose C, Kutlu MG, Wheeler MA, Kahan A, Ishikawa M, Wang Y-C, Loh Y-HE, Jiang JX, Surmeier
1098   DJ, Robson SC, Junger WG, Sebra R, Calipari ES, Kenny PJ, Eyo UB, Colonna M, Quintana FJ, Wake H,
1099   Gradinaru V, Schaefer A. Negative feedback control of neuronal activity by microglia. Nature.
1100   2020;586(7829). doi:10.1038/s41586-020-2777-8.

1101

1102   70. Davalos D, Grutzendler J, Yang G, Kim JV, Zuo Y, Jung S, Littman DR, Dustin ML, Gan WB. ATP
1103   mediates rapid microglial response to local brain injury in vivo. Nat Neurosci. 2005 Jun;8(6):752-8. Epub
1104   2005/05/17. doi:10.1038/nn1472. Cited in: Pubmed; PMID 15895084.

1105

1106   71. Yu T, Zhang X, Shi H, Tian J, Sun L, Hu X, Cui W, Du D. P2Y12 regulates microglia activation and
1107   excitatory synaptic transmission in spinal lamina II neurons during neuropathic pain in rodents. Cell
1108   Death Dis. 2019 Feb 18;10(3):165. Epub 2019/02/20. doi:10.1038/s41419-019-1425-4. Cited in: Pubmed;
1109   PMID 30778044.

1110

1111   72. Schmidt R, Beil T, Strahle U, Rastegar S. Stab wound injury of the zebrafish adult telencephalon: a
1112   method to investigate vertebrate brain neurogenesis and regeneration. J Vis Exp. 2014 Aug
1113   4;(90):e51753. Epub 2014/08/26. doi:10.3791/51753. Cited in: Pubmed; PMID 25146302.

1114

1115   73. Sieger D, Moritz C, Ziegenhals T, Prykhozhij S, Peri F. Long-range Ca2+ waves transmit brain-damage
1116   signals to microglia. Dev Cell. 2012 Jun 12;22(6):1138-48. Epub 2012/05/29.
1117   doi:10.1016/j.devcel.2012.04.012. Cited in: Pubmed; PMID 22632801.

1118

1119   74. Donat CK, Scott G, Gentleman SM, Sastre M. Microglial Activation in Traumatic Brain Injury. Front
1120   Aging Neurosci. 2017;9:208. Epub 2017/07/14. doi:10.3389/fnagi.2017.00208. Cited in: Pubmed; PMID
1121   28701948.

1122

1123   75. Newell EW, Schlichter LC. Integration of K+ and Cl- currents regulate steady-state and dynamic
1124   membrane potentials in cultured rat microglia. J Physiol. 2005 Sep 15;567(Pt 3):869-90. Epub
1125   2005/07/16. doi:10.1113/jphysiol.2005.092056. Cited in: Pubmed; PMID 16020460.

1126

1127   76. Baraban SC, Taylor MR, Castro PA, Baier H. Pentylenetetrazole induced changes in zebrafish
1128   behavior, neural activity and c-fos expression. Neuroscience. 2005;131(3):759-68. Epub 2005/02/26.
1129   doi:10.1016/j.neuroscience.2004.11.031. Cited in: Pubmed; PMID 15730879.

1130

1131   77. Colwill RM, Creton R. Imaging escape and avoidance behavior in zebrafish larvae. Rev Neurosci.
1132   2011;22(1):63-73. Epub 2011/05/17. doi:10.1515/RNS.2011.008. Cited in: Pubmed; PMID 21572576.

1133

1134   78. Mathuru AS, Jesuthasan S. The medial habenula as a regulator of anxiety in adult zebrafish. Front
1135   Neural Circuits. 2013;7:99. Epub 2013/06/12. doi:10.3389/fncir.2013.00099. Cited in: Pubmed; PMID
1136   23750127.

1137
1138   79. Okamoto H, Agetsuma M, Aizawa H. Genetic dissection of the zebrafish habenula, a possible
1139   switching board for selection of behavioral strategy to cope with fear and anxiety. Dev Neurobiol. 2012
1140   Mar;72(3):386-94. Epub 2011/05/14. doi:10.1002/dneu.20913. Cited in: Pubmed; PMID 21567982.

1141
1142   80. Marinelli S, Basilico B, Marrone MC, Ragozzino D. Microglia-neuron crosstalk: Signaling mechanism
1143   and control of synaptic transmission. Semin Cell Dev Biol. 2019 Oct;94:138-151. Epub 2019/05/22.
1144   doi:10.1016/j.semcdb.2019.05.017. Cited in: Pubmed; PMID 31112798.

1145
1146   81. Trede NS, Langenau DM, Traver D, Look AT, Zon LI. The use of zebrafish to understand immunity.
1147   Immunity. 2004 Apr;20(4):367-79. Epub 2004/04/16. doi:10.1016/s1074-7613(04)00084-6. Cited in:
1148   Pubmed; PMID 15084267.

1149
1150   82. Galindo-Villegas J. Recent findings on vertebrate developmental immunity using the zebrafish model.
1151   Mol Immunol. 2016 Jan;69:106-12. Epub 2015/11/22. doi:10.1016/j.molimm.2015.10.011. Cited in:
1152   Pubmed; PMID 26589453.

1153
1154   83. Levraud JP, Rawls JF, Clatworthy AE. Using zebrafish to understand reciprocal interactions between
1155   the nervous and immune systems and the microbial world. J Neuroinflammation. 2022 Jun 28;19(1):170.
1156   Epub 2022/06/29. doi:10.1186/s12974-022-02506-x. Cited in: Pubmed; PMID 35765004.

1157
1158   84. DeWitt JC, Blossom SJ, Schaider LA. Exposure to per-fluoroalkyl and polyfluoroalkyl substances leads
1159   to immunotoxicity: epidemiological and toxicological evidence. J Expo Sci Environ Epidemiol. 2019
1160   Mar;29(2):148-156. Epub 2018/11/30. doi:10.1038/s41370-018-0097-y. Cited in: Pubmed; PMID
1161   30482935.

1162
1163   85. Mollenhauer MA, Bradshaw SG, Fair PA, McGuinn WD, Peden-Adams MM. Effects of
1164   perfluorooctane sulfonate (PFOS) exposure on markers of inflammation in female B6C3F1 mice. J
1165   Environ Sci Health A Tox Hazard Subst Environ Eng. 2011;46(2):97-108. Epub 2010/12/21.
1166   doi:10.1080/10934529.2011.532418. Cited in: Pubmed; PMID 21170772.

1167
1168   86. Hou YF, Shan C, Zhuang SY, Zhuang QQ, Ghosh A, Zhu KC, Kong XK, Wang SM, Gong YL, Yang YY, Tao
1169   B, Sun LH, Zhao HY, Guo XZ, Wang WQ, Ning G, Gu YY, Li ST, Liu JM. Gut microbiota-derived propionate
1170   mediates the neuroprotective effect of osteocalcin in a mouse model of Parkinson's disease.
1171   Microbiome. 2021 Jan 31;9(1):34. Epub 20210131. doi:10.1186/s40168-020-00988-6. Cited in: Pubmed;
1172   PMID 33517890.

1173

1174   87. Clark R, Kupper T. Old meets new: the interaction between innate and adaptive immunity. J Invest
1175   Dermatol. 2005 Oct;125(4):629-37. Epub 2005/09/28. doi:10.1111/j.0022-202X.2005.23856.x. Cited in:
1176   Pubmed; PMID 16185260.

1177
1178   88. Ge J, Wang C, Nie X, Yang J, Lu H, Song X, Su K, Li T, Han J, Zhang Y, Mao J, Gu Y, Zhao J, Jiang S, Wu
1179   Q. ROS-mediated apoptosis of HAPI microglia through p53 signaling following PFOS exposure. Environ
1180   Toxicol Pharmacol. 2016 Sep;46:9-16. eng. Epub 2016/06/27. doi:10.1016/j.etap.2016.06.025. Cited in:
1181   Pubmed; PMID 27414741.

1182
1183   89. Zhang L, Li YY, Zeng HC, Li M, Wan YJ, Schluesener HJ, Zhang ZY, Xu SQ. Perfluorooctane sulfonate
1184   induces apoptosis in N9 microglial cell line. Int J Toxicol. 2011 Mar;30(2):207-15. Epub 2010/12/01.
1185   doi:10.1177/1091581810387832. Cited in: Pubmed; PMID 21115943.

1186
1187   90. Goldfarb S, Fainstein N, Ganz T, Vershkov D, Lachish M, Ben-Hur T. Electric neurostimulation
1188   regulates microglial activation via retinoic acid receptor alpha signaling. Brain Behav Immun. 2021
1189   Aug;96:40-53. Epub 2021/05/15. doi:10.1016/j.bbi.2021.05.007. Cited in: Pubmed; PMID 33989746.

1190
1191   91. Szalay G, Martinecz B, Lenart N, Kornyei Z, Orsolits B, Judak L, Csaszar E, Fekete R, West BL, Katona
1192   G, Rozsa B, Denes A. Microglia protect against brain injury and their selective elimination dysregulates
1193   neuronal network activity after stroke. Nat Commun. 2016 May 3;7:11499. Epub 2016/05/04.
1194   doi:10.1038/ncomms11499. Cited in: Pubmed; PMID 27139776.

1195
1196   92. Liu M, Jiang L, Wen M, Ke Y, Tong X, Huang W, Chen R. Microglia depletion exacerbates acute
1197   seizures and hippocampal neuronal degeneration in mouse models of epilepsy. Am J Physiol Cell Physiol.
1198   2020 Sep;319(3):C605-C610. eng. Epub 2020/08/12. doi:10.1152/ajpcell.00205.2020. Cited in: Pubmed;
1199   PMID 32783655.

1200
1201   93. Tynan RJ, Naicker S, Hinwood M, Nalivaiko E, Buller KM, Pow DV, Day TA, Walker FR. Chronic stress
1202   alters the density and morphology of microglia in a subset of stress-responsive brain regions. Brain
1203   Behav Immun. 2010 Oct;24(7):1058-68. Epub 2010/02/16. doi:10.1016/j.bbi.2010.02.001. Cited in:
1204   Pubmed; PMID 20153418.

1205
1206   94. Henstridge CM, Tzioras M, Paolicelli RC. Glial Contribution to Excitatory and Inhibitory Synapse Loss
1207   in Neurodegeneration. Front Cell Neurosci. 2019;13:63. Epub 2019/03/14.
1208   doi:10.3389/fncel.2019.00063. Cited in: Pubmed; PMID 30863284.

1209
1210   95. Silva NJ, Dorman LC, Vainchtein ID, Horneck NC, Molofsky AV. In situ and transcriptomic
1211   identification of microglia in synapse-rich regions of the developing zebrafish brain. Nat Commun. 2021
1212   Oct 8;12(1):5916. Epub 2021/10/10. doi:10.1038/s41467-021-26206-x. Cited in: Pubmed; PMID
1213   34625548.

1214
96. Liu X, Liu W, Jin Y, Yu W, Wang F, Liu L. Effect of gestational and lactational exposure to perfluorooctanesulfonate on calcium-dependent signaling molecules gene expression in rats' hippocampus. Arch Toxicol. 2010 Jan;84(1):71-9. Epub 2009/09/17. doi:10.1007/s00204-009-0467-2. Cited in: Pubmed; PMID 19756518.

1219
97. Liao CY, Li XY, Wu B, Duan S, Jiang GB. Acute enhancement of synaptic transmission and chronic inhibition of synaptogenesis induced by perfluorooctane sulfonate through mediation of voltage-dependent calcium channel. Environ Sci Technol. 2008 Jul 15;42(14):5335-41. Epub 2008/08/30. doi:10.1021/es800018k. Cited in: Pubmed; PMID 18754390.

1224
98. Armada-Moreira A, Gomes JI, Pina CC, Savchak OK, Goncalves-Ribeiro J, Rei N, Pinto S, Morais TP, Martins RS, Ribeiro FF, Sebastiao AM, Crunelli V, Vaz SH. Going the Extra (Synaptic) Mile: Excitotoxicity as the Road Toward Neurodegenerative Diseases. Front Cell Neurosci. 2020;14:90. Epub 2020/05/12. doi:10.3389/fncel.2020.00090. Cited in: Pubmed; PMID 32390802.

1229
99. Olloquequi J, Cornejo-Cordova E, Verdaguer E, Soriano FX, Binvignat O, Auladell C, Camins A. Excitotoxicity in the pathogenesis of neurological and psychiatric disorders: Therapeutic implications. J Psychopharmacol. 2018 Mar;32(3):265-275. Epub 2018/02/16. doi:10.1177/0269881118754680. Cited in: Pubmed; PMID 29444621.

1234
100. Rosen MB, Das KP, Rooney J, Abbott B, Lau C, Corton JC. PPARalpha-independent transcriptional targets of perfluoroalkyl acids revealed by transcript profiling. Toxicology. 2017 Jul 15;387:95-107. Epub 2017/06/01. doi:10.1016/j.tox.2017.05.013. Cited in: Pubmed; PMID 28558994.

1238
101. Takacs ML, Abbott BD. Activation of mouse and human peroxisome proliferator-activated receptors (alpha, beta/delta, gamma) by perfluorooctanoic acid and perfluorooctane sulfonate. Toxicol Sci. 2007 Jan;95(1):108-17. Epub 2006/10/19. doi:10.1093/toxsci/kfl135. Cited in: Pubmed; PMID 17047030.

1242
102. Villapol S. Roles of Peroxisome Proliferator-Activated Receptor Gamma on Brain and Peripheral Inflammation. Cell Mol Neurobiol. 2018 Jan;38(1):121-132. Epub 2017/10/05. doi:10.1007/s10571-017-0554-5. Cited in: Pubmed; PMID 28975471.

1246
103. Fei C, McLaughlin JK, Lipworth L, Olsen J. Prenatal exposure to perfluorooctanoate (PFOA) and perfluorooctanesulfonate (PFOS) and maternally reported developmental milestones in infancy. Environ Health Perspect. 2008;116.

1250
104. Sato I, Kawamoto K, Nishikawa Y, Tsuda S, Yoshida M, Yaegashi K, Saito N, Liu W, Jin Y. Neurotoxicity of perfluorooctane sulfonate (PFOS) in rats and mice after single oral exposure. J Toxicol

Sci. 2009 Oct;34(5):569-74. Epub 2009/10/03. doi:10.2131/jts.34.569. Cited in: Pubmed; PMID 19797866.

105. Randlett O, Wee CL, Naumann EA, Nnaemeka O, Schoppik D, Fitzgerald JE, Portugues R, Lacoste AM, Riegler C, Engert F, Schier AF. Whole-brain activity mapping onto a zebrafish brain atlas. Nat Methods. 2015 Nov;12(11):1039-46. Epub 2016/01/19. doi:10.1038/nmeth.3581. Cited in: Pubmed; PMID 26778924.

106. Kalueff AV, Stewart AM. Zebrafish protocols for neurobehavioral research. New York: Humana Press; 2012. ISBN: 9781617795961 (cased)

9781617795978 (eBook). 1850-9999 http://www.springer.com/gb/ BLDSS.

107. Ahmad F, Richardson MK. Exploratory behaviour in the open field test adapted for larval zebrafish: impact of environmental complexity. Behav Processes. 2013 Jan;92:88-98. Epub 2012/11/06. doi:10.1016/j.beproc.2012.10.014. Cited in: Pubmed; PMID 23123970.

108. Richendrfer H, Pelkowski SD, Colwill RM, Creton R. On the edge: pharmacological evidence for anxiety-related behavior in zebrafish larvae. Behav Brain Res. 2012 Mar 1;228(1):99-106. Epub 2011/12/14. doi:10.1016/j.bbr.2011.11.041. Cited in: Pubmed; PMID 22155488.

109. Skogheim TS, Weyde KVF, Aase H, Engel SM, Suren P, Oie MG, Biele G, Reichborn-Kjennerud T, Brantsaeter AL, Haug LS, Sabaredzovic A, Auyeung B, Villanger GD. Prenatal exposure to per- and polyfluoroalkyl substances (PFAS) and associations with attention-deficit/hyperactivity disorder and autism spectrum disorder in children. Environ Res. 2021 Nov;202:111692. Epub 2021/07/23. doi:10.1016/j.envres.2021.111692. Cited in: Pubmed; PMID 34293314.

110. Oh J, Bennett DH, Calafat AM, Tancredi D, Roa DL, Schmidt RJ, Hertz-Picciotto I, Shin HM. Prenatal exposure to per- and polyfluoroalkyl substances in association with autism spectrum disorder in the MARBLES study. Environ Int. 2021 Feb;147:106328. Epub 2021/01/03. doi:10.1016/j.envint.2020.106328. Cited in: Pubmed; PMID 33387879.

111. Shin HM, Bennett DH, Calafat AM, Tancredi D, Hertz-Picciotto I. Modeled prenatal exposure to per- and polyfluoroalkyl substances in association with child autism spectrum disorder: A case-control study. Environ Res. 2020 Jul;186:109514. Epub 2020/05/01. doi:10.1016/j.envres.2020.109514. Cited in: Pubmed; PMID 32353786.

112. Fuentes S, Vicens P, Colomina MT, Domingo JL. Behavioral effects in adult mice exposed to perfluorooctane sulfonate (PFOS). Toxicology. 2007 Dec 5;242(1-3):123-9. Epub 2007/10/24. doi:10.1016/j.tox.2007.09.012. Cited in: Pubmed; PMID 17950980.

1292

1293   113. Browne CA, Hammack R, Lucki I. Dysregulation of the Lateral Habenula in Major Depressive
1294   Disorder. Front Synaptic Neurosci. 2018;10:46. Epub 2018/12/26. doi:10.3389/fnsyn.2018.00046. Cited
1295   in: Pubmed; PMID 30581384.

1296

1297   114. Post GB, Cohn PD, Cooper KR. Perfluorooctanoic acid (PFOA), an emerging drinking water
1298   contaminant: a critical review of recent literature. Environ Res. 2012 Jul;116:93-117. Epub 2012/05/09.
1299   doi:10.1016/j.envres.2012.03.007. Cited in: Pubmed; PMID 22560884.

1300

1301   115. Rericha Y, Cao D, Truong L, Simonich M, Field JA, Tanguay RL. Behavior Effects of Structurally
1302   Diverse Per- and Polyfluoroalkyl Substances in Zebrafish. Chem Res Toxicol. 2021 Jun 21;34(6):1409-
1303   1416. Epub 2021/05/22. doi:10.1021/acs.chemrestox.1c00101. Cited in: Pubmed; PMID 34018735.

1304

1305   116. Tukker AM, Bouwman LMS, van Kleef R, Hendriks HS, Legler J, Westerink RHS. Perfluorooctane
1306   sulfonate (PFOS) and perfluorooctanoate (PFOA) acutely affect human alpha1beta2gamma2L GABAA
1307   receptor and spontaneous neuronal network function in vitro. Sci Rep. 2020 Mar 24;10(1):5311. Epub
1308   2020/03/27. doi:10.1038/s41598-020-62152-2. Cited in: Pubmed; PMID 32210279.

1309

1310 **FIGURES**

1311

Figure 1. Microglia morphology and response to minor brain injury in 3 dpf larvae exposed to 28 µM
PFOS. (A) Exposure paradigm: zebrafish embryos from *Tg(mpeg1:EGFP)* adults were collected after a 1-
hour timed spawn. At 4 hpf, embryos were dosed with 0.1% DMSO (Control) or 28 µM PFOS. Treatment
solutions were static and not renewed. Embryos were dechorionated at 24 hpf, and imaging and injury
experiments were conducted at 3 dpf. (B) Schematic of larval brain injury: a pulled glass needle (OD 9 um)
was used to puncture the right telencephalon of larval zebrafish. (C) Hypothetical schematic of microglia
response 4 hours post-injury (hpi). (D-E") Homeostatic, non-activated microglia of 3 dpf control larvae have
several projections emanating from their cell bodies. (F-H) Control microglia respond to brain injury, with
visible accumulation at the injury site at 4 hpi that mostly clears by 12 hpi. (I-J") Microglia of 3 dpf larvae
exposed to 28 µM PFOS become phenotypically activated, with rounder cell bodies and reduced
projections. (K-M) In PFOS-exposed larvae, microglia robustly respond to brain injury by 4 hpi and are still
maintained at the site at 12 hpi. Quantifications of the images reveal that PFOS-exposed microglia (N) area
(P = 0.0342) and (O) cell perimeter (P = 0.0006) are significantly reduced. (P) The perimeter-to-area ratio
(P = 0.0002) is also significantly decreased, indicative of a rounder cell morphology. n = 15 fish per group
(3-18 cells counted per fish). (Q) qRT-PCR for the microglia activation gene, *p2ry12*, was significantly
upregulated in isolated heads of PFOS-exposed larvae (P = 0.0211; n = 10 pooled heads per sample).
PFOS exposure resulted in a significant increase in microglia recruitment in the first 4.5 hpi (P = 0.0208; n
= 5-6 per group), as well as increased response area at (S) 4 hpi (P = 0.0037; n = 19-20 per group) and (T)
12 hpi (P = 0.0467; n = 4 per group). Confocal micrographs at 40x magnification (D-E" & I-J") or 20x
magnification (F-H; K-M). Box plot limits represent 25th to 27th percentile, with the midline representing the
median. See Table S5 for additional statistical details.

1333

Figure 2. Optogenetic modulation of PFOS-exposed microglia. (A) Schematic of halorhodopsin:
Optogenetic modulation of microglia electrical state is achieved via photo-stimulation of the light-gated
chloride pump, halorhodopsin (eNpHR3.0). eNpHR3.0 is most responsive to 589 nm wavelength light. (B)
eNpHR3.0 was driven under a pan-macrophage promoter (*Tg(mpeg1:GalFF;UAS:eNpHR3.0-mCherry)*) to
achieve optogenetic control of microglia in zebrafish larvae. (C) Experimental paradigm: At 72 hpf, injured
or uninjured zebrafish were stimulated for 4 hours with 589 nm light in the enclosed Noldus DanioVision
Behavior Unit. (D-E") Unstimulated halorhodopsin+ microglia of 3 dpf 28 µM PFOS-exposed larvae were
rounded with few projections. As shown in Figure 1, (F) unstimulated control microglia were responsive to
minor injury at 4 hpi, though (G) unstimulated PFOS-exposed microglia had a significantly heightened
response. (H-I") Following 4-hour stimulation of halorhodopsin+ in PFOS-exposed larvae, microglia became
more ramified. (J, K) Stimulation of halorhodopsin+ microglia following PFOS exposure also normalized the
microglia response to injury. (L) Stimulated microglia area was unchanged, but (M) they had significantly
increased cell perimeter (*P* = 0.007) and (N) an increased perimeter-to-area ratio (*P* < 0.0001), indicative
of a more ramified cell shape (n = 40-42 cells per group from 3 independent experiments). (O) The injury
response area was significantly increased between unstimulated control and 28 µM PFOS exposed larvae
(*P* = 0.0306). However, there was no significant difference in response area between the 4-hour stimulated

1350   control and PFOS-exposed larvae. n = 9-15 per group. Box plot limits represent 25^{th} to 27^{th} percentile, with
1351   the midline representing the median. See Table S5 for additional statistical details.
1352

1353

1354

Figure 3. Analysis of regional and global neuronal network activity following chronic exposure to PFOS. (A) Illustrative representation of a larval zebrafish brain with anatomical regions outlined: forebrain (FB), optic tectum (OT), cerebellum (Ce), hindbrain (HB), and whole brain (WB). (B) Illustrative representation of neuron-driven CaMPARI: neurons with low intracellular calcium (low Ca^{2+}) remain green following 1 minute exposure to 405 nm light, while neurons with high intracellular calcium (high Ca^{2+}) are photoconverted to red. (C) Confocal micrograph of a 3 dpf larvae expressing neuron-specific CaMPARI (*Tg(elavl1:CaMPARI)*) following 1 minute photoconversion. (D) Generated high intensity LUTs heat map of the red, photoconverted channel in C depicting high activity neurons. Low to high Intracellular calcium is depicted by a blue-red-white spectrum. (E) Micrographs of active neurons at 3 dpf in control, (F) 7 µM PFOS, and (G) 28 µM PFOS larvae following 1-minute photoconversion. Neuronal activity can be quantified by determining the ratio of fluorescent intensity in the red versus green channels (F_{Red}/F_{Green}). (H) 7 µM PFOS does not result in a regional or global (WB) change in neuron activity in 3 dpf larvae. (J) Brain activity was also determined in 5 dpf larvae exposed to control, (K) 7 µM PFOS, or (L) 14 µM PFOS. At 5 dpf (M) 7 µM PFOS-exposed larvae have significant increases in brain activity in the OT ($P = 0.0058$) and near significance globally ($P = 0.0788$). (N) Larvae exposed to 14 µM PFOS also had increases in the OT ($P = 0.0005$), Ce ($P = 0.0010$), HB ($P = 0.0074$), and globally ($P = 0.0025$). Confocal micrographs at 10x magnification. n = 21-23 fish per group. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

1373

1374

1375

Figure 4. Assessing anxiety-like swim behavior in PFOS-exposed larvae. (A) Larval swim behavior during a 30-minute light/dark behavioral assay was used to determine time spent in the well's center (crossing center; less anxious) versus time spent along the well's edge (center avoidance; more anxious). (B) Heat maps were generated to indicate mobile (blue) versus stationary (red) swim activity within the well, as well as the zones traversed within the well. (C) At 3 dpf, larvae exposed to PFOS spent less time in the center relative to the corresponding control group (7 µM, -49.5%, P = 0.2994; 14 µM, -66.5%, P = 0.0021; 28 µM, -55.6%, P = 0.6682). (D) At 5 dpf, both 7 µM PFOS (-26.7%; P = 0.0288) and 14 µM PFOS (-57.5%; P < 0.0001) spent significantly less time in the center of the well. n = 76-101 fish per group. Error bars represent SEM. See Table S5 for additional statistical details.

1386

1387

Figure 5. Light/dark behavioral assay and neuronal activity in microglia mutants exposed to PFOS.
Wildtype and microglia-deficient *irf8*^st96/st96^ larvae dosed with either control or 7 μM PFOS were assessed
for potential changes in swim behavior and neuronal activity. (A) Light/dark behavioral assay of 5 dpf
wildtype or *irf8* mutant control or PFOS-exposed larvae. (B) Assessing the distance moved following the
first minute of the second light "off" cycle (i.e., minute 15 of the assay), control-treated *irf8* mutant larvae
responded significantly less than wildtype controls (*P* = 0.0237). While not significant, PFOS-exposed *irf8*
mutants had a heightened response to the first minute of the dark cycle compared to PFOS-exposed
wildtype larvae. PFOS mutants also had twice the percent recovery in swim activity by the last minute of
the assay (i.e., minute 15 versus minute 30). (C-G) PFOS-exposed macrophage mutants did not show a
significant change in regional or global neuronal activity compared to PFOS-exposed wildtype larvae. (C)
Forebrain, control-treated wildtype versus PFOS-treated wildtype (*P* = 0.0060); control-treated mutant
versus PFOS-treated mutant (*P* = 0.1176). (D) Optic tectum, control-treated wildtype versus PFOS-treated
wildtype (*P* = 0.0008); control-treated mutant versus PFOS-treated mutant (*P* = 0.0867). (E) Cerebellum,
control-treated wildtype versus PFOS-treated wildtype (*P* = 0.0020); control-treated mutant versus PFOS-
treated mutant (*P* = 0.1333). (F) Hindbrain, control-treated wildtype versus PFOS-treated wildtype (*P* <
0.0001); control-treated mutant versus PFOS-treated mutant (*P* = 0.0100). (G) Whole brain, control-treated
wildtype versus PFOS-treated wildtype (*P* = 0.0001); control-treated mutant versus PFOS-treated mutant
(*P* = 0.0349). There were no significant differences in regional or global PFOS-induced neuronal
hyperactivity between the wildtype or mutant larvae expressing *Tg*(*elavl3:CaMPARI*). n = 14-22 per group
for behavior; n = 13-16 per group for neuroimaging. Box plot limits represent 25th to 27th percentile, with the
midline representing the median. See Table S5 for additional statistical details.

1409

1410

1411

Figure 6. Microglia response to brain injury following optogenetic silencing of neurons in PFOS-
exposed larvae. Larvae expressing *Tg(elavl3:Gal4;cryaa:RFP;UAS:eNpHR3.0;mpeg1:EGFP)* were dosed
with (A-D) control or (E-H) 28 µM PFOS and injured at the right telencephalon at 3 dpf and imaged 4 hpi.
(E&F) PFOS-exposed unstimulated larvae had a 33.5% increase in microglia response compared to
controls (*P* = 0.0879). (C&D) Optogenetic stimulation of halorhodopsin with 570 nm light for 4 hpi did not
affect microglia response to injury in control larvae (6.6% decrease). (G&H) However, neuronal silencing in
PFOS-exposed larvae normalized the microglia response, such that the response was equivalent to the
stimulated controls. (I) Quantification of microglia response area. n = 14-20 per group. Box plot limits
represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional
statistical details.

1422

1423

1424

Figure 7. Neuronal activity and microglia response to brain injury following exposure to PFOA, a known immunotoxic congener. PFOA is an 8-carbon PFAS compound with a carboxylate head group. To observe the effects of PFOA on neuronal activity, we again used larvae of the Tg(*elavl3:CaMPARI*) background. (A) At 5 dpf, control and (B) 64 μM PFOA-exposed larvae show no significant changes in regional or global neuronal activity, quantified in (C). (D) Larvae expressing Tg*(mpeg1:EGFP)* were also dosed with control solution or (E) 64 μM PFOA to determine if this non-excitatory PFAS compound resulted in microglia response differences at 3dpf. (F) At 4 hpi, 64 μM PFOA-exposed larvae did not have a significant change in microglia response to injury compared to sibling controls. n = 8-9 per group. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

1435

1436

1437

1438

1439

1440

1441

Evaluation of neural regulation and microglial responses to brain injury in larval zebrafish exposed to perfluorooctane sulfonate

Shannon E. Paquette[1#], Nathan R. Martin[1#], April Rodd[1], Katherine E. Manz[2], Manuel Camarillo[1], Eden Allen[1], Hannah I. Weller[3], Kurt Pennell[2], and Jessica S. Plavicki[1]

[1]Department of Pathology & Laboratory Medicine, Brown University, Providence, RI, USA

[2]School of Engineering, Brown University, Providence, RI, USA

[3]Department of Ecology, Evolution, and Organismal Biology, Brown University, Providence, RI, USA

[#]These authors contributed equally

Table of Contents

neurons with high intracellular calcium from green to red. (B-D) We modified our Noldus DanioVision Behavioral unit, outfitted with LEDs for optogenetic manipulation, to efficiently photoconvert CaMPARI in free-swimming zebrafish. (B) A 3D printed pedestal is used to reduce the working distance between the light source and zebrafish larvae. Applying the inverse-square law, the light intensity at the apex of the pedestal was calculated to be 135 mW/cm². (C) Illustration of the apex of the pedestal containing a single well dish (OD 15 mm). (D) The 405 nm light has an intensity of 5700 mW at the source. The light travels horizontally then is reflected off a dichroic mirror within the unit. The working distance was calculated as the distance from the light source to the dichroic mirror (α) plus the distance from the dichroic mirror to the apex of the pedestal (β). (E) Confocal micrographs of neuronal calcium following 1-minute photoconversion of CaMPARI in 3 dpf larvae treated with egg water or (E') after 10 minutes in 10 mM pentylenetetrazol (PTZ), a GABAA-inhibitor known to induce hyperactivity. (F, F') Green, non-active neurons with low intracellular calcium from E & E'. (G, G') Photoconverted, active neurons with high intracellular calcium, pseudo-labeled as magenta, from E & E'. (H, H') An intensity LUTs applied to the active neurons in G & G' spectrally maps the regional increases of neuronal activity on a scale from less active (blue) to more active (red-white). (I&J) Regional and global neuronal activity can be quantified by determining the ratio of fluorescent intensity in the red versus green channel (F_{Red}/F_{Green}). Exposure to 10 mM PTZ significantly increased brain activity in the optic tectum (OT; $P = 0.0496$) and trended toward increased activity in the cerebellum (Ce), hindbrain (HB), and whole brain (WB). Red/Green ratios were normalized relative to vehicle control. n = 16 fish per group. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S4. Regional and global brain morphology in larvae exposed to varying concentrations of PFOS. (A-D) Confocal micrographs of 3 dpf larvae or (J-L) 5 dpf larvae expressing *Tg(elavl3:CaMPARI)* exposed to (A&J) control, (B&K) 7 µM, (C&L) 14 µM, or (D) 28 µM PFOS. At 3 dpf, 7 µM PFOS larvae had a significant reduction in area of the (E) forebrain (Control vs 7 µM PFOS, $P = 0.0035$) and (F) optic tectum (Control vs 7 µM PFOS, $P = 0.0047$). (I) When measuring the area of the entire brain, only 7 µM and 14 µM PFOS exposure resulted in reduced global brain area (Control vs 7 µM & 14 µM, $P = 0.0174$ & $P = 0.0373$, respectively). At 5 dpf, there were no significant differences in regional area or global brain area with 7 µM nor 14 µM PFOS exposure. Confocal micrographs at 10x magnification. Control n = 52-54; treated n = 22-25. Scale bar = 100 um. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S5. MWAS of heads collected from control versus PFOS-exposed larvae. Heads were collected from 3 dpf control or 28 µM PFOS exposed larvae for an untargeted metabolome wide association study (MWAS). (A) Volcano plots of metabolites detected using a C18 column with negative ionization and (B) a HILIC column with positive ionization reveal several significantly down- or up-regulated metabolites in the PFOS-exposed brain (Green = downregulated; Blue = upregulated). Statistical significance was set at $P = 0.05$. (C) There are several significantly enriched pathways following both negative ionization (pink) and positive ionization (teal). The vertical line is at $P = 0.05$. Size of each dot represents the number of significant metabolites in that pathway. HILIC, hydrophilic interaction liquid chromatography column.

Figure S6. Light/dark behavioral assay in larvae exposed to varying concentrations of PFOS. (A) Exposure paradigm: at 4 hpf, zebrafish embryos were statically exposed to 0.1% DMSO (Control) or varying concentrations of PFOS (7, 14, or 28 µM). At 24 hpf, embryos were enzymatically dechorionated. Behavior was captured at 72, 96, and 120 hpf. (B) Light/dark behavioral assay: Zebrafish larvae were placed in individual wells of a 24-well plate containing the relevant toxicant solution 18-24 hours prior to the behavior assay to allow for well acclimation. A light/dark behavioral assay was performed using the Noldus DanioVision Behavioral Unit. The light/dark behavioral assay was as follows: 15-minute acclimation in the unit with the light off, 5 minutes with the light on, 5 minutes with the light off, 5 minutes with the light on, and

15 minutes with the light off. (C-H) At 3 dpf, larvae exposed to 28 µM PFOS had a significant increase in swim activity during the (F) light on cycles (P = 0.0003) and (H) over the course of the assay (P = 0.0123). (K) At 4 dpf, 7 µM PFOS larvae had a significant increase in swim distance when the light was on (P = 0.0277), as did 14 µM PFOS larvae (P = 0.0002). (L) At 4 dpf, 7 µM PFOS-exposed larvae also had a significant increase in swim behavior when the light was off (P = 0.0018) and (M) in total over the course of the assay (P < 0.0001). (P) At 5dpf, 14 µM PFOS had increased swim behavior when the light was on (P = 0.0006), and (Q) 7 µM PFOS larvae had increased swim behavior when the light was off (P < 0.0001). (R) Both 7 µM (P < 0.0001) and 14 µM PFOS (P = 0.0003) larvae had a significant increase in total distance moved. Control n = 130-190, treated n = 88-99. Error bars represent SEM. See Table S5 for additional statistical details.

Figure S7. Absolute and relative time spent in center throughout the light/dark behavioral response assay. (A) Total time spent in the center during the light on and light off cycles at 3dpf, control versus 7 µM (P = 0.0139, "OFF"). (B) Relative time spent in the center during the light on and light off cycles at 3 dpf, control versus 7 µM. (C) Total time spent in the center during the light on and light off cycles at 3 dpf, control versus 14 µM (P = 0.0075, "ON"; P < 0.0001, "OFF"). (D) Relative time spent in the center during the light on and light off cycles at 3 dpf, control versus 14 µM (P = 0.0015, "ON"; P = 0.0018, "OFF"). (E) Total time spent in the center during the light on and light off cycles at 3 dpf, control versus 28 µM (P = 0.0061, "OFF"). (F) Relative time spent in the center during the light on and light off cycles at 3 dpf, control versus 28 µM. (G) Total time spent in the center during the light on and light off cycles at 5 dpf, control versus 7 µM (P = 0.0004, "ON"). (H) Relative time spent in the center during the light on and light off cycles at 5 dpf, control versus 7 µM (P < 0.0001). (I) Total time spent in the center during the light on and light off cycles at 5 dpf, control versus 14 µM (P < 0.0001, "ON"; P < 0.0001, "OFF"). (J) Relative time spent in the center during the light on and light off cycles at 5 dpf, control versus 14 µM (P < 0.0001, "ON"; P = 0.0004, "OFF"). Control n = 88-102 for controls; treated n = 84-100. Error bars represent SEM. See Table S5 for additional statistical details.

Figure S8. CaMPARI analysis of the habenula in larvae exposed to varying concentrations to PFOS. (A) Illustrative representation of a larval zebrafish brain with anatomical regions outlined: forebrain (FB), habenula (Hb), and the optic tectum (OT). Activity quantification of neuron-driven CaMPARI was performed in the entire developing habenula following 1-minute exposure to 405 nm light. (B) At 3 dpf, there were no significant differences in habenular Ca2+ activity. (C) At 5 dpf, 14 µM PFOS-exposed larvae had a significant increase in habenular Ca^{2+} activity (P = 0.0170), but 7 µM PFOS was unchanged. n = 20-22 fish per treatment. Box plot limits represent 25^{th} to 27^{th} percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S9. Distance traveled in control versus PFOS-treated wildtype and macrophage mutant larvae. (A) 3 dpf light on (P = 0.0297, WT; P = 0.3036, *irf8*$^{-/-}$). (B) 3dpf light off (P < 0.0001, WT; P = 0.2162, *irf8*$^{-/-}$). (C) 3 dpf total distance moved throughout assay (P < 0.0001, WT; P = 0.1069, *irf8*$^{-/-}$). (D) Relative time spent in center normalized to the respective control groups. (E) 5 dpf light on (P < 0.0001, WT; P < 0.0001, *irf8*$^{-/-}$). (F) 5 dpf light off (P = 0.0200, WT; P = 0.0456, *irf8*$^{-/-}$). (G) 5 dpf total distance moved throughout assay (P = 0.0009, WT; P = 0.0011, *irf8*$^{-/-}$). (H) Relative time spent in center normalized to the respective control groups (P = 0.0213, WT; P = 0.1513, *irf8*$^{-/-}$). n = 14-22 per group. Error bars represent SEM. Table S5 for additional statistical details.

Figure S10. Microglia response to brain injury in control versus PFOS-exposed versus PTZ-exposed larvae at 3 dpf. Larvae with transgenic expression of macrophages (Tg(*mpeg1:EGFP*)) were dosed with either (A) control or (B) 28 μM PFOS at 4 hpf, or (C) 5 mM PTZ at 72 hpf, followed by brain injury. (D) At 4 hpi, PFOS-exposed larvae had a significant increase in microglia response area ($P = 0.0082$). In addition, neuronal excitation by PTZ also resulted in a significant increase in microglia response ($P = 0.0314$). n = 14-21 per group. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S11. Light/dark behavioral analysis in PFOA-exposed larvae. Zebrafish larvae were subjected to a 30-minute light/dark behavioral assay at 5 dpf, as previously described. (A) There were no apparent changes in distance moved during the 30-minute assay. (B) Total distance moved during light cycles, (C) dark cycles, and (D) in total throughout the assay were not significantly changed. (E-G) PFOA exposure also did not result in changes in center-avoidance behavior during the light on or light off cycles. n = 24 control and n = 46 treated for behavior analyses; n = 37 control and n = 35 treated for anxiety analyses. See Table S5 for additional statistical details.

Figure S12. Body length of 5 dpf exposed to PFOA. (A'-A'') Images of 5 dpf control-treated larvae. (B'-B'') Images of 5 dpf larvae chronically treated with 64 μM PFOA. (C) Body length measurements of control versus PFOA treatment at 5 dpf ($P < 0.0001$). n = 5-8 per group. Scale = 500 um. See Table S5 for additional statistical details.

Supplemental Tables

Table S1. Survival rate of zebrafish larvae exposed to PFOS over time

Age	Control	14 µM PFOS	28 µM PFOS	56 µM PFOS
24 hpf	97.93 ± 2.94	97.67 ± 1.75	98.15 ± 1.73	97.22 ± 2.84
48 hpf	97.74 ± 2.76	97.14 ± 1.21	97.09 ± 2.10	2.49 ± 2.84
72 hpf	97.43 ± 3.25	96.18 ± 1.81	95.62 ± 2.43	81.54 ± 13.23
96 hpf	96.41 ± 5.82	95.72 ± 1.20	**50.65 ± 10.06***	**25.00 ± 9.89****
120 hpf	95.93 ± 5.85	89.14 ± 3.65	**7.73 ± 3.03*****	**5.33 ± 2.67*****

Percent survival of larvae aged 24 hpf to 120 hpf exposed to control solution, 14 µM PFOS, 28 µM PFOS, or 56 µM PFOS. Statistics compared each timepoint to the 24 hpf timepoint of the same treatment group using unpaired t-tests. ± denotes standard deviation. *$P<0.05$; ***$P<0.001$; ****$P<0.0001$.

Table S2. Primers used for qRT-PCR and genotyping

Gene	TaqMan ID or Primer Sequences
il6 (m17)	Dr03098117_g1
il1b	Dr03114368_m1
tnfa	Dr03126850_m1
actb1 (b-actin)	Dr03432610_m1
pr2y12	F: 5'-CTTCAGGTCGTCGCTGTTTA-3' R: 5'-AGTGCGTTTCCCTGTTGAT-3'
b-actin	F: 5'-CGAGCAGGAGATGGGAACC-3' R: 5'-CAACGGAAACGCTCATTGC-3'
irf8	F: 5'-ACATAAGGCGTAGAGATTGGACG-3' R: 5'- GGATGAGGACCGCACTATGTTTC-3'.

Table S3. PFOS body burden in zebrafish larvae

Chemical	Formula	MW (g/mol)	Control		28 µM PFOS		56 µM PFOS	
			48 hpf	72 hpf	48 hpf	72 hpf	48 hpf	72 hpf
PFOS	$C_8HF_{17}O_3S$	500.13	0.12 ± 0.17	0.12 ± 0.16	32.22 ± 2.57	**70.46 ± 2.72******	46.04 ± 3.75	**109.15 ± 5.99******
PFNS	$C_9HF_{19}O_3S$	550.14	0.01 ± 0.02	0.00 ± 0.00	0.19 ± 0.05	**0.39 ± 0.07******	0.23 ± 0.04	**0.69 ± 0.20*****
PFHpS	$C_7HF_{15}O_3S$	450.12	0.00 ± 0.00	0.00 ± 0.00	0.35 ± 0.17	0.45 ± 0.13	0.40 ± 0.17	**0.55 ± 0.11***
PFHxS	$C_6HF_{13}O_3S$	400.12	0.006 ± 0.01	0.008 ± 0.01	0.008 ± 0.01	0.007 ± 0.01	0.008 ± 0.01	0.008 ± 0.01

PFOS body burden (ng/embryo) in 48 hpf and 72 hpf larvae dosed with control solution, 28 µM PFOS, or 56 µM PFOS. Our verification of PFOS stock solution revealed contamination by PFNS, PFHpS, and PFHxS. As such, we assessed the body burden of these compounds in the PFOS-exposed larvae, as well. Statistics compared 48 hpf to 72 hpf of the same treatment group using unpaired t-tests. ± denotes standard deviation. n = 10 pooled whole larvae from 3 experimental replicates. *P<0.05; ***P<0.001; ****P<0.0001.

Table S4. Significantly enriched metabolic pathways following 28 µM PFOS exposure in 3 dpf larvae.

Significantly Enriched Pathway	Physiological Relevance in CNS	Source
Vitamin B1 (thiamin) metabolism	- Microglia regulation - Neuronal K$^+$ channels regulation - Myelinogenesis - Improves cognitive function	(1, 2)
Valine-leucine and isoleucine degradation	- Neurotransmitter metabolism, including glutamate - Protein synthesis and energy production - Nitrogen homeostasis and neurotransmitter cycling	(3, 4)
Tyrosine metabolism	- Catecholamine synthesis - Regulator of neuronal longevity	(5, 6)
Propanoate metabolism	- Modulates glutamine synthetase in glia - Inhibits histone deacetylase in GABAergic neurons, increasing GABA levels	(7, 8)
Phosphatidylinositol phosphate metabolism	- Neurotransmitter receptor expression regulation - Synaptic vesicle regulation and recycling - Neurite and dendrite morphogenesis - Clathrin-dependent membrane trafficking - Ion channel and transporter activity regulation	(9, 10)
Linoleate metabolism	- Modulates astrocyte inflammatory response - Stimulates axonal growth of cortical neurons	(11, 12)
Limonene and pinene degradation	- Represses neuronal cell death - Decreases activated glial cell number - Involved in neuroprotection - Anti-inflammatory and analgesic roles	(13)
Heparan sulfate degradation	- Regulator of axon guidance and synapse development and specificity - Involved in regulating the dopamine system - Part of the core synaptic organizing complexes neurexin and neuroligin	(14, 15)
Glyoxylate and dicarboxylate metabolism	- Important for carbohydrate metabolism and energy availability	(16)

	- Altered following traumatic brain injury	
Glutamate metabolism	- Involved in nitrogen trafficking and ammonia homeostasis in brain - Excitatory neurotransmitter and immediate precursor for the neurotransmitter GABA	(17)
Chondroitin sulfate degradation	- Role in development, plasticity, and regulation of cortical circuitry - Modulates ion channel properties - Inhibits structural plasticity (scar after injury, axonal pathfinding, and synapse formation during development) - Neuronal excitability modulation	(18)
Butanoate metabolism	- Neuromodulator largely produced by gut microbiota - Attenuates neuronal apoptosis - Inhibits amyloidogenesis - Protects neurons from ischemic damage - Improves long-term memory - Neurodegenerative attenuation	(19-21)
Aspartate and asparagine metabolism	- Secondary excitatory neurotransmission - Amino acid involved in glutamate synthesis - Influences NMDAR-mediated transmission - Can evoke presynaptic release of endogenous l-glutamate release in selective brain regions - Synaptic strength and connectivity modulation	(22, 23)

Figure S1. Microglia quantification, cell death, and proinflammatory cytokine mRNA analysis in PFOS-exposed larvae. (A) Non-parenchymal, (B) parenchymal, and (C) total microglia on were quantified in 3 dpf control (n = 15), 28 µM PFOS (n = 16), or 56 µM PFOS (n = 16) exposed larvae. There was no significant difference in parenchymal or non-parenchymal microglia number in 28 µM PFOS-exposed larvae compared to controls, while 56 µM PFOS larvae had significantly fewer parenchymal (P = 0.0168) and total microglia (P = 0.0205). (D,E) To assess cell death, live 3 dpf larvae were stained with 5 ug/mL acridine orange during the first hour post injury or (F,G) between 3-4 hpi. (H) Quantification of acridine orange-positive clusters in 3 dpf control and 28 µM PFOS-exposed larvae show no significant change in cell death (n = 5-9 per group). (I) qRT-PCR for the inflammatory genes *il6*, *tnfα*, and *il1β* were not significantly changed in isolated heads of 3 dpf control or 28 µM PFOS-exposed larvae (n = 10 pooled heads per sample). Confocal micrographs at 20x magnification. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S2. Control for the optogenetic microglia stimulation experiment by stimulating halorhodopsin negative microglia. To ensure that 589 nm light alone does not alter microglia morphology, PFOS-exposed transgenic larvae expressing macrophage-driven mCherry (*Tg(mpeg1:gal4FF;UAS:nfsb-mCherry)*) were either (A-B") unstimulated or (C-E") subjected to the 589 nm light for 4 hours. (F) While microglia area was increased following light exposure (P = 0.0168), (G) the light stimulation did not result in significant changes in perimeter or (H) the perimeter-to-area ratio. Therefore, light stimulation alone does not result in the ramification of microglia. n = 12 cells. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S3. Functional neuroimaging to understand the effects of toxicant exposure on neuronal activity. (A) Schematic overview of CaMPARI photoconversion: free-swimming zebrafish larvae with neuron-specific expression of the genetically encoded calcium indicator CaMPARI protein (*Tg(elavl3:CaMPARI)*) are subjected to 405 nm light for 1 minute. Exposure to blue light photoconverts any neurons with high intracellular calcium from green to red. (B-D) We modified our Noldus DanioVision Behavioral unit, outfitted with LEDs for optogenetic manipulation, to efficiently photoconvert CaMPARI in free-swimming zebrafish. (B) A 3D printed pedestal is used to reduce the working distance between the light source and zebrafish larvae. Applying the inverse-square law, the light intensity at the apex of the pedestal was calculated to be 135 mW/cm². (C) Illustration of the apex of the pedestal containing a single well dish (OD 15 mm). (D) The 405 nm light has an intensity of 5700 mW at the source. The light travels horizontally then is reflected off a dichroic mirror within the unit. The working distance was calculated as the distance from the light source to the dichroic mirror (α) plus the distance from the dichroic mirror to the apex of the pedestal (β). (E) Confocal micrographs of neuronal calcium following 1-minute photoconversion of CaMPARI in 3 dpf larvae treated with egg water or (E') after 10 minutes in 10 mM pentylenetetrazol (PTZ), a GABAA-inhibitor known to induce hyperactivity. (F, F') Green, non-active neurons with low intracellular calcium from E & E'. (G, G') Photoconverted, active neurons with high intracellular calcium, pseudo-labeled as magenta, from E & E'. (H, H') An intensity LUTs applied to the active neurons in G & G' spectrally maps the regional increases of neuronal activity on a scale from less active (blue) to more active (red-white). (I&J) Regional and global neuronal activity can be quantified by determining the ratio of fluorescent intensity in the red versus green channel (F_{Red}/F_{Green}). Exposure to 10 mM PTZ significantly increased brain activity in the optic tectum (OT; *P* = 0.0496) and trended toward increased activity in the cerebellum (Ce), hindbrain (HB), and whole brain (WB). Red/Green ratios were normalized relative to

vehicle control. n = 16 fish per group. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S4. Regional and global brain morphology in larvae exposed to varying concentrations of PFOS. (A-D) Confocal micrographs of 3 dpf larvae or (J-L) 5 dpf larvae expressing *Tg(elavl3:CaMPARI)* exposed to (A&J) control, (B&K) 7 μM, (C&L) 14 μM, or (D) 28 μM PFOS. At 3 dpf, 7 μM PFOS larvae had a significant reduction in area of the (E) forebrain (Control vs 7 μM PFOS, *P* = 0.0035) and (F) optic tectum (Control vs 7 μM PFOS, *P* = 0.0047). (I) When measuring the area of the entire brain, only 7 μM and 14 μM PFOS exposure resulted in reduced global brain area (Control vs 7 μM & 14 μM, *P* = 0.0174 & *P* = 0.0373, respectively). At 5 dpf, there were no significant differences in regional area or global brain area with 7 μM nor 14 μM PFOS exposure. Confocal micrographs at 10x magnification. Control n = 52-54; treated n = 22-25. Scale bar = 100 um. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S5. MWAS of heads collected from control versus PFOS-exposed larvae. Heads were collected from 3 dpf control or 28 µM PFOS exposed larvae for an untargeted metabolome wide association study (MWAS). (A) Volcano plots of metabolites detected using a C18 column with negative ionization and (B) a HILIC column with positive ionization reveal several significantly down- or up-regulated metabolites in the PFOS-exposed brain (Green = downregulated; Blue = upregulated). Statistical significance was set at *P* = 0.05. (C) There are several significantly enriched pathways following both negative ionization (pink) and positive ionization (teal). The vertical line is at *P* = 0.05. Size of each dot represents the number of significant metabolites in that pathway. HILIC, hydrophilic interaction liquid chromatography column.

Figure S6. Light/dark behavioral assay in larvae exposed to varying concentrations of PFOS. (A) Exposure paradigm: at 4 hpf, zebrafish embryos were statically exposed to 0.1% DMSO (Control) or varying concentrations of PFOS (7, 14, or 28 μM). At 24 hpf, embryos were enzymatically dechorionated. Behavior was captured at 72, 96, and 120 hpf. (B) Light/dark behavioral assay: Zebrafish larvae were placed in individual wells of a 24-well plate containing the relevant toxicant solution 18-24 hours prior to the behavior assay to allow for well acclimation. A light/dark behavioral assay was performed using the Noldus DanioVision Behavioral Unit. The light/dark behavioral assay was as follows: 15-minute acclimation in the unit with the light off, 5 minutes with the light on, 5 minutes with the light off, 5 minutes with the light on, and 15 minutes with the light off. (C-H) At 3 dpf, larvae exposed to 28 μM PFOS had a significant increase in swim activity during the (F) light on cycles (P = 0.0003) and (H) over the course of the assay (P = 0.0123). (K) At 4 dpf, 7 μM PFOS larvae had a significant increase in swim distance when the light was on (P = 0.0277), as did 14 μM PFOS larvae (P = 0.0002). (L) At 4 dpf, 7 μM PFOS-exposed larvae also had a significant increase in swim behavior when the light was off (P = 0.0018) and (M) in total over the course of the assay (P < 0.0001). (P) At 5dpf, 14 μM PFOS had increased swim behavior when the light was on (P = 0.0006), and (Q) 7 μM PFOS larvae had increased swim behavior when the light was off (P < 0.0001). (R) Both 7 μM (P < 0.0001) and 14 μM PFOS (P = 0.0003) larvae had a significant increase in total distance moved. Control n = 130-190, treated n = 88-99. Error bars represent SEM. See Table S5 for additional statistical details.

Figure S7. Absolute and relative time spent in center throughout the light/dark behavioral response assay. (A) Total time spent in the center during the light on and light off cycles at 3dpf, control versus 7 µM (*P* = 0.0139, "OFF"). (B) Relative time spent in the center during the light on and light off cycles at 3 dpf, control versus 7 µM. (C) Total time spent in the center during the light on and light off cycles at 3 dpf, control versus 14 µM (*P* = 0.0075, "ON"; *P* < 0.0001, "OFF"). (D) Relative time spent in the center during the light on and light off cycles at 3 dpf, control versus 14 µM (*P* = 0.0015, "ON"; *P* = 0.0018, "OFF"). (E) Total time spent in the center during the light on and light off cycles at 3 dpf, control versus 28 µM (*P* = 0.0061, "OFF"). (F) Relative time spent in the center during the light on and light off cycles at 3 dpf, control versus 28 µM. (G) Total time spent in the center during the light on and light off cycles at 5 dpf, control versus 7 µM (*P* = 0.0004, "ON"). (H) Relative time spent in the center during the light on and light off cycles at 5 dpf, control versus 7 µM (*P* < 0.0001). (I) Total time spent in the center during the light on and light off cycles at 5 dpf, control versus 14 µM (*P* < 0.0001, "ON"; *P* < 0.0001, "OFF"). (J) Relative time spent in the center during the light on and light off cycles at 5 dpf, control versus 14 µM (*P* < 0.0001, "ON"; *P* = 0.0004, "OFF"). Control n = 88-102 for controls; treated n = 84-100. Error bars represent SEM. See Table S5 for additional statistical details.

Figure S8. CaMPARI analysis of the habenula in larvae exposed to varying concentrations to PFOS.
(A) Illustrative representation of a larval zebrafish brain with anatomical regions outlined: forebrain (FB), habenula (Hb), and the optic tectum (OT). Activity quantification of neuron-driven CaMPARI was performed in the entire developing habenula following 1-minute exposure to 405 nm light. (B) At 3 dpf, there were no significant differences in habenular Ca2+ activity. (C) At 5 dpf, 14 µM PFOS-exposed larvae had a significant increase in habenular Ca^{2+} activity (P = 0.0170), but 7 µM PFOS was unchanged. n = 20-22 fish per treatment. Box plot limits represent 25^{th} to 27^{th} percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S9. Distance traveled in control versus PFOS-treated wildtype and macrophage mutant larvae. (A) 3 dpf light on (P = 0.0297, WT; P = 0.3036, $irf8^{-/-}$). (B) 3dpf light off (P < 0.0001, WT; P = 0.2162, $irf8^{-/-}$). (C) 3 dpf total distance moved throughout assay (P < 0.0001, WT; P = 0.1069, $irf8^{-/-}$). (D) Relative time spent in center normalized to the respective control groups. (E) 5 dpf light on (P < 0.0001, WT; P < 0.0001, $irf8^{-/-}$). (F) 5 dpf light off (P = 0.0200, WT; P = 0.0456, $irf8^{-/-}$). (G) 5 dpf total distance moved throughout assay (P = 0.0009, WT; P = 0.0011, $irf8^{-/-}$). (H) Relative time spent in center normalized to the respective control groups (P = 0.0213, WT; P = 0.1513, $irf8^{-/-}$). n = 14-22 per group. Error bars represent SEM. See Table S5 for additional statistical details.

Figure S10. Microglia response to brain injury in control versus PFOS-exposed versus PTZ-exposed larvae at 3 dpf. Larvae with transgenic expression of macrophages (Tg(*mpeg1:EGFP*)) were dosed with either (A) control or (B) 28 µM PFOS at 4 hpf, or (C) 5 mM PTZ at 72 hpf, followed by brain injury. (D) At 4 hpi, PFOS-exposed larvae had a significant increase in microglia response area (P = 0.0082). In addition, neuronal excitation by PTZ also resulted in a significant increase in microglia response (P = 0.0314). n = 14-21 per group. Box plot limits represent 25th to 27th percentile, with the midline representing the median. See Table S5 for additional statistical details.

Figure S11. Light/dark behavioral analysis in PFOA-exposed larvae. Zebrafish larvae were subjected to a 30-minute light/dark behavioral assay at 5 dpf, as previously described. (A) There were no apparent changes in distance moved during the 30-minute assay. (B) Total distance moved during light cycles, (C) dark cycles, and (D) in total throughout the assay were not significantly changed. (E-G) PFOA exposure also did not result in changes in center-avoidance behavior during the light on or light off cycles. n = 24 control and n = 46 treated for behavior analyses; n = 37 control and n = 35 treated for anxiety analyses. See Table S5 for additional statistical details.

Figure S12. Body length of 5 dpf exposed to PFOA. (A'-A") Images of 5 dpf control-treated larvae. (B'-B") Images of 5 dpf larvae chronically treated with 64 μM PFOA. (C) Body length measurements of control versus PFOA treatment at 5 dpf ($P < 0.0001$). n = 5-8 per group. Scale = 500 um. See Table S5 for additional statistical details.

References

1. Ba A. Metabolic and structural role of thiamine in nervous tissues. Cell Mol Neurobiol. 2008 Nov;28(7):923-31. Epub 20080719. doi:10.1007/s10571-008-9297-7. Cited in: Pubmed; PMID 18642074.

2. Mkrtchyan G, Aleshin V, Parkhomenko Y, Kaehne T, Di Salvo ML, Parroni A, Contestabile R, Vovk A, Bettendorff L, Bunik V. Molecular mechanisms of the non-coenzyme action of thiamin in brain: biochemical, structural and pathway analysis. Sci Rep. 2015 Jul 27;5:12583. Epub 20150727. doi:10.1038/srep12583. Cited in: Pubmed; PMID 26212886.

3. Polis B, Samson AO. Role of the metabolism of branched-chain amino acids in the development of Alzheimer's disease and other metabolic disorders. Neural Regen Res. 2020 Aug;15(8):1460-1470. None. doi:10.4103/1673-5374.274328. Cited in: Pubmed; PMID 31997805.

4. Salcedo C, Andersen JV, Vinten KT, Pinborg LH, Waagepetersen HS, Freude KK, Aldana BI. Functional Metabolic Mapping Reveals Highly Active Branched-Chain Amino Acid Metabolism in Human Astrocytes, Which Is Impaired in iPSC-Derived Astrocytes in Alzheimer's Disease. Front Aging Neurosci. 2021;13:736580. The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. Epub 20210917. doi:10.3389/fnagi.2021.736580. Cited in: Pubmed; PMID 34603012.

5. Kobayashi K, Morita S, Sawada H, Mizuguchi T, Yamada K, Nagatsu I, Hata T, Watanabe Y, Fujita K, Nagatsu T. Targeted disruption of the tyrosine hydroxylase locus results in severe catecholamine depletion and perinatal lethality in mice. J Biol Chem. 1995 Nov 10;270(45):27235-43. Epub 1995/11/10. doi:10.1074/jbc.270.45.27235. Cited in: Pubmed; PMID 7592982.

6. Parkhitko AA, Ramesh D, Wang L, Leshchiner D, Filine E, Binari R, Olsen AL, Asara JM, Cracan V, Rabinowitz JD, Brockmann A, Perrimon N. Downregulation of the tyrosine degradation pathway extends Drosophila lifespan. Elife. 2020 Dec 15;9. AP, DR, LW, DL, EF, RB, AO, JA, VC, JR, AB, NP No competing interests declared. Epub 20201215. doi:10.7554/eLife.58053. Cited in: Pubmed; PMID 33319750.

7. Morland C, Froland AS, Pettersen MN, Storm-Mathisen J, Gundersen V, Rise F, Hassel B. Propionate enters GABAergic neurons, inhibits GABA transaminase, causes GABA accumulation and lethargy in a model of propionic acidemia. Biochem J. 2018 Feb 16;475(4):749-758. Epub 20180216. doi:10.1042/BCJ20170814. Cited in: Pubmed; PMID 29339464.

8. Nguyen NH, Morland C, Gonzalez SV, Rise F, Storm-Mathisen J, Gundersen V, Hassel B. Propionate increases neuronal histone acetylation, but is metabolized oxidatively by glia. Relevance for propionic acidemia. J Neurochem. 2007 May;101(3):806-14. Epub 20070205. doi:10.1111/j.1471-4159.2006.04397.x. Cited in: Pubmed; PMID 17286595.

9. Clayton EL, Minogue S, Waugh MG. Phosphatidylinositol 4-kinases and PI4P metabolism in the nervous system: roles in psychiatric and neurological diseases. Mol Neurobiol. 2013 Feb;47(1):361-72. Epub 20121010. doi:10.1007/s12035-012-8358-6. Cited in: Pubmed; PMID 23054682.

10. Raghu P, Joseph A, Krishnan H, Singh P, Saha S. Phosphoinositides: Regulators of Nervous System Function in Health and Disease. Front Mol Neurosci. 2019;12:208. Epub 2019/09/12. doi:10.3389/fnmol.2019.00208. Cited in: Pubmed; PMID 31507376.

11. Saba F, Sirigu A, Pillai R, Caria P, Cordeddu L, Carta G, Murru E, Sogos V, Banni S. Downregulation of inflammatory markers by conjugated linoleic acid isomers in human cultured astrocytes. Nutr Neurosci. 2019 Mar;22(3):207-214. Epub 20170829. doi:10.1080/1028415X.2017.1367130. Cited in: Pubmed; PMID 28847225.

12. Hennebelle M, Morgan RK, Sethi S, Zhang Z, Chen H, Grodzki AC, Lein PJ, Taha AY. Linoleic acid-derived metabolites constitute the majority of oxylipins in the rat pup brain and stimulate axonal growth in primary rat cortical neuron-glia co-cultures in a sex-dependent manner. J Neurochem. 2020 Jan;152(2):195-207. Epub 20191126. doi:10.1111/jnc.14818. Cited in: Pubmed; PMID 31283837.

13. Eddin LB, Jha NK, Meeran MFN, Kesari KK, Beiram R, Ojha S. Neuroprotective Potential of Limonene and Limonene Containing Natural Products. Molecules. 2021 Jul 27;26(15). Epub 20210727. doi:10.3390/molecules26154535. Cited in: Pubmed; PMID 34361686.

14. Condomitti G, de Wit J. Heparan Sulfate Proteoglycans as Emerging Players in Synaptic Specificity. Front Mol Neurosci. 2018;11:14. Epub 20180126. doi:10.3389/fnmol.2018.00014. Cited in: Pubmed; PMID 29434536.

15. De Risi M, Tufano M, Alvino FG, Ferraro MG, Torromino G, Gigante Y, Monfregola J, Marrocco E, Pulcrano S, Tunisi L, Lubrano C, Papy-Garcia D, Tuchman Y, Salleo A, Santoro F, Bellenchi GC, Cristino L, Ballabio A, Fraldi A, De Leonibus E. Altered heparan sulfate metabolism during development triggers dopamine-dependent autistic-behaviours in models of lysosomal storage disorders. Nat Commun. 2021 Jun 9;12(1):3495. Epub 20210609. doi:10.1038/s41467-021-23903-5. Cited in: Pubmed; PMID 34108486.

16. Baker EW, Henderson WM, Kinder HA, Hutcheson JM, Platt SR, West FD. Scaled traumatic brain injury results in unique metabolomic signatures between gray matter, white matter, and serum in a piglet model. PLoS One. 2018;13(10):e0206481. The authors have declared that no competing interests exist. Epub 20181031. doi:10.1371/journal.pone.0206481. Cited in: Pubmed; PMID 30379914.

17. Schousboe A, Scafidi S, Bak LK, Waagepetersen HS, McKenna MC. Glutamate metabolism in the brain focusing on astrocytes. Adv Neurobiol. 2014;11:13-30. doi:10.1007/978-3-319-08894-5_2. Cited in: Pubmed; PMID 25236722.

18. Hudson AE, Gollnick C, Gourdine JP, Prinz AA. Degradation of extracellular chondroitin sulfate delays recovery of network activity after perturbation. J Neurophysiol. 2015 Aug;114(2):1346-52. Epub 20150624. doi:10.1152/jn.00455.2015. Cited in: Pubmed; PMID 26108956.

19. Kim SY, Chae CW, Lee HJ, Jung YH, Choi GE, Kim JS, Lim JR, Lee JE, Cho JH, Park H, Park C, Han HJ. Sodium butyrate inhibits high cholesterol-induced neuronal amyloidogenesis by modulating NRF2 stabilization-mediated ROS levels: involvement of NOX2 and SOD1. Cell Death Dis. 2020 Jun 18;11(6):469. Epub 20200618. doi:10.1038/s41419-020-2663-1. Cited in: Pubmed; PMID 32555166.

20. Xu Y, Peng S, Cao X, Qian S, Shen S, Luo J, Zhang X, Sun H, Shen WL, Jia W, Ye J. High doses of butyrate induce a reversible body temperature drop through transient proton leak in mitochondria of brain neurons. Life Sci. 2021 Aug 1;278:119614. Epub 20210519. doi:10.1016/j.lfs.2021.119614. Cited in: Pubmed; PMID 34022200.

21. Zhou Z, Xu N, Matei N, McBride DW, Ding Y, Liang H, Tang J, Zhang JH. Sodium butyrate attenuated neuronal apoptosis via GPR41/Gbetagamma/PI3K/Akt pathway after MCAO in rats. J Cereb Blood Flow Metab. 2021 Feb;41(2):267-281. Epub 20200309. doi:10.1177/0271678X20910533. Cited in: Pubmed; PMID 32151222.

22. Pardo B, Rodrigues TB, Contreras L, Garzon M, Llorente-Folch I, Kobayashi K, Saheki T, Cerdan S, Satrustegui J. Brain glutamine synthesis requires neuronal-born aspartate as amino donor for glial glutamate formation. J Cereb Blood Flow Metab. 2011 Jan;31(1):90-101. Epub 20100825. doi:10.1038/jcbfm.2010.146. Cited in: Pubmed; PMID 20736955.

23. Errico F, Nuzzo T, Carella M, Bertolino A, Usiello A. The Emerging Role of Altered d-Aspartate Metabolism in Schizophrenia: New Insights From Preclinical Models and Human Studies. Front Psychiatry. 2018;9:559. Epub 20181106. doi:10.3389/fpsyt.2018.00559. Cited in: Pubmed; PMID 30459655.