What causes bird-building collision risk? Seasonal dynamics and weather drivers

Kara M. Scott[1], Attilla Danko, Paloma Plant[2], and Roslyn Dakin[1]

[1]Department of Biology
Carleton University
1125 Colonel By Drive
Ottawa, Ontario, Canada, K1S 5B6

[2]Fatal Light Awareness Program Canada
PO Box 430
Toronto, Ontario, Canada, M5C 2J5

To whom correspondence should be addressed:
Roslyn Dakin: roslyn.dakin@gmail.com

KEYWORDS: aeroconservation, birds, building collisions, conservation, migration, mortality, weather

Abstract

Bird-building collisions are a major source of wild bird mortality, with hundreds of millions of fatalities each year in the United States and Canada alone. Here, we use two decades of daily citizen-science monitoring to characterize day-to-day variation in building collisions and determine the factors that predict the highest risk times in two North American cities. We use these analyses to evaluate three potential causes of increased collision risk: heightened migration traffic during benign weather, increased navigational and flight errors during inclement weather, and increased errors in response to highly directional sunlight that enhances reflected images. The seasonal phenology of collisions was consistent across sites and years, with daily collision rates approximately two-fold higher in autumn as compared to spring. During both migration seasons, collision risk was best predicted by the weather conditions at dawn. In spring, peak collision risk occurs on days with warm temperatures, south winds, and a lack of precipitation at dawn. In autumn, peak collision occurs on days with cool temperatures, north winds, high atmospheric pressure, a lack of precipitation, and clear conditions with high visibility. Based on these results, we hypothesize that collisions are influenced by two main weather-driven mechanisms. First, benign weather at dawn with winds that are favorable for migration flight causes an increase in migration traffic in both spring and autumn, creating greater opportunity for collisions to occur. Second, for autumnal migrants, cold clear conditions may cause an additional increase in collision risk. We propose that these conditions may be particularly hazardous in autumn because of the high abundance of naïve and diurnal migrants at that time of year. Our analysis also establishes that a relatively small proportion of days (15%) are responsible for 50% of the total collision mortality within a season, highlighting the importance of targeting mitigation strategies to the most hazardous times.

1. Introduction

Birds have been in a marked decline, and North America has lost an approximate 29% of bird
50 abundance since the 1970s (Rosenberg et al., 2019). One factor that may contribute to these
losses is the widespread mortality caused by birds colliding with buildings and windows, which
52 are largely driven by collisions by migrating birds (Arnold & Zink, 2011; Elmore et al., 2021;
Loss, Will, Loss, & Marra, 2014; Loss, Will, & Marra, 2015). Indeed, meta-analyses have
54 estimated that 365–988 million birds are killed in collisions each year in the USA, while another
16–42 million are killed in collisions each year in Canada (Loss et al., 2014; Machtans et al.,
56 2013). Collision risk varies greatly across landscapes, buildings, seasons, days, and times of day.
Given that collisions occur during flight, weather may be an important driver of this variation
58 because atmospheric conditions influence migration decisions, flight and navigational errors, and
the visual appearance of glass. Understanding the temporal dynamics of these collisions is
60 important for two reasons: it allows us to better predict times that will have elevated risk, and it
provides insight into potential causal mechanisms that can inform mitigation. This knowledge is
62 important for aeroecology and aeroconservation, which are underrepresented areas of study
(Davy, Ford, & Fraser, 2017; Horton, Van Doren, Stepanian, Farnsworth, & Kelly, 2016).
64

Several previous studies have examined the question of how weather conditions may predict bird
66 collisions. An early study by Klem (1989) reported a sample of anecdotal descriptions of weather
conditions at 2 houses on days when 83 collisions occurred, and hypothesized that collisions at
68 houses may be higher during benign weather. A 2020 study by Loss et al. examined the weather
conditions associated with collisions of 15 individual American Woodcocks at 21 houses during
70 spring 2018, and found that these Woodcock collisions were associated with inclement weather
events involving strong north winds (which opposed the direction of their pre-breeding migration
72 at that time) (Loss et al., 2020). While these observations are valuable, they do not allow us to
draw generalized conclusions regarding collision dynamics. Most recently, Lao et al. (2022)
74 analyzed daily data on several hundred collisions that occurred at 21 buildings in Minneapolis,
and reported that collision risk was greater during conditions with wind directions that were
76 favorable for migration, as well as with changing weather patterns. Although Lao's (2022) study
is the most rigorous to date, it is still limited geographically. Here, we leverage information on
78 more than 62,000 migratory bird collisions, representing two decades of daily monitoring by
citizen scientists in two large metropolitan areas (Toronto, Canada, and Chicago, USA). Our goal
80 is to evaluate the influence of weather and atmospheric conditions on daily variation in avian
collision rates in these urban environments.
82

There are several non-exclusive mechanisms by which weather and atmospheric conditions
84 might influence collision risk for birds. The first possibility is that weather-induced changes in
migration traffic (defined as the abundance of migrating birds in the air) can influence the
86 occurrence of collisions. Consistent with this hypothesis, avian collision rates with buildings and
aircraft broadly correlate with times of higher migration traffic (Elmore et al., 2021; Lao et al.,
88 2022; Nilsson et al., 2021; van Doren et al., 2021). Increased migration traffic is commonly
driven by benign weather during both spring and autumn migration periods (Bozó, Csörgõ, &
90 Heim, 2018; Erni, Liechti, Underhill, & Bruderer, 2002; Panuccio, Dell'Omo, Bogliani, Catoni,
& Sapir, 2019; Welcker & Vilela, 2019). During spring, the strongest predictor of migration
92 traffic is often warmer temperatures (Bozó et al., 2018; Haest, Hüppop, & Bairlein, 2020; Hamer

et al., 2021; Schekler & Sapir, 2021; Van Doren & Horton, 2018; Welcker & Vilela, 2019). By contrast, in autumn, the effects of temperature on migration traffic are less clear, with different studies finding positive, negative, or no associations between temperatures and autumn migration traffic (Bozó et al., 2018; Erni et al., 2002; Loss et al., 2020; Nilsson et al., 2019; Schekler & Sapir, 2021; Welcker & Vilela, 2019). If weather-induced changes in migration traffic are a major driver of avian collisions, then we expect higher collision rates to occur on days with benign conditions in both spring and fall (i.e., favorable wind direction, low wind speeds, low precipitation, and high atmospheric pressure), with additional risk posed by warm temperatures in spring.

A second mechanism by which weather and atmospheric conditions might influence collision risk is that inclement weather may increase the probability that birds fail to avoid collisions during flight. For example, inclement weather may negatively impact visual guidance and the ability of birds to detect and respond appropriately to obstacles (Marques et al., 2014). Inclement weather could also increase physiological stress through depletion of energy reserves, making birds more likely to make fatal errors. This is occasionally observed in mass casualty events; for example, in the fall of 2020, winter storm conditions led to a mass bird casualty event in the southwestern USA (Johnson, 2020). Certain weather conditions may also cause flying migrants to descend to lower altitudes, where the risk of collisions is greater. For instance, it has been proposed that inclement weather and cloud cover may drive flying birds down to lower altitudes that have more artificial light at night (ALAN) and more obstacles (Cochran & Graber, 1958; Van Doren et al., 2021). In general, if weather-induced flight and navigational errors are a major driver of collisions, then we then expect higher collision rates to occur on days with high wind speeds, high precipitation, increased cloud cover, and cold temperatures, all of which may disrupt sensory performance, increase stress, and/or drive flying birds to change flight altitude.

A third mechanism by which atmospheric conditions might influence collision risk is by affecting the visual appearance of glass windows. The appearance of glass has been found to be an important proximate driver of collisions, based on positive associations between collision rates and the extent of glass on buildings (Riding, O'Connell, & Loss, 2020), and experiments demonstrating that visual deterrents and acoustic signals placed on or in front of windows can prevent collisions (Boycott, Mullis, Jackson, & Swaddle, 2021; Klem, 2009; Klem & Saenger, 2013). The appearance of glass is also strongly influenced by weather: on days with clear skies (i.e., little to no cloud cover and/or high visibility), sunlight is more directional, which creates specular reflections and brightly-coloured reflected images of the surroundings on glass. These clear-sky conditions may increase the probability that birds are deceived by reflected images. By contrast, on cloudy or overcast days, and on days with precipitation, sunlight is diffuse, and the brightness and coherence of reflections is reduced due to low coherence as well as water droplets on reflective surfaces. If weather-induced changes in the appearance of glass are an important driver of collisions, then we expect collision rates to be elevated on days with reduced cloud cover, high atmospheric visibility, and a lack of precipitation.

In addition to the mechanisms above, we also consider the possibility that the phase of the moon may influence daily variation in collision risk (Norevik, Åkesson, Andersson, Bäckman, & Hedenström, 2019). The moon can be a navigational beacon for some nocturnal migrants and may facilitate flight at night (Miles, Money, Luxmoore, & Furness, 2010; Syposz, Gonçalves,

Carty, Hoppitt, & Manco, 2018). In addition to its potential effects on migration traffic, it is also possible that lunar illumination may interact with ALAN and/or other sources of navigational disruption that may influence collision risk. We therefore include lunar illumination in our analysis, although we have no *a priori* predictions as to the direction of any potential lunar effects.

Here, we investigate the weather conditions that predict day-to-day variation in building collisions for a broad range of North American migratory birds. Our analysis considers nine variables: air temperature, wind speed, wind direction, precipitation, relative humidity, atmospheric pressure, cloud cover, atmospheric visibility, and lunar illumination. We examine these potential drivers using data on over 62,000 collisions involving more than 100 migratory bird species. Our goal is to identify the general drivers of collisions in urban environments, rather than species-specific effects. We first use these data to estimate day-to-day variation in collision rates, and compare spring and autumn seasons. Second, we quantify the extent to which collisions are concentrated over a small number of dates (Horton, Van Doren, Albers, Farnsworth, & Sheldon, 2021). Third, we evaluate if daily collision rates are best explained by weather conditions at a specific time of day (morning, dawn, overnight, or the previous evening). Finally, we describe the relationships between collision rates and weather during spring and autumn, respectively. Our overall aim is to identify the major drivers of collision risk as a first step to identifying causal mechanisms. Ultimately, with increased knowledge, improved mitigation and aeroconservation measures will be possible (Davy et al., 2017).

2. Methods

2.1 Collision rates

To estimate daily collision rates, we obtained data on the number of migratory bird collision casualties recorded per day from two large metropolitan areas: Toronto, Canada (43.74 N, 79.37 W) and Chicago, USA (41.88 N, 87.63 W). Both cities have a long-term daily collision monitoring programs that provide the basis for important continent-wide estimates of the extent and causes of collision mortality (Arnold & Zink, 2011; Colling, Guglielmo, Bonner, & Morbey, 2022; Cusa, Jackson, & Mesure, 2015; Evans Ogden, 1996; Loss et al., 2014; Machtans et al., 2013; Van Doren et al., 2021; Winger et al., 2019). The data from Toronto were collected by FLAP Canada (Fatal Light Awareness Program, www.flap.org) and in Chicago by the Chicago Bird Collision Monitors and McCormick Place monitors (www.birdmonitors.net) (Winger et al., 2019).

For the purpose of this study, we focused on a 10-year period from each city with daily records on the number of migratory bird collision casualties detected (years 2009 to 2018 in Toronto, and 2006 to 2015 in Chicago). We chose these particular years because they represent the contiguous 10-year time span within each city that had the largest number of collision records available. The total number of records in the data included 28,657 migratory bird casualties in Toronto and 33,493 in Chicago. The Chicago dataset focused on casualties from the passerine order (89 identified species), whereas the Toronto dataset was predominantly passerines, with some non-passerines also recorded (138 identified bird species, 92% of which were passerines). All data processing and analyses were performed in R 4.2.2.

The vast majority of building collisions occur during the periods from March–June, and August–November, representing spring and autumn migration seasons, respectively. During the months of July, December, January, and February, over 95% of dates had 0 collision records in our data. Moreover, only 0.2% of all collision records were attributed to dates within the predominantly non-migratory months. Given that our aim in this study was to evaluate sources of day-to-day variation in avian collision rates, we limited our analyses to the spring and autumn migration seasons of March–June and August–November, as day-to-day variation was not present at other times (Fig. 1). The total number of city-dates in our analysis was 1,465 days in spring (745 from Toronto and 720 from Chicago) and 1,855 in autumn (925 from Toronto and 930 from Chicago). For the 1,023 dates that overlapped between the two cities, daily log-counts of collisions were moderately correlated (Pearson's R = 0.52, p < 0.0001).

2.2 Weather and atmospheric conditions

We obtained hourly weather station data on nine weather and atmospheric parameters that are expected to influence flight and collision risk: air temperature (C), wind speed (km/h), wind direction (º), precipitation (mm/h), relative humidity (%), atmospheric pressure (kPa), cloud cover (%), atmospheric visibility (km), and lunar illumination (%). Hourly values were obtained for each city from www.openweathermap.org. All cloud cover estimates were expressed as a percent of full cloud cover, such that a value of 100% indicates a completely overcast sky. Lunar illumination values represent a measure of lunar phase and were obtained from the R package lunar 0.1-04; a value of 0% indicates the new moon and 100% indicates a full moon.

We assume that collisions recorded on a given day may have occurred at any time in the previous 24 hours, with weather during the previous 12-14 hours hypothesized to have the greatest influence (Evans Ogden, 1996; Lao et al., 2020). Therefore, we summarized the weather data into four time periods: (i) the previous evening, defined as the average of hourly weather conditions from 2000 to 2300 on the previous day; (ii) the overnight period, defined as the average of hourly conditions from 0000 to 0300; (iii) dawn, defined as the average of hourly conditions from 0400 to 0700; and (iv) morning, defined as the average of hourly conditions from 0800 to 1100 (all local times). As an example, for collisions recorded on April 10, 2009, the corresponding local morning temperature would be calculated as the average from 0800-1100 that day, whereas the corresponding previous evening temperature would be calculated as the average from 2000-2300 on April 9, 2009.

2.3 Migration collision phenology

Collision rates have a well-defined phenological peak within each spring and autumn migration season, corresponding with avian migration phenology (Fig. 1). To quantify this baseline temporal variation in migration collision phenology, we fit a cubic spline to the daily number of casualties as a function of day of the year within each city-year-season, with 0.6 as the smoothing parameter (Fig. 1B). The resulting splines represent the expected phenological pulse of casualties for a given season and year, at a given location. We used these phenology pulse estimates as a covariate in our statistical analysis of day-to-day variation in collision rates (see details in section 2.4 below). Hence, our analysis evaluates if local daily weather can explain day-to-day variation in collision rates, after accounting for the expected seasonal pulse due to the timing of migration in a given year (Fig. 1B).

2.4 Statistical analysis of collision rates and weather

232 We used mixed-effects regression models to evaluate sources of variation in daily collision rates. The response variable was the daily number of collisions recorded (per city-date), log-

234 transformed to meet the assumptions of linear regression. All models were fit in R using the lme4 1.1-30 and lmerTest 3.1-3 packages.

To provide descriptive statistics for seasonal differences in collision rates, we first fit a

238 regression model with season as the only fixed effect predictor (spring or autumn), and year and city as random effects.

Next, we evaluated which weather time period was the best predictor of daily collision rates

within each season. To do this, we compared five candidate mixed-effect regression models: a previous evening model, an overnight model, a dawn model, a morning model, and a null model.

All models included migration collision phenology as a fixed effect predictor. The time period models included as additional fixed effects the nine weather variables described in section 2.2

above. Hence, a morning model contained local weather conditions for the morning time period, whereas a dawn model contained local weather conditions for the dawn period, etc. Because

wind direction is a circular variable ($360º = 0º$), we modelled wind direction as a 2^{nd}-order orthogonal polynomial. This was important because it allows wind directions at the boundary ($0º$

and $360º$) to have equivalent effects. In the context our study sites in Toronto and Chicago, south winds are expected to be favorable in spring, and north winds in autumn. The null model

included migration collision phenology as a fixed effect predictor, but had no weather variables. In all models, year and city were included as random effects.

Regression models were fit with maximum likelihood and compared using Akaike's Information

Criterion (AIC). We also used the MuMIn package 1.46.0 to calculate Akaike weights. The Akaike weight is a measure of the relative strength of the evidence for a given model in a model

set, expressed as a probability ranging from 0 to 1; the weights for all models in a set sum to 1. Hence, if a model has an Akaike weight of 1, it indicates complete certainty that the model is

best supported within its model set. If two models receive similar Akaike weights, it indicates that they are equivalently well-supported. We performed this model selection analysis separately

for each season (spring and autumn).

Finally, we examined the coefficient estimates from the best-supported regression model for spring and autumn, respectively. We used mean-centred and standardized variables (response

and predictor) to allow comparison of effect sizes within and across models. We also obtained p-values using Satterthwaite's method in the lmerTest package. As a measure of the total variance

in collision rates explained by the fixed effects in each model, we calculated R^2_{GLMM} using the MuMIn package. As an additional check on our results, we verified that day of the week was not

a significant predictor of collision rates in our analyses, and we confirmed that all conclusions were unchanged when controlling for day of the week.

One possibility that we do not consider here is the potential that collision risk could be

274 influenced by interactive effects of several weather variables (wherein one weather parameter modulates the effect others). Similarly, we do not consider combinations of weather conditions

276 from different time periods in the present study. Instead, for simplicity and greatest

generalizability, we limit our analyses here to only considering the main effects of weather
278 variables within a given time period.

280 **3. Results**

282 *3.1 Seasonal and day-to-day variation in collision rates*
The average daily collision rate was approximately twice as high in autumn as compared to
284 spring (1.86-fold greater in autumn; p < 0.0001; Fig. 1A). Within each season, there was
substantial day-to-day variation in collision rates (Fig. 1B). Notably, a relatively small number of
286 days account for a large proportion of the seasonal total (Fig. 2). For example, the 12 worst days
typically account for about 50% of the seasonal total, and the three worst days typically account
288 for between 19-22% of the seasonal total. In spring, collision risk consistently peaked in the 17[th]
week of the year (the last days of April and first days of May). In autumn, collisions consistently
290 peaked in the 40[th] week of the year (the first week of October).

292 *3.2 Drivers of spring collisions*
Dawn weather was the best predictor of spring collision risk (Akaike weight > 0.57; Table S1).
294 The null model, which included phenology, but no weather effects, had an Akaike weight of 0.
The best-supported model using weather at dawn indicates that spring collisions are mainly
296 driven by warm temperatures, south winds (which are favorable for north-flying migrants), and
dry conditions with a lack of precipitation (Fig. 3A and Table 1).
298

3.3 Drivers of autumn collisions
300 Dawn weather was also the best predictor of autumn collision rates (Akaike weight > 0.99; Table
S2). The null model for autumn had an Akaike weight of 0. The best-supported model using
302 weather at dawn indicates that autumn collisions are mainly driven by cool temperatures, north
winds (which are favorable for south-flying migrants), a lack of precipitation, high relative
304 humidity, high atmospheric pressure, and high visibility (Fig. 3B and Table 2).

306 There was also a tendency for elevated collision rates after dark autumn nights with low lunar
illumination (Table 2). However, as shown in Fig. 3B, this lunar illumination effect does not
308 explain as much of the day-to-day variation in collision rates as the effects of temperature, wind,
precipitation/pressure and visibility.
310

4. Discussion
312

We show here that bird building collisions follow a consistent seasonal phenology with strong
314 peaks in the spring and autumn migration seasons (Fig. 1). Our analysis also shows that autumn
migration is particularly hazardous (Fig. 1A), with daily collisions rates approximately twice as
316 severe as spring. The increased risk of collision mortality in autumn is likely due to the presence
of numerous first-time, naïve migrants in autumn (Newton, 2007). Although post-breeding
318 autumn migration has cumulatively more individual migrants, studies of bird migration intensity
have shown that spring migration generally has greater daily peak levels of migration traffic (i.e.,
320 in spring, migration traffic is concentrated with higher peak traffic per day) (Dokter et al., 2018;
Ng et al., 2022). The idea that greater daily collision rates in autumn are mainly driven by first-
322 time migrants is also corroborated by recent studies finding that local birds (with greater

experience of the landscape) are better at avoiding collisions (Riding et al., 2021; Sabo,
324   Hagemeyer, Lahey, & Walters, 2016).

326   We also found that a relatively small number of days are responsible for a large burden of total
       collision mortality within a site. For example, 50% of the recorded collisions occur in just 15%
328   of days within a season (Fig. 2). This finding is consistent with previous studies monitoring the
       abundance of migrating birds, which have revealed that a relatively small number of days
330   accounts for the majority of individual movements (Horton et al., 2021; Van Doren & Horton,
       2018). This concentration of migration (and collisions) into peak days emphasizes the
332   importance of understanding drivers of day-to-day variation.

334   During spring, we found that day-to-day variation in collision rates was best explained by
       weather conditions at dawn, consistent with recent reports that collisions peak overnight and
336   during early morning hours (Colling et al., 2022; Riding et al., 2021; Van Doren et al., 2021).
       Specifically, we found that spring collision risk was greatly elevated on days with warmer
338   temperatures, a lack of precipitation, and south winds (which would be favorable for pre-
       breeding migration at our study sites; Fig. 3A). Based on these results in the context of migration
340   biology, we suggest that the biggest driver of spring collisions is weather-induced changes in
       migration traffic (Elmore et al., 2021; Hamer et al., 2021). On warm, dry spring days with
342   favorable winds, there is an increased abundance of birds in the air, and this increases the
       opportunity for collisions to occur.
344
       During autumn, collision rates were also best predicted by the weather at dawn. Specifically,
346   autumn collision mortality was elevated on days that were cold, with high atmospheric pressure,
       a lack of precipitation, north winds (favorable for post-breeding migration), and high
348   atmospheric visibility (Fig. 3B). These results suggest that autumn collisions may be driven by a
       combination of two major mechanisms: an increase in migration traffic at dawn during mornings
350   with favorable flight conditions, and an increase in potential errors in response to glass under
       conditions with highly directional sunlight. We found a weak association between collision rates
352   and times with a darkened moon phase in autumn, but lunar illumination explained much less of
       the day-to-day variation as compared to other predictors (Fig. 3B).
354
       We designed our analysis to focus on the most generalizable weather predictors for the two
356   broad, urban sites in our data. It is important to note that there may be differences across sites,
       and for other locations, additional site-specific predictors are possible depending on geography,
358   and the geometry of specific landscape features used by birds as well as that of the buildings
       themselves. For example, Van Doren et al (2021) found that Lake Michigan factored into
360   weather-collision interactions, where winds could cause higher avian concentrations along
       Chicago's shoreline. Inclement weather can also drive mass casualty events that are specific to
362   certain locations (Johnson, 2020; Newton, 2007) and/or species (Loss et al., 2020). It is
       important to note that even though inclement weather may not be associated with most collisions
364   at our study site, it can still cause many collisions and may even be a contributing factor to
       collisions that occur on benign days. For example, inclement weather may cause birds to make
366   suboptimal choices for stopover landing areas, e.g., in urban regions or other areas that pose
       greater mortality risk (Jenni & Schaub, 2003). This may indirectly influence collision rates if
birds later attempt to leave the urban area at the next benign weather opportunity (Clipp et al.,

2020; Van Doren et al., 2017). This could be due to the greater number of birds forced to initiate flights in a hazardous environment (Lao et al., 2022). It is also possible that ALAN may compound the effects of inclement weather, as both may be disorienting factors (Arnold & Zink,
2011; Evans Ogden, 1996; La Sorte & Horton, 2021; Lao et al., 2020; Van Doren et al., 2021).

Overall, our analysis reveals that the vast majority of urban bird-building collisions occur on fair weather days, consistent with migration traffic as a major driver. In autumn, collisions also peak
on days with high visibility, pointing to the importance of both migration traffic and the appearance of reflections. These immediate predictors of daily collision risk – benign weather,
favorable winds, and high visibility at peak migration times – are valuable for predicting days of greatest risk in the future. It is important to note that our study here is limited to urban
landscapes. A key question for future work is whether the same patterns occur in residential and suburban sites, which are responsible for a large burden collision mortality across North America
(Loss et al., 2014; Machtans et al., 2013). Increased understanding of the conditions and mechanisms that drive bird-building collisions is vital in working towards prevention. As pointed
out by Gelb and Delacretaz (2009), there are high-collision sites that are responsible for most of the mortality. New platforms that model and predict migration traffic, such as BirdCast
(https://birdcast.info/) and BirdFlow (Fuentes, Van Doren, Fink, & Sheldon, 2022), could be used for focused action (such as turning off lights) on these high-risk sites during the days with
highest migration traffic as well as high visibility, yielding the greatest improvements in mitigation measures (Van Doren et al., 2021).
390

References
392

Arnold, T. W., & Zink, R. M. (2011). Collision mortality has no discernible effect on population
394   trends of North American birds. *PLoS ONE*, *6*(9). doi: 10.1371/journal.pone.0024708
Boycott, T. J., Mullis, S. M., Jackson, B. E., & Swaddle, J. P. (2021). Field testing an "acoustic
396   lighthouse": Combined acoustic and visual cues provide a multimodal solution that reduces avian collision risk with tall human-made structures. *PLoS ONE*, *16*(4 April), 1–23. doi:
398   10.1371/journal.pone.0249826
Bozó, L., Csörgõ, T., & Heim, W. (2018). Weather conditions affect spring and autumn
400   migration of Siberian leaf warblers. *Avian Research*, *9*(1), 1–8. doi: 10.1186/s40657-018-0126-5
402 Clipp, H. L., Cohen, E. B., Smolinsky, J. A., Horton, K. G., Farnsworth, A., & Buler, J. J. (2020). Broad-scale weather patterns encountered during flight influence landbird stopover
404   distributions. *Remote Sensing*, *12*(3). doi: 10.3390/rs12030565
Cochran, W. W., & Graber, R. R. (1958). Attraction of nocturnal migrants by lights on a
406   television tower. *The Wilson Bulletin*, *70*(4), 378–380.
Colling, O. M., Guglielmo, C. G., Bonner, S. J., & Morbey, Y. E. (2022). Migratory songbirds
408   and urban window collision mortality: vulnerability depends on species, diel timing of migration, and age class. *Avian Conservation and Ecology*, *17*(1). doi: 10.5751/ACE-
410   02107-170122
Cusa, M., Jackson, D. A., & Mesure, M. (2015). Window collisions by migratory bird species:
412   urban geographical patterns and habitat associations. *Urban Ecosystems*, *18*(4), 1427–1446. doi: 10.1007/s11252-015-0459-3
414 Davy, C. M., Ford, A. T., & Fraser, K. C. (2017). Aeroconservation for the fragmented skies.

Conservation Letters, *10*(6), 773–780. doi: 10.1111/conl.12347

416   Dokter, A. M., Farnsworth, A., Fink, D., Ruiz-Gutierrez, V., Hochachka, W. M., La Sorte, F.
       A., … Kelling, S. (2018). Seasonal abundance and survival of North America's migratory
418   avifauna determined by weather radar. *Nature Ecology and Evolution*, *2*(10), 1603–1609.
       doi: 10.1038/s41559-018-0666-4

420   Elmore, J. A., Riding, C. S., Horton, K. G., O'Connell, T. J., Farnsworth, A., & Loss, S. R.
       (2021). Predicting bird-window collisions with weather radar. *Journal of Applied Ecology*,
422   1–9. doi: 10.1111/1365-2664.13832

       Erni, B., Liechti, F., Underhill, L. G., & Bruderer, B. (2002). Wind and rain govern the intensity
424   of nocturnal bird migration in central Europe - A log-linear regression analysis. *Ardea*,
       90(1), 155–166.

426   Evans Ogden, L. J. (1996). *Collision course: the hazards of lighted structures and windows to
       migrating birds*. Technical Report. World Wildlife Fund Canada, Toronto: World Wildlife
428   Fund Canada.

       Fuentes, M., Van Doren, B. M., Fink, D., & Sheldon, D. (2022). BirdFlow: learning seasonal
430   bird movements from citizen science data. *BioRxiv*. Retrieved from
       https://www.biorxiv.org/content/10.1101/2022.04.12.488057v2%0Ahttps://www.biorxiv.or
432   g/content/10.1101/2022.04.12.488057v2.abstract

       Gelb, Y., & Delacretaz, N. (2009). Windows and vegetation: primary factors in Manhattan bird
434   collisions. *Northeastern Naturalist*, *16*(3), 455–470. doi: 10.1656/045.016.n312

       Haest, B., Hüppop, O., & Bairlein, F. (2020). Weather at the winter and stopover areas
436   determines spring migration onset, progress, and advancements in Afro-Palearctic migrant
       birds. *Proceedings of the National Academy of Sciences of the United States of America*,
438   *117*(29), 17056–17062. doi: 10.1073/pnas.1920448117

       Hamer, T. E., Denis, N., Cardoso, T. P., Rocca, C. E., Luzenski, J. G., Harness, R. E., …
440   Landon, M. A. (2021). Influence of local weather on collision risk for nocturnal migrants
       near an electric power transmission line crossing Kittatinny Ridge, New Jersey. *Wilson
442   Journal of Ornithology*, *133*(2), 190–201. doi: 10.1676/19-00030

       Horton, K. G., Van Doren, B. M., Albers, H. J., Farnsworth, A., & Sheldon, D. (2021). Near-
444   term ecological forecasting for dynamic aeroconservation of migratory birds. *Conservation
       Biology*, *35*(6), 1777–1786. doi: 10.1111/cobi.13740

446   Horton, K. G., Van Doren, B. M., Stepanian, P. M., Farnsworth, A., & Kelly, J. F. (2016). Where
       in the air? Aerial habitat use of nocturnally migrating birds. *Biology Letters*, *12*(11), 4–8.
448   doi: 10.1098/rsbl.2016.0591

       Jenni, L., & Schaub, M. (2003). Behavioural and physiological reactions to environmental
450   variation in bird migration: a review. In *Avian Migration* (pp. 155–171).

       Johnson, K. (2020, December 16). Study: starvation and a freak snow storm caused the
452   southwest's mass bird die-off. *Audubon*. Retrieved from
       https://www.audubon.org/news/study-starvation-and-freak-snow-storm-caused-southwests-
454   mass-bird-die

       Klem, D. (1989). Bird-window collisions. *The Wilson Bulletin*, *101*(4), 606–620.

456   Klem, D. (2009). Preventing bird-window collisions. *Wilson Journal of Ornithology*, *121*(2),
       314–321.

458   Klem, D., & Saenger, P. G. (2013). Evaluating the effectiveness of select visual signals to
       prevent bird-window collisions. *Wilson Journal of Ornithology*, *125*(2), 406–411. doi:
460   10.1676/12-106.1

La Sorte, F. A., & Horton, K. G. (2021). Seasonal variation in the effects of artificial light at night on the occurrence of nocturnally migrating birds in urban areas. *Environmental Pollution*, *270*, 1–8. doi: 10.1016/j.envpol.2020.116085

Lao, S., Anderson, A. W., Blair, R. B., Eckles, J. W., Turner, R. J., & Loss, S. R. (2022). Bird–building collisions increase with weather conditions that favor nocturnal migration and with inclement and changing weather. *American Ornithological Society*, 1–76.

Lao, S., Robertson, B. A., Anderson, A. W., Blair, R. B., Eckles, J. W., Turner, R. J., & Loss, S. R. (2020). The influence of artificial night at night and polarized light on bird-building collisions. *Biological Conservation*, *241*, 1–8. doi: 10.1016/j.biocon.2019.108358

Loss, S. R., Lao, S., Anderson, A. W., Blair, R. B., Eckles, J. W., & Turner, R. J. (2020). Inclement weather and American woodcock building collisions during spring migration. *Wildlife Biology*, *2020*(1), 1–8. doi: 10.2981/wlb.00623

Loss, S. R., Will, T., Loss, S. S., & Marra, P. P. (2014). Bird–building collisions in the United States: Estimates of annual mortality and species vulnerability. *The Condor*, *116*(1), 8–23. doi: 10.1650/condor-13-090.1

Loss, S. R., Will, T., & Marra, P. P. (2015). Direct mortality of birds from anthropogenic causes. *Annual Review of Ecology, Evolution, and Systematics*, *46*, 99–120. doi: 10.1146/annurev-ecolsys-112414-054133

Machtans, C. S., Wedeles, C. H. R., & Bayne, E. M. (2013). A first estimate for Canada of the cumber of birds killed by colliding with building windows. *Avian Conservation and Ecology*, *8*(2), 1–6.

Marques, A. T., Batalha, H., Rodrigues, S., Costa, H., Pereira, M. J. R., Fonseca, C., … Bernardino, J. (2014). Understanding bird collisions at wind farms: An updated review on the causes and possible mitigation strategies. *Biological Conservation*, *179*, 40–52. doi: 10.1016/j.biocon.2014.08.017

Miles, W., Money, S., Luxmoore, R., & Furness, R. W. (2010). Effects of artificial lights and moonlight on petrels at St Kilda. *Bird Study*, *57*(2), 244–251. doi: 10.1080/00063651003605064

Newton, I. (2007). Weather-related mass-mortality events in migrants. *Ibis*, *149*(3), 453–467. doi: 10.1111/j.1474-919X.2007.00704.x

Ng, W. H., Fink, D., La Sorte, F. A., Auer, T., Hochachka, W. M., Johnston, A., & Dokter, A. M. (2022). Continental-scale biomass redistribution by migratory birds in response to seasonal variation in productivity. *Global Ecology and Biogeography*, *31*(4), 727–739. doi: 10.1111/geb.13460

Nilsson, C., Dokter, A. M., Verlinden, L., Shamoun-Baranes, J., Schmid, B., Desmet, P., … Liechti, F. (2019). Revealing patterns of nocturnal migration using the European weather radar network. *Ecography*, *42*(5), 876–886. doi: 10.1111/ecog.04003

Nilsson, C., La Sorte, F. A., Dokter, A., Horton, K., Van Doren, B. M., Kolodzinski, J. J., … Farnsworth, A. (2021). Bird strikes at commercial airports explained by citizen science and weather radar data. *Journal of Applied Ecology*, *58*(10), 2029–2039. doi: 10.1111/1365-2664.13971

Norevik, G., Åkesson, S., Andersson, A., Bäckman, J., & Hedenström, A. (2019). The lunar cycle drives migration of a nocturnal bird. *PLoS Biology*, *17*(10), 1–13. doi: 10.1371/journal.pbio.3000456

Panuccio, M., Dell'Omo, G., Bogliani, G., Catoni, C., & Sapir, N. (2019). Migrating birds avoid flying through fog and low clouds. *International Journal of Biometeorology*, *63*(2), 231–

239. doi: 10.1007/s00484-018-01656-z

508   Riding, C. S., O'Connell, T. J., & Loss, S. R. (2020). Building façade-level correlates of bird-window collisions in a small urban area. *The Condor: Ornithological Applications*, *122*(1),
510   1–14. doi: 10.1093/condor/duz065

       Riding, C. S., O'Connell, T. J., & Loss, S. R. (2021). Multi-scale temporal variation in bird-
512   window collisions in the central United States. *Scientific Reports*, *11*(1), 1–12. doi: 10.1038/s41598-021-89875-0

514   Rosenberg, K. V., Dokter, A. M., Blancher, P. J., Sauer, J. R., Smith, A. C., Smith, P. A., … Marra, P. P. (2019). Decline of the North American avifauna. *Science*, *366*(6461), 120–124.
516   doi: 10.1126/science.aaw1313

       Sabo, A. M., Hagemeyer, N. D. G., Lahey, A. S., & Walters, E. L. (2016). Local avian density
518   influences risk of mortality from window strikes. *PeerJ*, *4*, 1–15. doi: 10.7717/peerj.2170

       Schekler, I., & Sapir, N. (2021). Quantitative analysis of bird migration over Israel. *Meeting of*
520   *the Society for Integrative and Comparative Biology*. Washington D.C. Retrieved from http://sicb.burkclients.com/meetings/2021/abstracts/abstractdetails.php?id=690

522   Syposz, M., Gonçalves, F., Carty, M., Hoppitt, W., & Manco, F. (2018). Factors influencing Manx Shearwater grounding on the west coast of Scotland. *Ibis*, *160*(4), 846–854. doi:
524   10.1111/ibi.12594

       Van Doren, B. M., & Horton, K. G. (2018). A continental system for forecasting bird migration.
526   *Science*, *361*, 1115–1118. doi: 10.1101/293092

       Van Doren, B. M., Horton, K. G., Dokter, A. M., Klinck, H., Elbin, S. B., & Farnsworth, A.
528   (2017). High-intensity urban light installation dramatically alters nocturnal bird migration. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(42),
530   11175–11180. doi: 10.1073/pnas.1708574114

       van Doren, B. M., Willard, D. E., Hennen, M., Horton, K. G., Stuber, E. F., Sheldon, D., …
532   Winger, B. M. (2021). Drivers of fatal bird collisions in an urban center. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(24). doi:
534   10.1073/pnas.2101666118

       Van Doren, B. M., Willard, D. E., Hennen, M., Horton, K. G., Stuber, E. F., Sheldon, D., …
536   Winger, B. M. (2021). Drivers of fatal bird collisions in an urban center. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(24). doi:
538   10.1073/pnas.2101666118

       Welcker, J., & Vilela, R. (2019). *Weather-dependence of nocturnal bird migration and*
540   *cumulative collision risk at offshore wind farms in the German North and Baltic Seas*. Technical Report. Bio-Consult SH, Husum.

542   Winger, B. M., Weeks, B. C., Farnsworth, A., Jones, A. W., Hennen, M., & Willard, D. E. (2019). Nocturnal flight-calling behaviour predicts vulnerability to artificial light in
544   migratory birds. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1900). doi: 10.1098/rspb.2019.0364

DATA ACCESSIBILITY

548   All data and R scripts are available at: https://figshare.com/s/5c455f2a5c763682a737
The repository will be made public when the final version of the study is published.

AUTHOR CONTRIBUTIONS

552   RD and KMS designed the study; RD, KMS, PP and AA collected the data; RD and KMS
analyzed the data and wrote the manuscript.

COMPETING INTERESTS

556   We have no competing interests.

558   **FUNDING**
Supported by an NSERC Discovery Grant to RD and Carleton University.

ACKNOWLEDGEMENTS

562   We are grateful to all of the people who have contributed to the long-term collision monitoring
programs at FLAP Canada in Toronto (www.FLAP.org) and Chicago Bird Collision Monitors

564   (www.birdmonitors.net).

FIGURE LEGENDS

Figure 1. Seasonal phenology of bird collisions with buildings. (A) Daily collision rates
568 recorded in two cities, Toronto and Chicago, during the spring (green) and autumn (red)
migration periods (horizontal lines show the means). Daily collision rates are consistently higher
570 during autumn migration, as compared to spring. Note that the span of studied years differs for
the two sites. (B) To evaluate atmospheric and weather drivers of collision rates while
572 controlling for seasonal phenology, we first determined the expected rate of collisions by fitting
a cubic spline to each seasonal peak (shown by the green and pink lines in B). This procedure
574 was repeated for each site-season dataset; examples from Toronto are shown here. Next, we
analyzed the daily collision rate while controlling for the expected phenological pulse within
576 each season. Hence, the analysis investigates how weather conditions may explain fluctuations
above or below the seasonal expected rate of collisions.

Figure 2. Most collision risk is concentrated within a small number of days. These plots
580 show the cumulative proportion of total seasonal mortality for (A) Toronto and (B) Chicago. To
perform this analysis, days were first ranked by severity within each site-season (1^{st} = most
582 severe, etc.). Then, each day's cumulative proportion of the seasonal total was plotted with lines
connecting data from a given site-season. The steep increase in these cumulative proportion
584 curves illustrates how most collisions can be attributed to a small number of severe days within a
season. Each season represents four months: spring = March through June, and autumn = August
586 to November.

588 **Figure 3. Collision risk is driven by weather conditions that are largely benign for flight, as
well as high-visibility days in autumn.** (A) In spring, high collision rates are associated with
590 warm temperatures, south (favorable) winds, a lack of precipitation and low humidity. (B) In
autumn, high collision rates are associated with cool temperatures, north (favorable) winds, high
592 pressure, a lack of precipitation, and high atmospheric visibility. Each panel in (A) and (B)
shows the predictions of the best-supported model when all other predictors are set to the mean
594 value. The inner shaded region shows the 95% CI for the line-of-best fit. The outer shaded region
shows the 50% prediction interval of the model. Red is used for predictors that had a statistically
596 significant association with collision rates in the full model. For simplicity, only the weather
parameters that explain a substantial amount of collision variation in at least one season are
598 presented in this plot. See Tables 1-2 and Tables S1-S2 for additional details of this analysis.

600 **Table 1. Weather drivers of bird-building collisions in spring.** The spring weather conditions associated with the greatest collision risk include warm temperatures with a favorable (south)
602 wind, a lack of precipitation, and low relative humidity. To facilitate comparison of effect sizes, the response variable and all predictor variables were centered and standardized to have mean =
604 0 and SD = 1. Predictors are indicated in bold where the slope is statistically significant. See Tables S1 for additional details.

Predictor	Estimate	95% CI	p
Temperature	**0.14**	**[0.10, 0.18]**	**< 0.0001**
Wind speed	–0.02	[–0.07, 0.02]	0.27
Poly(Wind direction)-1	–0.56	[–2.13, –1.01]	0.48
Poly(Wind direction)-2	**–4.90**	**[–6.40, –3.40]**	**< 0.0001**
Precipitation	**–0.05**	**[–0.09, –0.01]**	**0.02**
Relative humidity	**–0.06**	**[–0.11, –0.02]**	**0.002**
Atmos. Pressure	–0.02	[–0.06, 0.03]	0.48
Cloud cover	0.04	[0.00, 0.09]	0.052
Visibility	–0.03	[–0.07, 0.01]	0.11
Lunar illum.	–0.01	[–0.05, 0.03]	0.62
Phenology	**0.62**	**[0.58, 0.66]**	**< 0.0001**

608 **Table 2. Weather drivers of bird-building collisions in autumn.** The autumn weather conditions associated with the greatest collision risk include cold temperatures with a favorable
610 (north) wind, a lack of precipitation, high visibility, and high atmospheric pressure. To facilitate comparison of effect sizes, the response variable and all predictor variables were centered and
612 standardized to have mean = 0 and SD = 1. Predictors are indicated in bold where the slope is statistically significant. See Tables S2 for additional details.

Predictor	Estimate	95% CI	p
Temperature	**–0.05**	**[–0.08, –0.02]**	**0.0006**
Wind speed	–0.03	[–0.06, 0.00]	0.06
Poly(Wind direction)-1	**3.42**	**[2.17, 4.69]**	**< 0.0001**
Poly(Wind direction)-2	**5.29**	**[4.06, 6.51]**	**< 0.0001**
Precipitation	**–0.08**	**[–0.10, –0.05]**	**< 0.0001**
Relative humidity	**0.05**	**[0.01, 0.09]**	**0.006**
Atmos. Pressure	**0.10**	**[0.06, 0.13]**	**< 0.0001**
Cloud cover	–0.02	[–0.05, 0.01]	0.16
Visibility	**0.14**	**[0.09, 0.18]**	**< 0.0001**
Lunar illum.	**–0.03**	**[–0.06, –0.01]**	**0.01**
Phenology	**0.78**	**[0.75, 0.81]**	**< 0.0001**

Figure 1

Figure 2

Figure 3

A) SPRING   B) AUTUMN

Collisions (/day)