1 **RESEARCH ARTICLE**
2

HY5 Regulates GLK and GNC Transcription Factors to Orchestrate Photomorphogenesis in *Arabidopsis thaliana*

7 **Ting Zhang[1,2,5], Rui Zhang[1,5], Xiyu Zeng[1,2,5], Luhuan Ye[1], Shilong Tian[2], Yijing**
8 **Zhang[3], Wenbin Zhou[4], Xin-Guang Zhu[1], Peng Wang[1]***
9
[1] Laboratory of Photosynthesis and Environmental Biology, CAS Center for
Excellence in Molecular Plant Sciences (CEMPS), Chinese Academy of Sciences,
Shanghai 200032, China
[2] University of Chinese Academy of Sciences, Shanghai 200032, China
[3] State Key Laboratory of Genetic Engineering, Collaborative Innovation Center of
Genetics and Development, Department of Biochemistry, Institute of Plant Biology,
School of Life Sciences, Fudan University, Shanghai 200438, China
[4] Institute of Crop Sciences, Chinese Academy of Agricultural Sciences, Beijing
100081, China
[5] These authors contributed equally to this work
20
21 * Corresponding author: wangpeng@cemps.ac.cn

22
23 **Short title:** GLK and GNC regulate photomorphogenesis
24
25 **One-sentence summary:** GLK and GNC act downstream of HY5, and cooperate
26 with HY5 and DET1, to regulate both chloroplast development and hypocotyl
27 elongation during the transition from skotomorphogenesis to
28 photomorphogenesis.
29

34 **ABSTRACT**

35 Light induced de-etiolation is an important aspect of seedling
36 photomorphogenesis. GOLDEN2 LIKE (GLK) and GATA NITRATE-INDUCIBLE

CARBON-METABOLISM-INVOLVED (GNC) are master transcriptional regulators involved in chloroplast development, but whether they participate in photomorphogenesis is largely unknown. Here we show that ELONGATED HYPOCOTYL5 (HY5) binds to *GLK* and *GNC* promoters and activates their expression. Chlorophyll content in the de-etiolating seedlings of the *hy5 glk2* and *hy5 gnc* double mutants was lower than that in the single mutants. It was further found that GLK inhibited hypocotyl elongation, whereas GNC promoted hypocotyl elongation, and that both could superimpose on the *hy5* phenotype. Correspondingly, GLK and GNC differentially regulate the expression of cell elongation genes. DE-ETIOLATED 1 (DET1) destabilises GLK proteins. The light-grown traits exhibited by etioplast of *det1* mutant were attenuated in *det1 glk2* double mutant, while similar to *det1* mutant, photosystem genes were up-regulated to different extent in etiolated seedlings overexpressing *GLK2* or *GNC*. Our study reveals that GLK and GNC act downstream of HY5 and likely cooperate with HY5 and DET1 to orchestrate multiple developmental traits during the light-induced skotomorphogenesis to photomorphogenesis transition.

INTRODUCTION

Light serves as an indispensable resource and the essential regulator for plant growth. In addition to photosynthesis, light also play roles in regulating seed germination, seedling de-etiolation, organ development, flowering and other important physiological processes (Jiao et al., 2007). After germination in darkness, seeds first grow into etiolated seedlings, displaying long hypocotyls with hooked tops and closed cotyledons. These characteristics are called skotomorphogenesis. Once the etiolated seedlings break through the soil surface and receive light, hypocotyl elongation is inhibited, cotyledons open, chloroplasts develop, and photosynthesis begins. This series of processes is referred to as

photomorphogenesis. Skotomorphogenesis and photomorphogenesis together ensure that etiolated seedlings emerge from the soil and develop into normal healthy seedlings, and are both a prerequisite for plant growth and development (Sullivan and Deng, 2003; Jiao et al., 2007).

As a positive regulator of photomorphogenesis, HY5 acts downstream of photoreceptors, such as cryptochromes, phytochromes and UV RESISTANCE LOCUS8 (UVR8), and plays important roles in seedling de-etiolation (Jiao et al., 2007; Stracke et al., 2010). Studies have shown that over 3000 genes can be directly regulated by the basic leucine zipper (bZIP) transcription factor HY5 (Jakoby et al., 2002; Lee et al., 2007; Zhang et al., 2011). It regulates many basic plant growth and development processes, including cell elongation, cell proliferation, chloroplast development, pigment accumulation, and nutrient uptake (Oyama et al., 1997; Shin et al., 2007; Jing et al., 2013). HY5 binds to promoters through G-box elements and inhibits the expression of cell elongation genes, such that *hy5* mutants show a typical long hypocotyl phenotype (Jing et al., 2013; Xu et al., 2016b). HY5 is important for chlorophyll and carotenoid accumulation under light as it regulates the expression of key enzymes such as PHYTOENE SYNTHASE (PSY) and PROTOCHLOROPHYLLIDE OXIDOREDUCTASE C (PORC). It also regulates the expression of photosynthetic pigment proteins, such as light-harvesting chlorophyll-protein complex I subunit A4 (LHCA4) (Toledo-Ortiz et al., 2014). Moreover, HY5 can bind to promoters of flavonoid synthesis genes, activating their expression and promoting anthocyanin synthesis (Shin et al., 2007; Song et al., 2008). HY5 functions through many different regulatory interactions. For example, it interacts with HY5-HOMOLOGY (HYH) or CALMODULIN7 (CAM7) protein to binds to the HY5 promoter and activate its own expression (Kushwaha et al., 2008; Abbas et al., 2014; Binkert et al., 2014). The interaction between HY5 and B-box-containing (BBX) protein BBX21-BBX22

enhances the promoting effect of HY5 on photomorphogenesis, whereas the

interaction between HY5 and BBX24-BBX25 does the opposite, suggesting that

BBX protein acts as coregulator of HY5, for fine-tuning of photomorphogenesis

(Datta et al., 2007; Datta et al., 2008; Xu et al., 2016a). Furthermore, HY5

interacts with PIF1 and PIF3 proteins to coordinate the production of reactive

oxygen species and the temperature control of photosynthetic gene expression

(Chen et al., 2013; Toledo-ortiz et al., 2014). In the case of excess HY5 protein, it

can activate the expression of COP1, thus inducing the degradation of HY5

protein and maintaining it at a relatively stable level (Huang et al., 2012).

GOLDEN2 (G2)-LIKE (GLK) and GATA NITRATE-INDUCIBLE CARBON

METABOLISM-INVOLVED (GNC) are two types of transcription factors directly

involved in the regulation of chloroplast development. GLK transcription factors

belong to the GARP superfamily (Fitter et al., 2002), which consists of G̲LK

(initially from maize), Arabidopsis RESPONSE REGULATOR-B (A̲R̲R-B) and

PHOSPHATE STARVATION RESPONSE 1 (P̲SR1) from Chlamydomonas (Hall et

al., 1998; Imamura et al., 1999; Wykoff et al., 1999). GLKs were first found to

regulate chloroplast development in maize, with a pair of GLK homologs ZmG2

and ZmGLK1 expressed in bundle sheath and mesophyll cells respectively (Hall

et al., 1998; Rossini et al., 2001). When maize G2 and GLK1 were overexpressed

in rice, chloroplasts in the bundle sheath of transgenic leaves were larger than

those in the wild type (Wang et al., 2017). In Arabidopsis, GLK1 and GLK2 are

regulated by light, and function redundantly in regulating chlorophyll biosynthesis

and chloroplast development. The leaves and siliques of the double mutant are

pale green, and thylakoid development in the chloroplasts is retarded (Fitter et al.,

2002). Overexpression of GLK1 or GLK2 in *glk1 glk2* double mutants can restore

the defect in chloroplast development (Waters et al., 2008). Ectopic expression of

GLKs was also able to induce increased chloroplast numbers in non-green

tissues such as roots and callus of Arabidopsis and rice (Nakamura et al., 2009;

Kobayashi et al., 2012a), and in tomato fruits (Nguyen et al., 2014). GLK target

genes are mainly related to chlorophyll biosynthesis, light-harvesting, and

electron transport (Waters et al., 2009; Kobayashi et al., 2012b). It also regulates

the expression of thylakoid membrane galactolipids monogalactosyldiacylglycerol

(MGDG) and digalactosyldiacylglycerol (DGDG) synthesis related genes

(Kobayashi et al., 2014). In the process of seedling de-etiolation, GLKs interact

with and are phosphorylated by BRASSINOSTEROID INSENSITIVE2 (BIN2)

protein, which promotes the stability and transcriptional activity of GLK1 (Zhang et

al., 2021). GLK also interacts with ORESARA 1 (ORE1) protein during

senescence, with ORE1 antagonizing GLK induction of target genes (Rauf et al.,

2013). The degradation of GLK protein itself is related to ubiquitination, with GLK2

shown to interact with DET1 protein in tomato prior to processing by CUL4-DDB1

ubiquitin E3 ligase for degradation (Tang et al., 2016).

The *GATA NITRATE-INDUCIBLE CARBON-METABOLISM-INVOLVED*

(*GNC*) gene was identified by screening mutants on nitrogen-deficient medium,

with *gnc* mutants accumulating lower chlorophyll than wild type (Bi et al., 2005).

The *CYTOKININ-RESPONSIVE GATA FACTOR1* (*CGA1*) gene is strongly

induced by cytokinin, and its expression in Arabidopsis histidine kinase mutants

ahk2 or *ahk3* is significantly reduced, suggesting that CGA1 is involved in the

cytokinin signaling (Naito et al., 2007). The *gnc cga1* double mutant exhibits

decreased chlorophyll content and decreased chloroplast number.

Overexpression of *GNC* or *CGA1* in Arabidopsis resulted in obvious chloroplast

development in seedling roots and hypocotyls, and increased chloroplast number

in cotyledon and leaf epidermal cells (Chiang et al., 2012). Overexpression of

CGA1 in rice increases chloroplast number as well as starch biosynthesis in the

leaves (Hudson et al., 2013; Lee et al., 2021). In spite of the well-known

downstream targets of GLK and GNC, their upstream regulation, particularly in the context of light signaling is not clear. As HY5 is a regulatory hub in the light signaling pathway, and it has been reported that GLK and GNC are dependent on HY5 in regulating root greening (Kobayashi et al., 2012a, 2017), the molecular relationships between HY5, GLK, and GNC are in need of systematic analysis. In addition, as the stability of GLK and GNC proteins is crucial for their activities, it is also important to consider the role of the COP/DET/FUS complex in regulating GLK and GNC function. DE-ETIOLATED 1 (DET1) is involved in the ubiquitination and degradation of positive regulators of photomorphogenesis in darkness. Apart from shorter hypocotyls and opened cotyledons, the dark-germinated *det1* mutant seedlings show part of light-grown traits, such as expression of photosynthetic genes and development of thylakoid membranes (Chory et al., 1989). The molecular mechanism behind such phenotypes in the dark is not fully understood and the potential involvement of GLK or GNC is worth considering.

Therefore, this study focused on the molecular and functional linkages among HY5, GLK, and GNC/CGA1 transcription factors, aiming to analyze and clarify the local regulatory network connecting them, so as to better understand the relationships between light signaling, seedling formation, and chloroplast development during photomorphogenesis. At the same time, relationships between GLK, GNC, and DET1, and their effects on the expression of photosynthetic genes in etiolated seedlings were examined. The results shed light on the concerted involvement of HY5, GLK, and GNC/CGA1 transcription factors in both skotomorphogenesis and photomorphogenesis.

RESULTS

HY5, GLK1/2, and GNC/CGA1 Regulate Chlorophyll Content to Different Extents in De-etiolating Arabidopsis Seedlings

HY5 and GLK both regulate the expression of chlorophyll biosynthesis and

photosynthesis related genes (Fitter et al., 2002; Lee et al., 2007; Waters et al.,

2009; Toledo-ortiz et al., 2014). To explore the significance of these regulations

during photomorphogenesis, we generated combined mutations in Arabidopsis,

and systematically investigated the effects of HY5 and GLK on chlorophyll

biosynthesis in de-etiolating seedlings. All plants were grown in darkness for 4

days, and then sampled at different time points after white light irradiation to

determine the content of chlorophyll. It was found that the chlorophyll content in

glk1 mutant had no or little difference from the wild type, while the chlorophyll

content in *glk2* mutant was significantly lower than that in the wild type, and the

chlorophyll content in *glk1 glk2* double mutant decreased more significantly than

that in *glk2* mutant **(Figures 1A and 1C; Supplemental Figures 1A and 1C)**.

This is consistent with the decrease of chlorophyll content in mature leaves of *glk1*

glk2 double mutant reported in previous literature (Fitter et al., 2002), indicating

that both GLKs affect chlorophyll biosynthesis in seedling de-etiolation process

and GLK2 may play a dominant role. Overexpression of GLK1 or GLK2 in double

mutants can compensate for the phenotype of decreased chlorophyll

(Supplemental Figures 1A and 1C). Compared with the above, we measured

the changes in the chlorophyll of *gnc* and *cga1* mutants during seedling

de-etiolation, and found that chlorophyll content of *cga1* mutants was not notably

different from the wild type, chlorophyll content of *gnc* and *gnc cga1* mutants was

lower than that of the wild type, while the difference was much smaller than that

between *glk2* related mutants and the wild type **(Figures 1B and 1D;**

Supplemental Figure 1B and 1D). The chlorophyll content of *35S:GNC*

overexpressed material was significantly higher than that of the wild type in the

greening process, while the chlorophyll content of *35S:CGA1* overexpressed

material showed no difference with that of the wild type at this stage

(Supplemental Figures 1B and 1D). The chlorophyll content of *hy5* mutant was

lower than that of the wild type, and overexpression of HY5 in the mutant could

effectively compensate for the phenotype of reduced chlorophyll **(Figure 1;**

Supplemental Figure 1), which supports the function of *hy5* regulating

chlorophyll biosynthesis (Tolede-Ortiz et al., 2014). In fact, we checked the

expression of *GLK1*, *GLK2*, *GNC*, and *CGA1* genes in the de-etiolation process of

wild type seedlings, and found that the expression levels of *GLK2* and *GNC* were

more prominent in response to light **(Figure 1E)**, which again demonstrates the

relative greater importance of them to seedling greening. Therefore, we conclude

here that HY5, GLKs and GNC/CGA1 all positively regulate chlorophyll

biosynthesis, but to different extents, during the greening process of seedling

photomorphogenesis.

Loss of *GLK1/2* or *GNC/CGA1* Function Combined with *HY5* Mutation Has

Limited Further Effects on Seedling Chlorophyll Content

In order to elucidate the genetic upstream or downstream relationships between

HY5, GLKs and GNC/CGA1, we constructed double and triple mutants, and

investigated chlorophyll content in seedlings during de-etiolation. It was found that

the chlorophyll content of *hy5 glk1* double mutant was not different from that of the

wild type and *glk1* during de-etiolation. Similarly, there was no notable difference

between *glk1* and wild type during de-etiolation from previous measurements

(Figures 1A and 1C). We noticed that the *glk1* single mutant used was caused by

an insertion in the promoter region, and thus the phenotypes in *glk1* single mutant

and *hy5 glk1* double mutant may not be representative of null alleles. On the other

hand, the chlorophyll content of *hy5 glk2* double mutant was lower than that of

glk2 or *hy5* mutant at 12 h in the process of greening. However, there was no

difference in chlorophyll content between *hy5 glk1 glk2* triple mutant and *glk1 glk2*

double mutant during de-etiolation **(Figures 1A and 1C)**. Compared with *hy5*

mutant, *hy5 gnc* and *hy5 cga1* mutants showed little or no difference in chlorophyll

content during de-etiolation, and only *hy5 gnc cga1* triple mutant exhibited notably

lower chlorophyll content than *gnc cga1* and *hy5* mutants at 12 h during greening.

Moreover, chlorophyll content of these mutants was generally higher than that of

glk related double or triple mutants **(Figures 1B and 1D)**.

In order to clarify the chlorophyll phenotype of *hy5* and *glk2* mutants, we

verified the expression levels of some chlorophyll biosynthesis and photosystem

genes in de-etiolating seedlings. At 12 h of light exposure, the expression levels of

chlorophyll biosynthesis genes (*HEMA1*, *PORB*, *PORC* and *CAO*) in *hy5*, *glk2*

and *hy5 glk2* mutants were all remarkably lower than those in the wild type. The

decrease in *glk2* and *hy5 glk2* mutants was not lower than that in the *hy5* mutant,

except for *PORB* **(Figure 2A)**. The expression levels of photosystem light

harvesting genes (*LHCB1.2*, *LHCB2.2*, *LHCA1* and *LHCA4*) was also significantly

lower in *hy5*, *glk2*, and *hy5 glk2* mutants than in the wild type, with their

expression in *hy5* mutant considerably higher than those in *glk2* and *hy5 glk2*

mutants **(Figure 2B)**. These results indicate that during de-etiolation of seedlings

GLK2 has stronger regulation on photosystem light harvesting genes than HY5,

whereas their regulation on chlorophyll biosynthesis may be comparable.

To further complement the observed phenotype, we moved on to analyse the

changes of chlorophyll in the mutant seedlings grown under continuous white light

for 4 days, and found that the trend of chlorophyll content among single mutant

and double mutant was similar to that in the de-etiolation process of seedlings.

However, chlorophyll content in *hy5 glk1 glk2* triple mutant was lower than that in

glk1 glk2 double mutant **(Figures 2C and 2D)**. We further investigated chloroplast

development in wild type, *glk1 glk2*, *hy5*, and *hy5 glk1 glk2* mutants under

sustained light exposure. Transmission electron microscopy showed that there

was no significant difference in mesophyll cell chloroplast ultrastructure between

hy5 and wild type at this stage. The chloroplast size of *glk1 glk2* and *hy5 glk1 glk2*

mutants was smaller than that of the wild type and *hy5* mutant, the accumulation

of starch grains was decreased, and the thylakoid stacking of *glk1 glk2* mutants

was sparser. Even worse, *hy5 glk1 glk2* mutant showed obvious fractured stromal

thylakoids **(Figure 2E)**. These results suggest that under certain conditions, *HY5*

mutation combined with *GLK1* and *GLK2* double mutations can lead to more

severe defects in chloroplast development. Because of the functional redundancy

of *GLK1* and *GLK2*, the double mutant shows severe perturbations in chlorophyll

biosynthesis and chloroplast development, with the superimposed mutation of

HY5 causing limited further defects. As such, we hypothesized that HY5 is either

located upstream of the GLK1 and GLK2 co-regulatory module, or that HY5 has

only a minor role in the regulation of chlorophyll biosynthesis.

HY5 Directly Regulates *GLK1*, *GLK2*, *GNC* and *CGA1* Gene Expression

According to ChIP-Seq analysis for HY5, its target genes include *GLK2*, *GNC*,

and *CGA1* (Lee et al., 2007). Through RT-qPCR verification, we found that the

expression levels of *GLK1*, *GLK2*, *GNC* and *CGA1* in *hy5* mutants were

significantly lower than those in the wild type **(Figures 1F)**, indicating that HY5

positively regulates their expression. Typical *cis*-elements to which HY5 binds to

regulate target gene expression include G-box (CACGTG), T/G-box (CACGTT),

E-box (CAATTG), ACE-Box (ACGT), and Z-Box (ATACGGT) (Shin et al., 2007;

Chang et al., 2008; Catala et al., 2011; Abbas et al., 2014). We performed

promoter analysis for *GLK1* and *GLK2*, and found that their promoter regions

contained HY5-binding *cis*-elements. These regions were synthesized to conduct

electrophoretic mobility shift assays (EMSA). It was shown that HY5 could bind to

GLK1 and *GLK2* promoter sequences, competitive probes weakened such

binding, while the binding ability of mutant probes was much lower than that of

normal probes **(Figure 3A)**. In parallel, by transforming protoplasts dual-luciferase

assay displayed that HY5 could activate *GLK1* and *GLK2* expression **(Figures 3B**

and 3C). While obtaining the above *in vitro* results, we also validated the binding

in vivo via Chromatin immunoprecipitation (ChIP) coupled with quantitative PCR

(qPCR) assay. Three or two regions containing HY5 binding elements were

selected from the promoters of *GLK1* and *GLK2* respectively, and the results

demonstrated that the P2 region from *GLK1* promoter and the P1 region from

GLK2 promoter were most strongly bound by HY5 **(Figure 3D)**. *In vitro*

experiment EMSA and dual-luciferase assays confirmed that HY5 could also bind

to *GNC* and *CGA1* promoters to activate their transcription, again followed by *in*

vivo experiment ChIP-qPCR showing that the highest enrichment of P1 on *GNC*

and *CGA1* promoters by HY5 **(Figure 3)**. In conclusion, HY5 directly binds to the

promoters of *GLK1*, *GLK2*, *GNC*, and *CGA1* to activate their gene expression.

GLK1/2 and GNC/CGA1 oppositely Regulate Hypocotyl Elongation of

Light-grown Seedlings

HY5 is known to inhibit hypocotyl elongation during photomorphogenesis, as *hy5*

seedlings exhibit longer hypocotyls (Sullivan and Deng, 2003; Jiao et al., 2007).

Since we found that HY5 directly regulates the expression of *GLK*, *GNC*, and

CGA1 genes, whether GLK, GNC and CGA1 also play roles in seedling

morphological development upon light regulation became the next question.

Therefore, we systematically investigated the hypocotyl phenotypes of a series of

glk, *gnc*, and *cga1* mutants. The hypocotyl length of *glk* mutants was measured

under different light quality, and those in *glk1*, *glk2* and *glk1 glk2* mutants were

longer than in wild type, while shorter than in the *hy5* mutant, after germinating

and growing under white, blue or red light conditions for 4 days. When *GLK1* or

GLK2 were overexpressed in *glk1 glk2* mutants, the hypocotyl length was

restored to the level of wild type. There was no significant difference in hypocotyl

length between the mutants and wild type after germinating and growing for 4

days in darkness **(Supplemental Figure 3)**. Together the above results indicate

that GLK inhibits hypocotyl elongation during light-induced seedling development.

We further investigated the hypocotyl phenotype in *hy5* and *glk* double and

triple mutants. Under white or blue light, compared with *hy5*, hypocotyls of *hy5*

glk1 were slightly longer, while hypocotyls of *hy5 glk2* and *hy5 glk1 glk2* were

significantly longer than that of *hy5*. The hypocotyls of all materials were generally

longer under red light, and those of *hy5 glk1*, *hy5 glk2* and *hy5 glk1 glk2* were still

longer than those of *hy5*, but with smaller differences. No notable differences in

hypocotyl length were observed in dark-grown seedlings between wild type and

these mutants **(Figures 4A and 4B)**. We measured the cell length of wild type,

glk1 glk2, *hy5*, and *hy5 glk1 glk2* materials in the middle segment of hypocotyls

growing under white light for 4 days, and found that the hypocotyl cell length of

glk1 glk2 mutant was longer than that of the wild type, and that of *hy5 glk1 glk2*

mutant was longer than that of *hy5* mutant **(Figures 4C and 4D)**. In general, there

is an obvious trend of *hy5* < *hy5 glk1* < *hy5 glk2* < *hy5 glk1 glk2* in hypocotyl

length, indicating that GLK2 may work independent (at least partly) of HY5 to

inhibit hypocotyl elongation, but again GLK2 plays a more important role than

GLK1.

Different from the hypocotyl phenotype of *glk* series mutants, after

germinating and growing under white light for 4 days, the hypocotyl length of *gnc*,

cga1 and *gnc cga1* mutants was shorter than that of the wild type, while the

hypocotyl length of *35S:GNC* and *35S:CGA1* overexpressed materials was longer

than that of the wild type. There was no difference in hypocotyl length between

these mutants and wild type after germinating and growing in darkness

(Supplemental Figure 4), indicating that GNC and CGA1 promoted hypocotyl

elongation under white light. In addition, the hypocotyl length of *hy5 gnc*, *hy5 cga1*

and *hy5 gnc cga1* mutants was shorter than that of *hy5* under white light, but

significantly longer than that of *gnc*, *cga1* and *gnc cga1* mutants **(Figures 4E and**

4F), indicating that GNC and CGA1 were able to reduce the inhibition of HY5 on

hypocotyl elongation. Therefore, during seedling development GLK transcription

factors can inhibit hypocotyl elongation, while GNC and CGA1 are able to

promote hypocotyl elongation, such that their functions as photomorphogenesis

regulators are expanded.

Transcriptome and RT-qPCR Analysis Reveals Differential Regulation of

Cell Elongation Genes by HY5, GLK2, and GNC

The above results showed that the phenotype of the *hy5 glk2* mutant was more

severe than that of the *hy5 glk1* mutant, both in regulating chlorophyll content

during seedling de-etiolation and inhibiting hypocotyl elongation during

light-induced seedling development. Hence, we selected wild type, *glk2*, *hy5* and

hy5 glk2 seedlings that germinated and grew for 4 days under white light for

transcriptome sequencing, and analysed the genes jointly or differentially

regulated by HY5 and GLK2. As shown in a Venn diagram, *glk2*, *hy5* and *hy5 glk2*

mutants jointly affected the expression of 142 genes, including 138

down-regulated genes and only 2 up-regulated genes **(Figure 5A)**. Deletion of

HY5 or *GLK2* in most cases resulted in down-regulation of genes, and K-means

clustering showed the same trend **(Figure 5B)**, in line with the expectation of

them as positive regulators of photomorphogenesis. Most up-regulated genes

were expressed in *hy5* and *hy5 glk2* mutants. Among the down-regulated genes,

385 were down-regulated together in *hy5* and *hy5 glk2* mutants, while 201 were

down-regulated together in *glk2* and *hy5 glk2* mutants. This is consistent with

HY5's upstream position in activating *GLK* gene expression mentioned above.

The remaining 91 down-regulated genes only in *hy5 glk2* mutants may represent

targets of combined HY5 and GLK activity **(Figure 5A)**. According to the trend of

glk2 < hy5 < hy5 glk2 in hypocotyl length, K4 and K7 groups from K-means

clustering could be considered for investigating GLK2 and HY5 regulated

hypocotyl elongation **(Figure 5B)**. In addition, a heat map was generated from the

differential expression of a list of known cell elongation related genes, and a

series of up-regulated genes were considered for subsequent validation **(Figure**

5C).

Elongation genes from **Figure 5C** were selected for RT-qPCR verification.

Many of them expressed higher in *glk2* than in the wild type, and higher in *hy5*

than in *glk2*, but the expression in *hy5 glk2* double mutant was more variable.

Among them, the expression levels of *EXPA5*, *IAA19*, *SAUR20*, and *PME16* were

detected higher in *hy5 glk2* than in *hy5* **(Figure 5D)**. We hypothesized that the

interaction between GLK2 and HY5 proteins could enhance the inhibitory effect on

these elongation genes, but GLK2 may independently inhibit other elongation

genes. We also analysed the expression levels of selected elongation genes from

Figure 5C in the *gnc* and *hy5 gnc* materials. The expression level of *IAA7* in *gnc*

mutant was lower than that in wild type, and the expression levels of several other

genes in *gnc* mutant tended to be lower than in wild type, although not significant

(Figure 5E). In addition, the expression levels of these elongation genes,

including *IAA7*, *IAA19*, *XTH4*, and *PME16*, appeared lower in the *hy5 gnc* double

mutant than in *hy5* mutant **(Figure 5E)**. These results indicate that GNC could

promote the expression of these elongation genes to less and varying degrees,

both independently or on top of HY5. It also suggests that the targeted hypocotyl

elongation genes by GNC/CGA1 may be largely different from those by GLKs.

Potential Protein Interaction of HY5 and GLK, and the Effect on Photosystem and Cell Elongation Genes

388
389

Apart from directly targeting many genes to promote their expression, HY5

also interacts with other proteins, HYH or HFR1 for instance, to mutually enhance

protein stability, or to bind to *HY5* or downstream gene promoters (Lee et al., 2007;

Ciolfi et al., 2013; Jang et al., 2013). The interaction between HY5 and G-box

binding factor 1 (GBF1) attenuates the activation of *RBCS-1A* by HY5. They share

a common G-box *cis*-element, which enables them to bind to the same target

promoter. This may enable more precisely regulated expression of downstream

genes and better adapts to the environment. Since HY5 interacts with GBF1, and

GBF1 interacts with GLK (Tamai et al., 2002; Singh et al., 2012; Ram et al., 2014),

we then examined the possibility of HY5 interacting with GLK proteins. In

pull-down experiments, the experimental group GST-GLK1 and GST-GLK2 could

pull down HY5 protein, while the control group could not **(Supplemental Figure

7A)**, indicating that HY5 binds to GLK1 and GLK2. Through bimolecular

luminescence complementation assays, it was found that luciferase signal could

be detected when HY5 was co-transfected with GLK1 and GLK2 proteins, while

the signal was not present in the negative controls **(Supplemental Figure 7B)**,

indicating that HY5 interacts with GLK1 and GLK2. The above experiments

provide *in vitro* evidence of HY5 and GLK interaction. HY5 might first activate the

expression of *GLK*, and then interacts with GLK protein to regulate common

targets. Dual-luciferase assay displayed that both HY5 and GLK2 could activate

LHCB1.2 expression, and GLK2 could activate to higher extent, but HY5 and

GLK2 together did not show an additive effect **(Supplemental Figure 7C)**, which

is consistent with the result shown in **Figure 2B**. **Supplemental Figure 7D**

showed that GLK2 could slightly activate *PME16* while HY5 inhibited its

expression, as indicated in **Figure 5C**. When HY5 and GLK2 were applied

together, it was more similar to the effect of GLK2 on *PME16* expression. This

appears discordant with the gene expression result in *hy5 glk2* mutant **(Figures**

5C and 5D), which might indicate that extra factors exist to coordinate HY5 and

GLK2 on their effects on *PME16*.

GLK2 Content Affects the Plastid Ultrastructure and Photosystem Gene

Expression in Dark-grown *det1* Mutant of Arabidopsis

Compared to the traits of photomorphogenesis, the morphology and gene

expression changes of skotomorphogenesis are relatively less addressed by

previous researches, although seedling development before they were exposed

to light represents a preparative and important stage. One of the few but well

known observations is that, the stromal thylakoid membrane was found

sufficiently extended in the plastids of etiolated *det1* mutant seedlings rather than

the default form of prolamella body, and some photosynthetic genes were already

expressed in the dark (Chory et al., 1989). On account of the emphasis on GLK2

from above, we constructed *det1 glk2* mutant and comparative observations were

made on the plastid ultrastructure in the cotyledons of dark-grown *det1*, *glk2*, and

det1 glk2 mutants. As expected, the cotyledons were open in the etiolated *det1*

glk2 mutant, and paler in the light-grown *det1 glk2* mutant **(Figure 6A)**. We found

that the thylakoid lamellae in the plastids of *det1 glk2* mutant were less developed

than in *det1* mutant in the dark, although in both cases there was little evidence of

a prolamellar body. By contrast, there was no significant structural difference

between the prolamellar body of the *glk2* mutant, the *GLK2*-overexpressing line,

and the wild type, but the size of the prolamellar body was smaller and larger than

that of the wild type respectively **(Figure 6B)**. The changes of plastid

ultrastructure in *det1 glk2* mutant suggested that developed chloroplasts in

etiolated *det1* mutant were related to GLK2 function, while in the dark-grown wild

type, DET1 may affect GLK2 protein stability, thus preventing chloroplast development.

It is known that in the dark, HY5 ubiquitination and degradation are mediated by the COP/DET/FUS complex (Saijo et al., 2003). In order to study the stability of GLK proteins in Arabidopsis, we examined the dark-grown wild type and *det1* mutant using Arabidopsis GLK1 and GLK2 antibodies for Western-blots. The results showed that both GLK1 and GLK2 protein contents in *det1* mutant were higher than those in the wild type **(Figure 7A)**. The accumulation of GLK1 and GLK2 proteins after DET1 deletion suggests that DET1 may promote the degradation of GLK proteins. It should be noted that Rubisco large subunit (RbcL) content was higher in *det1* than in the wild type, which is consistent with the increased accumulation of *RbcL* transcript in dark-grown *det1* mutant (Chory et al., 1989). We also grew the wild type and *det1* mutant in darkness for 4 days and treated them with MG132, a proteasome inhibitor, for 12 h in darkness with DMSO as the control. After MG132 treatment, the accumulation of GLK proteins in the wild type was higher than that in the control group, but still much lower than in the MG132 treated *det1* mutant **(Figure 7A)**. These results suggest that MG132 is beneficial to the accumulation of GLK1 and GLK2 proteins in the wild type by inhibiting proteasome activity, but the deficiency of DET1 may have blocked the upstream steps of protein degradation such as ubiquitination, so that the significant accumulation of GLK proteins in *det1* mutant maybe less affected. We therefore assessed whether GLK1 and GLK2 interact with DET1 protein in Arabidopsis. *In vitro* protein binding experiments support the expectation that DET1 interacts with both GLK1 and GLK2 proteins **(Supplemental Figure 8A)**. We also checked whether DET1 interacted with GNC and CGA1 proteins, and again positive results were obtained from *in vitro* protein binding experiments **(Supplemental Figure 8B)**.

The expression of some photosynthetic genes in dark-grown *det1* mutant

seedlings is an interesting phenomenon (Chory et al., 1989), but the mechanism

of the dark activation of these genes, or the significance of being turned off in

skotomorphogenesis, has not been elucidated. We measured the expression of

photosystem genes in a series of dark-grown *det1* and *glk2* related mutants, and

found that the expression levels of *LHCB1.2* and *LHCA1* were much higher in

det1 mutant than those in wild type, but were lower in *glk2* mutant than in the wild

type. The down-regulation of these genes by GLK2 was also found in *det1 glk2*

double mutant comparing with the *det1* mutant **(Figure 7B)**. This indicates that

GLK2 contributes significantly to the expression of photosystem genes in

dark-grown seedlings, and the effects could be additive to the loss of DET1

function. At the same time, the notably elevated expression of photosynthetic

genes in *det1* mutant should be contributed by the increased amount of GLK1,

GNC, or CGA1 as well. As supportive results, in the etiolated materials

overexpressing *GLK1*, *GLK2*, *GNC* or *CGA1*, the expression of *LHCB1.2* and

LHCA1 genes were significantly promoted in *GLK1* or *GLK2* overexpression lines,

less promoted in the *GNC* overexpression line, and least changed in the *CGA1*

overexpression line **(Figure 7C)**. When grown this same set of materials in the

light, increased expression of *LHCB1.2* and *LHCA1* genes were still observed to

larger extent in *GLK1* or *GLK2* overexpression lines than in *GNC* and *CGA1*

overexpression lines relative to the wild type **(Figure 7D)**. Notably, expression

levels of *LHCB1.2* and *LHCA1* genes also increased in both dark and light-grown

HY5 overexpression lines, but were not higher than those in *GLK1* or *GLK2*

overexpression lines, indicating again that GLKs function at least partly

independent of HY5 during seedling photomorphogenesis.

DISCUSSION

In this work, we have constructed a series of mutant materials to systematically

study the regulatory relationships between HY5, GLK, and GNC/CGA1

transcription factors, as well as their stabilities and functions within the scenario of

skotomorphogenesis and photomorphogenesis. We find that: (1) HY5 directly

activates *GLK, GNC*, and *CGA1* expression, and together they positively regulate

chlorophyll biosynthesis and photosystem formation; (2) GLK proteins inhibit

hypocotyl elongation, whereas GNC and CGA1 promote hypocotyl elongation, by

differentially regulating the expression of cell elongation genes; (3) In the dark,

DET1 destabilises GLK proteins, and the light-grown traits exhibited in the

etioplast of *det1* mutant seedlings can be attributed to GLK2 activity; (4) GLK2

maybe the predominant regulator of photosynthetic genes, during both

skotomorphogenesis and photomorphogenesis. At present, the molecular

mechanism of GLK and GNC in regulating hypocotyl elongation is not entirely

clear. These need to be further investigated in combination with specific genetic

material construction. In addition, GLK2 function has been standing out in multiple

ways, partially recruited by HY5 in the light, and effectively compensates for HY5

function, so that it deserves targeted studies in order to guide the potential

application of GLK2 and HY5 in crop enhancement.

GLK and GNC Function Downstream but Partially Independent of HY5 in

Seedling Photomorphogenesis

HY5 and GLK both functions in regulating chlorophyll biosynthesis and chloroplast

development, but targeted studies and clear conclusions are missing on whether

or how the two are related, especially during photomorphogenesis. Studies have

shown that PHYTOCHROME-INTERACTING FACTOR 4 (PIF4) in Arabidopsis

binds to the promoters of *GLK1* and *GLK2* to inhibit their expression (Song et al.,

2014), while HY5 antagonises PIFs to regulate the biosynthesis of photosynthetic

pigments (Tolede-Ortiz et al., 2014). Besides, microarray studies showed that

both phytochrome A and phytochrome B regulate GLK transcription (Tepperman

et al., 2006). Considering that HY5 acts downstream of the photoreceptors, and is

a high-level regulator (many HY5 targets are other transcription factors) of

photomorphogenesis (Lee et al., 2007), there may be a direct positive regulation

of HY5 to *GLKs*. We proved in this work that HY5 indeed binds to the promoters of

GLK genes and activates their expression, through EMSA *in vitro* assay, dual

luciferase assay, and ChIP-qPCR *in vivo* assay. Similarly, *in vitro* and *in vivo*

experiments also confirmed that HY5 binds to the promoters of *GNC* and *CGA1*,

and regulates their expression **(Figure 3)**.

We have constructed double and triple mutants of HY5, GLK1/2, and

GNC/CGA1 to study their functional relationships. Only limited decrease of

chlorophyll content and related gene expression were observed when *hy5 glk2*

mutant was compared to *glk2* mutant, or *hy5 glk1 glk2* mutant was compared to

the pale green *glk1 glk2* mutant **(Figures 1 and 2)**. Our transcriptome data

indicate that, compared with HY5, GLK2 tends to positively affect more PSII and

PSI related genes **(Supplemental Data Set 1-3)**. Furthermore, gene expression

analysis reveals that GLK2 exhibits stronger regulation on light harvesting genes

than HY5, while their regulatory strength on chlorophyll biosynthesis may be

similar during seedling greening **(Figure 2A and 2B)**. When overexpressing

GLK2 in the *hy5* mutant, the chlorophyll level increased significantly, although it

was still lower than that in the *GLK2* overexpression line with a wild type

background (Liu et al., 2021). HY5 was suggested to play an important role for

GLK2 function, while our study clarifies the functional downstream location of

GLKs to HY5, and implies that the complementary function of GLK1 and GLK2

contributes more intensively than HY5, to chlorophyll biosynthesis and chloroplast

development during Arabidopsis seedling greening.

It has been reported that HY5 and GLK2 are involved in the greening of

detached root in Arabidopsis, and it was suggested that the GLK2 regulation of

root greening depends to some extent on the presence of HY5 (Kobayashi et al.,

2012a). Using *hy5 glk* series mutants, we observed that the detached roots could

still turn green in the absence of a single GLK or both GLKs, but not in the

absence of HY5 **(Supplemental Figure 5)**. Endogenous cytokinin promotes the

greening process of detached roots (Kobayashi et al., 2012a). Our results confirm

that HY5 is essential in the process of root greening induced by cytokinin

accumulation. In order to investigate the effect of HY5 on cytokinin regulated leaf

greening, we added 6-Benzylaminopurine (6-BA) during seedling greening of *hy5*

glk series mutants. Results showed that 6-BA treatment promoted the

accumulation of chlorophyll during this process, with or without GLKs or HY5

(Supplemental Figure 6). These results suggest that GLKs and HY5 play similar

roles in response to cytokinin to promote chlorophyll biosynthesis in leaves,

whereas HY5 plays a more significant role in detached roots.

GLKs seem to also exert stronger regulation than GNC/CGA1 in chlorophyll

biosynthesis, as *glk1 glk2* mutants are much paler than *gnc cga1* mutants relative

to wild type **(Figure 1; Supplemental Figure 1)**. No difference in chlorophyll

content between *glk1 glk2 gnc cga1* quadruple mutant and *glk1 glk2* mutant was

found, and *CGA1* overexpression in *glk1 glk2* mutant could not restore the

chlorophyll content, while overexpression of *GLK1* partially restores the

chlorophyll content in *gnc cga1* mutants (Bastakis et al., 2018). According to our

work, the reduction of chlorophyll contents in *hy5 gnc* and *hy5 gnc cga1* mutants

compared to *hy5* mutant, was less obvious than that of *hy5 glk2* and *hy5 glk1 glk2*

mutants compared with *hy5* mutant **(Figures 1B and 1D)**. Zubo et al. suggested

that GLK1/GLK2 and GNC/CGA1 play both overlapping and independent roles in

regulating chlorophyll biosynthesis and chloroplast development (Zubo et al.,

2018). We further propose that GLKs play more prominent roles than GNC/CGA1,

downstream but partly independent of HY5, to regulate various targets during the

de-etiolation process of photomorphogenesis.

Differential Participation of GLK and GNC in HY5 Regulated Hypocotyl

Elongation

The inhibition of hypocotyl elongation, opening of cotyledon, chlorophyll

biosynthesis and chloroplast development are continuous and inseparable

processes during seedling photomorphogenesis. Since HY5 inhibits hypocotyl

elongation, we were also curious about the potential GLK and GNC regulation of

hypocotyls. It has been mentioned that *glk1* seedlings displayed longer hypocotyls

and less separated cotyledons (Martín et al., 2016, Alem et al., 2022). By

systematically analysing the hypocotyl phenotype of mutants and overexpression

lines, we certified that both GLK1 and GLK2 inhibit hypocotyl elongation, with

additive effects on the function of HY5 **(Figures 4A and 4B; Supplemental**

Figure 3). Transcriptomic sequencing and RT-qPCR verification showed that

GLK2 itself could inhibit the expression of elongation genes, and different

elongation genes exhibited different regulatory responses to GLK2 and HY5, such

that the inhibition of hypocotyl elongation in photomorphogenesis was the result of

comprehensive regulations.

Contrary to GLK's inhibition of hypocotyl elongation, we found that GNC and

CGA1 promote hypocotyl elongation **(Figures 4E and 4F; Supplemental Figure**

4). We noticed that the expression levels of some elongation genes were reported

lower in *gnc cga1* mutants than those in the wild type (Xu et al., 2017). Other

clues were found in the *quintuple* mutant of *GNC, GNL,* and *B-GATA* genes, that

the seedlings had shorter hypocotyl than wild type seedlings (Ranftl et al., 2016).

We further found in this work that the mutation of GNC or CGA1 displayed

negative effects on the inhibition of hypocotyl lengths by HY5 **(Figures 4E and**

4F). The expression of some tested elongation genes in *hy5 gnc* mutant was

decreased relative to that of *hy5* mutant **(Figure 5E)**. Our results indicate that

GNC could promote the expression of elongation genes to varying degrees,

independently or on top of HY5. It should be pointed out that the tested elongation

genes for GNC were initially from differentially expressed genes in *glk2* mutants,

so that they may not be the best regulated genes by GNC, and larger scale gene

expression analysis is needed to uncover the GNC/CGA1 specific elongation

genes. In addition, the changes of elongation genes in *glk1 glk2*, *gnc cga1*, *hy5*

glk1 glk2, and *hy5 gnc cga1* mutants are worth exploring in the future, to clarify

the potential redundant or additive effect of these transcription factors on

hypocotyl length.

Significance   of   GLK2   and   GNC   through   the   Context   from

Skotomorphogenesis to Photomorphogenesis

Traits of photomorphogenesis are negatively regulated by the COP/DET/FUS

complex in the dark, which mediates the ubiquitination and proteasome

degradation of positive regulators such as HY5, HYH, LAF1, and HFR1

(Osterlund et al., 2000; Holm et al., 2002; Seo et al., 2003; Yang et al., 2005).

Interestingly, *det1* and *cop1* mutants exhibit some of the characteristics of

photomorphogenesis during skotomorphogenesis, with shorter hypocotyls, open

cotyledons, and expression of photosynthetic genes (Chory et al., 1989; Deng et

al., 1992). These phenotypes should be related to the accumulation of positive

regulators of photomorphogenesis in darkness, but the specific regulators and

their influence on the phenotypes are not clear. It has been reported that

long-term ABA treatment activates the activity of COP1, which then interacts with

GLK1 protein to mediate its ubiquitination and degradation (Tokumaru et al., 2017;

Lee et al., 2021). Further, brassinolide or darkness induces GLK1 degradation by

inhibiting the BIN2 phosphorylation of GLK1, and it was speculated that

BIN2-mediated phosphorylation prevents the interaction of GLK1 and E3 ligases

(Zhang et al., 2021). In this study we further showed that DET1 is responsible for

the protein degradation of GLK1 and GLK2 in Arabidopsis seedlings **(Figures 8A**

and 8B). As for GLK2, we have constructed *det1 glk2* mutant in Arabidopsis to

compare its characters with *det1* mutant during skotomorphogenesis. Extended

thylakoid lamellae were found in the cotyledon protoplast of both mutants, but in

det1 glk2 mutant the overall number of lamellae was significantly reduced **(Figure**

6B). This is likely related with the GLK2 regulation on photosynthetic genes in

etiolated seedlings **(Figures 8C)**. In fact, both GLK2 and GLK1 contribute

significantly to the expression of photosystem genes, not only in the light but also

in dark-grown seedlings, while GNC and CGA1 seem to contribute less than

GLKs **(Figures 8D and 8E)**. Thus, the appropriate function of GLKs and

GNC/CGA1 is important to the whole process from skotomorphogenesis to

photomorphogenesis. They could be regulated at protein level by DET1 or COP1

related pathway in etiolated seedlings, while upon light HY5 takes the

transcriptional control and at least partly coordinates their functions, in a

regulation cascade of light, HY5, and GLKs or GNC/CGA1.

We constructed a working model as shown in **Figure 8**. In dark-grown

seedlings, GLK, GNC, CGA1 and HY5 proteins are subjected to the

COP/DET/FUS complex mediated degradation. Photosynthetic genes are inactive

and plastid inner membranes stay at prolamellar body status. After exposure to

light, COP/DET/FUS complex activity is inhibited and GLK, GNC, CGA1 and HY5

proteins accumulate. HY5 activates *GLK*, *GNC* and *CGA1* genes in a

light-dependent manner, and may interact with their proteins to cooperatively

regulate downstream gene expression to promote photomorphogenesis. GLKs
are master regulators of chloroplast development and able to inhibit hypocotyl
elongation independent of HY5. GNC and CGA1 are less dominant in chloroplast
development and able to promote hypocotyl elongation. These could be the
compensation for the relatively weak regulation of HY5 on chloroplast
development, and an efficient way to orchestrate light induced seedling
development, by recruiting multitasking yet complementary regulators. Our study
systematically reveals the new function of GLK and GNC regulating hypocotyl
length under light, refines the local network of HY5, GLK, GNC and CGA1, and
provides new perspective for better understanding the transition between dark
and light-grown seedlings.

MATERIALS AND METHODS

Plant Materials and Growth Conditions

All mutants used in this study were in the Columbia background. The *glk1*, *glk2*,
glk1 glk2, *gnc*, *cga1*, *gnc cga1*, *hy5*, and *det1* mutants have been previously
described (Chory et al., 1989; Oyama et al., 1997; Fitter et al., 2002; Chiang et al.,
2012). *35S:GLK1 glk1 glk2*, *35S:GLK2 glk1 glk2*, *35S:GNC*, and *35S:CGA1*
overexpression plants were also described previously (Waters et al., 2008;
Chiang et al., 2012). Other mutants described in this study were created by
crossing from the above materials. Surface-sterilized seeds were sown onto
half-strength Murashige and Skoog (MS) medium, which contained 1% sucrose
and 0.8% agar, and were cold-treated at 4°C in darkness for 3 days to ensure
synchronized germination, followed by growing in dark or light chambers
maintained at 22°C.

Chlorophyll Measurements

685 To measure the chlorophyll content, 4 d-old seedlings grown in the darkness or

686 continuous light were used. Total chlorophylls were extracted by homogenizing

687 the seedlings in 80% acetone, and leaving the solutions at 4°C for overnight

688 incubation. The absorbance was detected at wavelength OD_{652} with a

689 spectrophotometer. Chlorophyll concentration Ca+b was calculated by OD_{652}/34.5

690 (mg/mL), and then converted according to the fresh weight of the material (Arnon,

691 1949; Zhao et al., 2016).

693 **Hypocotyl Length Measurement**

694 To measure the hypocotyl length of seedlings, seeds were sown on plates and

695 stratified at 4°C in darkness for 3 d, and then kept in white light for 4 h in order to

696 induce uniform germination. The seeds were then transferred to white, blue, red

697 light, or dark conditions and incubated at 22°C for 4 d. The hypocotyl length were

698 measured using ImageJ software after photographing the seedlings.

700 **Quantitative RT-PCR**

701 Total RNA was extracted from wild type and mutant seedlings using an RNAiso

702 Plus kit (Takara). Complementary DNAs (cDNA) were synthesized from 2 μg of

703 total RNA using a Prime Script RT Reagent Kit, with a genomic DNA Eraser

704 (Yeasen) according to the manufacturer's instructions. Then, cDNA was subjected

705 to RT-qPCR assays, using the Step One Plus RT-PCR detection system (Applied

706 Biosystems) and SYBR Green PCR Master Mix (Yeasen). PCR was performed in

707 triplicate for each sample, and the expression levels were normalized to that of

708 *ACT7* gene.

710 **Immunoblot Analysis**

711 For immunoblot analysis, wild type and mutant seedlings were homogenized in a

protein extraction buffer containing 100 mM Tris (pH 6.8), 10% Glycerol, 0.5% SDS, 0.1% Triton, 5 mM EDTA, 0.01 M DTT, and 1 × complete protease inhibitor cocktail (Roche). Equal amount of proteins were loaded and separated on a 10% SDS-PAGE gel and transferred to polyvinylidene fluoride membrane. The membrane was blocked with 5% milk and incubated with primary antibody overnight at 4°C. Primary antibodies anti-AtGLK1/2 were used with 1:2000 dilution. The membrane was then washed four times with 1 × Phosphate Buffered Saline Tween-20 (PBST) for 5 minutes, and incubated with secondary antibody (1:10000 dilution) for 1 h at room temperature. After four washes with PBST for 5 minutes, signal was detected with ECL kit (Millipore) according to manufacturer's protocol.

EMSA Assay

Synthetic complementary oligonucleotides of *GLK1*, *GLK2*, *GNC*, and *CGA1* (or mutant probe with HY5 binding motif mutation) were obtained, and probes were PCR amplified using Cy5-labeled primers. For proteins, the sequence encoding HY5 was cloned into the pCold-TF vector, expressed, and purified with Capturem™ His-Tagged Purification Maxiprep Kit (Takara). The binding reaction was performed in 20 µL binding buffer (10 mM Tris pH 8.0, 1 mM KCl, 4 mM $MgCl_2$, 0.5mM DTT, 5% glycerol, 0.2 mM EDTA, and 0.01% BSA) using 15 nM probes and 200 ng proteins, and incubated at room temperature for 30 min. The reactions were resolved by electrophoresis in a 6% (v/v) native polyacrylamide gel at 4°C. Cy5-labeled DNA in the gel was scanned by an Amersham Typhoon 5 Biomolecular Imager.

Dual-luciferase Reporter System

Promoter sub-fragments of *GLK1*, *GLK2*, *GNC*, and *CGA1* were cloned into the pGreenII 0800-LUC vector to drive the firefly luciferase gene. p2GWY7-HY5-YFP

was used as the effector construct. Arabidopsis mesophyll cell protoplasts were

isolated and transfected as described previously (Yoo et al., 2007). We used a

Dual Luciferase kit (Yeasen) for transient expression analysis to detect reporter

activity. The Renilla luciferase gene, driven by the cauliflower mosaic virus 35S

promoter, was used as an internal control. The ratio of LUC/REN was calculated

as an indicator of the final transcriptional activity.

Chromatin Immunoprecipitation (ChIP)

The ChIP assay was performed as described by Xu et al. (2016). Chromatin

isolation was performed using Col and *HA-HY5 hy5-215* transgenic seedlings

grown under constant white light for 4 d. The resuspended chromatin was

sonicated at 4°C to 250- to 500-bp fragments. The sheared chromatin was

immunoprecipitated, washed, reverse cross linked, and finally amplified. About 10%

of sonicated but non-immunoprecipitated chromatin was reverse cross linked and

used as an input DNA control. Both immunoprecipitated DNA and input DNA were

analyzed by RT-PCR (Applied Biorad). Monoclonal anti-HA antibody (ab9110,

Abcam, 1:100 dilution) was used for the immunoprecipitation. The level of binding

was calculated as the ratio between the IP and Input groups.

In vitro **Pull-downs**

The full-length coding sequences of *GLK1*, *GLK2*, *GNC*, and *CGA1* were cloned

into pGEX-6P-1 vector, and the full-length coding sequences of *HY5* and *nDET1*

(26-87 aa) were cloned into pCold-TF vector. Two mL of transformed bacteria

cultures were grown over night, diluted 1:100 in the next morning, and grown for

another 3 hours at 37°C. When the cells reached logarithmic phase, 0.1 mM IPTG

was added to induce the expression of proteins at 37°C for 4 hours. Then 0.5 mL

of each selected *E. coli* cultures was added in the same tube, resuspended in

lysis buffer (1 × PBS, pH 7.4, 1 × complete protease inhibitor cocktail), and the

cells were broken with a sonicator for 3 times (30 s on and 60 s off) on ice. After

the cell debris were removed by centrifugation at 18000 g for 30 min, 20 µL

Glutathione Sepharose 4B was added to the mixture sample and rotated at 4°C

for 6 hours. After 4 washes with PBST buffer (1 × PBS, pH 7.4, 0.1% Triton X-100),

the pellet fraction was boiled in 5 × SDS protein loading buffer, and the input and

pull-down prey proteins were detected by immunoblot analysis using anti-His

(M201, Takara, 1:3000 dilution) and anti-GST (G018, Abcam, 1:3000 dilution)

antibodies respectively.

Bi-luminescence Complementation (BiLC) Assay

GLK or HY5 was fused to the N- or C-terminus of firefly luciferase, and the

constructs were transformed into agrobacterium (*Agrobacterium tumefaciens*)

strain GV3101. Overnight cultures of agrobacteria were collected by

centrifugation at 4000 g for 10 min, resuspended in MES buffer (10 mM MES,10

mM $MgCl_2$, and 100 mM acetosyringone), mixed with GV3101 colonies

expressing pSoup-P19 to a final OD_{600} = 0.5, and incubated at room temperature

for 3 h in the dark before infiltration. The agrobacterium suspension in a 1 mL

syringe (without the metal needle) was carefully press-infiltrated onto healthy

leaves of 3-week-old *N. benthamiana*. The infiltrated plants were returned to

long-day conditions for 3 d. Leaves were infiltrated with luciferin solution and left

for 10 min, before observing luciferase activity by imaging with a CCD camera

(Tanon-5200, BioTanon, China).

RNA-seq Analysis

Total RNA was extracted from the 4-d-old Col-0, *glk2*, *hy5*, and *hy5 glk2* seedlings

grown in constant white light (100 µmol photons m^{-2} s^{-1}) conditions. Sequencing

libraries were generated with three independent biological replicates for each

material, and the sequencing was performed using Illumina platform by Biomarker

company (Beijing). Clean reads were mapped to the Arabidopsis reference

genome (TAIR11.0_NCBI) using Hisat2 (Version 2.2.1.0). Read counts per gene

and Fragments Per Kilobase exon per Million reads mapped (FPKM) values were

quantified using Cufflinks (Version 2.2.1), while read count values per transcript

(protein coding) were calculated using Bowtie2 and eXpress. Differentially

expressed genes (DEGs) were identified using DESeq (Version1.18.0) (Anders et

al., 2013). The R package functions 'Estimate Size Factors' and 'nbinom Test',

Padj values<0.05 and Fold Changes >1 were set as the threshold for significantly

different expression.

Accession Numbers

Sequence information from this article can be found in The Arabidopsis

Information Resource (TAIR) under the following accession numbers: *GLK1*

(AT2G20570), *GLK2* (AT5G44190), *GNC* (AT5G56860), *CGA1* (AT4G26150),

DET1 (AT4G10180), *HY5* (At5G11260), *ACT7* (At5G09810), *HEMA1*

(AT1G58290), *PORB* (AT4G27440), *PORC* (AT1G03630), *CAO* (AT1G44446),

LHCB1.2 (AT1G29910), *LHCB2.2* (AT2G05070), *LHCA1* (AT3G54890), *LHCA4*

(AT3G47470), *EXPA5* (AT3G29030), *IAA19* (AT3G15540), *SAUR20*

(AT5G18020), *PME16* (AT2G43050), *IAA7* (AT3G23050), and *XTH4*

(AT2G06850).

Supplemental Data

Supplemental Figure 1. HY5, GLK, GNC and CGA1 regulate the chlorophyll

biosynthesis.

Supplemental Figure 2. Phenotypes of the mutants and overexpression lines

under continuous white light conditions.

Supplemental Figure 3. GLK inhibits hypocotyl elongation.

Supplemental Figure 4. GNC and CGA1 promote hypocotyl elongation.

Supplemental Figure 5. HY5 but not GLKs is crucial for the greening of detached

roots.

Supplemental Figure 6. 6-BA promotes GLK and HY5 dependent chlorophyll

biosynthesis.

Supplemental Figure 7. *In vitro* Evidence of HY5 Interacting with GLK and

Potential Co-regulation of Target Genes.

Supplemental Figure 8. *In vitro* evidence of DET1 interacting with GLK and

GNC/CGA1.

Supplemental Dataset 1. List of 516 GLK2 regulated genes.

Supplemental Dataset 2. List of 656 HY5 regulated genes.

Supplemental Dataset 3. List of 716 HY5 and GLK2 regulated genes.

Supplemental Data 4. Primers used in this study.

Supplemental Data 5. Statistical analysis for the data shown in figures.

Acknowledgments

We thank Prof. Jane Langdale for comments and suggestions on the work. We

thank Xiaoyan Gao and Zhiping Zhang for technical support on the transmission

electron microscopy. We are grateful to Prof. Honghui Lin for kindly providing the

AtGLK1 antibody; Chunguang Chen from ORIZYMES for generating the AtGLK2

antibody; Prof. G. Eric Schaller for gift of Arabidopsis GNC/CGA1 mutants and

overexpression lines. This work was supported by the National Natural Science

Foundation of China (No. 31970257).

Author Contributions

TZ and PW designed the study. TZ and RZ performed most of the experiments.

XZ worked on the transcriptional activation and protein interaction. LY worked on

the ChIP-qPCR experiments. ST analyzed the RNA-seq data. PW, YZ, WZ, and

XZ wrote and revised the manuscript.

REFERENCES

Abbas, N., Maurya, J.P., Senapati, D., Gangappa, S.N., and Chattopadhyay, S.

(2014). Arabidopsis CAM7 and HY5 physically interact and directly bind to

the HY5 promoter to regulate its expression and thereby promote

photomorphogenesis. Plant Cell **26:** 1036-1052.

Alem, A.L., Ariel, F.D., Cho, Y., Hong, J.C., Gonzalez, D.H., and Viola, I.L.

(2022). TCP15 interacts with GOLDEN2-LIKE 1 to control cotyledon opening

in Arabidopsis. Plant J. **110(3):** 748-763.

Anders, S., and Hubre, W. (2013). Differential expression of RNA-Seq data at

the gene level-the DESeq package. European Molecular Biology Laboratory

(EMBL).

Arnon, D.I. (1949). Copper enzymes in isolated chloroplasts polypheoloxidase in

Bata Vulgaris. Plant Physiol. **24:** 1-15.

Bastakis, E., Hedtke, B., Klermund, C., Grimm, B., and Schwechheimer, C.

(2018). LLM-Domain B-GATA transcription factors play multifaceted roles in

controlling greening in Arabidopsis. Plant Cell **30:** 582-599.

Bi, Y.M., Zhang, Y., Signorelli, T., Zhao, R., Zhu, T., and Rothstein, S. (2005).

Genetic analysis of Arabidopsis GATA transcription factor gene family reveals

a nitrate-inducible member important for chlorophyll synthesis and glucose

sensitivity. Plant J. **44:** 680-692.

Binkert, M., Kozma-Bognár, L., Terecskei, K., De Veylder, L., Nagy, F., and

Ulm, R. (2014). UV-B-responsive association of the Arabidopsis bZIP

transcription factor ELONGATED HYPOCOTYL5 with target genes, including

875  its own promoter. Plant Cell **26:** 4200-4213.

876 **Catalá, R., Medina, J., and Salinas, J.** (2011). Integration of low temperature and

877  light signaling during cold acclimation response in Arabidopsis. Proc. Natl.

878  Acad. Sci. U. S. A. **108:** 16475-16480.

879 **Chang, C.S., Li, Y.H., Chen, L.T., Chen, W.C., Hsieh, W.P., Shin, J., Jane, W.N.,**

880  **Chou, S.J., Choi, G., Hu, J.M., Somerville, S., and Wu, S.H.** (2008). LZF1,

881  a HY5-regulated transcriptional factor, functions in Arabidopsis de-etiolation.

882  Plant J. **54:** 205-219.

883 **Chen, D., Xu, G., Tang, W., Jing, Y., Ji, Q., Fei, Z., and Lin, R.** (2013).

884  Antagonistic basic helix-loop-helix/bZIP transcription factors form

885  transcriptional modules that integrate light and reactive oxygen species

886  signaling in Arabidopsis. Plant Cell **25:** 1657-1673.

887 **Chiang, Y.H., Zubo, Y.O., Tapken, W., Kim, H.J., Lavanway, A.M., Howard, L.,**

888  **Pilon, M., Kieber, J.J., and Schaller, G.E.** (2012). Functional

889  characterization of the GATA transcription factors GNC and CGA1 reveals

890  their key role in chloroplast development, growth, and division in Arabidopsis.

891  Plant Physiol. **160:** 332-348.

892 **Chory, J., Peto, C., Feinbaum, R., Pratt, L., and Ausubel, F.** (1989).

893  Arabidopsis thaliana mutant that develops as a light-grown plant in the

894  absence of light. Cell **58:** 991-999.

895 **Ciolfi, A., Sessa, G., Sassi, M., Possenti, M., Salvucci, S., Carabelli, M.,**

896  **Morelli, G., and Ruberti, I.** (2013). Dynamics of the shade-avoidance

897  response in Arabidopsis. Plant Physiol. **163:** 331-353.

898 **Datta, S., Hettiarachchi, C., Johansson, H., and Holm, M.** (2007). SALT

899  TOLERANCE HOMOLOG2, a B-box protein in Arabidopsis that activates

900  transcription and positively regulates light-mediated development. Plant Cell

901  **19:** 3242-3255.

Datta, S., Johansson, H., Hettiarachchi, C., Irigoyen, M.L., Desai, M., Rubio,
V., and Holm, M. (2008). LZF1/SALT TOLERANCE HOMOLOG3, an
Arabidopsis B-box protein involved in light-dependent development and gene
expression, undergoes COP1-mediated ubiquitination. Plant Cell **20:**
2324-2338.

Deng, X.W., Matsui, M., Wei, N., Wagner, D., Chu, A.M., Feldmann, K.A., and
Quail, P.H. (1992). COP1, an Arabidopsis regulatory gene, encodes a protein
with both a zinc-binding motif and a G beta homologous domain. Cell **71:**
791-801.

Fitter, D.W., Martin, D.J., Copley, M.J., Scotland, R.W., and Langdale, J.A.
(2002). GLK gene pairs regulate chloroplast development in diverse plant
species. Plant J. **31:** 713-727.

Hall, L.N., Rossini, L., Cribb, L., and Langdale, J.A. (1998). GOLDEN 2: a
novel transcriptional regulator of cellular differentiation in the maize leaf.
Plant Cell **10:** 925-936.

Holm, M., Ma, L.G., Qu, L.J., and Deng, X.W. (2002). Two interacting bZIP
proteins are direct targets of COP1-mediated control of light-dependent gene
expression in Arabidopsis. Genes Dev. **16:** 1247-1259.

Huang, X., Ouyang, X., Yang, P., Lau, O.S., Li, G., Li, J., Chen, H., and Deng,
X.W. (2012). Arabidopsis FHY3 and HY5 positively mediate induction of
COP1 transcription in response to photomorphogenic UV-B light. Plant Cell
24: 4590-4606.

Hudson, D., Guevara, D.R., Hand, A.J., Xu, Z., Hao, L., Chen, X., Zhu, T., Bi,
Y.M., and Rothstein, S.J. (2013). Rice cytokinin GATA transcription Factor1
regulates chloroplast development and plant architecture. Plant Physiol. **162:**
132-144.

Imamura, A., Hanaki, N., Nakamura, A., Suzuki, T., Taniguchi, M., Kiba, T.,

Ueguchi, C., Sugiyama, T., and Mizuno, T. (1999). Compilation and
characterization of Arabidopsis thaliana response regulators implicated in
His-Asp phosphorelay signal transduction. Plant Cell Physiol. **40:** 733-742.

Jakoby, M., Weisshaar, B., Dröge-Laser, W., Vicente-Carbajosa, J.,
Tiedemann, J., Kroj, T., Parcy, F., and bZIP Research Group. (2002). bZIP
transcription factors in Arabidopsis. Trends Plant Sci. **7:** 106-111.

Jang, I.C., Henriques, R., and Chua, N.H. (2013). Three transcription factors,
HFR1, LAF1 and HY5, regulate largely independent signaling pathways
downstream of phytochrome A. Plant Cell Physiol. **54:** 907-916.

Jiao, Y., Lau, O.S., and Deng, X.W. (2007). Light-regulated transcriptional
networks in higher plants. Nat. Rev. Genet. **8:** 217-230.

Jing, Y., Zhang, D., Wang, X., Tang, W., Wang, W., Huai, J., Xu, G., Chen, D., Li,
Y., and Lin, R. (2013). Arabidopsis chromatin remodeling factor PICKLE
interacts with transcription factor HY5 to regulate hypocotyl cell elongation.
Plant Cell **25:** 242-256.

Kobayashi, K., Baba, S., Obayashi, T., Sato, M., Toyooka, K., Keränen, M.,
Aro, E.M., Fukaki, H., Ohta, H., Sugimoto, K., and Masuda, T. (2012a).
Regulation of root greening by light and auxin/cytokinin signaling in
Arabidopsis. Plant Cell **24:** 1081-1095.

Kobayashi, K., Fujii, S., Sasaki, D., Baba, S., Ohta, H., Masuda, T., and Wada,
H. (2014). Transcriptional regulation of thylakoid galactolipid biosynthesis
coordinated with chlorophyll biosynthesis during the development of
chloroplasts in Arabidopsis. Front Plant Sci. **5:** 272.

Kobayashi, K., Obayashi, T., and Masuda, T. (2012b). Role of the G-box
element in regulation of chlorophyll biosynthesis in Arabidopsis roots. Plant
Signal.Behav. **7:** 922-926.

Kobayashi, K., Ohnishi, A., Sasaki, D., Fujii, S., Iwase, A., Sugimoto, K.,

Masuda, T., and Wada, H. (2017). Shoot Removal Induces Chloroplast

Development in Roots via Cytokinin Signaling. Plant Physiol. **173:**

2340-2355.

Kushwaha, R., Singh, A., and Chattopadhyay, S. (2008). Calmodulin7 plays an

important role as transcriptional regulator in Arabidopsis seedling

development. Plant Cell **20:** 1747-1759.

Lee, D., Hua, L., Khoshravesh, R., Giuliani, R., Kumar, I., Cousins, A., Sage, T.

L., Hibberd, J. M., and Brutnell, T. P. (2021). Engineering chloroplast

development in rice through cell‑specific control of endogenous genetic

circuits. Plant Biotechnol. J. **19(11):** 2291-2303.

Lee, J., Choi, B., Yun, A., Son, N., Ahn, G., Cha, J.Y., Kim, W.Y., and Hwang, I.

(2021). Long-term abscisic acid promotes golden2-like1 degradation through

constitutive photomorphogenic 1 in a light intensity-dependent manner to

suppress chloroplast development. Plant Cell Environ. **44:** 3034-3048.

Lee, J., He, K., Stolc, V., Lee, H., Figueroa, P., Gao, Y., Tongprasit, W., Zhao,

H., Lee, I., and Deng, X.W. (2007). Analysis of transcription factor HY5

genomic binding sites revealed its hierarchical role in light regulation of

development. Plant Cell **19:** 731-749.

Liu, D., Zhao, D., Li, X., and Zeng, Y.J. (2022) AtGLK2, an Arabidopsis

GOLDEN2-LIKE transcription factor, positively regulates anthocyanin

biosynthesis via AtHY5-mediated light signaling. Plant Growth Regul. **96:**

79-90.

Martín, G., Leivar, P., Ludevid, D., Tepperman, J.M., Quail, P.H., and Monte, E.

(2016). Phytochrome and retrograde signalling pathways converge to

antagonistically regulate a light-induced transcriptional network. Nat.

Commun. **7:** 11431.

Naito, T., Kiba, T., Koizumi, N., Yamashino, T., and Mizuno, T. (2007).

Characterization of a unique GATA family gene that responds to both light
and cytokinin in Arabidopsis thaliana. Biosci. Biotechnol. Biochem. **71:**
1557-1560.

Nakamura, H., Muramatsu, M., Hakata, M., Ueno, O., Nagamura, Y., Hirochika,
H., Takano, M., and Ichikawa, H. (2009). Ectopic overexpression of the
transcription factor OsGLK1 induces chloroplast development in non-green
rice cells. Plant Cell Physiol. **50:** 1933-1949.

Nguyen, C.V., Vrebalov, J.T., Gapper, N.E., Zheng, Y., Zhong, S., Fei, Z., and
Giovannoni, J.J. (2014). Tomato GOLDEN2-LIKE transcription factors
reveal molecular gradients that function during fruit development and ripening.
Plant Cell **26:** 585-601.

Osterlund, M.T., Hardtke, C.S., Wei, N., and Deng, X.W. (2000). Targeted
destabilization of HY5 during light-regulated development of Arabidopsis.
Nature **405:** 462-466.

Oyama, T., Shimura, Y., and Okada, K. (1997). The Arabidopsis HY5 gene
encodes a bZIP protein that regulates stimulus-induced development of root
and hypocotyl. Genes Dev. **11:** 2983-2995.

Ram, H., Priya, P., Jain, M., and Chattopadhyay, S. (2014). Genome-wide DNA
binding of GBF1 is modulated by its heterodimerizing protein partners, HY5
and HYH. Mol. Plant **7:** 448-451.

Ranftl, Q.L., Bastakis, E., Klermund, C., and Schwechheimer, C. (2016).
LLM-Domain Containing B-GATA Factors Control Different Aspects of
Cytokinin-Regulated Development in Arabidopsis thaliana. Plant Physiol.
170(4): 2295-2311.

Rauf, M., Arif, M., Dortay, H., Matallana-Ramírez, L.P., Waters, M.T., Gil Nam,
H., Lim, P.O., Mueller-Roeber, B., and Balazadeh, S. (2013). ORE1
balances leaf senescence against maintenance by antagonizing

G2-like-mediated transcription. Embo Rep. **14:** 382-388.

Rossini, L., Cribb, L., Martin, D.J., and Langdale, J.A. (2001). The maize golden2 gene defines a novel class of transcriptional regulators in plants. Plant Cell **13:** 1231-1244.

Saijo, Y., Sullivan, J.A., Wang, H., Yang, J., Shen, Y., Rubio, V., Ma, L., Hoecker, U., and Deng, X.W. (2003). The COP1-SPA1 interaction defines a critical step in phytochrome A-mediated regulation of HY5 activity. Genes Dev. **17:** 2642-2647.

Seo, H.S., Yang, J.Y., Ishikawa, M., Bolle, C., Ballesteros, M.L., and Chua, N.H. (2003). LAF1 ubiquitination by COP1 controls photomorphogenesis and is stimulated by SPA1. Nature **423:** 995-999.

Shin, J., Park, E., and Choi, G. (2007). PIF3 regulates anthocyanin biosynthesis in an HY5-dependent manner with both factors directly binding anthocyanin biosynthetic gene promoters in Arabidopsis. Plant J. **49:** 981-994.

Singh, A., Ram, H., Abbas, N., and Chattopadhyay, S. (2012). Molecular interactions of GBF1 with HY5 and HYH proteins during light-mediated seedling development in Arabidopsis thaliana. J. Biol. Chem. **287:** 25995-26009.

Song, Y., Yang, C., Gao, S., Zhang, W., Li, L., and Kuai, B. (2014). Age-triggered and dark-induced leaf senescence require the bHLH transcription factors PIF3, 4, and 5. Mol. Plant **7:** 1776-1787.

Song, Y.H., Yoo, C.M., Hong, A.P., Kim, S.H., Jeong, H.J., Shin, S.Y., Kim, H.J., Yun, D.J., Lim, C.O., Bahk, J.D., Lee, S.Y., Nagao, R.T., Key, J.L., and Hong, J.C. (2008). DNA-binding study identifies C-box and hybrid C/G-box or C/A-box motifs as high-affinity binding sites for STF1 and LONG HYPOCOTYL5 proteins. Plant Physiol. **146:** 1862-1877.

Stracke, R., Favory, J.J., Gruber, H., Bartelniewoehner, L., Bartels, S.,

Binkert, M., Funk, M., Weisshaar, B., and Ulm, R. (2010). The Arabidopsis

bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene

in response to light and ultraviolet-B radiation. Plant Cell Environ. **33:** 88-103.

Sullivan, J.A., and Deng, X.W. (2003). From seed to seed: the role of

photoreceptors in Arabidopsis development. Dev. Biol. **260:** 289-297.

Tamai, H., Iwabuchi, M., and Meshi, T. (2002). Arabidopsis GARP transcriptional

activators interact with the Pro-rich activation domain shared by

G-box-binding bZIP factors. Plant Cell Physiol. **43:** 99-107.

Tang, X., Miao, M., Niu, X., Zhang, D., Cao, X., Jin, X., Zhu, Y., Fan, Y., Wang,

H., Liu, Y., Sui, Y., Wang, W., Wang, A., Xiao, F., Giovannoni, J., and Liu, Y.

(2016). Ubiquitin-conjugated degradation of golden 2-like transcription factor

is mediated by CUL4-DDB1-based E3 ligase complex in tomato. New Phytol.

209: 1028-1039.

Tepperman, J.M., Hwang, Y.-S., and Quail, P.H. (2006). phyA dominates in

transduction of red-light signals to rapidly responding genes at the initiation of

Arabidopsis seedling de-etiolation. Plant J. **48:** 728-742.

Tokumaru, M., Adachi, F., Toda, M., Ito-Inaba, Y., Yazu, F., Hirosawa, Y.,

Sakakibara, Y., Suiko, M., Kakizaki, T., and Inaba, T. (2017).

Ubiquitin-Proteasome Dependent Regulation of the GOLDEN2-LIKE 1

Transcription Factor in Response to Plastid Signals. Plant Physiol. **173:**

524-535.

Toledo-Ortiz, G., Johansson, H., Lee, K.P., Bou-Torrent, J., Stewart, K., Steel,

G., Rodríguez-Concepción, M., and Halliday, K. J. (2014). The HY5-PIF

regulatory module coordinates light and temperature control of

photosynthetic gene transcription. PLoS Genet. **10:** e1004416.

Wang, P., Khoshravesh, R., Karki, S., Tapia, R., Balahadia, C. P.,

Bandyopadhyay, A., Quick, W. P., Furbank, R., Sage, T. L., and Langdale,

J. A. (2017). Re-creation of a Key Step in the Evolutionary Switch from C (3)

to C (4) Leaf Anatomy. Curr. Biol. **27:** 3278-3287.e3276.

Waters, M.T., Moylan, E.C., and Langdale, J.A. (2008). GLK transcription

factors regulate chloroplast development in a cell-autonomous manner. Plant

J. **56:** 432-444.

Waters, M.T., Wang, P., Korkaric, M., Capper, R.G., Saunders, N.J., and

Langdale, J.A. (2009). GLK transcription factors coordinate expression of

the photosynthetic apparatus in Arabidopsis. Plant Cell **21:** 1109-1128.

Wykoff, D.D., Grossman, A.R., Weeks, D.P., Usuda, H., and Shimogawara, K.

(1999). Psr1, a nuclear localized protein that regulates phosphorus

metabolism in Chlamydomonas. Proc. Natl. Acad. Sci. U. S. A. **96:**

15336-15341.

Xu, D., Jiang, Y., Li, J., Lin, F., Holm, M., and Deng, X.W. (2016a). BBX21, an

Arabidopsis B-box protein, directly activates HY5 and is targeted by COP1 for

26S proteasome-mediated degradation. Proc. Natl. Acad. Sci. U. S. A. **113:**

7655-7660.

Xu, X., Chi, W., Sun, X., Feng, P., Guo, H., Li, J., Lin, R., Lu, C., Wang, H.,

Leister, D., and Zhang, L. (2016b). Convergence of light and chloroplast

signals for de-etiolation through ABI4-HY5 and COP1. Nat. Plants **2:** 16066.

Xu, Z., Casaretto, J.A., Bi, Y.M., and Rothstein, S.J. (2017). Genome-wide

binding analysis of AtGNC and AtCGA1 demonstrates their cross-regulation

and common and specific functions. Plant Direct **1:** e00016.

Yang, J., Lin, R., Sullivan, J., Hoecker, U., Liu, B., Xu, L., Deng, X. W., and

Wang, H. (2005). Light regulates COP1-mediated degradation of HFR1, a

transcription factor essential for light signaling in Arabidopsis. Plant Cell **17:**

804-821.

Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). Arabidopsis mesophyll protoplasts: a

versatile cell system for transient gene expression analysis. Nat. Protoc. **2(7):**
1565-1572.

Zhang, D., Tan, W., Yang, F., Han, Q., Deng, X., Guo, H., Liu, B., Yin, Y., and
Lin, H. (2021). A BIN2-GLK1 Signaling Module Integrates Brassinosteroid
and Light Signaling to Repress Chloroplast Development in the Dark. Dev.
Cell **56:** 310-324 e317.

Zhang, H., He, H., Wang, X., Wang, X., Yang, X., Li, L., and Deng, X.W. (2011).
Genome-wide mapping of the HY5-mediated gene networks in Arabidopsis
that involve both transcriptional and post-transcriptional regulation. Plant J.
65: 346-358.

Zhao, Y.R., Li, X., Yu, K.Q., Cheng, F., and He, Y. (2016). Hyperspectral Imaging
for Determining Pigment Contents in Cucumber Leaves in Response to
Angular Leaf Spot Disease. Sci. Rep. **6:** 27790.

Zubo, Y.O., Blakley, I.C., Franco-Zorrilla, J.M., Yamburenko, M.V., Solano, R.,
Kieber, J.J., Loraine, A.E., and Schaller, G.E. (2018). Coordination of
Chloroplast Development through the Action of the GNC and GLK
Transcription Factor Families. Plant Physiol. **178:** 130-147.

1108
1109

1110

1111

Figure 1. GLK1/2 and GNC/CGA1 Positively Regulate Chlorophyll Content Downstream of HY5 during De-etiolation.

(A) Phenotypes of 4-day-old etiolated seedlings of Col-0, *glk1*, *glk2*, *glk1 glk2*, *hy5*,

hy5 glk1, *hy5 glk2*, and *hy5 glk1 glk2* during transition from dark to light conditions

for 6 h, 12 h, 24 h. Scale bars = 1 mm.

(B) Phenotypes of 4-day-old etiolated seedlings of Col-0, *gnc*, *cga1*, *gnc cga1*,

hy5, *hy5 gnc*, *hy5 cga1*, and *hy5 gnc cga1* during transition from dark to light

conditions for 6 h, 12 h, 24 h. Scale bars = 1 mm.

(C, D) Chlorophyll contents of de-etiolating seedlings during transition from (A)

and (B) dark to light for 6 h and 12 h. The data represent means ± SD (n = 3) and

letters above the bars indicate significant differences ($P < 0.05$), as determined by

one-way ANOVA with Turkey's HSD test.

(E) The expression levels of *GLK1*, *GLK2*, *GNC* and *CGA1* in 4-d-old dark-grown

Col-0 upon being transferred to white light at indicated time points. The *ACT7*

gene was used as internal control. The data represent means ± SD (n = 3) and

letters above the bars indicate significant differences ($P < 0.05$), as determined by

one-way ANOVA with Turkey's HSD test.

(F) The expression levels of *GLK1*, *GLK2*, *GNC* and *CGA1* in 4-d-old dark-grown

Col-0 and *hy5* seedlings upon being transferred to white light at indicated time

points. The data represent means ± SD (n = 3), and asterisks indicate a significant

difference compared with Col-0 (*$P<0.05$, **$P<0.01$, paired samples *t*-test).

1134

1135

1136

1137

1138 **Figure 2.** Chloroplast Development is Further Impaired in *hy5* and *glk* Multiple
1139 Mutants.

1140

1141 **(A, B)** RT-qPCR analysis of chlorophyll biosynthesis and photosystem genes

expressed in 4-d-old etiolated seedlings of Col-0, *hy5, glk2* and *hy5 glk2* upon being transferred to white light for 12 h. The *ACT7* gene was used as internal control. The data represent means ± SD (n = 3) and letters above the bars indicate significant differences ($P < 0.05$), as determined by one-way ANOVA with Turkey's HSD test.

(C) Phenotypes of 4-day-old Col-0, *glk1, glk2, glk1 glk2, hy5, hy5 glk1, hy5 glk2,* and *hy5 glk1 glk2* seedlings grown on half-strength MS plates under continuous white light conditions. Scale bars = 1 mm.

(D) Chlorophyll contents of the genotypes shown in **(A)**. The data represent means ± SD (n = 3) and letters above the bars indicate significant differences ($P < 0.05$), as determined by one-way ANOVA with Turkey's HSD test.

(E) The ultrastructure of mesophyll cell chloroplasts of Col-0, *hy5, glk1 glk2*, and *hy5 glk1 glk2* seedlings grown under continuous white light for 7 days. Scale bar = 1 µm.

1158
1159
1160 **Figure 3.** Binding of HY5 to the Promoters and Transcriptional Activation of *GLK1,*
1161 *GLK2, GNC* and *CGA1.*
1162
(A) EMSA showing HY5 binds to the sub-fragments of promoters of *GLK1*, *GLK2*,
GNC and *CGA1 in vitro*. "-" and "+" indicate the absence and presence of

corresponding probes or proteins. FP means free probe.

(B) Schematic structures of effector and reporter constructs used in

dual-luciferase (LUC) reporter system. *REN*, renilla luciferase gene.

(C) Bar graphs showing HY5 induces the activation of *GLK1pro:LUC*,

GLK2pro:LUC, GNCpro:LUC and *CGA1pro:LUC*. The data represent means ± SD

(n = 4), and asterisks indicate a significant difference compared with control

(*P<0.05, **P<0.01, paired samples *t*-test).

(D) Illustration of *GLK1*, *GLK2*, *GNC* and *CGA1* promoter regions with the

indicated positions of primers used in ChIP-qPCR. Bar graphs showing

ChIP-qPCR assays that HY5 associates with the promoters in vivo. Col-0 material

and *ACTIN* gene were used as negative controls. The data represent means ± SD

(n = 3), and asterisks indicate a significant difference compared with Col-0

(*P<0.05, **P<0.01, paired samples *t*-test).

1178

1179

1180
1181
1182 **Figure 4.** GLK1/2 Negatively and GNC/CGA1 Positively Regulate Hypocotyl
1183 Elongation.
1184

(A) Phenotypes of 4-d-old Col-0, *glk1*, *glk2*, *glk1 glk2*, *hy5*, *hy5 glk1*, *hy5 glk2*, and
hy5 glk1 glk2 seedlings grown in white (100 µmol/m^2/s), blue (60 µmol/m^2/s), red
(90 µmol/m^2/s) light and dark conditions. Scale bar = 1 mm.
(B) Quantification of hypocotyl lengths indicated in (A). The data represent means
± SD (n ≥17) and letters above the bars indicate significant differences ($P < 0.05$),
as determined by one-way ANOVA with Turkey's HSD test.
(C) Quantification of hypocotyl cell length from 4-d-old Col-0, *glk1 glk2*, *hy5*, and
hy5 glk1 glk2 seedlings, grown in white light (100 µmol/m^2/s) conditions. The data
represent means ± SD (n = 15) and letters above the bars indicate significant
differences ($P < 0.05$), as determined by one-way ANOVA with Turkey's HSD test.
(D) Hypocotyl cell length phenotypes from 4-d-old Col-0, *glk1 glk2*, *hy5*, and *hy5*
glk1 glk2 seedlings. Scale bar = 20 µm.
(E) Phenotypes of 4-d-old Col-0, *gnc*, *cga1*, *gnc cga1*, *hy5*, *hy5 gnc*, *hy5 cga1*,
and *hy5 gnc cga1* seedlings grown in white light (100 µmol/m^2/s) and dark
conditions. Scale bar = 1 mm.
(F) Quantification of hypocotyl lengths indicated in (E). The data represent means
± SD (n ≥15) and letters above the bars indicate significant differences ($P < 0.05$),
as determined by one-way ANOVA with Turkey's HSD test.
1203
1204

1207 **Figure 5.** Different Expression Patterns of HY5, GLK2, and GNC Regulated
1208 Elongation Genes.

1209

(A) Venn diagram showing overlaps between sets of differentially expressed
genes in *glk2*, *hy5* and *hy5 glk2* mutants, relative to Col-0.
(B) K-means clustering of differentially expressed genes regulated by GLK2, HY5
and both. The scale bar shows fold changes (log2 value).
(C) Expression patterns of cell elongation genes in *glk2*, *hy5* and *hy5 glk2*
mutants, relative to Col-0. The scale bar shows fold changes (log2 value). Heat

map was aligned according to fold changes in *hy5 glk2* mutant.

(D, E) RT-qPCR analysis of cell elongation genes expressed in 4-d-old Col-0, *hy5,*

glk2, *hy5 glk2*, *gnc*, and *hy5 gnc* seedlings grown under continuous white light.

The *ACT7* gene was used as internal control. The data represent means ± SD (n

= 3) and letters above the bars indicate significant differences ($P < 0.05$), as

determined by one-way ANOVA with Turkey's HSD test.

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

A

B

Figure 6. GLK2 Is Responsible for the Developed Thylakoid Structures in the Etiolated *det1* Mutant.

(A) Phenotype of 4 d dark (upper panel) or light (lower panel) grown Col-0, *glk2*, *det1*, and *det1 glk2* seedlings. Scale bars represent 500 μm (upper panel) or 1 mm (lower panel).

(B) The ultrastructure of mesophyll chloroplasts of Col-0, *glk2*, *35S:GLK2 glk1 glk2*, *det1*, and *det1 glk2* seedlings grown under dark condition for 4 days. Scale bar = 1 μm. Black arrow heads in the bottom right image point to the thylakoid lamellae in very few amout.

1259

1260

1261 **Figure 7.** DET1 Promotes the Degradation of GLK Protein, and GLK Is
1262 Responsible for Photosystem Gene Expression in the Etiolated *det1* Mutant.

1263

(A) Immunoblot detection of GLK1 and GLK2 in Col-0, *glk1 glk2*, and *det1*
seedlings grown in the dark for 4 d (left). GLK protein levels in 4-d-old dark-grown
Col-0 and *det1* seedlings treated with 50 µM MG132 or DMSO (right). Coomassie

stained PAGE gel was shown as loading control, and the band of Rubisco large
subunit was indicated.
(**B**) RT-qPCR analysis of photosystem genes expressed in 4-d-old etiolated Col-0,
glk2, *35S:GLK2 glk1 glk2*, *det1*, and *det1 glk2* seedlings.
(**C, D**) RT-qPCR analysis of photosystem genes expressed in 4-d-old etiolated (D)
or light-grown (E) Col-0, *35S:GLK1 glk1 glk2*, *35S:GNC*, *35S:CGA1*, and *HA-HY5*
hy5 seedlings. The *ACT7* gene was used as internal control. The data represent
means ± SD (n = 3) and letters above the bars indicate significant differences ($P <$
0.05), as determined by one-way ANOVA with Turkey's HSD test.

1297

1298

Figure 8. A Working Model Depicting GLK, GNC/CGA1, and HY5 in the Regulation of Skotomorphogenesis and Photomorphogenesis.

GLK, GNC, CGA1 and HY5 undergo DET1-mediated degradation in the dark, so that the expression of photosynthesis associated nuclear genes (PhANGs), and the transformation of prolamellar body towards extended lamellae are inhibited. Light inhibits DET1 activity, and HY5 directs the promoter of *GLK*, *GNC* and *CGA1*, inducing their activities to promote chloroplast development. GLK inhibits hypocotyl elongation while GNC and CGA1 promote hypocotyl elongation. Thick green arrow indicates strong promotion of chloroplast development by GLK. Thick gray line indicates strong inhibition of hypocotyl elongation by HY5. Thinner arrows indicate less strong regulation. Dashed lines in the dark indicate GNC/CGA1 related regulations that pending to be further validated. Dashed double sided arrows in the light indicate potential protein interactions of HY5 with GLKs and GNC/CGA1.

1314

1315

Supplemental Figure 1. HY5, GLK, GNC and CGA1 Regulate the Chlorophyll Biosynthesis. (Supports Figure 1)

(A) Representative images of 4-day-old etiolated seedlings of Col-0, *glk1*, *glk2*, *glk1 glk2*, *35S:GLK1 glk1 glk2*, *35S:GLK2 glk1 glk2*, *hy5*, and *HA-HY5 hy5* during the transition from dark to light conditions for 6 h, 12 h, and 24 h. Scale bars = 1 mm.

(B) Representative images of 4-day-old etiolated seedlings of Col-0, *gnc*, *cga1*, *gnc cga1*, *35S:GNC*, *35S:CGA1*, *hy5*, and *HA-HY5 hy5* during the transition from dark to light conditions for 6 h, 12 h, and 24 h. Scale bars = 1 mm.

(C) Chlorophyll contents of 4-day-old etiolated seedlings from **(A)** during the transition from dark to light conditions for 6 h and 12 h. The data represent means ± SD (n=3) and letters above the bars indicate significant differences (*P* < 0.05), as determined by one-way ANOVA with Turkey's HSD test. The experiments were performed three times with similar results.

(D) Chlorophyll contents of 4-day-old etiolated seedlings from **(B)** during the transition from dark to light conditions for 6 h and 12 h. The data represent means ± SD (n=3) and letters above the bars indicate significant differences (*P* < 0.05), as determined by one-way ANOVA with Turkey's HSD test. The experiments were performed three times with similar results.

1342 **Supplemental Figure 2.** Phenotypes of the Mutants and Overexpression Lines
1343 under Continuous Light Conditions. (Supports Figure 2)

1344

(A) Representative images of 4-day-old Col-0, *glk1*, *glk2*, *glk1 glk2*, *35S:GLK1*
glk1 glk2, *35S:GLK2 glk1 glk2*, *hy5*, and *HA-HY5 hy5* seedlings grown on
half-strength MS plates for 4 days under continuous white light conditions. Scale
bars = 1 mm.
(B) Representative images of 4-day-old Col-0, *gnc*, *cga1*, *gnc cga1*, *35S:GNC*,
35S:CGA1, *hy5*, and *HA-HY5 hy5* seedlings grown on half-strength MS plates for
4 days under continuous white light conditions. Scale bars = 1 mm.
(C) Chlorophyll contents of the genotypes shown in **(A)**. The data represent
means ± SD (n=3) and letters above the bars indicate significant differences ($P <$
0.05), as determined by one-way ANOVA with Turkey's HSD test. The
experiments were performed three times with similar results.
(D) Chlorophyll contents of the genotypes shown in **(B)**. The data represent
means ± SD (n=3) and letters above the bars indicate significant differences ($P <$
0.05), as determined by one-way ANOVA with Turkey's HSD test. The
experiments were performed three times with similar results.

1360

1361

1362

1363 **Supplemental Figure 3.** GLK Inhibits Hypocotyl Elongation. (Supports Figure 4)

1364

(A) Hypocotyl phenotypes of 4-d-old Col-0, *glk1*, *glk2*, *glk1 glk2*, *hy5*, *35S:GLK1*
glk1 glk2, and *35S:GLK2 glk1 glk2* seedlings grown in white (100 μmol/m^2/s), blue
(60 μmol/m^2/s), red (90 μmol/m^2/s) light and dark conditions. Scale bar = 1 mm.
(B) Quantification of hypocotyl lengths indicated in **(A)**. The data represent means
± SD (n ≥18) and letters above the bars indicate significant differences (*P* < 0.05),
as determined by one-way ANOVA with Turkey's HSD test. The experiments were
performed three times with similar results.

1372

1373

1374

1375

1376 **Supplemental Figure 4.** GNC and CGA1 Promote Hypocotyl Elongation.
1377 (Supports Figure 4)

1378

(A) Hypocotyl phenotypes of 4-d-old Col-0, *gnc*, *cga1*, *gnc cga1*, *35S:GNC*, and
35S:CGA1 seedlings grown in white light (100 µmol/m^2/s) and dark conditions.
Scale bar = 1 mm.
(B) Quantification of hypocotyl lengths indicated in **(A)**. The data represent means
± SD (n ≥15) and letters above the bars indicate significant differences (P < 0.05),
as determined by one-way ANOVA with Turkey's HSD test. The experiments were
performed three times with similar results.

1386

1387

Supplemental Figure 5. HY5 but Not GLKs Is Crucial for the Greening of Detached Roots.

Root samples were detached from 7-d-old seedlings, and then cultured on hormone-free MS medium for 14 d. Scale bar = 5 mm.

1405
1406
1407 **Supplemental Figure 6.** 6-BA Promotes GLK and HY5 Dependent Chlorophyll
1408 Biosynthesis.
1409
Chlorophyll content of 4-day-old etiolated seedlings of Col-0, *glk1*, *glk2*, *glk1 glk2*,
hy5, *hy5 glk1*, *hy5 glk2*, and *hy5 glk1 glk2* grown on medium with or without 6-BA
during transition from dark to light conditions for 6 h **(A)** and 12 h **(B)**. The data
represent means ± SD (n=3) and letters above the bars indicate significant
differences ($P < 0.05$), as determined by one-way ANOVA with Turkey's HSD test.
The experiments were performed three times with similar results.

1426

1427 **Supplemental Figure 7.** *In vitro* Evidence of HY5 Interacting with GLK and
1428 Potential Co-regulation of Target Genes.

1429

1430 **(A)** *In vitro* pulldown assays showing the interaction of GLK1 and GLK2 with HY5.
1431 GST-GLK protein or GST protein were used to pull down His-HY5 protein using
1432 GST beads. Anti-GST and anti-His antibodies were used for immunoblot analysis.

"-" and "+" indicate the absence and presence of corresponding proteins.

(B) Bimolecular luminescence complementation assays showing GLK1 and GLK2 can interact with HY5. nLUC and cLUC served as negative controls.

(C, D) Illustration of *LHCB1.2* and *PME16* promoter regions used in dual-luciferase assay. Bar graphs showing HY5 and GLK2 induce the activation or inhibition of *LHCB1.2pro:LUC* and *PME16pro:LUC*. The data represent means ± SD (n = 4), and asterisks indicate a significant difference compared with control (*P<0.05, **P<0.01, paired samples *t*-test).

1464
1465
1466 **Supplemental Figure 8.** *In vitro* Evidence of DET1 Interacting with GLK and
1467 GNC/CGA1.

1468
1469 *In vitro* pulldown assays showing the interactions of GLK1, GLK2 (A), GNC and
1470 CGA1 (B) with DET1. GST-GLK1, GST-GLK2, GST-GNC, GST-CGA1 or GST
1471 proteins were used to pull down His-nDET1 protein using GST beads. Anti-GST
1472 and anti-His antibodies were used for immunoblot analysis. "-" and "+" indicate the
1473 absence and presence of corresponding proteins.
1474
1475

Parsed Citations

Abbas, N., Maurya, J.P., Senapati, D., Gangappa, S.N., and Chattopadhyay, S. (2014). Arabidopsis CAM7 and HY5 physically interact and directly bind to the HY5 promoter to regulate its expression and thereby promote photomorphogenesis. Plant Cell 26: 1036-1052.
Google Scholar: Author Only Title Only Author and Title

Alem, A.L., Ariel, F.D., Cho, Y., Hong, J.C., Gonzalez, D.H., and Viola, I.L. (2022). TCP15 interacts with GOLDEN2-LIKE 1 to control cotyledon opening in Arabidopsis. Plant J. 110(3): 748-763.
Google Scholar: Author Only Title Only Author and Title

Anders, S., and Hubre, W. (2013). Differential expression of RNA-Seq data at the gene level-the DESeq package. European Molecular Biology Laboratory (EMBL).
Google Scholar: Author Only Title Only Author and Title

Arnon, D.I. (1949). Copper enzymes in isolated chloroplasts polypheoloxidase in Bata Vulgaris. Plant Physiol. 24: 1-15.
Google Scholar: Author Only Title Only Author and Title

Bastakis, E., Hedtke, B., Klermund, C., Grimm, B., and Schwechheimer, C. (2018). LLM-Domain B-GATA transcription factors play multifaceted roles in controlling greening in Arabidopsis. Plant Cell 30: 582-599.
Google Scholar: Author Only Title Only Author and Title

Bi, Y.M., Zhang, Y., Signorelli, T., Zhao, R., Zhu, T., and Rothstein, S. (2005). Genetic analysis of Arabidopsis GATA transcription factor gene family reveals a nitrate-inducible member important for chlorophyll synthesis and glucose sensitivity. Plant J. 44: 680-692.
Google Scholar: Author Only Title Only Author and Title

Binkert, M., Kozma-Bognár, L., Terecskei, K., De Veylder, L., Nagy, F., and Ulm, R. (2014). UV-B-responsive association of the Arabidopsis bZIP transcription factor ELONGATED HYPOCOTYL5 with target genes, including its own promoter. Plant Cell 26: 4200-4213.
Google Scholar: Author Only Title Only Author and Title

Catalá, R., Medina, J., and Salinas, J. (2011). Integration of low temperature and light signaling during cold acclimation response in Arabidopsis. Proc. Natl. Acad. Sci. U. S. A. 108: 16475-16480.
Google Scholar: Author Only Title Only Author and Title

Chang, C.S., Li, Y.H., Chen, L.T., Chen, W.C., Hsieh, W.P., Shin, J., Jane, W.N., Chou, S.J., Choi, G., Hu, J.M., Somerville, S., and Wu, S.H. (2008). LZF1, a HY5-regulated transcriptional factor, functions in Arabidopsis de-etiolation. Plant J. 54: 205-219.
Google Scholar: Author Only Title Only Author and Title

Chen, D., Xu, G., Tang, W., Jing, Y., Ji, Q., Fei, Z., and Lin, R. (2013). Antagonistic basic helix-loop-helix/bZIP transcription factors form transcriptional modules that integrate light and reactive oxygen species signaling in Arabidopsis. Plant Cell 25: 1657-1673.
Google Scholar: Author Only Title Only Author and Title

Chiang, Y.H., Zubo, Y.O., Tapken, W., Kim, H.J., Lavanway, A.M., Howard, L., Pilon, M., Kieber, J.J., and Schaller, G.E. (2012). Functional characterization of the GATA transcription factors GNC and CGA1 reveals their key role in chloroplast development, growth, and division in Arabidopsis. Plant Physiol. 160: 332-348.
Google Scholar: Author Only Title Only Author and Title

Chory, J., Peto, C., Feinbaum, R., Pratt, L., and Ausubel, F. (1989). Arabidopsis thaliana mutant that develops as a light-grown plant in the absence of light. Cell 58: 991-999.
Google Scholar: Author Only Title Only Author and Title

Ciolfi, A., Sessa, G., Sassi, M., Possenti, M., Salvucci, S., Carabelli, M., Morelli, G., and Ruberti, I. (2013). Dynamics of the shade-avoidance response in Arabidopsis. Plant Physiol. 163: 331-353.
Google Scholar: Author Only Title Only Author and Title

Datta, S., Hettiarachchi, C., Johansson, H., and Holm, M. (2007). SALT TOLERANCE HOMOLOG2, a B-box protein in Arabidopsis that activates transcription and positively regulates light-mediated development. Plant Cell 19: 3242-3255.
Google Scholar: Author Only Title Only Author and Title

Datta, S., Johansson, H., Hettiarachchi, C., Irigoyen, M.L., Desai, M., Rubio, V., and Holm, M. (2008). LZF1/SALT TOLERANCE HOMOLOG3, an Arabidopsis B-box protein involved in light-dependent development and gene expression, undergoes COP1-mediated ubiquitination. Plant Cell 20: 2324-2338.
Google Scholar: Author Only Title Only Author and Title

Deng, X.W., Matsui, M., Wei, N., Wagner, D., Chu, A.M., Feldmann, K.A., and Quail, P.H. (1992). COP1, an Arabidopsis regulatory gene, encodes a protein with both a zinc-binding motif and a G beta homologous domain. Cell 71: 791-801.
Google Scholar: Author Only Title Only Author and Title

Fitter, D.W., Martin, D.J., Copley, M.J., Scotland, R.W., and Langdale, J.A. (2002). GLK gene pairs regulate chloroplast development in diverse plant species. Plant J. 31: 713-727.
Google Scholar: Author Only Title Only Author and Title

Hall, L.N., Rossini, L., Cribb, L., and Langdale, J.A. (1998). GOLDEN 2: a novel transcriptional regulator of cellular differentiation in the maize leaf. Plant Cell 10: 925-936.
Google Scholar: Author Only Title Only Author and Title

Holm, M., Ma, L.G., Qu, L.J., and Deng, X.W. (2002). Two interacting bZIP proteins are direct targets of COP1-mediated control of light-dependent gene expression in Arabidopsis. Genes Dev. 16: 1247-1259.
Google Scholar: Author Only Title Only Author and Title

Huang, X., Ouyang, X., Yang, P., Lau, O.S., Li, G., Li, J., Chen, H., and Deng, X.W. (2012). Arabidopsis FHY3 and HY5 positively mediate induction of COP1 transcription in response to photomorphogenic UV-B light. Plant Cell 24: 4590-4606.
Google Scholar: Author Only Title Only Author and Title

Hudson, D., Guevara, D.R., Hand, A.J., Xu, Z, Hao, L., Chen, X., Zhu, T., Bi, Y.M., and Rothstein, S.J. (2013). Rice cytokinin GATA transcription Factor1 regulates chloroplast development and plant architecture. Plant Physiol. 162: 132-144.
Google Scholar: Author Only Title Only Author and Title

Imamura, A, Hanaki, N., Nakamura, A, Suzuki, T., Taniguchi, M., Kiba, T., Ueguchi, C., Sugiyama, T., and Mizuno, T. (1999). Compilation and characterization of Arabidopsis thaliana response regulators implicated in His-Asp phosphorelay signal transduction. Plant Cell Physiol. 40: 733-742.
Google Scholar: Author Only Title Only Author and Title

Jakoby, M., Weisshaar, B., Dröge-Laser, W., Vicente-Carbajosa, J., Tiedemann, J., Kroj, T., Parcy, F., and bZIP Research Group. (2002). bZIP transcription factors in Arabidopsis. Trends Plant Sci. 7: 106-111.
Google Scholar: Author Only Title Only Author and Title

Jang, I.C., Henriques, R., and Chua, N.H. (2013). Three transcription factors, HFR1, LAF1 and HY5, regulate largely independent signaling pathways downstream of phytochrome A. Plant Cell Physiol. 54: 907-916.
Google Scholar: Author Only Title Only Author and Title

Jiao, Y., Lau, O.S., and Deng, X.W. (2007). Light-regulated transcriptional networks in higher plants. Nat. Rev. Genet. 8: 217-230.
Google Scholar: Author Only Title Only Author and Title

Jing, Y., Zhang, D., Wang, X., Tang, W., Wang, W., Huai, J., Xu, G., Chen, D., Li, Y., and Lin, R. (2013). Arabidopsis chromatin remodeling factor PICKLE interacts with transcription factor HY5 to regulate hypocotyl cell elongation. Plant Cell 25: 242-256.
Google Scholar: Author Only Title Only Author and Title

Kobayashi, K., Baba, S., Obayashi, T., Sato, M., Toyooka, K., Keränen, M., Aro, E.M., Fukaki, H., Ohta, H., Sugimoto, K., and Masuda, T. (2012a). Regulation of root greening by light and auxin/cytokinin signaling in Arabidopsis. Plant Cell 24: 1081-1095.
Google Scholar: Author Only Title Only Author and Title

Kobayashi, K., Fujii, S., Sasaki, D., Baba, S., Ohta, H., Masuda, T., and Wada, H. (2014). Transcriptional regulation of thylakoid galactolipid biosynthesis coordinated with chlorophyll biosynthesis during the development of chloroplasts in Arabidopsis. Front Plant Sci. 5: 272.
Google Scholar: Author Only Title Only Author and Title

Kobayashi, K., Obayashi, T., and Masuda, T. (2012b). Role of the G-box element in regulation of chlorophyll biosynthesis in Arabidopsis roots. Plant Signal.Behav. 7: 922-926.
Google Scholar: Author Only Title Only Author and Title

Kobayashi, K., Ohnishi, A, Sasaki, D., Fujii, S., Iwase, A, Sugimoto, K., Masuda, T., and Wada, H. (2017). Shoot Removal Induces Chloroplast Development in Roots via Cytokinin Signaling. Plant Physiol. 173: 2340-2355.
Google Scholar: Author Only Title Only Author and Title

Kushwaha, R., Singh, A, and Chattopadhyay, S. (2008). Calmodulin7 plays an important role as transcriptional regulator in Arabidopsis seedling development. Plant Cell 20: 1747-1759.
Google Scholar: Author Only Title Only Author and Title

Lee, D., Hua, L., Khoshravesh, R., Giuliani, R., Kumar, I., Cousins, A, Sage, T. L., Hibberd, J. M., and Brutnell, T. P. (2021). Engineering chloroplast development in rice through cell-specific control of endogenous genetic circuits. Plant Biotechnol. J. 19(11): 2291-2303.
Google Scholar: Author Only Title Only Author and Title

Lee, J., Choi, B., Yun, A, Son, N., Ahn, G., Cha, J.Y., Kim, W.Y., and Hwang, I. (2021). Long-term abscisic acid promotes golden2-like1 degradation through constitutive photomorphogenic 1 in a light intensity-dependent manner to suppress chloroplast development. Plant Cell Environ. 44: 3034-3048.
Google Scholar: Author Only Title Only Author and Title

Lee, J., He, K., Stolc, V., Lee, H., Figueroa, P., Gao, Y., Tongprasit, W., Zhao, H., Lee, I., and Deng, X.W. (2007). Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. Plant Cell 19: 731-749.
Google Scholar: Author Only Title Only Author and Title

Liu, D., Zhao, D., Li, X., and Zeng, Y.J. (2022) AtGLK2, an Arabidopsis GOLDEN2-LIKE transcription factor, positively regulates anthocyanin biosynthesis via AtHY5-mediated light signaling. Plant Growth Regul. 96: 79-90.
Google Scholar: Author Only Title Only Author and Title

Martín, G., Leivar, P., Ludevid, D., Tepperman, J.M., Quail, P.H., and Monte, E. (2016). Phytochrome and retrograde signalling pathways converge to antagonistically regulate a light-induced transcriptional network. Nat. Commun. 7: 11431.
Google Scholar: Author Only Title Only Author and Title

Naito, T., Kiba, T., Koizumi, N., Yamashino, T., and Mizuno, T. (2007). Characterization of a unique GATA family gene that responds to both light and cytokinin in Arabidopsis thaliana. Biosci. Biotechnol. Biochem. 71: 1557-1560.
Google Scholar: Author Only Title Only Author and Title

Nakamura, H., Muramatsu, M., Hakata, M., Ueno, O., Nagamura, Y., Hirochika, H., Takano, M., and Ichikawa, H. (2009). Ectopic overexpression of the transcription factor OsGLK1 induces chloroplast development in non-green rice cells. Plant Cell Physiol. 50: 1933-1949.
Google Scholar: Author Only Title Only Author and Title

Nguyen, C.V., Vrebalov, J.T., Gapper, N.E., Zheng, Y., Zhong, S., Fei, Z., and Giovannoni, J.J. (2014). Tomato GOLDEN2-LIKE transcription factors reveal molecular gradients that function during fruit development and ripening. Plant Cell 26: 585-601.
Google Scholar: Author Only Title Only Author and Title

Osterlund, M.T., Hardtke, C.S., Wei, N., and Deng, X.W. (2000). Targeted destabilization of HY5 during light-regulated development of Arabidopsis. Nature 405: 462-466.
Google Scholar: Author Only Title Only Author and Title

Oyama, T., Shimura, Y., and Okada, K. (1997). The Arabidopsis HY5 gene encodes a bZIP protein that regulates stimulus-induced development of root and hypocotyl. Genes Dev. 11: 2983-2995.
Google Scholar: Author Only Title Only Author and Title

Ram, H., Priya, P., Jain, M., and Chattopadhyay, S. (2014). Genome-wide DNA binding of GBF1 is modulated by its heterodimerizing protein partners, HY5 and HYH. Mol. Plant 7: 448-451.
Google Scholar: Author Only Title Only Author and Title

Ranftl, Q.L., Bastakis, E., Klermund, C., and Schwechheimer, C. (2016). LLM-Domain Containing B-GATA Factors Control Different Aspects of Cytokinin-Regulated Development in Arabidopsis thaliana. Plant Physiol. 170(4): 2295-2311.
Google Scholar: Author Only Title Only Author and Title

Rauf, M., Arif, M., Dortay, H., Matallana-Ramírez, L.P., Waters, M.T., Gil Nam, H., Lim, P.O., Mueller-Roeber, B., and Balazadeh, S. (2013). ORE1 balances leaf senescence against maintenance by antagonizing G2-like-mediated transcription. Embo Rep. 14: 382-388.
Google Scholar: Author Only Title Only Author and Title

Rossini, L., Cribb, L., Martin, D.J., and Langdale, J.A. (2001). The maize golden2 gene defines a novel class of transcriptional regulators in plants. Plant Cell 13: 1231-1244.
Google Scholar: Author Only Title Only Author and Title

Saijo, Y., Sullivan, J.A., Wang, H., Yang, J., Shen, Y., Rubio, V., Ma, L., Hoecker, U., and Deng, X.W. (2003). The COP1-SPA1 interaction defines a critical step in phytochrome A-mediated regulation of HY5 activity. Genes Dev. 17: 2642-2647.
Google Scholar: Author Only Title Only Author and Title

Seo, H.S., Yang, J.Y., Ishikawa, M., Bolle, C., Ballesteros, M.L., and Chua, N.H. (2003). LAF1 ubiquitination by COP1 controls photomorphogenesis and is stimulated by SPA1. Nature 423: 995-999.
Google Scholar: Author Only Title Only Author and Title

Shin, J., Park, E., and Choi, G. (2007). PIF3 regulates anthocyanin biosynthesis in an HY5-dependent manner with both factors directly binding anthocyanin biosynthetic gene promoters in Arabidopsis. Plant J. 49: 981-994.
Google Scholar: Author Only Title Only Author and Title

Singh, A., Ram, H., Abbas, N., and Chattopadhyay, S. (2012). Molecular interactions of GBF1 with HY5 and HYH proteins during light-mediated seedling development in Arabidopsis thaliana. J. Biol. Chem. 287: 25995-26009.
Google Scholar: Author Only Title Only Author and Title

Song, Y., Yang, C., Gao, S., Zhang, W., Li, L., and Kuai, B. (2014). Age-triggered and dark-induced leaf senescence require the bHLH transcription factors PIF3, 4, and 5. Mol. Plant 7: 1776-1787.
Google Scholar: Author Only Title Only Author and Title

Song, Y.H., Yoo, C.M., Hong, A.P., Kim, S.H., Jeong, H.J., Shin, S.Y., Kim, H.J., Yun, D.J., Lim, C.O., Bahk, J.D., Lee, S.Y., Nagao, R.T., Key, J.L., and Hong, J.C. (2008). DNA-binding study identifies C-box and hybrid C/G-box or C/A-box motifs as high-affinity binding sites for STF1 and LONG HYPOCOTYL5 proteins. Plant Physiol. 146: 1862-1877.
Google Scholar: Author Only Title Only Author and Title

Stracke, R., Favory, J.J., Gruber, H., Bartelniewoehner, L., Bartels, S., Binkert, M., Funk, M., Weisshaar, B., and Ulm, R. (2010). The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene in response to light and ultraviolet-B radiation. Plant Cell Environ. 33: 88-103.
Google Scholar: Author Only Title Only Author and Title

Sullivan, J.A., and Deng, X.W. (2003). From seed to seed: the role of photoreceptors in Arabidopsis development. Dev. Biol. 260: 289-297.
Google Scholar: Author Only Title Only Author and Title

Tamai, H., Iwabuchi, M., and Meshi, T. (2002). Arabidopsis GARP transcriptional activators interact with the Pro-rich activation domain shared by G-box-binding bZIP factors. Plant Cell Physiol. 43: 99-107.
Google Scholar: Author Only Title Only Author and Title

Tang, X., Miao, M., Niu, X., Zhang, D., Cao, X., Jin, X., Zhu, Y., Fan, Y., Wang, H., Liu, Y., Sui, Y., Wang, W., Wang, A., Xiao, F., Giovannoni, J., and Liu, Y. (2016). Ubiquitin-conjugated degradation of golden 2-like transcription factor is mediated by CUL4-DDB1-based E3 ligase complex in tomato. New Phytol. 209: 1028-1039.
Google Scholar: Author Only Title Only Author and Title

Tepperman, J.M., Hwang, Y.-S., and Quail, P.H. (2006). phyA dominates in transduction of red-light signals to rapidly responding genes at the initiation of Arabidopsis seedling de-etiolation. Plant J. 48: 728-742.
Google Scholar: Author Only Title Only Author and Title

Tokumaru, M., Adachi, F., Toda, M., Ito-Inaba, Y., Yazu, F., Hirosawa, Y., Sakakibara, Y., Suiko, M., Kakizaki, T., and Inaba, T. (2017). Ubiquitin-Proteasome Dependent Regulation of the GOLDEN2-LIKE 1 Transcription Factor in Response to Plastid Signals. Plant Physiol. 173: 524-535.
Google Scholar: Author Only Title Only Author and Title

Toledo-Ortiz, G., Johansson, H., Lee, K.P., Bou-Torrent, J., Stewart, K., Steel, G., Rodríguez-Concepción, M., and Halliday, K. J. (2014). The HY5-PIF regulatory module coordinates light and temperature control of photosynthetic gene transcription. PLoS Genet. 10: e1004416.
Google Scholar: Author Only Title Only Author and Title

Wang, P., Khoshravesh, R., Karki, S., Tapia, R., Balahadia, C. P., Bandyopadhyay, A, Quick, W. P., Furbank, R., Sage, T. L., and Langdale, J. A. (2017). Re-creation of a Key Step in the Evolutionary Switch from C (3) to C (4) Leaf Anatomy. Curr. Biol. 27: 3278-3287.e3276.
Google Scholar: Author Only Title Only Author and Title

Waters, M.T., Moylan, E.C., and Langdale, J.A. (2008). GLK transcription factors regulate chloroplast development in a cell-autonomous manner. Plant J. 56: 432-444.
Google Scholar: Author Only Title Only Author and Title

Waters, M.T., Wang, P., Korkaric, M., Capper, R.G., Saunders, N.J., and Langdale, J.A. (2009). GLK transcription factors coordinate expression of the photosynthetic apparatus in Arabidopsis. Plant Cell 21: 1109-1128.
Google Scholar: Author Only Title Only Author and Title

Wykoff, D.D., Grossman, A.R., Weeks, D.P., Usuda, H., and Shimogawara, K. (1999). Psr1, a nuclear localized protein that regulates phosphorus metabolism in Chlamydomonas. Proc. Natl. Acad. Sci. U. S. A. 96: 15336-15341.
Google Scholar: Author Only Title Only Author and Title

Xu, D., Jiang, Y., Li, J., Lin, F., Holm, M., and Deng, X.W. (2016a). BBX21, an Arabidopsis B-box protein, directly activates HY5 and is targeted by COP1 for 26S proteasome-mediated degradation. Proc. Natl. Acad. Sci. U. S. A. 113: 7655-7660.
Google Scholar: Author Only Title Only Author and Title

Xu, X., Chi, W., Sun, X., Feng, P., Guo, H., Li, J., Lin, R., Lu, C., Wang, H., Leister, D., and Zhang, L. (2016b). Convergence of light and chloroplast signals for de-etiolation through ABI4-HY5 and COP1. Nat. Plants 2: 16066.
Google Scholar: Author Only Title Only Author and Title

Xu, Z, Casaretto, J.A, Bi, Y.M., and Rothstein, S.J. (2017). Genome-wide binding analysis of AtGNC and AtCGA1 demonstrates their cross-regulation and common and specific functions. Plant Direct 1: e00016.
Google Scholar: Author Only Title Only Author and Title

Yang, J., Lin, R., Sullivan, J., Hoecker, U., Liu, B., Xu, L., Deng, X. W., and Wang, H. (2005). Light regulates COP1-mediated degradation of HFR1, a transcription factor essential for light signaling in Arabidopsis. Plant Cell 17: 804-821.
Google Scholar: Author Only Title Only Author and Title

Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. Nat. Protoc. 2(7): 1565-1572.
Google Scholar: Author Only Title Only Author and Title

Zhang, D., Tan, W., Yang, F., Han, Q., Deng, X., Guo, H., Liu, B., Yin, Y., and Lin, H. (2021). A BIN2-GLK1 Signaling Module Integrates Brassinosteroid and Light Signaling to Repress Chloroplast Development in the Dark. Dev. Cell 56: 310-324 e317.
Google Scholar: Author Only Title Only Author and Title

Zhang, H., He, H., Wang, X., Wang, X., Yang, X., Li, L., and Deng, X.W. (2011). Genome-wide mapping of the HY5-mediated gene networks in Arabidopsis that involve both transcriptional and post-transcriptional regulation. Plant J. 65: 346-358.
Google Scholar: Author Only Title Only Author and Title

Zhao, Y.R., Li, X., Yu, K.Q., Cheng, F., and He, Y. (2016). Hyperspectral Imaging for Determining Pigment Contents in Cucumber Leaves in Response to Angular Leaf Spot Disease. Sci. Rep. 6: 27790.
Google Scholar: Author Only Title Only Author and Title

Zubo, Y.O., Blakley, I.C., Franco-Zorrilla, J.M., Yamburenko, M.V., Solano, R., Kieber, J.J., Loraine, A.E., and Schaller, G.E. (2018). Coordination of Chloroplast Development through the Action of the GNC and GLK Transcription Factor Families. Plant Physiol. 178: 130-147.
Google Scholar: Author Only Title Only Author and Title