Actin cytoskeleton dynamics affect replication of Human Metapneumovirus

Pamela Elizabeth Rodríguez[a][§], Pedro Ignacio Gil[b][§], Jorge Augusto Cámara[a], Alicia Cámara[a]

and María Gabriela Paglini*[b,c]

[a]Laboratory of Influenza and other Respiratory Viruses, Virology Institute "Dr. J. M. Vanella".
Facultad de Ciencias Médicas. Universidad Nacional de Córdoba. Córdoba. Córdoba.
Argentina.
[b]Laboratory of Cell Biology of Viral Infection, Virology Institute "Dr. J. M. Vanella". Facultad
de Ciencias Médicas. Universidad Nacional de Córdoba. Córdoba. Córdoba. Argentina.
[c]Laboratory of Neurophysiology, Instituto de Investigación Médica Mercedes y Martín
Ferreyra, INIMEC-CONICET, Universidad Nacional de Córdoba, Córdoba, Argentina.

§: Pamela Elizabeth Rodriguez and Pedro Ignacio Gil contributed equally to this work.

***Corresponding author.**
María Gabriela Paglini, Ph.D. ORCID: 0000-0002-2565-8699
Laboratory of Cell Biology of Viral Infection, Virology Institute "Dr. J. M. Vanella". Facultad
de Ciencias Médicas. Universidad Nacional de Córdoba. Córdoba. Argentina,
Enfermera Gordillo s/n. Ciudad Universitaria, 5000 - Córdoba Argentina.
Phone: +54-351-4334022
E-mail: gpaglini@immf.uncor.edu

ORCID:

Pamela Elizabeth Rodríguez (rpamelaelizabeth@gmail.com): 0000-0003-3827-3760

Pedro Ignacio Gil (pedro.gil@unc.edu.ar): 0000-0002-5444-063X

Jorge Augusto Cámara (jcamara@fcm.unc.edu.ar): 0000-0002-5072-2795

Alicia Cámara (camaraalicia@yahoo.com.ar): 0000-0003-3988-2664

Word count manuscript: 3526

Word count abstract: 200

Author contribution

All authors were responsible for the study design. PER, PIG performed the experimental

procedures and collected/curated data. PER and PIG conducted statistical data analysis. PER,

PIG, AC and MGP performed imaging analysis. JC, AC and MGP contributed ideas and

comments. PER, PIG, JC, AC and MGP interpreted the findings and drafted the manuscript.

Supervision and project administration were done by AC and MGP and funding's acquisition

by MGP and AC. All authors have read and agreed to the published version of the manuscript.

Short title

Human Metapneumovirus replication dynamics

ABSTRACT

Human Metapneumovirus (hMPV) is responsible for viral respiratory infection with clinical

and epidemiological relevance in pediatric, immunocompromised, and elderly populations.

Little is known about hMPV *in vitro* replication processes and their relationship with cellular

structures such as the cytoskeleton. Our goal was to evaluate the role of the actin cytoskeleton

in hMPV replication at different stages of viral growth. hMPV was isolated in Vero cells from

a clinical sample and identified as A_2 genotype. The cytopathic effect was detected by the

appearance of cell rounding and refractory cell clusters. The growth curve showed that viral

replication maximum level was between 48 and 72 hpi. The highest percentage of infected cells

and intracellular hMPV-protein were detected at the early stages of the replication cycle.

Disruption of actin microfilaments with Cytochalasin D during the early events provoked an

increase in both intracellular and extracellular viruses. We demonstrate that the early phase of

the hMPV curve is crucial for viral replication; and the disruption of microfilaments during this

time increments both viral protein expression and release of viruses to the extracellular space.

This study contributes to elucidate wild-type hMPV growth kinetics, providing new insights on

the actin cytoskeleton role in viral replication mechanisms.

Keywords.

Virus isolation; Replication cycle; Actin filaments; Cytochalasin D; Respiratory disease.

BACKGROUND

Two members of the *Pneumoviridae* family [1] human Metapneumovirus (hMPV) and respiratory syncytial virus (RSV), cause severe respiratory diseases in infants, immunocompromised and older adults [2]. Currently, for RSV, a bivalent prefusion F vaccine was approved for use during pregnancy in 2023 to protect infants from severe RSV illness during their early months of life [3]. However, there is no licensed vaccine or approved antiviral therapy available for hMPV.

HMPV is an enveloped negative-stranded RNA virus with a non-segmented genome composed of eight genes encoding for nine proteins [4]. Two major hMPV subtypes have been described: A (A1, A2a, A2b1, A2b2) and B (B1, B2) [2,5].

Since the discovery of hMPV [5], several approaches for its *in vitro* isolation have been followed. Viral adaptation and spread in cell cultures are difficult since both require high viral loads and successive blind passages of about 14-21 days to visualize the cytopathic effect (CPE), characterized by changes in cell morphology or by the formation of syncytia [6,7].

The viral infection cycle begins with the entry of particles into susceptible cells through the interaction with receptors that trigger the internalization process. Several studies have reported an important role for the cytoskeleton, particularly actin microfilaments (MFLs) dynamics, in entry and replication of numerous viruses, including hMPV, into the host cell [8]. In this sense, it has been shown that MFLs are essential cofactors for RSV replication, spread, and morphogenesis [9]. Moreover, previous studies from our laboratory demonstrated that the disruption of MFLs during early stage of Pixuna virus (PIXV) replication, increased the extracellular viral yields, probably promoting endocytosis and thus increasing the entry of viral particles [10].

Despite hMPV's importance as an etiological agent of respiratory pathologies, the mechanisms

of interaction between virus and host cell to secure infection remain largely unexplored. We

examined the role of MFLs in hMPV replication at different stages of viral growth. To this end,

we isolated and identified hMPV circulating in Cordoba, Argentina. Furthermore, we describe

the hMPV replication curve and characterize the subcellular expression of viral proteins at

different times after infection. We demonstrate that actin depolymerization at the early phase

of the hMPV replication curve is crucial for a successful replication. The presence of

Cytochalasin D (CytD) during this period increased viral protein expression in the cytoplasm

and the release of viruses to the extracellular space.

Considering that hMPV causes a severe disease, identifying the molecular mechanisms

underlying its replication cycle broadens our understanding of the cell biology of viral infection.

Moreover, it provides possible targets to develop new antiviral treatments.

METHODS

Cell culture and Virus isolation

Vero cells (ATCC® CCL-81) were grown in Minimum Essential Medium (MEM, GIBCO-

BRL®) with 5-10% fetal bovine serum (FBS, Natocor), 1% antibiotic-antimycotic (Pen-Strept

100X, GIBCO), at 37°C and 5% CO_2.

Human Metapneumovirus was isolated from nasopharyngeal aspirates (NPA) of hospitalized

children in Cordoba, Argentina. All procedures were complied with the principles outlined by

the Declaration of Helsinki and were approved by an Independent Ethics Committee of Hospital

de Niños "Santísima Trinidad" (CIEIS) Protocol: 05/2011. The volunteers who offered samples,

signed written assent/consent and their personal data was kept anonymous. The infection

protocol was adapted from Van den Hoogen [5]. As infection medium: 0.00125% trypsin

(Trypsin Solution 10X, SIGMA®), 0.3% bovine serum albumin (BSA, SIGMA®) and 1%

antibiotic-antimycotic. The NPA positive was diluted in 2 ml of MEM without FBS and

centrifuged at 200Xg for 5 min (HSR Centrifuge Eppendorf Presvac) (NPA supernatant). Viral

inoculum consisted of 150 μl of NPA supernatant plus 50 μl of infection medium. Vero cells

(70-80% confluence) were grown in a 24-well plate with MEM. 24 hours later, the plate was

washed three times with PBS, 200 μl of the viral inoculum was added in each well and

centrifuged at 800Xg for 15 min (IEC International Refrigerated Centrifuge Model-PR-2).

Afterwards, the plate was incubated at 37°C and 5% CO_2 for 2 hours, the monolayers were

washed three times with PBS and 1 ml of infection medium per well was added. The infection

medium was changed at 4, 7, 10, 14, 17, and 21 days post-infection (dpi) and the culture

supernatants were stored until processed at -70°C. Cultures were observed daily for CPE by

phase-contrast microscopy. Three blind passages for 21 days each were performed. The hMPV

positive culture supernatants were used as an infection inoculum.

RT-PCR

Viral RNA was extracted from 140 μl of culture supernatants using the QIAamp® Viral RNA

Mini Kit (Qiagen, GmbH, Hilden, Germany) following the manufacturer's instructions. RT-

PCR, adapted from Bouscambert-Duchamp [11], was carried out as described by Rodriguez

[12].

Indirect immunofluorescence

Infected cells were grown at 60% confluence on glass coverslips during 24 hours. At different

times post-infection, cells were washed three times with PBS, fixed with 4% paraformaldehyde

and 120 mM sucrose (Sigma-Aldrich Laborchemikalien GmbH, Germany) for 20 min at room

temperature (RT). Immunofluorescence (IF) was performed according to Gil [10]. Cells were

incubated overnight with mouse anti-F protein/hMPV primary antibody (1/500) (IMAGEN™ hMPV. Oxoid-Ltd.) in 1% BSA/PBS at 4°C. After washed, samples were incubated with goat anti-mouse secondary antibody Alexa-Fluor 488 (1/1600) (Thermo Fisher Scientific Inc., USA) for 1 hour at RT. To visualize the actin filaments and nuclei, cells were labeled with Phalloidin-Tetramethyl rhodamine-B (1/1000) (Sigma-Aldrich) for 1 hour at RT and Hoechst (1X) for 5 min. Then, samples were washed and mounted using Fluorsave (Calbiochem).

Sequence and phylogenetic analysis

The RT-PCR protocol, adaptaded from Van den Hoogen [13] to amplify a 696 bp fragment of the F gene, was carried out as described by Rodriguez [12]. PCR product was purified with QIAquick Gel Extraction Kit (Qiagen, Germany) following the manufacturer's instructions. Nucleotide sequencing reactions were performed in both directions using the internal PCR primers by Macrogen, Inc. (Seoul, Korea). Sequences were edited with MEGA-4.0.2 and aligned with sequences available in the GenBank, using the ClustalW. A Maximum Likelihood Tree, based on the amplified region of the fusion protein (F) gene, was generated using PhyML 3.0 software (Université de Montpellier, France). Branch support was evaluated via non-parametric bootstrapping with 1000 pseudoreplicates. The nucleotide substitution model was choosen based on the AIC implemented in ModelTest-3.7 software (University of Vigo, Spain). The sequence was deposited in GenBank (accession no. **MN117139**).

Viral quantification by RT-qPCR

Viral RNA copies from different supernatants were determined by absolute quantification in an Applied Biosystems 7500 Fast-Real-time PCR system. The reaction mix was prepared by adapting the manufacturer's protocol for AgPath-ID™ One-Step RT-PCR Reagents (Applied Biosystems™): 2X RT-PCR Buffer= 1X; Forward and reverse PCR primers= 50μM each;

Syber Green (SYBR™ Green 10000X-Invitrogen) dilution 1/100= 5.0 and -5; 25X RT-PCR

Enzyme Mix (ArrayScript™ Reverse Transcriptase and AmpliTaq Gold®DNA Polymerase)=

1X. The final volume was 25 µl, using 2.5 µl of viral RNA. Cycling conditions: 50°C for 30

min; 45 cycles of 94°C for 2 min 95°C for 10 sec, 60°C for 30 sec and 72°C for 30 sec and a

final step of 72°C for 10 min. Data were analyzed in real-time using the Virtual Curve qPCR

program from Applied Biosystems (Thermo Fisher Connect™). Viral load was calculated from

a standard curve generated with serial dilutions (10^{-1} to 10^{-12}) of an hMPV N protein synthetic

oligonucleotide (199 bp) (Ultramer® DNA Oligo) with a known concentration (6.0219×10^{13}

copies of RNA/µl).

hMPV Replication Curve

Vero cells were grown (60% confluence) on glass coverslips (12mm in 24-well plate) for 24

hours and then infected with isolated hMPV (1.97×10^6 copies of RNA/µl). The culture

supernatants were harvested and the monolayers were fixed at different time points (8, 16, 24,

48, 72, and 96 hpi). Viral RNA was extracted from harvested supernatants, then the extracted

RNA was employed for the detection and quantification of hMPV. The mean number of copies

of extracellular viral RNA/µl of culture supernatant was then calculated (Applied biosystems.

Thermo-Fisher-Connect™).

Cells were processed for IF and the number of fluorescent dots/infected cell were calculated

through the analysis of fluorescence images using the ImageJ-Fiji software (ImageJ 1.50c

Wayne Rasband. NIH-USA).

Pharmacological treatment

Vero cells at 60% confluence cultured on glass coverslips were infected with hMPV. Infected

cells treated with CytD [2.5µM] [10] or vehicle (infection medium whitout CytD) were analized

at different times during the replication cycle. Cells were exposed to CytD or vehicle during

the 2 infection hours, as well as throughout the first 8 hpi (experimental period between 0-24

hpi); during the first and the last 24 hours (48-72 hpi) in an experimental period between 0-72

hpi. In each case, the supernatant was removed, the cells were washed three times with PBS

and the infection medium was added according to each treatment (Fig. 3B). Culture

supernatants and harvested cells were processed for RNA extraction, viral quantification by

RT-qPCR and IF, respectively.

Microscopy and digital image analysis

Immunostained cells were visualized using an inverted epi-fluorescence microscope (Olympus

IX81, Olympus). Images were taken with a regular fluorescence microscopy with a CCD

camera (Orca 1000, Hamamatsu-Corp.) and with an inverted confocal microscope FV1000

(Olympus). The collection and processing of data from the images were analyzed with the

ImageJ-Fiji software. Quantification of intracellular infection was carried out by the percentage

of infected cells and the number of fluorescent dots/ infected cell.

Statistical analysis

All tests and graphs were performed using GraphPad Prism 8.0 (GraphPad Software). Data

represent results from at least three independent experiments, and values are presented as the

mean ± standard error of the mean (SEM). Assumptions of homoscedasticity and normality

were tested using Brown-Forsythe and Barlett´s tests and D´Agostino & Pearson test,

respectively. All assumptions were met, allowing the use of Student's unpaired t-test or one-

way ANOVA test for specific group comparisons. A significance level of $p < 0.05$ was

considered statistically significant.

RESULTS

Human-Metapneumovirus isolation and identification

Human Metapneumovirus was isolated from NPA of hospitalized children in Cordoba, Argentina.

The virus replicated in Vero cells cultures after three blind passages of 21 days each one. The CPE was characterized by cell rounding, detachment, and formation of refractory cell clusters (Fig. 1A). IF detection confirmed that the CPE was caused by hMPV. The presence of specific hMPV proteins was observed at 4, 7, and 14 dpi located and distributed in the cell cytoplasm presenting a dotted pattern (Fig. 1B). Simultaneously, virus genome detection by nucleic acid amplification assays (RT-PCR) from supernatants of infected cultures collected at 4, 7, and 14 dpi were positive for the amplification of the 199 bp N protein (Fig. 1C), confirming the presence of hMPV in infected culture. Viral isolation was typified, the Cordoba/ARG/3864/2015 local isolated strain (no: MN117139) was identified as A_2 hMPV and grouped with Canada and The Netherlands prototype strains, and with sequences from Argentina [12,14], Peru, Brazil, United States, China, and Italy (Fig. 1F).

Different subcellular localization of large fluorescent dots were observed several times post-infection (Fig. 1B and D), while confocal microscopy detected small dots between 0 and 96 hpi in the subcortical area enriched in actin filaments, below the plasma membrane (Fig. 1E, low panel).

Figure 1. Human Metapneumovirus (hMPV) isolation and identification. A and B: Representative images of uninfected and infected VERO cells with hMPV. **A-** Images were obtained with phase contrast microscopy at low magnification, showing the viral infection cytopathic effect (CPE), which is visible at 14 d.p.i as cell rounding and shedding. Uninfected cells (top panel) and infected cells (bottom panel) at 4, 7 and 14 days dpi. Scale bar: 20 μm. **B-** The localization of the hMPV protein was determined using indirect immunofluorescence with specific anti-hMPV antibodies targeting protein F (green). MFLs were labeled with Phalloidin-Rho (red), and nuclei were stained with Hoechst (blue). A temporary sequence of 4 to 14 days in non infected (top panel) and infected cells (bottom panel). White arrows indicate the fluorescent label of the F protein. Scale bar: 20 μm. **C-** Amplification products (RT-PCR) of the hMPV N protein region (199 bp) by agarose gel electrophoresis. Bands corresponding to culture supernatants of infected cells collected at 4, 7 and 14 dpi and positive controls (C+) are shown. Uninfected cells collected at the same times are included (C-). MPM: molecular weight marker. **D-** Images obtained with epi-fluorescence microscopy, showing that the hMPV F protein accumulates near the nucleus at 48 hpi. Scale bar: 100 μm. **E-** Confocal microscopy images where viral proteins are observed at 0 and 96 hpi. The bottom panel shows a transverse optical section where the viral label is evidenced in the cell cytoplasm decorating the MFLs. Scale bar: 20 μm. **F-** A maximum-likelihood tree (PhyML software) was constructed, using the GTR+G model with parameters suggested by JModelTest 3.7 with bootstraps and 1000 pseudoreplicates. A1: strains to hMPV subgroup A1. A2: strains to hMPV subgroup A2. B1: strains to hMPV subgroup B1. B2: strains to hMPV subgroup B2. Avian MPV (aMPV) was

used to root the tree. The strain of interest is marked in bold. The scale bar indicates the changes
between nucleotides. All data were obteinded from 5 independet experiments.

The hMPV replication cycle

To characterize hMPV replication kinetics, we evaluated the production and release of the virus

over time (8, 16, 24, 48, 72, and 96 hpi) using qRT-PCR (RNA copies/μl of culture

supernatant). RNA copies increased progressively until 72 hpi, peaking at 4,961 copies/μl of

supernatant (Fig. 2A). The time course of hMPV infection was also studied by IF and viral

proteins was quantified as number of fluorescent dots/infected cell (Fig. 2B and D).

Interestingly, the number of fluorescent dots/infected cell decreased while the number of copies

of RNA/μl of culture supernatant increased. At 72 hpi the maximum extracellular viral yield

(4,961 copies) corresponded to the minimum number of cytoplasmic dots (2.2 dots), as well as

to the lowest percentage of infected cells (32.8%) (Fig. 2C). It is important to highlight that the

percentage of infected cells decreased significantly and progressively from 24 hpi until the end

of the experiment (Fig. 2C and D).

Taken together, our results demonstrate that the highest percentage of infected cells and

intracellular hMPV-protein contents were detected at the early phase of the replication cycle.

While the cycle progressed, hMPV immunostaining decreased whereas extracellular viral RNA

increased due to the release of the viral progeny.

Figure 2. HMPV viral replication curve. A- Quantification curve of extracellular viral production through the number of copies of viral RNA in the extracellular medium. Quantification was performed at 8, 16, 24, 48, 72 and 96 hpi. **B-** Quantification of intracellular viral production through the number of fluorescent dots/infected cell (cytoplasmic viral F protein). Quantification was performed at 8, 16, 24, 48, 72 and 96 hpi **C-** Percentage of quantified infected cells was determined by the number of cells with or without fluorescent dots at 8, 24, 48, and 72 hpi. **D-** Representative images of the temporal sequence of infection (0-96 hpi). The hMPV F protein label (green), MFLs (red) and nuclei (Hoechst, blue) are observed. Scale bar: 20 μm. More than 400 cells were screened in each time point. Data represent the mean ± SEM from 3 independent experiments (one-way ANOVA with Tukey test post hoc test). **** <0.0001.

Actin microfilament perturbation at different viral stages affects hMPV replication.

Previous studies highlight the involvement of the cytoskeleton, particularly MFLs, in the entry

and replication of numerous viruses [8]. Therefore, we studied the role of MFLs at early and

late stages of hMPV infection cycle. For this, disruption of actin polymerization was performed

using CytD at different times of hMPV infection. We first determined the participation of MFLs

during the entry and at its early stages of hMPV replication. To this end, CytD was applied

during virus infection (2 hours), during the first 8 hpi and during the first 24 hpi (Fig. 3B).

Treatment with CytD during early replication stages prevented the significant decrease in the

number of infected cells, maintaining initial percentage, regardless of drug exposure timing

(Fig. 3A and C). Interestingly, CytD treatment during the first 24 hpi, also prevented the

dramatic diminution in the percentage of infected cells at 72 hpi (75.3%) compared to control

(33.3%) (Fig. 3C). To obtain a quantitative measure of CytD effect on F viral protein expression

at the different stages of the replication cycle, we quantified the number of fluorescent

dots/infected cell. CytD treatment during infection or during the first 8 hpi significantly

increased fluorescent dots/infected cell compared to controls (Fig. 3A and D). Particularly,

during the first 8 hpi of treatment a 2 to 2.5 fold increase was observed at both analyzed times

(8 hpi and 24 hpi) (Fig. 3A and D). When CytD was applied during the first 24 hpi, and

quantified at the end of the treatment, no significant differences in fluorescent dots/infected

cells was observed compared to control, but prevented viral protein loss at 72 hpi (Fig. 3D).

Finally, we quantified viral-RNA copies in culture supernatant to assess extracellular virus

yields (Figure 3E). CytD treatment during the infection significantly increased viral production

at 24 hpi. Treatment during the first 8 hpi significantly increased RNA copies at that time. This

effect was also observed in the accumulated values, this value is the sum of the mean of RNA

copies values at 8 plus 24 hpi (for vehicle: 6,674; CytD treatment: 10,727; see accumulated

value at 24 hpi). On the other hand, CytD treatment during the first 24 hpi, caused a significant

decrease in extracellular RNA viral copies at 24 hpi, remaining lower up to 72 hpi, as can be

observed in the accumulated values (vehicle: 4,510; CytD treatment: 2,930; see accumulated

value at 72 hpi) (Fig. 3E).

In addition, we focused on the late stage of the replication curve and observed that CytD applied

during the last 24 hpi (48-72 hpi) caused no changes in the percentage of infected cells, in the

number of fluorescent dots/infected cell or in the number of extracellular RNA-viral copies

compared to controls (data not shown).

These results underscore the importance of the initial 8 hours of hMPV infection for replication,

indicating that CytD presence during this period boots viral protein expression in the cell

cytoplasm and enhances virus release into extracellular space.

Figure 3. Effect of disruption of MFLs with CytD at different stages of the viral cycle. A- Images of CytD treatments in hMPV infected cells at different times of the growth curve: CytD during infection (2 h), during 8 hpi and during 24 hpi. Immunofluorescence performed with specific hMPV antibodies anti-protein F (green), MFLs (Phalloidin-Rho, red) and nuclei (Hoechst, blue). **B-** The scheme illustrates the pharmacological treatment with CytD at different times during the viral replication cycle and the time points of experimental procedures. In red color: cells exposed to vehicle or CytD during the 2 infection hours. In green color: cells exposed to vehicle or CytD throughout the first 8 hpi. In violet color: cells exposed to vehicle or CytD during the first 24 hpi. **C-** Percentage of infected cells after CytD treatment in relation to vehicle control. CytD treatment during infection (2 h): a significant increase was observed at 24 hpi (p< 0.0001). CytD treatment during the first 8 hpi: a significant increase was observed at 24 hpi (p<0.0001). In CytD treatment during first 24 hpi a significant increase was observed both at 24 and 72 hpi (24 hpi: p<0.0001; 72 hpi: p<0.0001). **D-** Number of fluorescent dots per infected cell in CytD treatments and in controls. CytD treatment during infection (2 h): a significant increase was observed at 24 hpi (p< 0.0001). CytD treatment during the first 8 hpi: a significant increase was observed both at 8 and 24 hpi (8 hpi: p< 0.0001; 24 hpi: p< 0.0001). CytD treatment during the first 24 hpi: a significant increase was observed at 72 hpi (p<0.0001). **E-** Quantification of extracellular viral RNA. CytD treatment during infection (2 h): a significant increase in RNA copies in the culture supernatant was observed at 24 hpi (p<0.0001). CytD treatment during the first 8 hpi caused a significant increase in RNA copies in culture supernatant at 8 hpi (p<0.0001); this was also observed in the accumulated values

(sum of the means of RNA copies at 8 plus 24 hpi). CytD treatment during the first 24 hpi
caused a significant decrease in extracellular viral RNA copies ($p < 0.0001$), remaining constant
until 72 hpi; this was also observed in the accumulated values (sum of the means of RNA copies
at 24 and 72 hpi). Scale bar: 20 µm. More than 400 cells were screened in each case. Data
represent the mean $\pm$ SEM from 3 independent experiments. Student unpaired t test was
performed (**** <0.0001).

DISCUSSION

In this study we report the first successful isolation of hMPV performed in Vero cell line from

a positive clinical sample in Cordoba, Argentina. We identified this local strain and described

the replication curve showing that the maximum viral production takes place at 72 hpi.

Moreover, we demonstrate that the disruption of MFL at the early stages of infection increases

intracellular viral production and the release of viruses into the extracellular space.

The viral isolate was identified as the A_2 hMPV subtype through phylogenetic analysis,

consistent with our previous studies of A_2 hMPV subtype circulation in Argentina [12].

Additionally, co-circulation with other genotypes like A_1, B_1, and B_2, has been reported in our

country [14]. Genotype predominance varies based on factors like epidemiological year and

region, host immunity, viral load, and susceptibility. Competitive replication was observed

between hMPV genotype A and B. A recent study by van den Hoogen's group [15]

demonstrated that lineage A replicates more efficiently than lineage B in an organoid-derived

human bronchial-epithelium model, with increased viral-RNA and infectious virus particles.

They also highlight the use of viruses with a history of controlled passage, instead of extensively

passaged hMPV strains. These data are important because the A_2 genotype is clinically relevant,

as it may cause diseases of varying severity. In studies where verification was possible,

genotype A has been associated with severe conditions such as pneumonia and hypoxia, leading

to more ICU admissions compared to the other subtypes [16].

In several studies, hMPV replication is limited to certain cell lines, with varied and mild CPE.

Isolation of the A_2 hMPV viral strain in this study showed initial CPE signs at the third blind

passage, at 14 dpi, characterized by cell rounding and refractive cells clusters, consistent with

previous reports [7,17]. Conversely, other authors have described that hMPV can appear as a

syncytium [5,6,18], similar to RSV, and this difference may be related to the cell lines that were

infected (e.g. LLC-MK2, tMK, MNT-1, A549) or to the viral genotype [19–21]. Numerous

authors note the challenge of hMPV isolation and the absence of an obvious CPE [5,19,20,22].

Isolating respiratory viruses from clinical specimens often exhibits low efficiency *in vitro* due

to factors like cell type and virus subtype [17,22–24]. LLC-MK2 and Vero cells are commonly

used for all hMPV strains. Although, this study did not compare different cell lines and isolated

only the A2 hMPV subtype, efficient viral replication was achieved, consistent with findings

by Nao [21] who demonstrated high infectivity with different hMPV strains in VeroE6 and

Vero-ATCC cells.

The hMPV replication curve showed an exponential phase between 48 to 72 hpi, with maximum

intracellular viral protein concentration at 8 hpi which is consistent with results obtained by

other groups. Tollefson et al. [19] described hMPV kinetics in LLC-MK2 cells, with the eclipse

phase at 24 hpi and the exponential phase between 48 and 72 hpi with a significant increase in

viral titer. El Najjar [25] showed maximum viral production at 72 hpi in BEAS-2B cells, both

intra- and extracellularly. Recent studies using recombinant viruses in Vero cells, three-

dimensional cultures and organoid-derived bronchial culture have described similar results,

both in the CPE caused by the virus and in the replicative cycle [15,16,26]. All these results

show that our model of infection in the Vero-CCL cell line with this wild-type virus (A$_2$ hMPV),

despite its isolation difficulties, is very efficient and representative of *in vivo* infection.

It is known that viruses such as RSV [9] and PIXV [10], among others, take advantage of

structures such as the cytoskeleton for entry, replication and exit from the cell. Our study aimed

to assess the role of actin cytoskeleton in hMPV replication. For this, we used CytD, known to

disrupt actin polymerization. CytD concentration [2.5μM] was effective and consistent with

prior studies [24,25] as well as previously in our laboratory with another viral model [10]. The

rapid reversible action of CytD facilitated our experimental designs. Intracellularly, CytD not

only prevented the decrease in the percentage of hMPV-infected cells (during 2 hours of

infection, 8 hpi and 24 hpi) but it also caused a significant increase in the number of fluorescent

dots/infected cell, indicative of protein accumulations, in all treatments. In line with this, El

Najjar [25] postulates that hMPV proteins assembly during early infection does not require

actin polymerization. These protein accumulations (fluorescent dots), resembling "inclusion

bodies" [27,28] persist despite actin alteration, hindering viral release. Cifuentes-Muñoz [24]

note these structures as primary replication sites, particulary at 24 hpi. In *Pneumovirus*, actin

polymerization partially aids inclusion body formation [17]. Unlike Cifuentes-Muñoz [24], our

early CytD treatment (8 hpi) boosts extracellular viral load, suggesting that MFLs interruption

could facilitate viral entry. This represents an increase in intracellular viral load and a decrease

in virus release times, compared to the control. Nevertheless, our results are consistent with

those of Cifuentes-Muñoz [24] and El Najjar [25] regarding reduced extracellular viral load

from 24 hpi, particularly with continuous 24-hour treatment. Several studies suggest that hMPV

relies on cell-cell transmission for efficient spread and infection [8,28–30]. Actin disruption

may hinder this process, redirecting virions to alternate exit routes and leading to extracellular

accumulation. Therefore, actin dynamics is essential for hMPV infection, enabling cell-to-cell

spread regardless of the extracellular viral load [25]. This could explain cytoplasmic viral

protein accumulation and the decrease of extracellular viral load after 24 hpi CytD exposure.

Given MFLs are involved in viral particle internalization and release [24], it is reasonable to

think that short-term disruption of MFLs with CytD would facilitate viral particles

internalization. Conversely, prolonged CytD treatment (24 hpi), could prevent the transport of

viral proteins and/or nucleocapsid, reducing cytoplasmic accumulation and extracellular viral

loading. Despite the years since its discovery, the cell biology of hMPV infection remains

poorly understood. This study broadens our knowledge on the isolation and growth curve characterization of a wild-type hMPV obtained from a clinical sample. Furthermore, these findings provide new insights on hMPV infection and the role of actin cytoskeleton in viral replication, enhancing understanding of hMPV internalization and spreading mechanisms.

ACKNOWLEDGEMENTS

We would like to thank Bernadette van den Hoogen for receiving PER in her laboratory and for advising on the viral isolation protocol; Guadalupe Carballal[†] and Cristina Videla for generously providing hMPV wild type positive control; Celia Frutos and Lucia Ghietto for technical assistance in Real Time RT-qPCR; Santiago Mirazo for generously providing polyclonal antibodies; Carlos Mas for technical assistance in confocal microscopy at the Centro de Micro y Nanoscopía de Córdoba, CEMINCO-CONICET-Universidad Nacional de Córdoba, Córdoba, Argentina. We particularly want to thank Miss Stella Ascheri for general technical assistance.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author [gpaglini@immf.uncor.edu] upon reasonable request.

DECLARATIONS

Conflict of Interest. The authors have no competing interests to declare that are relevant to the content of this article.

Funding source. This work was supported by grants from Secretaria de Ciencia y Tecnología-

Universidad Nacional de Córdoba (SECYT-Consolidar-I Nº 336 20180100191CB to A.C. and

SECYT-Consolidar-C Nº 33620180100091CB to M.G.P.) and from Fundación A. J. Roemmers

(2018-2020 to A.C. and 2017-2019 to P.E.R). The funding sources had no involvement in the

study design, data collection, analysis and interpretation, writing of the manuscript or in the

decision to submit the article for publication.

REFERENCES

1.   Afonso CL, Gaya KA, Bányai K, et al.: TAXONOMY OF THE ORDER

MONONEGAVIRALES: UPDATE 2016. Arch Virol **2016**; 161:2351–2360.

2.   Shafagati N, Williams J: Human metapneumovirus - what we know now.

F1000Research **2018**; 7:1–11.

3.   Kampmann B, Madhi SA, Munjal I, et al.: Bivalent Prefusion F Vaccine in Pregnancy

to Prevent RSV Illness in Infants. N Engl J Med **2023**; 388:1451–1464.

4.   Ballegeer M, Saelens X: Cell-Mediated Responses to Human Metapneumovirus

Infection. Viruses **2020**; 12:3–22.

5.   Van Den Hoogen BG, De Jong JC, Groen J, et al.: A newly discovered human

pneumovirus isolated from young children with respiratory tract disease. Nat Med

2001; 7:719–724.

6.   Sato K, Watanabe O, Ohmiya S, et al.: Efficient isolation of human metapneumovirus

using MNT-1, a human malignant melanoma cell line with early and distinct cytopathic

effects. Microbiol Immunol **2017**; 61:497–506.

7.   Deffrasnes C, Côté S, Boivin G: Analysis of Replication Kinetics of the Human

Metapneumovirus in Different Cell Lines by Real-Time PCR. J Clin Microbiol **2005**;

43:488–490.

8.   Cifuentes-Munoz N, El Najjar F, Dutch RE: Viral cell-to-cell spread: Conventional and

non-conventional ways. In: Advances in Virus Research. **2020**. p. 85–125.

9.   Shahriari S, Wei KJ, Ghildyal R: Respiratory syncytial virus matrix (M) protein

interacts with actin in vitro and in cell culture. Viruses **2018**; 10:7–12.

10. Gil PI, Albrieu-Llinás G, Mlewski EC, et al.: Pixuna virus modifies host cell

cytoskeleton to secure infection. Sci Rep **2017**; 7:1–9.

11. Bouscambert-Duchamp M, Lina B, Trompette A, Moret H, Motte J, Andréoletti L:

Detection of human metapneumovirus RNA sequences in nasopharyngeal aspirates of

young french children with acute bronchiolitis by real-time reverse transcriptase PCR

and phylogenetic analysis. J Clin Microbiol **2005**; 43:1411–1414.

12. Rodriguez PE, Frutos MC, Adamo MP, et al.: Human Metapneumovirus:

Epidemiology and genotype diversity in children and adult patients with respiratory

infection in Córdoba, Argentina. PLoS One **2020**; 15:e0244093.

13. Van Den Hoogen BG, Osterhaus DME, Fouchier RAM: Clinical impact and diagnosis

of human metapneumovirus infection. Pediatr Infect Dis J **2004**; 23:25–32.

14. Velez Rueda AJ, Mistchenko AS, Viegas M: Phylogenetic and Phylodynamic Analyses

of Human Metapneumovirus in Buenos Aires (Argentina) for a Three-Year Period

(2009-2011). PLoS One **2013**; 8:1–10.

15. Ribó-Molina P, van Nieuwkoop S, Mykytyn AZ, et al.: Human metapneumovirus

infection of organoid-derived human bronchial epithelium represents cell tropism and

cytopathology as observed in in vivo models. Virology **2024**; 9:1–13.

16. Arnott A, Vong S, Sek M, et al.: Genetic variability of human metapneumovirus

amongst an all ages population in Cambodia between 2007 and 2009. Infect Genet

Evol **2013**; 15:43–52.

17. Kinder JT, Moncman CL, Barrett C, Jin H, Kallewaard N, Dutch RE: Respiratory

Syncytial Virus and Human Metapneumovirus Infections in Three-Dimensional

Human Airway Tissues Expose an Interesting Dichotomy in Viral Replication, Spread,

and Inhibition by Neutralizing Antibodies. J Virol **2020**; 94:1–20.

18.   Bernal LJ, Velandia-Romero M, Guevara C, Castellanos JE: Human

Metapneumovirus: Laboratory Methods for Isolation, Propagation, and Plaque

Titration. Intervirology **2019**; 61:301–306.

19.   Tollefson SJ, Cox RG, Williams J V: Studies of culture conditions and environmental

stability of human metapneumovirus. Virus Res **2010**; 151:54–59.

20.   Jumat MR, Nguyen Huong T, Wong P, et al.: Imaging analysis of human

metapneumovirus-infected cells provides evidence for the involvement of F-actin and

the raft-lipid microdomains in virus morphogenesis. Virol J **2014**; 11:1–12.

21.   Nao N, Sato K, Yamagishi J, et al.: Consensus and variations in cell line specificity

among human metapneumovirus strains. PLoS One **2019**; 14:1–21.

22.   Lee H, Woo HM, Kim K, et al.: Improving pneumovirus isolation using a

centrifugation and AZD1480 combined method. J Microbiol Biotechnol **2019**;

29:2006–2013.

23.   Cox RG, Mainou BA, Johnson M, et al.: Human Metapneumovirus Is Capable of

Entering Cells by Fusion with Endosomal Membranes. PLoS Pathog **2015**; 11:1–29.

24.   Cifuentes-muñoz N, Branttie J, Slaughter KB, Dutch RE: Human Metapneumovirus

Induces Formation of Inclusion Bodies for Efficient Genome Replication and

Transcription. J Virol **2017**; 91:1–18.

25.   El Najjar F, Cifuentes-Muñoz N, Chen J, et al.: Human metapneumovirus Induces

Reorganization of the Actin Cytoskeleton for Direct Cell-to-Cell Spread. PLoS Pathog

2016; 12:1–30.

26.   Geiser J, Boivin G, Huang S, et al.: RSV and HMPV Infections in 3D Tissue Cultures:

Mechanisms Involved in Virus-Host and Virus-Virus Interactions. Viruses **2021**; 13:1–15.

27. Derdowski A, Peters TR, Glover N, et al.: Human metapneumovirus nucleoprotein and phosphoprotein interact and provide the minimal requirements for inclusion body formation. J Gen Virol **2008**; 89:2698–2708.

28. Najjar F El, Castillo SR, Moncman CL, et al.: Imaging analysis reveals budding of filamentous human metapneumovirus virions and direct transfer of inclusion bodies through intercellular extensions. MBio **2023**; 14:1–18.

29. Merwaiss F, Czibener C, Alvarez DE: Cell-to-cell transmission is the main mechanism supporting bovine viral diarrhea virus spread in cell culture. J Virol **2018**; 93:1–18.

30. Boggs KB, Edmonds K, Cifuentes-Munoz N, et al.: Human Metapneumovirus Phosphoprotein Independently Drives Phase Separation and Recruits Nucleoprotein to Liquid-Like Bodies. MBio **2022**; 13:.