Homeostatic Control of Deep Sleep in *Drosophila*: Implications for Discovering

Correlates of Sleep Pressure

Budhaditya Chowdhury[1]*, Lakshman Abhilash[1]*, Antonio Ortega[2], Sha Liu[2], Orie Shafer[1#]

[1] The Advanced Science Research Center, The City University of New York; The Graduate Center

at the City University of New York

[2] VIB-KU Leuven Center for Brain & Disease Research, Leuven, Belgium

*These authors contributed equally

Running title: Drosophila sleep homeostasis

[#]Correspondence: oshafer@gc.cuny.edu

Abstract

Homeostatic control of sleep is typically addressed through mechanical stimulation-induced forced wakefulness and the measurement of subsequent increases in sleep. A major confound attends this approach: biological responses to deprivation may reflect a direct response to the mechanical insult rather than to the loss of sleep. Similar confounds accompany all forms of sleep deprivation and represent a major challenge to the field. Here we describe a new paradigm for sleep deprivation in *Drosophila* that fully accounts for sleep-independent effects. Our results reveal that deep sleep states are the primary target of homeostatic control and establish the presence of multi-cycle sleep rebound following deprivation. Furthermore, we establish that specific deprivation of deep sleep state results in state-specific homeostatic rebound. Finally, by accounting for the molecular effects of mechanical stimulation during deprivation experiments, we show that serotonin levels track sleep pressure in the fly's central brain. Our results illustrate the critical need to control for sleep-independent effects of deprivation when examining the molecular correlates of sleep pressure and call for a critical reassessment of work that has not accounted for such non-specific effects.

Keywords: Drosophila, sleep, Ethoscope, yoked controls, mechanical deprivation, sleep rebound, homeostasis, MALDI-TOF MS

Introduction

Sleep is a fundamental biological need and is homeostatically controlled: when sleep is disrupted an elevated drive to sleep (i.e., sleepiness) is produced that is associated with decreased performance, well-being, and safety (Dijk and von Schantz, 2005; Holding et al., 2020; Krause et al., 2017; Lane et al., 2022; Meyer et al., 2022). Sleep disruption is widespread in the modern age and contributes significantly to a wide array of negative health consequences in humans (Colten et al., 2006; Medic et al., 2017). It is therefore critical to understand the homeostatic regulatory mechanisms of sleep. Sleep is ubiquitous in the animal kingdom and sleep-like states likely evolved at least 740 million years ago (Campbell and Tobler, 1984; Freiberg, 2020; Kanaya et al., 2020; Keene and Duboue, 2018; Nath et al., 2017; Webb, 1974). Being so deeply conserved, mechanisms of sleep regulation uncovered in animals with relatively simple nervous systems can be highly relevant to sleep in mammalian species, including humans. The relative simplicity of invertebrate nervous systems and the existence of invertebrate models for which there are well-established methods of genetic and neuronal manipulations make them amenable to experimental approaches that would be technically and logistically difficult or prohibitively expensive in mammalian model systems. Work in such model systems is therefore an important complement to studies in mammals and other vertebrate models (Davis and Raizen, 2017).

The vinegar fly *Drosophila melanogaster* has proved to be a useful model organism for understanding the molecular and cellular mechanisms regulating sleep (Hendricks et al., 2000; Shaw et al., 2000). The powerful genetic tools available in flies, along with the ability to manipulate and measure neural signaling within small groups of defined cell types, have revealed genes and brain networks that mediate the homeostatic regulation of sleep (De Jesús-Olmo et al.,

2020; Donlea et al., 2014; Ni et al., 2019; Seidner et al., 2015). Furthermore, such work in the fly

has revealed mechanisms that also regulate sleep in mammals (Liu et al., 2016; Shafer and Keene,

2021). Just as in humans and other mammalian species, fly sleep is controlled by both a homeostat

and a circadian clock, which, during normal sleep cycles, largely account for the timing and

amount of daily sleep (Abhilash and Shafer, 2022; Borbély, 1982; Borbély et al., 2016; Deboer,

2018). Future work in the fly will undoubtedly continue to enrich our understanding of sleep

regulation in all animals.

Nevertheless, there are significant challenges to the investigation of sleep regulation in

Drosophila. For example, the magnitude of sleep rebound in flies is quite modest (Huber et al.,

2004; Klose and Shaw, 2021; Oh et al., 2014; Vaccaro et al., 2020). Indeed, some forms of

optogenetic or thermogenetic sleep deprivation produce no apparent homeostatic sleep rebound at

all (Seidner et al., 2015; Vaccaro et al., 2020). As for other animal models of sleep, the action of

the fly's sleep homeostat is assayed by depriving flies of sleep and measuring subsequent increases

of sleep (compared to baseline), which reflect a homeostatic discharge of the sleep debt accrued

during deprivation (Hendricks et al., 2000; Huber et al., 2004; Melnattur et al., 2020; Shaw et al.,

2000). In contrast to mammalian species, however, sleep rebound following deprivation is often

quite modest and brief. For example, male flies display variable levels of increased sleep following

nighttime sleep deprivation of around 0-25 minutes, compared to 200-500 minutes of sleep lost (in

6- and 12-h deprivation windows) and this is usually detectable only during the first few hours of

the day after deprivation has ended (Geissmann et al., 2019, 2017; Hendricks et al., 2000; Huber

et al., 2004; Klose and Shaw, 2021). Furthermore, homeostatic sleep rebound in *Drosophila*

appears to repay a relatively small proportion of the sleep lost during deprivation; 800 minutes of

sleep loss in a full diurnal cycle was followed by only 100 minutes of recovery sleep (~12.5%)

(Huber et al., 2004). In contrast, mammalian species tend to display higher amounts of increased

sleep following deprivation that is observable over more than one circadian cycle (Berger and

Oswald, 1962; Mistlberger et al., 1983; Nakazawa et al., 1978; Tobler et al., 1983; Williams et al.,

1964), and such rebound usually accounts for a much larger proportion of lost sleep than the

rebound detectable in flies. For example, humans can reclaim around 40% of the sleep lost to

deprivation (Stroemel-Scheder and Lautenbacher, 2021). The relatively small rebound measured

in the fly is a major challenge to the study of sleep regulation in this species, as it provides a small

dynamic range over which to detect changes in homeostatic responses in experimental subjects.

These features of fly sleep are somewhat surprising, given the highly conserved nature of sleep in

the animal kingdom.

One possible cause of the relatively small homeostatic responses to sleep deprivation in flies are

the sleep-independent effects of the deprivation techniques used by the field. The most commonly

used method of fly sleep deprivation is frequent mechanical perturbation, typically shaking or

slamming (Hendricks et al., 2000; Melnattur et al., 2020; Shaw et al., 2000), to force movement

and prevent the attainment of sleep, which in *Drosophila* is defined as any bout of inactivity that

is five minutes or longer (Hendricks et al., 2000; Shaw et al., 2000). Though effective for the

prevention of sleep, these methods are likely to produce many physiological and behavioral effects

that are independent of sleep loss. Indeed, mechanical shaking is known to produce the

biochemical hallmarks of stress, even when delivered during times of wakefulness (Harbison and

Sehgal, 2009).

Recently, neurogenetic methods have been employed to deprive flies of sleep (Dubowy et al.,

2016; Vaccaro et al., 2020). This method involves the strong and chronic excitation of wake-

promoting neurons in the brain. Though this method avoids the sleep-independent effects of

physical perturbation, the strong non-physiological excitation of central brain neurons is likely to

be attended by its own sleep-independent effects. A fundamental challenge, therefore, is to

differentiate sleep-pressure-driven changes in behavior and physiology from sleep-independent

changes driven by the manipulation used to prevent sleep. We predict that accounting for such

sleep-independent effects would improve the sensitivity with which homeostatic sleep responses

can be detected in the fly.

In mammals, sleep consists of multiple, physiologically discrete stages, which differentially

respond to sleep deprivation (Achermann et al., 1993; Berger and Oswald, 1962; Endo et al., 1997;

Everson et al., 1989; Loomis et al., 1937; Rechtschaffen et al., 1999; Takahashi et al., 1978).

Following long bouts of wakefulness during normal sleep cycles, slow-wave Non-Rapid-Eye-

Movement (NREM) sleep stages dominate the initial bouts of sleep, suggesting that the daily rise

of sleep pressure is discharged first by slow-wave (i.e., deep) sleep (Borbély and Achermann,

1999; Webb, 1974). Slow-wave sleep is also preferentially increased following sleep deprivation

when compared to Rapid-Eye-Movement (REM) sleep (Berger and Oswald, 1962; Borbély and

Neuhaus, 1979; Dijk et al., 1990; Williams et al., 1964). Thus, differentiating between distinct

sleep stages is likely important for assessing the homeostatic responses to sleep deprivation.

A growing body of evidence indicates the existence of a deep sleep state in flies, which is

characterized by distinct patterns of brain signaling (Nitz et al., 2002; van Alphen et al., 2013),

reduced metabolic rate (Stahl et al., 2017), and waste clearance from the brain (van Alphen et al.,

2021). The physiological and metabolic correlates of this deep sleep state are associated with bouts

of inactivity that are significantly longer than the five-min inactivity criterion used to define fly

sleep (Nitz et al., 2002; Stahl et al., 2017; van Alphen et al., 2021, 2013). Furthermore, recent

studies have suggested that periods of inactivity briefer than the 5-min criterion may represent an

"active sleep" state in the fly (Niki et al., 2023; Tainton-Heap et al., 2021). The existence of

multiple sleep states in the fly suggests that treating sleep as a unitary state in this species might

obscure homeostatic sleep responses, particularly if, as in mammals, specific states more strongly

linked to homeostatic sleep pressure than shallower stages of sleep. Furthermore, our recent

analyses suggest that longer bouts of sleep are a better reflection of sleep homeostasis than

traditional definitions of sleep in the fly (Abhilash and Shafer, 2022).

In this study we address the sleep-independent behavioral effects of the most commonly used form

of sleep deprivation in the fly and develop a method of accounting for them. We also attempt to

address the extent to which the treatment of sleep as a single unitary state in the fly might explain

the apparently modest and brief nature of sleep rebound in the fly compared to such rebound in

mammalian species. We present evidence that the extent of mechanical perturbation employed

during sleep deprivation has significant effects on the amount of subsequent sleep displayed by

deprived flies. We describe the development of a yoked-control paradigm for flies, based on

previous work in rats (Rechtschaffen et al., 1983), that allows us to produce two sets of flies that

have experienced identical levels of mechanical perturbation while suffering significantly different

amounts of sleep restriction. Furthermore, by differentiating between long-bouts of sleep from the

traditional unitary definition of sleep, we show that flies display significant and lasting homeostatic

sleep increases following sleep deprivation that are only detectable when controlling for the sleep-

independent effects of mechanical deprivation. Finally, we illustrate the importance of yoked

controls for examining the molecular correlates of sleep pressure and identify serotonin as a

molecule that is increased within the brain in proportion to sleep loss but not mechanical

perturbation. Our work introduces methodological approaches that are likely to support the

discovery of new mechanisms of sleep regulation in the fly, calls for the reevaluation of previous

work identifying the molecular, physiological, and cellular correlates of sleep pressure, and

suggests that serotonin may act as a sleep substance within the central brain.

Results

Sleep rebound is shaped significantly by the frequency of the mechanical stimulus used to prevent sleep.

The drive to sleep in the face of prolonged sleep deprivation is so strong that significant experimental intervention is necessary to keep animals awake. Such intervention is known to produce sleep-independent changes in behavior and physiology in addition to changes caused by increased sleep pressure (Nollet et al., 2020). Traditionally, sleep-deprived flies, which have undergone forced wakefulness by means of mechanical or neurogenetic perturbation, are compared to unperturbed flies (e.g., Hendricks et al., 2000). Thus, it is not possible to differentiate between sleep-dependent and sleep-independent effects of deprivation, as the experimental flies have experienced both sleep deprivation and experimental insults (typically, physical agitation), whereas control flies have experienced neither. A further complication in assessing sleep rebound is the duration and time of day during which forced wakefulness stimuli are applied (Huber et al., 2004) as both levels of sleep pressure and diurnal/circadian changes in physiology can have complex interactions with the perturbing stimuli.

Though the minimum criterion for a sleep-like state in *Drosophila* is five continuous minutes of inactivity, the most common means of mechanical sleep deprivation in the field consists of the delivery of mechanical stimulation (shaking or slamming) for two seconds randomly within every 20 second interval (Geissmann et al., 2019; Hendricks et al., 2000; Shaw et al., 2000). Once the permissive window of sleep opportunity is open after the period of sleep deprivation, usually timed to the beginning of the day, sleep is measured across next few hours and the amount of sleep

displayed by sleep-deprived flies is compared to unperturbed controls (Beckwith et al., 2017;

Geissmann et al., 2019, 2017). The amount of baseline sleep displayed by sleep-deprived and

control flies before deprivation is also often accounted for in the assessment of rebound to control

for small but significant differences in baseline sleep that are often observed between experimental

and control flies, even when the two groups are genetically identical (Geissmann et al., 2019)

(Figure 1A). Given the substantial discrepancy between the inactivity duration criterion for sleep

(five minutes or more) and the frequency of mechanical stimulation used to deprive flies of sleep

(multiple times per minute) we first asked how rebound sleep might differ between flies that were

sleep deprived with different frequencies of mechanical stimulation. Therefore, we varied the

duration of mechanical deprivation to compare our results with previous deprivation studies, which

have employed a various durations of sleep deprivation (Huber et al., 2004; Nall and Sehgal, 2013;

Seidner et al., 2015).

As previously described in multiple studies by others using the *Drosophila* Activity Monitor

(DAM) system, flies deprived of sleep for 6-, 12-, and 24-h using two seconds of mechanical

stimulation delivered at random times within each 20-s interval displayed immediate increases in

total sleep compared to undisturbed flies (Figure 1A, D, G) (Cirelli et al., 2005; Hendricks et al.,

2000; Huber et al., 2004). In each of these instances a significant 24-h rebound was observed when

compared to undisturbed flies (Figure 1C, F, I). This reproducibility with previous studies

confirmed that our lab population of *Canton-S* flies display homeostatic sleep increases like those

described in previous studies (Hendricks et al., 2000; Huber et al., 2004; Melnattur et al., 2020;

Nall and Sehgal, 2013; Shafer and Keene, 2021; Shaw et al., 2000). To ask how the extent of

mechanical stimulation shapes homeostatic sleep rebound, we decreased the frequency of such

stimulation while maintaining the three durations (6-h, 12-h, and 24-h) of sleep deprivation, and

ensuring effective prevention of sleep. In order to examine the effects of stimulation frequency on

sleep rebound, we decreased the mechanical simulation frequency to once every 220 seconds. This

low-frequency trigger produced sleep deprivation for all the deprivation durations tested (Figure

1B, E, H). Surprisingly, no sleep rebound was observed for 6-h, 12-h, or 24-h deprivation

experiments, despite the fact that the lower frequency stimulations effectively deprived sleep

(Figure 1C, F, I). Lowering the intensity of mechanical stimulation therefore appeared to eliminate

sleep rebound, despite having prevented sleep during deprivation.

Examination of the magnitudes of sleep rebound following high frequency mechanical deprivation

revealed that the 6-h deprivation protocol produced higher sleep rebound than did 12-h, and 24-h

deprivation, despite the fact that this duration produced the smallest loss of sleep (Figure 1J). To

further confirm the role of mechanical disturbance in recovery sleep assessment, we added an

additional trigger frequency of random shaking for two seconds within every 120 second interval

for 24-h of sleep deprivation. We compared the sleep rebound produced by 24-h of sleep

deprivation with stimulation frequencies of 20-s, 120-s, and 220-s using a two-way ANOVA and

found that there was a significant interaction effect of treatment × trigger frequency on the amount

of sleep rebound (total sleep post-deprivation minus total sleep pre-deprivation). Remarkably, we

found significant rebound only for the flies deprived with 20-s trigger frequencies (Figure 1K). It

is also important to note that although the higher trigger frequency produced a significant rebound,

the amount of recovered sleep was very small (Figure 1L).

Thus, lower frequencies of mechanical stimulation failed to produce significant sleep rebounds,

despite having produced similar extents of sleep deprivation. To address the extent to which our

sleep deprived flies recovered lost sleep, we estimated how much lost sleep was recovered across

the 24-h following the end of the deprivation window (Donlea et al., 2014; Shaw et al., 2002). We

found that, as expected from previous work in many labs, the flies deprived of sleep using the most

frequent mechanical stimulation (every 20s) displayed apparent homeostatic sleep increases in the

hours immediately following deprivation (Figure 1A, D, G). In contrast, flies that experienced less

frequent stimulation displayed little if any such apparent rebound (Figure 1B, E, H). The fact that

lower levels of mechanical stimulation failed to produce rebound over one full cycle, despite

having produced similar amounts of sleep deprivation, suggests that high frequency physical

perturbation produces sleep-independent behavioral effects that might mask *bona-fide* homeostatic

sleep increases. Alternatively, it is also possible that during lower stimulation frequencies, flies

may increase their levels of the recently hypothesized "active sleep" state (inactivity of between

one to five minutes in duration) (Niki et al., 2023; Tainton-Heap et al., 2021) and that this may

prevent the build-up of sleep pressure and rebound. We examined the levels of such sleep during

and after deprivation using 220-s stimulation frequencies and found that the flies do display

substantial increases in this active sleep state throughout deprivation using 220-s stimulation

frequencies (Supplementary Figure 1). However, given the strong convergent evidence that

periods of inactivity of five minutes or more represent a physiologically distinct sleep-like state, it

is surprising to see the absence of rebound following deprivation of this sleep state in our

experiments using 220-s stimulation frequencies and standard sleep definitions. The absence of

rebound in these experiments supports the hypothesis that the apparent homeostatic rebound of fly

sleep is strongly shaped by the frequency of mechanical stimulation. Accounting for the sleep-

independent effects of mechanical deprivation on sleep and examining the homeostatic control of

distinct sleep states are likely to reveal previously hidden characteristics of sleep homeostasis.

Accounting for mechanical stimulus using a yoked-controlled design uncovers significant

sleep rebound in sleep-deprived flies when low-frequency triggers are used.

The mechanical shaking employed to sleep deprived flies loaded into the DAM system use near

constant agitation without regard to the flies' sleep state, delivering mechanical stimulation to both

active and inactive flies. To further examine the potential sleep-independent effects of mechanical

perturbation during sleep deprivation experiments, we adopted the Ethoscope, a video-based

system for recording fly behavior (Geissmann et al., 2017). In contrast to the DAM system, which

requires the simultaneous mechanical perturbation of all flies within the same DAM monitor, the

Ethoscope allows the user to deliver mechanical stimuli to individual flies only when a fly is

inactive for a specified amount of time (Geissmann et al., 2017). The ability to track single flies

and selectively stimulate individual flies to prevent sleep means that flies can be sleep deprived

with significantly fewer mechanical perturbations, and only when flies have been inactive for

defined durations.

We programmed Ethoscopes to track single flies and rotate their tubes at ~420 rpm for one second

each time a fly had been inactive for 220 seconds, which resulted in 24-h of sleep deprivation when

using the standard inactivity criterion for fly sleep (5-min or more) (Figure 2A, C). For both

Canton-S and w^{1118}, when compared to undisturbed flies, the sleep-deprived flies failed to display

significant increases in sleep during the 24-h following deprivation (Figure 2B, D); a surprising

result given the clear effectiveness of sleep deprivation (Figure 2A, C), but consistent with the lack

of rebound in similar situations using the vortexer (see Figure 1). This result is also surprising on

another account – the use of ethoscopes to measure sleep does not overestimate and saturate sleep

as much as the IR-beam crossing based method in the DAMs (compare Figures 1 and 2), thereby

providing a larger dynamic range wherein rebound could have been observed.

The failure to produce clear homeostatic sleep increases following deprivation using a 220-s

inactivity criterion could be the product of two causes. As suggested by our observations above,

the mechanical insult experienced by flies during sleep deprivation are likely to exert sleep-

independent effects on the behavior observed after deprivation. That is, the mechanical disturbance

experienced by the sleep-deprived flies may produce effects on the fly that could obscure sleep-

pressure-driven behavioral changes. Second, the 220-s inactivity triggers used to deprive flies of

sleep may have allowed brief inactivity bouts recently hypothesized to represent an "active sleep"

state (Niki et al., 2023; Tainton-Heap et al., 2021) and such a state prevented the build-up of

sufficient sleep pressure to produce a homeostatic response. This would suggest that the five-min

inactivity criterion for sleep may be too long to capture all forms of homeostatically-controlled fly

sleep. To begin to address these two possibilities, we first considered the effects of mechanical

perturbation and therefore sought to produce flies that differed in the extent of their sleep

deprivation, despite experiencing identical mechanical stimulation. Comparing sleep in such flies

would allow us to examine the specific contribution of sleep pressure to sleep behavior following

deprivation.

To accomplish this, we adopted a paradigm of "yoked" sleep deprivation that was first introduced

in the rat model, which produces pairs of rats that have experienced perfectly matched mechanical

stimulation, in the form of a rotating platform over water, that nevertheless differed significantly

in the amount of sleep deprivation experienced (Rechtschaffen et al., 1983). This approach

involved a "focal" animal which was placed on one side of the platform and monitored via

electroencephalogram recording. Whenever the focal animal displayed the physiological

hallmarks of sleep, the platform would begin to rotate, thereby keeping the focal animal awake. A

paired rat, the "yoked" control, was placed on the opposite side of the platform and could sleep

when the focal rat was awake. This arrangement produced a pair of rats that experienced identical,

time-matched mechanical stimulation while experiencing significantly different amounts of sleep

loss (Rechtschaffen et al., 1983).

We programmed the Ethoscope platform to produce a paired yoked-control fly for every sleep-

deprived focal fly. We conducted these experiments for a 24-h period commencing at lights ON

to avoid the circadian confounds of mechanical stimulations given at specific times of the day or

night. For each fly that was tracked and stimulated upon being inactive for 220-s, a second fly

received matched, time-locked tube rotations, which were delivered independently of its

sleep/wake state. This ensured that each focal and yoked pair received identical mechanical stimuli

(Figure 2E). 220-s inactivity criteria were necessary to provide sufficient time for our yoked

controls to attain substantial amounts of sleep, as defined by the standard criterion of 5-min or

more of inactivity. Indeed, using the 220-s inactivity trigger allowed yoked flies to attain sleep

while the focal fly was active (Figure 2F and G). Yoked controls, therefore, suffered only partial

sleep loss compared to focal flies (Figure 2H and J). This approach, therefore, succeeded in

producing two sets of flies that had experienced the same level of physical perturbation while

suffering significantly different levels of sleep loss.

A comparison of total sleep across the entire 24-h of the post-deprivation day revealed significantly

higher levels of sleep in the focal flies compared to yoked controls for both *Canton-S* and *w[1118]*

(Figure 2I and K). Thus, the inclusion of yoked controls revealed a homeostatic sleep response

that was not apparent when comparing sleep-deprived flies to unperturbed controls, supporting the

notion that sleep-independent effects of mechanical perturbation can mask homeostatic sleep

responses. Both *Canton-S* (Figure 2H and I) and *w[1118]* (Figure 2J and K) strains displayed such

rebound, exhibiting increases in both daytime and nighttime sleep during the 24-h following

deprivation. Mechanical stimulation during deprivation significantly increased the activity levels

of sleep-deprived flies a day after mechanical perturbation, when compared to undisturbed

controls. This increase was seen immediately after the light-to-dark transition, i.e., at the time

associated with the transition to night-time sleep (Supplementary Figure 2A and B). This

difference was not observed when comparing focal flies with yoked controls (Supplementary

Figure 2C and D). Though yoked controls revealed significant sleep-pressure specific increases in

sleep, this homeostatic increase was quite modest, resulting in very little discharge of the sleep

pressure built during the 24-h of sleep deprivation (Figure 2I and K). Given that most animals

display homeostatic sleep increases over multiple sleep/wake cycles (Mistlberger et al., 1983;

Nakazawa et al., 1978; Rechtschaffen et al., 1999), we wondered if the rebound we detected using

yoked controls might persist over multiple diurnal cycles.

Sleep rebound accumulates over multiple cycles in response to infrequent mechanical deprivation, but little sleep debt is paid off.

The circadian control of sleep thresholds ensures that sleep is gated to occur at the appropriate/adaptive time of day, even in the presence of increased sleep drive. The action of the circadian system therefore prevents sleep debt from being discharged completely upon the first opportunity to sleep, ensuring that wakefulness will occur during the next diurnal cycle despite the presence of homeostatic sleep pressure. For this reason, sleep debt is typically discharged over several sleep/wake cycles (Mistlberger et al., 1983; Nakazawa et al., 1978; Rechtschaffen et al., 1999). Current methods in the *Drosophila* sleep field do not typically produce multi-cycle rebound, but instead, reveal fairly modest sleep rebound during the first hours of the day-time following deprivation (Beckwith et al., 2017; Geissmann et al., 2019, 2017). In our yoked-controlled experiment, we observed only modest sleep rebound that discharged very little of the sleep lost during the previous day's deprivation, which may have been a product of the smaller but significant deprivation experienced by yoked controls. However, in this case, sleep rebound was apparent across the diurnal cycle, rather than being limited to the first few hours of the recovery day (Figure 2H and J). We wondered if the comparison of deprived flies with yoked controls might reveal persistent increases in sleep for deprived flies across subsequent sleep/wake cycles. We therefore assessed the behavior of flies deprived of sleep using the Ethoscope for three full diurnal cycles following 24-h of sleep deprivation (Figure 3A and B) and compared the sleep of deprived flies with sleep observed in both undisturbed and paired yoked controls.

When compared to yoked controls, focal sleep-deprived flies displayed modest but significant

increases in cumulative sleep over two successive cycles (Figure 3D). Once again, these increases

were not detectable when deprived flies were compared to undisturbed controls (Figure 3C). In

this case, baseline sleep on the days before deprivation was not different between the three groups

of flies (see Figure 2B and G). We examined how much lost sleep was recovered across the three

post-deprivation cycles by quantifying sleep gain in focal flies, which we calculated as the increase

in total post-deprivation sleep compared to baseline sleep and normalized to sleep in undisturbed

controls to account for normal changes in sleep during the duration of the experiment (see

methods). Despite this apparent two-cycle rebound, focal flies failed to discharge a substantial

portion of the sleep debt accrued during deprivation (Figure 3E). This result is surprising because

when we estimated p(Doze), a recently introduced conditional probability metric for sleep pressure

(Wiggin et al., 2020), we found that the focal flies show sustained increases in sleep pressure for

at least three days post sleep deprivation (Supplementary Figure 3).

We wondered if the apparent absence of substantial homeostatic payback of sleep lost by focal

flies during deprivation might be explained by the fact that current sleep analysis methods do not

recognize distinct sleep states, treating sleep as a unitary state in which all bouts of inactivity of

five minutes or more are considered to be the same sleep state. Given the growing evidence of

physiologically and metabolically distinct sleep states in the fly (Stahl et al., 2017; van Alphen et

al., 2021, 2013) and the fact that deep sleep stages are most strongly and immediately affected by

sleep deprivation in mammals, we turned our attention to how the architecture of sleep might differ

between focally deprived flies and their yoked controls across the three days following deprivation.

Though the number of sleep bouts was not significantly different between focal and yoked flies

across any of the three post-deprivation cycles (Supplementary Figure 4A), there was a significant

increase in bout duration in the focal flies compared to yoked controls across the first two days

following deprivation (Supplementary Figure 4B). These two groups of flies displayed no

differences in these metrics during baseline sleep before deprivation (Supplementary Figure 4C).

This result suggested that focal flies, which had higher sleep pressures than yoked controls,

displayed increased sleep bout durations.

The magnitude of homeostatic sleep rebound in the fly is masked by short-bout sleep.

Recent work in our lab revealed that long-bout sleep (~30 minutes or more of inactivity) is a better

reflection of sleep homeostat action than shorter bouts of inactivity (Abhilash and Shafer, 2022).

Furthermore, such long-bout sleep has been shown by others to reflect a deep sleep stage that is

physiologically and metabolically distinct from shorter bouts of sleep (Stahl et al., 2017; van

Alphen et al., 2021, 2013). We therefore hypothesized that the extent of sleep rebound and payback

of lost sleep might be more readily apparent if we focused our analysis on long-bout sleep alone,

eliminating the contribution of shorter, presumably shallower, bouts of sleep.

We reanalyzed the 24-h sleep deprivation experiment reported in Figure 3, iteratively varying the

inactivity duration criterion for sleep from one to 30-min, a range, which includes "active sleep"

(defined as any bout of inactivity lasting one to five minutes), to ask how varying the sleep

definition might shape the apparent homeostatic recovery of lost sleep. We found that inactivity

criteria of 25-min or more resulted in the largest payback of lost sleep when examined on the third

day of recovery from deprivation (Figure 4B). This suggested that longer bouts of sleep are most

strongly influenced by homeostatic sleep regulation and that sleep bouts lasting one to 24 minutes

are characterized by substantially lower rebound following deprivation. We found no evidence of

post-deprivation increases in brief, "active sleep" (periods of inactivity lasting between one and

five minutes). This was not surprising given our finding that 220-s inactivity triggers produce

increases in such active sleep bouts during deprivation (Supplemental Figure 1). When we

employed an inactivity criterion of 25-min, a duration that likely reflects a relatively deep sleep

state (Stahl et al., 2017) sleep recovery approached ~50% of lost sleep by the third day of recovery,

a much higher proportion than was apparent using currently employed unitary definition of sleep

as bouts of inactivity lasting 5-min or more.

Based on this result, we, re-examined our 24-h sleep deprivation data using 25 minutes of inactivity

as our criterion for sleep, thereby focusing on the longer bouts of sleep that likely correspond to a

deep sleep state and omitting shorter, presumably shallower, bouts of inactivity from our analysis.

Five-day long bout sleep time-series for focal, yoked, and undisturbed flies are shown in Figure

4A. Unperturbed controls displayed two daily peaks of long bout sleep, with approximately twice

as much deep sleep taking place at night, as previously described (Hendricks et al., 2000; Stahl et

al., 2017). Focal flies were, as expected from 220-s inactivity triggers, completely deprived of

sleep bouts consisting of 25-min or more. Yoked flies also, as expected, displayed significantly

reduced levels of such long bout sleep compared to unperturbed controls, but such sleep was in

fact attained by these controls, though at levels significantly reduced from those displayed by

unperturbed controls (Figure 4A).

When total long bout sleep across three successive cycles was quantified, focal flies displayed

significant increases in such sleep when compared only to yoked controls on days two and three

of recovery (Figure 4C and D). The absence of a difference with yoked flies on day one is likely a

product of the substantial reduction in long-bout sleep that yoked flies experienced during the

deprivation period. Nevertheless, focal and yoked flies revealed differential long-bout sleep

rebounds across three days of recovery. Remarkably, this rebound was not seen when focal flies

were compared to unperturbed controls (Figure 4C). Once again, this was likely due to the sleep-

pressure-independent effects of mechanical disturbance. To confirm that frequent 25-min bouts of

inactivity were not an abnormal occurrence induced by the Ethoscope (i.e., that this state was not

induced by the rotations delivered by the Ethoscopes), we confirmed that such long bouts of sleep

are frequently observed in unperturbed control flies. Indeed, all unperturbed control flies routinely

displayed 25-min or longer bouts of inactivity (Supplementary Figure 5A). The sleep architecture

of focal flies after sleep deprivation also changed significantly compared to yoked controls where

instances of longer and consolidated sleep bouts were significantly increased in the focal flies,

although long bout sleep durations were not significantly different (Supplementary Figure 5B, C).

These results suggested that longer bouts of sleep are a more sensitive reflection of

homeostatically-controlled sleep in flies. This conclusion is supported by the significant recovery

of lost long bout sleep that continues across all three days of the observed recovery period (Figure

4E). Thus, when the analysis focused on long and presumably deep sleep states, a much larger

percentage such sleep is recovered compared to a consideration of all epochs of inactivity lasting

longer than one minute. At least half of all sleep-deprived flies paid back more than 50% of the

sleep they lost during the 24-h of deprivation, and over 25 percent of flies overshot their sleep-

debt, thereby displaying a defining hallmark of homeostatic control (Figure 4E). These results lend

further support for the idea that long bouts of inactivity represent a sleep state that is distinct from

shorter bout sleep (Stahl et al., 2017; Tainton-Heap et al., 2021; van Alphen et al., 2013).

Furthermore, robust homeostatic recovery of long-bout suggests that our initial failure to detect

rebound when 220-s inactivity triggers were used for deprivation (see Figure 1) was not simply

due to increases in "active sleep" preventing the build-up of homeostatic sleep drive

(Supplementary Figure 1), as long bout sleep displays clear homeostatic rebound even when 220-

s triggers allow for substantial levels of "active sleep" during deprivation.

Long Bout Sleep is Under Potent Homeostatic Control.

In mammals, both REM and NREM stages of sleep are controlled homeostatically, with stage-

specific deprivation leading to stage-specific rebound (Borbély, 1982; Borbély and Neuhaus, 1979;

Endo et al., 1998, 1997). If long-bout sleep indeed reflects a distinct sleep stage in *Drosophila*,

selectively depriving flies of it should produce a homeostatic rebound in long-bout sleep, despite

the presence of abundant short-bout sleep. To test this prediction, we set the Ethoscope's

inactivity-dependent trigger to rotate tubes only when focal flies had been immobile for 22-min as

a means to prevent flies from attaining 25-min of uninterrupted inactivity. When using the standard

inactivity duration criterion for sleep (5-min or more), this long trigger latency supported what

appeared to be largely normal sleep rhythms in both focal and yoked flies (Figure 5A).

Unsurprisingly, normal sleep patterns were also observed in unperturbed flies (data not shown).

Employing a 25-min inactivity duration criterion to remove short bout sleep from our analysis

revealed that 22-min triggers effectively deprived flies of long-duration sleep while permitting

yoked flies to achieve appreciable amounts of such sleep during focal deprivation (Figure 5B).

Unperturbed control flies once again displayed the bimodal long-bout sleep rhythms described

above, with substantially more nighttime than daytime deep sleep (Supplementary Figure 6). When

behavior was analyzed using the standard five-min inactivity criterion, which would include short-

bout sleep, focal flies displayed no significant increases in total sleep when compared to yoked

controls (Figure 5C). In contrast, when short-bout sleep was eliminated from the analysis through

the use of a 25-min inactivity criterion, focal flies displayed significantly higher levels of long-

bout sleep across all three days following deprivation (Figure 5D). Remarkably, the relative levels

of post-deprivation sleep displayed by the focal flies appeared to increase over the course of the

three days of observation, suggesting that homeostatic responses to long-bout sleep deprivation

may persist for an extended period of time (Figure 5D). This rebound produced a significant

recovery of lost sleep (Figure 5E), one that may have continued to build over additional sleep/wake

cycles. No such patterns were observed when focal flies were compared to unperturbed controls

(data not shown), thereby once again highlighting the importance of yoked controls.

Long-bout-specific deprivation also produced significant effects on sleep architecture. Focal flies

were characterized by an increased number of bouts that were longer than 25-mins compared to

yoked controls (Supplementary Figure 7B), although the durations of these longer sleep bouts did

not increase (Supplementary Figure 7D). No differences in the number of long-bout sleep between

focal and yoked flies were observed prior to the deprivation cycle (Supplementary Figure 7A). Our

results suggest that long-bout sleep, previously shown by others to represent a relatively deep sleep

state, is under strong homeostatic control and highlights the importance of controlling for the sleep-

independent effects of mechanical sleep deprivation on behavior.

Yoked controls are necessary to differentiate effects of sleep deprivation from effects driven

by mechanical perturbation.

A major goal of sleep science is to identify the molecular, cellular, and physiological processes

mediating the rise and fall of sleep pressure and how these changes operate in the brain to promote

sleep or wakefulness. In the fly, previous work by others has described cellular (Blum et al., 2021;

Donlea et al., 2014; Liu et al., 2016) and physiological (Pimentel et al., 2016; Tainton-Heap et al.,

2021) correlates of sleep pressure, based on the effects of mechanical sleep deprivation. Given our

behavioral results indicating significant sleep-independent effects of mechanical deprivation that

appear to mask homeostatic sleep responses, we sought to examine the potential utility of yoking

for differentiating sleep-pressure driven changes from the sleep-independent effects of mechanical

perturbation. Toward this end we conducted a simple molecular screen in the fly brain using

matrix-assisted laser desorption/ionization – time-of-flight (MALDI-TOF) mass spectrometry

(Veerasammy et al., 2020). This method supports the mass spectrometric analysis of fly head

sections (Figure 6A; Salisbury et al., 2013) and consists of three major steps: the crystallization of

a matrix and analytes within the tissue section, the laser ionization of analytes within this matrix,

and time-of-flight mass spectrometry (MS) to analyze the molecules present (Veerasammy et al.,

2020). This method provides sufficient spatial resolution to detect the presence of molecules

defined by their weight over charge ratios (m/z), specifically within the central brain (Figure 6B).

Using our behavioral sleep deprivation approach (Figure 2E-G), we examined the molecular

responses within the central brains of focally deprived flies, paired yoked controls, and

unperturbed flies following 24-h of sleep deprivation using 220-s inactivity triggers.

Using 2,5-dihydrobenzoic acid (DHB) as a positive-ion matrix, which is considered useful for

"universal analysis" and the detection of diverse molecular types (Snovida et al., 2006), we

sampled m/z value peaks of up to 1,300 and detected 188 distinct peaks (in the central brain regions

of *Drosophila* head sections. Each m/z ratio peak measurement was made specifically for the

central brain by registering the MALDI-TOF signal directly over the histological image of our

tissue sections (dashed square, Figure 6B). When we compared the heights of these peaks between

focal, sleep-deprived flies and unperturbed controls, we found 127 peaks that displayed

statistically significantly higher amplitudes in sections from focal flies compared to those of

unperturbed flies (Figure 6C). Thus, when compared only to unperturbed flies, sleep deprivation

in focal flies was accompanied by increases in a large proportion of the molecules examined.

However, it is highly likely that the mechanical stimulation associated with sleep deprivation

causes many of these molecular changes within the brain and such increases might not track sleep

pressure. We therefore asked the extent to which controlling for mechanical insult using yoked

controls might allow us to differentiate increases explained by sleep pressure from those that were

simply produced by mechanical stimulation.

If a molecule's abundance is increased simply as consequence of mechanical insult during sleep

deprivation, we would expect head sections from both focal and yoked flies to display similar

concentrations of that molecule that are elevated compared to unperturbed controls (Figure 6D,

left-top). In contrast, if a molecule's abundance is increased specifically as consequence of sleep

pressure, we predict that it will be present at relatively low levels in sections from unperturbed

controls, significantly higher levels in sections from focal flies, and with intermediate levels in

yoked controls that are statistically distinguishable from the other two conditions (Figure 6D, left-

bottom). Example m/z peaks that show these characteristic patterns in the central brain are shown

in Figure 6D (right; m/z values of 104.1 and 553). When yoked controls are included in the

analysis, approximately half of the discernable m/z value peaks displayed statistically

indistinguishable levels in the central brain across the undisturbed, yoked, and focal samples

(Figure 6E, "N.S.", 45%). Seventy-three of the m/z peaks displayed the relative abundances

expected of molecules tracking mechanical perturbation (38.8%), whereas only eight displayed the

relative abundances expected of molecules tracking sleep pressure (~4%, Figure 6E). Thus, the

standard method of comparing sleep-deprived flies to unperturbed controls produced 127

candidate sleep-tracking molecules in two replicate experiments, whereas the inclusion of yoked

controls produced only eight candidates. Thus, many of molecular differences between sleep-

deprived flies and unperturbed controls were likely explained as responses to mechanical

stimulation rather than to the loss of sleep.

We replicated this molecular screen in an independent MALDI-TOF experiment. This run detected

slightly fewer m/z peaks (170 compared to 188 in the first replicate). Despite the markedly lower

sensitivity of this replicate, we found that two of the eight molecules identified in the first screen

as tracking sleep pressure, once again displayed the relative abundances expected of a sleep

pressure tracking molecule. One of these had an m/z peak at ~553, and the other at 177.1 m/z in

the DHB matrix. The 177.1 m/z peak has previously been shown to correspond to serotonin (Cao

et al., 2019; Moriarty et al., 2011) and fits the molecular weight range of serotonin (176.2) in a

positive ion matrix like DHB, which adds a cation with a single positive charge to the m/z peak,

which for serotonin would result in a peak at ~177 m/z. Furthermore, we confirmed that this m/z

value reflected serotonin by spotting serotonin on slides imaged by MALDI-TOF MS and

confirmed strong peaks at m/z = 177 for regions of interest corresponding to these spots

(Supplementary Figure 8). Thus, serotonin would appear to be a strong candidate for a molecule

whose abundance reflects sleep pressure in the fly's central brain (see Discussion), that follows

the expected pattern across unperturbed, yoked and focal flies (Figure 6F). An unknown molecule

with an m/z value of ~553 is likewise also a strong candidate for another such molecule.

Discussion

The impact of the sleep-independent effects of mechanical sleep deprivation on the measurement of homeostatic sleep rebound.

In order to study the sleep homeostat, one must prevent sleep through the imposition of prolonged wakefulness (Bentivoglio and Grassi-Zucconi, 1997; Henri, 1895; Patrick and Gilbert, 1896; Rechtschaffen et al., 1983). However, the very act of sleep disruption is expected to cause not just the build-up of sleep pressure, but also other potentially confounding responses to the stimulus used to prevent sleep. The specific confounds attending sleep deprivation will depend on the method, duration, and time-of-day of the sleep-depriving stimulus employed, and all forms of experimental sleep deprivation are expected to produce such confounds, including methods based on chemo-, thermo-, or opto-genetics. Any sleep deprivation method that produces increased locomotion will increase muscular activity, which will be attended by sleep-independent effects. Chronic mechanical agitation also produces stress (Meerlo et al., 2008; Sgoifo et al., 2006; Tartar et al., 2009), and even gentle handling of rodent subjects, a means of deprivation thought to cause little stress, has significant effects on feeding behavior following deprivation (Đukanović et al., 2022). Such confounding, sleep-independent effects can mask homeostatic sleep responses and must therefore be accounted for whenever possible. Here we have attempted to account for the sleep-independent effects of mechanical sleep deprivation in *Drosophila,* the most commonly used method in the field of fly sleep research.

To assess the degree to which the frequency of mechanical agitation affects subsequent sleep

rebound in *Drosophila*, we examined flies that were sleep deprived for 24-h with different

frequencies of mechanical perturbation. The less frequent agitation produced significantly less

sleep rebound, despite having produced similar amounts of sleep loss. Thus, a sleep-pressure

independent effect of mechanical perturbation appeared to be the primary determinant of post-

deprivation sleep. An alternative explanation for this result was that short bouts of inactivity may

have provided brief but restorative sleep that the longer perturbation frequencies were not effective

at depriving. Indeed, 220-s inactivity triggers produced increases in brief (between one and five

minute) periods of inactivity during deprivation (Supplementary Figure 1). Such "active sleep" is

proposed to be a distinct sleep state from the standard sleep traditionally measured in flies (Niki et

al., 2023; Tainton-Heap et al., 2021). However, based on our estimates of p(Doze), a metric of

sleep pressure (Supplementary Figure 3), which is persistently high for three cycles post sleep

deprivation, we do not think that this "active sleep" prevented homeostatic sleep drive, at least in

this context. Furthermore, based on a large body of work conducted on humans and other

mammals, we would not necessarily expect increases in one sleep state to prevent the homeostatic

regulation of another.

Though it is possible that increases in the amount of brief "active sleep" bouts may have reduced

the build-up of sleep pressure during deprivation experiments employing 220-s inactivity triggers,

our results strongly support the conclusion that mechanical stimulation produces sleep-

independent effects that can mask homeostatic sleep rebound. Six, 12-, and 24-h of nighttime sleep

deprivation using frequent (20-s) triggers failed to produce the expected relationship between the

amount of sleep lost and the magnitude of rebound during the recovery day (Figure 1J). The failure

of rebound to track the amount of sleep loss indicates that the comparison of sleep deprived flies

and unperturbed controls is not sufficiently sensitive to detect the differential sleep rebounds

expected from flies that have suffered significantly different levels of sleep loss (Figure 1).

A previous attempt to control for the sleep-independent effects of mechanical perturbation

delivered mechanical stimuli matched to sleep deprived flies, but delivered during periods of high

activity (i.e., wakefulness), thereby exposing the control flies to mechanical shaking while

resulting in negligible amounts of sleep loss (Harbison and Sehgal, 2009). However, mechanical

shaking only during times of wakefulness increased molecular signatures of stress, even when

given for relatively short periods (Harbison and Sehgal, 2009). To minimize the extent of

mechanical perturbation necessary to prevent sleep and to control for the sleep-independent effects

of such perturbation, we adopted the Ethoscope, a video based system that tracks single flies and

delivers a mechanical stimulus, the rapid rotation of the glass tube containing the fly, only when a

fly meets a user-defined inactivity duration criterion (Geissmann et al., 2017). Thus, rather than

agitating all the flies of the sleep-deprived condition at a given frequency throughout the entire

deprivation period, the Ethoscope prevents single flies from attaining specific durations of

inactivity, just before meeting a user defined duration. Previous work using this system has

revealed remarkably low levels of homeostatic rebound and no sleep deprivation-associated death

(Geissmann et al., 2019).

Though the Ethoscope produces sleep deprivation using fewer mechanical perturbations, it

nevertheless delivers significant mechanical perturbation whose sleep-independent effects might

mask homeostatic sleep responses (Figure 2A-D). For this reason, we introduced yoked controls

to the Ethoscope platform, allowing us for the first time to disentangle effects associated with sleep

pressure from those best explained by the mechanical insult delivered during sleep deprivation

(Figure 2E-G). When flies were prevented from attaining five-min bouts of inactivity - the standard

criteria for a sleeplike state in the fly – using 220-s inactivity triggers in Ethoscopes, they failed to

show significant homeostatic rebound when their sleep was compared to unperturbed controls.

Comparison of these deprived flies to their yoked controls, in contrast, revealed a significant, albeit

small, increase in sleep following deprivation (Figure 2F, G). Thus, yoking revealed homeostatic

rebound that was not apparent when deprived flies were compared to controls that had not

experienced mechanical stimulation. However, when compared to yoked controls, the homeostatic

response of sleep deprived flies was quite small, amounting to very little recovery of lost sleep

(Figure 3E).

Sleep as a non-unitary state in *Drosophila*: Long-bout sleep is a distinctly regulated sleep

stage.

In mammals, sleep is not a unitary state but rather consists of distinguishable sleep stages (Davis

et al., 1937; Loomis et al., 1937). For example, REM and NREM sleep represent distinct

physiological states that differ in their relationship to sleep pressure and the circadian system

(Borbély, 1982; Borbély and Neuhaus, 1979; Czeisler et al., 1980; Endo et al., 1997). Slow-wave

NREM sleep appears to be the mammalian sleep state most strongly controlled by the daily

homeostatic control of sleep: initial sleep cycles following prolonged wakefulness are rich in slow

wave sleep, and this form of sleep is also prevalent following sleep deprivation (Berger and

Oswald, 1962; Ursin, 1971). For the majority of studies using *Drosophila*, sleep has been treated

as unitary state, with all inactivity bouts of five minutes or more being treated as sleep. However,

a growing body of works suggests the existence of distinct sleep stages in the fly. For example, in

one of the very first papers describing sleep-like states in *Drosophila*, Hendricks et al., reported

differences in durations of sleep bouts consisting of epochs of inactivity lasting five minutes or

more (Hendricks et al., 2000). The longest bout lengths occurred at the beginning of the subjective

night under constant conditions between circadian time 13.5 and 16. More recently, quantitative

approaches using Hidden Markov models have suggested the existence of light and deep sleep

states in the fly (Wiggin et al., 2020; Xu et al., 2021).

Flies display both longer and deeper bouts of sleep at night (Hendricks et al., 2000; van Alphen et

al., 2013) and deeper sleep is associated with distinct patterns of global brain activity when

compared to brain activity during wakefulness or during more brief and shallow bouts sleep (Nitz

et al., 2002; Tainton-Heap et al., 2021). Longer sleep bouts are also attended by lower metabolic

rates, where bout lengths of 60-min were shown to be associated with the lowest rates (Stahl et al.,

2017), a phenomenon similar to that observed for deeper NREM sleep stages in humans (Berger

and Phillips, 1995). Finally, flies display a stereotyped pattern of proboscis movements that appear

to drive waste clearance (van Alphen et al., 2021), reminiscent of the changes in fluid dynamics

observed in the cerebral spinal fluid of sleeping mammals, that occur at times corresponding to

deep sleep stages (Fultz et al., 2019). The time of peak occurrence of these proboscis movements

were characterized by a significant reduction of brain wave activity compared to sleep at other

times (van Alphen et al., 2021).

Physiological and metabolic signatures of deep sleep in the fly both suggest that they correspond

to sleep bouts that are significantly longer than the standard sleep definition of five minutes of

inactivity and multiple lines of evidence suggest that deeper sleep states are likely reached upon

attaining 15-30 minutes of inactivity (Tainton-Heap et al., 2021; van Alphen et al., 2013). Finally,

recent work from our lab has suggested that bouts of inactivity of between 30 and 60 minute are

stronger reflections of the daily homeostatic control sleep than shorter bouts of sleep (Abhilash

and Shafer, 2022). How might such long bouts of inactivity relate to homoeostatic sleep rebound?

Homeostatic sleep rebound, as traditionally measured in flies, appears to differ from mammalian

sleep rebound, in that it is relatively modest in magnitude and appears to largely run to completion

during the first few hours following the offset of deprivation (Huber et al., 2004). In contrast,

mammalian rebound appears to play-out over several sleep-wake cycles (Coborn et al., 2019;

Mistlberger et al., 1983; Nakazawa et al., 1978; Rechtschaffen et al., 1999). Remarkably, when we

biased our analysis to bouts of inactivity lasting 25-min or more, clear and prolonged homeostatic

rebound of sleep was observed following deprivation. This rebound lasted several days (Figure 4)

and discharged a large proportion (>50%) of the long-bout sleep debt suffered during deprivation

(compare Figures 3E to 4E). This result further supports the notion that long durations of inactivity

more sensitively reflect the action of the sleep homeostat.

When we deprived flies only of inactivity bouts that were longer than 22 minutes (see methods), a

duration that we chose based on its effectiveness at reliably preventing 25-min epochs of inactivity,

we observed a homeostatic increase in long-bout sleep, while seeing no such homeostatic response

in short-bout sleep (Figure 5). Moreover, the homeostatic rebound of long bout-sleep was apparent

on the very first day of recovery (Figure 5B, D) as opposed to the rebound seen following

deprivation using shorter inactivity triggers (Figure 4A, D). This is probably due to the fact that

long bout sleep deprivation engaged mechanical triggers only 12-14 times on average over the 24

hours of deprivation, and thereby likely reduced significantly the sleep-independent effects of

mechanical perturbation.

Yoked controls are critical for the examination of molecular and physiological correlates of

sleep pressure.

A major goal of sleep science is to identify the molecular, cellular, and physiological mechanisms

underlying the build-up of sleep pressure and how such pressure promotes sleep. As has been

recognized since the founding of sleep science (Kleitman, 1923, 1939), inescapable confounds

attend sleep deprivation experiments. The increased activity caused by most forms of experimental

sleep deprivation is likely to have myriad behavioral, physiological, and metabolic effects.

Mechanical sleep deprivation through vortexing, the most commonly used method of sleep

deprivation in *Drosophila* research, is particularly problematic in this regard, as it appears to cause

significant physical stress (Harbison and Sehgal, 2009).

The sleep-independent effects of deprivation protocols represent a significant confound and pose

a major challenge to the identification of the molecular and physiological correlates of sleep

pressure. If a molecular or physiological response to deprivation truly represents an increase in

sleep pressure, it must be established that it is not simply a response to the stimuli used to prevent

sleep. A fundamental challenge for the sleep scientist working in the fly system therefore is to

differentiate between the effects of increased sleep pressure from the sleep-independent effects of

the stimuli used to prevent sleep. The use of yoked controls and longer duration inactivity triggers

in our study appears to be a useful means of accounting for the sleep-independent effects of

mechanical sleep disruption on behavior (Figures 2-5). We therefore wondered if this approach

would allow us to differentiate molecular changes that were specific to sleep pressure from changes

produced directly by the effects of mechanical perturbation.

Given the extent of disturbance necessary to keep flies awake during prolonged sleep deprivation,

we predicted that a large number of molecules whose abundance increases in response to sleep

deprivation would reflect responses to mechanical stimulation rather than sleep pressure driven

increases. Indeed, when we compared the brains of sleep-deprived files to unperturbed controls,

around 70% of all detected molecules showed a significant increase in abundance. However, when

we employed yoked controls as an additional control to identify which molecular weights were

higher in deprived flies compared to controls that had experienced identical mechanical

perturbation but significantly less sleep deprivation, we found that only approximately four percent

of detectable molecules reflected sleep pressure (Figure 6). Thus, many of the molecular changes

in response to sleep deprivation can likely be explained by the effects of mechanical disturbance

and not by sleep pressure. This result highlights the importance of yoked controls for confirming

that specific molecules, genes, and cells mediate sleep homeostasis rather than the behavioral or

physiological responses to prolonged mechanical stimulation. Such controls will be critical for the

vetting of physiological or cellular correlates of sleep pressure. More specifically, the limited

molecular screen we describe here supports the feasibility of using yoked controls to discover sleep

substances within the fly brain, molecules whose abundance rises and falls with sleep pressure that

are used by sleep control centers to exert homeostatic control of sleep (Borbély, 1986).

Serotonin is a candidate sleep substance in the *Drosophila* central brain.

The inclusion of yoked controls in our experiments revealed that very few of the molecular changes

produced by mechanical sleep deprivation tracked sleep loss. One molecule that reliably tracked

sleep pressure in independent experiments was serotonin, which, based on our results, can be

considered a strong candidate sleep substance in the fly's central brain. A previous study using

HPLC to examine serotonin levels in dissected brains failed to observe serotonin increases in sleep

deprived flies (Davla et al., 2020). However, the spatial resolution provided by MALDI-TOF

imaging suggests that sleep loss driven serotonin increases take place in the central brain and not

the optic lobes (e.g., Figure 6B, F). We therefore suspect that serotonin does not build-up

throughout the optic lobes and that whole-brain assays might therefore fail to detect the sleep loss

induced changes we describe here. The identification of serotonin as a candidate sleep substance

is notable, as convergent evidence in the fly strongly supports the conclusions that serotonin

signaling promotes sleep in *Drosophila*.

The identification of serotonin as a candidate sleep substance is notable, as convergent evidence

in the fly strongly supports the conclusion that serotonin signaling promotes sleep in *Drosophila*.

Pharmacological increases in serotonin and the excitation of serotonergic neurons produce

increases in baseline sleep in the fly (Haynes et al., 2015; Pooryasin and Fiala, 2015; Yuan et al.,

2006). Similarly, blocking synaptic transmission in single pairs of serotonergic neurons have been

reported to reduce baseline sleep (Alekseyenko et al., 2014). Genetic loss of serotonin synthesis

decreases baseline sleep, as does the loss of 5HT-R-1A and 5HT-R-2B serotonin receptors (Qian

et al., 2017; Yuan et al., 2006). Loss of serotonin and the serotonin receptor 5HT-R-2B also results

in a reduction in sleep rebound following deprivation (Qian et al., 2017). Furthermore, the genetic

loss of arylalkylamine N-acetyltransferase (ANAAT), which normally functions to inactivate

serotonin and dopamine, is accompanied by both increased homeostatic sleep rebound and brain

serotonin levels following sleep deprivation, consistent with serotonin acting as a sleep substance

during mechanical deprivation (Davla et al., 2020). Finally, loss of the serotonin transporter

dSERT, which normally functions to remove extracellular serotonin, is associated with significant

increases in sleep (Knapp et al., 2022). Thus, convergent evidence supports the conclusion that

serotonin promotes both basal sleep and homeostatic increases in sleep following deprivation.

According to the two-process model of sleep regulation, basal seep and homeostatic increases in

sleep are driven by "process S," which is thought to be mediated by somnogens, molecules that

rise and fall with sleep pressure (Borbély, 1986, 1982; Borbély and Achermann, 1999). Our

MALDI results, in conjunction with previous work establishing a sleep-promoting role for

serotonin in the fly, suggest that serotonin levels build in the central brain during sleep deprivation

and increase the pressure to sleep through actions of serotonin receptors. Remarkably, this very

model was proposed by Jouvet to reconcile conflicting results regarding serotonin's role in the

control of mammalian sleep (Jouvet, 1999). A significant body of work has established that

increasing serotonin in the mammalian brain produces sleep increases and that the loss of serotonin

results in insomnia. These observations were the basis of the serotonergic theory of sleep (Jouvet,

1999). Serotonergic neurons were subsequently shown to be active during wakefulness and this

was taken as evidence against the hypothesis that serotonin promotes sleep (Jouvet, 1999).

However, neurons producing a sleep substance would be expected to be active during wakefulness

and to thereby promote sleep pressure during prolonged bouts of wakefulness. Serotonin neurons

were hypothesized to act in this way by Jouvet in order to reconcile the wake-active nature of

serotonin neurons with the serotonergic theory of sleep (Jouvet, 1999). Our work strongly

implicates serotonergic signaling as a component of process S in the *Drosophila* brain and sets the

stage for future work to test the predictions of the serotonergic theory of sleep in the fly.

Conclusion

The work described here provides both new insights into sleep homeostasis in flies and new

methodologies that are likely to improve our ability to detect homeostatic sleep responses and

identify molecular correlates of sleep pressure in this species. Our work indicates that the

mechanical perturbation most commonly used to deprive flies of sleep produces both behavioral

and molecular changes that potentially mask the sleep-pressure-specific changes that underly

homeostatic sleep control. The results of our work also reveal that long-bouts of sleep, which have

been shown by several studies to represent a deep sleep state (Tainton-Heap et al., 2021; van

Alphen et al., 2013), are a more sensitive indicator of sleep pressure and rebound than the

previously used unitary definition of sleep as five minutes or more of inactivity. Remarkably, when

analysis is focused on longer (~30-min) bouts of inactivity, homeostatic control of sleep in the fly

appears much more similar to that seen in mammals, in that a larger proportion sleep debt is repaid

and is discharged over multiple sleep wake cycles. We predict that this larger magnitude of sleep

rebound will be of great utility to the field, increasing the dynamic range over which the effects of

molecular and physiological alterations can be examined and thereby increasing the likelihood of

discovering new molecular and cellular components of sleep homeostasis.

Methods

Fly stocks and Husbandry

Flies were reared on Corn Syrup/Soy media made by Archon Scientific (Durham, North Carolina) under a 12h:12h light:dark (LD) cycle at 25 °C and 60-70% humidity. Male wild-type *Canton-S* (*CS*; BDSC stock number: 64349) and w^{1118} (BDSC stock number: 3605) flies were used for the experiments.

Sleep assay and analysis

One- to three-day old male flies were collected in groups of 30 into Corn Syrup/Soy containing vials and were subsequently isolated under CO_2 anesthesia and loaded into glass tubes [70 mm × 5mm × 3 mm (length × external diameter × internal diameter)] containing 5% sucrose and 2% agar when they were five to seven days old. The loading was done at least 24-h prior to the beginning of behavioral experiments. Flies were allowed to adjust to the tubes for the remainder of the loading day and no data from this day was subject to analysis, which began using data from the next LD cycle. Locomotor activity was measured using two independent methodologies: *Drosophila* Activity Monitors (DAM, TriKinetics, Waltham, MA), and Ethoscopes built in our lab based on resources provided by the Gilestro Lab (Imperial College London) (Geissmann et al., 2017).

The DAM system employs a single infrared beam and a sensor for each tube and records beam crossing at the tube's mid-point as a measure of locomotor activity. Beam crossings were recorded every 20-s and from these data, inactivity bouts of five minutes or more were used as the standard definition of sleep (Hendricks et al., 2000; Shaw et al., 2000). For Ethoscope experiments,

continuous video tracking via infrared cameras were done for individual flies using Raspberry Pi

(https://github.com/raspberrypi/documentation/). Ethoscopes quantified maximal velocity in

mm/s in 10-s epochs. The maximal velocity data was analyzed using a custom R package (available

on request) to quantify sleep. Maximal velocities of <1 mm/s for any epoch were considered an

instance of immobility (Geissmann et al., 2017), and sleep was then computed using the standard

five-min inactivity criterion, with immobility lasting 300 seconds (30 epochs) or more considered

a bout of sleep. This criterion was subsequently adjusted to bias our analysis toward longer bouts

of sleep.

Homeostatic sleep rebound analysis: Total sleep displayed by sleep-deprived flies on days one,

two, and three following sleep deprivation (see below) were calculated across each 24-h day,

subtracted from the baseline sleep displayed on the day before deprivation. Normalization was

carried out using the following formula $[(\text{Focal}_{\text{PostDeprivationSleep}} - \text{Focal}_{\text{PreDeprivationSleep}}) -$

$(\text{Unperturbed}_{\text{PostDeprivationSleep}} - \text{Unperturbed}_{\text{PreDeprivationSleep}})]$. Sleep debt accrued in sleep deprived

flies was quantified by subtracting baseline 24-h sleep on the day before deprivation from total

sleep measured during the 24-h deprivation window.

Mechanical Sleep deprivation

24-h mechanical deprivation was carried under 12/12 L:D cycles with mechanical deprivation

starting at ZT00 (lights-on) following one day of baseline sleep measurement. 12-h deprivation

commenced at lights-off (ZT12) following one day of baseline sleep measurement. 6-h

deprivation commenced at ZT 18 (six hours after lights-off) following one day of baseline sleep

measurement. The temperature was kept constant at 25 °C with 60-70% humidity.

Mechanical sleep deprivation using DAM and mechanical shakers:

Vortexers (Fisherbrand™ Analog MultiTube Vortexer, Catalog # 02-215-450) were used as previously described to sleep deprive animals in DAM monitors (Kayser et al., 2015). DAM monitors housing 32 flies within glass tubes were shaken randomly for 2-s every 20-s, 120-s, or 220-s across the entire period of deprivation. The Multitube Vortexer was set at a shaking intensity of four.

Inactivity dependent sleep deprivation using Ethoscopes:

Using the rotational module, we sleep deprived flies with a 220-s inactivity threshold. DAM tubes were rotated at ~420 rpm for one second every time an individual fly was inactive for 220 seconds.

Yoked-controlled mechanical deprivation in the Ethoscope:

The Ethoscope rotational module(Geissmann et al., 2017) was used to develop a yoked-controlled mechanical deprivation platform. The Ethoscope has been used previously to rotate single tubes only when the animal has been inactive for specific durations of time (as opposed to shaking randomly and without regard to behavioral state as is the case of vortexing DAMs). We altered the closed loop feedback control (see below) in the Ethoscope to rotate two tubes at the same time based on the inactivity state of only one of the two flies. The fly whose behavior determined the rotation of the tubes is referred to here as the focal fly, whereas the paired fly that received identical time-matched rotations without regard to its behavioral state is called the yoked fly. This approach was adapted from one used for the sleep deprivation of rats (Rechtschaffen et al., 1983). Two inactivity dependent rotational triggers were used in this study: 220 seconds, which prevented the

focal fly from attaining the standard definition of sleep (five minutes of inactivity or more) and

1320 seconds, which deprived focal flies of inactivity durations of 22 minutes or more. Due to

logistical issues relating to video-based feedback, the inactivity duration was set to 22 minutes –

this was to make sure that no bouts of inactivity that are longer than 25 minutes are allowed to

occur.

Development of Ethoscope code to Support Yoked-Controls:

Yoking was implemented at the level of the Ethoscope's "TrackingUnit," which is where the

connection between the video-based motion tracker and the stimulator (i.e., tube rotator) is

located [self._stimulator.bind_tracker(self._tracker)]. The Ethoscope's program was modified so

that the stimulator controlling the tubes containing yoked controls was bound to the tracker of

their paired focal flies. This results in the stimulator of the yoked animals responding to the

behavior of the focal animals at the same time as the stimulators of the focal flies. The code

supporting yoked controls is available at https://github.com/shaliulab/shaferlab-ethoscope (a fork

of https://github.com/gilestrolab/ethoscope) with tag v1.0.999, which can be acquired via pip

with the command pip install shaferlab-ethoscope.

MALDI-TOF analysis

After 24-h of sleep deprivation using 220-s immobility triggers for focal flies with their yoked and

unperturbed controls, heads from all three conditions were embedded in Optimal Cutting

Temperature (Tissue-Tek, Product code: 4583, Sakura Finetek USA Inc.) compound, frozen with

liquid nitrogen, cryo-sectioned, and prepared and assayed by MALDI-TOF as previously described

(Veerasammy et al., 2020). 2,5-Dihydrobenzoc acid (Sigma-Aldrich, Cat #149357) was used as

our matrix, which supports the ionization of metabolites and peptides, and is considered a useful

matrix for "universal analysis" of a diverse variety of molecular types (Snovida et al., 2006)

MALDI mass spectra were acquired in a Bruker Autoflex Speed TOF Machine (Bruker, Germany)

and peaks were identified in SCiLS MALDI Imaging software (Bruker Daltonics, Germany) by a

manual peak walk across the spectrum of m/z values between 0 and 1300 for each discernible m/z

ratio for unperturbed, focal, and yoked flies.

Statistical analysis

In order to test the effect of trigger frequency and mechanical disturbance on the amount of

rebound sleep, we used a two-way fixed factor ANOVA. We used trigger frequency as one fixed

factor with three levels, i.e., 20-s, 120-s and 220-s. The second fixed factor was treatment with two

levels, i.e., mechanically disturbed fly and unperturbed controls. Whether the difference in total

sleep post-deprivation between unperturbed controls and sleep deprived flies was dependent on

trigger frequency was inferred based on the interaction effect between the two factors described

above. In all other analyses reported throughout the manuscript, we either used the Mann Whitney-

U (also referred to as the Wilcoxon's Rank Sum test) or the Kruskal-Wallis test, depending on the

number of groups being compared. In all cases involving more than two groups, multiple

comparisons were done using a Bonferroni correction. In case of the MALDI-TOF experiment,

owing to the large number of peaks detected, all the *p*-values from individual tests were treated to

a Benjamini-Hochberg correction to adjust for inflated False Discovery Rates. All statistical

analyses were done and figures made using custom R scripts. Specific tests, sample sizes and *p*-

values are reported in the figure legends.

Acknowledgements

This work was supported by grants from the National Institute of Neurological Disorders and

Stroke (R01NS077933 and R21NS131939) and start-up funds provided by the State of New York.

We are grateful to Drs. Giorgio Gilestro, Bill Joiner, and Maria de la Paz Fernández for useful

discussions of the work presented in this study. Dr. Ye He and Dr. Rinat Abzalimov, the Co-

Directors of MALDI-TOF MS Imaging Facility, and Kelly Veerasammy at the Advanced Science

Research Center, City University of New York provided technical support with MALDI-TOF

experiments and imaging. Finally, we thank, Matthew Ciolkowski, and Robert Veline for

providing useful feedback on the manuscript.

 Figure legends

Figure 1. Homeostatic sleep rebound is driven by the frequency of mechanical agitation when

the amount of sleep deprivation is held constant. A-B. 6-h of sleep deprivation was induced

with two different mechanical trigger frequencies using vortexers: 2-s randomized shaking every

20-s or 220-s (each producing similar levels of sleep deprivation). **C.** A significant increase in

sleep was observed for the 20-s trigger, but the 220-s trigger failed to show such increases when

compared with undisturbed controls (*Kruskal-Wallis critical value* = 40.43, *df* = 2, *p* = 1.66 x 10^{-}

^{9}, followed by Bonferroni corrections for pairwise comparisons). **D-E.** Time course of 12-h of

nighttime sleep deprivation using two trigger frequencies (20-s, 220-s) results in nighttime sleep

loss. **F**. Sleep rebound using for 20-s trigger produced significant increases compared to controls,

but we did not observe any significant changes when the 220-s trigger was used (*Kruskal-Wallis*

critical value = 24.59, *df* = 2, *p* = 4.5 x 10^{-6}, followed by Bonferroni corrections for pairwise

comparisons). **G-H.** We increased the vortexing induced sleep deprivation to 24-h using 20-s and

220-s trigger frequencies and calculated sleep rebound for 24 hours post deprivation window

represented in the time course. **I.** Once again, the 20-s trigger produced significant sleep rebound

compared to control flies whereas the 220-s trigger induced deprivation did not, even though both

trigger frequencies effectively deprived sleep (*Kruskal-Wallis critical value* = 23.63, *df* = 2, *p* =

7.41 x 10^{-6}, followed by Bonferroni corrections for pairwise comparisons). **J.** We compared the

amount of sleep rebound during the first recovery day caused by different durations of deprivation

using 20-s (left) and 220-s (right) inactivity triggers and found significant rebound only for the 20-

s trigger. However, despite producing significantly different amounts of sleep loss, we observed

no increases in rebound between associated with increased duration of deprivation. Surprisingly,

the 6-h deprivation condition, which produced the least amount of sleep loss displayed

significantly higher rebound sleep compared to the 12-h, and 24-h deprivation conditions (20-s

trigger: *Kruskal-Wallis critical value* = 13.71, *df* = 2, *p* = 0.0011, followed by Bonferroni

corrections for pairwise comparisons; 220-s trigger: *Kruskal-Wallis critical value* = 1.96, *df* = 2, *p*

= 0.37, followed by Bonferroni corrections for pairwise comparisons). **K.** We included an

intermediate trigger frequency of 120-s for sleep deprivation and found that like the 220-s trigger,

it failed to produce statistically significant increases in sleep for 24 hours post-deprivation analysis

(ANOVA $F_{2,250}$ = 3.93, *p* = 0.02). **L.** Sleep levels observed during the 24-h following deprivation

reveal that flies recover very little of total sleep lost. Previously measured *Drosophila* (Huber et

al., 2004) and Human sleep (Stroemel-Scheder and Lautenbacher, 2021) recovery amounts are

indicated by dashed lines. 6-H SD Control n = 44, Sleep-Deprived$_{20 sec}$ n = 62, Sleep-Deprived$_{220}$

$_{sec}$ n = 57; 12-H SD Control n = 32, Sleep-Deprived$_{20 sec}$ n = 64, Sleep-Deprived$_{220 sec}$ n = 64; 24-

H SD Control n = 64, Sleep-Deprived$_{20 sec}$ n = 32, Sleep-Deprived$_{220 sec}$ n = 64, Sleep-Deprived$_{120}$

$_{sec}$ n = 64; * < 0.05, and N.S. (Not Significant).

Figure 2. Controlling for mechanical stimulation is necessary to observe 24-h sleep rebound

when low frequency mechanical triggers are used for deprivation. A. Inactivity dependent

sleep deprivation in Ethoscopes with 220-s inactivity triggers fails to generate post deprivation

sleep rebound in wild-type (*Canton-S*) flies when compared with undisturbed controls. **B.** Focal

flies do not show differences in total sleep compared to undisturbed flies across the baseline day

(Wilcoxon's *W* = 2523, *p* = 0.54) or across the first post deprivation day (Wilcoxon's *W* = 2340,

p = 0.87). **C.** *w^{1118}* flies also failed to display post-deprivation sleep rebound compared to

undisturbed controls. **D.** Comparison of total sleep on the day before and after sleep deprivation

for w^{1118} flies. There were no significant differences in sleep amount on the baseline day (Baseline sleep: Wilcoxon's $W = 351$, $p = 0.07$) or on the post deprivation day (Post deprivation sleep: Wilcoxon's $W = 519$, $p = 0.71$). **E.** The Ethoscope platform was modified to include "yoked-controls" where inactivity dependent sleep depriving triggers for experimental (focal) flies are matched to their yoked controls, thereby generating paired flies with identical mechanical experience but significantly different amounts of sleep loss. **F.** Sleep and mechanical perturbation in an example of a focal/yoked pair of flies: Sleep episodes for one baseline day and the day of deprivation are shown. Sleep episodes are indicated by red (focal) and blue (yoked) hash marks. Black hashmarks represent tube rotations triggered by the focal fly (Left panel). **G.** Averaged sleep profiles for ten focal-yoked paired flies on baseline, deprivation, and post-deprivation days shows how yoked flies (blue) were able to sleep during the deprivation cycle, whereas focal (red) flies lost all sleep. **H.** Average traces of focal and yoked flies 24-h before, during and after sleep deprivation for wild-type *Canton-S* flies. **I.** When compared to yoked flies that have undergone same levels of mechanical perturbation, focal flies increased sleep after 24-h deprivation (Wilcoxon's $W = 4074.5$, $p = 0.01$) while not showing any baseline differences on the day before deprivation (Wilcoxon's $W = 3829$, $p = 0.12$). **J.** To confirm reproducibility of yoking we used the w^{1118} line, a common genetic background, to replicate the sleep rebound finding observed for *Canton-S*. **K.** Compared to yoked flies that have undergone same levels of mechanical perturbation, focal flies show increased sleep following 24-h deprivation (Post deprivation sleep: Wilcoxon's $W = 277$, $p = 0.037$) while not showing any baseline differences on the deay before deprivation (Baseline sleep: Wilcoxon's $W = 176$, $p = 0.53$). For *Canton-S*, $n = 82$ each for focal and yoked categories, and $n = 58$ for unperturbed controls. For w^{1118}, $n = 20$ for focal and yoked categories, and $n = 49$ for unperturbed controls * < 0.05, and N.S. (Not Significant).

1025

Figure 3. Sleep pressure is discharged as cumulative sleep recovery across multiple cycles.

A. Average sleep time-courses of baseline, deprivation (220-s inactivity dependent trigger), and 3

post deprivation days in focal and undisturbed flies, and **B.** focal-yoked pairs for the same days

shown in A. **C.** Focal flies do not show significant differences in cumulative sleep gain across 3

post-deprivation days when compared to undisturbed controls (Post1: Wilcoxon's $W = 2323.5$, p

$= 0.81$; Post 2: Wilcoxon's $W = 2351$, $p = 0.91$; Post 3: Wilcoxon's $W = 2460.5$, $p = 0.72$). **D**.

When compared to paired yoked controls, focal animals showed significant increases in

cumulative sleep across two cycles (Post1: Wilcoxon's $W = 4088$, $p = 0.01$; Post 2: Wilcoxon's W

$= 4009$, $p = 0.03$). In the third post-deprivation day, even though the focal flies showed a tendency

to have higher cumulative sleep gain, this was not statistically significant (Wilcoxon's $W = 3932$,

$p = 0.06$). All comparisons are made using the Wilcoxon rank sum test, and $n = 82$ each for focal

and yoked categories. * < 0.05, and N.S. (Not Significant). **E.** Compared to total sleep lost, focal

flies do not pay back a significant amount of lost sleep, perhaps owing to normalization with

unperturbed controls (see methods). $n = 82$ each for focal and yoked categories, and n $= 58$ for

unperturbed controls * < 0.05, and N.S. (Not Significant).

1041

Figure 4. Examination of long-bout sleep reveals a multi-cycle sleep rebound that discharges

a significant proportion of lost sleep. A. Averaged sleep time-series of long-bout sleep (25-min

inactivity criterion) in baseline, deprivation (220-s inactivity dependent trigger), and 3 post-

deprivation windows in focal, yoked, and undisturbed flies. **B.** Sleep recovery as a function of

sleep definition (minutes of inactivity) visualized as a heatmap. When sleep is defined as

progressively longer bouts of inactivity, deprived flies paid back a majority of their sleep debt

cumulatively over three days as the inactivity criterion approaches 25-min. When either the

standard sleep definition (5 min or more of inactivity) or active sleep criteria (1- to 5-min) are

used, little to no sleep debt is repaid. **C.** In three post-deprivation days, focal flies showed trends

of increased cumulative sleep recovery in 25-min sleep but were not significantly different from

undisturbed controls (Post1: Wilcoxon's $W = 2458$, $p = 0.73$; Post 2: Wilcoxon's $W = 2689$, $p =$

0.18; Post 3: Wilcoxon's $W = 2766$, $p = 0.10$). **D.** Accounting for mechanical stimulation resulted

in significant and substantial multi-cycle total sleep increases for focal flies. Interestingly, the first

cycle after deprivation did not show significant increases in long bout sleep in focal flies (Post1:

Wilcoxon's $W = 3847$, $p = 0.11$; Post 2: Wilcoxon's $W = 3987$, $p = 0.03$; Post 3: Wilcoxon's $W =$

4001, $p = 0.03$). **E.** Normalized sleep rebound for sleep bouts of 25-min or longer across three days

shows the discharge of sleep debt accrued during the 24 hours of deprivation. Median recovery

was more than 50 percent, while 25 percent of flies overshoot total sleep loss, thereby showing

hallmarks of homeostatic control. $n = 82$ each for focal and yoked categories, and $n = 58$ for

unperturbed controls * < 0.05, and N.S. (Not Significant).

1062

Figure 5. Deprivation of only long bout sleep is sufficient to produce homeostatic responses

and elicit multiday sleep rebound. A. 5-day somnogram using the standard inactivity criterion

for sleep (5-min or more) in focal-yoked flies during baseline, long-bout sleep deprivation (22-

min inactivity triggers), and three days of recovery. There were no discernable changes in the

average standard sleep measurements between experimental and control flies. **B**. 5-day

somnograms of long-bout sleep (inactivity durations of 25-min or more) during baseline,

deprivation, and three post deprivation days in focal and yoked flies. **C.** In the three post-

deprivation days there were no significant differences in standard sleep between focal and yoked

flies (Post1: Wilcoxon's $W = 269$, $p = 0.06$; Post 2: Wilcoxon's $W = 258$, $p = 0.12$; Post 3:

Wilcoxon's $W = 263$, $p = 0.09$). **D.** Long bout sleep displayed significant increases in the total

amount of sleep over three post-deprivation cycles when compared to yoked controls (Post1:

Wilcoxon's $W = 288$, $p = 0.01$; Post 2: Wilcoxon's $W = 284$, $p = 0.02$; Post 3: Wilcoxon's $W =$

290, $p = 0.01$). **E.** Long bout sleep rebound showed over 25 percent of flies paying back total sleep

debt and overshooting in three post SD cycles. Focal sleep debt reflects the total sleep lost during

deprivation, which was normalized to sleep amounts in undisturbed controls. $n = 20$ each for focal

and yoked categories * < 0.05, and N.S. (Not Significant).

1079

Figure 6. Identifying sleep substances using MALDI-TOF. A. Representative hematoxylin-

eosin stained cryosection of a fly head section. **B**. Distribution of two representative molecules

(104.1 m/z and 175.1 m/z) within a single head section. Note the near zero relative intensity outside

the brain region. Images are overlays of the high intensity signals on the cryosection used to detect

the signal and highlights brain specific distribution of these molecules. The white dotted box

represents the area, corresponding to the central brain, from which signal intensity was quantified.

C. Post-deprivation increases in the relative abundance of molecules in focal (deprived) flies when

mechanical perturbations are not accounted for yields 127 positive hits (out of 188 molecules –

68%) in first replicate. We also found 59 peaks that were not significantly different between focal

and undisturbed flies, and 2 peaks that were significantly different, but lower in the focal than in

undisturbed flies, which we termed as "Others". **D**. Search patterns of molecules tracking

mechanical perturbation signals (top) and sleep pressure specific signals (bottom). Example peaks

(m/z 104.1 and m/z 553) showing each of the two categories. Values plotted are mean±SEM. **E.**

Pie chart representing the number molecules in various categories when yoking is employed. The

number of molecules reflecting sleep pressure was eight (4%) for the first replicate. We identified

an additional category of molecules that may track sleep and refer to it as "Sleep Absence".

Molecules that track the complete absence of sleep (as in the case of focal flies, but not in partial

sleep loss situations as in the yoked flies) are classified under this category, and we found nine

such candidates. These nine molecules had indistinguishable values between yoked and

undisturbed flies which were both significantly lower than focal flies. Focal brain sections $n = 6$;

Yoked brain sections $n = 3$; Unperturbed brain sections $n = 4$. **F.** One of the detected sleep-tracking

molecules was serotonin, detected at 177.1 m/z, which showed significant sleep pressure tracking

changes in both replicate runs (Replicate 1: *Kruskal Wallis critical value* = 9.32, *df* = 2, *p* = 0.009;

Replicate 2: *Kruskal Wallis critical value* = 11.56, *df* = 2, *p* = 0.003). Bars with the same letters

are not significantly different from each other.

1105

References

Abhilash L, Shafer OT. 2022. Circadian Regulation of Sleep in Drosophila: Insights from a Two-Process Model. doi:10.1101/2022.08.12.503775

Achermann P, Dijk DJ, Brunner DP, Borbély AA. 1993. A model of human sleep homeostasis based on EEG slow-wave activity: quantitative comparison of data and simulations. *Brain Res Bull* **31**:97–113. doi:10.1016/0361-9230(93)90016-5

Alekseyenko OV, Chan Y-B, Fernandez M de la P, Bülow T, Pankratz MJ, Kravitz EA. 2014. Single serotonergic neurons that modulate aggression in Drosophila. *Curr Biol* **24**:2700–2707. doi:10.1016/j.cub.2014.09.051

Beckwith EJ, Geissmann Q, French AS, Gilestro GF. 2017. Regulation of sleep homeostasis by sexual arousal. *eLife* **6**:e27445. doi:10.7554/eLife.27445

Bentivoglio M, Grassi-Zucconi G. 1997. The pioneering experimental studies on sleep deprivation. *Sleep* **20**:570–576. doi:10.1093/sleep/20.7.570

Berger RJ, Oswald I. 1962. Effects of sleep deprivation on behaviour, subsequent sleep, and dreaming. *J Ment Sci* **108**:457–465. doi:10.1192/bjp.108.455.457

Berger RJ, Phillips NH. 1995. Energy conservation and sleep. *Behav Brain Res* **69**:65–73. doi:10.1016/0166-4328(95)00002-b

Blum ID, Keleş MF, Baz E-S, Han E, Park K, Luu S, Issa H, Brown M, Ho MCW, Tabuchi M, Liu S, Wu MN. 2021. Astroglial Calcium Signaling Encodes Sleep Need in Drosophila. *Current Biology* **31**:150-162.e7. doi:10.1016/j.cub.2020.10.012

Borbély AA. 1986. Endogenous sleep-substances and sleep regulation. *J Neural Transm Suppl* **21**:243–254.

Borbély AA. 1982. A two process model of sleep regulation. *Hum Neurobiol* **1**:195–204.

Borbély AA, Achermann P. 1999. Sleep homeostasis and models of sleep regulation. *J Biol Rhythms* **14**:557–568. doi:10.1177/074873099129000894

Borbély AA, Daan S, Wirz-Justice A, Deboer T. 2016. The two-process model of sleep regulation: a reappraisal. *J Sleep Res* **25**:131–143. doi:10.1111/jsr.12371

Borbély AA, Neuhaus HU. 1979. Sleep-deprivation: Effects on sleep and EEG in the rat. *J Comp Physiol* **133**:71–87. doi:10.1007/BF00663111

Campbell SS, Tobler I. 1984. Animal sleep: a review of sleep duration across phylogeny. *Neurosci Biobehav Rev* **8**:269–300. doi:10.1016/0149-7634(84)90054-x

Cao Q, Wang Y, Chen B, Ma F, Hao L, Li G, Ouyang C, Li L. 2019. Visualization and Identification of Neurotransmitters in Crustacean Brain via Multifaceted Mass Spectrometric Approaches. *ACS Chem Neurosci* **10**:1222–1229. doi:10.1021/acschemneuro.8b00730

Cirelli C, Bushey D, Hill S, Huber R, Kreber R, Ganetzky B, Tononi G. 2005. Reduced sleep in Drosophila Shaker mutants. *Nature* **434**:1087–1092. doi:10.1038/nature03486

Coborn JE, Lessie RE, Sinton CM, Rance NE, Perez-Leighton CE, Teske JA. 2019. Noise-induced
sleep disruption increases weight gain and decreases energy metabolism in female rats. *Int*
J Obes (Lond) **43**:1759–1768. doi:10.1038/s41366-018-0293-9

Colten HR, Altevogt BM, Research I of M (US) C on SM and. 2006. Extent and Health
Consequences of Chronic Sleep Loss and Sleep Disorders, Sleep Disorders and Sleep
Deprivation: An Unmet Public Health Problem. National Academies Press (US).

Czeisler CA, Weitzman E d, Moore-Ede MC, Zimmerman JC, Knauer RS. 1980. Human sleep: its
duration and organization depend on its circadian phase. *Science* **210**:1264–1267.
doi:10.1126/science.7434029

Davis H, Davis PA, Loomis AL, Harvey EN, Hobart G. 1937. CHANGES IN HUMAN BRAIN
POTENTIALS DURING THE ONSET OF SLEEP. *Science* **86**:448–450.
doi:10.1126/science.86.2237.448

Davis KC, Raizen DM. 2017. A mechanism for sickness sleep: lessons from invertebrates. *J*
Physiol **595**:5415–5424. doi:10.1113/JP273009

Davla S, Artiushin G, Li Y, Chitsaz D, Li S, Sehgal A, van Meyel DJ. 2020. AANAT1 functions
in astrocytes to regulate sleep homeostasis. *Elife* **9**:e53994. doi:10.7554/eLife.53994

De Jesús-Olmo LA, Rodríguez N, Francia M, Alemán-Rios J, Pacheco-Agosto CJ, Ortega-Torres
J, Nieves R, Fuenzalida-Uribe N, Ghezzi A, Agosto JL. 2020. Pumilio Regulates Sleep
Homeostasis in Response to Chronic Sleep Deprivation in Drosophila melanogaster. *Front*
Neurosci **14**:319. doi:10.3389/fnins.2020.00319

Deboer T. 2018. Sleep homeostasis and the circadian clock: Do the circadian pacemaker and the
sleep homeostat influence each other's functioning? *Neurobiol Sleep Circadian Rhythms*
5:68–77. doi:10.1016/j.nbscr.2018.02.003

Dijk DJ, Brunner DP, Borbély AA. 1990. Time course of EEG power density during long sleep in
humans. *Am J Physiol* **258**:R650-661. doi:10.1152/ajpregu.1990.258.3.R650

Dijk D-J, von Schantz M. 2005. Timing and consolidation of human sleep, wakefulness, and
performance by a symphony of oscillators. *J Biol Rhythms* **20**:279–290.
doi:10.1177/0748730405278292

Donlea JM, Pimentel D, Miesenböck G. 2014. Neuronal machinery of sleep homeostasis in
Drosophila. *Neuron* **81**:860–872. doi:10.1016/j.neuron.2013.12.013

Dubowy C, Moravcevic K, Yue Z, Wan JY, Van Dongen HPA, Sehgal A. 2016. Genetic
Dissociation of Daily Sleep and Sleep Following Thermogenetic Sleep Deprivation in
Drosophila. *Sleep* **39**:1083–1095. doi:10.5665/sleep.5760

Đukanović N, La Spada F, Emmenegger Y, Niederhäuser G, Preitner F, Franken P. 2022.
Depriving Mice of Sleep also Deprives of Food. *Clocks & Sleep* **4**:37–51.
doi:10.3390/clockssleep4010006

Endo T, Roth C, Landolt H-P, Werth E, Aeschbach D, Achermann P, Borbély AA. 1998. Selective
REM sleep deprivation in humans: effects on sleep and sleep EEG. *American Journal of*
Physiology-Regulatory, Integrative and Comparative Physiology **274**:R1186–R1194.
doi:10.1152/ajpregu.1998.274.4.R1186

Endo T, Schwierin B, Borbély AA, Tobler I. 1997. Selective and total sleep deprivation: effect on
the sleep EEG in the rat. *Psychiatry Res* **66**:97–110. doi:10.1016/s0165-1781(96)03029-6

Everson CA, Gilliland MA, Kushida CA, Pilcher JJ, Fang VS, Refetoff S, Bergmann BM,
Rechtschaffen A. 1989. Sleep Deprivation in the Rat: IX. Recovery. *Sleep* **12**:60–67.
doi:10.1093/sleep/12.1.60

Freiberg AS. 2020. Why We Sleep: A Hypothesis for an Ultimate or Evolutionary Origin for Sleep
and Other Physiological Rhythms. *J Circadian Rhythms* **18**:2. doi:10.5334/jcr.189

Fultz NE, Bonmassar G, Setsompop K, Stickgold RA, Rosen BR, Polimeni JR, Lewis LD. 2019.
Coupled electrophysiological, hemodynamic, and cerebrospinal fluid oscillations in human
sleep. *Science* **366**:628–631. doi:10.1126/science.aax5440

Geissmann Q, Beckwith EJ, Gilestro GF. 2019. Most sleep does not serve a vital function:
Evidence from Drosophila melanogaster. *Sci Adv* **5**:eaau9253. doi:10.1126/sciadv.aau9253

Geissmann Q, Garcia Rodriguez L, Beckwith EJ, French AS, Jamasb AR, Gilestro GF. 2017.
Ethoscopes: An open platform for high-throughput ethomics. *PLoS Biol* **15**:e2003026.
doi:10.1371/journal.pbio.2003026

Harbison ST, Sehgal A. 2009. Energy Stores Are Not Altered by Long-Term Partial Sleep
Deprivation　in　Drosophila　melanogaster. *PLOS ONE* **4**:e6211.
doi:10.1371/journal.pone.0006211

Haynes PR, Christmann BL, Griffith LC. 2015. A single pair of neurons links sleep to memory
consolidation in Drosophila melanogaster. *eLife* **4**:e03868. doi:10.7554/eLife.03868

Hendricks JC, Finn SM, Panckeri KA, Chavkin J, Williams JA, Sehgal A, Pack AI. 2000. Rest in
Drosophila Is a Sleep-like State. *Neuron* **25**:129–138. doi:10.1016/S0896-6273(00)80877-
6

Henri V. 1895. Marie de Manacéine, Quelques observations sur l'insomnie absolue. *L'Année
psychologique* **2**:849–850.

Holding BC, Sundelin T, Schiller H, Åkerstedt T, Kecklund G, Axelsson J. 2020. Sleepiness, sleep
duration, and human social activity: An investigation into bidirectionality using
longitudinal time-use data. *Proceedings of the National Academy of Sciences* **117**:21209–
21217. doi:10.1073/pnas.2004535117

Huber R, Hill SL, Holladay C, Biesiadecki M, Tononi G, Cirelli C. 2004. Sleep homeostasis in
Drosophila melanogaster. *Sleep* **27**:628–639. doi:10.1093/sleep/27.4.628

Jouvet M. 1999. Sleep and serotonin: an unfinished story. *Neuropsychopharmacology* **21**:24S-
27S. doi:10.1016/S0893-133X(99)00009-3

Kanaya HJ, Park S, Kim J, Kusumi J, Krenenou S, Sawatari E, Sato A, Lee J, Bang H, Kobayakawa
Y, Lim C, Itoh TQ. 2020. A sleep-like state in Hydra unravels conserved sleep mechanisms
during the evolutionary development of the central nervous system. *Science Advances*
6:eabb9415. doi:10.1126/sciadv.abb9415

Kayser MS, Mainwaring B, Yue Z, Sehgal A. 2015. Sleep deprivation suppresses aggression in
Drosophila. *eLife* **4**:e07643. doi:10.7554/eLife.07643

Keene AC, Duboue ER. 2018. The origins and evolution of sleep. *Journal of Experimental Biology*
221:jeb159533. doi:10.1242/jeb.159533
Kleitman N. 1939. Sleep and wakefulness as alternating phases in the cycle of existence. Chicago:
Univ. of Chicago Press.
Kleitman N. 1923. Studies on the physiology of sleep. I. The effects of prolonged sleeplessness on
man. *American Journal of Physiology* **66**:67–92.
Klose MK, Shaw PJ. 2021. Sleep drive reconfigures wake-promoting clock circuitry to regulate
adaptive behavior. *PLoS Biol* **19**:e3001324. doi:10.1371/journal.pbio.3001324
Knapp EM, Kaiser A, Arnold RC, Sampson MM, Ruppert M, Xu L, Anderson MI, Bonanno SL,
Scholz H, Donlea JM, Krantz DE. 2022. Mutation of the Drosophila melanogaster
serotonin transporter dSERT impacts sleep, courtship, and feeding behaviors. *PLoS Genet*
18:e1010289. doi:10.1371/journal.pgen.1010289
Krause AJ, Simon EB, Mander BA, Greer SM, Saletin JM, Goldstein-Piekarski AN, Walker MP.
2017. The sleep-deprived human brain. *Nat Rev Neurosci* **18**:404–418.
doi:10.1038/nrn.2017.55
Lane JM, Qian J, Mignot E, Redline S, Scheer FAJL, Saxena R. 2022. Genetics of circadian
rhythms and sleep in human health and disease. *Nat Rev Genet* 1–17. doi:10.1038/s41576-
022-00519-z
Liu S, Liu Q, Tabuchi M, Wu MN. 2016. Sleep Drive Is Encoded by Neural Plastic Changes in a
Dedicated Circuit. *Cell* **165**:1347–1360. doi:10.1016/j.cell.2016.04.013
Loomis AL, Harvey EN, Hobart GA. 1937. Cerebral states during sleep, as studied by human brain
potentials. *Journal of Experimental Psychology* **21**:127–144. doi:10.1037/h0057431
Medic G, Wille M, Hemels ME. 2017. Short- and long-term health consequences of sleep
disruption. *Nat Sci Sleep* **9**:151–161. doi:10.2147/NSS.S134864
Meerlo P, Sgoifo A, Suchecki D. 2008. Restricted and disrupted sleep: effects on autonomic
function, neuroendocrine stress systems and stress responsivity. *Sleep Med Rev* **12**:197–
210. doi:10.1016/j.smrv.2007.07.007
Melnattur K, Morgan E, Duong V, Kalra A, Shaw PJ. 2020. The Sleep Nullifying Apparatus: A
Highly Efficient Method of Sleep Depriving Drosophila. *J Vis Exp* 10.3791/62105.
doi:10.3791/62105
Meyer N, Harvey AG, Lockley SW, Dijk D-J. 2022. Circadian rhythms and disorders of the timing
of sleep. *The Lancet* **400**:1061–1078. doi:10.1016/S0140-6736(22)00877-7
Mistlberger RE, Bergmann BM, Waldenar W, Rechtschaffen A. 1983. Recovery sleep following
sleep deprivation in intact and suprachiasmatic nuclei-lesioned rats. *Sleep* **6**:217–233.
doi:10.1093/sleep/6.3.217
Moriarty M, Lee A, O'Connell B, Kelleher A, Keeley H, Furey A. 2011. Development of an LC-
MS/MS method for the analysis of serotonin and related compounds in urine and the
identification of a potential biomarker for attention deficit hyperactivity/hyperkinetic
disorder. *Anal Bioanal Chem* **401**:2481–2493. doi:10.1007/s00216-011-5322-7

Nakazawa Y, Kotorii M, Ohshima M, Kotorii T, Hasuzawa H. 1978. Changes in sleep pattern after
sleep deprivation. *Folia Psychiatr Neurol Jpn* **32**:85–93. doi:10.1111/j.1440-
1819.1978.tb02782.x
Nall AH, Sehgal A. 2013. Small-molecule screen in adult Drosophila identifies VMAT as a
regulator of sleep. *J Neurosci* **33**:8534–8540. doi:10.1523/JNEUROSCI.0253-13.2013
Nath RD, Bedbrook CN, Abrams MJ, Basinger T, Bois JS, Prober DA, Sternberg PW, Gradinaru
V, Goentoro L. 2017. The jellyfish Cassiopea exhibits a sleep-like state. *Curr Biol* **27**:2984-
2990.e3. doi:10.1016/j.cub.2017.08.014
Ni JD, Gurav AS, Liu W, Ogunmowo TH, Hackbart H, Elsheikh A, Verdegaal AA, Montell C.
2019. Differential regulation of the Drosophila sleep homeostat by circadian and arousal
inputs. *eLife* **8**:e40487. doi:10.7554/eLife.40487
Niki A, A.l T-HL, Hang L, Eleni N, Qiongyi Z, Trent P, Philip B, J SP, Bruno van S. 2023.
Experimentally induced active and quiet sleep engage non-overlapping transcriptomes in
Drosophila. *eLife* **12**. doi:10.7554/eLife.88198
Nitz DA, van Swinderen B, Tononi G, Greenspan RJ. 2002. Electrophysiological correlates of rest
and activity in Drosophila melanogaster. *Curr Biol* **12**:1934–1940. doi:10.1016/s0960-
9822(02)01300-3
Nollet M, Wisden W, Franks NP. 2020. Sleep deprivation and stress: a reciprocal relationship.
Interface Focus **10**:20190092. doi:10.1098/rsfs.2019.0092
Oh Y, Yoon S-E, Zhang Q, Chae H-S, Daubnerová I, Shafer OT, Choe J, Kim Y-J. 2014. A
homeostatic sleep-stabilizing pathway in Drosophila composed of the sex peptide receptor
and its ligand, the myoinhibitory peptide. *PLoS Biol* **12**:e1001974.
doi:10.1371/journal.pbio.1001974
Patrick GTW, Gilbert JA. 1896. Studies from the psychological laboratory of the University of
Iowa: On the effects of loss of sleep. *Psychological Review* **3**:469–483.
doi:10.1037/h0075739
Pimentel D, Donlea JM, Talbot CB, Song SM, Thurston AJF, Miesenböck G. 2016. Operation of
a homeostatic sleep switch. *Nature* **536**:333–337. doi:10.1038/nature19055
Pooryasin A, Fiala A. 2015. Identified Serotonin-Releasing Neurons Induce Behavioral
Quiescence and Suppress Mating in Drosophila. *J Neurosci* **35**:12792–12812.
doi:10.1523/JNEUROSCI.1638-15.2015
Qian Y, Cao Y, Deng B, Yang G, Li J, Xu R, Zhang D, Huang J, Rao Y. 2017. Sleep homeostasis
regulated by 5HT2b receptor in a small subset of neurons in the dorsal fan-shaped body of
drosophila. *Elife* **6**:e26519. doi:10.7554/eLife.26519
Rechtschaffen A, Bergmann BM, Gilliland MA, Bauer K. 1999. Effects of method, duration, and
sleep stage on rebounds from sleep deprivation in the rat. *Sleep* **22**:11–31.
doi:10.1093/sleep/22.1.11
Rechtschaffen A, Gilliland MA, Bergmann BM, Winter JB. 1983. Physiological correlates of
prolonged sleep deprivation in rats. *Science* **221**:182–184. doi:10.1126/science.6857280

Salisbury JP, Boggio KJ, Hsu Y-WA, Quijada J, Sivachenko A, Gloeckner G, Kowalski PJ, Easterling ML, Rosbash M, Agar JN. 2013. A rapid MALDI-TOF mass spectrometry workflow for Drosophila melanogaster differential neuropeptidomics. *Mol Brain* **6**:60. doi:10.1186/1756-6606-6-60

Seidner G, Robinson JE, Wu M, Worden K, Masek P, Roberts SW, Keene AC, Joiner WJ. 2015. Identification of neurons with a privileged role in sleep homeostasis in Drosophila melanogaster. *Curr Biol* **25**:2928–2938. doi:10.1016/j.cub.2015.10.006

Sgoifo A, Buwalda B, Roos M, Costoli T, Merati G, Meerlo P. 2006. Effects of sleep deprivation on cardiac autonomic and pituitary-adrenocortical stress reactivity in rats. *Psychoneuroendocrinology* **31**:197–208. doi:10.1016/j.psyneuen.2005.06.009

Shafer OT, Keene AC. 2021. The Regulation of Drosophila Sleep. *Curr Biol* **31**:R38–R49. doi:10.1016/j.cub.2020.10.082

Shaw PJ, Cirelli C, Greenspan RJ, Tononi G. 2000. Correlates of Sleep and Waking in Drosophila melanogaster. *Science* **287**:1834–1837. doi:10.1126/science.287.5459.1834

Shaw PJ, Tononi G, Greenspan RJ, Robinson DF. 2002. Stress response genes protect against lethal effects of sleep deprivation in Drosophila. *Nature* **417**:287–291. doi:10.1038/417287a

Snovida SI, Chen VC, Perreault H. 2006. Use of a 2,5-dihydroxybenzoic acid/aniline MALDI matrix for improved detection and on-target Derivatization of glycans: A preliminary report. *Anal Chem* **78**:8561–8568. doi:10.1021/ac061375r

Stahl BA, Slocumb ME, Chaitin H, DiAngelo JR, Keene AC. 2017. Sleep-Dependent Modulation of Metabolic Rate in Drosophila. *Sleep* **40**. doi:10.1093/sleep/zsx084

Stroemel-Scheder C, Lautenbacher S. 2021. Assessment of effects of total sleep deprivation and subsequent recovery sleep: a methodological strategy feasible without sleep laboratory. *BMC Psychology* **9**:141. doi:10.1186/s40359-021-00641-3

Tainton-Heap LAL, Kirszenblat LC, Notaras ET, Grabowska MJ, Jeans R, Feng K, Shaw PJ, van Swinderen B. 2021. A Paradoxical Kind of Sleep in Drosophila melanogaster. *Curr Biol* **31**:578-590.e6. doi:10.1016/j.cub.2020.10.081

Takahashi Y, Ebihara S, Nakamura Y, Takahashi K. 1978. Temporal distributions of delta wave sleep and rem sleep during recovery sleep after 12-h forced wakefulness in dogs; similarity to human sleep. *Neurosci Lett* **10**:329–334. doi:10.1016/0304-3940(78)90248-3

Tartar JL, Ward CP, Cordeira JW, Legare SL, Blanchette AJ, McCarley RW, Strecker RE. 2009. Experimental sleep fragmentation and sleep deprivation in rats increases exploration in an open field test of anxiety while increasing plasma corticosterone levels. *Behav Brain Res* **197**:450–453. doi:10.1016/j.bbr.2008.08.035

Tobler I, Borbély AA, Groos G. 1983. The effect of sleep deprivation on sleep in rats with suprachiasmatic lesions. *Neuroscience Letters* **42**:49–54. doi:10.1016/0304-3940(83)90420-2

Ursin R. 1971. Differential effect of sleep deprivation on the two slow wave sleep stages in the cat. *Acta Physiol Scand* **83**:352–361. doi:10.1111/j.1748-1716.1971.tb05088.x

Vaccaro A, Kaplan Dor Y, Nambara K, Pollina EA, Lin C, Greenberg ME, Rogulja D. 2020. Sleep
Loss Can Cause Death through Accumulation of Reactive Oxygen Species in the Gut. *Cell*
181:1307-1328.e15. doi:10.1016/j.cell.2020.04.049

van Alphen B, Semenza ER, Yap M, van Swinderen B, Allada R. 2021. A deep sleep stage in
Drosophila with a functional role in waste clearance. *Sci Adv* **7**:eabc2999.
doi:10.1126/sciadv.abc2999

van Alphen B, Yap MHW, Kirszenblat L, Kottler B, van Swinderen B. 2013. A dynamic deep
sleep stage in Drosophila. *J Neurosci* **33**:6917–6927. doi:10.1523/JNEUROSCI.0061-
13.2013

Veerasammy K, Chen YX, Sauma S, Pruvost M, Dansu DK, Choetso T, Zhong T, Marechal D,
Casaccia P, Abzalimov R, He Y. 2020. Sample Preparation for Metabolic Profiling using
MALDI Mass Spectrometry Imaging. *J Vis Exp*. doi:10.3791/62008

Webb WB. 1974. Sleep as an adaptive response. *Percept Mot Skills* **38**:1023–1027.
doi:10.2466/pms.1974.38.3c.1023

Wiggin TD, Goodwin PR, Donelson NC, Liu C, Trinh K, Sanyal S, Griffith LC. 2020. Covert
sleep-related biological processes are revealed by probabilistic analysis in Drosophila.
Proc Natl Acad Sci U S A **117**:10024–10034. doi:10.1073/pnas.1917573117

Williams HL, Hammack JT, Daly RL, Dement WC, Lubin A. 1964. Responses to auditory
stimulation, sleep loss and the EEG stages of sleep. *Electroencephalography and Clinical*
Neurophysiology **16**:269–279. doi:10.1016/0013-4694(64)90109-9

Xu X, Yang W, Tian B, Sui X, Chi W, Rao Y, Tang C. 2021. Quantitative investigation reveals
distinct phases in Drosophila sleep. *Commun Biol* **4**:364. doi:10.1038/s42003-021-01883-
y

Yuan Q, Joiner WJ, Sehgal A. 2006. A sleep-promoting role for the Drosophila serotonin receptor
1A. *Curr Biol* **16**:1051–1062. doi:10.1016/j.cub.2006.04.032

1365

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

A

0.2 mm

B

Whole brain signal

104.1 m/z

Matrix background

Brain overlay

Central brain signal

175.1 m/z

Matrix background

Brain overlay

0%   Signal intensity   100%

C

Sleep pressure (68%)

Others (1%)

N.S. (31%)

D

Search patterns

Mech. perturbation signals

High

Low

Sleep pressure signals

High

Low

104.1 m/z

Pixel intensity (a.u.)

a

b   b

553 m/z

0.08

0.04

a   b

c

Unperturbed   Yoked   Focal

E

Mechanical perturbation (38.8%)

Sleep absence (4.8%)

Sleep pressure (4%)

Others (6.9%)

N.S. (45%)

F

Serotonin (177.1 m/z)

Pixel intensity (a.u.)

a

b

c

Replicate 1

a

b

c

Replicate 2

Unperturbed   Yoked   Focal