Transcriptional regulation and repressive condensates modulate a proliferative-invasive cellular switch *in vivo*

Authors: Taylor N. Medwig-Kinney[1†*], Brian A. Kinney[2†], Michael A. Q. Martinez[1], Callista Yee[3], Sydney S. Sirota[1‡], Angelina A. Mullarkey[1], Neha Somineni[1§], Justin Hippler[1,4#], Wan Zhang[1], Kang Shen[3], Christopher M. Hammell[2], Ariel M. Pani[5], and David Q. Matus[1¶*]

Affiliations:

1. Department of Biochemistry and Cell Biology, Stony Brook University, Stony Brook, NY 11794, United States.
2. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, United States.
3. Howard Hughes Medical Institute, Department of Biology, Stanford University, Stanford, CA 94305, United States.
4. Science and Technology Research Program, Smithtown High School East, St. James, NY 11780, United States.
5. Departments of Biology and Cell Biology, University of Virginia, Charlottesville, VA 22904, United States.

*For correspondence: tmkinney@unc.edu; david.matus@stonybrook.edu

Present address: †Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, United States; ‡Touro College of Osteopathic Medicine, Middletown, NY 10940, United States; §Integra LifeSciences, Princeton, NJ 08540, United States; #Northeastern University, Boston, MA 02115, United States; ¶Arcadia Science, Berkeley, CA 94704, United States.

Competing interest: David Q. Matus and Neha Somineni are paid employees of Arcadia Science and Integra LifeSciences, respectively.

Abstract

A growing body of evidence suggests that cell division and basement membrane invasion are mutually exclusive cellular behaviors. How cells switch between proliferative and invasive states is not well understood. Here, we investigated this dichotomy *in vivo* by examining two cell types that derive from equipotent progenitors, but exhibit distinct cell behaviors, in the developing *Caenorhabditis elegans* somatic gonad: the post-mitotic, invasive anchor cell and the neighboring proliferative, non-invasive ventral uterine (VU) cells. We report that the default invasive cellular state is suppressed in the VU cells through two distinct modes of regulation of the pro-invasive transcription factor NHR-67 (NR2E1/TLX). Levels of NHR-67 are important for discriminating between invasive and proliferative behavior, and *nhr-67* transcription is downregulated following post-translational degradation of its direct upstream regulator, HLH-2 (E/Daughterless) in VU cells. Residual NHR-67 protein is organized into discrete punctae in the nuclei of VU cells that are dynamic over the cell cycle and exhibit liquid-like properties. Strikingly, these NHR-67 punctae are not spatiotemporally associated with active transcription, but instead associate with homologs of the transcriptional co-repressor Groucho (UNC-37 and LSY-22), as well as the TCF/LEF homolog POP-1, likely mediated by a direct interaction between UNC-37 and the intrinsically disordered region of NHR-67. Further, perturbing UNC-37, LSY-22, or POP-1 results in ectopic invasive cells. We propose a model in which these proteins together form repressive condensates to suppress a default invasive state in non-invasive cells, which complements transcriptional regulation to add robustness to the proliferative-invasive cellular switch *in vivo*.

Introduction

Cellular proliferation and invasion are key aspects of development (reviewed in Medwig & Matus, 2017), and are also two of the defining hallmarks of cancer (reviewed in Hanahan and Weinberg, 2000). A growing body of evidence suggests that cell cycle progression and invasion through a basement membrane are mutually exclusive cellular

behaviors in both development and disease states (reviewed in Kohrman and Matus,

2017). Switching between invasive and proliferative phenotypes has been observed in

melanoma and recently in breast cancer (Hoek et al., 2008; Mondal et al., 2021), but

how these cell states are regulated in the context of development is not well

understood. To investigate how this dichotomy in cellular behavior is controlled *in vivo*,

we used *C. elegans*, leveraging its highly stereotypical development (Sulston and

Horvitz, 1977), as well as its genetic and optical tractability. During development of the

hermaphroditic reproductive system, the proximal granddaughters of the Z1 and Z4

somatic gonad progenitors, Z1.pp and Z4.aa, give rise to four cells that will adopt one of

two cellular fates: a proliferative ventral uterine (VU) cell or the terminally differentiated,

invasive anchor cell (AC) (Figure 1A) (Kimble and Hirsh, 1979). The distal cells of this

competency group, Z1.ppa and Z4.aap, quickly lose their bipotentiality and become VU

cells (Seydoux et al., 1990). In contrast, the proximal cells, Z1.ppp and Z4.aaa, undergo

a stochastic Notch-mediated cell fate decision, giving rise to another VU cell and the

post-mitotic AC (Figure 1A,B) (Greenwald et al., 1983; Seydoux and Greenwald, 1989).

Following fate specification, the AC undergoes invasive differentiation and breaches the

underlying basement membrane, connecting the uterus to the vulval epithelium to

facilitate egg-laying (Figure 1B) (Sherwood and Sternberg, 2003).

Our previous work has shown that AC invasion is dependent on G_0 cell cycle

arrest, which is coordinated by the pro-invasive transcription factor NHR-67

(NR2E1/TLX) (Figure 1–figure supplement 1A) (Matus et al., 2015). NHR-67 functions

within a gene regulatory network comprised of four conserved transcription factors

whose homologs have been implicated in several types of metastatic cancer (Liang and

Wang, 2020; Milde-Langosch, 2005; Nelson et al., 2021; Wang and Baker, 2015). We

previously reported that NHR-67 is regulated by a feed-forward loop formed by EGL-43

(Evi1) and HLH-2 (E/Daughterless), which functions largely in parallel to a cell cycle-

independent subcircuit controlled by FOS-1 (Fos) (Figure 1–figure supplement 1A)

(Medwig-Kinney et al., 2020). EGL-43, HLH-2, and NHR-67 are reiteratively used within

the Z lineage of the somatic gonad, in that they also function to independently regulate

LIN-12 (Notch) signaling during the initial AC/VU cell fate decision (Medwig-Kinney et

al., 2020). Despite its role in lateral inhibition between Z1.ppp and Z4.aaa, expression of

LIN-12 is not absolutely required for VU cell fate (Sallee et al., 2015). Cell cycle state

also cannot explain the difference between AC and VU cell fates, as arresting VU cells

in G_0 through ectopic expression of CKI-1 (p21/p27) does not make them invasive

(Smith et al., 2022). Thus, the mechanisms responsible for maintaining AC and VU

cellular identities following initial cell fate specification remain unclear.

Maintenance of differentiated cell identity is essential for ensuring tissue integrity

during development and homeostasis, and the inability to restrict phenotypic plasticity is

now being recognized as an integral part of cancer pathogenesis (Hanahan, 2022). *In*

vitro studies have identified several factors that safeguard differentiated cell identity

(reviewed in Brumbaugh et al., 2019). Despite its largely autonomous modality of

development, *C. elegans* has emerged as an ideal model system to study cell fate

maintenance *in vivo*. There have been several reports of cell fate transformations that

occur naturally, including two epithelial-to-neural transdifferentiation events (Jarriault et

al., 2008; Riva et al., 2022), or following fate challenges (reviewed in Rothman and

Jarriault, 2019). In such contexts, several epigenetic factors, including chromatin

remodelers and histone chaperones, have been identified for their roles in restricting

cell fate reprogramming (Hajduskova et al., 2019; Kagias et al., 2012; Kolundzic et al.,

2018; Patel et al., 2012; Rahe and Hobert, 2019; Zuryn et al., 2014). However, in some

cases, ectopic expression of a specific transcription factor is sufficient to overcome

these barriers, as was first shown through pioneering work in mouse embryonic

fibroblasts (Davis et al., 1987). Indeed, there are several examples in *C. elegans* where

ectopic expression of single lineage-specific transcription factors induces cell fate

transformations (Fukushige and Krause, 2005; Gilleard and McGhee, 2001; Horner et

al., 1998; Jin et al., 1994; Kiefer et al., 2007; Quintin et al., 2001; Richard et al., 2011;

Riddle et al., 2013; Tursun et al., 2011; Zhu et al., 1998). Moreover, *C. elegans* uterine

tissue may be particularly amenable to fate transformations, as ectopic expression of a

single GATA transcription factor, ELT-7, is sufficient to induce transorganogenesis of

the somatic gonad into gut by reprogramming the mesodermally-derived tissue into

endoderm (Riddle et al., 2016). Valuable insights have been made into how the function

of fate-specifying transcription factors can be tuned through means such as

autoregulation and dynamic heterodimerization (Leyva-Díaz and Hobert, 2019; Sallee et

al., 2017). We are just beginning to understand how an additional layer of control over

transcriptional regulators can be achieved through the formation of higher order

associations (Boija et al., 2018; Lim and Levine, 2021).

Here, in our endeavor to understand how AC and VU cellular fates are

maintained, we identified two mechanisms of NHR-67 regulation that together modulate

the invasive-proliferative switch in *C. elegans*. We found that high levels of NHR-67

expression are sufficient to drive invasive differentiation and, accordingly, *nhr-67* is

transcriptionally downregulated in the non-invasive VU cells following the post-

translational degradation of its direct upstream regulator, HLH-2. Additionally, we

observed that remaining NHR-67 protein forms discrete punctae in the nuclei of non-

invasive cells that exhibit liquid-like properties including dynamic assembly, fusion, and

dissolution over cell cycle as well as rapid recovery kinetics after photobleaching. These

NHR-67 punctae colocalize *in vivo* with UNC-37 and LSY-22, homologs of the

transcriptional co-repressor Groucho, as well as with POP-1 (TCF/LEF), which are

mediated through a direct interaction between UNC-37 and the intrinsically disordered

C-terminal region of NHR-67. Through functional perturbations, we demonstrate that

UNC-37, LSY-22, and POP-1 normally function in repressing the default invasive state

in VU cells. We propose a model in which the interaction between NHR-67 and Groucho

coordinates formation of repressive condensates that, combined with transcriptional

downregulation of *nhr-67*, suppress invasive differentiation. This work provides new

insights into how repressive nuclear condensates may coordinate cellular behaviors *in*

vivo and highlights how transcription factors can exhibit duality in functions depending

on cellular context.

<u>Results</u>

Levels of NHR-67 expression are important for distinguishing AC and VU cell

identity

Despite arising from initially equipotent cells, the differentiated AC and VU cells

exhibit very distinct cellular behaviors. The AC terminally differentiates to invade the

underlying basement membrane while the VU cells remain proliferative, undergoing

several rounds of division before terminally differentiating into the pi (π) and rho (ρ) cells

that function in uterine-vulval attachment. One potential explanation for this difference in

cell behavior is asymmetric expression of pro-invasive transcription factors. To

investigate this possibility, we examined endogenous expression levels of four

transcription factors that function in the gene regulatory network coordinating AC

invasion (EGL-43, FOS-1, HLH-2, and NHR-67) using previously generated GFP-

tagged alleles (Medwig-Kinney et al., 2020). While FOS-1 levels of expression in the AC

are nearly twice that of the VU cells (Figure 1–figure supplement 1B,C), FOS-1 has no

identified role in regulating cell cycle in the AC so we did not pursue this protein further

(Medwig-Kinney et al., 2021). EGL-43 also did not appear to be a promising candidate,

as it is expressed in both cell types at comparable levels, with VU cells exhibiting

approximately 89% of AC expression (Figure 1–figure supplement 1B,C). In contrast,

HLH-2 exhibits significant asymmetry in expression, as VU cells express merely 17% of

HLH-2 levels observed in the AC on average (Figure 1C,D). Previous studies have

shown that post-translational, dimerization-driven degradation of HLH-2 is responsible

for its downregulation in the VU cells (Benavidez et al., 2022; Karp and Greenwald,

2003; Sallee and Greenwald, 2015). Endogenously tagged NHR-67::GFP exhibits a

similar pattern of expression with over three-fold enrichment in the AC, consistent with

prior observations of transgenic reporters (Figure 1C,D) (Verghese et al., 2011). Given

the known role of NHR-67 in regulating cell cycle arrest and invasion, we hypothesized

that its differential expression between the AC and VU cells could potentially be

contributing to their distinct cellular behaviors.

To assess if NHR-67 plays a role in regulating uterine cell identities, we

manipulated its expression levels. We found that strong depletion of NHR-67 through

RNA interference (RNAi) treatment results in ACs adopting VU-like characteristics.

During AC/VU cell fate specification, LIN-12/Notch normally becomes restricted to the

VU cells while the Delta-like ligand LAG-2 (visualized by LAG-

2::P2A::H2B::mTurquoise2 (Medwig-Kinney et al., 2022)) accumulates in the AC

(Wilkinson et al., 1994). Here, we observe that NHR-67 deficient ACs not only

proliferated and failed to invade, as reported previously (Matus et al., 2015), but also

ectopically expressed membrane-localized Notch (visualized by LIN-12::mNeonGreen

(Pani et al., 2022)) (Figure 1E). Notably, NHR-67-deficient ACs expressed both LIN-12

and LAG-2, potentially indicating an intermediate state between AC and VU cell fate

(Figure 1E). Next, we ectopically expressed NHR-67 ubiquitously using a heat shock

inducible transgene (*hsp*::NHR-67::2x-BFP) (Medwig-Kinney et al., 2020). Intriguingly,

we observed that ectopic expression of NHR-67 following initial AC/VU specification

resulted in the presence of multiple invasive ACs at a low penetrance (approximately

5%, n > 50), denoted by ectopic expression of an AC marker (*cdh-*

3p::mCherry::moeABD) and expansion of the basement membrane gap (Figure 1F).

Since it has been previously demonstrated that proliferative ACs cannot invade (Matus

et al., 2015), we concluded that these invasive ectopic ACs most likely arose from fate

conversion of neighboring VU cells. Taken together, these pieces of evidence suggest

that high and low levels of NHR-67 expression correlate to properties of AC and VU cell

identities, respectively (Figure 1G).

NHR-67 is enriched in the AC through direct transcriptional regulation by HLH-2

Next, we investigated how NHR-67 expression levels become asymmetric

between the AC and VU cells. We and others have previously shown that HLH-2

positively regulates NHR-67 expression in the context of the AC (Figure 1–figure

supplement 1A) (Bodofsky et al., 2018; Medwig-Kinney et al., 2020). If this regulatory

interaction exists in the context of the VU cells as well, it could explain why the relative

expression pattern of NHR-67 in the AC and VU cells mirrors that of HLH-2. In support

of this hypothesis, we found that initial onset of HLH-2, which has shown to be

asymmetric in Z1.pp and Z4.aa (Attner et al., 2019), correlates to that of NHR-67 onset

(Figure 2–figure supplement 1A). To test whether HLH-2 degradation is responsible for

NHR-67 downregulation in the VU, we drove ectopic expression of HLH-2 using a

transgene under the control of a heat shock inducible promoter (*hsp*::HLH-2::2x-BFP)

(Medwig-Kinney et al., 2020). We observed that ectopic expression of HLH-2 resulted in

elevated NHR-67 expression in VU cells (43% increase; n > 30) (Figure 2A,B). To

control against potential dimerization-driven degradation of HLH-2 in the VU cells, which

the heat shock inducible transgene would still be susceptible to, we disrupted UBA-1, an

E1 ubiquitin-activating enzyme that has recently been shown to be necessary for HLH-2

degradation in VU cells (Benavidez et al., 2022). Following perturbation of UBA-1

through RNAi treatment, HLH-2 expression in the VU cells increased more than four-

fold and NHR-67 expression increased by nearly 60% compared to the empty vector

control (Figure 2–figure supplement 1B-D). Both experiments suggest that *nhr-67*

expression in the VU cells is at least partially regulated by levels of HLH-2.

It has previously been proposed that the interaction between HLH-2 and *nhr*-67

is direct. This is based on the identification of E binding motifs within a 276 bp region of

the *nhr-67* promoter that is required for NHR-67 expression in the uterine tissue and

encompasses the location of several hypomorphic mutations (*pf2*, *pf88*, *pf159*) (Figure

2C) (Bodofsky et al., 2018; Verghese et al., 2011). We confirmed this interaction

through a yeast one-hybrid assay after generating a bait strain containing this *nhr-67*

promoter region and pairing it with an HLH-2 Gal4-AD prey plasmid from an existing

yeast one-hybrid library (Reece-Hoyes et al., 2005). Yeast growth on the selective SC-

HIS-TRP plates containing the competitive inhibitor 3-aminotriazole (3-AT)

demonstrated that HLH-2 does indeed bind directly to this 276 bp region of the *nhr-67*

promoter (Figure 2D). Together, these results suggest that direct transcriptional

regulation of *nhr-67* by HLH-2 contributes to the asymmetry in NHR-67 expression

between the AC and VU cells.

NHR-67 forms dynamic punctae in VU cell nuclei that exhibit liquid properties

Upon closer examination of GFP-tagged NHR-67, it became evident that the AC

and VU cells not only exhibit differences in overall NHR-67 levels, but also in

localization of the protein. While NHR-67 localization is fairly uniform throughout the AC

nucleus (excluding the nucleolus), we often observed discrete punctae throughout the

nuclei of VU cells (Figure 3A,B). These punctae were observed with NHR-67

endogenously tagged with several different fluorescent proteins, including GFP,

mNeonGreen, mScarlet-I, and TagRFP-T (Figure 3–figure supplement 1A,B).

Furthermore, by utilizing a live-cell imaging approach, we would not expect to encounter

artificial puncta formation that can result from tissue fixation methods (Irgen-Gioro et al.,

2022). Thus, NHR-67 puncta formation in the VU cells does not appear to be an artifact

of the fluorophore or sample preparation.

To characterize dynamics of these punctae during interphase states of the cell

cycle, we paired GFP-tagged NHR-67 with a CDK activity sensor. The CDK activity

sensor is comprised of a fragment of DNA Helicase B (DHB) fused to a fluorophore (2x-

mKate2), expressed under a ubiquitous promoter (Figure 3C) (Adikes et al., 2020). DHB

contains a strong nuclear localization signal (NLS), flanked by four serine sites, as well

as a weaker nuclear export signal (NES). As CDK activity increases over the cell cycle,

the CDK sensor is translocated from the nucleus to the cytoplasm, allowing for

correlation of its relative subcellular localization to cell cycle state (Figure 3C) (Adikes et

al., 2020; Spencer et al., 2013). Time-lapse microscopy revealed that the number of

NHR-67 punctae was dynamic over the course of the cell cycle, with punctae first

appearing shortly after mitotic exit in the G1 phase, and then reducing in number to two

large punctae prior to nuclear envelope breakdown before disappearing (Figure 3D,E).

We collected additional recordings with finer time resolution and captured fusion, or

condensation, of punctae prior to their dissolution (representative of 6 biological

replicates) (Figure 3F). These punctae also exhibit relatively rapid diffusion kinetics, as

observed by fluorescence recovery following photobleaching ($t_{1/2}$ = 46 seconds; n = 8)

at a rate within the same order of magnitude as P granule proteins PGL-1 and PGL-3

(Figure 3G,H) (Putnam et al., 2019). These properties of NHR-67 punctae are

consistent with those observed with proteins that form nuclear condensates.

Groucho homologs UNC-37 and LSY-22 associate with NHR-67 punctae and

contribute to VU cell fate

In a first step towards defining the role of putative NHR-67 condensates, we

tested the extent to which NHR-67 punctae colocalized with homologs of other proteins

known to form nuclear condensates by pairing GFP- and mScarlet-I-tagged NHR-67

with other endogenously tagged alleles. As NHR-67 is a transcription factor, it is

reasonable to speculate that its punctae may represent clustering around sites of active

transcription, which would be consistent with data showing RNA Polymerase II and the

Mediator complex can associate with transcription factors through phase separation

(Cho et al., 2018). To test this hypothesis, we co-visualized NHR-67 with a GFP-tagged

allele of *ama-1*, the amanitin-binding subunit of RNA polymerase II (Hills-Muckey et al.,

2021) and failed to observe significant colocalization between NHR-67 and AMA-1

punctae (Manders' overlap coefficient, M = 0.066) compared to negative controls where

a single channel was compared to its 90-degree rotation (M = 0.108) (Figure 4A,B).

Another possibility considered is that NHR-67 localization is indicative of chromatin

organization, as heterochromatin has been shown to be compartmentalized in the

nucleus through phase separation (Larson et al., 2017; Strom et al., 2017). However,

we did not observe significant colocalization of NHR-67 with the endogenously tagged

HP1 heterochromatin proteins (Patel and Hobert, 2017) HPL-1 (M = 0.076) or HPL-2 (M

= 0.083) (Figure 4A,B). Recent work in *Ciona* embryos has shown that the

transcriptional co-repressor Groucho forms repressive condensates in nuclei through

phase separation (Treen et al., 2021). The *C. elegans* genome encodes one Groucho

homolog, UNC-37, as well as a Groucho-like protein, LSY-22. To examine their

localization compared to NHR-67, we tagged LSY-22 with TagRFP-T (Figure 4–figure

supplement 1) and acquired a mNeonGreen-tagged allele of *unc-37* (Ma et al., 2021).

Strikingly, we observed significant colocalization of NHR-67 punctae with both LSY-22

(M = 0.686) and UNC-37 (M = 0.741), comparable to colocalization measures in

heterozygous NHR-67::mScarlet-I/NHR-67::GFP animals (M = 0.651), which were used

as positive controls (Figure 4A,B). This evidence suggests that NHR-67 punctae do not

localize to sites of active transcription or chromatin compaction, but instead associate

with transcriptional co-repressors.

Since the AC is the default state of the AC/VU cell fate decision (Seydoux and

Greenwald, 1989), we hypothesized that the punctae including NHR-67, UNC-37, and

LSY-22 may function in repressing invasive differentiation. To test this hypothesis, we

depleted UNC-37 and LSY-22 utilizing the auxin inducible degron (AID) protein

degradation system, in which a protein of interest is tagged with an AID that is

recognized by TIR1 in the presence of auxin and ubiquitinated by the SCF E3 ubiquitin

ligase complex (Figure 4C) (Martinez et al., 2020; Zhang et al., 2015). We re-tagged

LSY-22 with mNeonGreen::AID (Figure 4–figure supplement 1) and acquired a

BFP::AID-tagged allele of *unc-37* (Kurashina et al., 2021). Each AID-tagged allele was

paired with a transgene encoding *Arabidopsis thaliana* TIR1 ($_{At}$TIR1) that was co-

expressed with a nuclear-localized mCherry::HIS-11. Following auxin treatment, we

observed ectopic expression of an AC marker (*cdh-3p*::mCherry::moeABD) in 28% of

LSY-22::AID animals and 59% of UNC-37::AID animals (n = 64 for both) (Figure 4D).

These results are consistent with phenotypes we observed in genetic backgrounds with

unc-37 hypomorphic (*unc-37(e262wd26)*) and null (*unc-37(wd17wd22)*) mutant alleles

(Figure 4–figure supplement 2). It is likely that dual depletion of UNC-37 and LSY-22

would result in a higher penetrance of ectopic ACs given their partial redundancy in

function (Flowers et al., 2010), but animals possessing both AID-tagged alleles were not

viable when paired with the $_{At}$TIR1 transgene.

TCF/LEF homolog POP-1 associates with NHR-67/Groucho punctae and is

necessary for VU cell maintenance post-specification

While UNC-37/LSY-22 colocalization with NHR-67 punctae and ectopic AC

phenotypes are consistent with roles in coordinating AC and VU cell fates, both genes

are broadly expressed and exhibit comparable levels (<10% difference) between the

two cell types (Figure 5A,C; Figure 5–figure supplement 1A,B). Therefore, we

hypothesized that another factor must be involved that confers VU cell specificity to

Groucho-mediated repression of invasiveness. It had previously been reported that the

sole TCF/LEF homolog in *C. elegans*, POP-1, forms a repressive complex with UNC-37

in the early embryo to restrict expression of the endoderm-determining gene, END-1

(Calvo et al., 2001). Additionally, POP-1 has a known role in development of the

somatic gonad, as perturbing its function results in ectopic ACs (Siegfried and Kimble,

2002). Examination of an eGFP-tagged *pop-1* allele (van der Horst et al., 2019),

showed significant enrichment in the VU cells (>20%) compared to the AC (Figure 5B,C;

Figure 5–figure supplement 1A,B). We also observed that POP-1 forms punctae in the

nuclei of VU cells, which had previously been observed during interphase in non-Wnt

signaled embryonic cells (Maduro et al., 2002). We found that these POP-1 punctae

colocalize with NHR-67 (M = 0.547), although to a lesser degree than UNC-37 and LSY-

22, likely because the strong POP-1 fluorescence outside of punctae made them more

difficult to segment (Figure 5D,E). Additionally, *nhr-67(RNAi)* treatment resulted in a

significant increase in AC expression of eGFP::POP-1 compared to empty vector

controls (225%, n > 30), a pattern we observed following depletion of other transcription

factors (Medwig-Kinney et al., 2020) and chromatin modifiers (Smith et al., 2022)

required for AC arrest and invasion (Figure 5F,G; Figure 5–figure supplement 2A,B).

This negative regulation of POP-1 by NHR-67 may explain why the proteins have

opposite patterns of enrichment.

It has previously been suggested that POP-1 may be functioning as an activator

in the VU precursors Z1.ppa and Z4.aap based on the relative expression of a POP-1

transgene (Sallee et al., 2015). This view is largely dependent on the notion that high

levels of POP-1 correlate to repressive function and low levels are conducive for

activator roles (Shetty et al., 2005). In contrast, we did not find evidence of

transcriptional activation by POP-1 in the AC/VU precursors nor their differentiated

descendants using an established POPTOP (POP-1 and TCF optimal promoter)

reporter, which contains seven copies of POP-1/TCF binding sites and the *pes-10*

minimal promoter (Figure 5F; Figure 5–figure supplement 3A,B) (Green et al., 2008).

The growing consensus regarding the Wnt/β-catenin asymmetry pathway is that relative

levels of POP-1 and β-catenin are more important than absolute protein levels of POP-1

(Phillips and Kimble, 2009). Our proposed model of POP-1 acting as a repressor in the

proximal gonad is consistent with the finding that SYS-1 (β-catenin) expression is

restricted to the distal gonad early in somatic gonad development and is not detectable

in the AC or VU cells (Figure 5–figure supplement 3C) (Phillips et al., 2007; Sallee et al.,

2015). It is also supported by recent evidence suggesting that UNC-37/LSY-22 mutant

alleles phenocopy *pop-1* knockdown, which produces ectopic distal tip cells (Bekas and

Phillips, 2022).

One aspect that makes studying the repressive role of POP-1 in cell fate

maintenance challenging is that its activator function is required for distal cell fate

specification in the somatic gonad earlier in development. Loss of either POP-1 and

SYS-1 results in a Sys (symmetrical sister cell) phenotype, where all somatic gonad

cells adopt the default proximal fate and thereby give rise to ectopic ACs (Siegfried and

Kimble, 2002; Siegfried et al., 2004). This likely occluded previous identification of the

repressive role of POP-1 in maintaining VU cell fates. To achieve temporal control over

POP-1 expression to tease apart its two opposing roles, we inserted an AID tag into the

N-terminus of the *pop-1* locus; however, this resulted in gonadal defects even in the

absence of both TIR1 and auxin. Instead, using tools at hand, we paired eGFP-tagged

POP-1 with a uterine-specific anti-GFP nanobody (Smith et al., 2022; Wang et al.,

2017). The anti-GFP nanobody is fused to ZIF-1 and serves as an adapter, recognizing

GFP-tagged proteins and promoting their ubiquitination by the Cullin2-based E3

ubiquitin ligase, which ultimately targets them for degradation via the proteasome

(Figure 6–figure supplement 1A) (Wang et al., 2017). This anti-GFP nanobody,

visualized by nuclear expression of mCherry, was not detectable prior to or even shortly

after the AC/VU cell fate decision, which allowed us to bypass disruption of initial cell

specification (Figure 6–figure supplement 1B). While this method only produced a mild

knockdown of POP-1 in the VU cells, we still observed the ectopic AC phenotype at low

penetrance (7%, n = 60) (Figure 6–figure supplement 1C). To achieve stronger

depletion, we used RNAi for further POP-1 perturbations.

To interrogate the phenotypic consequences of POP-1 perturbation, we utilized a

strain expressing two markers of AC fate (*cdh-3p*::mCherry::moeABD and LAG-

2::P2A::H2B::mTurquoise2). Following treatment with *pop-1(RNAi)*, we observed

several animals with two or more bright *cdh-3/lag-2*+ ACs, consistent with known

phenotypes caused by cell fate misspecification in the somatic gonad (17%, n = 30)

(Figure 6A). We also observed animals with invasive cells that express AC markers at

inconsistent levels (53%, n = 30), suggesting that the cells did not adopt AC fate at the

same time (Figure 6A). To test whether the subset of dim *cdh-3/lag-2*+ ACs are the

result of VU-to-AC cell fate conversion, we visualized AC and VU fates simultaneously

using the AC markers previously described along with an mNeonGreen-tagged allele of

lag-1 (CSL), a protein downstream of Notch signaling whose expression becomes

restricted to the VU cells following AC/VU cell fate specification. Following treatment

with *pop-1(RNAi)*, we found that a subset of ectopic ACs co-express AC markers and

LAG-1, likely indicating an intermediate state between the two cell types (Figure 6–

figure supplement 2). To visualize this process live, we used time-lapse microscopy and

were able to capture ectopic ACs gradually upregulating LAG-2 (+51%, n = 3) and

downregulating LAG-1 (-16%, n = 3) over time (Figure 6B,C), consistent with VU-to-AC

cell fate conversion.

IDR of NHR-67 facilitates protein-protein interaction with UNC-37

Given that UNC-37, LSY-22, and POP-1 phenocopy each other with respect to

AC/VU fates and all three colocalize with NHR-67 punctae, we next sought to further

characterize the interactions among these proteins. Previous work has either directly

identified or predicted protein-protein interactions among POP-1, UNC-37, and LSY-22

(Boxem et al., 2008; Calvo et al., 2001; Flowers et al., 2010; Reece-Hoyes et al., 2005;

Simonis et al., 2009; Zhong and Sternberg, 2006). Using a yeast two-hybrid assay with

UNC-37 Gal4-AD prey, we confirmed that UNC-37 directly interacts with both POP-1

and LSY-22 after observing yeast growth on the selective SC-HIS-TRP-LEU plates

containing 3-AT (Figure 7–figure supplement 1). Using the same technique, we found

that NHR-67 binds directly to UNC-37, as previously predicted (Li et al., 2004; Simonis

et al., 2009), but found no evidence of it directly interacting with LSY-22 or POP-1

(Figure 7–figure supplement 1).

To further characterize the protein-protein interaction between NHR-67 and

UNC-37, we assessed the protein structure of NHR-67 using AlphaFold, an artificial

intelligence-based protein structure prediction tool (Jumper et al., 2021; Varadi et al.,

2022), and PONDR, a predictor of intrinsic disorder (Peng and Zhang, 2006). Both

identify an intrinsically disordered region (IDR) at the C-terminus of NHR-67 (Figure

7A,B). IDRs are low complexity domains that lack fixed three-dimensional structure and

have been shown to support dynamic protein-protein interactions (Chong et al., 2018).

To determine if the IDR of NHR-67 is important for facilitating its interaction with UNC-

37, we repeated the yeast two-hybrid experiment using UNC-37 Gal4-AD prey, pairing it

with different fragments of the NHR-67 protein: full-length, without its IDR (ΔIDR), and

its IDR alone (Figure 7C,D). Yeast growth on the selective SC-HIS-TRP-LEU plates

containing the competitive inhibitor 3-aminotriazole (3-AT) demonstrates that the 108

amino acid IDR sequence of NHR-67 is necessary and sufficient to bind with UNC-37

(Figure 7C,D).

Thus, our current model supported by the data shown here proposes that NHR-

67 levels are controlled through two distinct mechanisms. First, transcription of *nhr-67* is

directly regulated by HLH-2, resulting in enrichment in the AC compared to the VU.

Second, in the AC, where NHR-67 levels are high and POP-1 is repressed, NHR-67 is

free to activate genes promoting invasive differentiation. In the VU cells, where NHR-67

levels are low and POP-1 levels are high, POP-1 assembles with LSY-22, UNC-37, and

NHR-67 to repress NHR-67 targets (Figure 7E). It is possible that POP-1 negatively

regulates NHR-67 at the transcriptional level as well, as the *nhr-67* promoter contains 7

putative TCF binding sites (Zacharias et al., 2015). Taken together, our findings reveal a

dual mechanism for repressing NHR-67 activity in the proliferative VU cells, maintaining

their proliferative fates while suppressing the acquisition of an invasive phenotype.

Discussion

In summary, here we provide evidence that activity of the pro-invasive

transcription factor, NHR-67, is simultaneously regulated by two distinct processes,

which together modulate the proliferative-invasive switch in *C. elegans*. We show that

NHR-67 is a potent fate-specifying transcription factor, in that its expression is sufficient

for invasive differentiation of ACs in the somatic gonad. This could explain why NHR-67

needs to be post-translationally sequestered in addition to being transcriptionally

downregulated in non-invasive cells to fully suppress its function in activating the pro-

invasive program. We also discovered that NHR-67 forms nuclear foci in non-invasive

cells, which exhibit liquid-like properties, indicated by observations of their

condensation, dissolution, and relatively rapid recovery from photobleaching, similar to

what has been described with P granules (Brangwynne et al., 2009). These NHR-67

punctae associate with Groucho homologs, UNC-37 and LSY-22, through a direct

protein-protein interaction with UNC-37 mediated by the C-terminal IDR of NHR-67. We

postulate that this association leads to formation of repressive condensates, as has

recently been described in *Ciona* embryos (Treen et al., 2021). Furthermore, Groucho-

mediated repression of the default invasive state appears to be dependent on

expression of the TCF/LEF homologs POP-1, which clarifies our understanding of the

dual roles this protein plays during development of the somatic gonad. It is also

interesting to note that the dynamic punctae formed by POP-1 in non-Wnt signaled cells

was first described 20 years ago (Maduro et al., 2002), but their function are only now

being appreciated in light of recent advances in our understanding of the formation of

higher order associations in the nucleus.

With regard to phase separation in the nucleus, most research has been through

the lens of transcriptional activation through RNA Polymerase II and the mediator

complex (Boija et al., 2018; Cho et al., 2018; Sabari et al., 2018) or repression through

condensate formation with HP1 heterochromatin proteins (Larson et al., 2017; Strom et

al., 2017). Rather than functioning through either of these established mechanisms, we

have identified here the second observed case of Groucho-mediated repressive

condensates (Treen et al., 2021), which suggests that Groucho proteins may have

evolutionarily conserved roles in repressing transcription through the formation of

nuclear condensates within the Metazoa. Further, we demonstrate how this repressive

mechanism can control cell biology *in vivo* using endogenously tagged alleles and a

clear cell biological read-out of fate and function: invasion versus proliferation.

Still, as this is one of the first studies into the role of repressive condensates *in*

vivo, there is much left to learn. For example, it is unknown whether DNA binding is

necessary for nuclear puncta formation. The interaction between UNC-37 and NHR-67

does not appear to depend on DNA binding, as the C-terminal IDR region of NHR-67

(excluding its zinc finger domains) was sufficient for binding with UNC-37 *in vitro*, but it

is possible that DNA binding is needed for oligomerization *in vivo*. Furthermore, it

remains unclear if suppression of invasive differentiation is achieved by simply

sequestering the pro-invasive transcription factor NHR-67 away from its transcriptional

targets or through direct repression of transcription. If the latter, another question that

arises is how the repressive complex gets recruited to specific genomic sites, since

POP-1 and NHR-67 are both capable of binding to DNA, and whether repression is

achieved through competition with transcriptional activators or recruitment of histone

deacetylases. Direct targets of NHR-67 have not yet been discovered, which makes it

difficult to investigate this specific aspect of the repressive mechanism at present. We

see this as a promising avenue of future study as technologies advance, allowing for

transcriptional profiling and target identification in specific tissues or cells (Gómez-

Saldivar et al., 2020; Katsanos and Barkoulas, 2022).

In this work, we have also identified several perturbations (i.e., increasing levels

of NHR-67, decreasing levels of UNC-37/LSY-22) that result in incompletely penetrant

transdifferentiation phenotypes and/or intermediate cell fates. We foresee these being

ideal cell fate challenge backgrounds in which to perform screens to identify regulators

of cellular plasticity, as has been done in other contexts (Rahe and Hobert, 2019).

Additionally, these induced fate transformations can be paired with tools to visualize and

manipulate cell cycle (Adikes et al., 2020) to determine if any cell cycle state is

particularly permissive for cell fate plasticity. While G1 arrest has been shown to

enhance conversion of human fibroblasts to dopaminergic neurons (Jiang et al., 2015),

mitosis is required for the natural K-to-DVB transdifferentiation event in *C. elegans* (Riva

et al., 2022). As control of proliferation and invasion, as well as maintenance of

differentiated cellular identities, are important for both homeostatic and disease states, it

is our hope that this work will shed light on how cells switch between these states in the

context of cancer growth and metastasis.

Materials and methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (*C. elegans*)	DQM335	Medwig-Kinney et al. (2020)		*egl-43*(*bmd88*[*egl-43p*::EGL-43::loxP::GFP::EGL-43]) II; *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM350	Medwig-Kinney et al. (2020)		*hlh-2*(*bmd90*[*hlh-2p*::loxP::GFP::HLH-2]) I; *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM354	This paper		*nhr-67*(*syb509*[*nhr-67p*::NHR-67::GFP]) IV; *bmd66*[loxP::*egl-43p*::GFP-nanobody::P2A::HIS-58::mCherry] I; *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM368	Medwig-Kinney et al. (2020)		*nhr-67*(*syb509*[*nhr-67p*::NHR-67::GFP]) IV; *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM444	Medwig-Kinney et al. (2020)		*bmd121*[*hsp*::NHR-67::2x-BFP] I; *qyIs227*[*cdh-3p*::mCherry::moeABD] I; *qyIs7*[laminin::GFP] X.

Strain, strain background (*C. elegans*)	DQM515	Medwig-Kinney et al. (2020)		*fos-1*(*bmd138*[*fos-1p*::loxP::GFP::FOS-1]) V; *qyIs227*[*cdh-3p*::mCherry::moeABD] I; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM704	Medwig-Kinney et al. (2021)		*nhr-67*(*bmd212*[*nhr-67p*::NHR-67::TagRFP-T::AID]) IV; *hlh-2*(*bmd90*[*hlh-2p*::LoxP::GFP::HLH-2]) I.
Strain, strain background (*C. elegans*)	DQM800	This paper		*pop-1*(*he335*[*pop-1p*::eGFP::loxP::POP-1]) I; *syIs187*[*pes-10*::7XTCF-mCherry-let-858(3'UTR) + unc-119(+)].
Strain, strain background (*C. elegans*)	DQM811	This paper		*qyIs227*[*cdh-3p*::mCherry::moeABD] I; *lam-2*(*qy20*[*lam-2p*::LAM-2::mNeonGreen]) X; *lag-2*(*bmd202*[*lag-2p*::LAG-2::P2A::H2B::mTurquoise2^lox511^2xHA]) V.
Strain, strain background (*C. elegans*)	DQM853	This paper		*hlh-2*(*bmd90*[*hlh-2p*::loxP::GFP::HLH-2]) I; *stIs11476*[*nhr-67p*::NHR-67::H1-wCherry + *unc-119(+)*].
Strain, strain background (*C. elegans*)	DQM957	This paper		*csh128*[*rpl-28p*::TIR1::T2A::mCherry::his-11] II; *qyIs225*[*cdh-3p*:: mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM958	This paper		*csh140*[*rpl-28p*::TIR1(F79G)::T2A::mCherry::his-11]] II; *qyIs225*[*cdh-3p*:: mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM971	This paper		*pop-1*(*he335*[*pop-1p*::eGFP::loxP::POP-1]) I; *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM989	This paper		*unc-37*(*devKi218*[*unc-37p*::mNeonGreen::UNC-37]) I; *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM990	This paper		*unc-37*(*e262wd26*) I; *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM1003	This paper		*nhr-67*(*syb509*[*nhr-67p*::NHR-67::GFP]) IV; *bmd168*[*rps-27p*::DHB::2x-mKate2] II.
Strain, strain background (*C. elegans*)	DQM1006	This paper		*lsy-22*(*bmd275*[*lsy-22p*::loxP::mNeonGreen::AID::LSY-22]) I; *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM1008	This paper		*pop-1*(*he335*[*pop-1p*::eGFP::loxP::POP-1]) I; *bmd277*[loxP::*egl-43p*::GFP-nanobody::P2A::HIS-58::mCherry] I;

				qyIs225[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM1009	This paper		*unc-37*(*devKi218*[*unc-37p*::mNeonGreen::UNC-37]) I; *nhr-67*(*wy1633*[*nhr-67p*::NHR-67::mScarlet-I::AID*::3xFLAG]) IV.
Strain, strain background (*C. elegans*)	DQM1010	This paper		*hpl-2*(*ot860*[*hpl-2p*::HPL-2::mKate2::HPL-2]) III; *nhr-67*(*syb509*[*nhr-67p*::NHR-67::GFP]) IV.
Strain, strain background (*C. elegans*)	DQM1011	This paper		*hpl-1*(*ot841*[*hpl-1p*::HPL-1::mKate2::HPL-1]) X; *nhr-67*(*syb509*[*nhr-67p*::NHR-67::GFP]) IV.
Strain, strain background (*C. elegans*)	DQM1012	This paper		*lsy-22*(*bmd214*[*lsy-22p*::lox2272::TagRFP-T::AID::LSY-22]) I; *nhr-67*(*syb509*[*nhr-67p*::NHR-67::GFP]) IV.
Strain, strain background (*C. elegans*)	DQM1013	This paper		*pop-1*(*he335*[*pop-1p*::eGFP::loxP::POP-1]) I; *nhr-67*(*syb509*[*nhr-67p*::NHR-67::GFP]) IV.
Strain, strain background (*C. elegans*)	DQM1014	This paper		*unc-37*(*wd17wd22*)/hT2[*bli-4*(*e937*) *let-?*(*q782*) *qIs48*] (I, III); *qyIs225*[*cdh-3p*::mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM1017	This paper		*ama-1*(*ers49*[*ama-1p*::AMA-1::AID::GFP]) IV; *nhr-67*(*wy1633*[*nhr-67p*::NHR-67::mScarlet-I::AID*::3xFLAG]) IV.
Strain, strain background (*C. elegans*)	DQM1051	This paper		*lin-12*(*ljf31*[*lin-12*::mNeonGreen[C1]^loxP^3xFlag]) III; *lag-2*(*bmd202*[*lag-2p*::LAG-2::P2A::H2B::mTurquoise2^lox511^2xHA]) V.
Strain, strain background (*C. elegans*)	DQM1081	This paper		*bmd168*[*rps-27p*::DHB::2x-mKate2] II; *egl-13*(*devKi199*[*egl-13p*::EGL-13::mNeonGreen]) X; *lag-2*(*bmd202*[*lag-2p*::LAG-2::P2A::H2B::mTurquoise2]) V.
Strain, strain background (*C. elegans*)	DQM1101	This paper		*lsy-22*(*bmd275*[*lsy-22p*::^loxP^mNeonGreen::AID::LSY-22]) I; *csh128*[*rpl-28p*::TIR1::P2A::mCherry::his-11]) II; *qyIs225*[*cdh-3p*:: mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.
Strain, strain background (*C. elegans*)	DQM1115	This paper		*unc-37*(*miz36*[*unc-37p*::UNC-37::AID::BFP]) I; *csh128*[*rpl-28p*::TIR1::P2A::mCherry::his-11]) II; *qyIs225*[*cdh-3p*:: mCherry::moeABD] V; *qyIs7*[laminin::GFP] X.

Strain, strain background (*C. elegans*)	DQM1127	This paper		*nhr-67(syb509[nhr-67p*::NHR-67::GFP]) IV; *stIs11476[nhr-67p*::NHR-67::H1-wCherry + unc-119(+)].
Strain, strain background (*C. elegans*)	DQM1129	This paper		*bmd143[hsp*::HLH-2::2xBFP] I; *nhr-67(syb509[nhr-67p*::NHR-67::GFP]) IV.
Strain, strain background (*C. elegans*)	DQM1135	This paper		*qyIs227[cdh-3p*::mCherry::moeABD] I; *lam-2(qy20[lam-2p*::LAM-2::mNeonGreen]) X; *lag-2(bmd202[lag-2p*::LAG-2::P2A::H2B::mTurquoise2^lox511^2xHA]) V; *lag-1(devKi208[lag-1*::mNeonGreen]) IV.
Strain, strain background (*C. elegans*)	JK3791	Phillips et al. (2007)		*qIs95[sys-1p::*Venus::SYS-1 + *pttx-3::DsRed]*
Strain, strain background (*C. elegans*)	NK1034	Matus et al. (2015)		*qyIs225[cdh-3p*::mCherry::moeABD] V; *qyIs7[laminin*::GFP] X.
Strain, strain background (*C. elegans*)	PHX509	Medwig-Kinney et al. (2020)		*nhr-67(syb509[nhr-67p*::NHR-67::GFP]) IV.
Strain, strain background (*C. elegans*)	PS5332	Green et al. (2008)		*syIs187[pes-10::7XTCF*-mCherry-*let-858(3'UTR) + unc-119(+)]*
Strain, strain background (*C. elegans*)	RW11476	Gerstein et al. (2010)		*unc-119(tm4063)* III; *stIs11476[nhr-67*::H1-wCherry + unc-119(+)].
Strain, strain background (*C. elegans*)	SV2114	van der Horst et al. (2019)		*pop-1(he335[*eGFP::loxP::*pop-1])* I.
Strain, strain background (*C. elegans*)	TV27467	This paper		*nhr-67(wy1632[nhr-67p*::NHR-67::mNeonGreen::AID*::3xFLAG]) IV.
Strain, strain background (*C. elegans*)	TV27468	This paper		*nhr-67(wy1633[nhr-67p*::NHR-67::mScarlet-I::AID*::3xFLAG]) IV.
Recombinant DNA reagent	Plasmid: pTNM087	This paper		*lsy-22* sgRNA plasmid (AAACGAAGTGGATCAGCCAG)
Recombinant DNA reagent	Plasmid: pTNM088	This paper		*lsy-22*^SEC^TagRFP-T::AID repair plasmid
Recombinant DNA reagent	Plasmid: pTNM140	This paper		*lsy-22*^SEC^mNeonGreen::AID repair plasmid
Chemical compound, drug	1-Naphthaleneacetic acid, potassium salt (K-NAA)	PhytoTech Labs	N610	
Chemical compound, drug	Hygromycin B	Omega Scientific, Inc.	HG-80	
Chemical compound, drug	Levamisole hydrochloride	Sigma-Aldrich	31742	

Chemical compound, drug	Sodium azide	Sigma-Aldrich	S2002	
Software, algorithm	Adobe Illustrator	Adobe	Version 26.0.2	
Software, algorithm	Alpha Fold	Jumper et al. (2021); Varadi et al. (2021)	Version 2	
Software, algorithm	ApE – A Plasmid Editor	M. Wayne Davis	Version 2.0.61	
Software, algorithm	Fiji/ImageJ	Schindelin et al. (2012)	Version 2.0.0-rc-69/1.53e	
Software, algorithm	ggplot2	Tidyverse	Version 3.3.5	
Software, algorithm	Exon-Intron Graphic Maker	Nikhil Bhatla	Version 4	
Software, algorithm	JACoP (Just Another Colocalization Plugin)	Bolte and Cordelières (2006)	Version 2.1.1	
Software, algorithm	Metamorph	Molecular Devices	Version 7.10.3.279	
Software, algorithm	Rstudio	R	Version 1.4.1717	

C. elegans strains, culture, and nomenclature

Methods for *C. elegans* culture and genetics were followed as previously
described (Brenner, 1974). Developmental synchronization for experiments was
achieved through alkaline hypochlorite treatment of gravid adults to isolate eggs (Porta-
de-la-Riva et al., 2012). L1 stage animals were plated on nematode growth media
plates and subsequently cultured at 20°C or 25°C. Heat shock-inducible transgenes
were activated by incubating animals on plates sealed with Parafilm in a 33°C water
bath for 2-3 hours. In the text and in figures, promoter sequences are designated with a
"*p*" following the gene name and gene fusions are represented by a double-colon (::)
symbol.

CRISPR/Cas9 injections

New alleles and single-copy transgenes were generated by homology directed
repair using CRISPR-based genome engineering. mScarlet::AID and mNeonGreen::AID
were inserted into the C-terminus of the *nhr-67* locus by injecting adult germlines with

Cas9 guide-RNA ribonucleoprotein complexes and short single-stranded

oligodeoxynucleotide donors, as previously described (Ghanta and Mello, 2020).

Successful integrants were identified through screening for fluorescence and by PCR.

The *lsy-22* locus was edited by injecting a Cas9 guide RNA plasmid and repair template

plasmid containing a self-excising cassette with selectable markers to facilitate

screening (Dickinson et al., 2015; Dickinson and Goldstein, 2016; Huang et al., 2021).

Repair templates used to tag *lsy-22* with TagRFP-T::AID and mNeonGreen::AID were

generated by cloning ~750-850 bp homology arms into pTNM063 and pDD312,

respectively (Ashley et al., 2021; Dickinson et al., 2015). All guide and repair sequences

used can be found in Supplemental Table 1.

Existing alleles

The GFP-tagged alleles of the pro-invasive transcription factors *(egl-43, fos-1,*

hlh-2, and *nhr-67)* and the TagRFP-T::AID-tagged *nhr-67* allele were generated in

preceding work (Medwig-Kinney et al., 2021, 2020). Recent micropublications describe

the P2A::H2B::mTurquoise2-tagged *lag-2* and mNeonGreen-tagged *lin-12* alleles used

in this study (Medwig-Kinney et al., 2022; Pani et al., 2022). The eGFP-tagged *pop-1*

allele and POPTOP reporter were previously published (Green et al., 2008; van der

Horst et al., 2019), as were the AID::BFP and mNeonGreen tagged alleles of *unc-37*

(Kurashina et al., 2021; Ma et al., 2021). GFP-tagged *ama-1* (Hills-Muckey et al., 2021)

as well as mKate2-tagged *hpl-1* and *hpl-2* (Patel and Hobert, 2017) were also

disseminated in prior publications. The single-copy transgenes expressing the CDK

sensor and TIR1 variants under ubiquitously expressed ribosomal promoters (*rps-27*

and *rpl-28*, respectively) as well as the tissue-specific GFP-targeting nanobody are

described in previous work (Adikes et al., 2020; Hills-Muckey et al., 2021; Smith et al.,

2022; Wang et al., 2017) and are located at neutral genomic sites, ttTi4348 or ttTi5605

(Frøkjær-Jensen et al., 2013). The same is true for the heat shock inducible constructs

for HLH-2 and NHR-67 (Medwig-Kinney et al., 2020). The cadherin *(cdh-3)* anchor cell

reporter and basement membrane (laminin) markers have already been characterized

(Keeley et al., 2020; Matus et al., 2010). The following mutant alleles were obtained

from the *Caenorhabditis* Genetics Center: *unc-37(e262wd26)* and *unc-37(wd17wd22)*

(Pflugrad et al., 1997), the latter of which was maintained using the chromosome I/III

balancer *hT2* (McKim et al., 1993). The genotypes of all strains used in this study can

be found within the Key Resources Table.

Auxin inducible protein degradation

The auxin inducible degron (AID) system was utilized to strongly deplete proteins

of interest (Zhang et al., 2015). AID-tagged alleles were paired with the *Arabidopsis*

thaliana F-box protein, transport inhibitor response 1 ($_{At}$TIR1), and treated with the

water-soluble auxin 1-Naphthaleneacetic acid (K-NAA) at 1 mM concentration (Martinez

et al., 2020). Auxin was added to nematode growth media plates according to

previously published protocols (Martinez and Matus, 2020), which were then seeded

with OP50 *E. coli*. To achieve robust depletion, synchronized L1 stage animals were

directly plated on auxin plates.

RNA interference

The RNAi clones targeting *pop-1* and *uba-1* as well as the corresponding empty

vector control (L4440) were obtained from the Vidal library (Rual et al., 2004). The RNAi

constructs targeting the pro-invasive transcription factors (*egl-43*, *fos-1*, *hlh-2*, and *nhr-*

67) and chromatin modifiers (*pbrm-1*, *swsn-4*, and *swsn-8*) are derived from the highly

efficient RNAi vector T444T (Sturm et al., 2018) and were generated in preceding work

(Medwig-Kinney et al., 2020; Smith et al., 2022). To avoid known AC/VU cell fate

specification defects caused by *hlh-2* perturbations, synchronized animals were grown

on OP50 until the L2 stage when they were shifted to *hlh-2* RNAi plates.

Live cell imaging

With the exception of the FRAP experiments shown in Figure 3, all micrographs

were collected on a Hamamatsu Orca EM-CCD camera mounted on an upright Zeiss

AxioImager A2 with a Borealis-modified CSU10 Yokagawa spinning disk scan head

(Nobska Imaging) using 405 nm, 440 nm, 488 nm, 514 nm, and 561 nm Vortran lasers

in a VersaLase merge and a Plan-Apochromat 100×/1.4 (NA) Oil DIC objective.

MetaMorph software (Molecular Devices) was used for microscopy automation. Several

experiments were scored using epifluorescence visualized on a Zeiss Axiocam MRM

camera, also mounted on an upright Zeiss AxioImager A2 and a Plan-Apochromat

100×/1.4 (NA) Oil DIC objective. For static imaging, animals were mounted into a drop

of M9 on a 5% Noble agar pad containing approximately 10 mM sodium azide

anesthetic and topped with a coverslip. For long-term time-lapse imaging, animals were

first anesthetized in 5 mM levamisole diluted in M9 for approximately 20 minutes, then

transferred to a 5% Noble agar pad and topped with a coverslip sealed with VALAP

(Kelley et al., 2017). For short-term time-lapse imaging, the pre-anesthetization step

was omitted, and animals were transferred directly into a drop of 5 mM levamisole

solution on the slide.

Fluorescence recovery after photobleaching

FRAP experiments were performed using an Acal BFi UV Optimicroscan

photostimulation device mounted on a spinning disk confocal system consisting of a

Nikon Ti2 inverted microscope with Yokogawa CSU-W1 SoRa spinning disk. Data were

acquired using a Hamamatsu ORCA Fusion camera, 60x 1.27 NA water immersion

objection, SoRa disk, and 2.8x SoRa magnifier. Single plane images were collected

every 1 second.

Yeast one-hybrid

The 276 bp fragment of the *nhr-67* promoter (Bodofsky et al., 2018) was cloned

into the pMW2 vector, linearized by BamHI digestion. Linearized plasmid was

transformed into the Y1H yeast strain (as described in Reece-Hoyes and Walhout,

2018). Transformed yeast was plated on SC-HIS plates for three days before being

transformed with the HLH-2 Gal4-AD plasmid. Three colonies from each transformation

plate were streaked onto SC-HIS-TRP +3-aminotriazole (3-AT) plates. Protein-DNA

interactions were determined by visible growth on 3-AT conditions with negative growth

in empty vector controls after three days. Plates were imaged on a Fotodyne

FOTO/Analyst Investigator/FX darkroom imaging station.

Yeast two-hybrid

Plasmids containing target proteins fused to GAL-4 DNA-binding-domain + LEU

and GAL-4 Activation Domain + TRP were co-transformed into the pJ69-4a Y2H yeast

strain as previously described (Reece-Hoyes and Walhout, 2018). Transformed yeast

was plated on SC-TRP-LEU plates for three days. Three colonies from each

transformation plate were streaked onto SC-HIS-TRP-LEU 3-AT plates. Protein

interactions were determined by visible growth on 3-AT conditions with negative growth

in empty vector controls after three days. Plates were imaged as described in the

previous section.

Quantification of protein expression and cell cycle state

Image quantification was performed in Fiji/ImageJ (Schindelin et al., 2012).

Protein expression was quantified by drawing a region of interest around the nucleus of

the cell of interest and measuring the mean gray value, then manually subtracting the

mean gray value of a background region of similar area to account for camera noise.

The CDK sensor was quantified as previously described (Adikes et al., 2020). Following

rolling ball subtraction (50 pixels), mean gray value is measured in a region of interest

drawn within the cytoplasm and one around the nucleus excluding the nucleolus. The

cytoplasmic-to-nuclear ratio correlates to CDK activity and is used to assess cell cycle

state (Adikes et al., 2020; Spencer et al., 2013). Movies were collected by acquiring z-

stacks at 5-minute intervals. Samples were time-aligned relative to anaphase. Cells that

did not undergo anaphase during the acquisition period were aligned based on their

DHB ratios. Animals that arrested in development (i.e., did not show evidence of

progressing through the cell cycle) were excluded from analysis.

Colocalization analyses

For colocalization analyses, single plane images were collected to avoid z drift

during acquisition and prevent photobleaching, which was often non-uniform between

red and green fluorophores. Micrographs were subject to background subtraction

(rolling ball radius = 50) followed by thresholding to segment punctae. Manders' overlap

coefficients (M) were calculated by measuring the extent that segmented punctae of

NHR-67 overlapped with that of other proteins using Just Another Colocalization Plugin

(JACoP) in Fiji/ImageJ (Bolte and Cordelières, 2006; Schindelin et al., 2012).

Heterozygous animals for *nhr-67*::mScarlet and *nhr-67*::GFP were used as positive

controls. These images were then re-analyzed following 90-degree rotation of one of the

two channels being compared, resulting in random colocalization that served as a

negative control.

Data visualization and statistical analyses

Representative images were processed using Fiji/ImageJ (Schindelin et al.,

2012). Heat maps were generated using the Fire lookup table. Tests to determine

statistical significance of data were conducted in RStudio and plots were generated

using the R package ggplot2 (Wickham, 2016). Error bars represent the mean ±

standard deviation. Schematics of gene loci were generated using sequences from

WormBase (Harris et al., 2020) and the Exon-Intron Graphic Maker

(http://wormweb.org/exonintron). Figures were assembled in Adobe Illustrator.

<u>**Acknowledgments**</u>

We are grateful to Dr. Derek Applewhite and Aidan Teran for advice on quantification of

protein colocalization. Additionally, we thank Chris Zhao for constructive comments on

the manuscript. Some strains were provided by the Caenorhabditis Genetics Center,

which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

<u>**Additional information**</u>

Funding

Funder	Grant reference number	Author
National Institutes of Health	R01GM121597	David Q. Matus
Damon Runyon Cancer Research Foundation	DRR-47-17	David Q. Matus

National Institutes of Health	F31HD100091	Taylor N. Medwig-Kinney
Stony Brook University	Presidential Critical Research Funds	Taylor N. Medwig-Kinney
National Institutes of Health	F30CA257383	Michael A. Q. Martinez
Human Frontier Science Program	LTF000127/2016-L	Callista Yee
Howard Hughes Medical Institute	Investigator	Kang Shen
National Institutes of Health	R01GM117406	Christopher M. Hammell
National Science Foundation	2217560	Christopher M. Hammell
National Institutes of Health	R35GM142880	Ariel M. Pani

Author contributions

Taylor N. Medwig-Kinney, Conceptualization, Formal Analysis, Investigation, Writing—

original draft, Visualization, Supervision, Funding acquisition; Brian A. Kinney,

Investigation, Writing—review and editing; Michael A. Q. Martinez, Investigation,

Resources, Writing—review and editing; Callista Yee, Resources, Writing—review and

editing; Sydney S. Sirota, Formal Analysis, Investigation, Resources; Angelina A.

Mullarkey, Formal Analysis, Writing—review and editing; Neha Somineni, Formal

Analysis, Resources; Justin Hippler, Formal Analysis, Resources; Wan Zhang,

Resources; Kang Shen, Supervision; Christopher M. Hammell, Resources, Supervision,

Writing—review and editing; Ariel M. Pani, Investigation, Writing—review and editing;

David Q. Matus, Conceptualization, Supervision, Funding acquisition, Writing—review

and editing

Author ORCIDs

Taylor N. Medwig-Kinney, https://orcid.org/0000-0001-7989-3291

Brian A. Kinney, https://orcid.org/0000-0001-5628-1436

Michael A. Q. Martinez, https://orcid.org/0000-0003-1178-7139

Callista Yee, https://orcid.org/0000-0002-2928-492X

Sydney S. Sirota, https://orcid.org/0000-0001-9990-3266

Angelina A. Mullarkey, https://orcid.org/0000-0002-5830-5347

Neha Somineni, https://orcid.org/0000-0001-5702-1695

Kang Shen, https://orcid.org/ 0000-0003-4059-8249

Christopher M. Hammell, https://orcid.org/0000-0002-5961-0976

Ariel M. Pani, https://orcid.org/0000-0002-9338-9750

David Q. Matus, https://orcid.org/0000-0002-1570-5025

Figure legends (supplementary figure legends are indented)

Figure 1: Invasive AC fate correlates to high levels of NHR-67. (A) Schematic of *C.*

elegans anchor cell (AC, magenta) and ventral uterine (VU, blue) cell fate specification

from the Z1 and Z4 somatic gonad precursor cell lineages (p, posterior daughter; a,

anterior daughter). (B) Micrographs depicting AC and VU cell differentiation over

developmental time. AC/VU precursors express LAG-2 (H2B::mTurquoise), which

eventually becomes restricted to the AC, whereas VU cells express LAG-1

(mNeonGreen) post-specification. The differentiated AC (*cdh-3p*::mCherry::moeABD)

then invades through the underlying basement membrane (LAM-2::mNeonGreen). (C-

D) Representative heat map micrographs (C) and quantification (D) of GFP-tagged

HLH-2 and NHR-67 expression in the AC and VU cells at the time of AC invasion. (E)

Expression of Notch (*lin-12*::mNeonGreen) and Delta (*lag-2*::P2A::H2B::mTurquoise2)

following RNAi-induced knockdown of *nhr-67* compared to empty vector control. (F)

Micrographs depicting the ectopic invasive ACs (*cdh-3p*::mCherry::moeABD,

arrowheads) and expanded basement membrane (*laminin*::GFP, arrows) gap observed

following heat shock induced expression of NHR-67 (*hsp*::NHR-67::2x-BFP) compared

to non-heat shocked controls. (G) Schematic summarizing AC and VU cell fates that

result from perturbations of NHR-67 levels. For all figures: asterisk (*), AC/VU

precursor; plus (+), VU precursor; solid arrowhead, AC; open arrowhead, VU cell;

arrows, basement membrane breach. Statistical significance determined by Student's t-

test (*$p > 0.05$, **$p > 0.01$, ***$p > 0.001$). Scale bars, 5 µm.

Figure 1–figure supplement 1: Expression of pro-invasive transcription

factors EGL-43 and FOS-1 in the somatic gonad. (A) Schematic of the AC

pro-invasive gene regulatory network comprised of four transcription factors:

EGL-43, FOS-1, HLH-2, and NHR-67. (B-C) Representative heat-map

micrographs (B) and quantification (C) of GFP-tagged EGL-43 and FOS-1

expression in the AC and VU cells.

Figure 2: NHR-67 expression is downregulated in VU cells through direct

transcriptional regulation by HLH-2. (A-B) Representative heat map micrographs (A)

and quantification (B) of NHR-67::GFP expression in VU cells following heat shock

induced expression of HLH-2 (2x-BFP) compared to non-heat shocked controls. (C)

Schematic of a 276 bp putative regulatory element within the promoter of *nhr-67*

(Bodofsky et al., 2018), annotated with the location of three hypomorphic mutations

(*pf2*, pf88, and *pf159*). (D) Yeast one-hybrid experiment pairing HLH-2 Gal4-AD prey

with the 276 bp fragment of the *nhr-67* promoter as bait on SC-HIS-TRP plates with and

without competitive inhibitor 3-AT (175 mM).

Figure 2–figure supplement 1: Onset of expression and regulatory

interaction between NHR-67 and HLH-2 in the somatic gonad. (A)

Micrographs depicting onset of GFP-tagged HLH-2 and a wCherry-labeled NHR-

67 transgene (inverted to aid visualization) in Z1.pp and Z4.aa cells at early (top)

and late (bottom) stages. (B-D) Representative micrographs (B) and

quantification (C-D) of GFP-tagged HLH-2 and TagRFP-T-tagged NHR-67 in AC

(C) and VU cells (D) following *uba-1(RNAi)* compared to control. Insets depict

different z planes of the same image.

Figure 3: NHR-67 forms dynamic punctae in nuclei of VU cells. (A) Heat-map

maximum intensity projection of NHR-67::GFP showing protein localization in the AC and VU cells. (B) Spatial color coded projection of NHR-67::GFP punctae in VU, with nuclear border indicated with a dotted line. (C) Schematic of DNA Helicase B (DHB) based CDK sensor and its dynamic localization over the cell cycle. (D) Graphs depicting CDK activity levels and corresponding cell cycle state (top), and percentage of cells exhibiting NHR-67::GFP punctae (bottom) over time, aligned to anaphase. (E) Representative time-lapse of NHR-67::GFP over the course of a cell cycle, with cell membranes indicated with dotted lines. (F) Time-lapse depicting NHR-67::GFP punctae fusion prior to cell division. Bottom panels are pseudo-colored. (G-H) Quantification (G) and representative images (H) depicting fluorescence recovery of NHR-67::GFP following photobleaching of individual punctae (arrow).

Figure 3–figure supplement 1: Knock-in alleles of *nhr-67*. (A) Representative images of VU cells exhibiting punctae formed by NHR-67 tagged with GFP, mNeonGreen, mScarlet-I, and TagRFP-T. (B) Schematics of the new endogenously tagged loci generated in this paper for *nhr-67*. Scale bar, 100 base pairs (bp).

Figure 4: Groucho homologs LSY-22 an UNC-37 colocalize with NHR-67 punctae and repress invasive differentiation in VU cells. (A) Co-visualization of NHR-67 with RNA Polymerase II (GFP::AMA-1), HP1 heterochromatin proteins (HPL-1::mKate2 and HPL-2::mKate2), and Groucho homologs (TagRFP-T::LSY-22 and mNeonGreen::UNC-37) in VU cells using endogenously tagged alleles. (B) Quantification of colocalization, with plot reporting Manders' overlap coefficients compared to negative controls (90 degree rotation of one channel) and positive controls. (C) Schematic of the auxin inducible degron (AID) system, where $_{At}$TIR1 mediates proteasomal degradation of AID-tagged proteins in the presence of auxin. (D) Representative images of phenotypes observed following individual AID-depletion of UNC-37 and LSY-22 compared to control animals without AID-tagged alleles. Insets depict different z planes of the same image.

Figure 4–figure supplement 1: Knock-in alleles of *lsy-22*. Schematics of the

new endogenously tagged loci generated in this paper for *lsy-22*. Scale bar, 100
base pairs (bp).

Figure 4–figure supplement 2: UNC-37 mutants show ectopic expression of
AC markers. Ectopic expression of AC marker (*cdh-3p*::mCherry::moeABD) in
hypomorphic (*unc-37(e262wd26)*) and null (*unc-37(wd17wd22)*) alleles of *unc-37*
compared to wild-type *unc-37*. Insets depict different z planes of the same image.

Figure 5: POP-1 associates with repressive condensates in the absence of Wnt
signaling. (A-B) Expression of mNeonGreen::UNC-37 and mNeonGreen::LSY-22 (A)
and eGFP::POP-1 (B) in the AC/VU precursors pre-specification (left), as well as the AC
and VU cells post-specification (right). (C) Quantification of UNC-37, LSY-22, and POP-
1 expression at the time of AC invasion. (D) Co-visualization of NHR-67::mScarlet-I and
EGFP::POP-1 in the VU. (E) Quantification of POP-1 and NHR-67 colocalization, with
plot reporting Manders' overlap coefficient compared to negative and positive controls.
(F) Representative micrographs showing expression of POPTOP, a synthetic *pop-1-*
activated reporter construct, in wild-type ACs, VU cells, and their precursors. Insets
depict different z planes of the same image. (G-H) Micrographs (G) and quantification
(H) of eGFP-tagged POP-1 expression in proliferative ACs following RNAi depletion of
nhr-67 compared to empty vector control.

Figure 5–figure supplement 1: Expression of LSY-22, UNC-37, and POP-1
over developmental time. (A-B) Developmental series (A) and quantified
expression (B) of mNeonGreen::UNC-37, mNeonGreen::LSY-22, and
eGFP::POP-1 expression in the AC/VU precursors, AC, and VU cells over time.
Following AC/VU cell specification, animals are staged by the division of the
underlying primary vulval precursor cells (1° VPCs).

Figure 5–figure supplement 2: POP-1 repressive function in VU cells is
distinct from activator function in distal somatic gonad. (A) Schematics
representing the dual functions of POP-1. In the presence of Wnt signaling, POP-

1 binds to its co-activator β-catenin (e.g., SYS-1) and activates transcription of its

target genes. In the absence of Wnt signaling, POP-1 binds to its co-repressor

Groucho (UNC-37) and represses transcription of its target genes. (B)

Representative micrographs of eGFP::POP-1 and POPTOP (*pes-10*::7x-

TCF::mCherry) expression in the AC, dorsal uterine cells (DU),

spermatheca/sheath cells (SS), and VU cells. (C) Schematic of SYS-1 (β-

catenin) expression in the Z1/Z4 lineage (based on Philips et al., 2007).

Figure 5–figure supplement 3: POP-1 expression is regulated by the cell

cycle-dependent pro-invasion pathway. (A-B) Representative micrographs of

eGFP::POP-1 and POPTOP (*pes-10*::7x-TCF::mCherry) following RNAi-induced

knockdown of pro-invasive transcription factors and chromatin modifiers

compared to control AC and VU cells. (B) Quantification of eGFP::POP-1

expression in ACs following RNAi treatments. Here, the presence of multiple ACs

are the result of failure of the AC to exit the cell cycle.

Figure 6: Ectopic ACs arise through VU-to-AC cell fate transformation. (A)

Representative images of ectopic AC (*cdh-3p*::mCherry::moeABD; LAG-

2::P2A::H2B::mTurquoise2) phenotypes observed following RNAi depletion of POP-1.

Schematics (right) depict potential explanations for observed phenotypes. (B)

Expression of AC markers and a VU cell marker (LAG-1::mNeonGreen, inverted to aid

visualization) in *pop-1(RNAi)* treated animals over time. (C) Quantification of LAG-2

(magenta) and LAG-1 (blue) expression in transdifferentiating cells produced by *pop-*

1(RNAi) over time.

Figure 6–figure supplement 1: POP-1 functions to regulate AC/VU cell fates

post-specification. (A) Schematic of the anti-GFP nanobody protein

degradation system (based on Wang et al., 2017). (B) Micrographs

demonstrating that the anti-GFP nanobody (driven under the *egl-43L* promoter) is

not expressed pre-specification or even shortly after when the presumptive AC

begins to express its differentiated cell reporter (*cdh-3*). (C) With decreased

levels of *pop-1*, a low penetrance (~7%) of multi-AC phenotypes were observed.

Figure 6–figure supplement 2: Ectopic ACs resulting from *pop-1*

perturbation express VU cell markers. Expression of AC markers (*cdh-*

3p::mCherry::moeABD; LAG-2::P2A::H2B::mTurquoise2) and VU marker (LAG-

1::mNeonGreen) in *pop-1(RNAi)* treated animals compared to empty vector

control.

Figure 7: NHR-67 binds to UNC-37 through IDR-mediated protein-protein

interaction. (A) Predicted structure of NHR-67 generated by AlphaFold. (B) Measure of

intrinsic disorder of NHR-67 using the PONDR VSL2 prediction algorithm. (C)

Schematic of NHR-67 protein coding sequences used for Yeast two-hybrid experiments

with reference to its intrinsically disordered region (IDR, magenta), DNA binding domain

(DBD, green), and ligand binding domain (LBD, cyan). Scale bar, 10 amino acids. (D)

Yeast two-hybrid experiment shows pairing of UNC-37 with either full-length NHR-67 or

the IDR alone allows for yeast growth in the presence of competitive inhibitor 3-AT (20

mM). (E) Summary model of the roles of NHR-67, UNC-37, LSY-22, and POP-1 in

maintenance of AC and VU cell fate.

Figure 7–figure supplement 1. NHR-67 exhibits protein-protein interaction

with UNC-37. Yeast two-hybrid experiment pairing UNC-37 and NHR-67 Gal4-

AD prey with LSY-22, NHR-67, and POP-1 Gal4-DBD bait on SC-HIS-TRP-LEU

plates with and without competitive inhibitor 3-AT (20 mM).

Supplemental Table 1: Sequences

Method	Reagent	Sequence
CRISPR	*lsy-22* sgRNA	AAACGAAGTGGATCAGCCAG
CRISPR	*lsy-22* left homology arm	GAGACCAGGCAGCATATTTTTGTGCAAACTCCTTTGAAATTATTGTTATT ACAGATTTTTTGAAATTAGTAATTTCGGAAGTCTCCAAATTCTTAAAATGT CCTTTCCTTCCTTCGTCTCCTTCTCACTTTCCCGGCGGTAATTGGTTTTG GCGTAGAACATGGCGCCGGCGCAAACACATTGACGCAAAAGTTCACAA TTTAAAAGACCGTGTAAACAAGACCTCCTCCCCCTGTCCGCTCGTCTTC

		TCAAAGAACTTTTTTATCTTTTCCGTTGTTTTATCGCTTTTCTCATAATAA AAATAAAAATCGTCGTTGAAGAAATAAAATGGGCCCCGCCGTTGGTGTG TGTGTGCCATCACAACCATCAATTCAAACTTTGTTTTATTTCTACTTTTTT GCAGCTCATTGCTCCGTGACGGATTCAAATTGTTGTTTTTCTTCGCGAC AAACAAGAAAGACTCTGGAATTGTATCGTTTTCGAATTTTTAAAATTTGG AATTTTTGATCGCCTGTCGATCAACCCCCGTCTACAATCACGTCGGACC AACGAAGATTCGCGAATCCGAACTTCTCGCATCCTAGCTCGAGTAGTTC AAGGGAAAAGTAGGTTTTTATTATATTTTAAAAGTAAAAGATTTGTTGAC AAATAGTTGAAAATGGTATTCTAAAAGTTGAGAAAATTAAAAATATGAAA GGTTCACTAGAGGCATATTTTTCATTTTGAAAGTATCAAAATAAATTCAA AAACAGCGTTTAATTTCTTTTAGAAAACTAGAAATTCGAAAACTAGAAAT TGGTGAATATGAACAGTATTTAATTTATATTTTAGTAGGAAACGAAGTGG ATCAGCCAGCGCA
CRISPR	*lsy-22* right homology arm	ATGTCACTAGCAGATCATATTGATGCAATCAAAAAAGAAATGACAAGTCT ATCGAGCCAATTCAACTCGAATAAATCAGAACTTGAACGATCAAAACAG GATTTATCATCAATGCAACTACAAATGCAACAACAAATGAACGATTCAGT ACAGCAGGACCTCCAGAAAGCTCATGATACATCGAAGAAATTAGATAAT CTTGCAAGGACGTTTCTTGCATTAATTACTGATCCATCGCAACAATCACA ATTATTAGCGGAATTTGAATCAATTTCCAAGAACGGAACACCTCAAATG CCAGTGATGACACCTCCAATGATGAATGCAGTTGCTGCAATGCAACCAC ATATGATGATGAGGAATCCAATGCTGAATGCAATGGGTGGAGCATCAC CACGTGTTTCTGGAAAAGGTGGAGCTCCACAAAACGGTGGAATGAATG GGATGTCGATGATGCAGCACATGCAAATGCAAGCATTCCAGGCACAAA TGATGCAAGCTCAAATGTTTCAGCAGATGCCGATGATGATGCCAGGGA TGGTTCCTGGAATGCCACCGGGAATGGCTGGATTGCCTGGAATGATGC CACCGAATAATATGGCAGCTGCAATGCAACATTTTAATGCGGTAAGTGT TAACTATTATTCAAAATGTTGAGATTAAATTATGTTTATCGTAATTTTCTC TTAGGGGCACGGATTTATCCAGATGGGGTCGGCGGGGAAAAATCTATG ATAGT
CRISPR	*nhr-67* sgRNA	AGAGAGTGTTAATGTTGAAG
CRISPR	*nhr-67* left homology arm	TTCCTCAACTCAGCCTTCATCGGCATCATCCCCTTCCTCTTCAAGACCA CGTCATTCGATTCGATCAATAACTGAATTATTATCAATTCAAGAAGAGGA AAGCGTGAACGTGGAGGAAGTG
CRISPR	*nhr-67* right homology arm	TAAATAGTAAATTCATGTTTCATATACAGTAACTCAATTATTCTAAGTATC TCTTTTCATTGTCTTTTTCACTCCGTTTCTTGCCTCGCCCGGATTTTCATT GGATTTTGATTTATACTT
Y1H	276 bp *nhr-67* promoter element	ACCATTTCGCTAAGTTTTCCAAAAGTTCACACCTGTATGACCCTCGTCTG TCTATCTCTCTTCAACGACCTTTTCCTGCATCGTTCTCGATATTGCGTAA AATCCCAACACATTGTCTTACTCTTGGGTAAAATCGAGAAAAAAAGTAT TTTTGATTGCGTAAGCATCAAAATGAAATGCAGGTGTTCACCACTAAAAT GAAATGGAGGTCATGTTCGTACCCTGTGGTACAAGCCACAGAGAGTTC AATTTATATGATGAGCATTTGAAGAATT
Y2H	LSY-22 cDNA	ATGTCACTAGCAGATCATATTGATGCAATCAAAAAAGAAATGACAAGTCT ATCGAGCCAATTCAACTCGAATAAATCAGAACTTGAACGATCAAAACAG GATTTATCATCAATGCAACTACAAATGCAACAACAAATGAACGATTCAGT ACAGCAGGACCTCCAGAAAGCTCATGATACATCGAAGAAATTAGATAAT CTTGCAAGGACGTTTCTTGCATTAATTACTGATCCATCGCAACAATCACA ATTATTAGCGGAATTTGAATCAATTTCCAAGAACGGAACACCTCAAATG

		CCAGTGATGACACCTCCAATGATGAATGCAGTTGCTGCAATGCAACCAC ATATGATGATGAGGAATCCAATGCTGAATGCAATGGGTGGAGCATCAC CACGTGTTTCTGGAAAAGGTGGAGCTCCACAAAACGGTGGAATGAATG GGATGTCGATGATGCAGCACATGCAAATGCAAGCATTCCAGGCACAAA TGATGCAAGCTCAAATGTTTCAGCAGATGCCGATGATGATGCCAGGGA TGGTTCCTGGAATGCCACCGGGAATGGCTGGATTGCCTGGAATGATGC CACCGAATAATATGGCAGCTGCAATGCAACATTTTAATGCGTTTCAGCA AATGGCCGCCGCTGCATCGGTAGCAGTTTCTACACCATCAAGGAATCC ATCAACATCTGGAGCCGCTTCTCGAACACGAACTCCATTAACCACATCA GCCACGAATTCACCTCGACCCACCACAGAACCAACGATTAAAGAAGAA GAACCACAGACAATGGAACATGATGTAGCTGCTGCTGTGGCCGCAACT ACTGCTACTGCAACAATCAAACAAGAAGAAAATTCGACAGTGGCGGTTG CTTGA
Y2H	NHR-67 cDNA	ATGATGACCGCTGTCTCTCAAATGTCTGTCCCATCCTCCCGTATCCTTC TTGACGTCGACTGCCGTGTTTGCGAGGACCACTCTTCTGGAAAGCACT ACTCTATCTTCTCTTGCGATGGATGCGCCGGATTCTTCAAGCGCTCTAT CCGCCGCCACCGCCAGTACGTTTGTAAGAACAAGGGATCTCCATCTGA GGGACAATGCAAGGTTGATAAGACTCACCGTAACCAATGCCGCGCCTG CCGTCTTCGCAAGTGCCTTGAGATCGGAATGAACAAGGACGCCGTTCA ACACGAGCGCGGACCACGCAACTCTTCTCTTCGCCGCCAACAAATGAT GTTCGACCACGGATCTTCCCCAAACTCTCCAGAGATGGGATCTGAGTC CGACGCTATCATCCTCCCAACCTCTTCTATGAACCGTGATACCGTCGCT GGAACCGCCGCCCGTATCTTCTTCGCTCTCGTCGGATTCTGCCAAAAC CCACTTAACGGAGTTCCAAAGGAGCGCCAGATGACTATGTTCCAGCAA AACTGGGCTGCCCTCCTCGTCCTTCACGCTACCGAGACCCGCGCTATC ACCTCCAAGCAAATCCGCACCGAGACCATCTCTGGATCTTCTGAGCAA CGTAACGCTGTCGCCAACGCTTTCGAGATCATCGAGCGTCTTCAACTTG ACAACCGCGAGTACATGATGCTTAAGCACTTCACCATGTGGCGCGACA CCCCATCTGCTATCCAAATCGTTTTCCAACTTGCTTCCATCCAAAACTTC ACTCACCGCACTGAGCCAACTCGCTACATCCAGTGCATCAACGCTATC GCTGCCATCCCAACTACCTCTATCATCGACGTCCTTTTCCGTCCATCTA TCGGATCTGCCTCTATGCCACGTCTCATCCAGGATATGTTCAAGCCACC ACAGCAGCCAACCCCAACCTCTCTTTTCCCAATGGCTAATTTCAACCTT AACTTCCTTCTTAAGCAAGAGAAGACTGAGACCGAGGAAGGAGAGGAC ATCGAGGAGGAGGACGACGCTACCTCTTCTAACCAATTCGACGAGAAC TCCTCCACCGACGACCGTTCCGTCGGAGAGCTTGACCCAGTCCAACTC TTCCTTGCCCTCAACTCTTCTACCCAACCATCTTCTGCTTCTTCTCCATC TTCCTCTCGCCCACGTCACTCTATCCGTTCTATCACCGAGCTTCTTTCTA TCCAAGAGGAGGAGTCTGTCAACGTCGAGGAGGTCTAG
Y2H	NHR-67 (ΔIDR) cDNA	ATGATGACCGCTGTCTCTCAAATGTCTGTCCCATCCTCCCGTATCCTTC TTGACGTCGACTGCCGTGTTTGCGAGGACCACTCTTCTGGAAAGCACT ACTCTATCTTCTCTTGCGATGGATGCGCCGGATTCTTCAAGCGCTCTAT CCGCCGCCACCGCCAGTACGTTTGTAAGAACAAGGGATCTCCATCTGA GGGACAATGCAAGGTTGATAAGACTCACCGTAACCAATGCCGCGCCTG CCGTCTTCGCAAGTGCCTTGAGATCGGAATGAACAAGGACGCCGTTCA ACACGAGCGCGGACCACGCAACTCTTCTCTTCGCCGCCAACAAATGAT GTTCGACCACGGATCTTCCCCAAACTCTCCAGAGATGGGATCTGAGTC CGACGCTATCATCCTCCCAACCTCTTCTATGAACCGTGATACCGTCGCT GGAACCGCCGCCCGTATCTTCTTCGCTCTCGTCGGATTCTGCCAAAAC CCACTTAACGGAGTTCCAAAGGAGCGCCAGATGACTATGTTCCAGCAA AACTGGGCTGCCCTCCTCGTCCTTCACGCTACCGAGACCCGCGCTATC ACCTCCAAGCAAATCCGCACCGAGACCATCTCTGGATCTTCTGAGCAA CGTAACGCTGTCGCCAACGCTTTCGAGATCATCGAGCGTCTTCAACTTG

		ACAACCGCGAGTACATGATGCTTAAGCACTTCACCATGTGGCGCGACA CCCCATCTGCTATCCAAATCGTTTTCCAACTTGCTTCCATCCAAAACTTC ACTCACCGCACTGAGCCAACTCGCTACATCCAGTGCATCAACGCTATC GCTGCCATCCCAACTACCTCTATCATCGACGTCCTTTTCCGTCCATCTA TCGGATCTGCCTCTATGCCACGTCTCATCCAGGATATGTTCAAGCCACC ACAGCAGTAG
Y2H	NHR-67 IDR cDNA	ATGCCAACCCCAACCTCTCTTTTCCCAATGGCTAATTTCAACCTTAACTT CCTTCTTAAGCAAGAGAAGACTGAGACCGAGGAAGGAGAGGACATCGA GGAGGAGGACGACGCTACCTCTTCTAACCAATTCGACGAGAACTCCTC CACCGACGACCGTTCCGTCGGAGAGCTTGACCCAGTCCAACTCTTCCT TGCCCTCAACTCTTCTACCCAACCATCTTCTGCTTCTTCTCCATCTTCCT CTCGCCCACGTCACTCTATCCGTTCTATCACCGAGCTTCTTTCTATCCA AGAGGAGGAGTCTGTCAACGTCGAGGAGGTCTAG
Y2H	POP-1 cDNA	ATGATGGCCGACGAAGAGCTCGGCGATGAGGTGAAGGTGTTCCGTCG GGATGAGGATGCTGACGATGATCCAATGATTAGTGGTGAAACGTCAGA ACAACAGTTAGCCGATGATAAAAAGAAGCTGTAATGGAAGCAGAATTA GACGGTGCCGGTCGAAATCCATCGATTGATGTGTTAAAAAGTGCATTTC CAAAAGTCGAACCAATGTCACCATCGTTTCCCGGTTTAATGTCACACTT CAGTCCTGGATACTCGGCAGCTGCTTTACCCATGTTTATGCCTCTATTT ATGAATCCATACGCAGCAGCACTACGATCACCAAGCCTGATGTTTCCAA TGGGAGCAATGAGCCCCACATTTCCAATGTTCCCGCCAAGTCCTGTCTA TGGAGCAGCAATCGCTGCGGCAGCCGCCAAACAACACTTTGAGAATAT GGCTCCACTGAACATGCGAGCCGGTCATCCAATGAATCAGATGGGAAT GCCACCATACATGCATCCATCATCAATGGCTCCACAGAATGTCGATCGA AGGGCTCAAGGAGGTGGAAAAGCGAAGAAGGATGATCATGTGAAGAAG CCATTAAACGCGTTCATGTGGTTTATGAAGGAAAATCGAAAAGCACTGC TGGAAGAGATTGGAAATAATGAGAAACAGAGTGCAGAGTTGAATAAAGA GCTTGGAAAGAGGTGGCATGATTTGTCGAAGGAAGAACAGGCGAAATA TTTTGAAATGGCAAAGAAGGATAAGGAAACACACAAGGAACGGTATCC GGAGTGGTCGGCGCGGGAAAATTATGCGGTTAATAAGAAAAAGACGAA GAAACGAAGGGATAAGAGTATTCCATCGGAGAACAACGATCAGAAGAA GTGCCGAGCCAGATTCGGAGTTAACAACACAGAAATGTGGTGTAAATTC TGCAAGCGGAAGAAGAAGTGCGAGTACGCAACTGATCGTTCGGGCGG TTCCGATATAACTGACAGTCAGGATGGACGAGGTACAAGTGGTGCGTA TAGCAGTAGCTCGGAGAGCCCATCACCAAAGGCAAACGCTGGAATTGC ACTGACCACACAGCAGCAGCAAGCAGCAATGATGCATACGATGTTGAT GCAAATGCGTCTAGGATCGACGACGGGAGCATCGACGCACGTTCCATC ACCACTGGCGTCCTCGTCGGCAGGCAGGAGTCCGCTGGATGCGAACG CGTCGGATTCGGAATCTGATGTTGAGGAGGAGGAAGACGAGCAGATTG ATCCGACGGTTATGCAGCAGACACATGATATGCTTATGCAGGAATCGAT GTGTACTATTTAA
Y2H	UNC-37 cDNA	ATGAAGGCATCGTATCTGGAAACCCTCGATCGAATCAAAGACGAGCAT GCGGAAATGTCGAAGCATGTCAACCAGCAACGATCGGATATCGAAAAG GTTGCGTTAGAGAAAGAAAATATGAATAGATCGTATATGACGTACGCTG AAGTATCAAATACTCTTCGTAGCGATCTTCGAAAAGCAGAAGAAATCAA CAAACGCCTTCAGGAATTCATCATGCAATCGTTGGCTCCTCAACTATCT CAAGATAATCAAGCAAACTGCCTCGCCGCACTGGAAGCTTTCAAAACC GCATCACCACGTGAAAATGGAAATGGAGCACCAGCACTTCCACCAGGT TTTCCTCCCGGTGCAGCTGGAATGCTTGGAATGATGCCAAATATGCCAT TCGGAATGAGCCCTGCAATGAGTCAACTATTCAATCAATTCGCATCACC ACATGTCAACGGTGGAGATGGAGCCGGTGGAAGTTCTGGAGGTGCAA GTGAAGCGAAAAAGGCCAAGCTGGAAGATCCAGATGATGGAGAACTTG

| | | AGATTGATGTCACAAATGATGATCATCCAAGTACTGCATCAAATGGAGG
CGCAGCTAATAAAAATGGAAGAGATAGCACAAATAGTGTTGCCTCATCG
GGAGCATCTACACCGAGCATCGCATCAAATTCTAGAGCAAGACAACAA
CAGCAGCCACTCGCTGGGCTTCAAGGATTGGAGCAAATGAACTTTTTA
GCTGGATTCAATCCAAATCTTCTCCGACAAGCTAGTGCTGCTGGAGGAT
TCAACTTTTTGAATGATCCCCATGCACAGGCCCGTCTTGCAGCTGCCAT
CGGGCAAATCGGTAGCAGACCGGCCTATTCATTCAAAATTGTGGACGG
TGGAGTTCCAACACCTACATCCTTCCCTCCAGATGCACAGAAAGGTCCC
GGAATTCCGACAGGTCTCAAGAAGAAATGGAATTAAATCACGGAGAA
GTAGTTTGTGCGGCCACAATTTCCCGTGATAACAGTCGTGTTTATACTG
GTGGAAAAGGATGTGTTAAGATCTGGGATGTCAAAGAATCAGATATTTC
TGGCGCAACGGTTGTGAATCGGCCTCCAATTGCATCCTTGGATTGTCTC
AAAGAGAACTATATTAGATCTTGTAAACTCTTTGAAGATGGAAATACACT
TCTGATTGGAGGAGAAGCCAGTACTGTTGCTCTTTGGGATCTTACAACT
GAAACAAAAACTTTGGACTTGGAAACTGATTCACAAGCATGCTACGCGT
TGGCGATGTCACCAGATGAGAAATTATTGTTCGCATGTCTTGCTGATGG
AAATATCCTCATTTATGATATCCATAATAAGGTGAAAGTTGGAACCCTAC
CTGGACATCAAGATGGAGCATCATGTCTCGATCTCTCAAAAGATGGTAC
AAAGCTCTGGTCAGGTGGTCTTGACAATTCTGTTAGATGCTGGGATCTT
GCACAACGAAAGGAAGTTGCGAAGCATGATTTTGCGAGCCAAGTTTTCT
CTCTTGGATGTTGTCCAAATGATGAATGGGTGGCTGTTGGTATGGAGAA
TAATTATGTCGAGGTTCTGTCAACAACTGGAAAAGAAAGTATCAATTGA
CACAACACGAATCATGCGTCCTTTCGCTCAAATTTGCGCATTCCGGAAA
GTTCTTCATTTCAACTGGAAAGGACAATGCTCTCAACGCCTGGCGTACT
CCATACGGAGCATCACTCTTTCAACTAAAAGAGAACAGCTCCGTTCTCT
CATGTGACATCTCATTCGACGACTCACTCATTGTCACTGGATCAGGGGA
GAAGAAGGCAACTCTCTATGCAGTTGAATATTAA |

References

Adikes RC, Kohrman AQ, Martinez MAQ, Palmisano NJ, Smith JJ, Medwig-Kinney TN, Min M, Sallee MD, Ahmed OB, Kim N, Liu S, Morabito RD, Weeks N, Zhao Q, Zhang W, Feldman JL, Barkoulas M, Pani AM, Spencer SL, Martin BL, Matus DQ. 2020. Visualizing the metazoan proliferation-quiescence decision in vivo. *Elife* **9**:1–74. doi:10.7554/eLife.63265

Ashley G, Duong T, Levenson MT, Martinez MAQ, Johnsen LC, Hibshman JD, Saeger HN, Palmisano NJ, Doonan R, Martinez-Mendez R, Davidson B, Zhang W, Ragle JM, Medwig-Kinney TN, Sirota SS, Goldstein B, Matus DQ, Dickinson DJ, Reiner DJ, Ward JD. 2021. An expanded auxin-inducible degron toolkit for Caenorhabditis elegans . *Genetics*. doi:10.1093/genetics/iyab006

Attner MA, Keil W, Benavidez JM, Greenwald I. 2019. HLH-2/E2A Expression Links Stochastic and Deterministic Elements of a Cell Fate Decision during C. elegans Gonadogenesis. *Curr Biol* **29**:1–7. doi:10.1016/j.cub.2019.07.062

Bekas KN, Phillips BT. 2022. unc-37/Groucho and lsy-22/AES repress Wnt target genes in C. elegans asymmetric cell divisions. *bioRxiv*.

Benavidez JM, Kim JH, Greenwald I. 2022. Influences of HLH-2 stability on anchor cell fate specification during Caenorhabditis elegans gonadogenesis. *G3: Genes|Genomes|Genetics*.

Bodofsky S, Liberatore K, Pioppo L, Lapadula D, Thompson L, Birnbaum S, McClung G, Kartik A, Clever S, Wightman B. 2018. A tissue-specific enhancer of the C. elegans

nhr-67/tailless gene drives coordinated expression in uterine stem cells and the differentiated anchor cell. *Gene Expr Patterns* **30**:71–81. doi:10.1016/j.gep.2018.10.003

Boija A, Klein IA, Sabari BR, Dall'Agnese A, Coffey EL, Zamudio A V., Li CH, Shrinivas K, Manteiga JC, Hannett NM, Abraham BJ, Afeyan LK, Guo YE, Rimel JK, Fant CB, Schuijers J, Lee TI, Taatjes DJ, Young RA. 2018. Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell* **175**:1842-1855.e16. doi:10.1016/j.cell.2018.10.042

Bolte S, Cordelières FP. 2006. A guided tour into subcellular colocalization analysis in light microscopy. *J Microsc* **224**:213–232. doi:10.1111/j.1365-2818.2006.01706.x

Boxem M, Maliga Z, Klitgord N, Li N, Lemmens I, Mana M, de Lichtervelde L, Mul JD, van de Peut D, Devos M, Simonis N, Yildirim MA, Cokol M, Kao HL, de Smet AS, Wang H, Schlaitz AL, Hao T, Milstein S, Fan C, Tipsword M, Drew K, Galli M, Rhrissorrakrai K, Drechsel D, Koller D, Roth FP, Iakoucheva LM, Dunker AK, Bonneau R, Gunsalus KC, Hill DE, Piano F, Tavernier J, van den Heuvel S, Hyman AA, Vidal M. 2008. A Protein Domain-Based Interactome Network for C. elegans Early Embryogenesis. *Cell* **134**:534–545. doi:10.1016/j.cell.2008.07.009

Brangwynne CP, Eckmann CR, Courson DS, Rybarska A, Hoege C, Gharakhani J, Jülicher F, Hyman AA. 2009. Germline P Granules Are Liquid Droplets That Localize by Controlled Dissolution/Condensation. *Science (80-)* **324**:1729–1732.

Brenner S. 1974. The genetics of *Caenorabditis elegans*. *Genetics* **77**:71–94. doi:10.1016/S0047-2484(78)80101-8

Brumbaugh J, Stefano B Di, Hochedlinger K. 2019. Reprogramming : identifying the mechanisms that safeguard cell identity. *Development* **146**:1–17. doi:10.1242/dev.182170

Calvo D, Victor M, Gay F, Sui G, Po-Shan Luke M, Dufourcq P, Wen G, Maduro M, Rothman J, Shi Y. 2001. A POP-1 repressor complex restricts inappropriate cell type-specific gene transcription during Caenorhabditis elegans embryogenesis. *EMBO J* **20**:7197–7208. doi:10.1093/emboj/20.24.7197

Cho WK, Spille JH, Hecht M, Lee C, Li C, Grube V, Cisse II. 2018. Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science (80-)* **361**:412–415. doi:10.1126/science.aar4199

Chong S, Dugast-Darzacq C, Liu Z, Dong P, Dailey GM, Cattoglio C, Heckert A, Banala S, Lavis L, Darzacq X, Tjian R. 2018. Imaging dynamic and selective low-complexity domain interactions that control gene transcription. *Science (80-)* **361**. doi:10.1126/science.aar2555

Davis RL, Weintraub H, Lassar AB. 1987. Expression of a single transfected cDNA converts fibroblasts to myoblasts. *Cell* **51**:987–1000. doi:10.1016/0092-8674(87)90585-X

Dickinson DJ, Goldstein B. 2016. CRISPR-based methods for caenorhabditis elegans genome engineering. *Genetics* **202**:885–901. doi:10.1534/genetics.115.182162

Dickinson DJ, Pani AM, Heppert JK, Higgins CD, Goldstein B. 2015. Streamlined genome engineering with a self-excising drug selection cassette. *Genetics* **200**:1035–1049. doi:10.1534/genetics.115.178335

Flowers EB, Poole RJ, Tursun B, Bashllari E, Pe'er I, Hobert O. 2010. The Groucho ortholog UNC-37 interacts with the short Groucho-like protein LSY-22 to control

developmental decisions in C. elegans. *Development* **137**:1799–1805. doi:10.1242/dev.046219

Frøkjær-Jensen C, Davis MW, Ailion M, Jorgensen EM. 2013. Improved Mos1-mediated transgenesis in C. elegans. *Nat Meth* **9**:117–118. doi:10.1038/mp.2011.182.doi

Fukushige T, Krause M. 2005. The myogenic potency of HLH-1 reveals wide-spread developmental plasticity in early C. elegans embryos. *Development* **132**:1795–1805. doi:10.1242/dev.01774

Ghanta KS, Mello CC. 2020. Melting dsDNA donor molecules greatly improves precision genome editing in caenorhabditis elegans. *Genetics* **216**:643–650. doi:10.1534/genetics.120.303564

Gilleard JS, McGhee JD. 2001. Activation of Hypodermal Differentiation in the Caenorhabditis elegans Embryo by GATA Transcription Factors ELT-1 and ELT-3 . *Mol Cell Biol* **21**:2533–2544. doi:10.1128/mcb.21.7.2533-2544.2001

Gómez-Saldivar G, Osuna-Luque J, Semple JI, Glauser DA, Jarriault S, Meister P. 2020. Tissue-Specific Transcription Footprinting Using RNA. *Genetics* **216**:931–945.

Green JL, Inoue T, Sternberg PW. 2008. Opposing Wnt pathways orient cell polarity during organogenesis. *Cell* **134**:646–656. doi:10.1016/j.cell.2008.06.026.Opposing

Greenwald IS, Sternberg PW, Robert Horvitz H. 1983. The lin-12 locus specifies cell fates in Caenorhabditis elegans. *Cell* **34**:435–444. doi:10.1016/0092-8674(83)90377-X

Hajduskova M, Baytek G, Kolundzic E, Gosdschan A. 2019. MRG-1 / MRG15 Is a Barrier for Germ Cell to Neuron reprogramming. *Genetics* **211**:121–139.

Hanahan D. 2022. Hallmarks of Cancer: New Dimensions. *Cancer Discov* 31–47. doi:10.1158/2159-8290.CD-21-1059

Hanahan D, Weinberg RA. 2000. The hallmarks of cancer. *Cell* **100**:57–70. doi:10.1007/s00262-010-0968-0

Harris TW, Arnaboldi V, Cain S, Chan J, Chen WJ, Cho J, Davis P, Gao S, Grove CA, Kishore R, Lee RYN, Muller HM, Nakamura C, Nuin P, Paulini M, Raciti D, Rodgers FH, Russell M, Schindelman G, Auken K V., Wang Q, Williams G, Wright AJ, Yook K, Howe KL, Schedl T, Stein L, Sternberg PW. 2020. WormBase: A Modern Model Organism Information Resource. *Nucleic Acids Res* **48**:D762–D767. doi:10.1093/nar/gkz920

Hills-Muckey K, Martinez MAQ, Stec N, Hebbar S, Saldanha J, Medwig-Kinney TN, Moore FEQ, Ivanova M, Moraro A, Jordan D, Moss EG, Ercan S, Zinovyeva AY, Matus DQ, Christopher M. 2021. An engineered, orthogonal auxin analog/AtTIR1(F79G) pairing improves both specificity and efficacy of the auxin degradation system in Caenorhabditis elegans. *Genetics* iyab174. doi:https://doi.org/10.1093/genetics/iyab174

Hoek KS, Eichhoff OM, Schlegel NC, Döbbeling U, Kobert N, Schaerer L, Hemmi S, Dummer R. 2008. In vivo switching of human melanoma cells between proliferative and invasive states. *Cancer Res* **68**:650–656. doi:10.1158/0008-5472.CAN-07-2491

Horner MA, Quintin S, Domeier ME, Kimble J, Labouesse M, Mango SE. 1998. pha-4, an HNF-3 homolog, specifies pharyngeal organ identity in Caenorhabditis elegans. *Genes Dev* **12**:1947–1952. doi:10.1101/gad.12.13.1947

Huang G, Jesus B De, Koh A, Blanco S, Rettmann A, Demott E, Sylvester M, Meng C,
Waterland S, Rhodes A, Alicea P, Flynn A, Dickinson DJ. 2021. Improved CRISPR
/ Cas9 knock-in efficiency via the self-excising cassette (SEC) selection method in
C . elegans. *Micropublication Biol*.
Irgen-Gioro S, Walling V, Chong S. 2022. Fixation can change the appearance of phase
separation in living cells. *bioRxiv*.
Jarriault S, Schwab Y, Greenwald I. 2008. A Caenorhabditis elegans model for
epithelial-neuronal transdifferentiation. *Proc Natl Acad Sci* **105**:3790–3795.
doi:10.1073/pnas.0712159105
Jiang H, Xu Z, Zhong P, Ren Y, Liang G, Schilling HA, Hu Z, Zhang Y, Wang X, Chen
S, Yan Z, Feng J. 2015. Cell cycle and p53 gate the direct conversion of human
fibroblasts to dopaminergic neurons. *Nat Commun* **6**. doi:10.1038/ncomms10100
Jin Y, Hoskins R, Horvitz HR. 1994. Control of type-D GABAergic neuron differentiation
by C. elegans UNC-30 homeodomain protein. *Nature* **372**:780–783.
doi:10.1038/372780a0
Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool
K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ,
Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S,
Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T,
Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis
D. 2021. Highly accurate protein structure prediction with AlphaFold. *Nature*
596:583–589. doi:10.1038/s41586-021-03819-2
Kagias K, Ahier A, Fischer N, Jarriault S. 2012. Members of the NODE (Nanog and
Oct4-associated deacetylase) complex and SOX-2 promote the initiation of a
natural cellular reprogramming event in vivo. *Proc Natl Acad Sci U S A* **109**:6596–
6601. doi:10.1073/pnas.1117031109
Karp X, Greenwald I. 2003. Post-transcriptional regulation of the E/Daughterless
ortholog HLH-2, negative feedback, and birth order bias during the AC/VU decision
in C. elegans. *Genes Dev* **17**:3100–3111. doi:10.1101/gad.1160803
Katsanos D, Barkoulas M. 2022. Targeted DamID in C . elegans reveals a direct role for
LIN-22 and NHR-25 in antagonizing the epidermal stem cell fate. *Sci Adv*
8:eabk3141.
Keeley DP, Hastie E, Jayadev R, Kelley LC, Chi Q, Payne SG, Jeger JL, Hoffman BD,
Sherwood DR. 2020. Comprehensive Endogenous Tagging of Basement
Membrane Components Reveals Dynamic Movement within the Matrix Scaffolding.
Dev Cell **54**:60-74.e7. doi:10.1016/j.devcel.2020.05.022
Kelley LC, Wang Z, Hagedorn EJ, Wang L, Shen W, Lei S, Johnson SA, Sherwood DR.
2017. Live-cell confocal microscopy and quantitative 4D image analysis of anchor-
cell invasion through the basement membrane in Caenorhabditis elegans. *Nat*
Protoc **12**:2081–2096. doi:10.1038/nprot.2017.093
Kiefer JC, Smith PA, Mango SE. 2007. PHA-4/FoxA cooperates with TAM-1/TRIM to
regulate cell fate restriction in the C. elegans foregut. *Dev Biol* **303**:611–624.
doi:10.1016/j.ydbio.2006.11.042
Kimble J, Hirsh D. 1979. The postembryonic cell lineages of the hermaphrodite and
male gonads in Caenorhabditis elegans. *Dev Biol* **70**:396–417. doi:10.1016/0012-
1606(79)90035-6

Kohrman AQ, Matus DQ. 2017. Divide or Conquer: Cell Cycle Regulation of Invasive
Behavior. *Trends Cell Biol*. doi:10.1016/j.tcb.2016.08.003
Kolundzic E, Ofenbauer A, Bulut SI, Uyar B, Baytek G, Sommermeier A, Seelk S, He M,
Hirsekorn A, Vucicevic D, Akalin A, Diecke S, Lacadie SA, Tursun B. 2018. FACT
Sets a Barrier for Cell Fate Reprogramming in Caenorhabditis elegans and Human
Cells. *Dev Cell* **46**:611-626.e12. doi:10.1016/j.devcel.2018.07.006
Kurashina M, Wang J, Lin J, Lee KK, Johal A, Mizumoto K. 2021. Sustained expression
of unc-4 homeobox gene and unc-37/groucho in postmitotic neurons specifies the
spatial organization of the cholinergic synapses in c. Elegans. *Elife* **10**:1–26.
doi:10.7554/eLife.66011
Larson AG, Elnatan D, Keenen MM, Trnka MJ, Johnston JB, Burlingame AL, Agard DA,
Redding S, Narlikar GJ. 2017. Liquid droplet formation by HP1α suggests a role for
phase separation in heterochromatin. *Nature* **547**:236–240.
doi:10.1038/nature22822
Leyva-Díaz E, Hobert O. 2019. Transcription factor autoregulation is required for
acquisition and maintenance of neuronal identity. *Development* **146**.
doi:10.1242/dev.177378
Li S, Armstrong CM, Bertin N, Ge H, Milstein S, Boxem M, Vidalain PO, Han JDJ,
Chesneau A, Hao T, Goldberg DS, Li N, Martinez M, Rual JF, Lamesch P, Xu L,
Tewari M, Wong SL, Zhang L V., Berriz GF, Jacotot L, Vaglio P, Reboul J,
Hirozane-Kishikawa T, Li Q, Gabel HW, Elewa A, Baumgartner B, Rose DJ, Yu H,
Bosak S, Sequerra R, Fraser A, Mango SE, Saxton WM, Strome S, Van Den
Heuvel S v., Piano F, Vandenhaute J, Sardet C, Gerstein M, Doucette-Stamm L,
Gunsalus KC, Harper JW, Cusick ME, Roth FP, Hill DE, Vidal M. 2004. A Map of
the Interactome Network of the Metazoan C. elegans. *Science (80-)* **303**:540–543.
doi:10.1126/science.1091403
Liang B, Wang J. 2020. EVI1 in leukemia and solid tumors. *Cancers (Basel)* **12**:1–17.
doi:10.3390/cancers12092667
Lim B, Levine MS. 2021. Enhancer-promoter communication: hubs or loops? *Curr Opin*
Genet Dev **67**:5–9. doi:10.1016/j.gde.2020.10.001
Ma X, Zhao Z, Xiao L, Xu W, Kou Y, Zhang Y, Wu G, Wang Y, Du Z. 2021. A 4D single-
cell protein atlas of transcription factors delineates spatiotemporal patterning during
embryogenesis. *Nat Methods* **18**:893–902. doi:10.1038/s41592-021-01216-1
Maduro MF, Lin R, Rothman JH. 2002. Dynamics of a developmental switch: Recursive
intracellular and intranuclear redistribution of Caenorhabditis elegans POP-1
parallels Wnt-inhibited transcriptional repression. *Dev Biol* **248**:128–142.
doi:10.1006/dbio.2002.0721
Martinez M, Matus D. 2020. Auxin-mediated Protein Degradation in Caenorhabditis
elegans. *Bio-Protocol* **10**:1–13. doi:10.21769/bioprotoc.3589
Martinez MAQ, Kinney BA, Medwig-Kinney TN, Ashley G, Ragle JM, Johnson L,
Aguilera J, Hammell CM, Ward JD, Matus DQ. 2020. Rapid Degradation of
Caenorhabditis elegans Proteins at Single-Cell Resolution with a Synthetic Auxin.
G3 Genes, Genomes, Genet **10**:267–280. doi:10.1534/g3.119.400781
Matus DQ, Li X-Y, Durbin S, Agarwal D, Chi Q, Weiss SJ, Sherwood DR. 2010. In vivo
identification of regulators of cell invasion across basement membranes. *Sci Signal*
3:ra35. doi:10.1126/scisignal.2000654

Matus DQ, Lohmer LL, Kelley LC, Schindler AJ, Kohrman AQ, Barkoulas M, Zhang W, Chi Q, Sherwood DR. 2015. Invasive Cell Fate Requires G1 Cell-Cycle Arrest and Histone Deacetylase-Mediated Changes in Gene Expression. *Dev Cell* **35**:162–174. doi:10.1016/j.devcel.2015.10.002

McKim KS, Peters K, Rose AM. 1993. Two Types of Sites Required for Meiotic Chromosome Pairing in Caenorhabditis elegans. *Genetics* **134**:749–768.

Medwig-Kinney TN, Palmisano NJ, Matus DQ. 2021. Deletion of a putative HDA-1 binding site in the hlh-2 promoter eliminates expression in C. elegans dorsal uterine cells. *Micropublication Biol* **1**.

Medwig-Kinney TN, Sirota SS, Gibney T V., Pani AM, Matus DQ. 2022. An in vivo toolkit to visualize endogenous LAG-2/Delta and LIN-12/Notch signaling in C. elegans. *microPublication Biol*.

Medwig-Kinney TN, Smith JJ, Palmisano NJ, Tank S, Zhang W, Matus DQ. 2020. A developmental gene regulatory network for C. elegans anchor cell invasion. *Development* **147**:dev185850. doi:https://doi.org/10.1101/691337

Medwig TN, Matus DQ. 2017. Breaking down barriers: the evolution of cell invasion. *Curr Opin Genet Dev* **47**:33–40. doi:10.1016/j.gde.2017.08.003

Milde-Langosch K. 2005. The Fos family of transcription factors and their role in tumourigenesis. *Eur J Cancer* **41**:2449–2461. doi:10.1016/j.ejca.2005.08.008

Mondal C, Gacha-Garay MJ, Larkin K, Adikes R, Di Martino J, Chien C-C, Fraser M, Eni-aganga I, Agullo-Pascual E, Ozbek U, Naba A, Gaitas A, Fu T-M, Upadhyayula S, Betzig E, Matus D, Martin BL, Bravo-Cordero JJ. 2021. A Proliferative to Invasive Switch is Mediated by srGAP1 Downregulation Through the Activation of TGFβ2 Signaling. *SSRN Electron J* **40**:111358. doi:https://doi.org/10.1016/j.celrep.2022.111358 ll

Nelson AT, Wang Y, Nelson ER. 2021. TLX, an Orphan Nuclear Receptor with Emerging Roles in Physiology and Disease. *Endocrinol (United States)* **162**:1–13. doi:10.1210/endocr/bqab184

Pani AM, Gibney T V., Medwig-Kinney TN, Matus DQ, Goldstein B. 2022. A new toolkit to visualize and perturb endogenous LIN-12/Notch signaling. *Micropublication Biol*.

Patel T, Hobert O. 2017. Coordinated control of terminal differentiation and restriction of cellular plasticity. *Elife* **6**:1–26. doi:10.7554/elife.24100

Patel T, Tursun B, Rahe DP, Hobert O. 2012. Removal of Polycomb Repressive Complex 2 Makes C. elegans Germ Cells Susceptible to Direct Conversion into Specific Somatic Cell Types. *Cell Rep* **2**:1178–1186. doi:10.1016/j.celrep.2012.09.020

Peng QY, Zhang QF. 2006. Precise positions of Phoebe determined with CCD image-overlapping calibration. *Mon Not R Astron Soc* **366**:208–212. doi:10.1186/1471-2105-7-208

Pflugrad A, Meir JYJ, Barnes TM, Miller DM. 1997. The Groucho-like transcription factor UNC-37 functions with the neural specificity gene unc-4 to govern motor neuron identity in C. elegans. *Development* **124**:1699–1709. doi:10.1242/dev.124.9.1699

Phillips BT, Kidd AR, King R, Hardin J, Kimble J. 2007. Reciprocal asymmetry of SYS-1/β-catenin and POP-1/TCF controls asymmetric divisions in Caenorhabditis elegans. *Proc Natl Acad Sci U S A* **104**:3231–3236. doi:10.1073/pnas.0611507104

Phillips BT, Kimble J. 2009. A New Look at TCF and β-Catenin through the Lens of a

Divergent C. elegans Wnt Pathway. *Dev Cell* **17**:27–34.
doi:10.1016/j.devcel.2009.07.002

Porta-de-la-Riva M, Fontrodona L, Villanueva A, Cerón J. 2012. Basic Caenorhabditis
elegans Methods: Synchronization and Observation. *J Vis Exp* 1–9.
doi:10.3791/4019

Putnam A, Cassani M, Smith J, Seydoux G. 2019. A gel phase promotes condensation
of liquid P granules in Caenorhabditis elegans embryos. *Nat Struct Mol Biol* **26**.

Quintin S, Michaux G, McMahon L, Gansmuller A, Labouesse M. 2001. The
Caenorhabditis elegans gene lin-26 can trigger epithelial differentiation without
conferring tissue specificity. *Dev Biol* **235**:410–421. doi:10.1006/dbio.2001.0294

Rahe DP, Hobert O. 2019. Restriction of Cellular Plasticity of Differentiated Cells
Mediated by Chromatin Modifiers, Transcription Factors and Protein Kinases.
G3: Genes|Genomes|Genetics **9**:2287–2302.
doi:10.1534/g3.119.400328

Reece-Hoyes JS, Deplancke B, Shingles J, Grove CA, Hope IA, Walhout AJM. 2005. A
compendium of Caenorhabditis elegans regulatory transcription factors: A resource
for mapping transcription regulatory networks. *Genome Biol* **6**. doi:10.1186/gb-
2005-6-13-r110

Reece-Hoyes JS, Walhout AJM. 2018. High-efficiency yeast transformation. *Cold
Spring Harb Protoc* **2018**:563–568. doi:10.1101/pdb.prot094995

Richard JP, Zuryn S, Fischer N, Pavet V, Vaucamps N, Jarriault S. 2011. Direct in vivo
cellular reprogramming involves transition through discrete, non-pluripotent steps. *J
Cell Sci* **124**:e1–e1. doi:10.1242/jcs.090043

Riddle MR, Spickard EA, Jevince A, Nguyen KCQ, Hall DH, Joshi PM, Rothman JH.
2016. Transorganogenesis and transdifferentiation in C. elegans are dependent on
differentiated cell identity. *Dev Biol* **420**:136–147. doi:10.1016/j.ydbio.2016.09.020

Riddle MR, Weintraub A, Nguyen KCQ, Hall DH, Rothman JH. 2013.
Transdifferentiation and remodeling of post-embryonic C. elegans cells by a single
transcription factor. *Development* **140**:4844–4849. doi:10.1242/dev.103010

Riva C, Hajduskova M, Gally C, Suman SK, Ahier A, Jarriault S. 2022. A natural
transdifferentiation event involving mitosis is empowered by integrating signaling
inputs with conserved plasticity factors. *Cell Rep* **40**:111365.
doi:10.1016/j.celrep.2022.111365

Rothman J, Jarriault S. 2019. Developmental Plasticity and Cellular Reprogramming.
Genetics **213**:723–757.

Rual JF, Ceron J, Koreth J, Hao T, Nicot AS, Hirozane-Kishikawa T, Vandenhaute J,
Orkin SH, Hill DE, van den Heuvel S, Vidal M. 2004. Toward improving
Caenorhabditis elegans phenome mapping with an ORFeome-based RNAi library.
Genome Res **14**:2162–2168. doi:10.1101/gr.2505604

Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ,
Hannett NM, Zamudio A V., Manteiga JC, Li CH, Guo YE, Day DS, Schuijers J,
Vasile E, Malik S, Hnisz D, Lee TI, Cisse II, Roeder RG, Sharp PA, Chakraborty
AK, Young RA. 2018. Coactivator condensation at super-enhancers links phase
separation and gene control. *Science (80-)* **361**. doi:10.1126/science.aar3958

Sallee MD, Aydin T, Greenwald I. 2015. Influences of LIN-12/Notch and POP-1/TCF on
the Robustness of Ventral Uterine Cell Fate Specification in Caenorhabditis

elegans Gonadogenesis. *G3: Genes|Genomes|Genetics* **5**:2775–2782.
doi:10.1534/g3.115.022608

Sallee MD, Greenwald I. 2015. Dimerization-driven degradation of C. elegans and
human E proteins. *Genes Dev* **29**:1356–1361. doi:10.1101/gad.261917.115

Sallee MD, Littleford HE, Greenwald I. 2017. A bHLH Code for Sexually Dimorphic
Form and Function of the C. elegans Somatic Gonad. *Curr Biol* **27**:1853-1860.e5.
doi:10.1016/j.cub.2017.05.059

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
analysis. *Nat Methods* **9**:676–82. doi:10.1038/nmeth.2019

Seydoux G, Greenwald I. 1989. Cell autonomy of lin-12 function in a cell fate decision in
C. elegans. *Cell* **57**:1237–1245. doi:10.1016/0092-8674(89)90060-3

Seydoux G, Schedl T, Greenwald I. 1990. Cell-cell interactions prevent a potential
inductive interaction between soma and germline in C. elegans. *Cell* **61**:939–951.
doi:10.1016/0092-8674(90)90060-R

Sherwood DR, Sternberg PW. 2003. Anchor cell invasion into the vulval epithelium in C.
elegans. *Dev Cell* **5**:21–31. doi:10.1016/S1534-5807(03)00168-0

Shetty P, Lo MC, Robertson SM, Lin R. 2005. C. elegans TCF protein, POP-1, converts
from repressor to activator as a result of Wnt-induced lowering of nuclear levels.
Dev Biol **285**:584–592. doi:10.1016/j.ydbio.2005.07.008

Siegfried K, Kimble J. 2002. POP-1 controls axis formation during early gonadogenesis
in C. elegans. *Development* **453**:443–453.

Siegfried KR, Kidd AR, Chesney MA, Kimble J. 2004. The sys-1 and sys-3 Genes
Cooperate with Wnt Signaling to Establish the Proximal-Distal Axis of the
Caenorhabditis elegans Gonad. *Genetics* **166**:171–186.
doi:10.1534/genetics.166.1.171

Simonis N, Rual JF, Carvunis AR, Tasan M, Lemmens I, Hirozane-Kishikawa T, Hao T,
Sahalie JM, Venkatesan K, Gebreab F, Cevik S, Klitgord N, Fan C, Braun P, Li N,
Ayivi-Guedehoussou N, Dann E, Bertin N, Szeto D, Dricot A, Yildirim MA, Lin C, de
Smet AS, Kao HL, Simon C, Smolyar A, Ahn JS, Tewari M, Boxem M, Milstein S,
Yu H, Dreze M, Vandenhaute J, Gunsalus KC, Cusick ME, Hill DE, Tavernier J,
Roth FP, Vidal M. 2009. Empirically controlled mapping of the Caenorhabditis
elegans protein-protein interactome network. *Nat Methods* **6**:47–54.
doi:10.1038/nmeth.1279

Smith JJ, Xiao Y, Parsan N, Medwig-Kinney TN, Martinez MAQ, Moore FEQ, Palmisano
NJ, Kohrman AQ, Chandhok Delos Reyes M, Adikes RC, Liu S, Bracht SA, Zhang
W, Wen K, Kratsios P, Matus DQ. 2022. The SWI/SNF chromatin remodeling
assemblies BAF and PBAF differentially regulate cell cycle exit and cellular
invasion in vivo, PLOS Genetics. doi:10.1371/journal.pgen.1009981

Spencer SL, Cappell SD, Tsai FC, Overton KW, Wang CL, Meyer T. 2013. The
proliferation-quiescence decision is controlled by a bifurcation in CDK2 activity at
mitotic exit. *Cell* **155**:369–383. doi:10.1016/j.cell.2013.08.062

Strom AR, Emelyanov A V., Mir M, Fyodorov D V., Darzacq X, Karpen GH. 2017. Phase
separation drives heterochromatin domain formation. *Nature* **547**:241–245.
doi:10.1038/nature22989

Sturm Á, Saskoi É, Tibor K, Weinhardt N, Vellai T. 2018. Highly efficient RNAi and
Cas9-based auto-cloning systems for C. elegans research. *Nucleic Acids Res*
46:e105. doi:10.1093/nar/gky516
Sulston JE, Horvitz HR. 1977. Post-embryonic cell lineages of the nematode,
Caenorhabditis elegans. *Dev Biol* **56**:110–156. doi:10.1016/0012-1606(77)90158-0
Treen N, Shimobayashi SF, Eeftens J, Brangwynne CP, Levine M. 2021. Properties of
repression condensates in living Ciona embryos. *Nat Commun* **12**:1–9.
doi:10.1038/s41467-021-21606-5
Tursun B, Patel T, Kratsios P, Hobert O. 2011. Direct Conversion of C. elegans Germ
Cells into Specific Neuron Types. *Science (80-)* **331**:304–309.
van der Horst SEM, Cravo J, Woollard A, Teapal J, Van den Heuvel S. 2019. C .
elegans Runx / CBFβ suppresses POP-1 (TCF) to convert asymmetric to
proliferative division of stem cell-like seam cells. *Development* **146**:1–34.
doi:10.1242/dev.180034
Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe
O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper
J, Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N, Kohli P,
Kleywegt G, Birney E, Hassabis D, Velankar S. 2022. AlphaFold Protein Structure
Database: massively expanding the structural coverage of protein-sequence space
with high-accuracy models. *Nucleic Acids Res* **50**:D439–D444.
doi:10.1093/nar/gkab1061
Verghese E, Schocken J, Jacob S, Wimer AM, Royce R, Nesmith JE, Baer GM, Clever
S, McCain E, Lakowski B, Wightman B. 2011. The tailless ortholog nhr-67 functions
in the development of the C. elegans ventral uterus. *Dev Biol* **356**:516–528.
doi:10.1016/j.ydbio.2011.06.007
Wang LH, Baker NE. 2015. E Proteins and ID Proteins: Helix-Loop-Helix Partners in
Development and Disease. *Dev Cell* **35**:269–280. doi:10.1016/j.devcel.2015.10.019
Wang S, Tang NH, Lara-Gonzalez P, Zhao Z, Cheerambathur DK, Prevo B, Chisholm
AD, Desai A, Oegema K. 2017. A toolkit for GFP-mediated tissue-specific protein
degradation in C. elegans. *Development* **144**:2694–2701. doi:10.1242/dev.150094
Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York.
Wilkinson HA, Fitzgerald K, Greenwald I. 1994. Reciprocal changes in expression of the
receptor lin-12 and its ligand lag-2 prior to commitment in a C. elegans cell fate
decision. *Cell* **79**:1187–1198. doi:10.1016/0092-8674(94)90010-8
Zacharias AL, Walton T, Preston E, Murray JI. 2015. Quantitative Differences in Nuclear
β-catenin and TCF Pattern Embryonic Cells in C. elegans. *PLoS Genet* **11**.
doi:10.1371/journal.pgen.1005585
Zhang L, Ward JD, Cheng Z, Dernburg AF. 2015. The auxin-inducible degradation (AID)
system enables versatile conditional protein depletion in C. elegans. *Development*
142:4374–4384. doi:10.1242/dev.129635
Zhong W, Sternberg PW. 2006. Genome-Wide Prediction of C. elegans Genetic
Interactions. *Science (80-)* **311**:1481–1484.
Zhu J, Fukushige T, McGhee JD, Rothman JH. 1998. Reprogramming of early
embryonic blastomeres into endodermal progenitors by a Caenorhabditis elegans
GATA factor. *Genes Dev* **12**:3809–3814. doi:10.1101/gad.12.24.3809

Zuryn S, Ahier A, Portoso M, White ER, Morin MC, Margueron R, Jarriault S. 2014.
Sequential histone-modifying activities determine the robustness of
transdifferentiation. *Science (80-)* **345**:826–829. doi:10.1126/science.1255885
1266

Figure 2

Medwig-Kinney et al. (2022)

Medwig-Kinney et al. (2022)

mNeonGreen::3xFLAG::AID

loxP

lsy-22

TagRFP-T::3xFLAG::AID

lox2272

Key:

- ■ exon
- ⋁ intron
- ■ spacer
- ■ TEV site
- ■ mNeonGreen flurophore
- ■ TagRFP-T fluorophore
- ■ auxin inducible degron (AID)
- ■ epitope tag (3x FLAG)

100 bp ─